

The physical basis of mollusk shell chiral coiling

Régis Chirat^{a,2}, Alain Goriely^b, and Derek E. Moulton^b

^aCNRS 5276, LGL-TPE, Université Lyon 1, 69622 Villeurbanne Cedex, France; ^bMathematical Institute, University of Oxford, Oxford, UK

This manuscript was compiled on October 7, 2021

Snails are model organisms for studying the genetic, molecular and developmental bases of left-right asymmetry in Bilateria. However, the development of their typical helicospiral shell, present for the last 540 million years in environments as different as the abyss or our gardens, remains poorly understood. Conversely, ammonites typically have a bilaterally symmetric, planispirally coiled shell, with only 1% of 3000 genera displaying either a helicospiral or a meandering asymmetric shell. A comparative analysis suggests that the development of chiral shells in these mollusks is different, and that unlike snails, ammonites with asymmetric shells probably had a bilaterally symmetric body diagnostic of cephalopods. We propose a mathematical model for the growth of shells, taking into account the physical interaction during development between the soft mollusk body and its hard shell. Our model shows that a growth mismatch between the secreted shell tube and a bilaterally symmetric body in ammonites can generate mechanical forces that are balanced by a twist of the body, breaking shell symmetry. In gastropods, where a twist is intrinsic to the body, the same model predicts that helicospiral shells are the most likely shell forms. Our model explains a large diversity of forms and shows that although molluscan shells are incrementally secreted at their opening, the path followed by the shell edge and the resulting form are partly governed by the mechanics of the body inside the shell, a new perspective that explains many aspects of their development and evolution.

coiling | symmetry breaking | chirality | mathematical model | mollusk

Among metazoans, Bilateria are organized along an antero-posterior and a dorso-ventral axis that both define the plane of bilateral symmetry, and the left and right sides of the animal. Although bilaterian animals are externally mostly symmetric, they usually show a consistent left-right asymmetry in internal organs. How left-right symmetry is broken during development raises fundamental questions, such as the functional implications of asymmetry, defective left-right asymmetry leading to severe pathologies in humans; the developmental stage at which asymmetry is initiated; the dominance in most cases of a given direction (e.g. our heart most often to the left side, liver to the right) rather than a random 50/50 ratio; the extent to which left-right symmetry breaking processes have been evolutionarily conserved among Bilateria; how multilevel asymmetries, from molecular, cellular to organismal level, are related to each other; and how consistent asymmetry is generated in a world where no macroscopic process of chemistry or physics can be used to define unequivocally left from right (1–4).

In contrast to most Bilateria, snails display a conspicuous outward asymmetry manifested by a typically dextral (with an opening on the right side when the tip is up) or rarely sinistral helicospiral shell together with marked left-right anatomical asymmetries. The characteristic helicospiral shape of snail shells is a particular kind of chirality, a form being *chiral* if it cannot be superimposed on its mirror image, like our

left and right hands. Shell chirality has intrigued biologists for centuries, and snails have emerged as model organisms to address the genetic and developmental bases of left-right symmetry breaking in Bilateria (5). Chirality in snails is in direct contrast with the shape of most ammonites*, a group of extinct mollusk cephalopods with an external chambered shell that populated the seas for 340 million years and became extinct 66 million years ago. Like the extant chambered *Nautilus*, about 99% of 3000 ammonite genera have *non-chiral*, bilaterally symmetric shells, most often a planispiral or more rarely a straight shell, or a combination of both forms, despite the fact that *Nautilus* and gastropods share the same basic structure of the shell-secreting system (6, 7), and that both empirical and theoretical evidences suggest it was shared by ammonites as well (8, 9). That is, ammonites were likely secreting their shells in the same way as gastropods, and yet producing in the vast majority of cases symmetric shells. The remaining 1% of ammonites are represented by some 40 genera mostly belonging to seven Cretaceous families displaying, at least during a part of their development, an asymmetric, often helicospiral shell (10). Two asymmetric genera are also known in the upper Triassic (11). These rare heteromorph ammonites display the most stunning shell shapes (Fig. 1), generated by a combination of different modes of shell coiling during development. For a long time considered as "aberrant", these forms have marveled and puzzled paleontologists for years. In addition to numerous taxonomic studies, special attention has been paid to the inference of their hydrostratic properties, lifestyle and paleocology (12–14). However a key question of developmental biology remains: what are the symmetry-

Significance Statement

A theoretical model suggests that a mechanically induced twist of the soft body underlies the formation of helicospiral shells in snails and ammonites, and also accounts for the startling and unique meandering shells observed in certain species. This theory addresses fundamental developmental issues of chirality and symmetry breaking: in the case of ammonites, how a bilaterally symmetric body can sometimes secrete a non-symmetric shell; for gastropods, how an intrinsic twist possibly due to the asymmetric development of musculature can provide a mechanical motor for generating a chiral shell. Our model highlights the importance of physical forces in biological development, and sheds light on shell coiling in snails, which have been used for a century as model organisms in genetic research.

DEM and RC conceived the study. DEM and AG devised the mathematical model. Computations were performed by DEM. Comparative approach was performed by RC. DEM collected data on shell coiling. All authors contributed to the writing of the paper.

We have no conflict of interest.

²To whom correspondence should be addressed. E-mail: regis.chirat@univ-lyon1.fr

breaking processes involved in the development of asymmetric shells among representatives of a group overwhelmingly characterized by a well-marked bilateral symmetry diagnostic of the cephalopod body plan?

The relative simplicity of the shell growth process in molluscs – an accretionary process occurring at the current shell opening by the secreting mantle edge – and the diverse and distinct forms that are generated, as described above, make molluscs an excellent case study for investigating symmetry breaking during development, notably in light of recent progress made in developmental biology on this question in the model organism, the pond snail *Lymnaea*. Here we present a comparative analysis between gastropods and ammonites and propose a new unifying model of shell coiling based on the interaction of the animal's soft body with its secreted hard shell. Our model provides a physical explanation for how a bilaterally symmetric ammonite body may secrete on occasion an asymmetric shell, and also addresses within the same framework the ubiquitous formation of helicospiral shells in gastropods, in light of the exception of bilaterally symmetric shells of limpets.

1. Background

A. How snails got their handedness. A direction of shell coiling in snails is overwhelmingly predominant in a given species with more than 90% of snails exhibiting dextral shells (15). For example, only six specimens of sinistral *Cerion* have ever been found among probably millions of specimens examined (16). Pond snails of the genus *Lymnaea* have become model organisms to study the genetic and developmental basis of left-right asymmetry, leading to a model of maternal inheritance in which offspring's handedness is dictated by the mother's genotype (17, 18) by a single maternal locus (19). Gastropods display a spiral cleavage mode of early cell divisions, as do most representatives of the Lophotrochozoa (one of the three super-phyyla of Bilateria). The first sign of chirality in snails is distinguishable in the orientation of the cleavage planes, and handedness may be defined as early as in the first or second blastomere divisions. But temporal and spatial cytoskeletal dynamics for dextral and sinistral embryos are not mirror images of each other, and show a bias towards dextral forms from the early stages of spiral cell division (20). Strikingly, inverting genetically specified third-cleavage directions by mechanically altering the relative orientation of cells leads to snails with inverted handedness, manipulated embryos growing to 'dextralized' sinistral or 'sinistralized' dextral snails (21). This handedness in cleavage acts upstream of the Nodal signalling pathway long known to be involved in left-right asymmetry in vertebrates, and involved in snails too (22). In the quest to discover the long-sought maternally expressed gene determining handedness, a diaphanous-related formin gene has been identified (23, 24), providing a proof for the role of an actin cytoskeleton-regulating protein in determining the arrangement of blastomeres. In summary, left-right asymmetry in snails anatomy originates in cellular architecture. The dynamics of the inherently chiral cytoskeleton governs mechanically the asymmetric behavior of dividing cells at the earliest stage of development and, ultimately, the body and shell handedness. We will show however that if the link between spiral cleavage, body and shell handedness is obvious in the model

organism *Lymnaea* (and probably many other gastropods), the link between cleavage pattern and helicospiral coiling itself is not straightforward and with a single explanation.

B. Ammonites took a weird turn. In contrast to gastropods that display a spiral cleavage typical and ancestral of the molluscan phylum, cephalopods show a bilateral cleavage. In this case, the first cleavage furrow fixates the plane of bilateral symmetry of the animal while the second furrow separates the future anterior and posterior areas (25). Moreover, unlike gastropods, a well-marked bilateral symmetry of the body organization both external and internal (e.g. symmetry and position of paired organs, such as gills, retractor muscles) is a diagnostic feature of the cephalopod body plan (26, 27). A bilateral symmetry characterizes also the shell of about 99% of ammonite genera, and though their soft body organization remains poorly known, muscle attachment marks are also bilaterally symmetric (28), unlike those of snails. We also know that the embryonic shell (ammonitella) is bilaterally symmetric (29), even in heteromorph ammonites with a post-embryonic helicospiral shell (30, 31). Moreover, while chirality in snails is visible at the earliest embryonic stages, the shell of heteromorph ammonites only becomes chiral at a much later stage of development, sometimes well after hatching and organogenesis, i.e. well after the stage at which the anatomical symmetries are established. Note that we reserve the term "heteromorph" for species displaying a non-planar shell despite the fact that a number of bilaterally symmetric species (but with non-overlapping whorls) have been called heteromorphs.



Fig. 1. Heteromorph ammonites with chiral shells. (A) *Turritites costatus* (Cenomanian, France). (B) *Colchidites breistrofferi* (Barremian, Columbia); note the inner helicospiral shell followed by a planispiral stage. (C) *Nipponites mirabilis* (Turonian, Japan). (D) CT scan of a *N. mirabilis* (Upper Cretaceous, Japan) showing the inner planispiral whorls. (E) *Didymoceras stevensoni* (Upper Cretaceous, USA). (F) Two enantiomorphs, sinistral and dextral, of *D. stevensoni* (Upper Cretaceous, USA). (Scale bars, 10 mm. Specimens numbers are given in SI Appendix B).

*When we use the vernacular term "ammonites" we refer to representatives of the cephalopod subclass Ammonoidea.

Heteromorph ammonites with helicospiral shells have

evolved repeatedly from ancestors with bilaterally symmetric, planispiral shells (11, 32, 33). One particularly intriguing feature is the modifications of their shell symmetry during development (Fig. 1B–F). For example, *Didymoceras*, shown in Fig. 1E, displays a bilaterally symmetric shell at the juvenile stage (straight or planispiral), a middle growth stage of asymmetric, helicospiral shell, and a bilaterally symmetric shell portion at maturity. Therefore, the shell shifts from bilaterally symmetric to asymmetric and then back to symmetric. It is difficult to conceive how the anatomical symmetry of the body itself could have shifted in the same way during development. In fact, the morphology of the shell in *Didymoceras* (and genera of other families) shows that during the asymmetric part, the ventral side of the shell runs along the longer helicospiral and the dorsal side on the shorter one (which results in shell edge and ribs oblique to the growth direction), while the left and the right sides run along helicospirals of the same length, i.e., grow at the same rate as in planispiral shells, contrarily to gastropods in which helicospiral shells display a clear left-right asymmetry in growth rate.

In contrast to snails, in which the direction of shell coiling is overwhelmingly predominant in a species, a study of about 1500 specimens of *Didymoceras* shows roughly an equal percentage of dextral and sinistral shells (*D. stevensoni*, n=264, d/s ratio: 47/53; *D. nebrascense*, n=882, d/s ratio: 49/51; *D. cheyennense*, n=338, d/s ratio: 52/48) (34). This roughly 50/50 ratio in handedness has also been reported in other genera of Nostoceratidae (35, 36), Heteroceratidae (37) or Turritidae (38), which suggests that the direction of asymmetry was randomly determined and non-heritable. Indeed, in the case of asymmetry induced by mechanical twisting, as will form the premise of our model, the twisting is equally likely to occur in either direction, and the actual observed directionality would be determined by ‘noise’ in the system and thus unpredictable and non-heritable. Likewise, in the known cases of existing Bilateria displaying a random direction of asymmetry in some anatomical traits, the direction of asymmetry is non-heritable (39).

The genus *Nipponites* displays some of the most startling shapes observed in Nature (Fig. 1C–D). While it seems to be irregularly convoluted at first sight, it is not (40) and the shell actually follows a precise and reproducible developmental sequence. At juvenile stages, *Nipponites* has a planispiral, logarithmically coiled shell with non-overlapping whorls. Then the shell unfolds in a succession of meandering oscillations on each side of the plane of bilateral symmetry of the first planispiral stage, forming alternating dextral and sinistral helical sections of increasing wavelength and amplitude. We refer to this inversion of handedness as a *perversion* following the nomenclature introduced by the mathematician Listing, and used by Maxwell and d’Arcy Thompson (41–44). *Nipponites* is thought to derive from *Eubosttrychoceras* (35), a genus that displays bilaterally symmetric planispiral whorls in the early stages, a middle growth stage with a helicospiral shell (dextral or sinistral in a 50/50 ratio) and a bilaterally symmetric shell segment at maturity. An important contribution in the geometric description of these heteromorph ammonites was made by Okamoto (45–47) who showed that these shapes could be modeled by varying the curvature and torsion of a centerline curve. However, this author assumed that shell coiling was controlled by the orientation of these ammonites in the water

column through an unknown regulatory mechanism.

In summary, comparative data present us with a conundrum: unlike snails, evidence suggests that ammonites had a bilaterally symmetric body diagnostic of the cephalopod body plan but nevertheless sometimes secreted an asymmetric shell. Our goal here is to devise a mathematical model that can elucidate the developmental mechanism of shell coiling and symmetry breaking, and show under what circumstances the different shell forms observed in ammonites and gastropods can be expected, under what conditions a symmetric body can give rise to an asymmetric shell, and how these asymmetric shells can change during development.

2. Model

Shell-building mollusks face strong geometric constraints associated with accretionary growth of their shell: they secrete during their development a shell to which the growing body will have to fit in subsequent stages sometimes several months or years later. For instance, a mean shell growth rate of 0.061mm/day in an immature *Nautilus* (48) implies that the rear of the growing body may be enclosed in a part of the shell tube secreted about 5 years earlier. Our main hypothesis is that any growth mismatch between the soft body and the secreted shell tube in which it resides can generate mechanical stresses balanced by an overall deformation of the body, impacting the geometry of future secretion. A mismatch between different growing parts of an organism plays a fundamental role in the genesis of mechanical forces underlying development and morphogenesis of plants and animals (49, 50), a mechanism involved in molluscan shell morphogenesis as well (8, 9, 51–54). That tissue growth and shell growth may be decoupled from each other is well known in bivalves and gastropods (55, 56). This question has rarely been addressed in *Nautilus*, though the allometric relationships between body and shell growth during sexual maturation has been reported in *N. pompilius* (57). In ammonites the allometric relationships between body and shell growth may be manifested by sometimes considerable variations in body chamber length during development (28), that however did not prevent these animals from regulating their buoyancy, probably due to a flexibility of the mechanisms of buoyancy regulation as in *Nautilus* (58). Our objective is to first investigate whether a mismatch between body and shell growth might account for the symmetry breaking observed in some ammonites, and then to analyse whether the same methodology can consistently explain the helicospiral shell form in gastropods.

As shown in Fig. 2, we model the mollusk body by two elastic rods, one for the ventral side of the animal, and one for the dorsal side. The reference shape of the growing body, i.e. the shape that the soft animal would take if it were removed from its shell, is given by the unstressed shape of these elastic rods, defined by their stress-free reference length and curvature that both evolve throughout development. For ammonites, the natural choice is to assume that the stress-free shape is a planar logarithmic spiral, for which the *growth rates* of the ventral and dorsal sides must be unequal: the ventral side is always growing at a higher rate than the dorsal side to ensure that they form a spiral. However, when the body occupies the shell, the elastic rods are constrained to match the shape of the shell tube that has been so far secreted. The shell shape is determined by both the orientation of the




Fig. 2. Model schematic. A. A mismatch between the growth rate of the body (idealized by dorsal and ventral elastic rods) and the secretion of the shell generates mechanical stress in the animal's body that may be partially alleviated by twisting of the body within the shell tube. B. Three self-similar (i.e. with isometric growth) shell types may be generated from the same secretion parameters: if no twist, a planispiral shell; with constant twist, a helicospiral shell; with oscillatory twist, a meandering shell. C. Planar coiling geometry is captured by two parameters, an expansion rate c_1 (red arrows) and coiling gradient c_2 (blue arrows). The coiling gradient (solid line) follows the dorso-ventral axis, and generates coiling around the orthogonal axis (dashed line). D. Representative shells for the 3 shell types.

animal within the shell and the *secretion rates*. Naturally, the secretion rate on the ventral side is higher than on the dorsal side. If the secretion rates exactly match the body growth rates, then the shell shape will exactly match the logarithmic spiral shape of the body – in this case the body is always in its reference shape, and no stress will be induced in the animal. If, however, the secretion rates do not exactly match the body growth rates, mechanical stress will be induced in the mollusc body, potentially forcing the body to twist within the shell to partially relieve these stresses (Fig. 2A – see also Supplementary Information (SI) Sec 1). If the animal twists within the shell, then the dorso-ventral axis will rotate about

the center line of the shell. Since the growth gradient follows the dorso-ventral axis, the axis about which the shell coils (dashed line in Fig 2C) will also rotate and thus the shell shape will change; in particular, any twist will by construction generate a non-planarity to the center line curve of the shell, i.e. twisting of the animal creates torsion[†] in the shell shape.

There is an interesting feedback at work: the shape of the shell that has been so far secreted dictates the stress in the animal within the tube; mechanical stresses generate a twist of the animal body; and the orientation of the animal dictates the subsequent shape of the shell, which will, in turn create stresses on the growing animal. This two-way coupling between body shape and shell shape makes the problem particularly difficult to solve in general. Here, our approach is to exploit self-similarity, (i.e. isometric growth), which enables us to decouple the influence of mechanical stress on shell shape, and to examine the conditions under which the animal may be predicted to secrete one of three classes of shell: (i) planispiral, (ii) helicospiral, or (iii) meandering. These three shell types can be produced with equivalent secretion rates; the only difference being the orientation of the secretion given by the twisting of the animal within the shell. In particular, there is no twist in the case of the planispiral shell, a constant twist rate (with respect to whorl) for the helicospiral, and an oscillatory twist rate in the case of the meandering shell – see Fig. 2B. Therefore, assuming that the secretion rates and body growth rates are given, and that the only degree of freedom is the twisting, we can posit that the degree and form of twist by the animal will be the one that minimizes the mechanical energy of the contorted body; and thus the shell actually produced by the animal is the one corresponding to that energy-minimizing twist. With the assumption of self-similarity, we do not need to solve for the shape at each point in time based on the current orientation, rather we find an energy-minimizing twist for a single (arbitrary) time point, and the self-similar assumption implies that the same twist will be selected throughout development.

The analysis above requires a description of the shell geometry, a characterisation of the internal energy for the soft body, and a procedure for energy minimization. Full details are provided in the SI; below we briefly outline the modeling components.

A. Geometry. The geometry of the shell can be described by a set of only 5 parameters (see SI Sec. 2), $\{c_1, c_2, d_0, d_1, d_2\}$ illustrated in Fig. 2C,D. Here c_1 describes the aperture expansion rate, c_2 describes the growth/secretion gradient, i.e. the difference in growth/secretion between the ventral and dorsal sides, that produces coiling, and the parameters d_i characterise twisting. In particular, d_0 describes a constant twist, while d_1 and d_2 are, respectively, the amplitude and frequency of an oscillatory twist – equivalently, these correspond to an oscillation in the torsion of the shell centerline. In terms of these parameters, a planispiral shell corresponds to setting $d_0 = d_1 = d_2 = 0$; a helicospiral shell is constructed by setting $d_1 = d_2 = 0$, with $d_0 \neq 0$, and a meandering shell is constructed by setting $d_0 = 0$, with d_1 and d_2 both non-zero (in each case c_1 and c_2 should be non-zero). Representative shells are shown in Fig. 2D, with parameters provided in SI

[†] By torsion we refer to the mathematical definition of a measure of the twisting out of the plane of curvature of a space curve.

Sec. 2E. In summary, the shell shape is characterised by one of the following parameter sets:

- Planispiral: $\mathcal{S}_p = \{c_1, c_2\}$
- Helicospiral: $\mathcal{S}_h = \{c_1, c_2, d_0\}$
- Meandering: $\mathcal{S}_m = \{c_1, c_2, d_1, d_2\}$

Our model is premised on the distinction between the geometry of the shell and the geometry of the body. When considering a bilaterally symmetric body, as with ammonites, the reference shape of the body is assumed to be planar, i.e. there is no intrinsic twist; thus the body is described by only two parameters, $\hat{\mathcal{S}} = \{\hat{c}_1, \hat{c}_2\}$, where we use the overhats to denote a reference quantity for the body. A mismatch between shell and body shape is then captured by any difference between $\{c_1, c_2\}$ and $\{\hat{c}_1, \hat{c}_2\}$, while any twist of the body is described by the parameters $\{d_0, d_1, d_2\}$. However, another type of mismatch between body and shell may occur: the animal may be growing in such a way as to match the shell shape it is secreting, but at a faster or slower rate. This type of mismatch is accounted for by a scaling parameter ξ of arc length between the body and shell:

$$\hat{t} = \xi t, \quad [1]$$

where $\hat{t}$ is the arc length of the centerline corresponding to the body, t is arc length attached to the shell, $\xi > 1$ means the body is growing faster than the shell and conversely, for $\xi < 1$, the shell is growing faster.

B. Mechanical energy. Given a set of parameters for both the shell and the body, we constrain the body to fit in the shell with the dorsal and ventral elastic rods situated on opposing sides of the shell tube and the ventral rod following the point of longest arc length. We then compute the mechanical energy in each of the rods by summing the energy contributions due to stretching, bending, and twisting (for details see SI Sec. 3), employing a standard quadratic energy, with particular care required to account for the difference in arc length between the centerline of the shell tube and the ventral and dorsal sides.

C. Energy minimization. Initially, we assume that the body growth and secretion rates are fixed through development for a given specimen, with the only degree of freedom being the twisting of the animal within the shell. This assumption is a sufficient and necessary condition for construction of a self-similar shell. Though we note this is at best an approximation: the growth and secretion vary to some degree in most shells (We explore in Section E below the consequence of a variation through development in these rates). Therefore, we fix the body parameters $\hat{\mathcal{S}} = \{\hat{c}_1, \hat{c}_2\}$, the shell parameters $\{c_1, c_2\}$, and the stretch mismatch factor ξ . We also require defining the values of stiffness moduli $\{K_1, K_2, K_3\}$ which characterize the resistance to stretching, bending, and twisting of the body, respectively. The energy $\mathcal{E}$ can then be expressed as a function only of twisting (SI Sec. 3B), i.e.

$$\mathcal{E} = \mathcal{E}(d_0, d_1, d_2).$$

For the planispiral shell, there is no twist, and the energy is $\mathcal{E}_p = \mathcal{E}(0, 0, 0)$. The helicospiral shell has energy $\mathcal{E}_h(d_0) = \mathcal{E}(d_0, 0, 0)$. As discussed in SI Sec. 3C, the most consistent

approach to energy minimization is to fix d_2 ; based on geometric considerations we fix the oscillation frequency as $d_2 = 0.8$ and define the meandering shell energy $\mathcal{E}_m(d_1) = \mathcal{E}(0, d_1, 0.8)$. The energy landscape is complex, varying both with the shell type and degree of mismatch imposed between body growth and secretion. Conceptually, the case that is of most interest is when the body growth rate exceeds the secretion rate, which causes the animal's body to be in compression. In this case, by examining the 3 components of the energy (see SI Sec. 3E), a general trend emerges that shows there are values of the stiffness parameters K_i for which any of the three shell types can be an energy minimizer if sufficient compression is generated. The other case, secretion outpacing the body growth, requires the body to stretch during shell secretion; then the body will be in tension, and in such cases the planar shell was always found to be the energy minimizer.

To demonstrate this range of energy minimizers, we portray the energy landscape via a morphological phase space in Fig. 3, constructed by fixing the geometric parameters, with a small degree of imposed mismatch, then sweeping over a range of mechanical parameters and determining for each parameter choice the shell with the minimum energy. Two such plots appear in Fig. 3, with the energy-minimizing shell type denoted by color: green for planispiral, blue for meandering, and red for helicospiral. The coiling parameters $\{\hat{c}_1, \hat{c}_2, c_1, c_2, \xi\}$ are chosen to correspond to sample values for a typical planispiral (Fig. 3A) and meandering (Fig. 3B) shell. We then sweep over the mechanical stiffness ratios K_1/K_2 and K_3/K_2 . For instance, a point in the lower right corner denotes a body with mechanical structure that has high resistance to stretching but low resistance to twisting. In both cases, stiffness ratios exist for which each of the three shell types is predicted. In particular, when the parameters correspond to a typical planispiral shell, the planispiral shell type is the energy minimizer for most stiffness ratios, while when the coiling parameters correspond to sample values for a meandering shell (Fig. 3B) with two different coiling gradients, the heteromorph shell types are energy minimizers for a much wider range of parameter space.

It is important to note that we did not include the steric constraint that prohibits self intersection of the shell tube with previous whorls. This means that some of the mechanically favorable shells are not geometrically possible. In particular, in Fig. 3B the planispiral shell has significant overlap. While some degree of overlap is a feature found in almost all planispiral ammonites (see Sec. 3), it is interesting to observe that with the secretion rates such that the overlap leaves little room for the mollusk body in the planispiral shell tube, there is a significant increase in mechanical favorability of the twisted, non-overlapping, shells.

D. Data comparison. Our model assumes that heteromorph ammonites emerge due to a mechanically induced twisting of the body, meaning that at the level of body growth and secretion, there is no difference between these shells and the far more typical planispiral ammonite. This feature enables us to test the model quantitatively: for each choice of coiling parameters $\{\hat{c}_1, \hat{c}_2, c_1, c_2, \xi\}$, we define the likelihood of finding a meandering or helicospiral shell by sweeping over possible stiffness ratios, and determining the percentage of parameter space for which each shell type is an energy minimizer. Such a calculation appears in Fig. 4A. Here we have




Fig. 3. Morphological phase space, sweeping over stiffness ratios K_1/K_2 (stretching to bending) and K_3/K_2 (twisting to bending). For each value of stiffness ratios, the energy minimizing shell type – planispiral (green), helicospiral (red), or meandering (blue) – is computed, with energy minimizing coiling values and corresponding shell forms indicated by colorbar. The shells in each phase space have equivalent coiling and expansion parameters, matching those in the planar green shell, differing only in the type and degree of twist. Body growth and secretion values are (A): $\hat{c}_1 = 0.02$, $\hat{c}_2 = 0.2$, $\hat{d}_0 = 0$, $c_1 = 0.02$, $c_2 = 0.25$, $\xi = 1.0$, and (B): $\hat{c}_1 = 0.02$, $\hat{c}_2 = 0.323$, $\hat{d}_0 = 0$, $c_1 = 0.02$, $c_2 = 0.4$, $\xi = 1.0$.

set $\hat{c}_1 = c_1$, $\hat{c}_2 = c_2$, and fixed $\xi = 1.075$, corresponding to an undersecreting shell (compressed body), but with body and shell shape matching. For each point in the c_1 - c_2 plane, we compute the energy minimizer over a range of 100 values of stiffness ratios, using the same range as in Fig 3. We then color that point with RGB value corresponding to the percentage of helicospiral (red), planispiral (green), and meandering (blue) energy minimizers. The dashed lines separate regions where each shell type is the overall winner. This plot gives an indication of where we would expect to find (and not find) meandering and helicospiral shells. In particular, the model predicts that coiling gradient is far more relevant than expansion rate, with meandering shells most likely in the coiling gradient range $c_2 \in [0.4, 0.6]$, and helicospiral most likely for




Fig. 4. (A) A phase space of coiling parameters is created by sweeping over mechanical stiffness ratios and computing the energy minimizing shell, then coloring the point using RGB value corresponding to the percent of energy minimizers of each type – planispiral=green, helicospiral=red, meandering=blue. Data points for 19 meandering and 17 helicospiral shells are plotted using extracted coiling parameters. Inset: a phase space with decreased compression factor. (B)-(E): sample shell images (Scale bars, 10 mm) and simulated shells with extracted parameters, corresponding to the indicated points.

$c_2 \lesssim 0.4$, while for large coiling gradient, the planispiral shell is by far the most likely shape. To test these predictions, we have extracted the coiling parameters (c_1, c_2) from a set of 19 meandering (*Nipponites mirabilis*) and 17 helicospiral shells (*Eubostyrchoceras japonicum*). These appear as the red and blue data points in Fig. 4A, and show broad agreement with the model prediction. The best fit shells for the indicated data points appear in Fig. 4B-E; shells (real and simulated) for all data points and all extracted parameter values are provided in SI Sec. 4.

While Fig. 4 provides strong evidence in favour of the mechanical twisting hypothesis, we must be careful with its interpretation. It would be incorrect to conclude that planispiral ammonite shells are only likely to be found on the right side of the diagram, as in fact planispiral ammonites may be found over the entire range of the coiling parameters. Here we emphasize that the twisting only occurs if there is a mismatch between body growth and shell secretion, characterized in this calculation by setting $\xi = 1.075$, meaning that the reference shape of the body is 7.5% longer than the shell tube it is secreting. Without some form of mismatch the body is stress free in the planar state and thus the planispiral shell is always mechanically favorable. Even with a reduced mismatch, the regions in which meandering and helicospiral shells are predicted become much smaller: a sample morphospace with ξ decreased to 1.025 is shown as the inset in Fig. 4; here the planispiral shell is the most favorable shape for all values of coiling parameters. The model thus predicts that most ammonites secreted a planispiral shell due to low or no mismatch. It is for this reason that we do not include data points for planispiral shells in Fig. 4; the point of the computation is not to predict the presence of planispiral shells, but rather to predict where heteromorph shells will appear when the

necessary ingredient of a mismatch is present.

E. Varying shell type through development. Observe that the twisting parameters do not appear in Fig. 4. Thus, while the helicospiral and meandering shells occupy much of the same region of the coiling parameter space, the difference in form comes from the simple difference between a constant twist rate in the case of the helicospiral shell and an oscillatory twist in the case of the meandering shell. It is worth highlighting that such distinctively different forms may be mechanically favorable in the same region of this (2D) coiling space; which may explain why some shells display both types of coiling at different life stages (59) and why *Nipponites* shares many diagnostic characteristics with the coexistent *Eubostriochoceras* from which it derives (35).




Fig. 5. Simulation and images of (A) *Didymoceras nebrascense* and (B) *Nostoceras malagasyense*, obtained by varying the mismatch and stiffness parameters during development, causing transitions in development between planispiral (green), meandering (blue), and helicospiral (red). Scale bars, 10 mm. Shell specimen info and other model parameters provided in SI.

In our model, a transition in shell form can be generated by a change in mismatch and/or stiffness parameters during development. This is demonstrated in Fig. 5, in which we provide simulations of *Didymoceras nebrascense* (A) and *Nostoceras malagasyense* (B). The shell in (A) was obtained by first varying the arc length mismatch parameter ξ , causing a transition in the juvenile stage from planispiral to meandering to helicospiral; and then varying the coiling and stiffness parameters in the late stage of development, which generates the reverse transition from helicospiral to meandering to planispiral (see details in SI Sec. 5). A similar transition produces the shell in (B), though without the juvenile transition missing due to breakage in our specimen. Although we can only speculate on the biological significance of these parameter changes, it should be noted that shell coiling changes in the last stages of development of many ammonites are associated with sexual maturation, which in *Nautilus*, are associated with modifications of growth of the shell and body parts (60). Our study of heteromorph ammonites illustrates also the clear difference between a purely geometric simulation of shell coiling and a model that includes explicitly developmental mechanisms and physical constraints. Indeed, while it is possible to simulate a developmental transition between a helicospiral and planispiral

stage with coaxial coiling, our model shows that this coaxiality is mechanically unlikely. As evident in Figs 3 and 4A, and demonstrated more thoroughly in SI Sec. 6, the regions of parameter space in which the helicospiral and planispiral shells are mechanically favorable are always separated by a region in which the meandering shell is favorable. Therefore, if a change in shell type occurs during development due to a continuous change in parameters, our model predicts that a transition from helicospiral to planispiral must always pass through an intermediate meandering stage which, by construction, will reorient the coiling axis. This rule is consistent with the fact that, to our knowledge, helicospiral and planispiral stages are never strictly coaxial in heteromorph ammonites, the coiling axes can even be at right angles to each other (Fig. 1B,E). This prediction is an example of a developmental constraint imposed by mechanics of morphogenesis (see (9) for a discussion of this concept).

3. A new twist on shell coiling

Although heteromorph ammonites with chiral shells represent only about 1% of 3000 genera, their geometric diversity surpasses that of the other ammonites, which probably lies in the fact that they have non-overlapping whorls. In gastropods, a whorl partially dictates the growth path of the next overlapping whorl (61, 62). The mantle secretes an overlapping layer on the previous whorl to which it adheres and when this attachment zone is partially or totally lost, the coiling geometry is quantitatively modified (63). Whorl overlap played a role in ammonites too (64) and in some way, constrained the range of possible morphologies in restricting the degrees of freedom of the growing system. For instance, a shell of the kind of *Didymoceras* generated by different coiling geometry during development, could not be achieved with overlapping whorls. But then, what are the regulation mechanisms of shell coiling in the non-overlapping case?

Since mollusk shells are incrementally secreted along their opening edge, it seems logical that their coiling geometry could be fully understood in light of growth regulating processes localized at the secreting mantle edge only. This idea has motivated all theoretical models of shell coiling and experimental approaches as well, but is confronted with an issue especially obvious in the case of heteromorph ammonites. One puzzling aspect of their morphogenesis is indeed the mechanisms that govern the three-dimensional path followed by the secreting mantle edge, resulting in highly convoluted forms. Theoretical models predict that an incremental rotation of the growing front underlies the development of helicospiral shells (45, 65, 66). Yet, to our knowledge, no mechanism localized at the mantle edge can trigger this movement. Our model suggests that this incremental rotation may be naturally triggered by a mechanical twist of the body, resulting from a mismatch between body and shell growth. An important conclusion may then be drawn from the study of these heteromorph ammonites: although the form of the shell corresponds only to a spatiotemporal record of accretionary growth at its edge, the three dimensional path followed by the secreting mantle edge is partly governed by the mechanics of the body inside the shell. Whereas, it is now clear that some ornamentation patterns in mollusk shells emerge as the result of mechanical forces at the secreting mantle margin (8, 9, 51–54), our study shows that the mechanical interactions between body and shell

may also play a key-role in the regulation of shell coiling.

This mechanical hypothesis explains a number of puzzling characteristics of these ammonites, notably how they secreted asymmetric shells while keeping a bilaterally symmetric body diagnostic of cephalopods. With the same bilaterally symmetric growth gradient at the shell edge, an asymmetric or symmetric shell may be secreted depending on whether the bilaterally symmetric body is twisted or not. This mechanical twist is recorded by the angular offset between the ventral siphuncle in the posterior part of the body chamber and the anterior ventral zone toward the shell edge (67). In a sinistral *Turrilites*, the ventral siphuncle is shifted toward the right side of the shell tube, while dorsal muscle scars are shifted toward the opposing left side (see (68) pl.18 fig.1-3). This mechanical twist explains also why asymmetric shells may develop after hatching, well after organogenesis and the formation of the plane of bilateral symmetry of the body. Further, the modifications of shell symmetry during development such as the shifts seen in some genera from bilaterally symmetric, to asymmetric and to symmetric again, reflect changes in mechanical strains affecting the bilaterally symmetric body. The fact that heteromorph ammonites with asymmetric shells have repeatedly evolved from ancestors with bilaterally symmetric shells is also consistent with this ahistorical generic mechanism.

In our model, the mechanical energy is equivalent for twisting in either direction. A twist of a bilaterally symmetric body is thus consistent with a random, non-heritable direction of shell handedness, with right and left-handed coiling arising with equal probability. However, representatives of the family Turritidae (Fig. 1A) show another puzzling evolutionary trend to our knowledge unique in the fossil record, and that may be interpreted for the first time in light of our approach. In the genus *Mariella* from South Africa and Texas, Albian species are dextral or sinistral in a 50/50 ratio while all Cenomanian species are sinistral (38, 69). Thus, directional asymmetry arose from ancestors where left-right asymmetry was random. Similar evolutionary patterns in current phallostethid fishes and fiddler crabs have been interpreted as an “unconventional mode” of evolution (“phenotype precedes genotype”), the idea being that phenotypic variation (right or left-handed) arose before genetic mechanisms controlling a given direction of asymmetry (39). But this interpretation depends on the way phenotypic characters are defined. Mechanical forces may generate helicospiral coiling, and though they are growth dependent and modulated by genetic and molecular processes so that their outcome cannot be described as “phenotype first”, they may equiprobably produce dextral or sinistral forms. Directional asymmetry, on the other hand, requires a consistent bias toward one side, as in physical systems generating helices (70). In light of randomization of visceral asymmetry in mutant mice, an original two components abstract system has been proposed to explain how left-right asymmetry might arise in Bilateria (71): a generic process (a reaction-diffusion system in the original hypothesis) producing random asymmetry at the cellular and multicellular level, can be consistently biased toward a direction by a mechanism that converts molecular to cellular asymmetry. Likewise the fixation of sinistral shells in Turritidae can be interpreted in light of a two components process: a generic, mechanical process generating helicospiral shells with no preferred handedness in ancestral forms, and another one (that unfortunately

will remain unknown) introducing a bias toward the leftward coiling in descendant species. An analogous situation has been described in the case of cardiac development in amniotes in which a buckling instability twists the straight cardiac tube into a helical loop with random handedness, while molecular and cellular mechanisms introduce a bias that, except in rare mutants, consistently triggers a rightward looping (72).

Finally, our approach may explain the development of rare paleozoic nautiloids with helicospiral shells (73). It may also shed new light on abnormal shell growth in ammonites with whorls overlapping, although the mechanical influence of this trait, probably dependant on the degree of overlapping, is not included in our model due to the additional theoretical difficulties that it would raise. In *Nautilus*, epizoans growing fixed on the outer surface of the shell may perturbate the growth of the next whorl, slowing or inhibiting the forward movement of the animal’s body (74), a process that could generate compression in the growing body. Our model suggests that this compression may generate meandering or helicospiral shells, similar to the abnormal forms described in slightly overlapping planispiral ammonites encrusted by epizoans (75).

4. How snails coil their shell

Much progress has been made on the genetic and molecular processes that set the left or right handedness of the asymmetric body in snails, but an important point rarely acknowledged is that the mechanisms underlying the development of helicospiral shells themselves remain poorly understood. First, the link between the body and shell handedness is not straightforward, contrarily to what may be reported in light of the development of the model organism, the pond snail *Lymnaea*. This genus is orthostrophic, which means that the body handedness corresponds to the shell handedness. But in hyperstrophic species, anatomically dextral animals have sinistral shell and *vice versa*, while in more complex cases called heterostrophy, shell handedness changes after hatching (76). Moreover, although limpets show a dextral cleavage pattern, a right expression of *nodal* (22), and are right-handed in their body anatomy, both their embryonic and post-embryonic shell is cone-shaped and bilaterally symmetric (77). The asymmetric development of gastropods is further complicated by a rotation which occurs during larval development and which moves the visceral mass, mantle, and shell at 180° with respect to the head and foot (this rotation is, confusingly, referred to as “torsion” in the literature but does not describe the coiling torsion of the shell). But this process cannot be unequivocally linked to helicospiral coiling since limpets also experience such a rotation (78). Furthermore, a left/right asymmetric gradient of the Dpp (decapentaplegic) protein (79) or an asymmetric cellular growth pattern in the mantle edge (80) cannot explain the incremental rotation of the growing front generating helicospiral shells. Comparative anatomy of limpets and helicospiral species suggests a possible mechanism.

In helicospiral species, the shell-muscle system is helically coiled around and anchored to the axial columella of the shell three-quarters to two whorls back from the aperture, and extends into the foot (81). In cone-like limpets, muscle runs dorso-ventrally and attaches to the inner shell surface in a horseshoe-shaped muscle scar bilaterally symmetric on both sides of the body (82). While “true limpets” belong to the order Patellogastropoda, limpet-shaped shells have

convergently evolved in not-closely-related species belonging to the four other gastropod orders predominantly helicospiral. In all cases, evolutionary changes from helicospiral to bilaterally symmetric limpet-shaped shells are correlated with a drastic modification of the shell-muscle system, from a coiled muscle attached on one side to the axial columella of the helicospiral shells, to a horseshoe-shaped muscle bilaterally symmetric on both sides of the body typical of true limpets (82–84). These repeated modifications of both shell coiling and muscle-shell system during evolution suggest that both characters could be developmentally correlated, and that bilateral asymmetry of the muscle-shell system could induce a twist of the body in species with helicospiral shells. Although this hypothesis remains to be tested experimentally, our theoretical framework already allows us to explore the effect of the intrinsic twist of the body on the shell form.




Fig. 6. Morphological phase space for gastropods for varying intrinsic twist $\hat{d}_0 \in \{0.1, 0.3\}$ and mismatch parameter $\xi \in \{0.95, 1.05\}$; with a sweep over stiffness ratios K_1/K_2 (stretching to bending) and K_3/K_2 (twisting to bending) at each point. Energy minimizing shell type is indicated by color, with red helicospiral shell the energy minimizer in every single case. Energy minimizing twisting value d_0 and corresponding shell type indicated by colorbar. Body growth and secretion values are (A): $\hat{c}_1 = c_1 = 0.06$, $\hat{c}_2 = c_2 = 0.9$.

A. Modeling gastropod form. The observations above point to the possible role of intrinsic twist, $\hat{d}_0$. This parameter is geometrically equivalent to the helicospiral shell parameter d_0 , but is intrinsic to the animal’s body. Due to the bilateral symmetry of ammonites, $\hat{d}_0 = 0$ for all shells, while we explore in this section the mechanical consequences of $\hat{d}_0 \neq 0$ for gastropods (note that anatomically, the blue rod is anterior and the red rod is posterior in the case of a gastropod). For given body parameters $\{\hat{c}_1, \hat{c}_2, \hat{d}_0\}$ and secretion parameters $\{c_1, c_2, \xi\}$, we compute as before the effect of a mismatch by determining the energy minimizing shell. The energy minimization proceeds in the same way as outlined above, with the appropriate variation to the twisting energy (see SI Sec. 7). Naturally, in this case if there is no mismatch the animal will secrete a helicospiral shell that matches its helicospiral body shape. The question then is whether other shell types might be mechanically favorable, given a mismatch. To answer this question, we proceed as before, sweeping over a range of mechanical stiffness ratios and comparing the total mechanical energy in the planispiral

shell with that of the helicospiral and meandering shells for which the energy is minimum. The result appears in Fig. 6. Here we have fixed the coiling parameters, and varied both the degree of mismatch via the parameter ξ and the degree of intrinsic twist via $\hat{d}_0$ and plotted the resulting phase space. We find that in every single case, the helicospiral shell is the energy minimizer. We have colored each point by the energy minimizing twist value d_0 , with sample shells appearing next to the colorbar.

An analysis of different base coiling parameters shows that the planispiral shell can also be the energy minimizer, but only in cases of a stretched body, whereas meandering shells are never found to be favorable (see SI Sec. 7). In Fig. 7, we show the morphological phase space in the case of low intrinsic twist and $\xi = 0.95$ (stretched body) for coiling parameters matching Planorbidae, a small-sized aquatic pulmonate gastropod family. As shown in Fig. 7, the model predicts a greater likelihood of planispiral shells at low expansion rate, with the helicospiral shell being the dominant form at higher expansion rate. This trend in variation is consistently observed among Planorbidae, and was already described in 1867 in the first phylogenetic tree based on fossil evidence (85), just eight years after Darwin’s Origin of Species. Comparing Figs 3 and 7 also highlights an interesting mechanical duality: tension can cause an asymmetric body to take on a symmetric shape, while compression can cause a symmetric body to take an asymmetric shape.

Our mechanical model predicts that in the presence of an intrinsic body twist, helicospiral shells are strongly favoured. It is well known that, once pulled out of its helicospiral shell, the body of gastropods remains helicospiral. This might seem logical since the body has grown and had to fit inside the shell. We suggest however that the body is helicospiral because it is intrinsically twisted, possibly due to the asymmetric development of the muscle-shell system, and that this intrinsic twist provides the motor for the incremental rotation of the secreting mantle edge required to generate the helicospiral shell.

Our approach may help to explain many aspects of shell coiling that are difficult to interpret in terms of relative growth rates at the shell edge only, such as the development of hyperstrophic or heterostrophic species (76), of heteromorph snails involving a rotation of the body inside the shell (86), of pharmaceutically induced “banana-shaped” Planorbidae (87), or of abnormal helicospiral individuals of this family found in the wild (63), among other examples. Further, our study may inform the long standing debate about the assignment of Bellerophonitida to gastropods (88, 89). These constitute an extinct order of mollusk of uncertain systematic position (Cambrian-Triassic) characterized by a planispiral, rapidly expanding shell, but interestingly displaying a pair of muscle scars symmetric on both sides of the shell (90) suggesting an untwisted body.

5. Conclusion

The natural world is overflowing with strikingly regular spiral, helical, and helicospiral shapes, such as keratin fibers, collagen assembly, DNA molecules, spiral bacteria, tendrils, climbing vines, seed pods, sheep horn, the cochlea and umbilical cords among others (70, 91–96). Such structures often develop as the result of fundamental mechanical forces generated by




Fig. 7. A sequence of morphological phase spaces for varying expansion rate in the case of a low intrinsic twist ($\hat{d}_0 = 0.1$) and stretched body ($\xi = 0.95$) with energy minimizing shells indicated by color. The model predicts an increased likelihood of planispiral shell at lower expansion rate, characteristic of trends in Planorbidae. Coiling parameters (c_1, c_2) from left to right: (0.01, 0.2), (0.02, 0.37), (0.08, 0.48), (0.1, 0.64). Shell images are (left to right): *Anisus leucostoma*, *Planorbis planorbis*, *Menetus dilatatus*, *Bulinus albus*.

a mismatch between different parts (97–99). While most previous theoretical or experimental approaches have tried to relate global shell geometry only to the growth occurring at the shell edge, our study highlights how the position and mechanics of the body inside the shell can serve to regulate its morphogenesis. Through consideration of the orientation and mechanical energy of the soft body constrained by the shell in which it resides, we have identified a basic physical mechanism that explains the origin and diversity in form of shell coiling in mollusks. This includes, in the case of ammonites, a natural explanation for the development of an asymmetric shell by a bilaterally symmetric animal, and in the case of gastropods a mechanical motor for the generation of helicospiral shells due to an intrinsic twist possibly connected to the asymmetric development of musculature. Our model also explains the meandering shells of *Nipponites*, one of the most startling forms in Nature. It would be tempting to see in this unique morphology an arbitrary quirk of evolution. In fact, similar geometric forms consisting of alternating helical sections of opposite handedness separated by multiple perversions are known to occur in bacterial shape and flagella, cellulose fibres,

vine tendrils and also telephone cords due to a combination of curvature-induced instability and geometric constraints (44, p. 150). In our study, the meandering form of *Nipponites* emerges as the energetically favorable path of an oscillatory twist of the animal's body in the shell. As ammonites have been extinct for 66 million years, it is of course impossible to confirm with certainty their body symmetry. However, as we have shown that it is mechanically unfavorable for an animal with an asymmetric body to secrete a symmetric or meandering shell, our study provides strong new evidence for a bilaterally symmetric body, including in these heteromorph ammonites; this highlights the potential value of mechanics in deciphering the form of the soft body parts of a long extinct animal. Likewise, snails, which have long been used as model organisms for genetic studies, could also constitute an excellent model for studying the canalizing role of mechanics in the genesis, variation, and evolution of biological forms.

Data availability. A Mathematica notebook containing model details and calculations has been deposited in the Oxford University Research Archive

Acknowledgements We are grateful to the people who provided us access to paleontological data: E. Robert, C. Salaviale (Univ. Lyon1, France), F. Giraud (Univ. Grenoble, France), D. Berthet (Mus. confluentes, Lyon, France), H. Châtelier (ammonites.fr, France), C. Scheskie (Denver MNS, USA), S. Luallin, S. Jorgensen (Western Interior Pal. Soc., USA), C. Chen (JAMSTEC, Japan), D. Aiba (Mikasa Mus., Japan), Y. Shigeta (NMNS, Japan), H. Maeda, Y. Ito (Kyushu Univ. Mus., Japan), Y. Shiino (Pal. Soc. Japan), and C. Serra (Mus. CosmoCaixa, Spain). This work was supported by the Engineering and Physical Sciences Research Council grant EP/R020205/1

1. Vandenberg LN, Lemire JM, Levin M (2013) It's never too early to get it right: a conserved role for the cytoskeleton in left-right asymmetry. *Communicative & integrative biology* 6(6):12586–91.

2. Inaki M, Liu J, Matsuno K (2016) Cell chirality: its origin and roles in left–right asymmetric development. *Philosophical Transactions of the Royal Society B: Biological Sciences* 371(1710):20150403.

3. Lebreton G, et al. (2018) Molecular to organismal chirality is induced by the conserved myosin 1D. *Science* 362(6417):949–952.

4. Juan T, et al. (2018) Myosin1D is an evolutionarily conserved regulator of animal left–right asymmetry. *Nature communications* 9(1):1–12.

5. Davison A (2020) Flipping shells! unwinding Lr asymmetry in mirror-image molluscs. *Trends in Genetics* 36(3):189–202.

6. Westermann B, Schmidtberg H, Beuerlein K (2005) Functional morphology of the mantle of *Nautilus pompilius* (mollusca, cephalopoda). *Journal of morphology* 264(3):277–285.

7. Jackson DJ, et al. (2006) A rapidly evolving secretome builds and patterns a sea shell. *BMC biology* 4(1):1–10.

8. Moulton D, Goriely A, Chirat R (2015) The morpho-mechanical basis of ammonite form. *Journal of theoretical biology* 364:220–230.

9. Erlich A, Moulton DE, Goriely A, Chirat R (2016) Morphomechanics and developmental constraints in the evolution of ammonites shell form. *Journal of Experimental Zoology Part B: Molecular and Developmental Evolution* 326(7):437–450.

10. Wright CW (1996) Cretaceous ammonioidea. *Treatise on Invertebrate Paleontology, Part L, Mollusca 4, Revised*.

11. Wiedmann J (1969) The heteromorphs and ammonoid extinction. *Biological Reviews* 44(4):563–602.

12. Landman NH, et al. (2012) Methane seeps as ammonite habitats in the us western interior seaway revealed by isotopic analyses of well-preserved shell material. *Geology* 40(6):507–510.

13. Peterman DJ, Yacobucci MM, Larson NL, Ciampaglio C, Linn T (2020) A method to the madness: ontogenetic changes in the hydrostatic properties of *Didymoceras* (Nostoceratidae: Ammonoidea). *Paleobiology* 46(2):237–258.

14. Peterman DJ, Mikami T, Inoue S (2020) The balancing act of *Nipponites mirabilis* (Nostoceratidae, Ammonoidea): Managing hydrostatics throughout a complex ontogeny. *Plos one* 15(8):e0235180.

15. Van Batenburg F, Gittenberger E (1996) Ease of fixation of a change in coiling: computer experiments on chirality in snails. *Heredity* 76(3):278–286.

16. Gould SJ, Young ND, Kasson B (1985) The consequences of being different: sinistral coiling in cerion. *Evolution* 39(6):1364–1379.

17. Sturtevant AH (1923) Inheritance of direction of coiling in *Limnaea*. *Science* 58(1501):269–270.

18. Freeman G, Lundelius JW (1992) Evolutionary implications of the mode of d quadrant specification in coelomates with spiral cleavage. *Journal of Evolutionary Biology* 5(2):205–247.

19. Hosoi Y, Harada Y, Kuroda R (2003) Construction of a backcross progeny collection of dextral and sinistral individuals of a freshwater gastropod, *Limnaea stagnalis*. *Development genes and evolution* 213(4):193–198.

20. Shibazaki Y, Shimizu M, Kuroda R (2004) Body handedness is directed by genetically determined cytoskeletal dynamics in the early embryo. *Current Biology* 14(16):1462–1467.

21. Kuroda R, Endo B, Abe M, Shimizu M (2009) Chiral blastomere arrangement dictates zygotic left–right asymmetry pathway in snails. *Nature* 462(7274):790–794.

22. Grande C, Patel NH (2009) Nodal signalling is involved in left–right asymmetry in snails. *Nature* 457(7232):1007–1011.

23. Davison A, et al. (2016) Formin is associated with left-right asymmetry in the pond snail and the frog. *Current Biology* 26(5):654–660.

24. Abe M, Kuroda R (2019) The development of CRISPR for a mollusc establishes the formin *Lsdia1* as the long-sought gene for snail dextral/sinistral coiling. *Development* 146(9).

25. Boletzky S (1988) Cephalopod development and evolutionary concepts. *The Mollusca* 12:185–202.

26. Shigeno S, Takeno S, Boletzky Sv (2010) The origins of cephalopod body plans: a geometrical and developmental basis for the evolution of vertebrate-like organ systems. *Cephalopods: Present and Past* 1:23–34.

27. Hanlon R, Vecchione M, Allcock L (2018) *Octopus, squid, and cuttlefish: a visual, scientific guide to the oceans, of most advanced invertebrates..* University of Chicago Press.

28. Doguzhaeva LA, Mapes RH (2015) The body chamber length variations and muscle and mantle attachments in ammonoids. In *Ammonoid Paleobiology: From anatomy to ecology*. (Springer), pp. 545–584.

29. Landman NH, Tanabe K, Shigeta Y (1996) Ammonoid embryonic development. In *Ammonoid paleobiology*. (Springer), pp. 343–405.

30. Tanabe K, Obata I, Futakami M (1981) Early shell morphology in some upper cretaceous heteromorph ammonites. *Transactions and proceedings of the Paleontological Society of Japan. New series*. (Palaeontological Society of Japan), Vol. 124, pp. 215–234.

31. Druschits V (1977) The structure of the ammonitella and the direct development of ammonites. *Paleont. Jour.* 2: 188–199.

32. Monks N (1999) Cladistic analysis of albian heteromorph ammonites. *Paleontology* 42(5):907–925.

33. Hoffmann R, et al. (2021) Recent advances in heteromorph ammonoid palaeobiology. *Biological Reviews* 96(2):576–610.

34. Kennedy WJ, Landman N, Cobban WA, Scott G (2000) Late campanian (cretaceous) heteromorph ammonites from the western interior of the united states. *Bulletin of the American Museum of Natural History* 2000(251):1–86.

35. Okamoto T (1989) Comparative morphology of *Nipponites* and *Eubostrioceras* (cretaceous nostoceratids). *Transactions and proceedings of the Paleontological Society of Japan. New series*. (Palaeontological Society of Japan), Vol.154, pp. 117–139.

36. Klinger HC, Kennedy WJ, Grulke WE (2007) New and little-known Nostoceratidae and Diplomoceratidae (cephalopoda: Ammonoidea) from Madagascar. *African Natural History* 3(1):89–115.

37. Klinger HC, Kakabadze MV, Kennedy WJ (1984) Upper barremian (cretaceous) heteroceratid ammonites from south africa and the caucasus and their-paleobiogeographic significance. *Journal of Molluscan Studies* 50(1):43–60.

38. Klinger H, Kennedy W (1978) Turrititidae (Cretaceous Ammonoidea) from South Africa, with a discussion of the evolution and limits of the family. *Journal of Molluscan Studies* 44:1–48.

39. Palmer AR (2004) Symmetry breaking and the evolution of development. *Science* 306(5697):828–833.

40. Yabe H (1903) Cretaceous Cephalopoda from the Hokkaido *Journal of the College of Science Tokyo* Vol. 20, 2: 1–45.

41. Listing JB (1847) Vorstudien über topologie. *Göttinger Studien* 1.

42. Clark WELG, Medawar PB (1945) *Essays on Growth and Form presented to D'Arcy Wentworth Thompson*. (Oxford University Press).

43. Maxwell JC (1892) *A treatise on electricity and magnetism*. (Clarendon, Oxford).

44. Goriely A (2017) *The Mathematics and Mechanics of Biological Growth*. (Springer Verlag, New York).

45. Okamoto T (1988) Analysis of heteromorph ammonoids by differential geometry. *Paleontology* 31(1):35–52.

46. Okamoto T (1988) Changes in life orientation during the ontogeny of some heteromorph ammonoids. *Paleontology* 31:281–294.

47. Okamoto T (1988) Developmental regulation and morphological saltation in the heteromorph ammonite *Nipponites*. *Paleobiology* pp. 272–286.

48. Dunstan AJ, Ward PD, Marshall NJ (2011) *Nautilus pompilius* life history and demographics at the osprey reef seamount, coral sea, Australia. *PLoS One* 6(2):e16312.

49. Ambrosi D, et al. (2019) Growth and remodelling of living tissues: perspectives, challenges and opportunities. *Journal of the Royal Society Interface* 16(157):20190233.

50. Hamant O, Saunders T (2020) Shaping organs: shared structural principles across kingdoms. *Annual Review of Cell and Developmental Biology* 36:385–410.

51. Chirat R, Moulton DE, Goriely A (2013) Mechanical basis of morphogenesis and convergent evolution of spiny seashells. *Proceedings of the National Academy of Sciences* 110(15):6015–6020.

52. Erlich A, Howell R, Goriely A, Chirat R, Moulton D (2018) Mechanical feedback in seashell growth and form. *ANZIAM Journal* 59(4).

53. Rudraraju S, Moulton DE, Chirat R, Goriely A, Garikipati K (2019) A computational framework for the morpho-elastic development of molluscan shells by surface and volume growth. *PLoS computational biology* 15(7):e1007213.

54. Moulton DE, Goriely A, Chirat R (2020) Mechanics unlocks the morphogenetic puzzle of interlocking bivalved shells. *Proceedings of the National Academy of Sciences* 117(1):43–51.

55. Lewis DE, Cerrato RM (1997) Growth uncoupling and the relationship between shell growth and metabolism in the soft shell clam *Mya arenaria*. *Marine Ecology Progress Series* 158:177–189.

56. Bourdeau PE, Johansson F (2012) Predator-induced morphological defences as by-products of prey behaviour: a review and prospectus. *Oikos* 121(8):1175–1190.

57. Hayasaka S, Ōki K, Tanabe K, Saisho T, Shinomiya A (2010) On the habitat of *Nautilus pompilius* in Tanon Strait (Philippines) and the Fiji Islands. In *Nautilus The Biology and Paleobiology of a Living Fossil*. (Springer), pp. 179–200.

58. Ward PD (1987) *The natural history of Nautilus*. (Allen & Unwin).

59. Matsumoto T (1977) Some heteromorph ammonites from the cretaceous of Hokkaido. *Mem. Fac. Sci., Kyushu Univ., Ser. D, Geol.* 23:303–366.

60. Collins D, Ward PD (2010) Adolescent growth and maturity in *Nautilus*. In *Nautilus The Biology and Paleobiology of a Living Fossil*. (Springer), pp. 421–432.

61. Hutchinson J (1989) Control of gastropod shell shape; the role of the preceding whorl. *Journal of Theoretical Biology* 140(4):431–444.

62. Checa AG, Jiménez-Jiménez AP, Rivas P (1998) Regulation of spiral coiling in the terrestrial gastropod *Sphincterochila*: An experimental test of the road-holding model. *Journal of Morphology* 235(3):249–257.

63. Clewing C, Riedel F, Wilke T, Albrecht C (2015) Ecophenotypic plasticity leads to extraordinary gastropod shells found on the "roof of the world". *Ecology and Evolution* 5(14):2966–2979.

64. Ubukata T, Tanabe K, Shigeta Y, Maeda H, Mapes RH (2008) Piggyback whorls: a new theoretical morphologic model reveals constructional linkages among morphological characters in ammonoids. *Acta Paleontologica Polonica* 53(1):113–128.

65. Moulton DE, Goriely A, Chirat R (2012) Mechanical growth and morphogenesis of seashells. *Journal of theoretical biology* 311:69–79.

66. Moulton DE, Goriely A (2014) Surface growth kinematics via local curve evolution. *Journal of mathematical biology* 68(1):81–108.

67. Ward P (1979) Functional morphology of cretaceous helically-coiled ammonite shells. *Paleobiology* 5(4):415–422.

68. Crick GC (1898) On the muscular attachment of the animal to its shell in some fossil Cephalopoda (Ammonoidea). *Transactions of the Linnean Society of London. 2nd Series. Zoology* 7(4):71–113.

69. Clark DL (1965) *Heteromorph ammonoids from the Albian and Cenomanian of Texas and adjacent areas*. (Geological Society of America) Vol. 95.

70. Armon S, Efrati E, Kupferman R, Sharon E (2011) Geometry and mechanics in the opening of chiral seed pods. *Science* 333(6050):1726–1730.

71. Brown NA, Wolpert L (1990) The development of handedness in left/right asymmetry. *Development* 109(1):1–9.

72. Desgrange A, Le Garrec JF, Meilhac SM (2018) Left-right asymmetry in heart development and disease: forming the right loop. *Development* 145(22).

73. Teichert C (1964) Mollusca 3. Cephalopoda General Features - Endoceratoidea - Actinoceratoidea - Nautiloidea - Bactritoidea. *Treatise on invertebrate paleontology. Part K*, pp. 1–519.

74. Arnold J (1985) Shell growth, trauma, and repair as an indicator of life history for *Nautilus*. *The Veliger* 27(4):386–396.

75. Checa AG, Okamoto T, Keupp H (2002) Abnormalities as natural experiments: a morphogenetic model for coiling regulation in planispiral ammonites. *Paleobiology* 28(1):127–138.

76. Okumura T, et al. (2008) The development and evolution of left-right asymmetry in invertebrates: Lessons from drosophila and snails. *Developmental dynamics* 237(12):3497–3515.

77. Kay MC, Emlet RB (2002) Laboratory spawning, larval development, and metamorphosis of the limpets *Lottia digitalis* and *Lottia asmi* (Patellogastropoda, Lottiidae). *Invertebrate Biology* 121(1):11–24.

78. Wanninger A, Ruthensteiner B, Haszprunar G (2000) Torsion in *Patella caerulea* (Mollusca, Patellogastropoda): ontogenetic process, timing, and mechanisms. *Invertebrate Biology* 119(2):177–187.

79. Shimizu K, et al. (2013) Left-right asymmetric expression of dpp in the mantle of gastropods correlates with asymmetric shell coiling. *EvoDevo* 4(1):1–7.

80. Johnson AB, Fogel NS, Lambert JD (2019) Growth and morphogenesis of the gastropod shell. *Proceedings of the National Academy of Sciences* 116(14):6878–6883.

81. Price RM (2003) Columellar muscle of neogastropods: muscle attachment and the function of columellar folds. *The Biological Bulletin* 205(3):351–366.

82. Heller J (2015) *Sea snails. A natural history*. (Springer).

83. Yamamori L, Kato M (2018) Morphological and ecological adaptation of limpet-shaped top shells (Gastropoda, Trochidae, Fossarininae) to wave-swept rock reef habitats. *PLoS one* 13(8):e0197719.

84. Tseng R, Dayrat BA (2014) Anatomical redescription of the limpet-like marine pulmonate *Trimusculus reticulatus* (Sowerby, 1835). *Veliger* 51(4):194–207.

85. Rasser MW (2013) Darwin's dilemma: the steinheim snails' point of view. *Zoosystematics and Evolution* 89(1):13–20.

86. Liew TS, Kok AC, Schilthuizen M, Urdy S (2014) On growth and form of a heteromorphic terrestrial snail: *Plectostoma concinnum* (Fulton, 1901) (Mollusca, Gastropoda, Diplommatinidae). *PeerJ PrePrints* 2:e289v1.

87. Baynes A, et al. (2019) Early embryonic exposure of freshwater gastropods to pharmaceutical 5-alpha-reductase inhibitors results in a surprising open-coiled "banana-shaped" shell. *Scientific reports* 9(1):1–12.

88. Ponder WF, Lindberg DR (1997) Towards a phylogeny of gastropod molluscs: an analysis using morphological characters. *Zoological Journal of the Linnean society* 119(2):83–265.

89. Bouchet P, Rocroi J-P (2005) Classification and nomenclator of gastropod families. *Malacologia* 147: 1-397.

90. Knight JB (1947) Bellerophon muscle scars. *Journal of Paleontology* pp. 264–267.

91. Goriely A (2004) Knotted umbilical cords. In *Physical and Numerical Models in Knot Theory Including Their Application to the Life Sciences*, eds. Calvo JA, Stasiak A, Rawdon E. (World Scientific, Singapore), pp. 109–126.

92. McMillen T, Goriely A (2002) Tendril perversion in intrinsically curved rods. *Journal of Nonlinear Science* 12(3):241–281.

93. Goldstein RE, Goriely A, Hubber G, Wolgemuth C (2000) Bistable helices. *Phys. Rev. Lett.* 84.

94. Wolgemuth CW, et al. (2005) How to make a spiral bacterium. *Physical biology* 2(3):189.

95. Goriely A, Neukirch S (2006) Mechanics of climbing and attachment in twining plants. *Phys. Rev. Lett.* 97(18):184302.

96. Moulton DE, Oliveri H, Goriely A (2020) Multiscale integration of environmental stimuli in plant tropism produces complex behaviors. *Proceedings of the National Academy of Sciences* 117(51):32226–32237.

97. Lessinnes T, Moulton DE, Goriely A (2017) Morphoelastic rods. Part II: Growing birods. *J. Mech. Phys. Solids* 100:147–196.

98. van Manen T, Janbaz S, Zadpoor AA (2018) Programming the shape-shifting of flat soft matter. *Materials Today* 21(2):144–163.

99. Moulton DE, Lessinnes T, Goriely A (2020) Morphoelastic rods iii: Differential growth and curvature generation in elastic filaments. *Journal of the Mechanics and Physics of Solids* p. 104022.