

Callose Deposition during Pollen Development

Callose is a cell wall component that is dynamically deposited and degraded during pollen development. Thanks to a new article investigating pollen formation in cotton (*Gossypium hirsutum*), we now know that a pollen-specific protein regulates callose deposition by inhibiting the action of a transcription factor, WRKY15 (Li et al., 2020).

Callose is a polymer formed primarily of Glc units connected by β 1-3-linkages. It is transiently deposited at sites of wounding, plasmodesmata, and pollen cell walls (Chen and Kim, 2009). Callose is also frequently present as a transient wall component in newly formed cell plates during cytokinesis of mitotically dividing plant cells (Scherp et al., 2001).

In the anthers of many angiosperms, a transient callose-rich cell wall is laid down, surrounding and separating each microsporocyte and callose is also deposited on the outer pollen wall (Blackmore et al., 2007; De Storme and Geelen, 2013). Callose is thought to isolate microspores from one another and is considered an impermeable barrier. Additionally, in many species that form these callose-rich walls, mutants with defective callose production do not create viable pollen (De Storme and Geelen, 2013). Nevertheless, the essential role of callose is somewhat contested, as there is some evidence that large molecules can still cross the callose wall and some species do not deposit callose at this early meiotic stage (Scott et al., 2004). Despite this, callose deposition during meiosis has been observed across land plants including in mosses, liverworts, hornworts, and lycophytes (Flowers, 2018).

Li et al. (2020) have identified a gene expressed in cotton pollen, *POLLEN SPECIFIC PROTEIN 231* (*PSP231*). They uncovered a sequence of regulatory interactions leading from *PSP231* through to callose

deposition phenotypes. *PSP231* is part of the *SKU5-SIMILAR* (*SKS*) family of genes, which are frequently glycosylated but otherwise have unknown biochemical functions. Some *SKS* genes are involved in root development, and others have been shown to be expressed in pollen (Albani et al., 1992; Wittink et al., 2000; Zhou, 2019b).

Studying gametogenesis is notoriously difficult as mutants are frequently sterile and it is even more difficult in species such as *Gossypium* spp., in which established tools for pollen germination research are not yet available. Despite this, the authors persevered by using in vitro methods as well as RNA interference (RNAi) and heterologous expression.

Li et al. (2020) found that *PSP231*-RNAi lines were defective in male gametogenesis, forming pollen grains that were frequently withered and misshapen. Over-expressing *PSP231* in tobacco (*Nicotiana tabacum*) pollen also resulted in reduced pollen germination. These findings illustrated an essential role for *PSP231* and suggested that its expression had to be carefully balanced to achieve its correct effects.

To understand what *PSP231* might be doing in cotton, Li et al. (2020) analyzed the transcriptome of wild-type and *PSP231*-RNAi pollen. They found that two callose synthase genes were downregulated in the knockdown line compared to wild type. Furthermore, in the pollen of the tobacco *PSP231* overexpressors, aniline-blue staining highlighted increased callose deposition both in the newly formed wall dividing the vegetative and germ cells and in mature pollen.

In the absence of *PSP231*, *GhWRKY15* showed increased expression. The *GhWRKY15* protein is a transcriptional repressor and is able to bind to the promoters of the two callose synthase genes, consistent with the decreased expression levels of these genes.




Figure 1. Schematic illustrating features of male gametogenesis and genetic interactions. Adapted from figure 8 in Li et al. (2020).

Li et al. (2020) also used a variety of in vitro and in vivo methods to establish that PSP231 forms a protein–protein complex with an RNA-binding protein, GhRBPL1. To complete the links in the chain, Li et al. (2020) then demonstrated that GhRBPL1 is able to specifically bind to the sense strand of *GhWRKY15* mRNA. Taking the data together, they suggested a model in which PSP231 binds to and inhibits the actions of GhRBPL1. When de-repressed, GhRBPL1 is thought to bind to and stabilize *GhWRKY15* mRNA. In turn, GhWRKY15 represses transcription of callose synthase genes. When PSP231 is present, GhWRKY15 is therefore downregulated and callose synthesis can take place (Fig. 1). As *PSP231* is largely expressed at later stages of pollen development, it is likely that additional regulators also exist that influence the early-stage deposition of callose.

Remarkably, altered callose deposition was not only seen in plants, but also in yeast (*Schizosaccharomyces pombe*) cells overexpressing *PSP231*. In fission yeast, dividing cells have callose in their new cell plates. When *S. pombe* overexpressed *PSP231*, cells were more often seen at late stages of cell division compared to wild type, exhibiting callose-rich cell plates. These cells also showed growth retardation and decreased viability, likely due to the inability to degrade the excessively callosic septa.

While *SKS-like* genes have been known for some time, their roles have not been extensively studied. In *Arabidopsis* (*Arabidopsis thaliana*), *SKU5* regulates root growth and mutants show twisted roots (Sedbrook et al., 2002). Similarly to *PSP231*, *Arabidopsis sks* mutants show modified cell geometry phenotypes with increased cell wall thickness in seedling roots (Zhou, 2019b). Previous research has demonstrated that *SKS* genes are frequently glycosylphosphatidylinositol (GPI)-anchored. Supporting this, Li et al. (2020) found that *PSP231* is glycosylated. It is therefore likely that the *PSP231* is anchored to lipid membranes within the cell, as has been demonstrated for other *SKS* proteins (Sedbrook et al., 2002; Wittink et al., 2000; Zhou, 2019a). In *Arabidopsis*, disruption of GPI reduces the prevalence of GPI-anchored proteins including *SKU5*, and results in nonviable pollen and general disruption to cell wall synthesis (Gillmor et al., 2005). It will be interesting to see whether this type of post-translational modification is also essential for *PSP231* function.

This is the one of the first instances in which a regulatory network with an *SKS-like* gene has been established, and it may open up further avenues of research for the roles of *SKS* genes in other species and tissues. The ability of *PSP231* to substantially impact not only plant pollen development but also yeast cell division

suggests that there may be common regulatory processes for callose synthesis across species and kingdoms. This has already been hinted at by the observation that both the plant and fungal genes that synthesize callose share significant sequence similarity (Latgé, 2007). Li et al. (2020) have established an important set of regulatory interactions governing the development and cellular structure of pollen, which will lead to a greater understanding of male reproductive development in plants.

Madeleine Seale^{1,2}

ORCID ID: 0000-0002-8924-3943

Department of Plant Sciences,
University of Oxford, Oxford, OX1 3RB,
United Kingdom

LITERATURE CITED

- Albani D, Sardana R, Robert LS, Altosaar I, Arnison PG, Fabijanski SF (1992) A *Brassica napus* gene family which shows sequence similarity to ascorbate oxidase is expressed in developing pollen: Molecular characterization and analysis of promoter activity in transgenic tobacco plants. *Plant J* 2: 331–342
- Blackmore S, Wortley AH, Skvarla JJ, Rowley JR (2007) Pollen wall development in flowering plants. *New Phytol* 174: 483–498
- Chen X-Y, Kim J-Y (2009) Callose synthesis in higher plants. *Plant Signal Behav* 4: 489–492
- De Storme N, Geelen D (2013) Cytokinesis in plant male meiosis. *Plant Signal Behav* 8: e23394
- Flowers N (2018) Sporogenesis and callose localization in Anthocerotophyta. PhD Thesis. Illinois University, Carbondale, IL
- Gillmor CS, Lukowitz W, Brininstool G, Sedbrook JC, Hamann T, Poindexter P, Somerville C (2005) Glycosylphosphatidylinositol-anchored proteins are required for cell wall synthesis and morphogenesis in *Arabidopsis*. *Plant Cell* 17: 1128–1140
- Latgé JP (2007) The cell wall: A carbohydrate armour for the fungal cell. *Mol Microbiol* 66: 279–290
- Li Y, Li L, Wang Y, Wang Y-C, Wang NN, Lu R, Wu YW, Li XB (2020) Pollen-specific protein *PSP231* activates callose synthesis to govern male gametogenesis and pollen germination. *Plant Physiol* 184: 1024–1041
- Scherp P, Grotha R, Kutschera U (2001) Occurrence and phylogenetic significance of cytokinesis-related callose in green algae, bryophytes, ferns and seed plants. *Plant Cell Rep* 20: 143–149
- Scott RJ, Spielman M, Dickinson HG (2004) Stamen structure and function. *Plant Cell* 16: S46 LP-S60
- Sedbrook JC, Carroll KL, Hung KF, Masson PH, Somerville CR (2002) The *Arabidopsis* *SKU5* gene encodes an extracellular glycosyl phosphatidylinositol-anchored glycoprotein involved in directional root growth. *Plant Cell* 14: 1635–1648
- Wittink FRA, Knuiman B, Derksen J, Čapková V, Twell D, Schrauwen JAM, Wullems GJ (2000) The pollen-specific gene *Ntp303* encodes a 69-kDa glycoprotein associated with the vegetative membranes and the cell wall. *Sex Plant Reprod* 12: 276–284
- Zhou K (2019a) Glycosylphosphatidylinositol-anchored proteins in *Arabidopsis* and one of their common roles in signaling transduction. *Front Plant Sci* 10: 1022
- Zhou K (2019b) GPI-anchored *SKS* proteins regulate root development through controlling cell polar expansion and cell wall synthesis. *Biochem Biophys Res Commun* 509: 119–124

¹Senior author.

²Author for contact: maddy.seale@plants.ox.ac.uk.
www.plantphysiol.org/cgi/doi/10.1104/pp.20.01143