

1 Human influence on climate in the 2014 Southern 2 England winter floods and their impacts

3 Nathalie Schaller^{1,2}, Alison L. Kay³, Rob Lamb^{4,10}, Neil R. Massey², Geert Jan van
4 Oldenborgh⁵, Friederike E. L. Otto², Sarah N. Sparrow², Robert Vautard⁶, Pascal
5 Yiou⁶, Ian Ashpole², Andy Bowery⁷, Susan M. Crooks³, Karsten Haustein², Chris
6 Huntingford³, William J. Ingram^{1,8}, Richard G. Jones^{2,8}, Tim Legg⁸, Jonathan Miller⁷,
7 Jessica Skeggs⁹, David Wallom⁷, Antje Weisheimer^{1,11,12}, Simon Wilson⁸, Peter A.
8 Stott⁸ & Myles R. Allen^{2,1}

9 1: Department of Physics, Atmospheric Oceanic and Planetary Physics, University of
10 Oxford, Oxford OX1 3PU, UK

11 2: Environmental Change Institute, University of Oxford, South Parks Road, Oxford
12 OX1 3QY, UK

13 3: Centre for Ecology and Hydrology, Benson Lane, Wallingford OX10 8BB, UK

14 4: JBA Trust, South Barn, Broughton Hall, Skipton BD23 3AE, UK

15 5: Koninklijk Nederlands Meteorologisch Instituut, 3730 AE De Bilt, The Netherlands

16 6: Laboratoire des Sciences du Climat et de l'Environnement & IPSL, UMR CEA-
17 CNRS-UVSQ, 91191 Gif-sur-Yvette, France

18 7: Oxford e-Research Centre, 7 Keble Road, Oxford OX1 3QG, UK

19 8: Met Office Hadley Centre, FitzRoy Road, Exeter EX1 3PB, UK

20 9: JBA Risk Management Ltd., South Barn, Broughton Hall, Skipton BD23 3AE, UK

21 10: Lancaster Environment Centre, Lancaster University, Lancaster LA1 4YQ, UK

22 11: Department of Physics, National Centre for Atmospheric Science (NCAS),
23 University of Oxford, Oxford OX1 3PU, UK

24 12: European Centre for Medium-Range Weather Forecasts (ECMWF), Reading
25 RG2 9AX, UK

26 **A succession of storms reaching Southern England in the winter of**
27 **2013/2014 caused severe floods and £451 million insured losses. In a**
28 **large ensemble of climate model simulations, we find that, as well as**
29 **increasing the amount of moisture the atmosphere can hold,**
30 **anthropogenic warming caused a small but significant increase in the**
31 **number of January days with westerly flow, both of which increased**
32 **extreme precipitation. Hydrological modelling indicates this increased**
33 **extreme 30-day-average Thames river flows, and slightly increased daily**
34 **peak flows, consistent with the understanding of the catchment's**
35 **sensitivity to longer-duration precipitation and changes in the role of**
36 **snowmelt. Consequently, flood risk mapping shows a small increase in**
37 **properties in the Thames catchment potentially at risk of riverine**
38 **flooding, with a substantial range of uncertainty, demonstrating the**
39 **importance of explicit modelling of impacts and relatively subtle**
40 **changes in weather-related risks when quantifying present-day effects**
41 **of human influence on climate.**

42 The winter of 2013/2014, and January in particular, saw above-average
43 precipitation over England and Wales^{1,2} and below-average sea level
44 pressure (SLP) in the North Atlantic north and west of the British Isles (Fig.
45 1a-b). This persistent synoptic situation was associated with a near-
46 continuous succession of low-pressure systems moving in from the Atlantic
47 and across Southern England¹. Like the very wet autumn of 2000 in England
48 and Wales³, this winter was characterized by an anomalous eastward
49 extension of the jet stream (Fig. 2a). This persistent atmospheric circulation
50 pattern resulted in extreme precipitation (Supplementary Fig. 1), flooding and
51 storm surges in large parts of Southern England and Wales, with serious
52 consequences for infrastructure and livelihoods¹. 18,700 flood insurance
53 claims were reported⁴, leading to £451 million insured losses in Southern
54 England. Although not unprecedented, this was a significant event;
55 comparative UK insurance losses⁵ in recent history include flooding in the
56 summer of 2007, which cost £3 billion, the 2005 floods in Carlisle (£272
57 million) and Cumbrian floods in November 2009 (£174 million). Daily total
58 precipitation, recorded since 1767 at the Radcliffe Observatory in Oxford
59 (continuously since 1827), shows January 2014, as well as winter 2013/2014,
60 precipitation set a record (Fig. 3a). Sustained high precipitation amounts
61 during the whole winter led to this record, rather than a few very wet days,
62 and none of the 5-day precipitation averages over the three winter months
63 was a record (Fig. 3b). Similarly, while Thames' daily peak river flows were
64 not exceptional, the 30-day peak flow was the second highest since
65 measurements began in 1883 (Supplementary Fig. 10). Whether
66 anthropogenic climate change contributed to this event was much discussed
67 at the time, with the British Prime Minister David Cameron telling Parliament "I
68 very much suspect that it is"⁶. Although in a chaotic system a single extreme
69 event cannot be attributed to changes in boundary conditions⁷, the change in
70 risk of a class of extremes in the current climate relative to a climate unaltered
71 by anthropogenic greenhouse gas (GHG) emissions can be estimated⁸. This
72 study uses a range of models and observations to estimate anthropogenic
73 influence on the risk of experiencing such atmospheric flow and precipitation,

separating thermodynamic and dynamic factors. To estimate the impacts of climate change, we use a hydrological model to calculate the anthropogenic changes in risk in peak flows of the river Thames. Finally, with detailed flood maps of the Thames basin we estimate the number of properties put at additional risk of flooding by anthropogenic GHG emissions.

1. Experimental setup and model evaluation

We use the citizen-science project “weather@home”⁹ to produce an ensemble of 134,354 simulations of possible weather under current climate and under counterfactual conditions as might have been without human influence on atmospheric composition. This project uses spare CPU time on volunteers’ personal computers to run the regional climate model (RCM) HadRM3P nested in the HadAM3P atmospheric general circulation climate model (AGCM)⁹ driven with prescribed sea surface temperatures (SSTs) and sea ice concentration (SIC). The RCM covers Europe and the Eastern North Atlantic Ocean, at a spatial resolution of about 50 km. 17,367 winters (December, January and February: DJF) were simulated under observed 2013/2014 GHG concentrations, SSTs and SIC (“Actual Conditions”). Initial conditions are perturbed slightly for each ensemble member on December 1 to give a different realisation of the winter weather⁹. The remaining simulations (“Natural”) represent different estimates of conditions that might have occurred in a world without past emissions of GHGs and other pollutants including sulphate aerosol precursors. In the Natural simulations, atmospheric composition is set to pre-industrial, the maximum well-observed SIC is used (DJF 1986/1987, the precise choice is unimportant: Supplementary Fig. 5) and estimated anthropogenic SST change patterns are removed from observed DJF 2013/2014 SSTs. To account for the uncertainty in our estimates of a world without anthropogenic influence, 11 different patterns are calculated from GCM simulations of the Coupled Model Intercomparison Project phase 5 (CMIP5)¹⁰ (Supplementary Information Section 2). We include all CMIP5 models with at least 3 ensemble members available regardless of

105 how well their simulated trends fit observed SST trends in the North Atlantic,
106 to provide a conservative estimate of uncertainty.

107 We consider January precipitation and SLP, with Southern England
108 Precipitation (SEP) averaged over land grid points in 50°–52°N, 6.5°W–2°E.
109 Simulated anomalies for Actual Conditions ensemble members with the
110 wettest 1% SEP, i.e. return periods of 1-in-100-year and rarer, are
111 comparable to observations of January 2014, consistent with previous model
112 evaluation⁹ (Fig. 1c-d). The mean climate of the RCM has a wet bias of ~0.4
113 mm day⁻¹ in January over Southern England⁹ but most RCM simulations for
114 January 2014 show smaller anomalies than observed, and show a weaker
115 SLP pattern for the same precipitation anomaly (Fig. 1c-d). On average, the
116 Actual Conditions simulations reproduce a stronger jet stream, compared to
117 the 1986-2011 climatology, of January 2014 in the North Atlantic (ERA-
118 Interim¹¹, Fig. 2a-b), suggesting some potential predictability for the enhanced
119 jet stream of January 2014. The differences in SSTs, SICs and atmospheric
120 composition between Actual Conditions and Natural simulations lead to an
121 increase of up to 0.5 mm day⁻¹ in the wettest 1% ensemble members for
122 January SEP (Supplementary Fig. 8). While a warmer atmosphere holds more
123 water vapour, causing an increase in risk of heavy winter rainfall, a dynamic
124 effect where anthropogenic forcings altered probability of occurrence of the
125 atmospheric circulation that favoured the winter 2013/2014 conditions¹² is also
126 possible. Disentangling whether a change in precipitation extremes is caused
127 by anthropogenic forcing via thermodynamic or dynamic processes remains a
128 major challenge^{3,13}, which we now address.

130 2. Relationships between atmospheric circulation and precipitation

131 To investigate the joint changes in precipitation and circulation, the observed
132 and modelled Atlantic flows are classified into four main weather regimes
133 using a classical cluster analysis¹⁴⁻¹⁶ (Supplementary Information Section 3).
134 During January 2014, the atmospheric circulation was classified on 26 out of
135 31 days as “zonal regime” (ZO). This is the highest ZO occupancy in January

since 1871 (Supplementary Fig. 7f). The winter as a whole also set a record (70% of days in ZO), in both cases with record low pressure northwest of Scotland (20°W, 60°N, the centre of the anomaly associated with the ZO regime, Supplementary Fig. 7b, and where SLP is strongly associated with SEP, Supplementary Fig. 2a). In the following we use these two circulation indices - the January average sea level pressure Northwest of Scotland and the number of days spent in the ZO regime - to characterize the circulation and its changes. In the RCM simulations, anthropogenic forcing is found to affect the joint distribution of precipitation in Southern England with both low pressure and ZO occupancy (Figs 4a-b). The joint distribution of the Actual Conditions ensemble is stretched towards lower pressures (higher ZO occupancies) and higher precipitation compared to the pooled Natural ensemble, while the other end of the joint distribution (lower precipitation and higher pressure) is unaffected. The model shows more low-pressure systems and days in the ZO regime in the current climate than in the counterfactual world without human influence on climate, with correspondingly higher monthly precipitation amounts in Southern England. Fig. 5a shows the return period (i.e. the inverse of the tail probability) of the pressure index values for all ensembles. Comparing return periods in the Actual Conditions and Natural ensembles gives the change in risk. The risk of experiencing a 1-in-100-year low-pressure event Northwest of Scotland in the Actual Conditions ensemble increases by a best estimate of 55% due to climate change (with an uncertainty range of no change to over 120% increase). We have used all ensemble members available from the individual Natural simulations as our best estimate (Supplementary Information Section 2 discusses this choice and sensitivity of our results to it).

This change in risk is of similar amplitude to the difference from the 1986-2011 climatology (grey dots) and implies that the anomalous circulation in January 2014 was both a response to the January 2014 SSTs and sea ice concentration, hence potentially predictable, and influenced by anthropogenic forcing.

Even with these SSTs, however, it still appears to have been relatively unlikely: monthly ZO occupancy of 24 days have on average a return period of 1-in-151-year in the pre-industrial climate (uncertainty range: 1-in-104-year to 1-in-230-year), which changes to 1-in-113-year due to climate change (Fig. 5b). Flows under the ZO regime have an eastward-extended jet stream towards European coasts. A higher frequency of ZO regimes is thus consistent with recent studies of the effect of climate change on limiting large latitudinal fluctuations of the jet-stream¹⁷, thereby favouring occupancy of regimes like ZO, in line with Ref 18. Our results are not inconsistent with studies reporting insignificant future mean changes of the North Annular Mode or North Atlantic Oscillation (NAM/NAO)^{17,19} because we are detecting a weak signal in extremes, in a much larger ensemble than previously used.

To examine changes in the frequency of extreme precipitation events, we use RCM outputs for the Southern England region and average observations from 8 stations in this region with long records in Met Office archives. Using the time series from 1912-2013 for these 8 stations alone (Supplementary Fig. 1) and treating individual months as independent, the best estimate of the return period of January 2014 SEP is around 85 years (90% confidence interval of 35-550 years; Fig. 5c). Observed Southern England monthly winter precipitation amounts show no statistically significant change in extreme values between the recent period and a century ago using a simple statistical model, although the sensitivity of the test is low (Supplementary Information Section 4).

In the large RCM ensemble, the best estimate for the overall change in risk of a 1-in-100-year January precipitation event pooling all the Natural simulations is an increase of 43%, with a range from no change to 164% increase associated with uncertainty in the pattern of anthropogenic warming (Fig. 5d). Supplementary Fig. 5 shows that this uncertainty is mainly caused by the difference in SSTs and is not affected by the exact choice of sea ice conditions. The potential predictability identified for the pressure index (Fig. 5a) does not appear to extend to precipitation for which the climatological

distribution is consistent with the Actual Conditions ensemble. The Natural ensemble with the smallest change in risk of 1-in-100-year precipitation between Actual and Natural conditions (with the SST pattern from the HadGEM2-ES model) also shows a similar jet stream anomaly to the Actual Conditions ensemble (Fig. 2c). There is no such anomaly in the Natural ensemble showing the greatest change in this risk (with the SST pattern from the CCSM4 model, Fig. 2d).

The 11 estimates of the SST response to anthropogenic forcing allow a statistical investigation into the drivers of the dynamic response. The obvious candidate indices are the global-mean warming and the anthropogenic change in meridional SST gradient upstream (since mid-latitude cyclones are forced by the atmospheric meridional temperature gradient). We represent the latter by the difference between the regions 30°N–50°N, 40°W–0°W and 50°N–70°N, 40°W–0°W. Correlations across the 11 anthropogenic SST change patterns of the change in 1-in-100-year SEP with the global-mean warming and the anthropogenic change in meridional SST gradient upstream are 0.73 and 0.74 (in line with previous studies^{20,21}) respectively (notional *p*-value of 0.01 using a *t*-test). As expected, these two indices are themselves correlated, but only at 0.44 (*p*-value of 0.17). Dividing the change in gradient by the global-mean warming to leave only the pattern of change, not of its magnitude, still gives a correlation of 0.69 (*p*-value of 0.02). Thus both large-scale warming and local dynamical changes play a role.

We estimate the relative importance of thermodynamic and dynamic effects by using the pressure index as a proxy for the changes in circulation between Actual Conditions and Natural simulations. By weighting the Natural ensemble members to match the distribution of the Actual Conditions pressure index values (Fig. 4c and Supplementary Information Section 5) and applying this weighting to the precipitation index to remove the effect of circulation (Fig. 4d), we estimate that the increase in risk of the 1-in-100-year precipitation event due to anthropogenic forcing is caused approximately 2/3 by

thermodynamic changes, and approximately 1/3 by circulation changes.
Previous studies such as Ref 3 found only a thermodynamic influence.

3. Attributing changes in impacts

Modelled precipitation and temperature are fed into the CLASSIC hydrological model of the Thames catchment²², spun up with observed data from January 2010 to early December 2013 (Supplementary Information Section 6).

For a 1-in-100-year event in the hydrological model, anthropogenic climate change increased the modelled risk of 30-day peak river flows at Kingston by a best estimate value of 21% (uncertainty range: -12% to 133%) (Fig. 5e). For daily peak flows however, the increase was a best estimate of 4% (uncertainty range: -17% to 30%). The impacts on daily peak flows are moderated by changes in snow (Supplementary Section 6.4). Snow has historically been one of the primary flood-generating mechanisms in the lower Thames (typically via rapid melt of large accumulations coincident with heavy rainfall, as occurred to cause the major flooding of March 1947), but has been less common in recent years²³. However, the other primary flood-generating mechanism in the lower Thames is sustained heavy rainfall (typically over 4-7 days) on saturated ground²³. Thus differences in the anthropogenic influence on extreme 5-day and 30-day rainfall accumulations (Supplementary Fig. 14) further explain the more modest impacts on daily peak flows compared to 30-day peak flows. These differences between 30-day and 5-day rainfall accumulations are correlated with the SST gradients of the 11 Natural ensembles at 0.65 (p-value of 0.03). Thus the anthropogenic increase in rainfall that we simulate is less on timescales that dominate flooding in this catchment, consistent with the mechanism being an increase in the frequency of the zonal regime, and so, successions of strong but fast-moving storms.

Outputs from CLASSIC are combined with information about the location of properties at risk of flooding in the Thames catchment, for flood events of various magnitudes, in order to estimate the change in risk of numbers of

properties (Supplementary Information Section 7). These estimates are derived using a method previously applied in the production of official government flood zone maps in England²⁴ (incorporating subsequent improvements in data and modelling). The Ordnance Survey, the government agency responsible for mapping of Great Britain, supplied property location data. Changes in risk are reported here based on the daily peak flows, which represent the closest available approximation to the instantaneous peak flow rates that determine river water levels, even though the effects of changes in forcing are greater for flow volumes integrated over longer durations.

For events with around a 100-year return period, the best estimate is that about 1,000 more properties are placed at risk of flooding in a human-altered climate (Fig. 5f). Again, the results span a range of possible outcomes from around 4,000 fewer to 8,000 more properties at risk. The average flood insurance claim during the period DJF 2013/2014 (which predominantly reflects flooding in Southern England, especially around the Thames) is reported by industry sources⁴ to be approximately £24,000. Therefore the best estimate additional exposure to flood risk in an event similar to DJF 2013/2014 would be about £24 million in terms of potential losses (uncertainty range -£96 million to £192 million) suggesting a non-negligible contribution to risk when taking account of the ensemble uncertainty around the central estimate. Although there is only a small (ensemble average) increase in daily peak flows the results suggest that when winter flooding of the Thames does occur, it could be lasting longer which has implications both for damages and civil emergency management.

The only human influence considered here is the change in atmospheric composition. In both Actual and hypothetical Natural conditions, the flood risk would have been affected by anthropogenic interventions, in particular flood defences, although only a relatively small proportion of floodplain properties benefit from significant defences (Supplementary Information Section 7) and it is not known how that infrastructure might have evolved in the counterfactual world represented in the Natural ensembles.

290 4. Conclusions

291 This is the first end-to-end attribution study from anthropogenic changes in
292 atmospheric composition, through a meteorological extreme event and its
293 hydrological impacts to an estimate of the value of those impacts in terms of
294 flood damages. It illustrates how even relatively subtle changes in weather-
295 related risks could potentially have significant monetary impacts. In summary
296 we find that human influence:

- 297 • Increased the risk of low pressure Northwest of Britain and the number
298 of days with zonal flow over the North Atlantic
- 299 • Increased the risk of heavy precipitation in Southern England
- 300 • Increased the chance of extreme 30-day flows for the river Thames
- 301 • Had more modest effects on peak daily flows for the river Thames and
302 the risk of flooding to properties in its basin.

303 All these cases have large uncertainties due to sensitivity to the uncertain
304 geographical pattern of anthropogenic SST warming. We further estimate that
305 while thermodynamic effects cause most of the increase in precipitation,
306 around 1/3 is caused by changes in circulation.

307 Our results illustrate the importance of considering changing risks of extreme
308 weather in quantifying climate change impacts and highlights that a holistic
309 assessment of the risk requires the consideration of both the thermodynamic
310 and dynamic response of the climate system to human-induced changes in
311 the atmospheric composition^{25,26}.

312 Although the central estimate of increase in the number of properties at risk is
313 small, the ensemble uncertainty spans a range of changes in flood damages
314 that includes some chance of reductions, and also a substantial chance of
315 increased damages that would be significant relative to total flood claims
316 during DJF 2013/2014. A broader assessment could include the risks from

storm surge in the Thames estuary and from a wider range of extreme weather and flood events. It should be noted that this analysis does not take into account other factors that influence the risk of flooding to properties in southern England, such as continuing development on flood plains and levels of spending on flood defences that have been criticized as inadequate²⁷, and that some residual risk of flooding will need to be managed under investment strategies regarded as economically optimal^{28,29}. It should also be noted that the impacts on flows and damages for other catchments are likely to differ from those estimated for the Thames catchment draining to Kingston, because of differences in catchment characteristics and potential spatial differences in rainfall patterns.

This study is based on one particular atmospheric model where physical model uncertainty is represented only by the differing SST patterns representing the difference between current and pre-industrial obtained from 11 different climate models. It would clearly be desirable to replicate these results with a broader range of climate models to better understand the sensitivities to model formulations as well as biases and forcings, including model resolution and the pattern and magnitude of the anthropogenic SST signal used to simulate the 'climate that might have been' without human influence. Similarly, potential sensitivity of results to the choice of hydrological model should be assessed, although this is likely to be less important than choice of climate model³⁰. More studies of this nature are needed if loss and damage from anthropogenic climate change are to be quantified objectively³¹ and future assessments of the impacts of climate change are to progress from attributing them simply to changes in climate which are not themselves explained³², to attributing them specifically to human influence³³.

Correspondence should be addressed to Nathalie Schaller
(Nathalie.Schaller@physics.ox.ac.uk)


Acknowledgements

The authors thank the *climateprediction.net* participants whose generous donation of their spare computer processing power has enabled the large model ensembles to be created. Thanks to Tim Palmer for suggesting Fig. 2, to Sarah Kew for assistance with the kernel density estimates, and to Maliko Tanguy and Virginie Keller for producing the CEH-GEAR data for 2013/2014 ahead of schedule. We further thank JBA Risk Management Ltd. for permission to use data derived from their GB Comprehensive Flood Map, based on Astrium digital terrain data. Property locations were derived from AddressPoint data, used with kind permission of Ordnance Survey. NS, NRM, GJvO, RV, PY, AW, PAS and MRA were supported by the EUCLEIA project funded by the European Union's Seventh Framework Programme [FP7/2007-2013] under grant agreement no. 607085. NS received additional support from the Swiss National Science Foundation. NRM, FELO, SNS, WJI, AB, JM & DW also received support from the NERC HYDRA Changing Water Cycle project. ALK, SMC and CH were supported by the CEH/NERC National Capability fund. PAS, WJI and RGJ were also supported by the UK Joint Department for Energy and Climate Change (DECC), Department for Environment, Food and Rural Affairs (Defra) MOHC Climate Programme (GA01101).

Author contributions

NS, AK, RL, GJvO, RV, PY, PAS and MRA designed the study, NS, AK, RL, NRM, AB, JM, JS set up and performed model experiments, NS, AK, RL, NRM, GJvO, FELO, SNS, RV, PY, KH, CH, TL and JS provided analyses and all authors wrote the paper.

372 **Figures**



373
374 **Figure 1:** Precipitation³⁴ (colours, in mm day⁻¹) and mean sea level pressure¹¹ (contours, in hPa) as
375 observed for January 2014 absolute values in **a** and as anomalies from the observed 1981-2010
376 climatology in **b**, and in the wettest 1% of the Actual Conditions ensemble as absolute values in **c** and
377 as anomalies from the model 1986-2011 climatology in **d**.




Figure 2: Anomalies of zonal wind at 200 hPa for January 2014 **a** in ERA-interim¹¹, relative to the 1986-2011 ERA-interim climatology, and **b** in the ensemble mean of the Actual Conditions simulations, relative to the model 1986-2011 climatology. **c** and **d**, as **b**, but for the ensemble means of the Natural simulations with the HadGEM2-ES and CCSM4 models respectively.




Figure 3: a Time series of monthly mean rain/precipitation for January 1768-2014 at the Radcliffe Observatory, Oxford. Above/below overall average values are plotted in blue/brown. January 2014 is highlighted in red. The black line is the 20-year Lowess-smoothed monthly mean precipitation. The measurements are rain only until around 1867 (dotted thin vertical line), but include snow since then. **b** Comparison of all the 5-day mean precipitation for all winter months from 1827/28-2013/14. The 5 wettest years are highlighted in dark grey. Winter 2013/14 is plotted in red.




Figure 4: **a** Relationship between modelled January monthly average Southern England precipitation and mean sea level pressure at 20°W, 60°N. The 50th, 75th, 95th and 99th percentiles of the distribution of the Actual Conditions and all Natural simulations are estimated using a Gaussian bivariate kernel density estimator. Grey dots represent January averages for each individual Actual Conditions simulations and the black dots show values from observations ("8 stations" refers to the average of 8 stations in Southern England for the precipitation index and the NCEP reanalysis³⁵ for the pressure index, "E-OBS" refers to the same definition as the modelled precipitation index using the gridded E-OBS dataset³⁶ also with NCEP pressure index). The Actual Conditions and Natural joint distributions are significantly different at the 0.05 level based on a two-sided bivariate version of the Kolmogorov-Smirnov test³⁷. **b** As a but showing the relationship between modelled January Southern England precipitation binned in 7 categories and the January ZO index binned in three categories of number of days per month. For all three categories, the distributions of Actual Conditions and Natural are statistically different at the 0.05 level, according to both a two-sided Kolmogorov-Smirnov and a two-sided Cramer-von Mises test. The number of ensemble members in each of the three categories is given on the bottom-right corner of each sub-panel. **c** Return periods for pressure for the Actual Conditions and pooled Natural simulations along with pooled Natural weighted to make its pressure values match the Actual Conditions simulation. **d** as **c** but for precipitation, using the same weights as in **c**.




Figure 5: Return periods for **a** modelled January pressure index (each dot represents an ensemble member) with 5-95% confidence intervals for 1-in-50-year events and 1-in-100-year events in Actual Conditions estimated by resampling the distribution 100 times represented as horizontal lines. Red represents Actual Conditions simulations, grey a similar ensemble but for 1986-2011 (the model climatology), dark blue the pooled Natural simulations, and light blue individual Natural (sub-) ensembles, with solid circles for the 6 of the 11 Natural ensembles with around 15,000 simulations, and empty circles for the other 5 with around 7,000 simulations. Only four 5-95% confidence intervals for 1-in-50-year events and 1-in-100-year events (red: Actual Conditions, grey: Climatology, light blue: Natural ensembles with around 15,000 ensemble members and dashed light blue: Natural ensembles with around 7,000 simulations) are shown because the confidence intervals represent only the sampling uncertainty, not the uncertainty in the estimation of the model simulations. **b** as **a** but modelled

frequency of the ZO regime. No confidence intervals are shown due to the categorical nature of return values. **c** observed monthly precipitation averaged for 8 stations across Southern England for the months of November to February individually for the years 1912-2013 fitted to a Generalised Pareto Distribution with location and scale parameters linearly dependent on the low-pass filtered global mean temperature. Red lines indicate the fit and 90% confidence interval for the current temperature (2013/2014), blue for a temperature representative of pre-industrial conditions (1912/1913). The red (blue) crosses show the observations shifted up (down) to these years using the fitted trend. The horizontal grey line represents the observed value for January 2014. The fit has been performed for monthly means of four calendar months to increase the sample size, the return period is given per month for comparison with the other results. **d** as **a** for modelled January mean precipitation in Southern England, **e** as **a** for modelled 30-day peak flows for the Thames at Kingston, and **f** difference between the Natural and the Actual Conditions simulations in number of properties individually at risk of flooding with annual probability $1/T$, where T is the return period.

References

- 1 Huntingford, C. *et al.* Potential influences on the United Kingdom's floods of winter 2013/14. *Nature Climate Change* **4**, 769-777, doi:10.1038/nclimate2314 (2014).
- 2 Matthews, T., Murphy, C., Wilby, R. L. & Harrigan, S. Stormiest winter on record for Ireland and UK. *Nature Climate Change* **4**, 738-740 (2014).
- 3 Pall, P. *et al.* Anthropogenic greenhouse gas contribution to flood risk in England and Wales in autumn 2000. *Nature* **470**, 382-385 (2011).
- 4 Association of British Insurers, <https://www.abi.org.uk/Insurance-and-savings/Topics-and-issues/Flooding/2014-floods-in-numbers> (Accessed September 2015)
- 5 Association of British Insurers, <https://www.abi.org.uk/News/News-releases/2010/11/massive-rise-in-britains-flood-damage-bill-highlights-the-need-for-more-help-for-flood-vulnerable-communities-says-the-abi.aspx> (Accessed September 2015)
- 6 <http://www.bbc.co.uk/news/uk-politics-25656426>
- 7 Stott, P. A., Stone, D. A. & Allen, M. R. Human contribution to the European heatwave of 2003. *Nature* **432**, 610-614, doi:10.1038/nature03089 (2004).
- 8 Kay, A. L., Crooks, S. M., Pall, P. & Stone, D. A. Attribution of Autumn/Winter 2000 flood risk in England to anthropogenic climate change: A catchment-based study. *Journal of Hydrology* **406**, 97-112, doi:10.1016/j.jhydrol.2011.06.006 (2011).
- 9 Massey, N. *et al.* weather@home - development and validation of a very large ensemble modelling system for probabilistic event attribution. *Quarterly Journal Of The Royal Meteorological Society*, doi:10.1002/qj.2455 (2014).
- 10 Taylor, K. E., Stouffer, R. J. & Meehl, G. A. An Overview of CMIP5 and the Experiment Design. *Bull. Amer. Meteorol. Soc.* **93**, 485-498 (2012).
- 11 Dee, D. P. *et al.* The ERA-Interim reanalysis: configuration and performance of the data assimilation system. *Quarterly Journal of the Royal Meteorological Society* **137**, 553-597 (2011).
- 12 van Haren, R., van Oldenborgh, G. J., Lenderink, G. & Hazeleger, W. Evaluation of modeled changes in extreme precipitation in Europe and the Rhine basin. *Environ. Res. Lett.* **8**, 7, doi:10.1088/1748-9326/8/1/014053 (2013).
- 13 van Haren, R., van Oldenborgh, G. J., Lenderink, G., Collins, M. & Hazeleger, W. SST and circulation trend biases cause an underestimation of European precipitation trends. *Climate Dynamics* **40**, 1-20, doi:10.1007/s00382-012-1401-5 (2013).

470 14 Vautard, R. Multiple weather regimes over the North Atlantic - Analysis of precursors
471 and successors. *Mon. Weather Rev.* **118**, 2056-2081, doi:10.1175/1520-
472 0493(1990)118<2056:mwrotn>2.0.co;2 (1990).

473 15 Michelangeli, P. A., Vautard, R. & Legras, B. Weather regimes - Recurrence and
474 quasi stationarity. *J. Atmos. Sci.* **52**, 1237-1256, doi:10.1175/1520-
475 0469(1995)052<1237:wrraqs>2.0.co;2 (1995).

476 16 Yiou, P., Goubanova, K., Li, Z. X. & Nogaj, M. Weather regime dependence of
477 extreme value statistics for summer temperature and precipitation. *Nonlinear Process*
478 *Geophys.* **15**, 365-378 (2008).

479 17 Barnes, E. A. & Polvani, L. Response of the Midlatitude Jets, and of Their Variability,
480 to Increased Greenhouse Gases in the CMIP5 Models. *Journal of Climate* **26**, 7117-
481 7135 (2013).

482 18 Zappa, G., Hoskins, B. J. & Shepherd, T. G. Improving Climate Change Detection
483 through Optimal Seasonal Averaging: The Case of the North Atlantic Jet and
484 European Precipitation. *Journal of Climate* **28** (16) (2015).

485 19 Cattiaux, J. & Cassou, C. Opposite CMIP3/CMIP5 trends in the wintertime Northern
486 Annular Mode explained by combined local sea ice and remote tropical influences.
487 *Geophysical Research Letters* **40** (2013).

488 20 Rodwell, M. J., Rowell, D. P. & Folland, C. K. Oceanic forcing of the wintertime North
489 Atlantic Oscillation and European climate. *Nature* **398**, 320-323, doi:10.1038/18648
490 (1999).

491 21 Haarsma, R. J., Selten, F. & van Oldenborgh, G. J. Anthropogenic changes of the
492 thermal and zonal flow structure over Western Europe and Eastern North Atlantic in
493 CMIP3 and CMIP5 models. *Climate Dynamics* **41**, 2577-2588, doi:10.1007/s00382-
494 013-1734-8 (2013).

495 22 Crooks, S. M. & Naden, P. S. CLASSIC: a semi-distributed rainfall-runoff modelling
496 system. *Hydrol. Earth Syst. Sci.* **11**, 516-531 (2007).

497 23 Marsh, T. & Harvey, C.L. 2012. The Thames flood series: a lack of trend in flood
498 magnitude and a decline in maximum levels. *Hydrology Research*, **43**(3), 203-214

499 24 Bradbrook, K., Waller, S., & Morris, D. National floodplain mapping: Datasets and
500 methods - 160,000 km in 12 months. *Natural Hazards*, 36(1-2), 103-123 (2005).

501 25 Trenberth, K., Fasullo, J. T. & Shepherd, T. G. Attribution of climate extreme events.
502 *Nature Climate Change* **5**, 725-730, doi:10.1038/nclimate2657 (2015).

503 26 Hansen, J., Sato, M. & Ruedy, R. Perception of climate change. *PNAS* **109** (37),
504 E2415-2423, doi:10.1073/pnas.1205276109 (2012).

505 27 Crichton D. Flood Risk and Insurance in England and Wales: Are there lessons to be
506 learned from Scotland? (Benfield Hazard Research Centre, UCL, London, 2005).

507 28 Committee on Climate Change. Managing climate risks to well-being and the
508 economy. (Adaptation Sub-Committee Progress Report, Committee on Climate
509 Change, London, 2014). [http://www.theccc.org.uk/wp-](http://www.theccc.org.uk/wp-content/uploads/2014/07/Final_ASC-2014_web-version-4.pdf)
510 [content/uploads/2014/07/Final_ASC-2014_web-version-4.pdf](http://www.theccc.org.uk/wp-content/uploads/2014/07/Final_ASC-2014_web-version-4.pdf) (Accessed September
511 2015)

512 29 Environment Agency. Flood and coastal erosion risk management. (Long-term
513 investment scenarios, Report No. LIT10045, Environment Agency, Bristol UK, 2014).
514 [https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/381939](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/381939/FCRM_Long_term_investment_scenarios.pdf)
515 [/FCRM_Long_term_investment_scenarios.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/381939/FCRM_Long_term_investment_scenarios.pdf) (Accessed September 2015)

516 30 Kay, A. L., Davies, H. N., Bell, V. A. & Jones, R. G. Comparison of uncertainty
517 sources for climate change impacts: flood frequency in England. *Climatic Change* **92**,
518 41-63, doi:10.1007/s10584-008-9471-4 (2009).

519 31 James, R. *et al.* Characterizing loss and damage from climate change. *Nature Clim.*
520 *Change* **4**, 938-939, doi:10.1038/nclimate2411 (2014).

521 32 Cramer, W. *et al.* in *Climate Change 2014: Impacts, Adaptation, and Vulnerability.*
522 *Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth*
523 *Assessment Report of the Intergovernmental Panel on Climate Change* (eds C. B.
524 Field *et al.*) (Cambridge University Press, 2014).

525 33 Bindoff, N. L. *et al.* in *Climate Change 2013: The Physical Science Basis.*
526 *Contribution of Working Group I to the Fifth Assessment Report of the*
527 *Intergovernmental Panel on Climate Change* (eds T. F. Stocker *et al.*) (Cambridge
528 University Press, 2013).

529 34 Perry, M. & Hollis, D. The generation of monthly gridded datasets for a range of
530 climatic variables over the UK. *Int. J. Climatol.* **25**, 1041-1054, doi:10.1002/joc.1161
531 (2005).

532 35 Kistler, R. *et al.* The NCEP-NCAR 50-year reanalysis: Monthly means CD-ROM and
533 documentation. *Bull. Amer. Meteorol. Soc.* **82**, 247-267, doi:10.1175/1520-
534 0477(2001)082<0247:tnnyrm>2.3.co;2 (2001).

535 36 Haylock, M. R. *et al.* A European daily high-resolution gridded data set of surface
536 temperature and precipitation for 1950-2006. *J. Geophys. Res.-Atmos.* **113**, D20119-
537 D20119 (2008).

538 37 Peacock, J. A. Two-dimensional goodness-of-fit testing in astronomy. *Mon. Not. Roy.*
539 *Astron. Soc.* **202**, 615-627 (1983).