

**Avian influenza A (H5N1) virus infection in humans, 1997 – 2015: a
descriptive global epidemiological analysis**

Shengjie Lai*, Ying Qin*, Benjamin J. Cowling*, Xiang Ren, Nicola A. Wardrop,
Marius Gilbert, Tim K. Tsang, Peng Wu, Luzhao Feng, Hui Jiang, Zhibin Peng,
Jiandong Zheng, Qiaohong Liao, Sa Li, Peter W. Horby, Jeremy J. Farrar, George F.
Gao, Andrew J. Tatem, Hongjie Yu

* Authors contributed equally

**Division of Infectious Disease, Key Laboratory of Surveillance and Early-
warning on Infectious Disease, Chinese Center for Disease Control and
Prevention, Beijing, China** (S Lai MD, Y Qin PhD, X Ren MD, L Feng PhD, H Jiang
MD, Z Peng MD, J Zheng PhD, Q Liao MD, S Li MD, Prof H Yu PhD); **Department of
Geography and Environment, University of Southampton, Southampton, UK**
(S Lai MD, N A Wardrop PhD, Prof A J Tatem PhD); **WHO Collaborating Centre
for Infectious Disease Epidemiology and Control, School of Public Health, Li
Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong
Special Administrative Region, China** (B J Cowling PhD, T K Tsang PhD, P Wu
PhD); **Biological Control and Spatial Ecology, Universite' Libre de Bruxelles,
Brussels, Belgium** (M Gilbert PhD); **Fonds National de la Recherche
Scientifique, Brussels, Belgium** (M Gilbert PhD); **Oxford University Clinical**

Research Unit, Wellcome Trust Major Overseas Programme, Ho Chi Minh City, Vietnam (Prof P W Horby MD, Prof J J Farrar MD); **Centre for Tropical Medicine, Nuffield Department of Clinical Medicine, Oxford University, UK** (Prof P W Horby, Prof J J Farrar); **Singapore Infectious Disease Initiative, Singapore** (Prof P W Horby, Prof J J Farrar); **ISARIC, Centre for Tropical Medicine, University of Oxford, Churchill Hospital, United Kingdom** (Prof J J Farrar); **CAS Key Laboratory of Pathogenic Microbiology and Immunology, Institute of Microbiology, Chinese Academy of Sciences, Beijing, China** (Prof G F Gao PhD); **Office of Director-General, Chinese Center for Disease Control and Prevention, Beijing, China** (Prof G F Gao); **Flowminder Foundation, Stockholm, Sweden** (Prof A J Tatem); **Spatial Ecology and Epidemiology Group, Tinbergen Building, Department of Zoology, University of Oxford, UK** (Prof A J Tatem).

Correspondence to:

Prof Hongjie Yu, Division of Infectious Disease, Key Laboratory of Surveillance and Early-warning on Infectious Disease, Chinese Center for Disease Control and Prevention, No. 155 Changbai Road, Changping district, Beijing, China
yuhj@chinacdc.cn

Word count (abstract): 325

Word count (main text): 3,497

Running head: Global human cases of H5N1 in 1997-2015

SUMMARY

Background: From November 2014 through April 2015, there were a total of 165 confirmed human cases of avian influenza A(H5N1) virus infection, including 48 deaths that were reported to the World Health Organization (WHO). This is the highest number of human cases ever reported globally over a similar period since the first human case of H5N1 was reported in Hong Kong in 1997. This study describes the epidemiology of human cases of H5N1 in 1997-2015.

Methods: We describe the epidemiology of global human H5N1 cases spanning 18 years between 1 May 1997 and 30 April 2015, focusing on the characteristics of cases, seasonality, geographical distribution, and outcome.

Findings: A total of 907 human H5N1 cases were reported globally from May 1997 to April 2015, with no cases reported between 1998 and 2002, and the largest annual numbers recorded in 2015 (152 cases). The number of affected countries rose between 2003 and 2008, with expansion from East and Southeast Asia, then to West Asia and Africa. The overall case fatality risk (CFR) was 53.5%, and this varied across geographical regions ($p < 0.001$) with the highest (69.4%) in East and Southeast Asia and the lowest in North Africa (32.1%). The age of cases differed significantly by outcome ($p < 0.001$), with a median age of 22 years for fatal cases and 10 years for non-fatal cases. The majority (95.8%, 748/781) of cases reported exposure to poultry. Although the incidence of human cases in

Egypt has increased dramatically from November 2014 through April 2015, but compared to the cases before November 2014 there were no significant differences in the CFR, history of exposure to poultry, history of human case contact, and time from onset to hospitalization.

Interpretation: The geographic distribution of countries affected has expanded, especially between 2003 and 2008, with variations in outcome, demography, seasonality and virus clade across regions. The incidence of human infection has increased dramatically in Egypt since November 2014, but the CFR has not changed significantly.

Funding: The Ministry of Science and Technology of China, the Ministry of Health of China, the US National Institutes of Health, the Harvard Center for Communicable Disease Dynamics from the National Institute of General Medical Sciences, the University Grants Committee of the Hong Kong Special Administrative Region of China, the Wellcome Trust, the Flowminder Foundation, the Bill & Melinda Gates Foundation, and the UK Medical Research Council.

Key words: Avian influenza; H5N1; Epidemiology; Seasonality; Egypt

INTRODUCTION

Highly pathogenic avian influenza (HPAI) A(H5N1) viruses emerged in poultry in Southeast Asia in 1996, and outbreaks have since been reported in domestic poultry, wild birds and humans in over 60 countries.^{1,2} The spread of HPAI H5N1 in poultry populations increases the risk of human infections.³⁻⁵ The first reported case of human illness from infection with H5N1 virus occurred in May 1997 in Hong Kong Special Administrative Region (SAR) of China.⁶⁻⁸ After an apparent five-year absence, two cases were reported in February 2003 in Hong Kong SAR.⁹ Following the pattern of spread and persistence of the virus in poultry, human cases of H5N1 virus infection with high mortality were subsequently detected in China, south-east Asia,^{10,11} West Asia and most recently Africa, with cases detected in Egypt every year from 2006 to 2015. Compared to previous years, the incidence of human H5N1 cases has remained at a low level globally between October 2012 and October 2014,¹² while attention has been focused on the more recent emergence of variant swine influenza A(H3N2) in North America,¹³ novel avian influenza A(H7N9) in China,¹⁴⁻¹⁶ other avian influenza A(H5) subtypes in Asia, Europe and North America,¹⁷ and other emerging infections.

However, between 1 November 2014 and 30 April 2015, a total of 165 cases, including 48 deaths were reported to the World Health Organization (WHO).¹⁸

This is by far the highest number of human cases ever reported globally over a

similar period.¹⁹ Moreover, the number of human H5N1 cases reported in Egypt with onset in February 2015 is the highest number reported by any country in a single month.²⁰ The emergence of a novel cluster of H5N1 virus clade 2.2.1.2 has been found in poultry in Egypt since mid-2014 and has quickly become predominant.²¹ It is not yet known if this emerging phylotype has increased zoonotic potential and, thus, can be transmitted more efficiently to humans.²⁰⁻²²

In order to improve understanding of the epidemiology of HPAI H5N1, we describe the magnitude and distribution of all human H5N1 cases globally with illness onset between 1 May 1997 and 30 April 2015. We focus on the characteristics of cases, seasonal and geographical patterns, and examine in more detail the epidemiology of human H5N1 cases in Egypt.

METHODS

Sources of Data

Human H5N1 case data were identified and compiled according to the probable and confirmed case definitions described in the next section. Data on all human H5N1 cases in mainland China were downloaded from the online National Notifiable Infectious Disease Reporting Information System at the Chinese Center for Disease Control and Prevention (China CDC).^{14,23} Data on human H5N1 cases in Vietnam and Azerbaijan as of 30 April 2014 were provided by the Vietnam National Institute of Hygiene and Epidemiology and the Azerbaijan

Ministry of Health, respectively. Data on human H5N1 cases in all other affected countries or regions were obtained from publicly available sources (Appendix Table 1), which were also described in a previously published report,²⁴ and include the WHO's Disease Outbreak News of the Global Alert and Response (GAR);²⁵ the WHO's Weekly Epidemiological Record;²⁶ the FluTrackers website (<https://flutrackers.com/>);²⁷ and published literature in PubMed on May 1, 2015 using the queries "H5N1[Title] AND human[Title]" including all articles published in English between May 1, 1997 and April 30, 2015, then excluding articles without individual human case data.

We used WHO GAR updates and the WHO statistics on cumulative number of confirmed human H5N1 cases from November 2003 to April 2015 to establish a line list of human H5N1 cases.²⁸ All cases from sources other than WHO updates were matched with the initial line list. The latest cases, which were not yet officially announced by WHO, were identified through ProMed-mail posts and FluTrackers, and confirmed by the announcements of Ministries of Health in individual countries. When critical information was missing, additional information was sought from published literature, ProMed posts and English language news releases from the regional office of WHO and the relevant Ministry of Health.²⁹⁻³²

Case definition

The WHO case definition was used.³³ A ***confirmed case*** was defined as an human case of influenza A(H5N1) virus infection reported by WHO and with laboratory confirmation, i.e. a patient with defined clinical signs, epidemiological linkage and laboratory confirmation by an influenza laboratory accepted by WHO, as specified in the WHO case definition. Other reported cases were considered as ***probable cases*** if they had exposure to other confirmed human cases, or to sick or dead poultry, or the H5N1 infection was confirmed by the country or local institutions but not meeting WHO criteria.

Data variables and extraction

Individual data on cases included age, sex, country, type of diagnosis, year, month and day of onset, date of hospitalization, final outcome (fatal or non-fatal), date of outcome, and potential risk factors (Appendix Table 2). Information on exposure potentially related to the acquisition of H5N1 infections was collected (Box 1). Where missing variables were identified in the line list of human H5N1 cases, we extracted data from reports by the WHO^{12,20,25} and FluTrackers.²⁷ The websites of national Ministries of Health, ProMed-mail posts, and other online sources were also searched to identify missing information. Where contradictory information was found for a given variable, WHO and Ministry of Health data were given priority over journal articles, and journal articles were given priority over other sources of information (Appendix Table 1). All the data used in this

study were anonymized.

Data on the clade or subclade of H5N1 virus isolated from human cases were also collated from the regular WHO reports.³⁴ In total, 17 reports issued between August 2006 and February 2015 were reviewed, which provided information on H5N1 virus clades circulating and characterized from 1997 to February 2015.³⁴ However, not all individual cases were reported with laboratory results of the clade or subclade, and thus, where this information was not available we presumed infection was with the clade or subclade of H5N1 virus in the same period and area.^{11,21,35-37}

Ethical approval

The National Health and Family Planning Commission of China, the Ministry of Health of Vietnam, and the Ministry of Health of Azerbaijan determined that the collection of data from human cases of avian influenza A(H5N1) virus infection was part of the public health investigation of an outbreak and was exempt from institutional review board assessment. All data were supplied and analyzed in an anonymous format, without access to personal identifying information.

Data analysis

We included all probable and confirmed cases with illness onset by 30 April 2015 in the analysis. We plotted epidemic curves and summarized demographic

characteristics by final outcome and geographical region. A heat map illustrating the proportion of cases reported each month, standardized by the total number of cases in each country over the 18-year period and grouped by region, was created to analyze the occurrence of H5N1 cases. To describe seasonal patterns, we created a heat map of the mean value of the proportion of cases in each week by country from May 1997 through April 2015. The distribution of time intervals from onset of symptoms to hospital admission was presented and smoothed using kernel density estimation.

The *R* statistical software (version 3.1.2, R Foundation for Statistical Computing, Vienna, Austria) with the package *Epicalc* (version 2.15.1.0) was used to produce the graphs, heat maps and perform statistical analyses with a significance level of $\alpha = 0.05$. *ArcGIS* 10.2 software (ESRI, Redlands, CA, USA) software was employed to plot the geographical distribution of cases. Mann–Whitney U tests and Chi-squared tests were used to test for differences between continuous and dichotomous variables respectively. Crude odds ratios (OR) with exact 95% confidence intervals (CI) were estimated for potential risk factors.

Role of the funding source

The sponsor of the study had no role in the study design, data collection, data analysis, data interpretation, writing of the report, or the decision to publish. The corresponding author had full access to all the data in the study and had final

responsibility for the decision to submit for publication. The views expressed are those of the authors and do not necessarily represent the policy of the China CDC.

RESULTS

A total of 907 human H5N1 cases were reported globally during the 18-year period from May 1, 1997 through April 30, 2015, of which 94.6% were confirmed cases and 5.4% were probable cases (Table 1 and Figure 1). Annual case numbers displayed striking variations, with the highest numbers recorded in 2015 (Figure 2, Appendix Figures 1-2). The total number of cases (213) in 2014-2015 was greater than that (181 cases) of the four years from 2010 –2013 (Appendix Figure 3), with the highest monthly number occurring in February 2015 (55 cases in Egypt and one in China).

The overall male-to-female ratio was almost even (1:1.2) from 1997 to 2014, although this pattern was not uniform across regions (Table 1). The median age of cases was 19 years, with an inter quartile range (IQR) from 5 to 32 years, and 41.2% (363/881) were children under 15 years of age and 80.3% (707/881) were in people under 35y. The median age of non-fatal cases was younger in North Africa and older in East and Southeast Asia (median and IQR: 6 and 3-31 years vs. 18.5 and 6-30 years; $p=0.001$), but the median age of fatal cases was older in North Africa than in East and Southeast Asia (median and IQR: 30 and

20-36 years vs. 19 and 9-30 years; $p < 0.0001$) (Figure 3).

In total, 16 countries reported human cases between 1997 and 2015. The number of affected countries has risen between 2003 and 2008, with range expansion from East Asia to Southeast Asia, then West Asia, North Africa and other regions, with an apparent ongoing transmission and cases reported almost every year in China, Vietnam, Cambodia, Indonesia and Egypt (Figure 4A). The incidence in Asia remained at low levels in 2013-2015, while the number of cases in Egypt has increased in 2014-2015. During 1997-2015, 67.2% (594/884) of cases were reported from December to March, with a peak in January (20.9%). However, countries in East Asia and West Asia had fewer cases occurring in the warm/hot season from April to September (8.1% vs. 26.2%; $p < 0.0001$), and showed earlier peaks (December vs. January) and shorter epidemic periods than countries in Southeast Asia and North Africa at lower latitudes (Figure 4B, Appendix Figure 4).

The overall case fatality risk (CFR) was 53.5% (483/903), with a decrease from 70.7% (275/420) in 2003-2008 to 43.4% (202/465) in 2009-2015, and varied across geographical regions ($p < 0.0001$), with a CFR (69.4%, 349/503) in East and Southeast Asia was more than twice that in North Africa (32.1%, 116/361) (Table 1). Compared with Egypt, there was a higher risk of death in cases in Indonesia, Vietnam, Cambodia, mainland China and Thailand ($p < 0.0001$). The

age distribution of cases also differed significantly by outcome ($p < 0.0001$), with a median age of 22 years (IQR: 11.5-32 years) for fatal cases and 10 years (IQR: 3-30 years) for cases who recovered (Figure 2C-D). The majority (95.8%, 748/781) of cases reported exposure to poultry including: 85.7% (439/512) exposed to sick or dead poultry; 61.4% (188/306) exposed to backyard poultry; 26.4% (82/311) exposed to LBMs; 4.7% (15/321) occupational exposure to live poultry. In addition, 6.2% (49/792) reported contact with a human H5N1 case before the onset of illness (Table 1, Appendix Table 3).

Time from onset of illness to hospitalization was available for 79.7% (723/907) cases with a median of 4 days (IQR: 2-6 days). Generally, the survived cases had an earlier hospitalization than fatal cases (median and IQR: 3 and 1-6 days vs. 5 and 3-7; $p < 0.0001$) (Figure 5). The cases with ≥ 3 days from onset of illness to hospitalization were more likely to be fatal than those admitted within 3 days of onset (OR 3.6; 95%CI: 2.5-5.1). Additionally, the cases in North Africa had a shorter time from onset to hospitalization than cases in East and Southeast Asia (median and IQR: 3 and 1-6 days vs. 5 and 3-7; $p < 0.0001$), but the time from onset to outcome was not significantly different between cases in North Africa and cases in East and Southeast Asia, with the same median of 10 days ($p = 0.067$).

The A(H5N1) viruses in human cases have been characterized belong to clade or

subclade 0, 1, 2.1, 2.1, 2.3, and 7 (Table 1-2, Appendix Table 3). Clade 1 was first reported in Hong Kong SAR in 2003, and then reported in Southeast Asia each year from 2003 to 2014, but subclade 2.1 was only reported in Indonesia since 2005, and subclade 2.2 has circulated in Egypt since 2006 with sporadically reported in Africa and West Asia. In addition, subclade 2.3 has been reported in East and Southeast Asia since 2005. Compared to clade 0, the cases with clade 1, subclade 2.1 and 2.3 were more likely to result in death with an OR of 2.8 (95%CI: 0.93, 9.6), 11.0 (95%CI: 3.5, 37.8) and 3.2 (95%CI: 1.0, 11.4) respectively. However, the risk of death between cases with clade 0 and subclade 2.1 was not significantly different (OR: 1.0; 95%CI: 0.3, 3.3).

Human cases of H5N1 in Egypt

During March 2006 – April 2015, a total of 363 human cases with influenza A(H5N1) virus infection were reported with 116 deaths (32%) (Figure 6), of which more than half (51%) of cases were reported during the 6 months of November 2014 – April 2015 (Appendix Table 4). The male-to-female ratio was not significantly different between cases before November 2014 and cases in November 2014 – April 2015, but the latter had an older median age (median and IQR: 26; 4-38 years) than the former (median and IQR: 16; 3.25-30 years) ($p < 0.0001$), which was also different for both non-fatal ($p = 0.04$) and fatal cases ($p < 0.0001$) (Figure 3E-F). However, the CFR was not significantly different at 36% (64/178) before November 2014 and 28.4% (52/183) during November 2014 –

April 2015. For fatal cases, the median time and IQR (5; 3-6 days) for onset to hospital admission was the same between March 2006 – October 2014 and November 2014 – April 2015, but the time was different for non-fatal cases ($p < 0.0001$) with a median of one day (IQR: 1-3 days) before November 2014 and four days (IQR: 2-6 days) during November 2014 – April 2015. Most cases reported a history of poultry exposure, 96.1% before November 2014 and 90.8% in November 2014 – April 2015.

DISCUSSION

In this study, a global, longitudinal dataset spanning 18 years was collated to investigate changes in the epidemiological characteristics of human H5N1 cases, and also focused on Egypt, given its unique situation of increasing incidence since November 2014.

We found that the geographic extent of human H5N1 cases has expanded from East Asia to Southeast Asia, then to West Asia and North Africa during 2003-2009, which may be related to the global dispersal of the virus via bird migration.³⁸⁻⁴⁰ The bird migration network was shown to better reflect the observed viral gene sequence data than other networks and contributes to seasonal H5N1 epidemics in local regions.^{3,5} In addition, previous evidence demonstrated Siberia as a major hub for the dispersal of the virus via bird populations, and Southeast Asia and Africa as areas of local virus persistence and

the major sources of genetically and antigenically novel strains.^{3,5,41} Therefore, the increasing virus dispersal and outbreaks among birds may also increase the risk of human exposure.^{2,42} However, some of the apparent geographical dispersal in cases may also be attributed to enhanced clinical and laboratory surveillance capacity in the past 15-20 years.

We also found that human H5N1 cases occurred with seasonality related to the cooler season from December to March and across diverse climate zones in the Northern Hemisphere, which may correlate with the migration patterns of wild birds and the activity of virus in winter or cooler seasons (Figure 4).^{2,5,43} A recent study found that the timing of H5N1 outbreaks and viral migrations were closely associated with bird migration networks in Asia.³ In addition, the lower temperatures in Asia and North Africa across diverse climates was associated with increasing human H5N1 virus infection in winter, which is consistent with increased poultry outbreaks and H5N1 virus transmission during cold and dry conditions, and also overlapped with human seasonal influenza epidemics.^{2,43-45}

Although most human populations are thought to have little or no immunity to influenza A(H5N1) viruses, we found that most cases were children and younger adults, and these age groups were also more likely to recover, whereas the fatality risk was higher in adults, which might be related to the immunological reaction of virus in different age groups.²² Consistent with previous reports,^{46,47}

cases hospitalized earlier (<3 days after onset) were more likely to survive than other cases, which might be related to the early administration of antiviral treatment, or because of selection bias where the cases admitted later after onset were more likely to be severe.¹¹ Compared with other regions, the lower fatality risk in Egypt might be related to a less virulent virus clade, less severe clinical disease, and earlier identification, hospitalization and treatment for H5N1 cases.^{48,49} Additionally, we found that almost all human cases of H5N1 infection had a recent history of close contact with infected live or dead birds, other human cases, or H5N1-contaminated environments, which reaffirmed reports that human H5N1 virus infection is typically preceded by exposure to sick or dead poultry in backyards, LBMs or farms.⁵⁰⁻⁵⁵

We found that an increased number of animal-to-human infections has been reported by Egypt during November 2014 – April 2015 with the number of cases more than the total of the last 8 years from 2006-2014. The increase in the number of human cases in Egypt since November 2014 can be attributed to a mixture of factors, including increased circulation of H5N1 viruses in poultry, lower public health awareness of risks in middle and upper Egypt and seasonal factors, such as closer proximity to poultry because of cold weather and possible longer survival of the viruses in the environment.⁵⁶ Nevertheless, H5N1 viruses do not currently appear to transmit easily from person to person, and the risk of community level spread of these viruses remains low.^{12,20} A new candidate

vaccine virus has been proposed to protect against the current circulating H5N1 virus clade 2.2.1.2 viruses, covering all the recent H5N1 virus isolates from Egypt.²²

H5N1 virus has evolved from the 1996 progenitor strain and comprises at least 10 clades, through a complexity of genetic changes, which have infected domestic poultry and wild birds in many countries.^{38,39,57,58} In this study, we found that 4 clades (0, 1, 2, and 7) and 3 subclades (2.1, 2.2, and 2.3) of H5N1 virus strains have infected human, all of which have been reported in human cases before 2006.^{22,59} Based on available information, the clades of viruses isolated from humans were the same as the clades circulating in local poultry.⁴⁶ During the period from late 2003 to mid-2005, most H5N1 virus infections in humans were caused by clade 1 strains in Southeast Asia (i.e., Vietnam, Thailand, and Cambodia).⁵⁹

Although the highly pathogenic H5N1 virus strains can be expected to continue evolving over time, preliminary laboratory investigation has not detected major genetic changes in the viruses isolated from the patients or animals in 2014-2015 compared to previously circulating isolates in same regions,^{22,60} and the genetic diversity of the H5N1 virus decreased significantly between 1996 and 2011 in China, presumably under strong selective pressure associated with vaccination in poultry.³² However, other influenza A(H5) subtypes, such as influenza A(H5N2), A(H5N3), A(H5N6) and A(H5N8), have recently been

detected in birds in Europe, North America, and Asia, and so far no human cases of infection have been reported, with the exception of three human infections with influenza A(H5N6) virus detected in China in 2014-15.^{20,56} However, the co-circulation of influenza A viruses in human and animal reservoirs can provide opportunities for these viruses to re-assort and acquire the genetic characteristics that facilitate sustained human-to-human transmission, a necessary trait of pandemic viruses.^{2,61}

There are some limitations to this study. First, the data used were collated from different sources. The data quality may be influenced by key steps in public health surveillance or reports including the case definitions, reporting methods, availability of health care and laboratory diagnostics, under reporting, and completeness and accuracy of data reported or announced by different countries or organizations. Second, detailed data on case characteristics and clinical management were unavailable to assess the association between clinical manifestation, treatment and outcome. Third, the findings might be influenced by missing data on exposure, outcome and hospitalization, and the misclassification of cases with presumed clade or subclade.

In conclusion, we found that the geographic distribution of countries affected has expanded, especially between 2003 and 2008, with variations in outcome, demography, seasonality and the clade or subclade of viruses across the region.

The incidence of human infections increased dramatically in Egypt from November 2014 to April 2015, but remained at a low level in other regions, and the CFR in Egypt has not significantly changed. This study has identified high-risk areas, population groups and seasons, providing evidence for the planning of human HPAI H5N1 prevention and control.

Contributors

HY conceived, designed, and supervised the study. SL and YQ designed the study, collected data, finalized the analysis, wrote the drafts of the manuscript, and interpreted the findings. XR participated in the analysis and mapped the case geographic distribution. TKT, LF, HJ, ZP, JZ, QL, SL participated in the literature search, data collection and analysis. BJC, NAW, MG, WP, PWH, JJF, GFG, AJT interpreted the findings and commented on and helped revise drafts of the manuscript. All authors read and approved the final manuscript.

Conflicts of interest

BJC has received research funding from MedImmune Inc. and Sanofi Pasteur, and consults for Crucell NV. The authors report no other potential conflicts of interest.

Acknowledgments

We thank staff members at county-, district-, prefecture-, and provincial- level CDCs at the provinces with human H5N1 cases occurred for providing assistance with field investigation, administration and data collection. We thank Dr Viktor Gasimov from Ministry of Health of Azerbaijan for data collections for Azerbaijan cases. We thank Dr Nahoko Shindo and Dr Elkhan Gasimov from WHO for assistance in coordination of data collection.

Funding

This study was supported by the grants from the Ministry of Science and Technology of China (2012 ZX10004-201, 2014BAI13B05); the Ministry of Health of China (201202006); China CDC's Key Laboratory of Surveillance and Early-warning on Infectious Disease; the US National Institutes of Health (Comprehensive International Program for Research on AIDS grant U19 AI51915), the Harvard Center for Communicable Disease Dynamics from the National Institute of General Medical Sciences (grant no. U54 GM088558), and the University Grants Committee of the Hong Kong SAR of China (project no. T11-705/14N). SL is supported by the Flowminder Foundation for his postgraduate research on population movement and infectious disease dynamics in the University of Southampton. PWH is funded by the Wellcome Trust (grants 089276/Z/09/Z and 089276/B/09/Z) and the EU FP7 project PREPARE (602525). AJT is supported by funding from NIH/NIAID (U19AI089674), the Bill & Melinda Gates Foundation (OPP1106427, 1032350) and the RAPIDD program of the Science and Technology Directorate, Department of Homeland Security, and the Fogarty International Center, National Institutes of Health. NAW is supported by the UK Medical Research Council (MR/J012343/1).

Authors' information

SL is an assistant investigator and medical epidemiologist of the Division of Infectious Diseases, China CDC, and is also a Ph.D. candidate of the Department of

Geography and Environment at the University of Southampton, UK. HY is the director and medical epidemiologist of the Division of Infectious Diseases, China CDC.

References:

1. Olsen B, Munster VJ, Wallensten A, Waldenstrom J, Osterhaus AD, Fouchier RA. Global patterns of influenza A virus in wild birds. *Science* 2006; **312**(5772):384-8.
2. Durand LO, Glew P, Gross D, et al. Timing of influenza A(H5N1) in poultry and humans and seasonal influenza activity worldwide, 2004-2013. *Emerg Infect Dis* 2015; **21**(2):202-8.
3. Tian H, Zhou S, Dong L, et al. Avian influenza H5N1 viral and bird migration networks in Asia. *Proc Natl Acad Sci U S A* 2015; **112**(1):172-7.
4. Chen H, Smith GJ, Zhang SY, et al. Avian flu: H5N1 virus outbreak in migratory waterfowl. *Nature* 2005; **436**(7048):191-2.
5. Li X, Zhang Z, Yu A, et al. Global and local persistence of influenza A(H5N1) virus. *Emerg Infect Dis* 2014; **20**(8):1287-95.
6. Shortridge KF. Poultry and the influenza H5N1 outbreak in Hong Kong, 1997: abridged chronology and virus isolation. *Vaccine* 1999; **17 Suppl 1**:S26-9.
7. Subbarao K, Klimov A, Katz J, et al. Characterization of an avian influenza A (H5N1) virus isolated from a child with a fatal respiratory illness. *Science* 1998; **279**(5349):393-6.
8. Chan PK. Outbreak of avian influenza A(H5N1) virus infection in Hong Kong in 1997. *Clin Infect Dis* 2002; **34 Suppl 2**:S58-64.
9. Peiris JS, Yu WC, Leung CW, et al. Re-emergence of fatal human influenza A subtype H5N1 disease. *Lancet* 2004; **363**(9409):617-9.
10. Beigel JH, Farrar J, Han AM, et al. Avian influenza A (H5N1) infection in humans. *N Engl J Med* 2005; **353**(13):1374-85.
11. Abdel-Ghaffar AN, Chotpitayasunondh T, Gao Z, et al. Update on avian influenza A (H5N1) virus infection in humans. *N Engl J Med* 2008; **358**(3):261-73.
12. World Health Organization. Influenza at the human-animal interface: Summary and assessment as of 2 October 2014. http://www.who.int/influenza/human_animal_interface/Influenza_Summary_IRA_HA_interface_04December2014.pdf?ua=1 (accessed Apr 28, 2015).
13. Bowman AS, Nelson SW, Page SL, et al. Swine-to-human transmission of influenza A(H3N2) virus at agricultural fairs, Ohio, USA, 2012. *Emerg Infect Dis* 2014; **20**(9):1472-80.
14. Cowling BJ, Jin L, Lau EH, et al. Comparative epidemiology of human infections with avian influenza A H7N9 and H5N1 viruses in China: a population-based study of laboratory-confirmed cases. *Lancet* 2013.
15. Gao R, Cao B, Hu Y, et al. Human infection with a novel avian-origin influenza A (H7N9) virus. *N Engl J Med* 2013; **368**(20):1888-97.

16. Uyeki TM, Cox NJ. Global concerns regarding novel influenza A (H7N9) virus infections. *N Engl J Med* 2013; **368**(20):1862-4.
17. de Vries E, Guo H, Dai M, Rottier PJ, van Kuppeveld FJ, de Haan CA. Rapid Emergence of Highly Pathogenic Avian Influenza Subtypes from a Subtype H5N1 Hemagglutinin Variant. *Emerg Infect Dis* 2015; **21**(5):842-6.
18. World Health Organization. Avian influenza A (H5N1) in Egypt update, 9 April 2015. <http://www.emro.who.int/surveillance-forecasting-response/surveillance-news/avian-influenza-ah5n1-in-egypt-9-april-2015.html> (accessed Apr 21, 2015).
19. World Health Organization. Egypt: upsurge in H5N1 human and poultry cases but no change in transmission pattern of infection. <http://www.emro.who.int/egy/egypt-news/upsurge-h5n1-human-poultry-cases-may-2015.html> (accessed May 21, 2015).
20. World Health Organization. Influenza at the human-animal interface: Summary and assessment as of 31 March 2015. http://www.who.int/influenza/human_animal_interface/Influenza_Summary_IRA_HA_interface_31_March_2015.pdf?ua=1 (accessed Apr 28, 2015).
21. Arafa AS, Naguib MM, Luttermann C, et al. Emergence of a novel cluster of influenza A(H5N1) virus clade 2.2.1.2 with putative human health impact in Egypt, 2014/15. *Euro Surveill* 2015; **20**(13).
22. World Health Organization. Antigenic and genetic characteristics of zoonotic influenza viruses and development of candidate vaccine viruses for pandemic preparedness, February 2015. http://www.who.int/influenza/vaccines/virus/201502_zoonotic_vaccinevirusupdate.pdf?ua=1 (accessed Apr 28, 2015).
23. Zhou L, Liao Q, Dong L, et al. Risk factors for human illness with avian influenza A (H5N1) virus infection in China. *J Infect Dis* 2009; **199**(12):1726-34.
24. Qin Y, Horby PW, Tsang TK, et al. Differences in the Epidemiology of Human Cases of Avian Influenza A(H7N9) and A(H5N1) Viruses Infection. *Clin Infect Dis* 2015; advance published.
25. World Health Organization. Global Alert and Response (GAR): Disease Outbreak News (DONs). <http://www.who.int/csr/don/en/> (accessed Apr 28, 2015).
26. World Health Organization. The Weekly Epidemiological Record (WER). <http://www.who.int/wer/en/>. (accessed Apr 28, 2015).
27. FluTrackers. Global WHO & Ministries of Health Confirmed H5N1 Human Cases List. <https://flutrackers.com/forum/forum/flutrackers-high-pathogenic-h5n1-h1n08-h5n8-h5n6-h5n3-tracking-outbreaks-spread/720310-flutrackers-2015-global-who-ministries-of-health-confirmed-h5n1-human-cases-list> (accessed Apr 28, 2015).
28. World Health Organization. Cumulative number of confirmed human cases of avian influenza A(H5N1) reported to WHO. http://www.who.int/influenza/human_animal_interface/H5N1_cumulative_table_archives/en/ (accessed Apr 28, 2015).
29. Centers for Disease Control and Prevention. Morbidity and Mortality Weekly Report. <http://www.cdc.gov/mmwr/> (accessed Apr 28, 2015).
30. International Society for Infectious Diseases. ProMed-mail. <http://www.promedmail.org/> (accessed Apr 28, 2015).
31. Oner AF, Bay A, Arslan S, et al. Avian influenza A (H5N1) infection in eastern Turkey in 2006.

- N Engl J Med* 2006; **355**(21):2179-85.
32. Mounts AW, Kwong H, Izurieta HS, et al. Case-control study of risk factors for avian influenza A (H5N1) disease, Hong Kong, 1997. *J Infect Dis* 1999; **180**(2):505-8.
 33. World Health Organization. Case definitions for human infections with influenza A(H5N1) virus, 29 August 2006.
http://www.who.int/influenza/resources/documents/case_definition2006_08_29/en/index.html
(accessed July 18, 2014).
 34. World Health Organization. Antigenic and genetic characteristics of zoonotic influenza viruses and candidate vaccine viruses developed for potential use in human vaccines.
http://www.who.int/influenza/vaccines/virus/characteristics_virus_vaccines/en/ (accessed Apr 28, 2015).
 35. Uyeki TM. Human infection with highly pathogenic avian influenza A (H5N1) virus: review of clinical issues. *Clin Infect Dis* 2009; **49**(2):279-90.
 36. Recombinomics. H5N1 Clade 7 Cases in China Raise Concerns, February 2009.
http://www.recombinomics.com/News/02050903/H5N1_China_7_Concerns.html (accessed Apr 28, 2015).
 37. Nguyen T, Rivaller P, Davis CT, et al. Evolution of highly pathogenic avian influenza (H5N1) virus populations in Vietnam between 2007 and 2010. *Virology* 2012; **432**(2):405-16.
 38. Smith GJ, Naipospos TS, Nguyen TD, et al. Evolution and adaptation of H5N1 influenza virus in avian and human hosts in Indonesia and Vietnam. *Virology* 2006; **350**(2):258-68.
 39. Wang J, Vijaykrishna D, Duan L, et al. Identification of the progenitors of Indonesian and Vietnamese avian influenza A (H5N1) viruses from southern China. *J Virol* 2008; **82**(7):3405-14.
 40. Le MT, Wertheim HF, Nguyen HD, et al. Influenza A H5N1 clade 2.3.4 virus with a different antiviral susceptibility profile replaced clade 1 virus in humans in northern Vietnam. *PLoS One* 2008; **3**(10):e3339.
 41. Tian H, Cui Y, Dong L, et al. Spatial, temporal and genetic dynamics of highly pathogenic avian influenza A (H5N1) virus in China. *BMC Infect Dis* 2015; **15**(1):54.
 42. Fang LQ, de Vlas SJ, Liang S, et al. Environmental factors contributing to the spread of H5N1 avian influenza in mainland China. *PLoS One* 2008; **3**(5):e2268.
 43. Mathur MB, Patel RB, Gould M, et al. Seasonal patterns in human A (H5N1) virus infection: analysis of global cases. *PLoS One* 2014; **9**(9):e106171.
 44. Chaichoune K, Wiriyarat W, Thitithyanont A, et al. Indigenous sources of 2007-2008 H5N1 avian influenza outbreaks in Thailand. *J Gen Virol* 2009; **90**(Pt 1):216-22.
 45. Rabinowitz PM, Galusha D, Vegso S, et al. Comparison of human and animal surveillance data for H5N1 influenza A in Egypt 2006-2011. *PLoS One* 2012; **7**(9):e43851.
 46. World Health Organization. Human cases of influenza at the human-animal interface, 2013. *Wkly Epidemiol Rec* 2014; **89**(28):309-20.
 47. Uyeki TM. Global epidemiology of human infections with highly pathogenic avian influenza A (H5N1) viruses. *Respirology* 2008; **13 Suppl 1**:S2-9.
 48. Kandeel A, Manoncourt S, Abd EKE, et al. Zoonotic transmission of avian influenza virus (H5N1), Egypt, 2006-2009. *Emerg Infect Dis* 2010; **16**(7):1101-7.
 49. Patel RB, Mathur MB, Gould M, et al. Demographic and clinical predictors of mortality from highly pathogenic avian influenza A (H5N1) virus infection: CART analysis of international cases.

- PLoS One* 2014; **9**(3):e91630.
50. Kuo HI, Lu CL, Tseng WC, Li HA. A spatiotemporal statistical model of the risk factors of human cases of H5N1 avian influenza in South-east Asian countries and China. *Public Health* 2009; **123**(2):188-93.
 51. Vong S, Ly S, Van Kerkhove MD, et al. Risk factors associated with subclinical human infection with avian influenza A (H5N1) virus--Cambodia, 2006. *J Infect Dis* 2009; **199**(12):1744-52.
 52. Zhou L, Liao Q, Dong L, et al. Risk factors for human illness with avian influenza A (H5N1) virus infection in China. *J Infect Dis* 2009; **199**(12):1726-34.
 53. Yupiana Y, de Vlas SJ, Adnan NM, Richardus JH. Risk factors of poultry outbreaks and human cases of H5N1 avian influenza virus infection in West Java Province, Indonesia. *Int J Infect Dis* 2010; **14**(9):e800-5.
 54. Lohiniva AL, Dueger E, Talaat M, et al. Poultry rearing and slaughtering practices in rural Egypt: an exploration of risk factors for H5N1 virus human transmission. *Influenza Other Respir Viruses* 2013; **7**(6):1251-9.
 55. Dinh PN, Long HT, Tien NT, et al. Risk factors for human infection with avian influenza A H5N1, Vietnam, 2004. *Emerg Infect Dis* 2006; **12**(12):1841-7.
 56. World Health Organization. Influenza at the human-animal interface: Summary and assessment as of 3 March 2015.
http://www.who.int/influenza/human_animal_interface/Influenza_Summary_IRA_HA_interface_3_March_2015.pdf?ua=1 (accessed Apr 29, 2015)
 57. Duan L, Bahl J, Smith GJ, et al. The development and genetic diversity of H5N1 influenza virus in China, 1996-2006. *Virology* 2008; **380**(2):243-54.
 58. Webster RG, Govorkova EA. H5N1 influenza--continuing evolution and spread. *N Engl J Med* 2006; **355**(21):2174-7.
 59. Smith GJ, Naipospos TS, Nguyen TD, et al. Evolution and adaptation of H5N1 influenza virus in avian and human hosts in Indonesia and Vietnam. *Virology* 2006; **350**(2):258-68.
 60. Ibrahim M, Eladl AF, Sultan HA, et al. Antigenic analysis of H5N1 highly pathogenic avian influenza viruses circulating in Egypt (2006-2012). *Vet Microbiol* 2013; **167**(3-4):651-61.
 61. Reperant LA, Kuiken T, Osterhaus AD. Adaptive pathways of zoonotic influenza viruses: from exposure to establishment in humans. *Vaccine* 2012; **30**(30):4419-34.

Box 1. Definition of exposures to poultry and humans

Type of exposure	Definition
<i>Occupational exposure to live poultry</i>	refers to poultry related professional exposure (e.g. poultry breeding, trafficking, sale, and quarantine) in the two weeks prior to the onset of illness.
<i>Visiting live bird market (LBM)</i>	refers to the visit of a wholesale or retail market of live poultry or birds within the two weeks prior to the onset of symptoms.
<i>Exposure to sick or dead poultry</i>	refers to direct physical contact with, or in proximity to, sick or dead poultry or its production or faeces in the two weeks prior to onset.
<i>Exposure to backyard poultry</i>	refers to whether the case had been exposed to poultry raised in the backyard within two weeks prior to onset.
<i>Any exposure to poultry</i>	refers to direct contact, indirect contact or proximity to healthy, sick or dead poultry (including any kind of poultry or birds, e.g. chicken, ducks, goose, pet birds, pigeons, et al.) in LBMs, backyard, farm, or neighborhood, or consumption of improperly processed poultry products.
<i>Human case contact</i>	refers to a patient with a history of close contact with

	<p>a confirmed or probable human H5N1 case (any time from the day before the onset of illness to death or during the period that case was hospitalized) in the two weeks before the onset of symptoms.</p>
--	--

Table 1. The characteristics of human case with H5N1 virus infection by geographic region, May 1997 – April 2015

Characteristics	Total (n=907)	East and Southeast Asia (n=505)	North Africa (n=363)	Other (n=39)
Type of cases				
Confirmed case	858 (94.6%)	479 (94.9%)	343 (94.5%)	36 (92.3%)
Probable case	49 (5.4%)	26 (5.1%)	20 (5.5%)	3 (7.7%)
Sex				
Female	476 (52.5%)	246 (48.7%)	213 (58.7%)	17 (43.6%)
Unknown	29 (3.2%)	21 (4.2%)	6 (1.7%)	2 (5.1%)
Age				
Median (yrs, range)	19 (0.25, 86)	19 (0.3, 75)	20 (0.25, 86)	15 (1.3, 52)
Final outcome				
Death	483 (53.3%)	349 (69.1%)	116 (32%)	18 (46.2%)
Unknown	4 (0.4%)	2 (0.4%)	2 (0.6%)	0 (0)
Hospitalization				
Yes	819 (90.3%)	438 (86.7%)	353 (97.2%)	28 (71.8%)
Unknown	82 (9%)	64 (12.7%)	9 (2.5%)	9 (23.1%)
Median of time delay (days, range)				
Time from onset to hospital admission	4 (0, 90)	5 (0, 90)	3 (0, 33)	2 (0, 13)

Unknown	184 (20.3%)	121 (24%)	46 (12.7%)	17 (43.6%)
---------	-------------	-----------	------------	------------

Time from hospital admission	5 (0, 116)	4 (0, 116)	5 (0, 28)	5 (2, 25)
------------------------------	------------	------------	-----------	-----------

to death or discharge

(recovery)

Unknown	403 (44.4%)	166 (32.9%)	219 (60.3%)	18 (46.2%)
---------	-------------	-------------	-------------	------------

Time from onset to death or	10 (0, 119)	10 (0, 119)	10 (2, 32)	9 (2, 32)
-----------------------------	-------------	-------------	------------	-----------

discharge (recovery)

Unknown	360 (39.7%)	124 (24.6%)	221 (60.9%)	15 (38.5%)
---------	-------------	-------------	-------------	------------

Predominant clade or subclade

0	18 (2%)	18 (3.6%)	0 (0)	0 (0)
---	---------	-----------	-------	-------

1	193 (21.3%)	193 (38.2%)	0 (0)	0 (0)
---	-------------	-------------	-------	-------

2.1	208 (22.9%)	208 (41.2%)	0 (0)	0 (0)
-----	-------------	-------------	-------	-------

2.2	393 (43.3%)	0 (0)	363 (100%)	30 (76.9%)
-----	-------------	-------	------------	------------

2.3	89 (9.8%)	84 (16.6%)	0 (0)	5 (12.8%)
-----	-----------	------------	-------	-----------

7	2 (0.2%)	2 (0.4%)	0 (0)	0 (0)
---	----------	----------	-------	-------

Unknown	4 (0.4%)	0 (0)	0 (0)	4 (10.3%)
---------	----------	-------	-------	-----------

Exposure history

Any exposure to poultry	748 (82.5%)	382 (75.6%)	339 (93.4%)	27 (69.2%)
-------------------------	-------------	-------------	-------------	------------

Unknown	126 (13.9%)	94 (18.6%)	24 (6.6%)	8 (20.5%)
---------	-------------	------------	-----------	-----------

Occupational exposure to live	15 (1.7%)	12 (2.4%)	2 (0.6%)	1 (2.6%)
-------------------------------	-----------	-----------	----------	----------

poultry

Unknown	586 (64.6%)	289 (57.2%)	286 (78.8%)	11 (28.2%)
Visit LBMs	82 (9%)	68 (13.5%)	11 (3%)	3 (7.7%)
Unknown	596 (65.7%)	296 (58.6%)	286 (78.8%)	14 (35.9%)
Exposure to sick or dead poultry	439 (48.4%)	242 (47.9%)	174 (47.9%)	23 (59%)
Unknown	395 (43.6%)	217 (43%)	166 (45.7%)	12 (30.8%)
Exposure to backyard poultry	188 (20.7%)	113 (22.4%)	64 (17.6%)	11 (28.2%)
Unknown	601 (66.3%)	301 (59.6%)	286 (78.8%)	14 (35.9%)
Human case contact	49 (5.4%)	35 (6.9%)	3 (0.8%)	11 (28.2%)
Unknown	115 (12.7%)	86 (17%)	21 (5.8%)	8 (20.5%)

Note: Data are presented as no. (%) of patients unless otherwise indicated. LBMs:

Live bird markets. East and Southeast Asia (505 cases): Indonesia (208), Viet Nam (134), Cambodia (58), mainland China (52), Thailand (27), Hong Kong SAR (23), Laos (2), and Myanmar (1); North Africa (363 cases): Egypt (363); Other (39 cases): Turkey (12), Azerbaijan (9), Bangladesh (7), Pakistan (4), Iraq (3), Nigeria (2), Djibouti (1), and Canada (1). Data on H5N1 clade or subclade of Human cases was based on the reports from WHO website, or the literature, and the known geographic distribution of the viruses. No all cases were laboratory confirmed and reported with clade results, so we presumed that the case in each area was infected by the reported predominant clade or subclade of H5N1 virus in the same period and area. The clade or subclade in each area were clade 0 in

Hong Kong SAR in 1997, clade 1 in Viet Nam, Cambodia, Thailand, and Hong Kong SAR, subclade 2.1 mainly in Indonesia, 2.2 in Egypt, Turkey, Azerbaijan, Bangladesh, Iraq, Nigeria and Djibouti, and 2.3 in Viet Nam, Viet Nam, Bangladesh, Laos, Canada and Myanmar, and 7 in mainland China. The data of clade was unavailable for 4 cases in Pakistan in 2007.

Table 2. The clade or subclade and fatality of human case with H5N1 virus infection, May 1997 – April 2015

Clade or subclade	Year first identified	Locations identified	Case fatality risk
0	1997	Hong Kong SAR	31.6% (6/18)
1	2003	Hong Kong SAR, Vietnam, Cambodia and Thailand	58.6% (112/191)
2.1	2005	Indonesia	84.6% (176/208)
2.2	2005	Turkey, Egypt, Azerbaijan, Djibouti, Iraq, Nigeria, and Bangladesh	33.2% (130/391)
2.3	2005	Mainland China, Laos, Myanmar, Vietnam, Hong Kong SAR, Bangladesh and Canada	61.8% (55/89)
7	2003	Mainland China	100% (2/2)

Note: Data on H5N1 clade or subclade of Human cases was based on the reports from WHO website, or the literature, and the known geographic distribution of the viruses. No all cases were laboratory confirmed and reported with clade results, so we presumed that the case was infected by the reported clade or subclade of H5N1 virus in the same period and area. The data of clade was unavailable for four cases in Pakistan in 2007, and four cases with unknown outcome (two of Viet Nam in 2005 and two of Egypt in 2015) were also excluded.

Panel: Research in Focus

Systematic Review

We searched PubMed on June 26, 2015, with the terms “H5N1” and “avian influenza” for all journal articles written in English between May 1, 1997 and April 30, 2015. In the past 18 years, avian influenza A(H5N1) virus infection in humans has been increasingly reported across the world. Most reports of human cases were case study within one country,⁹ with very few exceptions that reviewed or descriptively analyze global epidemiology before 2010 and just including the human cases reported to WHO since 2003,^{10,11} and few previous reports had compared the epidemiology of East and Southeast Asia with that in North Africa.⁴⁷ Following the emergence of a novel cluster of influenza A(H5N1) virus clade 2.2.1.2 in Egypt in 2014-15,²¹ the incidence of human infection has increased dramatically in Egypt since November 2014,¹⁸ but no previous scientific articles had compared the epidemiology of human infection in Egypt before and after November 2014.

Interpretation

Our study showed that a total of 907 human H5N1 cases were reported globally by the WHO, national health authorities or other open sources spanning 18 years between 1 May 1997 and 30 April 2015, with no cases reported between 1998 and 2002, and the largest annual numbers recorded in 2015 (Figure 2). The affected countries has expanded from East and Southeast Asia, then to West Asia

and Africa, especially between 2003 and 2008, with the CFR varied across geographical regions ($p < 0.001$) and an obvious seasonality (Figure 4). The age of cases differed significantly by outcome ($p < 0.001$), with a median age of 22 years for fatal cases and 10 years for non-fatal cases. Although the incidence of human cases in Egypt has increased dramatically from November 2014 through April 2015, but compared to the cases before November 2014 there were no significant differences in the CFR, history of exposure to poultry, history of human case contact, and time from onset to hospitalization. In view of the seasonal pattern in human infections of H5N1, we must prepare for this disease to reappear later this year, not just in North Africa and but also in East and Southeast Asia.

Figure Legends

Figure 1: The geographic distribution of human cases with H5N1 virus infection by outcome, May 1997 – April 2015 (n=907)

The data for China includes the cases reported by mainland China (52 cases) and Hong Kong SAR (23 cases).

Figure 2: Epidemic curve of human cases with H5N1 virus infection by region, May 1997 – April 2015

(A) The epidemic curve of H5N1 human cases reported globally (884 cases). (B) East and Southeast Asia (484 cases) includes Indonesia (187), Viet Nam (134), Cambodia (58), mainland China (52), Thailand (27), Hong Kong SAR (23), Laos (2), and Myanmar (1). (C) North Africa (363 cases) includes Egypt (363).

Twenty-three cases with unknown month of illness (21 cases of Indonesia in 2009 and two cases of Turkey in 2006) are excluded from this epidemic curve.

Figure 3: The age distribution of human cases with H5N1 virus infection by gender, geographic regions and outcome, May 1997 – April 2015

(A) The age distribution of all cases by male (n=401) and female (n=476). (B) The age distribution of all cases by death (n=463) and survive (n=416). (C) The age distribution of survive cases by North Africa (n=245), East and Southeast Asia (n=152). (D) The age distribution of death cases by North Africa (n=116),

East and Southeast Asia (n=329). (E) The age distribution of survive cases in Egypt before (n=114) and since 1 November 2014 (n=131). (F) The age distribution of death cases in Egypt before (n=64) and since 1 November 2014 (n=52).

Figure 4: Heat map of the reported data of human cases with H5N1 virus infection by country/district, sorted by geographical region and the date of the first cases illness onset, May 1997 – April 2015

(A) Time series of monthly cases, standardized by the total number of cases in each country/region. Twenty-three cases with unknown month of illness (21 cases of Indonesia in 2009 and two cases of Turkey in 2006) are excluded. (B) Seasonal distribution of cases, plotted as the mean value of the proportion of weekly case number in the total number of each country/region. Forty-three cases with unknown week of onset are excluded: Indonesia (21 cases), Egypt (18), Turkey (2), Cambodia (1) and Vietnam (1). HK SAR: Hong Kong Special Administrative Region of China.

Figure 5: The distribution of days from onset to hospital admission of human H5N1 cases by outcome and geographic region, May 1997 – April 2015

(A) The time distribution of all fatal (n=380) and non-fatal (n=343) cases. (B) The time distribution of fatal (n=268) and non-fatal (n=116) cases in East and

Southeast Asia. (C) The time distribution of fatal (n=99) and non-fatal (n=218) cases in North Africa.

Figure 6: The geographic distribution of human cases with H5N1 virus infection by outcome in Egypt, March 2006 - April 2015 (n=363)

Supporting Information

Appendix Table 1 The data source of human case with H5N1 virus infection in each country, May 1997 – April 2015.

Appendix Table 2 The list of variables in the individual dataset of human case with H5N1 virus infection, May 1997 – April 2015.

Appendix Table 3 Demographic and Epidemiologic characteristics of human case with H5N1 virus infection by outcomes, May 1997 – April 2015.

Appendix Table 4 The characteristics of human case with H5N1 virus infection in Egypt before and since 1 November 2014.

Appendix Figure 1 Epidemic curve of human cases with H5N1 virus infection by climate zones, May 1997 – April 2015.

Appendix Figure 2 The number of human cases with H5N1 virus infection by year and geographic region, May 1997 – April 2015 (N=907).

Appendix Figure 3 The number of human cases with H5N1 virus infection by year and country, May 1997 – April 2015 (N=907).

Appendix Figure 4 The seasonality of human cases with H5N1 virus infection by the month of illness onset, May 1997 – April 2015.