

**Comment on “Interplay of deformation and magmatism in the Pangong Transpressional Zone, Eastern Ladakh, India: Implications for remobilization of the trans-Himalayan magmatic arc and initiation of the Karakoram Fault” by K. Sen, B.K. Mukherjee and A.S. Collins, Journal of Structural Geology 62 (2014) 13-24**

D. Wallis<sup>\*1</sup>, A.J. Parsons<sup>1</sup>, R.J. Phillips<sup>1</sup>, M.P. Searle<sup>2</sup> & E.C. Ferré<sup>3</sup>

*\*Corresponding author*

<sup>1</sup>*School of Earth and Environment, University of Leeds, Leeds, UK, LS2 9JT*

<sup>2</sup>*Department of Earth Sciences, University of Oxford, Oxford, UK, OX1 3AN*

<sup>3</sup>*Department of Geology, Southern Illinois University, Carbondale IL 62901, USA*

**Keywords: Karakoram Fault, microstructure, anisotropy of magnetic susceptibility, leucogranite, magmatism, deformation**

**1. Introduction**

Sen *et al.* (2014) address the issue of the age of the Karakoram Fault Zone (KFZ), western Tibet, and suggest that the fault has accommodated significant eastward extrusion of the Tibetan plateau since initiating prior to *c.* 23 Ma. Anisotropy of magnetic susceptibility (AMS) data provided by Sen *et al.* potentially provide a new approach to investigating the relationship between the KFZ and the *c.* 22-16 Ma Tangtse-Darbuk leucogranite (TDL) ([Phillips \*et al.\*, 2004](#)). However, we show that misinterpretation of key microstructures has led Sen *et al.* to reach conclusions starkly at odds with the evidence that they present. Furthermore, we demonstrate that the study lacks the cautionary procedures and considerations that are crucial to the valid interpretation of AMS data. We show that the findings of Sen *et al.* are entirely consistent with pre-kinematic rather than syn-kinematic emplacement of the TDL. We conclude that the KFZ initiated after cooling of the TDL to ambient

23 lower amphibolite grade conditions after  $15.55 \pm 0.74$  Ma ([Phillips \*et al.\*, 2004](#)) and is a relatively  
24 recent structure of the India-Asia collision zone.

## 25 **2. Re-interpretation of field and microstructural observations**

26 Sen *et al.* recognise that the main bulk of the TDL is relatively undeformed and that strain increases  
27 dramatically towards the Tangtse fault strand. This structural relationship is consistent with the  
28 findings of previous studies of the TDL ([Phillips and Searle, 2007](#); [Wallis \*et al.\*, 2013](#)) and is widely  
29 recognised as indicative of pre-kinematic intrusion of granitic bodies ([Lamouroux \*et al.\*, 1980](#);  
30 Paterson and Tobisch, 1988). Sen *et al.* provide AMS data for fifteen KFZ samples and suggest that  
31 ten show magnetic fabrics consistent with the known kinematics of the Pangong Transpressional  
32 Zone (PTZ). Notably, two samples (7/4B and 7/5A) from the TDL interior show AMS fabrics  
33 discordant to the KFZ deformation, which Sen *et al.* suggest result from magma chamber convection.  
34 These discordant fabrics, preserved outside the marginal KFZ shear zones, fulfil another important  
35 criterion for identification of pre-kinematic granites (Miller and Paterson, 1995; Paterson *et al.*,  
36 1998).

37 The syn-kinematic interpretation of the TDL put forward by Sen *et al.* relies on their statement that  
38 samples “MST 7/3A and MST 7/3B, which are relatively undeformed, show a magnetic fabric that is  
39 emplacement related and not tectonized”, presumably meaning that the magnetic fabric formed after  
40 emplacement but before full crystallisation. The evidence that they put forward as showing that these  
41 samples are not ‘tectonized’ are their figures 4a-c. Figures 4a and 4b show only a large garnet in  
42 MST 7/3B and are insufficient to determine whether or not the sample is tectonized as the matrix is  
43 not shown. The quartz microstructure in the sample is shown however in Figure 4c, where the  
44 authors describe it as “devoid of dynamic recrystallization features and lacking any preferred  
45 orientation, showing regular grain boundaries and undulatory extinction.” This description is  
46 inconsistent with the microstructure shown in their Figure 4c. The highly irregular, amoeboid and

47 embayed quartz grain boundaries are indicative of deformation by grain boundary migration dynamic  
48 recrystallization. Sub-grain walls are also present within larger grains, suggesting a component of  
49 sub-grain rotation. These combined deformation mechanisms indicate deformation at *c.*500°C at  
50 typical geological shear zone strain rates of *c.*10<sup>-12</sup> s<sup>-1</sup> (Stipp *et al.*, 2002) and are consistent with  
51 previous microstructural observations and deformation temperature estimates within the Tangtse  
52 fault strand (Wallis *et al.*, 2013). Higher temperature quartz deformation microstructures such as  
53 “chess board” extinction, which form at near-solidus temperatures (Blumenfeld *et al.*, 1986), are  
54 absent from the images provided. An alternative explanation for the quartz deformation  
55 microstructures could be that they formed by deformation during, and resulting from, ascent and  
56 emplacement of partially crystallised granitic magma, as proposed by Sen and Collins (2013) for the  
57 Ladakh batholith. However, such an interpretation should be supported by evidence of  
58 ascent/emplacement related deformation microstructures (e.g. crystal tiling, magmatic growth in  
59 pressure shadows). Similarly, the interpretation by Sen *et al.* that the TDL magma chamber  
60 underwent significant convection supports emplacement of mobile fluid magma rather than a crystal  
61 mush undergoing dynamic recrystallization. Sen *et al.* note that the centre of the TDL preserves  
62 magmatic or high temperature microstructures such as fluid/melt inclusions in garnet and “melt”  
63 filled cleavage and fractures in feldspar. These microstructures alone are insufficient to support syn-  
64 ascent/emplacement deformation of the quartz or a syn-KFZ interpretation of the TDL as they may  
65 (most likely given the aforementioned evidence) result from normal magmatic processes including  
66 deformation during magmatic convection, as suggested by Sen *et al.* for the magnetic fabrics of the  
67 TDL interior. The concordance between the magnetic fabrics in samples MST 7/3A and MST 7/3B  
68 and the wider macroscopic, microstructural and magnetic deformation fabrics of the KFZ is therefore  
69 best explained by deformation of these samples at *c.*500°C in the margin of the Tangtse fault strand,  
70 not by deformation during ascent/emplacement or sub-solidus cooling. The evidence put forward by  
71 Sen *et al.* is therefore entirely consistent with a pre-KFZ interpretation of the TDL and lends no

72 support to the syn-KFZ interpretation. The KFZ must therefore have initiated after final solidification  
73 of the TDL at  $15.55 \pm 0.74$  Ma ([Phillips \*et al.\*, 2004](#)). The syn-kinematic migmatite structures formed  
74 at  $c.17.4$  Ma ([Phillips \*et al.\*, 2013](#)) therefore do not record anatexis and melt migration within the  
75 KFZ shear zone but provide valuable information on magma migration through the regionally  
76 deforming crust of the Karakoram terrane prior to KFZ initiation.

### 77 **3. Critique of AMS analysis**

78 AMS can serve as a proxy for deformation when (1) all magnetic carriers are identified, that (2) all  
79 possible controls on AMS are carefully taken into account and (3) a genetic relationship between the  
80 origin of AMS and deformation fabrics is confirmed ([Borradaile & Jackson, 2004, 2010](#)). Modern  
81 AMS investigations ([Borradaile \*et al.\*, 2011](#); [Kontny \*et al.\*, 2011](#); [Kruckenberg \*et al.\*, 2010](#)) include a  
82 variety of corroborative data (other than AMS data) to identify the magnetic carriers, and determine  
83 the grain size and extent of magnetostatic interactions of any ferromagnetic phases. These additional  
84 procedures include magnetic hysteresis measurements ([Tauxe \*et al.\*, 2002](#)), Isothermal Remanent  
85 Magnetisation (IRM) acquisition curve analysis ([Robertson & France, 1994](#)), FORC analysis  
86 ([Roberts \*et al.\*, 2000](#)) and thermomagnetic experiments ([Ferré \*et al.\*, 2003](#)). In some cases it is also  
87 necessary to determine the petrological relationship between the magnetic carriers and surrounding  
88 crystal fabric with microscopy and/or SPO or CPO measurements ([Kruckenberg \*et al.\*, 2010](#)). By  
89 failing to conduct most of these additional analyses, [Sen \*et al.\*](#) are unable to correctly identify the  
90 magnetic carriers and cannot rigorously evaluate the controls that may influence their AMS results.

91 [Sen \*et al.\*](#) present AMS data from 15 samples identified as leucogranite, dioritic gneiss, pink  
92 mylonitised granite and migmatite. [Sen \*et al.\*](#) first use bulk susceptibility ( $K_m$ ) and corrected degree  
93 of anisotropy ( $P'$ ) ([Jelínek, 1981](#)) to split their sample suite into paramagnetic and ferromagnetic  
94 samples (Figure 5a, [Sen \*et al.\*](#)). [Sen \*et al.\*](#) correctly note that many of the dioritic gneisses have high  
95 values of  $K_m (>10^{-3} SI)$  which are typical values for ferromagnetic materials. [Sen \*et al.\*](#) use a single

96 thermal magnetisation experiment on sample 7/3AX (Figure 5b, Sen *et al.*) to infer the presence of  
97 magnetite in all sample lithologies other than leucogranite. Whilst most of the values of  $K_m$  are high  
98 for these samples ( $>10^{-3} SI$ ), the variability in  $K_m$  and  $P'$  suggests that magnetite may not be the  
99 magnetic carrier in all of these samples and at least two of these samples (7/7, 26/8/4, 27/8/2A) are  
100 most likely to be paramagnetic. This variability in  $K_m$  and  $P'$  highlights the inappropriateness of  
101 using a single thermomagnetic result to determine the magnetic carriers of multiple samples with a  
102 variety of lithologies. Valid identification of the magnetic carriers requires further investigation,  
103 either via additional thermal magnetisation experiments or more suitably via magnetic hysteresis  
104 analysis (Tauxe *et al.*, 2002) and IRM acquisition curve analysis (Robertson & France, 1994). Where  
105 ferromagnetic phases are identified, the grain sizes and magnetostatic interactions between these  
106 grains, which also control AMS, should also be evaluated through magnetic hysteresis and FORC  
107 analysis (Roberts *et al.*, 2000; Dunlop, 2002).

108 There are other unexplained errors presented in the  $K_m$ ,  $P'$  and  $T$  plots in Figures 5a and 5d. Figure  
109 5a ( $K_m$  vs.  $P'$ ) displays seven leucogranite data points (white circles) and eight dioritic gneiss data  
110 points (black squares), whilst Figure 5d ( $T$  vs.  $P'$ ) displays eight leucogranite data points and seven  
111 dioritic gneiss data points. Figure 5a displays an anomalous data point ( $K_m = 17500 : P' = 1.7$ ) that is  
112 not presented in Table 2. Only three leucogranite data points with  $P' > 1.2$  are displayed in Figure 5a,  
113 whereas four leucogranite samples have  $P' > 1.2$  in Table 2. Sample 27/8/2A (migmatite) is  
114 represented by a white circle (i.e. leucogranite) in Figure 5a and a black square (diorite gneiss) in  
115 Figure 5d. We highlight this last error because if sample 27/8/2A had been displayed as a black  
116 square amongst the white circled leucogranite data cluster then Figure 5a would actually disagree  
117 with Sen *et al.*'s assumption that magnetite controls the AMS of all samples except leucogranite. The  
118 values of  $K_m$  displayed in Table 2 and Figure 5a should also be presented as an order of  $\times 10^{-6}$   
119 (Tarling & Hrouda, 1993).

120 Sen *et al.* state that the AMS fabrics of the diorite gneisses, migmatites and pink mylonitic granites  
121 are concordant with the local strike of the KFZ and must be representative of lateral shearing along  
122 the KFZ. However, the magnetic foliation of half of these samples is misaligned with the strike and  
123 dip of the local structural foliation by 15-90° and 9-38° respectively. Sen *et al.* claim that the  
124 discordances between the orientations of the structural and magnetic fabrics “can be attributed to  
125 scattered crystallization of very fine grained magnetite having no significant preferred orientation,”  
126 however, there is no clear evidence that magnetite controls the AMS of all of these samples and  
127 without knowing the nature of all magnetic carriers and their relation to the surrounding deformation  
128 fabric, it is not possible to justify a correlation between magnetic and deformation fabrics.  
129 Furthermore, Sen *et al.*’s explanation is actually incorrect. A scattering of magnetite grains with no  
130 significant preferred orientation would produce a very scattered AMS fabric with wide confidence  
131 ellipses surrounding the mean AMS axis orientations. Most of the AMS fabrics presented in Figure 6  
132 are actually very well defined, with tight clusters and small confidence ellipses around the mean  
133 AMS axes.

134 Sen *et al.* present good evidence to suggest that biotite is the magnetic carrier in the leucogranite  
135 samples and that their random orientations within the Durbuk Pluton are likely to represent a  
136 magmatic fabric. However, the authors’ interpretation that the AMS fabrics of leucogranite samples  
137 7/3A and 7/3B represent syn-kinematic magmatic fabrics is less convincing. Such interpretations  
138 require evidence of a KFZ aligned magmatic fabric in the absence of solid-state deformation. We  
139 have demonstrated that the microstructural evidence from sample 7/3B is actually typical of dynamic  
140 recrystallisation textures formed during solid-state deformation. It should also be noted that the  
141 accompanying AMS fabric from this sample is actually *misaligned* with the strike and dip of the  
142 KFZ by 28-43° and 27-42° respectively. We find the AMS orientation of 7/3B to be statistically  
143 indistinguishable from the random AMS orientations of the rest of the leucogranite samples and  
144 cannot justify a correlation between the AMS of this sample and KFZ-related deformation. Without

evidence for a lack of solid state deformation from 7/3A, it is not possible to suggest that this AMS fabric was formed during syn-kinematic magmatic flow. It is just as likely that the strong AMS fabric in 7/3A is due to localised shearing at the margin of the pluton after crystallisation.

#### 4. Concluding Remarks

Mis-interpretation of critical microstructures has led Sen *et al.* to a syn-kinematic emplacement interpretation of the TDL. Instead, the field and microstructural evidence that they observe is entirely consistent with a pre-KFZ interpretation of the TDL, and precludes syn-KFZ emplacement. Furthermore, the AMS results reported by Sen *et al.* lack the supporting magnetic data needed to determine and evaluate the AMS controls. Without these constraints, a valid correlation between AMS and deformation fabrics cannot be made. We conclude that KFZ deformation in this region commenced after solidification of the TDL at  $15.55 \pm 0.74$  Ma (sample P1, [Phillips \*et al.\*, 2004](#)) and cooling to ambient lower amphibolite grade conditions ([Wallis \*et al.\*, 2013](#)).

#### References

- [Blumenfeld, P., Mainprice, D., Bouchez, J.L., 1986. C-slip in quartz from subsolidus deformed granite. \*Tectonophysics\* 127, 97-115.](#)
- [Lamouroux, C., Soula, J.C., Deramond, J., Debut, P., 1980. Shear zones in the granodioritic massifs of the central Pyrenees and the behaviour of these massifs during Alpine orogenesis. \*J. Struct. Geol.\* 2, 49-53.](#)
- [Borradaile, G.J., Almqvist, B.S.G., Geneviciene, I., 2011. Anisotropy of magnetic susceptibility \(AMS\) and diamagnetic fabrics in the Durness Limestone, NW Scotland. \*J. Struct. Geol.\* 34, 54-60.](#)
- [Borradaile, G.J., Jackson, M., 2004. Anisotropy of magnetic susceptibility \(AMS\): magnetic petrofabrics of deformed rocks. \*Geol. Soc., Lond., Spec. Pub.\* 238, 299-360.](#)
- [Borradaile, G.J., Jackson, M., 2010. Structural geology, petrofabrics and magnetic fabrics \(AMS, AARM, AIRM\). \*J. Struct. Geol.\* 32, 1519-1551.](#)

169 [Dunlop, D.J., 2002. Theory and application of the Day plot \(Mrs/Ms versus Hcr/Hc\) 1. Theoretical](#)  
170 [curves and tests using titanomagnetite data. J. Geophys. Res.: S. Earth 107, EPM 4-1-EPM 4-22.](#)

171 [Ferré, E.C., Teyssier, C., Jackson, M., Thill, J.W., Rainey, E.S.G., 2003. Magnetic susceptibility](#)  
172 [anisotropy: A new petrofabric tool in migmatites. J. Geophys. Res.: S. Earth 108, 2086.](#)

173 [Jelínek, V., 1981. Characterization of the magnetic fabric of rocks. Tectonophys. 79, T63-T67.](#)

174 [Kontny, A., Engelmann, R., Grimmer, J., Greiling, R., Hirt, A., 2011. Magnetic fabric development](#)  
175 [in a highly anisotropic magnetite-bearing ductile shear zone \(Seve Nappe Complex, Scandinavian](#)  
176 [Caledonides\). Int. J. Earth Sci., 10.1007/s00531-011-0713-8, 1-22.](#)

177 [Kruckenberg, S.C., Ferré, E.C., Teyssier, C., Vanderhaeghe, O., Whitney, D.L., Seaton, N.C.A.,](#)  
178 [Skord, J.A., 2010. Viscoplastic flow in migmatites deduced from fabric anisotropy: An example](#)  
179 [from the Naxos dome, Greece. J. Geophys. Res, 115, B09401.](#)

180 [Miller, R.B., Paterson, S.R., 1995. Construction of mid-crustal magma chambers during regional](#)  
181 [contraction, North Cascades, Washington. In: Brown, M., Piccoli, P.M. \(Eds.\), The Origin of](#)  
182 [Granites and Related Rocks. U.S. Geol. Surv. Circ. 1129, 97-98.](#)

183 [Paterson, S.R., Fowler, T.K., Schmidt, K.L., Yoshinobu, A.Y., Yuan, E.S., Miller, R.B., 1998.](#)  
184 [Interpreting magmatic fabric patterns in plutons. Lithos 44, 53-82.](#)

185 [Paterson, S.R., Tobisch, O.T., 1988. Using pluton ages to date regional deformations: Problems with](#)  
186 [commonly used criteria. Geology 16, 1108-1111.](#)

187 [Phillips, R.J., Parrish, R.R., Searle, M.P., 2004. Age constraints on ductile deformation and long-](#)  
188 [term slip rates along the Karakoram fault zone, Ladakh. E. Plan. Sci. Lett. 226, 305-319.](#)

189 [Phillips, R.J., Searle, M.P., 2007. Macrostructural and microstructural architecture of the Karakoram](#)  
190 [fault: relationship between magmatism and strike-slip faulting. Tectonics 26, TC3017.](#)

191 [Phillips, R.J., Searle, M.P., Parrish, R.R., 2013. The geochemical and temporal evolution of the](#)  
192 [continental lithosphere and its relationship to continental-scale faulting: The Karakoram Fault,](#)  
193 [eastern Karakoram, NW Himalayas. Geochem., Geophys., Geosys. 14, 583-603.](#)



194 [Robertson, D.J., France, D.E., 1994. Discrimination of remanence-carrying minerals in mixtures,](#)  
195 [using isothermal remanent magnetisation acquisition curves. Phys. Earth Plan. Int. 82, 223-234.](#)

196 [Roberts, A.P., Pike, C.R., Verosub, K.L., 2000. First-order reversal curve diagrams: A new tool for](#)  
197 [characterizing the magnetic properties of natural samples. J. Geophys. Res.: S. Earth 105, 28461-](#)  
198 [28475.](#)

199 [Sen, K., Collins, A.S., 2013. Reply to comment on “Dextral transpression and late-Eocene](#)  
200 [magmatism in the trans-Himalayan Ladakh Batholith \(North India\): implications for tectono-](#)  
201 [magmatic evolution of the Indo-Eurasian collisional arc. Int. J. Earth Sci. 102, 973-975.](#)

202 Sen, K., Mukherjee, B.K., Collins, A.S., 2014. Interplay of deformation and magmatism in the  
203 Pangong Transpressional Zone, Eastern Ladakh, India: Implications for remobilization of the trans-  
204 Himalayan magmatic arc and initiation of the Karakoram Fault. J. Struct. Geol. 62, 13-24.

205 Stipp, M., Stünitz, H., Heilbronner, R., Schmid, S.M., 2002. Dynamic recrystallization of quartz:  
206 correlation between natural and experimental conditions. In: De Meer, S., Drury, M.R., De Bresser,  
207 J.H.P., Pennock, G.M. (Eds.), Deformation Mechanisms, Rheology and Tectonics: Current Status  
208 and Future Perspectives. Geol. Soc., Lond., Spec. Pub. 200, 171-190.

209 Tarling, D.H., Hrouda, F., 1993. The magnetic anisotropy of rocks, 1<sup>st</sup> ed., 217 pp, Chapman and  
210 Hall, London.

211 Tauxe, L., Bertram, H.N., Seberino, C., 2002. Physical interpretation of hysteresis loops:  
212 Micromagnetic modeling of fine particle magnetite. Geochem. Geophys. Geosys. 3, 1055.

213 [Wallis, D., Phillips, R.J., Lloyd, G.E., 2013. Fault weakening across the frictional-viscous transition](#)  
214 [zone, Karakoram Fault Zone, NW Himalaya. Tectonics 32, 1227-1246.](#)