

Volatility Forecast Comparison

Using Imperfect Volatility Proxies

Andrew J. Patton*

University of Oxford

First version: March 2004. This version: 11 August, 2007.

The use of a conditionally unbiased, but imperfect, volatility proxy can lead to undesirable outcomes in standard methods for comparing conditional variance forecasts. We motivate our study with analytical results on the distortions caused by some widely-used loss functions, when used with standard volatility proxies such as squared returns, the intra-daily range or realised volatility. We then derive necessary and sufficient conditions on the functional form of the loss function for the ranking of competing volatility forecasts to be robust to the presence of noise in the volatility proxy, and derive some useful special cases of this class of “robust” loss functions. The methods are illustrated with an application to the volatility of returns on IBM over the period 1993 to 2003.

Keywords: forecast evaluation, forecast comparison, loss functions, realised variance, range.

J.E.L. Codes: C53, C52, C22.

*The author would particularly like to thank Peter Hansen, Ivana Komunjer and Asger Lunde for helpful suggestions and comments. Thanks also go to Torben Andersen, Tim Bollerslev, Peter Christoffersen, Rob Engle, Christian Gourieroux, Tony Hall, Mike McCracken, Nour Meddahi, Roel Oomen, Adrian Pagan, Neil Shephard, Kevin Sheppard, and Ken Wallis. Runquan Chen provided excellent research assistance. The author gratefully acknowledges financial support from the Leverhulme Trust under grant F/0004/AF. Some of the work on this paper was conducted while the author was a visiting scholar at the School of Finance and Economics, University of Technology, Sydney. Contact address: Department of Economics, and Oxford-Man Institute of Quantitative Finance, University of Oxford, Manor Road, Oxford OX1 3UQ, United Kingdom. Email: andrew.patton@economics.ox.ac.uk. Matlab code used in this paper is available from <http://www.economics.ox.ac.uk/members/andrew.patton/code.html>.

1 Introduction

Many forecasting problems in economics and finance involve a variable of interest that is unobservable, even *ex post*. The most prominent example of such a problem is the forecasting of volatility for use in financial decision-making. Other problems include forecasting the true rates of inflation, GDP growth or unemployment (not simply the announced rates); forecasting trade intensities; and forecasting default probabilities or ‘crash’ probabilities. While evaluating and comparing economic forecasts is a well-studied problem, dating back at least to Cowles (1933), if the variable of interest is latent then the problem of forecast evaluation and comparison becomes more complicated¹.

This complication can be resolved, at least partly, if an unbiased estimator of the latent variable of interest is available. In volatility forecasting, for example, the squared return on an asset over the period t (assuming a zero mean return) can be interpreted as a conditionally unbiased estimator of the true unobserved conditional variance of the asset over the period t .² Many of the standard methods for forecast evaluation and comparison, such as the Mincer-Zarnowitz (1969) regression and the Diebold and Mariano (1995) and West (1996) tests, can be shown to be applicable when such a conditionally unbiased proxy is used, see Hansen and Lunde (2006) for example. However, it is not true that using a conditionally unbiased proxy will *always* lead to the same outcome as if the true latent variable was used: Andersen and Bollerslev (1998) and Andersen, *et al.* (2005), amongst others, study the reduction in finite-sample power of tests based on noisy volatility proxies; we focus, like Hansen and Lunde (2006), on distortions in the rankings of competing forecasts that can arise when using a noisy volatility proxy in some commonly-used tests for forecast comparison.

For example, in the volatility forecasting literature numerous authors have expressed concern that a few extreme observations may have an unduly large impact on the outcomes of forecast evaluation and comparison tests, see Bollerslev and Ghysels (1994), Andersen, *et al.* (1999) and Poon and Granger (2003) amongst others. One common response to this

¹For recent surveys of the forecast evaluation literature see Corradi and Swanson (2006), Clements (2005), and West (2006). For recent surveys of the volatility forecasting literature, see Andersen, *et al.* (2006), Poon and Granger (2003) and Shephard (2005).

²The high/low range and realised volatility, see Parkinson (1980) and Andersen, *et al.* (2003) for example, have also been used as volatility proxies. These are discussed in detail below.

concern is to employ forecast loss functions that are “less sensitive” to large observations than the usual squared forecast error loss function, such as absolute error or proportional error loss functions. In this paper we show analytically that such approaches can lead to incorrect inferences and the selection of inferior forecasts over better forecasts.

We focus on volatility forecasting as a specific case of the more general problem of latent variable forecasting. In Section 5 we discuss the extension of our results to other latent variable forecasting problems. Our research builds on work by Andersen and Bollerslev (1998), Meddahi (2001) and Hansen and Lunde (2006), who were among the first to analyse the problems introduced by the presence of noise in a volatility proxy. This paper is related to the paper of Hansen and Lunde (2006), and we extend their work in two important directions, discussed below.

This paper makes two main contributions. Firstly, we derive explicit analytical results for the distortions that may arise when some common loss functions are employed, considering the three most commonly-used volatility proxies: the daily squared return, the intra-daily range and a realised variance estimator. We show that these distortions can be large, even for favourable scenarios (such as Gaussianity). Further, we show that the distortions vary greatly with the choice of loss function, thus providing a theoretical explanation for the widespread finding of conflicting rankings of volatility forecasts when “non-robust” loss functions (defined precisely in Section 2) are used in applied work, see Lamoureux and Lastrapes (1993), Hamilton and Susmel (1994), Bollerslev and Ghysels (1994) and Hansen and Lunde (2005), amongst many others³.

Secondly, we provide necessary and sufficient conditions on the functional form of the loss function to ensure that the ranking of various forecasts is preserved when using a noisy volatility proxy. These conditions are related to those of Gouriéroux, *et al.* (1984) for quasi-maximum likelihood estimation. Interestingly, we find that there are an infinite number of

³All of the results in this paper apply directly to the problem of forecasting integrated variance (IV), which Andersen, *et al.* (2002), amongst others, argue is a more “relevant” notion of variability. We focus on the problem of conditional variance forecasting due to its prevalence in applied work in the past two decades. If we take expected IV rather than the conditional variance as the latent object of interest, then we only require that an unbiased realised variance estimator is available for the results to go through. In the presence of jumps in the price process, quadratic variation (QV) is a more appropriate measure of risk, and a similar extension is possible.

loss functions that satisfy these conditions, and that these loss functions differ in meaningful ways (such as the penalty applied to overprediction versus underprediction). Thus our class of “robust” loss functions is not simply the quadratic loss function or minor variations thereof.

The canonical problem in point forecasting is to find the forecast that minimises the expected loss, conditional on time t information. That is,

$$\hat{Y}_{t+h,t}^* \equiv \arg \min_{\hat{y} \in \mathcal{Y}} E [L(Y_{t+h}, \hat{y}) | \mathcal{F}_t] \quad (1)$$

where Y_{t+h} is the variable of interest, L is the forecast user’s loss function, \mathcal{Y} is the set of possible forecasts, and \mathcal{F}_t is the time t information set. Starting with the assumption that the forecast user is interested in the conditional variance, we effectively take the *solution* of the optimisation problem above (the conditional variance) as given, and consider the loss functions that will generate the desired solution. This approach is unusual in the economic forecasting literature: the more common approach is to take the forecast user’s loss function as given and derive the optimal forecast for that loss function; related papers here are Granger (1969), Engle (1993), Christoffersen and Diebold (1997), Christoffersen and Jacobs (2004) and Patton and Timmermann (2004), amongst others. The fact that we know the forecast user desires a variance forecast places limits on the class of loss functions that may be used for volatility comparison, ruling out some choices previously used in the literature. However we show that the class of “robust” loss functions still admits a wide variety of loss functions, allowing much flexibility in representing volatility forecast users’ preferences.

One of the main practical implications of this paper is that the stated goal of forecasting the conditional variance is not consistent with the use of some loss functions when an imperfect volatility proxy is employed. However, these loss functions are *not* inherently invalid or inappropriate: if the forecast user’s preferences are indeed described by an “non-robust” loss function, then this simply implies that the object of interest to that forecast user is not the conditional variance but rather some other quantity⁴. In academic research the preferences of the end-user of the forecast are often unknown, and a common response to this is to select forecasts based on their average distance, somehow measured, to the true latent conditional

⁴For example, the utility of realised returns on a portfolio formed using a volatility forecast, or the profits obtained from an option trading strategy based on a volatility forecast, see West, *et al.* (1993) and Engle, *et al.* (1993) for example, define economically meaningful loss functions, even though the optimal forecasts under those loss functions will not generally be the true conditional variance.

variance. In such cases, the methods outlined in this paper can be applied to identify the forecast that is closest to the true conditional variance by using imperfect volatility proxy and a “robust” loss function.

The remainder of this paper is as follows. In Section 2 we analytically consider volatility forecast comparison tests using an imperfect volatility proxy, showing the problems that arise when using some common loss functions. We initially consider using squared daily returns as the proxy, and then consider using the range and realised variance. In Section 3 we provide necessary and sufficient conditions on the functional form of a loss function for the ranking of competing volatility forecasts to be robust to the presence of noise in the volatility proxy, and derive some useful special cases of this class of robust loss functions. One of these special cases is a parametric family of loss functions that nests two of the most widely-used loss functions in the literature, namely the MSE and QLIKE loss functions (defined in equations (5) and (6) below). In Section 4 we present an illustration using two widely-used volatility models, and in Section 5 we conclude and suggest extensions. All proofs and derivations are provided in appendices.

1.1 Notation

Let r_t be the variable whose conditional variance is of interest, usually a daily or monthly asset return in the volatility forecasting literature. The information set used in defining the conditional variance of interest is denoted \mathcal{F}_{t-1} , which is assumed to contain $\sigma(r_{t-j}, j \geq 1)$, but may also include other variables and/or variables measured at a higher frequency than r_t (such as intra-daily returns). Denote $V[r_t|\mathcal{F}_{t-1}] \equiv V_{t-1}[r_t] \equiv \sigma_t^2$. We will assume throughout that $E[r_t|\mathcal{F}_{t-1}] \equiv E_{t-1}[r_t] = 0$, and so $\sigma_t^2 = E_{t-1}[r_t^2]$. Let $\varepsilon_t \equiv r_t/\sigma_t$ denote the ‘standardised return’. Let a forecast of the conditional variance of r_t be denoted h_t , or $h_{i,t}$ if there is more than one forecast under analysis. We will take forecasts as “primitive”, and not consider the specific models and estimators that may have generated the forecasts. The loss function of the forecast user is $L : \mathbb{R}_+ \times \mathcal{H} \rightarrow \mathbb{R}_+$, where the first argument of L is σ_t^2 or some proxy for σ_t^2 , denoted $\hat{\sigma}_t^2$, and the second is h_t . \mathbb{R}_+ and \mathbb{R}_{++} denote the non-negative and positive parts of the real line respectively, and \mathcal{H} is a compact subset of \mathbb{R}_{++} . Commonly used volatility proxies are the squared return, r_t^2 , realised volatility, RV_t , and the range, RG_t .

Optimal forecasts for a given loss function and proxy are denoted h_t^* and are defined as:

$$h_t^* \equiv \arg \min_{h \in \mathcal{H}} E [L (\hat{\sigma}_t^2, h) | \mathcal{F}_{t-1}] \quad (2)$$

2 Volatility forecast comparison using an imperfect volatility proxy

We consider volatility forecast comparisons based on expected loss, or distance to the true conditional variance. These comparisons can be implemented in finite samples using the tests of Diebold and Mariano (1995) and West (1996), (henceforth DMW). If we define $u_{i,t} \equiv L (\sigma_t^2, h_{i,t})$, where L is the forecast user's loss function, and let $d_t = u_{1,t} - u_{2,t}$, then a DMW test of equal predictive accuracy can be conducted as a simple Wald test that $E [d_t] = 0$.⁵

Of primary interest is whether the feasible ranking of two forecasts obtained using an imperfect volatility proxy is the same as the infeasible ranking that would be obtained using the unobservable true conditional variance. In such a case we are able to measure average forecast accuracy even though the variable of interest is unobservable. We define loss functions that yield such an equivalence as “robust”:

Definition 1 *A loss function, L , is “robust” if the ranking of any two (possibly imperfect) volatility forecasts, h_{1t} and h_{2t} , by expected loss is the same whether the ranking is done using the true conditional variance, σ_t^2 , or some conditionally unbiased volatility proxy, $\hat{\sigma}_t^2$. That is,*

$$E [L (\sigma_t^2, h_{1t})] \gtrless E [L (\sigma_t^2, h_{2t})] \Leftrightarrow E [L (\hat{\sigma}_t^2, h_{1t})] \gtrless E [L (\hat{\sigma}_t^2, h_{2t})] \quad (3)$$

for any $\hat{\sigma}_t^2$ s.t. $E [\hat{\sigma}_t^2 | \mathcal{F}_{t-1}] = \sigma_t^2$.

⁵The key difference between the approaches of Diebold and Mariano (1995) and West (1996) is that the latter explicitly allows for forecasts that are based on estimated parameters, whereas the null of equal predictive accuracy is based on population parameters, see West (2005). The problems we identify below arise even in the absence of estimation error in the forecasts, thus our treatment of the forecasts as primitive, and so for our purposes these two approaches coincide.

Meddahi (2001) showed that the ranking of forecasts on the basis of the R^2 from the Mincer-Zarnowitz regression:

$$\hat{\sigma}_t^2 = \beta_0 + \beta_1 h_{it} + e_{it} \quad (4)$$

is robust to noise in $\hat{\sigma}_t^2$. Hansen and Lunde (2006) showed that the R^2 from a regression of $\log(\hat{\sigma}_t^2)$ on a constant and $\log(h_t)$ is not robust to noise, and showed more generally that a sufficient condition for a loss function to be robust is that $\partial^2 L(\sigma^2, h) / \partial (\sigma^2)^2$ does not depend on h . In Section 3 we generalise this result by providing necessary and sufficient conditions for a loss function to be robust.^{6,7}

It is worth noting that although the ranking obtained from a robust loss function will be invariant to noise in the proxy, the actual *level* of expected loss obtained using a proxy will be larger than that which would be obtained when using the true conditional variance. This point was compellingly presented in Andersen and Bollerslev (1998) and Andersen, *et al.* (2004). Andersen, *et al.* (2005) provide a method to estimate the distortion in the level of expected loss and thereby obtain an estimator of the level of expected loss that would be obtained using the true latent variable of interest.

It follows directly from the definition of a robust loss function that the true conditional variance is the optimal forecast (we formally show this in the proof of Proposition 2), and thus a necessary condition for a loss function to be robust to noise is that the true conditional variance is the optimal forecast. In this section we determine whether this condition holds for some common loss functions, and analytically characterise the distortion for those cases where it is violated.

A common response to the concern that a few extreme observations drive the results of volatility forecast comparison studies is to employ alternative measures of forecast accuracy to the usual MSE loss function, see Pagan and Schwert (1990), Bollerslev and Ghysels (1994),

⁶Our use of the adjective “robust” is related, though not equivalent, to its use in estimation theory, where it applies to estimators that insensitive/less sensitive to the presence of outliers in the data, see Huber (1981) for example. A “robust” loss function, in the sense of Definition 1, will generally not be robust to the presence of outliers.

⁷In recent work Giacomini and White (2006) propose ranking forecasts by expected loss conditional on some information set \mathcal{G}_t , rather than by unconditional expected loss as in Definition 1. The numerical examples provided below will differ in this more general case, of course, however the theoretical results in this paper go through if $\mathcal{G}_t \subseteq \mathcal{F}_{t-1}$, which is true for all of the examples considered by Giacomini and White (2006).

Bollerslev, *et al.* (1994), Diebold and Lopez (1996), Andersen, *et al.* (1999), Poon and Granger (2003) and Hansen and Lunde (2005), for example. A collection of loss functions employed in the literature on volatility forecast evaluation is presented below⁸. In the next two sections we will study the properties of these loss functions when combined with common volatility proxies.

$$MSE : L(\hat{\sigma}^2, h) = (\hat{\sigma}^2 - h)^2 \quad (5)$$

$$QLIKE : L(\hat{\sigma}^2, h) = \log h + \frac{\hat{\sigma}^2}{h} \quad (6)$$

$$MSE-LOG : L(\hat{\sigma}^2, h) = (\log \hat{\sigma}^2 - \log h)^2 \quad (7)$$

$$MSE-SD : L(\hat{\sigma}^2, h) = (\hat{\sigma} - \sqrt{h})^2 \quad (8)$$

$$MSE-prop : L(\hat{\sigma}^2, h) = \left(\frac{\hat{\sigma}^2}{h} - 1\right)^2 \quad (9)$$

$$MAE : L(\hat{\sigma}^2, h) = |\hat{\sigma}^2 - h| \quad (10)$$

$$MAE-LOG : L(\hat{\sigma}^2, h) = |\log \hat{\sigma}^2 - \log h| \quad (11)$$

$$MAE-SD : L(\hat{\sigma}^2, h) = |\hat{\sigma} - \sqrt{h}| \quad (12)$$

$$MAE-prop : L(\hat{\sigma}^2, h) = \left|\frac{\hat{\sigma}^2}{h} - 1\right| \quad (13)$$

2.1 Using squared returns as a volatility proxy

In this section we will focus on the use of daily squared returns for volatility forecast evaluation, and in Section 2.2 we will examine the use of realised volatility and the range. We will derive our results under three assumptions for the conditional distribution of daily returns:

$$r_t | \mathcal{F}_{t-1} \sim \begin{cases} F_t(0, \sigma_t^2) \\ \text{Student's } t(0, \sigma_t^2, \nu) \\ N(0, \sigma_t^2) \end{cases}$$

where $F_t(0, \sigma_t^2)$ is some unspecified distribution with mean zero and variance σ_t^2 , and *Student's* $t(0, \sigma_t^2, \nu)$ is a Student's t distribution with mean zero, variance σ_t^2 and ν degrees of freedom. In all cases it is clear that $E_{t-1}[r_t^2] = \sigma_t^2$, and so the squared daily return is a valid volatility proxy.

⁸Some of these loss functions are called different names by different authors: MSE-prop is also known as “heteroskedasticity-adjusted MSE (HMSE)”; MAE-prop is also known as “mean absolute percentage error (MAPE)” or as “heteroskedasticity-adjusted MAE (HMAE)”.

It is trivial to show that the MSE loss function generates an optimal forecast equal to the conditional variance: $h_t^* = E_{t-1} [r_t^2] = \sigma_t^2$, and thus satisfies the necessary condition for robustness. Further, the MSE loss function also satisfies the sufficient condition of Hansen and Lunde (2006), and thus MSE is a “robust” loss function. Another commonly used loss function is the MSE loss function on standard deviations rather than variances, MSE-SD, see equation (8). The motivation for this loss function is that taking square root of the two arguments of the squared-error loss function shrinks the larger values towards zero, reducing the impact of the most extreme values of r_t . However it also leads to an incorrect volatility forecast being selected as optimal:

$$\begin{aligned}
h_t^* &\equiv \arg \min_{h \in \mathcal{H}} E_{t-1} \left[\left(|r_t| - \sqrt{h} \right)^2 \right] \\
\text{FOC } 0 &= \left. \frac{\partial}{\partial h} E_{t-1} \left[\left(|r_t| - \sqrt{h} \right)^2 \right] \right|_{h=h_t^*} \\
\text{so } h_t^* &= (E_{t-1} [|r_t|])^2 & (14) \\
&= \sigma_t^2 (E_{t-1} [|\varepsilon_t|])^2 \\
&= \begin{cases} \frac{\nu-2}{\pi} \left(\Gamma \left(\frac{\nu-1}{2} \right) / \Gamma \left(\frac{\nu}{2} \right) \right)^2 \sigma_t^2, & \text{if } r_t | \mathcal{F}_{t-1} \sim \text{Student's } t(0, \sigma_t^2, \nu), \nu > 2 \\ \frac{2}{\pi} \sigma_t^2 \approx 0.64 \sigma_t^2, & \text{if } r_t | \mathcal{F}_{t-1} \sim N(0, \sigma_t^2) \end{cases} & (15)
\end{aligned}$$

Excess kurtosis in asset returns exacerbates the distortion, which we can see in Figure 1 for returns that have the Student’s t distribution.

As mentioned in the Introduction, if the forecast user’s loss function truly is the square of the difference between the *absolute return* and the square root of the forecast, then the “distortion” in the optimal forecast above is desirable, as this is the forecast that minimises his/her expected loss. However, if the goal is to find the forecast that is closest to the *true conditional variance*, then this distortion in the optimal forecast can lead to an incorrect ranking of competing forecasts⁹. Thus the MSE-SD loss function is not consistent with the goal of ranking volatility forecast by their distance to the true conditional variance when using the squared return as the volatility proxy: either the proxy has to be re-scaled by a quantity that depends critically on the underlying conditional distribution of returns, or, more simply, a different loss function must be chosen.

⁹This distortion remains if the target is instead the conditional standard deviation, as the absolute return is not an unbiased proxy for that quantity. This is discussed in further detail in Section 2.3.

We provide the corresponding calculations for the remaining loss functions in equations (5) to (13) in the technical appendix, and summarise the results in Table 1. This table shows that the degree of distortion in the optimal forecast according to some of the loss functions used in the literature can be substantial. Under normality the optimal forecast under these loss functions ranges from about one quarter of the true conditional variance to three times the true conditional variance. If returns exhibit excess conditional kurtosis then the range of optimal forecasts from these loss functions is even wider.

Table 1 provides a theoretical explanation for the widespread finding of conflicting rankings of volatility forecasts when non-robust loss functions are used in applied work. Lamoureux and Lastrapes (1993), Hamilton and Susmel (1994), Bollerslev and Ghysels (1994) and Hansen and Lunde (2005), amongst many others, use some or all of the nine loss functions considered in Table 1 and find that the best-performing volatility model changes with the choice of loss function. Given that, for example, the MSE-prop loss function leads to an optimal forecast that is biased *upwards* by at least a factor of three, while the MAE loss function leads to an optimal forecast that is biased *downwards* by at least a factor of two, it is no surprise that different rankings of volatility forecasts are found.

2.2 Using better volatility proxies

It has long been known that squared returns are a rather noisy proxy for the true conditional variance. One alternative volatility proxy that has gained much attention recently is “realised volatility”, see Andersen, *et al.* (2001, 2003), and Barndorff-Nielsen and Shephard (2002, 2004). Another commonly-used alternative to squared returns is the intra-daily range. It is well-known that if the log stock price follows a Brownian motion then both of these estimators are unbiased and more efficient than the squared return. In this section we obtain the rate at which the distortion in the ranking of alternative forecasts disappears when using realised volatility as the proxy, as the sampling frequency increases, for a simple data generating process (DGP).

Assume that there are m equally-spaced observations per trade day, and let $r_{i,m,t}$ denote the i^{th} intra-daily return on day t . While recent work on realised volatility would enable us to consider a quite general class of DGPs, in order to obtain analytical results for problems involving the *range* as a volatility proxy we consider only a simple DGP: zero mean return,

no jumps, and constant conditional volatility within a trade day¹⁰. Patton and Sheppard (2007) present the corresponding results for a range of more realistic DGPs via simulation¹¹. Let

$$r_t = d \log P_t = \sigma_t dW_t \quad (16)$$

$$\sigma_\tau = \sigma_t \forall \tau \in (t-1, t] \quad (17)$$

$$r_{i,m,t} \equiv \int_{(i-1)/m}^{i/m} r_\tau d\tau = \sigma_t \int_{(i-1)/m}^{i/m} dW_\tau \quad (18)$$

$$\text{so } \{r_{i,m,t}\}_{i=1}^m \sim iid N\left(0, \frac{\sigma_t^2}{m}\right) \quad (19)$$

The “realised volatility” or “realised variance” is defined as:

$$RV_t^{(m)} \equiv \sum_{i=1}^m r_{i,m,t}^2$$

Realised variance, like the daily squared return (which is obtained in the above framework by setting $m = 1$), is a conditionally unbiased estimator of the daily conditional variance. Its main advantage is that it is more efficient estimator than the daily squared return: for this DGP it can be shown that $MSE_{t-1} [r_t^2] = 2\sigma_t^4$ while $MSE_{t-1} [RV_t^{(m)}] = 2\sigma_t^4/m$. Thus $RV_t^{(m)} \xrightarrow{p} \sigma_t^2$ as $m \rightarrow \infty$, under these assumptions, and we find in this case that σ_t^2 is observable. As expected, all distortions vanish in this case.

The range, or the high/low, estimator has been used in finance for many years, see Parkinson (1980), Garman and Klass (1980) and Ball and Torous (1984). The intra-daily log range is defined as:

$$RG_t \equiv \max_{\tau} \log P_\tau - \min_{\tau} \log P_\tau, \quad t-1 < \tau \leq t \quad (20)$$

¹⁰ Analytical and empirical results on the range and “realised range” under more flexible DGPs are presented in two recent papers by Christensen and Podolskij (2006) and Martens and van Dijk (2007).

¹¹ When the DGP is specified to be log-normal or GARCH stochastic volatility diffusions, Patton and Sheppard (2007) find results very similar to those obtained for the case below. Using the same parameterisations as those in the simulations of Gonçalves and Meddahi (2005), slightly larger biases from the non-robust loss functions are found, but they generally differ from those in Table 2 only in the second decimal place. In contrast, the biases are found to be much larger under the two-factor stochastic volatility diffusion considered by Gonçalves and Meddahi (2005).

Under the dynamics in equation (16) Feller (1951) presented the density of RG_t , and Parkinson (1980) presented a formula for obtaining moments of the range, which enable us to compute:

$$E_{t-1} [RG_t^2] = 4 \log(2) \cdot \sigma_t^2 \approx 2.7726\sigma_t^2 \quad (21)$$

Details on the distributional properties of the range under this DGP are presented in the technical appendix. The above expression shows that squared range is *not* a conditionally unbiased estimator of σ_t^2 ; we will thus focus below on the *adjusted range*:

$$RG_t^* \equiv \frac{RG_t}{2\sqrt{\log(2)}} \approx 0.6006RG_t \quad (22)$$

which, when squared, is an unbiased proxy for the conditional variance. Note that the adjustment factor depends critically on the assumed DGP, which is a potential drawback of the range as a volatility proxy. Using the results of Parkinson (1980) it is simple to determine that $MSE_{t-1} [RG_t^{*2}] \approx 0.4073\sigma_t^4$, which is approximately one-fifth of the MSE of the daily squared return.

We now determine the optimal forecasts obtained using the various loss functions considered above, when $\hat{\sigma}_t^2 = RV_t^{(m)}$ or $\hat{\sigma}_t^2 = RG_t^{*2}$ is used as a proxy for the conditional variance rather than r_t^2 . We initially leave m unspecified for the realised volatility proxy, and then specialise to three cases: $m = 1, 13$ and 78 , corresponding to the use of daily, half-hourly and 5-minute returns, on a stock listed on the New York Stock Exchange (NYSE).

For MSE and QLIKE the optimal forecast is simply the conditional mean of $\hat{\sigma}_t^2$, which equals the conditional variance, as $RV_t^{(m)}$ and RG_t^{*2} are both conditionally unbiased. The MSE-SD loss function yields $(E_{t-1} [\hat{\sigma}_t])^2$ as an optimal forecast. Under the set-up above,

$$\begin{aligned} RV_t^{(m)} &\equiv \sum_{i=1}^m r_{t,i}^2 = \frac{\sigma_t^2}{m} \sum_{i=1}^m \varepsilon_{t,i}^2 \\ \text{so } m\sigma_t^{-2}RV_t^{(m)} &\sim \chi_m^2 \\ \text{so } h_t^* &= \frac{\sigma_t^2}{m} \left(E \left[\sqrt{\chi_m^2} \right] \right)^2 \\ E \left[\sqrt{\chi_m^2} \right] &\approx \sqrt{m} - \frac{1}{4\sqrt{m}} \text{ by a Taylor series approximation} \\ \text{so } h_t^* &\approx \sigma_t^2 \left(1 - \frac{1}{2m} + \frac{1}{16m^2} \right) \end{aligned}$$

$$\approx \begin{cases} 0.5625 \cdot \sigma_t^2 & \text{for } m = 1 \\ 0.9619 \cdot \sigma_t^2 & \text{for } m = 13 \\ 0.9936 \cdot \sigma_t^2 & \text{for } m = 78 \end{cases}$$

The results for the MSE-SD loss function using realised volatility show that reducing the noise in the volatility proxy improves the optimal forecast, consistent with Hansen and Lunde (2006).¹² Using the range we find that

$$h_t^* = (E_{t-1} [RG_t^*])^2 = \frac{2}{\pi \log 2} \sigma_t^2 \approx 0.9184 \sigma_t^2$$

and so the distortion from using the range is approximately equal to that incurred when using a realised volatility constructed using 6 intra-daily observations. Calculations for the remaining loss functions are collected in the technical appendix, and the results are summarised in Table 2.

The results in Table 2 confirm that as the proxy used to measure the true conditional variance gets more efficient the degree of distortion decreases for all loss functions. Using half-hour returns (13 intra-daily observations) or the intra-daily range still leaves substantial distortions in the optimal forecasts, but using 5-minute returns (78 intra-daily observations) eliminates almost all of the bias, at least in this simple framework. While high frequency data is available and reliable for some assets (the most liquid assets on well-developed exchanges), for most assets it is not possible to obtain reliable high-frequency data, and thus the impact of noise in the volatility proxy cannot be ignored.

2.3 General comments on non-robust loss functions

The sources of the mismatches between the optimal forecast for a given loss function and the true conditional variance are easily identified. The MAE, MAE-SD and MAE-prop loss functions consider mean absolute distances rather than mean squared distances, which then naturally change the solution of the optimisation problem from an expectation to a median. For the MSE-log, MSE-SD and MSE-prop loss functions the distortion follows from the fact that the unbiasedness property is not invariant to nonlinear transformations.

¹²Note that the result for $m = 1$ is different to that obtained in Section 2, which was $h_t^* = \frac{2}{\pi} \sigma_t^2 \approx 0.6366 \sigma_t^2$. This is because for $m = 1$ we can obtain the expression exactly, using results for the normal distribution, whereas for arbitrary m we relied on a second-order Taylor series approximation.

In all of these cases the distortions can be remedied if one can obtain a conditionally unbiased estimator of the alternative quantity of interest (σ_t , $\log \sigma_t^2$, etc.) either exactly or approximately. When using the squared return as a proxy, this will generally require an assumption about the entire conditional distribution of returns. When using realised variance as a volatility proxy, one may obtain an approximate distribution of the volatility proxy under relatively mild assumptions, by drawing on the distribution theory for realised volatility developed in Barndorff-Nielsen and Shephard (2004) and extensions, as in Andersen, *et al.* (2005). On the other hand, when using a robust loss function only the assumption of conditional unbiasedness of the proxy is required, which is often satisfied under much weaker assumptions and requires no adjustment of the proxy.

We now seek to identify the reason why some non-robust loss function yield upward-biased forecasts, whilst others yield downward-biased forecasts. We do so by generalising the results from the previous sections to a broad class of arbitrary loss functions, making use of a mean-value expansion. This requires some differentiability assumptions on the loss function, which are not satisfied for some of the loss functions considered above.

G1: The volatility proxy is conditionally unbiased: $E_{t-1} [\hat{\sigma}_{t,m}^2] = \sigma_t^2$

G2: The loss function L is three times differentiable.

G3: The loss function L is such that $L(\sigma^2, h) = 0$ iff $h = \sigma^2$

G4: The loss function L is such that $\partial L(\sigma^2, h) / \partial h \begin{smallmatrix} \geq \\ \leq \end{smallmatrix} 0$ if $\sigma^2 \begin{smallmatrix} \leq \\ \geq \end{smallmatrix} h$

Proposition 1 *Let assumptions G1-G4 hold, and define*

$$h_{t,m}^* \equiv \arg \min_{h \in \mathcal{H}} E_{t-1} [L(\hat{\sigma}_{t,m}^2, h)]$$

then

$$\frac{\partial^3 L(\sigma^2, h)}{\partial (\sigma^2)^2 \partial h} \begin{smallmatrix} \geq \\ \leq \end{smallmatrix} 0 \text{ for all } (\sigma^2, h) \Rightarrow h_{t,m}^* \begin{smallmatrix} \leq \\ \geq \end{smallmatrix} \sigma_t^2$$

The above proposition shows that it is the sign of the third derivative of the loss function that determines whether the optimal forecast is above, below or equal to the true conditional variance. The case that this third derivative is equal to zero, and thus that the optimal forecast is the conditional variance, corresponds to a result of Hansen and Lunde (2006). The third derivative is always positive for the MSE-log and MSE-SD loss functions, and so part (i) above implies that $h_{t,m}^* < \sigma_t^2$ for these loss functions, which is consistent with the

results in Tables 1 and 2. Alternatively, for the MSE-prop loss function this third derivative is always negative, implying that $h_{t,m}^* > \sigma_t^2$ for this loss function, which is again consistent with Tables 1 and 2.

3 A class of robust loss functions

In the previous section we showed that amongst nine loss functions commonly used to compare volatility forecasts, only the MSE and the QLIKE loss functions lead to $h_t^* = E_{t-1} [\hat{\sigma}_t^2] = \sigma_t^2$, which is a necessary condition for a loss function to be robust to noise in the volatility proxy. The following proposition is the main theoretical contribution of the paper; it provides a necessary and sufficient class of robust loss functions, which are related to the class of linear-exponential densities of Gouriéroux, *et al.* (1984), and to the work of Gouriéroux, *et al.* (1987). We will show below that this class contains an infinite number of loss functions, and allows for asymmetric penalties to be applied to over- versus under-predictions, as well as for a symmetric penalty. We make the following assumptions:

A1: $E_{t-1} [\hat{\sigma}_t^2] = \sigma_t^2$ for all t

A2: $\hat{\sigma}_t^2 | \mathcal{F}_{t-1} \sim F_t \in \tilde{F}$, the set of all absolutely continuous distribution functions on \mathbb{R}_+ .

A3: L is twice continuously differentiable with respect to h and $\hat{\sigma}^2$, and has a unique minimum at $\hat{\sigma}^2 = h$.

A4: There exists some $h_t^* \in \text{int}(\mathcal{H})$ such that $h_t^* = E_{t-1} [\hat{\sigma}_t^2]$, where \mathcal{H} is a compact subset of \mathbb{R}_{++} .

A5: L and F_t are such that: (a) $E_{t-1} [L(\hat{\sigma}_t^2, h)] < \infty$ for some $h \in \mathcal{H}$; (b) $\left| E_{t-1} \left[\partial L(\hat{\sigma}_t^2, h) / \partial h \Big|_{h=\sigma_t^2} \right] \right| < \infty$; and (c) $\left| E_{t-1} \left[\partial^2 L(\hat{\sigma}_t^2, h) / \partial h^2 \Big|_{h=\sigma_t^2} \right] \right| < \infty$, for all t .

Proposition 2 *Let assumptions A1 to A5 hold. Then a loss function L is robust, in the sense of Definition 1, if and only if it takes the following form:*

$$L(\hat{\sigma}^2, h) = \tilde{C}(h) + B(\hat{\sigma}^2) + C(h)(\hat{\sigma}^2 - h) \quad (23)$$

where B and C are twice continuously differentiable, C is a strictly decreasing function on \mathcal{H} , and \tilde{C} is the anti-derivative of C .

Remark 1 *If we normalise the loss function to yield zero loss when $\hat{\sigma}^2 = h$, then $B(\hat{\sigma}^2) = -\tilde{C}(\hat{\sigma}^2)$.*

Given the widespread interest in economics and finance in loss functions that depend only on the forecast error or the standardised forecast error, we present below a surprising result on the subset of robust loss functions that satisfy one of these restrictions.

Proposition 3 *(i) The “MSE” loss function is the only robust loss function satisfying assumptions A1-A5 that depends solely on the forecast error, $\hat{\sigma}^2 - h$.*

(ii) The “QLIKE” loss function is the only robust loss function satisfying assumptions A1-A5 that depends solely on the standardised forecast error, $\hat{\sigma}^2/h$.

Given the arbitrariness of the choice of units in most economic and financial problems (for example, measuring prices in dollars versus cents, or measuring returns in percentages versus decimals) it is potentially interesting to consider the impact of a simple change in units on the ranking of two competing forecasts by expected loss. The class of loss functions presented in Proposition 2 guarantees that the true conditional variance will be chosen (subject to sampling variation) over any other forecast regardless of the choice units. However it does not guarantee that the ranking of two *imperfect* forecasts will be invariant to the choice of units. The following proposition shows that by using a *homogeneous* robust loss function, the ranking of any two (possibly imperfect) forecasts is invariant to a re-scaling of the data. It further provides an example where the ranking can be reversed simply with a re-scaling of the data if a non-homogeneous robust loss function is used.

Proposition 4 *Recall that a loss function L is homogeneous of order k if*

$$L(a\hat{\sigma}^2, ah) = a^k L(\hat{\sigma}^2, h) \quad \forall a > 0 \quad \text{for some } k$$

Then:

(i) The ranking of any two (possibly imperfect) volatility forecasts by expected loss is invariant to a re-scaling of the data if the loss function is homogeneous.

(ii) The ranking of any two (possibly imperfect) volatility forecasts by expected loss may not be invariant to a re-scaling of the data if the loss function is robust but not homogeneous.

With the above motivation for homogeneous loss functions, we now derive the subset of homogeneous, robust loss functions. It turns out that this subset of functions is indexed by a single parameter, which determines the both degree of homogeneity and the shape of the loss function. Naturally, the MSE loss function is nested in this case (homogeneous of order 2), as is the QLIKE loss function (homogeneous of order zero).

Proposition 5 *The following family of loss functions, indexed by the scalar parameter b , corresponds to the entire subset of robust and homogeneous loss functions. The degree of homogeneity is equal to $b + 2$.*

$$L(\hat{\sigma}^2, h; b) = \begin{cases} \frac{1}{(b+1)(b+2)}(\hat{\sigma}^{2b+4} - h^{b+2}) - \frac{1}{b+1}h^{b+1}(\hat{\sigma}^2 - h), & \text{for } b \notin \{-1, -2\} \\ h - \hat{\sigma}^2 + \hat{\sigma}^2 \log \frac{\hat{\sigma}^2}{h}, & \text{for } b = -1 \\ \frac{\hat{\sigma}^2}{h} - \log \frac{\hat{\sigma}^2}{h} - 1, & \text{for } b = -2 \end{cases} \quad (24)$$

The MSE loss function is obtained when $b = 0$ and the QLIKE loss function is obtained when $b = -2$, up to additive and multiplicative constants. In Figure 2 we present the above class of functions for various values of b , ranging from 1 to -5 , and including the MSE and QLIKE cases. This figure shows that this family of loss functions can take a wide variety of shapes, ranging from symmetric ($b = 0$, corresponding to the MSE loss function) to asymmetric, with heavier penalty either on under-prediction ($b < 0$) or over-prediction ($b > 0$). Figure 3 plots the ratio of losses incurred for negative forecast errors to those incurred for positive forecast errors, to make clearer the form of asymmetries in these loss functions.

4 Empirical application to forecasting IBM return volatility

In this section we consider the problem of forecasting the conditional variance of the daily return on IBM, using data from the TAQ database over the period from January 1993 to December 2003. We consider two simple volatility models that are widely-used in industry: a 60-day rolling window estimator, and the RiskMetrics volatility model based on daily returns:

$$\text{Rolling window} : h_{1t} = \frac{1}{60} \sum_{j=1}^{60} r_{t-j}^2 \quad (25)$$

$$\text{RiskMetrics} : h_{2t} = \lambda h_{2t-1} + (1 - \lambda) r_{t-1}^2, \lambda = 0.94 \quad (26)$$

We use approximately the first year of observations (272 observations) to initiate the RiskMetrics forecasts, and the remaining 2500 observations to compare the forecasts. A plot of the volatility forecasts is provided in Figure 4.

We employ a variety of volatility proxies in the comparison of these forecasts: the daily squared return, and realised variance computed using 65-minute, 15-minute and 5-minute returns¹³. In comparing these forecasts we present the results of Diebold-Mariano-West tests using the loss function presented in Proposition 5, for five different choices of the loss function parameter: $b = \{1, 0, -1, -2, -5\}$. MSE loss and QLIKE loss correspond to $b = 0$ and $b = -2$ respectively¹⁴.

In Table 3 we present tests comparing the RiskMetrics forecasts based on daily returns with the 60-day rolling window volatility forecasts. The only loss function for which the difference in forecast performance is significantly different from zero is the QLIKE loss function: the difference is significant at the 0.05 level using 65-minute, 15-minute and 5-minute realised variances as the volatility proxy, and significant at the 0.10 level using daily squared returns as the proxy. In all of these cases the t-statistic is positive, indicating that the rolling window forecasts generated larger average loss than the RiskMetrics forecasts.

Interestingly, under MSE loss, the differences in average loss favour the rolling window forecasts, though these differences are not statistically significant. Mincer-Zarnowitz tests (available upon request) revealed, unsurprisingly, that neither of these forecasts is optimal. Robust loss functions are designed to always select the true conditional variance over any competing forecast, but when comparing two *imperfect* forecasts the ranking can, as in this example, change depending on the choice of loss function. This emphasises the flexibility that remains even when we restrict attention to homogeneous, robust loss functions.

5 Conclusion

We analytically demonstrated some problems with volatility forecast comparison techniques used in the literature. These techniques invariably rely on a volatility proxy, which is some

¹³We use 65-minute returns rather than 60-minute returns so that there are an even number of intervals within the NYSE trade day, which runs from 9.30am to 4pm.

¹⁴Moment and memory conditions required for the DMW test to be implemented are detailed in an earlier version of this paper, Patton (2006).

imperfect estimator of the true conditional variance, and the presence of noise in the volatility proxy can lead an imperfect volatility forecast being selected over the true conditional variance for certain choices of loss function. Thus noisy volatility proxies not only reduce power, as discussed in Andersen and Bollerslev (1998) for example, they can also seriously affect the asymptotic size of commonly-used tests. We showed analytically that less noisy volatility proxies, such as the intra-daily range and realised volatility, lead to less distortion, though in many cases the degree of distortion is still large.

We derived necessary and sufficient conditions for the loss function to yield rankings of volatility forecasts that are robust to noise in the proxy. We also proposed a new parametric family of robust and homogeneous loss functions, which yield inference that is invariant to the choice of units of measurement. The new family of loss function nests both squared-error and the “QLIKE” loss functions, two of the most widely-used in the volatility forecasting literature. A small empirical study of IBM equity volatility illustrated the new loss functions in forecast comparison tests.

Whilst volatility forecasting is a prominent example of a problem in economics where the variable of interest is unobserved, there are many other such examples: forecasting the true rate of GDP growth (not simply the announced rate); forecasting default probabilities; and forecasting covariances or correlations. The derivations in this paper exploited the fact that the latent variable of interest in volatility forecasting (namely the conditional variance) is a positive random variable, and the proxy is non-negative and continuously distributed. Extending the results in this paper to handle latent variables of interest with support on the entire real line, as would be required for applications to studies of the “true” rates of growth in macroeconomic aggregates or to conditional covariances, should not be difficult. Extending our results to handle proxies with discrete support, such as those that would be used in default forecasting applications, may require a different method of proof. We leave such extensions to future research.

6 Appendix: Proofs of Propositions

Proof of Proposition 1. Use a second-order mean-value expansion of L :

$$L(\hat{\sigma}_{t,m}^2, h) = L(\sigma_t^2, h) + \frac{\partial L(\sigma_t^2, h)}{\partial \sigma_t^2} (\hat{\sigma}_{t,m}^2 - \sigma_t^2) + \frac{1}{2} \frac{\partial^2 L(\hat{\sigma}_{t,m}^2, h)}{\partial (\sigma_t^2)^2} (\hat{\sigma}_{t,m}^2 - \sigma_t^2)^2$$

$$\text{where } \hat{\sigma}_{t,m}^2 = \lambda_{t,m} \hat{\sigma}_{t,m}^2 + (1 - \lambda_{t,m}) \sigma_t^2, \quad \lambda_{t,m} \in [0, 1]$$

$$\text{so } E_{t-1}[L(\hat{\sigma}_{t,m}^2, h)] = L(\sigma_t^2, h) + \frac{1}{2m} E_{t-1} \left[\frac{\partial^2 L(\hat{\sigma}_{t,m}^2, h)}{\partial (\sigma_t^2)^2} (\hat{\sigma}_{t,m}^2 - \sigma_t^2)^2 \right]$$

by assumption G1. The first-order condition for forecast optimality is

$$\begin{aligned} 0 &= E_{t-1} \left[\frac{\partial L(\hat{\sigma}_{t,m}^2, h_{t,m}^*)}{\partial h} \right] \\ &= \frac{\partial L(\sigma_t^2, h_{t,m}^*)}{\partial h} + \frac{1}{2m} E_{t-1} \left[\frac{\partial^3 L(\hat{\sigma}_{t,m}^2, h_{t,m}^*)}{\partial (\sigma_t^2)^2 \partial h} (\hat{\sigma}_{t,m}^2 - \sigma_t^2)^2 \right] \end{aligned}$$

In the absence of noise in the volatility proxy (i.e. $\hat{\sigma}_{t,m}^2 = \sigma_t^2$ a.s.) the second term above would equal zero and the first-order condition would be the same as if the true conditional variance was observable. By assumption G4 this yields $h_{t,m}^* = \sigma_t^2$. One of the conditions of Hansen and Lunde (2006) was $\partial^3 L(\sigma^2, h) / \partial (\sigma^2)^2 \partial h = 0$, which implies that the second term above equals zero even in the presence of a noisy volatility proxy. For loss functions that yield $\partial^3 L(\sigma^2, h) / \partial (\sigma^2)^2 \partial h \neq 0$ the presence of noise in the volatility proxy distorts the first-order condition from what it would be in the absence of noise, and thus affects the optimal forecast. If $\partial^3 L(\sigma^2, h) / \partial (\sigma^2)^2 \partial h > (<) 0 \forall (\sigma^2, h)$, then the FOC implies that we must have $\partial L(\sigma_t^2, h_{t,m}^*) / \partial h < (>) 0$, which implies that $h_{t,m}^* < (>) \sigma_t^2$, by assumption G4. ■

Proof of Proposition 2. We prove this proposition by showing the equivalence of the following three statements:

- S1: The loss function takes the form given the statement of the proposition;
- S2: The loss function is robust in the sense of Definition 1;
- S3: The optimal forecast under the loss function is the conditional variance.

We will show that $\mathcal{S1} \Rightarrow \mathcal{S2} \Rightarrow \mathcal{S3} \Rightarrow \mathcal{S1}$. That $\mathcal{S1} \Rightarrow \mathcal{S2}$ follows from Hansen and Lunde (2006): their assumption 2 is satisfied given the assumptions for the proposition and noting that $\partial^2 L(\hat{\sigma}^2, h) / \partial (\hat{\sigma}^2)^2 = B''(\hat{\sigma}^2)$ does not depend on h .

We next show that $\mathcal{S}2 \Rightarrow \mathcal{S}3$: by the definition of h_t^* we have

$$E_{t-1} [L(\hat{\sigma}_t^2, h_t^*)] \leq E_{t-1} [L(\hat{\sigma}_t^2, \tilde{h}_t)]$$

for any other sequence of \mathcal{F}_{t-1} -measurable forecasts \tilde{h}_t . Then

$$\begin{aligned} E [L(\hat{\sigma}_t^2, h_t^*)] &\leq E [L(\hat{\sigma}_t^2, \tilde{h}_t)] \text{ by the LIE} \\ \text{and } E [L(\sigma_t^2, h_t^*)] &\leq E [L(\sigma_t^2, \tilde{h}_t)] \text{ since } L \text{ is robust under } \mathcal{S}2 \end{aligned}$$

But $L(\hat{\sigma}^2, h)$ has a unique minimum at $\hat{\sigma}^2 = h$, and if we set $\tilde{h}_t = \sigma_t^2$ then it must be the case that $h_t^* = \sigma_t^2$.

Proving $\mathcal{S}3 \Rightarrow \mathcal{S}1$ is more challenging. For this part we follow the proof of Theorem 1 of Komunjer and Vuong (2004), adapted to our problem. We seek to show that the functional form of the loss function given in the proposition is necessary for $h_t^* = E_{t-1} [\hat{\sigma}_t^2]$, for any $F_t \in \tilde{F}$. Notice that we can write

$$\frac{\partial L(\hat{\sigma}_t^2, h_t)}{\partial h} = c(\hat{\sigma}_t^2, h_t) (\hat{\sigma}_t^2 - h_t)$$

where $c(\hat{\sigma}_t^2, h_t) = (\hat{\sigma}_t^2 - h_t)^{-1} \partial L(\hat{\sigma}_t^2, h_t) / \partial h$, since $\hat{\sigma}_t^2 \neq h_t$ a.s. by assumption A2. Now decompose $c(\hat{\sigma}_t^2, h_t)$ into

$$c(\hat{\sigma}_t^2, h_t) = E_{t-1} [c(\hat{\sigma}_t^2, h_t)] + \varepsilon_t$$

where $E_{t-1} [\varepsilon_t] = 0$. Thus

$$\begin{aligned} E_{t-1} \left[\frac{\partial L(\hat{\sigma}_t^2, h_t^*)}{\partial h} \right] &= E_{t-1} [c(\hat{\sigma}_t^2, h_t^*) (\hat{\sigma}_t^2 - h_t^*)] \\ &= E_{t-1} [c(\hat{\sigma}_t^2, h_t)] E_{t-1} [\hat{\sigma}_t^2 - h_t^*] + E_{t-1} [\varepsilon_t (\hat{\sigma}_t^2 - h_t^*)] \end{aligned}$$

If $E_{t-1} [\partial L(\hat{\sigma}_t^2, h_t^*) / \partial h] = 0$ for $h_t^* = E_{t-1} [\hat{\sigma}_t^2]$, then it must be that $E_{t-1} [\hat{\sigma}_t^2 - h_t^*] = 0 \Rightarrow E_{t-1} [\varepsilon_t (\hat{\sigma}_t^2 - h_t^*)] = 0$ for all $F_t \in \tilde{F}$. Employing a generalised Farkas lemma, see Lemma 8.1 of Gourieroux and Monfort (1996), this implies that $\exists \lambda \in \mathbb{R}$ such that $\lambda (\hat{\sigma}_t^2 - h_t^*) = \varepsilon_t (\hat{\sigma}_t^2 - h_t^*)$ for every $F_t \in \tilde{F}$ and for all t . Since $\hat{\sigma}_t^2 - h_t^* \neq 0$ a.s. by assumption A2 this implies that $\varepsilon_t = \lambda$ a.s. for all t . Since $E_{t-1} [\varepsilon_t] = 0$ we then have $\lambda = 0$. Thus $c(\hat{\sigma}_t^2, h_t^*) = E_{t-1} [c(\hat{\sigma}_t^2, h_t^*)]$ for all t , which implies that $c(\hat{\sigma}_t^2, h_t^*) = c(h_t^*)$, and thus that $\partial L(\hat{\sigma}_t^2, h_t) / \partial h = c(h_t) (\hat{\sigma}_t^2 - h_t)$.

The remainder of the proof is straightforward: A necessary condition for h_t^* to minimise $E_{t-1} [L(\hat{\sigma}_t^2, h)]$ is that $E_{t-1} [\partial^2 L(\hat{\sigma}_t^2, h_t^*) / \partial h^2] \geq 0$, using A5 to interchange expectation and differentiation. Using the previous result we have:

$$E_{t-1} \left[\frac{\partial^2 L(\hat{\sigma}_t^2, h_t^*)}{\partial h^2} \right] = E_{t-1} [c'(h_t^*) (\hat{\sigma}_t^2 - h_t^*) - c(h_t^*)] = -c(h_t^*)$$

which is non-negative iff $c(h_t^*)$ is non-positive. From assumption A4 we know that the optimum is in the interior of \mathcal{H} and so we know that $c \neq 0$, and thus $c(h) < 0 \forall h \in \mathcal{H}$. To obtain the loss function corresponding to the given first derivative we simply integrate up:

$$\begin{aligned} L(\hat{\sigma}^2, h) &= \hat{\sigma}^2 \int c(h) dh - \int c(h) h dh \\ &= B(\hat{\sigma}^2) + \hat{\sigma}^2 C(h) - C(h) h + \int C(h) dh \\ &= \tilde{C}(h) + B(\hat{\sigma}^2) + C(h) (\hat{\sigma}^2 - h) \end{aligned}$$

where C is a strictly decreasing function (i.e. $C' \equiv c$ is negative) and \tilde{C} is the anti-derivative of C . By assumption A3 both B and C are twice continuously differentiable. Thus $\mathcal{S3} \Rightarrow \mathcal{S1}$, completing the proof. ■

Proof of Proposition 3. Without loss of generality, we work below with loss functions that have been normalised to imply zero loss when the forecast error is zero: $L(\hat{\sigma}^2, h) = \tilde{C}(h) - \tilde{C}(\hat{\sigma}^2) + C(h) (\hat{\sigma}^2 - h)$.

(i) We want to find the general sub-set of loss functions that satisfy $L(\hat{\sigma}^2, h) = \tilde{L}(\hat{\sigma}^2 - h) \forall (\hat{\sigma}^2, h)$ for some function \tilde{L} . This condition implies

$$\begin{aligned} \frac{\partial L(\hat{\sigma}^2, h)}{\partial \hat{\sigma}^2} &= -\frac{\partial L(\hat{\sigma}^2, h)}{\partial h} \forall (\hat{\sigma}^2, h) \\ -C(\hat{\sigma}^2) + C(h) + C'(h) (\hat{\sigma}^2 - h) &= 0 \forall (\hat{\sigma}^2, h) \end{aligned}$$

Taking the derivative of both sides w.r.t. $\hat{\sigma}^2$ we obtain:

$$\begin{aligned} -C'(\hat{\sigma}^2) + C'(h) &= 0 \forall (\hat{\sigma}^2, h) \\ \text{which implies } C'(h) &= \kappa_1 \forall h \end{aligned}$$

and since we know C is strictly decreasing, we also have $\kappa_1 < 0$.

$$\begin{aligned} \text{so } C(h) &= \kappa_1 h + \kappa_2 (\hat{\sigma}^2) \\ \tilde{C}(h) &= \frac{1}{2} \kappa_1 h^2 + \kappa_2 (\hat{\sigma}^2) h + \kappa_3 (\hat{\sigma}^2) \end{aligned}$$

where κ_2, κ_3 are constants of integration, and may be functions of $\hat{\sigma}^2$. Thus the loss function becomes

$$\begin{aligned} L(\hat{\sigma}^2, h) &= \frac{1}{2}\kappa_1 h^2 + \kappa_2(\hat{\sigma}^2)h + \kappa_3(\hat{\sigma}^2) \\ &\quad - \frac{1}{2}\kappa_1 \hat{\sigma}^4 - \kappa_2(\hat{\sigma}^2)\hat{\sigma}^2 - \kappa_3(\hat{\sigma}^2) \\ &\quad + (\kappa_1 h + \kappa_2(\hat{\sigma}^2))(\hat{\sigma}^2 - h) \\ &= -\frac{1}{2}\kappa_1(\hat{\sigma}^2 - h)^2 \end{aligned}$$

Since proportionality constants do not affect the loss function, we find that the only loss function that depends on $(\hat{\sigma}^2, h)$ only through the forecast error, $\hat{\sigma}^2 - h$, is the MSE loss function.

(ii) We next want to find the general sub-set of loss functions that satisfy $L(\hat{\sigma}^2, h) = \tilde{L}(\hat{\sigma}^2/h) \forall (\hat{\sigma}^2, h)$ for some function \tilde{L} . Note that this condition implies that L is homogeneous of degree zero. Using Proposition 5 below, this implies that the loss function must be of the form:

$$L(\hat{\sigma}^2, h) = \frac{\hat{\sigma}^2}{h} - \log \frac{\hat{\sigma}^2}{h} - 1$$

which is the QLIKE loss function up to additive and multiplicative constants. ■

Proof of Proposition 4. (i) If L is homogeneous then $E[L(a\hat{\sigma}_t^2, ah_{1t})] \geq E[L(a\hat{\sigma}_t^2, ah_{2t})] \Leftrightarrow E[a^k L(\hat{\sigma}_t^2, h_{1t})] \geq E[a^k L(\hat{\sigma}_t^2, h_{2t})] \Leftrightarrow E[L(\hat{\sigma}_t^2, h_{1t})] \geq E[L(\hat{\sigma}_t^2, h_{2t})]$, for any $a > 0$.

(ii) Here we need only provide an example. Consider the following stylised case: $\sigma_t^2 = 1$ a.s. $\forall t$, $(h_{1t}, h_{2t}) = (\gamma_1, \gamma_2) \forall t$, and $\hat{\sigma}_t^2$ is such that $E_{t-1}[\hat{\sigma}_t^2] = 1$ a.s. $\forall t$. As a robust but non-homogeneous loss we will use the one generated by the following specification for C' :

$$\begin{aligned} C'(h) &= -\log(1+h) \\ \text{so } C(h) &= h - (1+h)\log(1+h) \\ \text{and } \tilde{C}(h) &= \frac{1}{4} \left[h(3h+2) - 2(1+h)^2 \log(1+h) \right] \end{aligned}$$

For small h this loss function resembles the the $b = 1$ loss function from Proposition 5 (up to a scaling constant), but for medium to large h this loss function does not correspond to any in Proposition 5.

Given this set-up, we have

$$\begin{aligned} E [L (a\hat{\sigma}_t^2, ah_{it})] &= \frac{1}{4} \left[a\gamma_i (3a\gamma_i + 2) - 2(1 + a\gamma_i)^2 \log(1 + a\gamma_i) \right] - E [\tilde{C} (a\hat{\sigma}_t^2)] \\ &\quad + a [a\gamma_i - (1 + a\gamma_i) \log(1 + a\gamma_i)] (1 - \gamma_i) \end{aligned}$$

Then define

$$\begin{aligned} d_t (\gamma_1, \gamma_2, a) &\equiv L (a\hat{\sigma}_t^2, a\gamma_1) - L (a\hat{\sigma}_t^2, a\gamma_2) \\ E [d_t (\gamma_1, \gamma_2, a)] &= \frac{a}{4} (\gamma_1 - \gamma_2) (2 - 4a - a(\gamma_1 + \gamma_2)) \\ &\quad + \frac{1}{2} \left(a^2 (\gamma_1 - 1)^2 - (1 + a)^2 \right) \log(1 + a\gamma_1) \\ &\quad - \frac{1}{2} \left(a^2 (\gamma_2 - 1)^2 - (1 + a)^2 \right) \log(1 + a\gamma_2) \end{aligned}$$

Then note that $E [d_t (1/3, 3/2, 1)] = -0.0087$, and so the first forecast has lower expected loss than the second using the “original” scaling of the data. But $E [d_t (1/3, 3/2, 2)] = 0.0061$, and so if all variables are multiplied by 2 then the second forecast has lower expected loss than the first. ■

Proof of Proposition 5. We seek the subset of robust loss functions that are homogeneous of order k : $L (a\hat{\sigma}^2, ah) = a^k L (\hat{\sigma}^2, h) \forall a > 0$. Let

$$\begin{aligned} \lambda (\hat{\sigma}^2, h) &\equiv \partial L (\hat{\sigma}^2, h) / \partial h \\ &= C' (h) (\hat{\sigma}^2 - h) \text{ for robust loss functions.} \end{aligned}$$

Since L is homogeneous of order k , λ is homogeneous of order $(k - 1)$. This implies $\lambda (a\hat{\sigma}^2, ah) = a^{k-1} \lambda (\hat{\sigma}^2, h) = a^{k-1} C' (h) (\hat{\sigma}^2 - h)$, while direct substitution yields $\lambda (a\hat{\sigma}^2, ah) = a C' (ah) (\hat{\sigma}^2 - h)$. Thus $C' (ah) = a^{k-2} C' (h) \forall a > 0$, that is, C' is homogeneous of order $(k - 2)$.

Next we apply Euler’s theorem to C' : $C'' (h) h = (k - 2) C' (h) \forall h > 0$, and so

$$(2 - k) C' (h) + C'' (h) h = 0$$

We can solve this first-order differential equation to find:

$$C' (h) = \gamma h^{k-2}$$

where γ is an unknown scalar. Since $C' < 0$ we know that $\gamma < 0$, and as this is just a scaling parameter we set it to -1 without loss of generality.

$$\begin{aligned} C'(h) &= -h^{k-2} \\ C(h) &= \begin{cases} \frac{1}{1-k}h^{k-1} + z_1 & k \neq 1 \\ -\log h + z_1 & k = 1 \end{cases} \\ \tilde{C}(h) &= \begin{cases} z_1 h + \frac{1}{k(1-k)}h^k + z_2 & k \notin \{0, 1\} \\ z_1 h + h - h \log h + z_2 & k = 1 \\ z_1 h + \log h + z_2 & k = 0 \end{cases} \end{aligned}$$

where z_1 and z_2 are constants of integration. Finally, we substitute the expressions for C and \tilde{C} into equation (23), set $B = -\tilde{C}$, and simplify to obtain the loss functions in equation (24) with $k = b + 2$. ■

7 Technical Appendix: supporting calculations for Section 2

Section 2.1:

MSE-log:

$$\begin{aligned} h_t^* &= \exp \{ E_{t-1} [\log \varepsilon_t^2] \} \sigma_t^2 \\ &= \begin{cases} (\nu - 2) \exp \{ \Psi(\frac{1}{2}) - \Psi(\frac{\nu}{2}) \} \sigma_t^2, & \text{if } r_t | \mathcal{F}_{t-1} \sim \text{Student's } t(0, \sigma_t^2, \nu), \nu > 2 \\ \frac{1}{2} \exp \{ -\gamma_E \} \sigma_t^2 \approx 0.28 \sigma_t^2, & \text{if } r_t | \mathcal{F}_{t-1} \sim N(0, \sigma_t^2) \end{cases} \end{aligned}$$

where Ψ is the digamma function and $\gamma_E = -\Psi(1) \approx 0.58$ is Euler's constant, see Harvey, et al. (1994).

MSE-prop:

$$\begin{aligned} h_t^* &= \frac{E_{t-1} [r_t^4]}{E_{t-1} [r_t^2]} = Kurtosis_{t-1} [r_t] \sigma_t^2 \\ &= \begin{cases} 3 \left(\frac{\nu-2}{\nu-4} \right) \sigma_t^2, & \text{if } r_t | \mathcal{F}_{t-1} \sim \text{Student's } t(0, \sigma_t^2, \nu), \nu > 4 \\ 3 \sigma_t^2, & \text{if } r_t | \mathcal{F}_{t-1} \sim N(0, \sigma_t^2) \end{cases} \end{aligned}$$

QLIKE: $h_t^* = E_{t-1} [r_t^2] = \sigma_t^2$

MAE: As usual with an absolute-error loss function we obtain the median as the optimal forecast:

$$\begin{aligned}
h_t^* &= \text{Median}_{t-1} [r_t^2] \\
&= \sigma_t^2 \cdot \text{Median}_{t-1} [\varepsilon_t^2] \\
&= \begin{cases} \sigma_t^2 \cdot \frac{\nu-2}{\nu} \cdot \text{Median} [F_{1,\nu}], & \text{if } r_t | \mathcal{F}_{t-1} \sim \text{Student's } t(0, \sigma_t^2, \nu) \\ \sigma_t^2 \cdot \text{Median} [\chi_1^2] \approx 0.45\sigma_t^2, & \text{if } r_t | \mathcal{F}_{t-1} \sim N(0, \sigma_t^2) \end{cases} \quad (27)
\end{aligned}$$

where $\varepsilon_t \equiv r_t/\sigma_t$ and $\text{Median}_{t-1} [r_t^2]$ is the conditional median of r_t^2 given F_{t-1} .

MAE-log: $h_t^* = \exp \{ \text{Median}_{t-1} [\log \varepsilon_t^2] \} \sigma_t^2 = \text{Median}_{t-1} [\varepsilon_t^2] \sigma_t^2$, since $\text{Median} [\log (X)] = \log (\text{Median} [X])$ for any non-negative random variable X . Thus the optimal forecast is identical to that under MAE loss, given above.

MAE-SD: $h_t^* = (\text{Median}_{t-1} [|\varepsilon_t|])^2 \sigma_t^2 = \text{Median}_{t-1} [\varepsilon_t^2] \sigma_t^2$, since $\text{Median} [X]^2 = \text{Median} [X^2]$ for any non-negative random variable X . Thus the optimal forecast is identical to that under MAE loss, given above.

MAE-prop: If $r_t^2 | \mathcal{F}_{t-1} \sim F_t(\sigma_t^2)$ and $\varepsilon_t^2 \equiv r_t^2/\sigma_t^2 | \mathcal{F}_{t-1} \sim G_t(1)$ then

$$\begin{aligned}
\text{FOC} \quad 0 &= \int_0^{h_t^*} \frac{r_t^2}{h_t^*} f_t(r_t^2) dr_t^2 - \int_{h_t^*}^{\infty} \frac{r_t^2}{h_t^*} f_t(r_t^2) dr_t^2 \\
\text{so } \int_0^{h_t^*} \frac{r_t^2}{h_t^*} f_t(r_t^2) dr_t^2 &= \int_{h_t^*}^{\infty} \frac{r_t^2}{h_t^*} f_t(r_t^2) dr_t^2 \\
F_t(h_t^*) E_{t-1} \left[\frac{r_t^2}{h_t^*} \middle| r_t^2 \leq h_t^* \right] &= (1 - F_t(h_t^*)) E_{t-1} \left[\frac{r_t^2}{h_t^*} \middle| r_t^2 > h_t^* \right]
\end{aligned}$$

without loss of generality let $h_t^* \equiv \sigma_t^2 \gamma_t^*$, $\gamma_t^* > 0$, so

$$\begin{aligned}
F_t(\sigma_t^2 \gamma_t^*) E_{t-1} \left[\frac{\varepsilon_t^2}{\gamma_t^*} \middle| \varepsilon_t^2 \leq \gamma_t^* \right] &= (1 - F_t(\sigma_t^2 \gamma_t^*)) E_{t-1} \left[\frac{\varepsilon_t^2}{\gamma_t^*} \middle| \varepsilon_t^2 > \gamma_t^* \right] \\
G_t(\gamma_t^*) E_{t-1} [\varepsilon_t^2 | \varepsilon_t^2 \leq \gamma_t^*] &= (1 - G_t(\gamma_t^*)) E_{t-1} [\varepsilon_t^2 | \varepsilon_t^2 > \gamma_t^*]
\end{aligned}$$

If $\varepsilon_t^2 | \mathcal{F}_{t-1} \sim G(1)$, then $\gamma_t^* = \gamma^* \forall t$. Finding an explicit expression for h_t^* is difficult, and so we used 10,000 simulated draws for $\nu = \{4, 6, 10, 20, 30, 50, 100, 1000, \infty\}$ and numerically obtained h_t^* for each ν . We then used OLS to find the approximation given in Table 1, which yielded an R^2 of 0.9667.

Section 2.2:

Wherever possible we derived solutions or approximate solutions analytically. This was not always possible and so in some cases we had to resort to simulations to obtain solutions. Feller (1951) presents the density of the range:

$$f(RG_t; \sigma_t) = 8 \sum_{k=1}^{\infty} (-1)^{k-1} \frac{k^2}{\sigma_t} \phi\left(\frac{k \cdot RG_t}{\sigma_t}\right)$$

where ϕ is the standard normal *pdf*. For practical purposes the sum in the above expression needs to be truncated at some finite value; we truncate at $k = 1000$. Parkinson (1980) presented the *cdf* of the range, and a formula for obtaining moments:

$$\begin{aligned} F(RG_t; \sigma_t) &= \sum_{k=1}^{\infty} (-1)^{k-1} k \left\{ \operatorname{erfc}\left(\frac{(k+1)RG_t}{\sigma\sqrt{2}}\right) - 2\operatorname{erfc}\left(\frac{k \cdot RG_t}{\sigma\sqrt{2}}\right) + \operatorname{erfc}\left(\frac{(k-1)RG_t}{\sigma\sqrt{2}}\right) \right\} \\ E[RG_t^p] &= \frac{4}{\sqrt{\pi}} \Gamma\left(\frac{p+1}{2}\right) \left(2^{p/2} - 2^{2-p/2}\right) \zeta(p-1) \sigma_t^p, \text{ for } p \geq 1 \end{aligned}$$

where $\operatorname{erfc}(x) \equiv 1 - \operatorname{erf}(x)$, $\operatorname{erf}(x)$ is the ‘error function’: $\operatorname{erf}(x) \equiv 2/\sqrt{\pi} \int_0^x e^{-t^2} dt$. ζ is the Riemann zeta function. From this expression we can obtain the necessary moments for computing optimal forecasts when the range is used as a volatility proxy. For the first and second moments of RG_t we can obtain simple expressions, but the fourth moment involves $\zeta(3) = \sum_{k=1}^{\infty} k^{-3}$ which is an irrational number, and thus only a numerical expression is available. In addition to the moments of RG_t , we will need the mean of $\log RG_t$ and the median of RG_t . We used quadrature and OLS to obtain the expression¹⁵:

$$E_{t-1}[\log RG_t] = 0.4257 + \log \sigma_t \quad (28)$$

which is consistent with the expression given in Alizadeh, et al. (2002). We numerically inverted the *cdf* of the range, given in Parkinson (1980), and used OLS to determine the following relation¹⁶:

$$\begin{aligned} \operatorname{Median}_{t-1}[RG_t] &= 1.5145\sigma_t \\ \text{so } \operatorname{Median}_{t-1}[RG_t^2] &= 2.2938\sigma_t^2, \text{ since } RG_t \text{ is weakly positive.} \end{aligned}$$

MSE-LOG: $h_t^* = \exp\{E_{t-1}[\log \hat{\sigma}_t^2]\}$. A Taylor series approximation did not provide a good fit when considering realised variance as a proxy, and so we resorted to simulations. We simulated

¹⁵We used quadrature to estimate $E_{t-1}[\log RG_t]$ for $\sigma_t = 0.5, 1, 1.5, \dots, 10$. We then regressed these estimates on a constant and $\log \sigma_t$ to obtain the parameter estimates. The R^2 from this regression was 1.0000.

¹⁶The R^2 from this relation for $\sigma = 0.5, 1, 1.5, \dots, 10$ was 1.0000.

50,000 “days” worth of observations, where the number of observations per day considered was $m = \{1, 3, 5, 7, 10, 13, 20, 40, 60, 78, 100\}$. The following expression yielded an R^2 of 0.9959 : $E_{t-1} \left[\log RV_t^{(m)} \right] \approx -1.2741/m$, so the optimal forecast under our DGP assumption is $h_t^* \approx \sigma_t^2 e^{-1.2741/m}$.

For the range we find that

$$\begin{aligned} E_{t-1} [\log RG_t^{*2}] &= 2E_{t-1} [\log RG_t^*] \\ &= -0.1684 + \log \sigma_t^2 \\ \text{so } h_t^* &= e^{-0.1684} \sigma_t^2 \approx 0.8450 \sigma_t^2 \end{aligned}$$

MAE-log: The optimal forecast is $h_t^* = \text{Median}_{t-1} [\hat{\sigma}_t^2]$, since $\hat{\sigma}_t^2$ is weakly positive we know that $\log (\text{Median}_{t-1} [\hat{\sigma}_t^2]) = \text{Median}_{t-1} [\log \hat{\sigma}_t^2]$, and so the results for this loss function are identical to those for the MAE loss function.

MAE-SD: The optimal forecast is $h_t^* = \text{Median}_{t-1} [\hat{\sigma}_t^2]$. Since $\hat{\sigma}_t^2$ is weakly positive we know that $\text{Median}_{t-1} [\hat{\sigma}_t^2] = (\text{Median}_{t-1} [\hat{\sigma}_t])^2$, and so the results for this loss function are identical to those for the MAE loss function.

MSE-prop: $h_t^* = E_{t-1} [\hat{\sigma}_t^4] / E_{t-1} [\hat{\sigma}_t^2]$. When realised volatility is used as the proxy we find: $h_t^* = \left(\frac{1}{m} \text{Kurtosis}_{t-1} [r_{t,i}] + \frac{m-1}{m} \right) \sigma_t^2 = \left(1 + \frac{2}{m} \right) \sigma_t^2$. For the range we find that: $h_t^* = 10.8185 / \left((\log 16)^2 \right) \sigma_t^2 \approx 1.4073 \sigma_t^2$.

MAE-prop: For realised variance, like the daily squared return, obtaining an analytical, even approximate, solution to this problem is difficult and so we used simulations. In the set-up given in the text it is again possible to show that the optimal forecast is of the form $h_t^* = \gamma^* \sigma_t^2$. For realised volatility we simulated 50,000 “days” worth of observations, where the number of observations per day considered was $m = \{1, 3, 5, 7, 10, 13, 20, 40, 60, 78, 100\}$, and used numerical methods to locate the optimum forecast. The following expression yielded an R^2 of 0.9999 : $h_t^* \approx \left(1 + \frac{1.3624}{m} \right) \sigma_t^2$. For the range we again used a numerical minimisation algorithm combined with quadrature to compute the expectation in the optimisation problem: $h_t^* \approx 0.9941 \sigma_t^2$.

References

- [1] Alizadeh, S., Brandt, M.W., and Diebold, F.X., 2002, Range-Based Estimation of Stochastic Volatility Models, *Journal of Finance*, 57(3), 1047-1091.
- [2] Andersen, T.G., Benzoni, L., and Lund, J., 2002, An Empirical Investigation of Continuous-Time Equity Return Models, *Journal of Finance*, 57(3), 1239-1284.
- [3] Andersen, T.G., and Bollerslev, T., 1998, Answering the Skeptics: Yes, Standard Volatility Models Do Provide Accurate Forecasts, *International Economic Review*, 39, 885-905.
- [4] Andersen, T.G., Bollerslev, T., and Lange, S., 1999, Forecasting Financial Market Volatility: Sample Frequency Vis-à-vis Forecast Horizon, *Journal of Empirical Finance*, 6, 457-477.
- [5] Andersen, T.G., Bollerslev, T., Diebold, F.X., and Ebens, H., 2001, The Distribution of Realized Stock Return Volatility, *Journal of Financial Economics*, 61, 43-76.
- [6] Andersen, T.G., Bollerslev, T., and Diebold, F.X., 2002, "Parametric and Nonparametric Volatility Measurement," forthcoming in L.P. Hansen and Y. Aït-Sahalia (ed.s), *Handbook of Financial Econometrics*, North-Holland Press, Amsterdam.
- [7] Andersen, T.G., Bollerslev, T., Diebold, F.X. and Labys, P., 2003, Modeling and Forecasting Realized Volatility, *Econometrica*, 71(2), 579-625.
- [8] Andersen, T.G., Bollerslev, T., and Meddahi, N., 2004, Analytic Evaluation of Volatility Forecasts, *International Economic Review*, 45, 1079-1110.
- [9] Andersen, T.G., Bollerslev, T., and Meddahi, N., 2005, Correcting the Errors: Volatility Forecast Evaluation Using High-Frequency Data and Realized Volatilities, *Econometrica*, 73(1), 279-296.
- [10] Andersen, T.G., Bollerslev, T., Christoffersen, P.F., and Diebold, F.X., 2006, Volatility and Correlation Forecasting, in the *Handbook of Economic Forecasting*, G. Elliott, C.W.J. Granger and A. Timmermann ed.s, North Holland Press, Amsterdam.
- [11] Ball, C.A., and Torous, W.N., 1984, The Maximum Likelihood Estimation of Security Price Volatility: Theory, Evidence, and Application to Option Pricing, *Journal of Business*, 57(1), 97-112.
- [12] Barndorff-Nielsen, O.E., and Shephard, N., 2002, Econometric Analysis of Realised Volatility and Its Use in Estimating Stochastic Volatility Models, *Journal of the Royal Statistical Society, Series B*, 64, 253-280.
- [13] Barndorff-Nielsen, O.E., and Shephard, N., 2004, Econometric Analysis of Realized Covariation: High Frequency Based Covariance, Regression and Correlation in Financial Economics, *Econometrica*, 72(3), 885-925.
- [14] Bollerslev, T., and Ghysels, E., 1994, Periodic Autoregressive Conditional Heteroscedasticity, *Journal of Business and Economic Statistics*, 14(2), 139-151.

- [15] Bollerslev, T., Engle, R.F., and Nelson, D.B., 1994, ARCH Models, in the *Handbook of Econometrics*, R.F. Engle and D. McFadden ed.s, North Holland Press, Amsterdam.
- [16] Christensen, K., and Podolskij, M., 2006, Realized Range-Based Estimation of Integrated Variance, *Journal of Econometrics*, forthcoming.
- [17] Christoffersen, P.F., and Diebold, F.X., 1997, Optimal prediction under asymmetric loss, *Econometric Theory*, 13, 808-817.
- [18] Christoffersen, P.F., and Jacobs, K., 2004, The Importance of the Loss Function in Option Valuation, *Journal of Financial Economics*, 72, 291-318.
- [19] Clements, M.P., 2005, *Evaluating Econometric Forecasts of Economic and Financial Variables*, Palgrave MacMillan, United Kingdom.
- [20] Corradi, V., and Swanson, N.R., 2006, Predictive Density Evaluation, in the *Handbook of Economic Forecasting*, G. Elliott, C.W.J. Granger and A. Timmermann (ed.s), North Holland Press, Amsterdam.
- [21] Cowles, A., 1933, Can Stock Market Forecasters Forecast?, *Econometrica*, 1(3), 309-324.
- [22] Diebold, F.X., and Mariano, R.S., 1995, Comparing Predictive Accuracy, *Journal of Business and Economic Statistics*, 13(3), 253-263.
- [23] Diebold, F.X., and Lopez, J. A., 1996, Forecast Evaluation and Combination, in the *Handbook of Statistics*, G.S. Maddala and C.R. Rao (ed.s), Amsterdam: North-Holland, 241-268.
- [24] Engle, R.F., 1993, A Comment on Hendry and Clements on the Limitations of Comparing Mean Square Forecast Errors, *Journal of Forecasting*, 12, 642-644.
- [25] Engle, R.F., Hong, C.-H., Kane, A., and Noh, J., 1993, Arbitrage valuation of variance forecasts with simulated options, in: D. Chance and R. Tripp, eds., *Advances in Futures and Options Research*, JIA Press, Greenwich, U.S.A.
- [26] Feller, W., 1951, The Asymptotic Distribution of the Range of Sums of Random Variables, *Annals of Mathematical Statistics*, 22, 427-432.
- [27] Garman, M.B., and Klass, M.J., 1980, On the Estimation of Security Price Volatilities from Historical Data, *Journal of Business*, 53(1), 67-78.
- [28] Giacomini, R., and White, H., 2006, Tests of Conditional Predictive Ability, *Econometrica*, 74(6), 1545-1578.
- [29] Gonçalves, S., and Meddahi, N., 2005, Bootstrapping Realized Volatility, working paper, Université de Montréal.
- [30] Gouriéroux, C. and Monfort, A., 1996, *Statistics and Econometric Models, Volume 1*, translated from the French by Q. Vuong, Cambridge University Press, Great Britain.
- [31] Gouriéroux, C., Monfort, A., and Trognon, A., 1984, Pseudo Maximum Likelihood Methods: Theory, *Econometrica*, 52(3), 681-700.

- [32] Gouriéroux, C., Monfort, A. and Renault, E., 1987, Consistent M-Estimators in a Semi-Parametric Model, CEPREMAP working paper 8720.
- [33] Granger, C.W.J., 1969, Prediction with a generalized cost function, *OR*, 20, 199-207.
- [34] Hamilton, J.D., and Susmel, R., 1994, Autoregressive Conditional Heteroskedasticity and Changes in Regime, *Journal of Econometrics*, 64(1-2), 307-333.
- [35] Hansen, P.R., and Lunde, A., 2005, A Forecast Comparison of Volatility Models: Does Anything Beat a GARCH(1,1)?, *Journal of Applied Econometrics*, 20(7), 873-889.
- [36] Hansen, P.R., and Lunde, A., 2006, Consistent Ranking of Volatility Models, *Journal of Econometrics*, 131(1-2), 97-121.
- [37] Harvey, A., Ruiz, E., and Shephard, N., 1994, Multivariate Stochastic Volatility Models, *Review of Economic Studies*, 61, 247-264.
- [38] Huber, P.J., 1981, *Robust Statistics*, Wiley, New York, U.S.A.
- [39] Komunjer, I., and Vuong, Q., 2004, Efficient Conditional Quantile Estimation, working paper.
- [40] Lamoureux, C.G. and Lastrapes, W.D., 1993, Forecasting Stock Return Variance: Toward an Understanding of Stochastic Implied Volatilities, *Review of Financial Studies*, 6(2), 293-326.
- [41] Martens, M., and van Dijk, D., 2007, Measuring volatility with the realized range, *Journal of Econometrics* 138, 181-207.
- [42] Meddahi, N., 2001, A Theoretical Comparison between Integrated and Realized Volatilities, manuscript, Université de Montréal.
- [43] Mincer, J., and Zarnowitz, V., 1969, The Evaluation of Economic Forecasts, in Zarnowitz, J. (ed.) *Economic Forecasts and Expectations*, National Bureau of Economic Research, New York.
- [44] Pagan, A.R., and Schwert, G.W., 1990, Alternative Models for Conditional Volatility, *Journal of Econometrics*, 45, 267-290.
- [45] Parkinson, M., 1980, The Extreme Value Method for Estimating the Variance of the Rate of Return, *Journal of Business*, 53(1), 61-65.
- [46] Patton, A.J., 2006, Volatility Forecast Comparison Using Imperfect Volatility Proxies, Research Paper 175, Quantitative Finance Research Centre, University of Technology Sydney.
- [47] Patton, A.J., and Timmermann, A. 2004, Properties of Optimal Forecasts under Asymmetric Loss and Nonlinearity, *Journal of Econometrics*, forthcoming.
- [48] Patton, A.J., and Sheppard K., 2007, Evaluating Volatility and Correlation Forecasts, in the *Handbook of Financial Time Series*, T.G. Andersen, R.A. Davis, J.-P. Kreiss and T. Mikosch (ed.s), Springer Verlag, forthcoming.

- [49] Poon, S.-H., and Granger, C.W.J., 2003, Forecasting Volatility in Financial Markets, *Journal of Economic Literature*, 41, 478-539.
- [50] Shephard, N., 2005, *Stochastic Volatility: Selected Readings*, Oxford University Press, United Kingdom.
- [51] West, K.D., 1996, Asymptotic Inference About Predictive Ability, *Econometrica*, 64, 1067-1084.
- [52] West, K.D., 2006, Forecast Evaluation, in the *Handbook of Economic Forecasting*, G. Elliott, C.W.J. Granger and A. Timmermann (ed.s), North Holland Press, Amsterdam.
- [53] West, K.D., Edison, H.J., and Cho, D., 1993, A Utility-Based Comparison of Some Models of Exchange Rate Volatility, *Journal of International Economics*, 35, 23-45.

8 Tables and Figures

Table 1: Optimal forecasts under various loss functions

		Distribution of daily returns			
		<i>Student's t</i> ($0, \sigma_t^2, \nu$)			
Loss function	$F_t(0, \sigma_t^2)$	ν	$\nu = 6$	$\nu = 10$	$\nu \rightarrow \infty$
<i>MSE</i>	σ_t^2	σ_t^2	σ_t^2	σ_t^2	σ_t^2
<i>QLIKE</i>	σ_t^2	σ_t^2	σ_t^2	σ_t^2	σ_t^2
<i>MSE-LOG</i>	$\exp \{E_{t-1} [\log \varepsilon_t^2]\} \sigma_t^2$	$\exp \{ \Psi(\frac{1}{2}) - \Psi(\frac{\nu}{2}) \} (\nu - 2) \sigma_t^2$	$0.22\sigma_t^2$	$0.25\sigma_t^2$	$0.28\sigma_t^2$
<i>MSE-SD</i>	$(E_{t-1} [\varepsilon_t])^2 \sigma_t^2$	$\frac{\nu-2}{\pi} (\Gamma(\frac{\nu-1}{2}) / \Gamma(\frac{\nu}{2}))^2 \sigma_t^2$	$0.56\sigma_t^2$	$0.60\sigma_t^2$	$0.64\sigma_t^2$
<i>MSE-prop</i>	$Kurtosis_{t-1} [r_t] \sigma_t^2$	$3 \frac{\nu-2}{\nu-4} \sigma_t^2$	$6.00\sigma_t^2$	$4.00\sigma_t^2$	$3.00\sigma_t^2$
<i>MAE</i>	$Median_{t-1} [r_t^2]$	$\frac{\nu-2}{\nu} Median [F_{1,\nu}] \sigma_t^2$	$0.34\sigma_t^2$	$0.39\sigma_t^2$	$0.45\sigma_t^2$
<i>MAE-LOG</i>	$Median_{t-1} [r_t^2]$	$\frac{\nu-2}{\nu} Median [F_{1,\nu}] \sigma_t^2$	$0.34\sigma_t^2$	$0.39\sigma_t^2$	$0.45\sigma_t^2$
<i>MAE-SD</i>	$Median_{t-1} [r_t^2]$	$\frac{\nu-2}{\nu} Median [F_{1,\nu}] \sigma_t^2$	$0.34\sigma_t^2$	$0.39\sigma_t^2$	$0.45\sigma_t^2$
<i>MAE-prop</i> [†]	n/a	$(2.36 + \frac{1.00}{\nu} + \frac{7.78}{\nu^2}) \sigma_t^2$	$2.73\sigma_t^2$	$2.55\sigma_t^2$	$2.36\sigma_t^2$

Notes: This table presents the forecast that minimises the conditional expected loss when the squared return is used as a volatility proxy. That is, h_t^* minimises $E_{t-1} [L(r_t^2, h)]$, for various loss functions L . The first column presents the solutions when returns have an arbitrary conditional distribution $r_t | \mathcal{F}_{t-1} \sim F_t$ with mean zero and conditional variance σ_t^2 , the second, third, and fourth columns present results with returns have the standardised Student's t distribution, and the final column presents the solutions when returns are conditionally normally distributed. Γ is the gamma function and Ψ is the digamma function. [†]The expressions given for MAE-prop are based on a numerical approximation, see the technical appendix for details.

Table 2: Optimal forecasts under various loss functions, using realised volatility and range

Volatility proxy						
Loss function	Range	<i>Realised volatility</i>				
		Arbitrary m	$m = 1$	$m = 13$	$m = 78$	$m \rightarrow \infty$
<i>MSE</i>	σ_t^2	σ_t^2	σ_t^2	σ_t^2	σ_t^2	σ_t^2
<i>QLIKE</i>	σ_t^2	σ_t^2	σ_t^2	σ_t^2	σ_t^2	σ_t^2
<i>MSE-LOG</i> [†]	$0.85\sigma_t^2$	$e^{-1.2741/m}\sigma_t^2$	$0.28\sigma_t^2$	$0.91\sigma_t^2$	$0.98\sigma_t^2$	σ_t^2
<i>MSE-SD</i>	$0.92\sigma_t^2$	$\frac{1}{m} \left(E \left[\sqrt{\chi_m^2} \right] \right)^2 \sigma_t^2$	$0.56\sigma_t^2$	$0.96\sigma_t^2$	$0.99\sigma_t^2$	σ_t^2
<i>MSE-prop</i>	$1.41\sigma_t^2$	$\left(1 + \frac{2}{m} \right) \sigma_t^2$	$3.00\sigma_t^2$	$1.15\sigma_t^2$	$1.03\sigma_t^2$	σ_t^2
<i>MAE</i>	$0.83\sigma_t^2$	$\frac{1}{m} \text{Median} \left[\chi_m^2 \right] \sigma_t^2$	$0.45\sigma_t^2$	$0.95\sigma_t^2$	$0.99\sigma_t^2$	σ_t^2
<i>MAE-LOG</i>	$0.83\sigma_t^2$	$\frac{1}{m} \text{Median} \left[\chi_m^2 \right] \sigma_t^2$	$0.45\sigma_t^2$	$0.95\sigma_t^2$	$0.99\sigma_t^2$	σ_t^2
<i>MAE-SD</i>	$0.83\sigma_t^2$	$\frac{1}{m} \text{Median} \left[\chi_m^2 \right] \sigma_t^2$	$0.45\sigma_t^2$	$0.95\sigma_t^2$	$0.99\sigma_t^2$	σ_t^2
<i>MAE-prop</i> [†]	$1.19\sigma_t^2$	$\left(1 + \frac{1.3624}{m} \right) \sigma_t^2$	$2.36\sigma_t^2$	$1.10\sigma_t^2$	$1.02\sigma_t^2$	σ_t^2

Notes: This table presents the forecast that minimises the conditional expected loss when the range or realised volatility is used as a volatility proxy. That is, h_t^* minimises $E_{t-1} [L(\hat{\sigma}_t^2, h)]$, for $\hat{\sigma}_t^2 = RG_t^{*2}$ or $\hat{\sigma}_t^2 = RV_t$, for various loss functions L . In all cases returns are assumed to be generated as a zero mean Brownian motion with constant volatility within each trade day and no jumps. The cases of $m = 1, 13, 78$ correspond to the use of daily squared returns, realised variance with 30-minute returns and realised variance with 5-minute returns respectively. The case that $m \rightarrow \infty$ corresponds to the case where the conditional variance is observable ex-post without error. [†]For the MSE-LOG and MAE-prop loss functions we used simulations, numerical integration and numerical optimisation to obtain the expressions given. Details on the computation of the figures in this table are given in the technical appendix.

Table 3: Comparison of rolling window and RiskMetrics forecasts

Loss function	Volatility proxy			
	Daily <i>squared return</i>	65-min <i>realised vol</i>	15-min <i>realised vol</i>	5-min <i>realised vol</i>
$b = 1$	-1.58	-1.66	-1.30	-1.35
$b = 0$ (MSE)	-0.59	-0.80	-0.03	-0.13
$b = -1$	1.30	1.04	1.65	-1.55
$b = -2$ (QLIKE)	1.94	2.21*	2.73*	2.41*
$b = -5$	-0.17	0.25	1.63	0.65

Notes: This table presents the t-statistics from Diebold-Mariano-West tests of equal predictive accuracy for a 60-day rolling window forecast and a RiskMetrics forecast, for IBM over the period January 1994 to December 2003. A t-statistic greater than 1.96 in absolute value indicates a rejection of the null of equal predictive accuracy at the 0.05 level. These statistics are marked with an asterisk. The sign of the t-statistics indicates which forecast performed better for each loss function: a positive t-statistic indicates that the rolling window forecast produced larger average loss than the RiskMetrics forecast, while a negative sign indicates the opposite.




Figure 1: *Optimal forecasts under MSE-SD loss when true variance is 1, for various levels of kurtosis, using the standardised Student's t distribution. The dashed line represents the optimal forecast as $kurtosis \rightarrow \infty$.*




Figure 2: Loss functions for various choices of b . True $\sigma^2=2$ in this example, with the volatility forecast ranging between 0 and 4. $b=0$ and $b=-2$ correspond to the MSE and QLIKE loss functions respectively.




Figure 3: Ratio of losses from negative forecast errors to positive forecast errors, for various choices of b . True $\sigma^2=2$ in this example, with the volatility forecast ranging between 0 and 4. $b=0$ and $b=-2$ correspond to the MSE and QLIKE loss functions respectively.




Figure 4: *Conditional variance forecasts for IBM returns from 60-day rolling window and RiskMetrics models, January 1994 to December 2003.*