

Structural basis of fumarate hydratase deficiency

Sarah Picaud · Kathryn L. Kavanagh · Wyatt W. Yue ·
Wen Hwa Lee · Susanne Muller-Knapp ·
Opher Gileadi · James Sacchettini · Udo Oppermann

Received: 11 May 2009 / Revised: 27 January 2011 / Accepted: 31 January 2011 / Published online: 29 March 2011
© The Author(s) 2011. This article is published with open access at Springerlink.com

Abstract Fumarate hydratase catalyzes the stereospecific hydration across the olefinic double bond in fumarate leading to L-malate. The enzyme is expressed in mitochondrial and cytosolic compartments, and participates in the Krebs cycle in mitochondria, as well as in regulation of cytosolic fumarate levels. Fumarate hydratase deficiency is an autosomal recessive trait presenting as metabolic disorder with severe encephalopathy, seizures and poor neurological outcome. Heterozygous mutations are associated with a predisposition to cutaneous and uterine leiomyomas and to renal cancer. The crystal structure of human fumarate hydratase shows that mutations can be grouped into two distinct classes either affecting structural integrity of the core enzyme architecture, or are localized around the enzyme active site.

An interactive version of this manuscript (which may contain additional mutations appended after acceptance of this manuscript) may be found on the SSIEM website at:

<http://www.ssiem.org/resources/structures/FH>.

Abbreviations

FH	Fumarate hydratase
FHD	Fumarate hydratase deficiency
MCUL1	Multiple cutaneous and uterine leiomyomata
HLRC	Hereditary leiomyomatosis and renal cancer syndrome

Introduction

Fumarate hydratase (FH) and succinate dehydrogenase are two integral enzyme components of the Krebs cycle, and besides their essential role in the TCA cycle, can act as tumour suppressors (King et al. 2006). The *FH* gene codes for fumarate hydratase (or fumarase; EC 4.2.1.2), which catalyzes the stereospecific, reversible hydration of fumarate to L-malate. The *FH* gene localized at 1q42.1 codes for differentially processed, but sequence-wise identical cytosolic and mitochondrial forms. Whereas the mitochondrial enzyme is part of the TCA cycle, the cytosolic form is thought to utilize fumarate derived from different sources. Deficiency in FH activity causes an impaired energy production by interrupting the flow of metabolites through the Krebs cycle. Accumulation of fumarate is thought to competitively inhibit 2-oxo-glutarate dependent dioxygenases that regulate hypoxia inducible factor (HIF), thus activating oncogenic hypoxia pathways (Ratcliffe, 2007).

Due to their essential role in energy production, enzyme deficiencies result in early onset of severe encephalopathy (Kerrigan et al. 2000). Accordingly, autosomal recessive fumarate hydratase deficiency (FHD) caused by mutations in the *FH* gene results in fumaric aciduria, and common clinical

Communicated by: Johannes Zschocke

References to electronic databases: OMIM: 606812 150800, 605839, 136850; EC 4.2.1.2; Gene symbol: *FH*; GenBank: 19743875; URL to the interactive version of the article: <http://www.ssiem.org/resources/structures/FH/>; PDB code: 3E04

Competing interest: None declared.

S. Picaud · K. L. Kavanagh · W. W. Yue · W. H. Lee ·
S. Muller-Knapp · O. Gileadi · U. Oppermann (✉)
Structural Genomics Consortium, University of Oxford,
Old Road Campus,
Headington OX3 7DQ, UK
e-mail: udo.oppermann@sgc.ox.ac.uk

J. Sacchettini
Department of Biochemistry & Biophysics,
Texas A&M University,
College Station, TX 77843-2128, USA

U. Oppermann
Botnar Research Center, NIHR Oxford Biomedical Research Unit,
Oxford OX3 7LD, UK

Table 1 X-ray data collection and refinement statistics

Data collection	
Space group	P3 ₂ 21
a, b, c (Å)	188.5, 188.5, 114.6
γ	120°
Wavelength (Å)	1.000
Resolution (Å)*	25.0 – 1.95 (2.06 – 1.95)
R _{merge} (%)*	0.141 (0.732)
I/ σ I*	9.7 (2.0)
Completeness (%)*	99.3 (96.8)
Redundancy*	6.2 (5.0)
Refinement	
Resolution (Å)	41.27 – 1.90
No. reflections	168629
R _{work} /R _{free} (%)	19.7/24.4
No. atoms	
Protein	13160
Ligand/ion	12
Water	655
B-factors (Å ²)	
Main-chain	24.88
Side-chain and water	25.91
RMS deviations	
Bond lengths (Å)	0.010
Bond angles (°)	1.201
PDB code	3E04 (doi:10.2210/pdb3e04/pdb)

* Numbers in parentheses represent data in the highest resolution shell.

Fig. 1 Ribbon/surface diagram of human fumarate hydratase illustrating the tetrameric assembly of class II fumarases. Molecular surface representation is used to convey the overall shape of each monomer as well as the tetrameric assembly. Each monomer has been coloured distinctively, to facilitate visualization. Two monomers are represented using semi-transparent surfaces, to highlight the fold (represented as ribbons). One of the active sites is highlighted in red, showing contribution of three distinct subunits. The figures were generated using the program ICM (www.molsoft.com)

features observed are hypotonia, failure to thrive, severe psychomotor retardation, seizures, facial dysmorphism and brain malformations. Interestingly, whereas homozygous FH mutations predispose to fumaric aciduria, several heterozygous FH mutations are known to be involved in the autosomal dominant syndrome of multiple cutaneous and uterine leiomyomata (MCUL1) (Tomlinson et al. 2002). Affected individuals develop benign smooth muscle tumours of the skin, and females develop fibroids of the uterus. When co-existing with an aggressive form of renal cell carcinoma (papillary renal type II cancer or renal collecting duct cancer) it is also known as hereditary leiomyomatosis and renal cancer (HLRCC) syndrome. In MCUL1/HLRCC germline mutations in FH are detected in the majority of the screened cases. To date, 107 variants have been described, of which 93 are thought to be pathogenic (Bayley et al. 2008). The most common types are missense mutations (57%), followed by frameshift and nonsense mutations (27%), as well as diverse deletions, insertions and duplications.

Here we present the crystal structure of human fumarase at 1.95 Å resolution and summarize structure-activity correlation between observed mutations and clinical phenotypes.

Materials and methods

Expression, purification & crystallization

DNA fragment encoding the fumarase domain of human FH (aa 44–510; GenBank entry 19743875) was subcloned into





Table 2 Mutations observed in the human fumarase gene and association to disease. Abbreviations: CL: cutaneous leiomyoma; FHD: fumarate hydratase deficiency; HLRCC: hereditary leiomyomatosis and renal cell cancer; LCT: Leydig cell tumors; MCUL: multiple cutaneous and uterine leiomyomata; OMC: ovarian mucinous cystadenoma; RCC: renal cell carcinoma; STS: soft tissue sarcoma; UL: uterine leiomyomas; ULMS: uterine leiomyosarcoma

#	Mutation site	Mutated residue	Protein change	DNA change	Exon	Conservation	Localization	Reference	Phenotype
1	Arg51	Glu	R51E	c.152 G>A	2	Conserved	Surface	(Kiuru et al. 2002)	STS
2	Arg101	Pro	R101P	c.302 G>C	3	Semi-conserved	Surface	(Chan et al. 2005), (Heinritz et al. 2008)	HLRCC
3	Asn107	Thr	N107T	c.320A>C	3	Conserved	Active site	(Tomlinson et al. 2002), (Alam et al. 2005a), (Carvajal-Carmona et al. 2006)	MCUL, LCT
4	Ala117	Pro	A117P	c.349 G>C	3	Semi-conserved	Near active site	(Tomlinson et al. 2002)	MCUL
5	Leu132	Ser	L132S	c.395 T>C	4	Semi-conserved	Surface	(Wei et al. 2006)	HLRCC, reduced FH activity
6	His135	Arg	H135R	c.404A>G	4	Semi-conserved	Surface	(Chuang et al. 2005)	MCUL
7	Gln142	Lys	Q142K	c.424 C>A	4	Conserved	Near active site	(Badeloe et al. 2006)	MCUL
8	Ser158	Ile	S158I	c.473 G>T	4	Semi-conserved	Near active site	(Martinez-Mir et al. 2003)	MCUL
9	Arg160	Gly	R160G	c.478A>G	4	Conserved	Surface	(Wei et al. 2006)	MUCL, reduced FH activity
10	Pro174	Arg	P174R	c.521 C>G	4	Not conserved	Surface	(Alam et al. 2005b), (Zeng et al. 2006), (Pollard et al. 2005)	FHD
11	His180	Arg	H180R	c.539A>G	4	Semi-conserved	Active site	(Tomlinson et al. 2002), (Alam et al. 2005b)	MUCL
12	Gln185	Arg	Q185R	c.554A>G	4	Conserved	Active site	(Tomlinson et al. 2002)	MCUL
13	Ser187	Leu	S187L	c.560C>T	5	Conserved	Active site	(Toro et al. 2003)	MCUL
14	Asn188	Ser	N188S	c.563A>G	5	Conserved	Active site	(Toro et al. 2003)	MCUL
15	Pro192	Leu	P192L	c.575A>G	5	Conserved	In core helice	(Chuang et al. 2005)	MCUL
16	Met195	Thr	M195T	c.584 T>C	5	Conserved	In core helice	(Toro et al. 2003)	MCUL
17	His196	Arg	H196R	c.587A>G	5	Conserved	In core helice	(Kiuru et al. 2002), (Lehtonen et al. 2004)	RCC, ULMS
18	Ile229	Thr	I229T	c.686 T>C	5	Not conserved	Surface	(Alam et al. 2005b)	MCUL
19	Lys230	Arg	K230R	c.689A>G	5	Conserved	Subunit stabilization	(Tomlinson et al. 2002), (Coughlin et al. 1998), (Manning et al. 2000)	FHD
20	Arg233	Cys	R233C	c.697 C>T	5	Conserved	Active site	(Rustin et al. 1997), (Chuang et al. 2005), (Wei et al. 2006)	FHD, HLRCC, MCUL
21	Arg233	His	R233H	c.698 G>A	5	Conserved	Active site	(Tomlinson et al. 2002), (Alam et al. 2005b), (Wei et al. 2006), (Chuang et al. 2005), (Toro et al. 2003)	HLRCC, MCUL
22	Arg233	Leu	R233L	c.698 G>T	5	Conserved	Active site	(Chuang et al. 2005), (Toro et al. 2003)	MCUL
23	Ala239	Thr	A239T	c.715 G>A	5	Conserved	Near active site	(Lehtonen et al. 2004)	UL
24	Ala274	Thr	A274T	c.820 G>A	6	Not conserved	Active site	(Ylisaukko-oja et al. 2006)	OMC
25	Gly282	Val	G282V	c.845 G>T	6	Conserved	Active site	(Tomlinson et al. 2002), (Alam et al. 2005b)	MCUL
26	Ala308	Thr	A308T	c.922 G>A	7	Conserved	Surface	(Coughlin et al. 1998)	FHD
27	Asn310	Tyr	N310Y	c.928A>T	7	Conserved	Surface	(Alam et al. 2005b)	MCUL
28	Phe312	Cys	F312C	c.935 T>G	7	Conserved	Surface	(Coughlin et al. 1998)	FHD
29	His318	Tyr	H318Y	c.952 C>T	7	Semi-conserved	In core helice	(Toro et al. 2003), (Martinez-Mir et al. 2003)	HLRCC
30	His318	Leu	H318L	c.953A>T	7	Semi-conserved	In core helice	(Deschauer et al. 2006)	FHD
31	Val322	Asp	V322D	c.964 T>A	7	Conserved	In core helice (interaction with 1 other monomer)	(Toro et al. 2003)	MCUL
32	Thr330	Pro	T330P	c.988A>C	7	Semi-conserved	In core helice (interaction with 1 other monomer)	(Chuang et al. 2005)	MCUL
33	Cys333	Tyr	C333Y	c.998 G>A	7	Semi-conserved	In core helice (interaction with 1 other monomer)		MCUL
34	Ser334	Arg	S334R	c.1002 T>G	7	Conserved	In core helice (interaction with 1 other monomer)	(Badeloe et al. 2006)	CL
35	Leu335	Pro	L335P	c.1004 T>C	7	Conserved	In core helice	(Toro et al. 2003)	MCUL

Table 2 (continued)

#	Mutation site	Mutated residue	Protein change	DNA change	Exon	Conservation	Localization	Reference	Phenotype
36	Asn340	Lys	N340K	c.1020 T>A	7	Semi-conserved	In core helice	(Toro et al. 2003), (Wei et al. 2006)	MCUL
37	Glu355	Lys	E355K	c.1063 G>A	7	Conserved	Subunit stabilization	(Alam et al. 2005b)	MCUL
38	Asn361	Lys	N361K	c.1083 T>A	7	Conserved	Active site	(Alam et al. 2005b)	HLRCC-CDC
39	Glu362	Gln	E362Q	c.1084 G>C	7	Conserved	Active site	(Bourgeron et al. 1994)	FHD
40	Ser365	Gly	S365G	c.1093 G>A	7	Conserved	Active site	(Toro et al. 2003), (Wei et al. 2006)	MCUL
41	Ser366	Asn	S366N	c.1097 G>A	7	Conserved	Active site (but out)	(Toro et al. 2003), (Alam et al. 2005b)	MCUL
42	Met368	Thr	M368T	c.1103 T>C	7	Conserved	Active site	(Badeloe et al. 2006)	MCUL
43	Pro369	Ser	P369S	c.1105 C>T	7	Conserved	Active site (but out)	(Maradin et al. 2006)	FHD
44	Asn373	Ser	N373S	c.1118A>G	8	Conserved	Active site	(Lehtonen et al. 2004)	HLRCC/clear cell RCC
45	Gln376	Pro	Q376P	c.1127A>C	8	Conserved	In core helice (interaction with 1 other monomer)	(Zeman et al. 2000), (Remes et al. 2004), (Phillips et al. 2006)	FHD
46	Ala385	Asp	A385D	c.1154 C>A	8	Not conserved	In core helice (interaction with 2 other monomers)	(Wei et al. 2006)	MCUL
47	Val394	Leu	V394L	c.1180 G>C	8	Not conserved	In core helice	(Martinez-Mir et al. 2003)	MCUL
48	Gly397	Arg	G397R	c.1189 G>A	8	Semi- conserved	In core helice	(Alam et al. 2005b)	MCUL
49	His402	Cys	H402C	c.1207 C>T	8	Conserved	In core helice turn (interaction with 2 other monomers)	(Phillips et al. 2006)	FHD
50	Ser419	Pro	S419P	c.1255 T>C	9	Conserved	In core helice	(Wei et al. 2006)	HLRCC
51	Asp425	Val	D425V	c.1274A>T	9	Conserved	In core helice (interaction with 1 other monomer)	(Coughlin et al. 1998)	FHD
52	Gln439	Pro	Q439P	c.1316A>C	9	Not conserved	Surface	(Wei et al. 2006)	HLRCC
53	Met454	Ile	M454I	c.1362 G>A	9	Conserved	Subunit interaction	(Carvajal-Carmona et al. 2006)	LCT
54	Tyr465	Cys	Y465C	c.1394A>G	10	Semi- conserved	Surface	(Toro et al. 2003)	MCUL
55	Leu507	Pro	L507P	c.1520 T>C	10	Semi- conserved	Surface near opening active site	(Alam et al. 2005b)	MCUL

Fig. 2 Clustering of human fumarase missense mutations observed in FHD, MCUL1 and HLRCC. The active site is highlighted in cyan. Positions of amino acid mutations are indicated as small spheres and numbered according to Table 2. The positions around the active site are indicated in red, mutations affecting inter- or intrasubunit interactions are indicated in dark yellow. For clarity, one monomeric subunit is omitted



pNIC28-Bsa4 vector incorporating an N-terminal His₆-tag. The plasmid was transformed into BL21(DE3)-pRARE, cultured in Terrific Broth at 37°C, and induced with 0.1 mM IPTG overnight at 18°C. Cells were homogenized in lysis buffer (50 mM K-phosphate pH 7.5, 500 mM NaCl, 1 mM TCEP), centrifuged to remove cell debris, and the supernatant was purified by Nickel affinity (HisTrap Crude FF) and size exclusion (HiLoad 16/60 Superdex S200) chromatography. Purified protein was concentrated to 12.6 mg/ml and stored in 10 mM HEPES pH 7.5, 150 mM NaCl, 5% (w/v) glycerol and 0.5 mM TCEP at -80°C. Crystals were grown by vapour diffusion at 20°C in sitting drops mixing 150 nl protein and 150 nl reservoir solution containing 20% (w/v) PEG 3350, 0.2 M sodium acetate, 10% (w/v) ethylene glycol and 100 mM Bis-Tris propane pH 7.5. Crystals were cryo-protected in mother liquor containing 25% (w/v) glycerol and flash-frozen in liquid nitrogen.

Data collection & structure determination

Diffraction data to maximum resolution of 1.95 Å were collected on beamline X10A at the Swiss Light Source, and processed using the CCP4 Program suite (CCP4, 1994). FH crystallized in the trigonal space group P3₂21 ($a=180.5$ Å, $b=180.5$ Å, $c=114.6$ Å, $\alpha=90^\circ$, $\beta=90^\circ$, $\gamma=120^\circ$) with four molecules in the asymmetric unit. The structure of FH was solved by molecular replacement with PHASER (McCoy et al. 2005), using the yeast fumarase structure as search model (PDB code 1YFM). Initial automated model building was performed with ARP/wARP (Perrakis et al. 1999). This is followed by cycles of iterative manual model building using COOT (Emsley & Cowtan 2004) and restrained refinement using REFMAC5 with TLS parameters (Murshudov et al. 1997). The final structure was deposited in the Protein Data Bank (www.rcsb.org) under accession code 3E04 (Table 1).

Results and discussion

Fumarases are divided into two distinct groups. Class I fumarases are iron-dependent iron-sulfur cluster containing, dimeric enzymes, whereas the class II enzymes, including human and other eukaryotic fumarases, are homotetrameric enzymes with a molecular mass of about 200 kDa. Class II fumarases are evolutionarily highly conserved enzymes, e.g. the pairwise identity between *E. coli* and human fumarase is about 60%. Every monomer exhibits a typical tridomain structure, with a central domain involved in subunit interaction, thus forming a typical bundle comprised of 20 α -helices (Fig. 1A). Previous crystallographic analyses have revealed two distinct sites (A and B) in *E. coli* fumarase that can bind carboxylic acids. Site A is formed from three different monomer chains and likely to be the catalytic site,

whereas site B is thought to allosterically regulate activity (Rose and Weaver 2004).

A previous study correlated 27 distinct missense mutations to the *E. coli* fumarase structure (Alam 2005b), since then the list of mutations has doubled. To this end, 55 missense mutations in the human fumarase gene are now described. Here we correlate this updated list of mutations to fumarase deficiency, MCUL1 and HLRCC syndrome (Table 2) by using the human fumarase structure. Although not all of these novel mutations have been biochemically characterized, previous results suggest that FH activity is related to HLRCC (Alam 2005a), although other environmental or genetic factors likely play a role in the etiology of the disease. The clustering of mutational “hotspots” suggests enzyme activity relationships to phenotypic appearances. Figure 2 illustrates the clustering of FH mutations observed in FHD, MCUL1 and HLRCC. The large majority of mutations are located at evolutionarily highly conserved positions (Table 2) indicating that these mutations likely affect stability and/or activity of the enzyme. Two major clusters of mutations are observed; one is likely to affect structural integrity of the enzyme by interrupting inter or intrasubunit interactions (indicated in yellow in Fig. 2), whereas the other mutations are located around the active site and likely directly affect activity.

Acknowledgments Help in data collection at SLS (Swiss Light Source, Villigen, CH) by Frank von Delft, Annette Roos and Panagis Filippakopoulos is gratefully acknowledged. The Structural Genomics Consortium is a registered charity (Number 1097737) funded by the Canadian Institutes for Health Research, the Canadian Foundation for Innovation, Genome Canada through the Ontario Genomic institute, GlaxoSmithKline, Karolinska Institutet, the Knut and Alice Wallenberg Innovation, Merck and Co., Inc., the Novartis Research Foundation, the Swedish Agency for Innovation Systems, the Swedish Foundation for Strategic Research and the Wellcome Trust. The study was supported by the NIHR Oxford Biomedical Research Unit.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- Alam NA, Olpin S, Leigh IM (2005a) Fumarate hydratase mutations and predisposition to cutaneous leiomyomas, uterine leiomyomas and renal cancer. *Br J Dermatol* 153:11–17
- Alam NA, Olpin S, Rowan A, Kelsell D, Leigh IM, Tomlinson IP, Weaver T (2005b) Missense mutations in fumarate hydratase in multiple cutaneous and uterine leiomyomatosis and renal cell cancer. *J Mol Diagn* 7:437–443
- Badeloe S, van Geel M, van Steensel MA, Bastida J, Ferrando J, Steijlen PM, Frank J, Pobleto-Gutierrez P (2006) Diffuse and segmental variants of cutaneous leiomyomatosis: novel mutations in the fumarate hydratase gene and review of the literature. *Exp Dermatol* 15:735–741

- Bayley JP, Launonen V, Tomlinson IP (2008) The FH mutation database: an online database of fumarate hydratase mutations involved in the MCUL (HLRCC) tumor syndrome and congenital fumarase deficiency. *BMC Med Genet* 9:20
- Bourgeron T, Chretien D, Poggi-Bach J, Doonan S, Rabier D, Letouze P, Munnich A, Rotig A, Landrieu P, Rustin P (1994) Mutation of the fumarase gene in two siblings with progressive encephalopathy and fumarase deficiency. *J Clin Invest* 93:2514–2518
- Carvajal-Carmona LG, Alam NA, Pollard PJ, Jones AM, Barclay E, Wortham N, Pignatelli M, Freeman A, Pomplun S, Ellis I, Poulsom R, El-Bahrawy MA, Berney DM, Tomlinson IP (2006) Adult leydig cell tumors of the testis caused by germline fumarate hydratase mutations. *J Clin Endocrinol Metab* 91:3071–3075
- CCP4 (1994) The CCP4 suite: programs for protein crystallography. *Acta Crystallogr D Biol Crystallogr* 50:760–763
- Chan I, Wong T, Martinez-Mir A, Christiano AM, McGrath JA (2005) Familial multiple cutaneous and uterine leiomyomas associated with papillary renal cell cancer. *Clin Exp Dermatol* 30:75–78
- Chuang GS, Martinez-Mir A, Geyer A, Engler DE, Glaser B, Cserhalmi-Friedman PB, Gordon D, Horev L, Lukash B, Herman E, Cid MP, Brenner S, Landau M, Sprecher E, Garcia Muret MP, Christiano AM, Zlotogorski A (2005) Germline fumarate hydratase mutations and evidence for a founder mutation underlying multiple cutaneous and uterine leiomyomata. *J Am Acad Dermatol* 52:410–416
- Coughlin EM, Christensen E, Kunz PL, Krishnamoorthy KS, Walker V, Dennis NR, Chalmers RA, Elpeleg ON, Whelan D, Pollitt RJ, Ramesh V, Mandell R, Shih VE (1998) Molecular analysis and prenatal diagnosis of human fumarase deficiency. *Mol Genet Metab* 63:254–262
- Deschauer M, Gizatullina Z, Schulze A, Pritsch M, Knoppel C, Nape M, Zierz S, Gellerich FN (2006) Molecular and biochemical investigations in fumarase deficiency. *Mol Genet Metab* 88:146–152
- Emsley P, Cowtan K (2004) Coot: model-building tools for molecular graphics. *Acta Crystallogr D Biol Crystallogr* 60:2126–2132
- Heinritz W, Paasch U, Sticherling M, Wittekind C, Simon JC, Froster UG, Renner R (2008) Evidence for a founder effect of the germline fumarate hydratase gene mutation R58P causing hereditary leiomyomatosis and renal cell cancer (HLRCC). *Ann Hum Genet* 72:35–40
- Kerrigan JF, Aleck KA, Tarby TJ, Bird CR, Heidenreich RA (2000) Fumaric aciduria: clinical and imaging features. *Ann Neurol* 47:583–588
- King A, Selak MA, Gottlieb E (2006) Succinate dehydrogenase and fumarate hydratase: linking mitochondrial dysfunction and cancer. *Oncogene* 25:4675–4682
- Kiuru M, Lehtonen R, Arola J, Salovaara R, Jarvinen H, Aittomaki K, Sjoberg J, Visakorpi T, Knuutila S, Isola J, Delahunt B, Herva R, Launonen V, Karhu A, Aaltonen LA (2002) Few FH mutations in sporadic counterparts of tumor types observed in hereditary leiomyomatosis and renal cell cancer families. *Cancer Res* 62:4554–4557
- Lehtonen R, Kiuru M, Vanharanta S, Sjoberg J, Aaltonen LM, Aittomaki K, Arola J, Butzow R, Eng C, Husgafvel-Pursiainen K, Isola J, Jarvinen H, Koivisto P, Mecklin JP, Peltomaki P, Salovaara R, Wasenius VM, Karhu A, Launonen V, Nupponen NN, Aaltonen LA (2004) Biallelic inactivation of fumarate hydratase (FH) occurs in nonsyndromic uterine leiomyomas but is rare in other tumors. *Am J Pathol* 164:17–22
- McCoy AJ, Grosse-Kunstleve RW, Storoni LC, Read RJ (2005) Likelihood-enhanced fast translation functions. *Acta Crystallogr D Biol Crystallogr* 61:458–464
- Manning NJ, Olpin SE, Pollitt RJ, Downing M, Heeley AF, Young ID (2000) Fumarate hydratase deficiency: increased fumaric acid in amniotic fluid of two affected pregnancies. *J Inherit Metab Dis* 23:757–759
- Maradin M, Fumic K, Hansikova H, Tesarova M, Wenich L, Dorner S, Sarnavka V, Zeman J, Baric I (2006) Fumaricaciduria: mild phenotype in a 8-year-old girl with novel mutations. *J Inherit Metab Dis* 29:683
- Martinez-Mir A, Glaser B, Chuang GS, Horev L, Waldman A, Engler DE, Gordon D, Spelman LJ, Hatzibougias I, Green J, Christiano AM, Zlotogorski A (2003) Germline fumarate hydratase mutations in families with multiple cutaneous and uterine leiomyomata. *J Invest Dermatol* 121:741–744
- Murshudov GN, Vagin AA, Dodson EJ (1997) Refinement of macromolecular structures by the maximum-likelihood method. *Acta Crystallogr D Biol Crystallogr* 53:240–255
- Perrakis A, Morris R, Lamzin VS (1999) Automated protein model building combined with iterative structure refinement. *Nat Struct Biol* 6:458–463
- Phillips TM, Gibson JB, Ellison DA (2006) Fumarate hydratase deficiency in monozygotic twins. *Pediatr Neurol* 35:150–153
- Pollard PJ, Briere JJ, Alam NA, Barwell J, Barclay E, Wortham NC, Hunt T, Mitchell M, Olpin S, Moat SJ, Hargreaves IP, Heales SJ, Chung YL, Griffiths JR, Dagleish A, McGrath JA, Gleeson MJ, Hodgson SV, Poulsom R, Rustin P, Tomlinson IP (2005) Accumulation of Krebs cycle intermediates and over-expression of HIF1alpha in tumours which result from germline FH and SDH mutations. *Hum Mol Genet* 14:2231–2239
- Ratcliffe PJ (2007) Fumarate hydratase deficiency and cancer: activation of hypoxia signaling? *Cancer Cell* 11:303–305
- Remes AM, Filppula SA, Rantala H, Leisti J, Ruokonen A, Sharma S, Juffer AH, Hiltunen JK (2004) A novel mutation of the fumarase gene in a family with autosomal recessive fumarase deficiency. *J Mol Med* 82:550–554
- Rose IA, Weaver TM (2004) The role of the allosteric B site in the fumarase reaction. *Proc Natl Acad Sci U S A* 101:3393–3397
- Rustin P, Bourgeron T, Parfait B, Chretien D, Munnich A, Rotig A (1997) Inborn errors of the Krebs cycle: a group of unusual mitochondrial diseases in human. *Biochim Biophys Acta* 1361:185–197
- Tomlinson IP, Alam NA, Rowan AJ, Barclay E, Jaeger EE, Kelsell D, Leigh I, Gorman P, Lamlum H, Rahman S, Roylance RR, Olpin S, Bevan S, Barker K, Hearle N, Houlston RS, Kiuru M, Lehtonen R, Karhu A, Vilkki S, Laiho P, Eklund C, Vierimaa O, Aittomaki K, Hietala M, Sistonen P, Paetau A, Salovaara R, Herva R, Launonen V, Aaltonen LA (2002) Germline mutations in FH predispose to dominantly inherited uterine fibroids, skin leiomyomata and papillary renal cell cancer. *Nat Genet* 30:406–410
- Toro JR, Nickerson ML, Wei MH, Warren MB, Glenn GM, Turner ML, Stewart L, Duray P, Tourre O, Sharma N, Choyke P, Stratton P, Merino M, Walther MM, Linehan WM, Schmidt LS, Zbar B (2003) Mutations in the fumarate hydratase gene cause hereditary leiomyomatosis and renal cell cancer in families in North America. *Am J Hum Genet* 73:95–106
- Wei MH, Toure O, Glenn GM, Pithukpakorn M, Neckers L, Stolle C, Choyke P, Grubb R, Middleton L, Turner ML, Walther MM, Merino MJ, Zbar B, Linehan WM, Toro JR (2006) Novel mutations in FH and expansion of the spectrum of phenotypes expressed in families with hereditary leiomyomatosis and renal cell cancer. *J Med Genet* 43:18–27
- Ylisaukko-oja SK, Cybulski C, Lehtonen R, Kiuru M, Matyjasik J, Szymanska A, Szymanska-Pasternak J, Dyrskjot L, Butzow R, Orntoft TF, Launonen V, Lubinski J, Aaltonen LA (2006) Germline fumarate hydratase mutations in patients with ovarian mucinous cystadenoma. *Eur J Hum Genet* 14:880–883
- Zeman J, Krijt J, Stratilova L, Hansikova H, Wenich L, Kmoch S, Chastina P, Houstek J (2000) Abnormalities in succinylpurines in fumarase deficiency: possible role in pathogenesis of CNS impairment. *J Inherit Metab Dis* 23:371–374
- Zeng WQ, Gao H, Brueton L, Hutchin T, Gray G, Chakrapani A, Olpin S, Shih VE (2006) Fumarase deficiency caused by homozygous P131R mutation and paternal partial isodisomy of chromosome 1. *Am J Med Genet A* 140:1004–1009