

OPTIMAL ENERGY DECAY FOR THE WAVE-HEAT SYSTEM ON A RECTANGULAR DOMAIN*

CHARLES BATTY[†], LASSI PAUNONEN[‡], AND DAVID SEIFERT[†]

Abstract. We study the rate of energy decay for solutions of a coupled wave-heat system on a rectangular domain. Using techniques from the theory of C_0 -semigroups, and in particular a well-known result due to Borichev and Tomilov, we prove that the energy of classical solutions decays like $t^{-2/3}$ as $t \rightarrow \infty$. This rate is moreover shown to be sharp. Our result implies in particular that a general estimate in the literature, which predicts at least logarithmic decay and is known to be best possible in general, is suboptimal in the special case under consideration here. Our strategy of proof involves direct estimates based on separation of variables and a refined version of the technique developed in our earlier paper for a one-dimensional wave-heat system.

Key words. wave equation, heat equation, coupled, rectangular domain, energy, rates of decay, C_0 -semigroups, resolvent estimates

AMS subject classifications. Primary, 35M33, 35B40, 47D06; Secondary, 34K30, 37A25

DOI. 10.1137/18M1195796

1. Introduction. In this paper we study the rate of energy decay for solutions of a coupled wave-heat system on a rectangular domain. Let $\Omega_- = (-1, 0) \times (0, 1)$ and $\Omega_+ = (0, 1) \times (0, 1)$. We consider the following model:

$$(1.1) \quad \begin{cases} u_{tt}(x, y, t) = \Delta u(x, y, t), & (x, y) \in \Omega_-, \ t > 0, \\ w_t(x, y, t) = \Delta w(x, y, t), & (x, y) \in \Omega_+, \ t > 0, \\ u(-1, y, t) = w(1, y, t) = 0, & y \in (0, 1), \ t > 0, \\ u(x, 0, t) = u(x, 1, t) = 0, & x \in (-1, 0), \ t > 0, \\ w(x, 0, t) = w(x, 1, t) = 0, & x \in (0, 1), \ t > 0, \\ u_t(0, y, t) = w(0, y, t), & y \in (0, 1), \ t > 0, \\ u_x(0, y, t) = w_x(0, y, t), & y \in (0, 1), \ t > 0, \\ u(x, y, 0) = u_0(x, y), & (x, y) \in \Omega_-, \\ u_t(x, y, 0) = v_0(x, y), & (x, y) \in \Omega_-, \\ w(x, y, 0) = w_0(x, y), & (x, y) \in \Omega_+, \end{cases}$$

for suitable initial data u_0, v_0 defined on Ω_- and w_0 defined on Ω_+ . Given an initial vector $z_0 = (u_0, v_0, w_0)$ of initial data, the energy $E_{z_0}(t)$ at time $t \geq 0$ of the corresponding solution is defined as

$$E_{z_0}(t) = \frac{1}{2} \int_{\Omega} \left(|\nabla u(x, y, t)|^2 + |u_t(x, y, t)|^2 + |w(x, y, t)|^2 \right) d(x, y), \quad t \geq 0,$$

*Received by the editors June 20, 2018; accepted for publication (in revised form) January 15, 2019; published electronically March 26, 2019.

<http://www.siam.org/journals/sima/51-2/M119579.html>

Funding: The work of the second author was supported by Academy of Finland grants 298182 and 310489.

[†]St John's College, St Giles, Oxford OX1 3JP, UK (charles.batty@sjc.ox.ac.uk, david.seifert@sjc.ox.ac.uk).

[‡]Department of Mathematics, Tampere University of Technology, P.O. Box 553, 33101 Tampere, Finland (lassi.paunonen@tut.fi).

where u and w have been extended by zero to $\Omega = (-1, 1) \times (0, 1)$. For sufficiently regular solutions we see by a direct calculation that

$$\dot{E}_{z_0}(t) = - \int_{\Omega_+} |\nabla w(x, y, t)|^2 dx dy, \quad t \geq 0,$$

so the energy is non-increasing in time. The aim in this paper is to establish a sharp estimate for the *rate* at which the energy of sufficiently regular solutions decays to zero as $t \rightarrow \infty$.

Questions of this type, often for more general domains, have received a considerable amount of attention over the last decade or so, going back at least to [16]. The underlying motivation is to understand the physical phenomenon of *structure-fluid* interaction, for which the wave-heat model serves as a linear, and hence more tractable, approximate model retaining many of the key features of more realistic models. Some of the most general results in this area are obtained in [12, 19], and they show in particular that the rate of energy decay one can expect depends crucially on the geometry of the wave and heat domains. In fact, as is customary in the theory of damped wave equations, one may consider billiard ball trajectories beginning at points in the wave domain and reflecting according to the laws of optics along the Dirichlet part of the boundary. According to [12, 19], if there exist so-called trapped rays, that is to say, trajectories which never enter the heat domain, then one obtains a logarithmic rate of energy decay for classical solutions, and for general geometries this result is sharp. On the other hand, if every trajectory eventually enters the heat domain, then for classical solutions one obtains $E_{z_0}(t) = o(t^{-(2-\varepsilon)})$ as $t \rightarrow \infty$ for every $\varepsilon > 0$. In fact, it was shown recently in [5] using a delicate microlocal argument that if the heat domain completely surrounds the wave domain one may even take $\varepsilon = 0$, which gives the best possible estimate; see also [6]. Our main objective in the present paper is to obtain a sharp estimate for the rate of energy decay of classical solutions to our wave-heat system (1.1). Our result in particular illustrates that the theoretical logarithmic estimate for the rate of energy decay is in general not optimal in the case where, as in (1.1), we have a rectangular domain with trapped rays. Our main results concern the rate of energy decay of so-called classical solutions of (1.1), to be introduced formally in section 2 below, and may be summarized as follows.

THEOREM 1.1. *For classical solutions of the system (1.1) with initial data vector z_0 the energy satisfies $E_{z_0}(t) = o(t^{-2/3})$ as $t \rightarrow \infty$, and moreover this rate is sharp.*

Our approach is based on the theory of C_0 -semigroups, and in particular on the a result due to Borichev and Tomilov [10], which reduces the problem of determining the rate of energy decay to estimating the norm of the resolvent operator along the imaginary axis; see also [5, 6, 9]. Our argument divides naturally into three steps. First, in section 2, we show that the system in (1.1) is well posed in the sense of C_0 -semigroups and we describe the spectrum $\sigma(A)$ of the corresponding infinitesimal generator A , showing in particular that $\sigma(A)$ is contained in the open left-half plane. Then in section 3 we turn to estimating the resolvent of A along the imaginary axis. We first obtain an upper bound on the growth of the resolvent in Theorem 3.1 and then we prove in Theorem 3.3 that our upper bound is sharp. Finally, in section 4 we put together the pieces and apply the Borichev–Tomilov theorem to obtain a suitable form of Theorem 1.1. Throughout sections 2 and 3 we take advantage of the special rectangular geometry of our domain, which allows us to obtain optimal bounds by means of direct estimates as opposed, for instance, to indirect microlocal arguments. More specifically, we use separation of variables, thus decomposing the

two-dimensional problem into a family of one-dimensional problems which can be dealt with using techniques akin to those developed in our earlier paper [9]; see also [11, 18] for similar arguments in the context of damped wave equations.

Our notation is standard throughout. Given a closed operator A on a Hilbert space, which will always be assumed to be complex, we denote its domain by $D(A)$ and its kernel by $\text{Ker}(A)$. The spectrum of A is denoted by $\sigma(A)$, and given $\lambda \in \mathbb{C} \setminus \sigma(A)$ we write $R(\lambda, A)$ for the resolvent operator $(\lambda - A)^{-1}$. For real-valued quantities p and q , we use the notation $p \lesssim q$ to indicate that $p \leq Cq$ for some constant $C > 0$ which is independent of all the parameters that are free to vary in a given situation. We write $p \asymp q$ if $p \lesssim q$ and $q \lesssim p$. Furthermore, we make use where convenient of standard asymptotic notation such as “big O,” “little o,” and \sim . We let $\mathbb{N} = \{1, 2, 3, \dots\}$ and write \mathbb{C}_- for the open left half-plane $\{\lambda \in \mathbb{C} : \text{Re } \lambda < 0\}$.

2. Well-posedness. Our approach to studying (1.1) is based on the theory of C_0 -semigroups, so we first recast our system in the form of an abstract Cauchy problem. Let $\Gamma = \partial\Omega_- \cap \partial\Omega_+$, which is simply the vertical line segment $\{(0, y) : 0 \leq y \leq 1\}$, and let $H_\Gamma^1(\Omega_\mp)$ denote the set of functions in $H^1(\Omega_\mp)$ whose trace vanishes on $\partial\Omega_\mp \setminus \Gamma$, endowed with the equivalent Poincaré norm $\|u\| = \|\nabla u\|_{L^2}$, $u \in H_\Gamma^1(\Omega_\mp)$. We let Z denote the Hilbert space $H_\Gamma^1(\Omega_-) \times L^2(\Omega_-) \times L^2(\Omega_+)$, endowed with its natural inner product, and let Z_0 denote the subspace of Z consisting of all $(u, v, w) \in Z$ such that $\Delta u \in L^2(\Omega_-)$, $v \in H_\Gamma^1(\Omega_-)$, $w \in H_\Gamma^1(\Omega_+)$, and $\Delta w \in L^2(\Omega_+)$. Define the operator A by $A(u, v, w) = (v, \Delta u, \Delta w)$ for (u, v, w) in the domain

$$D(A) = \{(u, v, w) \in Z_0 : v|_\Gamma = w|_\Gamma \text{ and } u_x|_\Gamma = w_x|_\Gamma\}$$

of A , where the coupling conditions along Γ appearing in the definition of $D(A)$ are to be understood in the sense of traces. Then A is a closed and densely defined operator on Z but does not have compact resolvent; see [19, Theorem 2]. Letting $z(t) = (u(\cdot, t), u_t(\cdot, t), w(\cdot, t))$, $t \geq 0$, we may rewrite the wave-heat system (1.1) as

$$(2.1) \quad \begin{cases} \dot{z}(t) = Az(t), & t \geq 0, \\ z(0) = z_0, \end{cases}$$

where $z_0 \in Z$. Our first result, Theorem 2.1 below, establishes among other things that the operator A is the infinitesimal generator of a C_0 -semigroup $(T(t))_{t \geq 0}$ of contractions on Z . It follows that the unique solution of (2.1) is given by $z(t) = T(t)z_0$, $t \geq 0$. This solution in general satisfies (2.1) only in the so-called mild sense, but it is a solution in the classical sense if, and in fact only if, $z_0 \in D(A)$; see, for instance, [4] for details on the theory of C_0 -semigroups. It is for such classical solutions that we shall establish, in section 4 below, a sharp estimate on the rate of energy decay in the sense described in section 1. In what follows we choose the square root function with a branch cut along the negative real axis.

THEOREM 2.1. *The operator A generates a C_0 -semigroup $(T(t))_{t \geq 0}$ of contractions on Z and $\sigma(A) \subseteq \mathbb{C}_-$. Moreover, if we let $p_k(\lambda) = (k^2\pi^2 + \lambda^2)^{1/2}$ and $q_k(\lambda) = (k^2\pi^2 + \lambda)^{1/2}$ for $k \in \mathbb{N}$ and $\lambda \in \mathbb{C}$, then the point spectrum of A satisfies $\sigma_p(A) = \bigcup_{k \in \mathbb{N}} \Sigma_k$, where, for $k \in \mathbb{N}$,*

$$\Sigma_k = \left\{ \lambda \in \mathbb{C}_- : p_k(\lambda), q_k(\lambda) \neq 0 \text{ and } \lambda \frac{\tanh p_k(\lambda)}{p_k(\lambda)} + \frac{\tanh q_k(\lambda)}{q_k(\lambda)} = 0 \right\}.$$

Proof. For $z_0 = (u, v, w) \in D(A)$ a simple calculation using Green's theorem on each of the regions Ω_- and Ω_+ shows that

$$(2.2) \quad \operatorname{Re}\langle Az_0, z_0 \rangle = - \int_{\Omega_+} |\nabla w(x, y)|^2 dx dy \leq 0,$$

so the operator A is dissipative. Moreover, proceeding as in the proof of [19, Theorem 1] we see that A is invertible. Since the resolvent set is open we may find $\lambda > 0$ such that $\lambda \notin \sigma(A)$, so that $\lambda - A$ is invertible and in particular surjective. Hence A generates a C_0 -semigroup $(T(t))_{t \geq 0}$ of contractions on Z by the Lumer–Phillips theorem, and it follows from standard semigroup theory that $\sigma(A) \subseteq \{\lambda \in \mathbb{C} : \operatorname{Re} \lambda \leq 0\}$. In order to see that $\sigma(A)$ contains no purely imaginary points let $s \in \mathbb{R}$. We wish to show that for every $z_0 = (f, g, h) \in Z$ there exists $(u, v, w) \in D(A)$ such that $(is - A)(u, v, w) = z_0$ and $\|(u, v, w)\| \leq C\|z_0\|$ for some $C > 0$ which is independent of z_0 . Let $e_k(y) = \sqrt{2} \sin(k\pi y)$ for $k \in \mathbb{N}$ and $y \in (0, 1)$, recalling that $\{e_k : k \in \mathbb{N}\}$ is an orthonormal basis for $L^2(0, 1)$. We may expand u into a series of the form

$$u(x, y) = \sum_{k=1}^{\infty} u_k(x) e_k(y), \quad (x, y) \in \Omega_-,$$

with convergence in the norm of $L^2(\Omega_-)$, and we may similarly decompose v, w, f, g , and h . This gives rise to functions $u_k \in H^2(-1, 0) \cap H^1_-(1, 0)$, $v_k \in H^1_-(1, 0)$, $w_k \in H^2(0, 1) \cap H^1_+(0, 1)$, $f_k \in H^1_-(1, 0)$, $g_k \in L^2(-1, 0)$, and $h_k \in L^2(0, 1)$, $k \in \mathbb{N}$, where $H^1_-(1, 0) = \{u_0 \in H^1(-1, 0) : u_0(-1) = 0\}$ and $H^1_+(0, 1) = \{w_0 \in H^1(0, 1) : w_0(1) = 0\}$. Using orthonormality of the set $\{e_k : k \in \mathbb{N}\}$, our problem turns into the system of one-dimensional equations

$$(2.3) \quad \begin{cases} v_k(x) = is u_k(x) - f_k(x), & x \in (-1, 0), \\ u_k''(x) = (k^2\pi^2 - s^2)u_k(x) - is f_k(x) - g_k(s), & x \in (-1, 0), \\ w_k''(x) = (k^2\pi^2 + is)w_k(x) - h_k(x), & x \in (0, 1), \end{cases}$$

with the boundary conditions $u_k(-1) = w_k(1) = 0$, $v_k(0) = w_k(0)$, and $u_k'(0) = w_k'(0)$, $k \in \mathbb{N}$. These equations can be solved for each $k \in \mathbb{N}$, and in fact we shall do so explicitly in the proof of Theorem 3.1 below. In order to show that $\|(u, v, w)\| \leq C\|z_0\|$ for some $C > 0$ we note first that

$$\|z_0\|^2 = \sum_{k=1}^{\infty} \left(k^2\pi^2 \|f_k\|_{L^2}^2 + \|f_k'\|_{L^2}^2 + \|g_k\|_{L^2}^2 + \|h_k\|_{L^2}^2 \right),$$

and we obtain an analogous expression for the norm of (u, v, w) . Let us write $z_k = (f_k, g_k, h_k)$ and

$$\|z_k\| = \left(k^2\pi^2 \|f_k\|_{L^2}^2 + \|f_k'\|_{L^2}^2 + \|g_k\|_{L^2}^2 + \|h_k\|_{L^2}^2 \right)^{1/2}, \quad k \in \mathbb{N}.$$

It suffices to show that

$$(2.4) \quad k\|u_k\|_{L^2}, \|u_k'\|_{L^2}, \|v_k\|_{L^2}, \|w_k\|_{L^2} \lesssim \|z_k\|, \quad k \in \mathbb{N},$$

where the implicit constant is independent of z_0 and k . Since we know that the resolvent set contains a neighborhood of zero we may assume that $|s| \geq s_0$ for some $s_0 > 0$. We omit this argument here, since a more careful version of it is presented

in the proof of Theorem 3.1 below, where we moreover keep track of how the implicit constant in (2.4) depends on $|s|$.

It remains to describe the point spectrum $\sigma_p(A)$ of A . Let $\lambda \in \mathbb{C}_-$ and suppose that $z_0 = (u, v, w) \in \text{Ker}(\lambda - A)$. Expanding the components of z_0 as above we obtain the system

$$(2.5) \quad \begin{cases} v_k(x) = \lambda u_k(x), & x \in (-1, 0), \\ u_k''(x) = (k^2\pi^2 + \lambda^2)u_k(x), & x \in (-1, 0), \\ w_k''(x) = (k^2\pi^2 + \lambda)w_k(x), & x \in (0, 1), \end{cases}$$

together with the boundary conditions $u_k(-1) = w_k(1) = 0$, $v_k(0) = w_k(0)$, and $u_k'(0) = w_k'(0)$, $k \in \mathbb{N}$. Since $\lambda \in \mathbb{C}_-$ we have $p_k(\lambda) \neq 0$ for all $k \in \mathbb{N}$. Let us assume for the moment that we also have $q_k(\lambda) \neq 0$, $k \in \mathbb{N}$. It is then straightforward to show that the ordinary differential equations in (2.5) together with the boundary conditions at $x = \pm 1$ imply

$$\begin{aligned} u_k(x) &= a_k(\lambda) \sinh(p_k(\lambda)(1+x)), & x \in (-1, 0), \\ w_k(x) &= b_k(\lambda) \sinh(q_k(\lambda)(1-x)), & x \in (0, 1), \end{aligned}$$

for some constants $a_k(\lambda), b_k(\lambda) \in \mathbb{C}$, $k \in \mathbb{N}$, and now the coupling conditions at $x = 0$ can be formulated as

$$(2.6) \quad \begin{pmatrix} \lambda \sinh p_k(\lambda) & -\sinh q_k(\lambda) \\ p_k(\lambda) \cosh p_k(\lambda) & q_k(\lambda) \cosh q_k(\lambda) \end{pmatrix} \begin{pmatrix} a_k(\lambda) \\ b_k(\lambda) \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \quad k \in \mathbb{N}.$$

Writing $M_k(\lambda)$ for the 2×2 matrix appearing in this equation we see that $z_0 \neq 0$ if and only if $\det M_k(\lambda) = 0$ for some $k \in \mathbb{N}$. On the other hand, if $q_k(\lambda) = 0$ for some $k \in \mathbb{N}$, then $w_k(x) = b_k(\lambda)(x-1)$, $x \in (0, 1)$, for some constant $b_k(\lambda) \in \mathbb{C}$, and it is straightforward to verify that the coupling conditions at $x = 0$ imply $a_k(\lambda) = b_k(\lambda) = 0$. The result now follows. \square

3. Resolvent estimates. We now study the growth behavior of the resolvent norms $\|R(is, A)\|$ as $|s| \rightarrow \infty$. First, in section 3.1 we obtain an upper bound for the resolvent norms, which we then prove to be optimal in section 3.2. These results will eventually allow us to obtain a sharp estimate on the rate of energy decay for classical solutions, which is done in section 4 below.

3.1. An upper bound. We establish the following result.

THEOREM 3.1. *We have $\|R(is, A)\| = O(|s|^3)$ as $|s| \rightarrow \infty$.*

Proof. We use the notation introduced in section 2. Fix $z_0 \in Z$ and $s \in \mathbb{R}$ with $|s| \geq s_0$, where $s_0 > 0$ is to be chosen in due course. Let us write $z_0 = (f, g, h)$ and $R(is, A)z_0 = (u, v, w)$, so that $z_0 = (is - A)(u, v, w)$. We may decompose each of the entries of z_0 and $R(is, A)z_0$ as in the proof of Theorem 2.1 to obtain the system (2.3) with the boundary conditions listed there. Using the same notation as in the proof of Theorem 2.1 but dropping the subscript L^2 from now on, we have $\|v_k\| \lesssim |s|\|u_k\| + \|z_k\|$, so our objective is to show that

$$k\|u_k\|, |s|\|u_k\|, \|u_k'\|, \|w_k\| \lesssim |s|^3\|z_k\|, \quad k \in \mathbb{N},$$

where the implicit constant is independent of z_0 , s , and k . Our approach depends crucially on the relationship between the parameters s and k , and we distinguish

between two main cases. In what follows we write $p_k(s)$ for $p_k(\lambda)$ when $\lambda = is$, and similarly for q_k .

Case 1. $s^2 \leq k^2\pi^2 + 1$. We begin with some preliminary maneuvers. Taking the inner product in $L^2(-1, 0)$ of the second equation of (2.3) with v_k and integrating by parts we obtain

$$(3.1) \quad \langle u'_k, v'_k \rangle - u'_k(0)\overline{v_k(0)} = \langle is f_k + g_k - p_k(s)^2 u_k, v_k \rangle.$$

Similarly, taking the inner product in $L^2(0, 1)$ of the third equation of (2.3) with w_k and integrating by parts we obtain

$$(3.2) \quad \|w'_k\|^2 + w'_k(0)\overline{w_k(0)} = \langle h_k - q_k(s)^2 w_k, w_k \rangle.$$

Adding (3.1) and (3.2) we find, after using the boundary conditions, that

$$(3.3) \quad \langle u'_k, v'_k \rangle + \|w'_k\|^2 = \langle is f_k + g_k - p_k(s)^2 u_k, v_k \rangle + \langle h_k - q_k(s)^2 w_k, w_k \rangle.$$

Expanding this equation using the identity $v_k = is u_k - f_k$ and then taking real parts we find, after some crude estimates, that

$$(3.4) \quad \|w_k\|^2 \lesssim \|u'_k\| \|z_k\| + \|z_k\|^2.$$

Here we have used that $\|u_k\| \leq \frac{2}{\pi} \|u'_k\|$ by the Poincaré inequality and also that $|s| \lesssim k$. Similarly, if we take imaginary parts in (3.3) and make use of (3.4), then after some standard estimates we obtain

$$(3.5) \quad \|u'_k\|^2 + p_k(s)^2 \|u_k\|^2 \lesssim \|u'_k\| \|z_k\| + \|z_k\|^2.$$

Here we have used that $|s| \geq s_0$ for some $s_0 > 0$. From the Poincaré inequality and the fact that $p_k(s)^2 \geq -1$ we see that $\|u'_k\|^2 \lesssim \|u'_k\|^2 + p_k(s)^2 \|u_k\|^2$, so by (3.5) we may find a constant $C > 0$ such that

$$\|u'_k\|^2 - C \|u'_k\| \|z_k\| - C^2 \|z_k\|^2 \leq 0.$$

By computing the roots of the polynomial $t \mapsto t^2 - C \|z_k\| t - C^2 \|z_k\|^2$ it follows easily that $\|u'_k\| \lesssim \|z_k\|$. By (3.4) we also have $\|w_k\| \lesssim \|z_k\|$. Next we estimate $k \|u_k\|$. Supposing for the moment that $p_k(s)^2 \geq k^2$ we have $k \|u_k\| \lesssim \|z_k\|$ from (3.5) and the estimate for $\|u'_k\|$. On the other hand, if $p_k(s)^2 < k^2$, then $k \lesssim |s|$ and hence by the Poincaré inequality $k \|u_k\| \lesssim |s| \|u'_k\| \lesssim |s| \|z_k\|$. Finally, we have $|s| \|u_k\| \lesssim k \|u_k\| \lesssim |s| \|z_k\|$.

Case 2. $s^2 > k^2\pi^2 + 1$. Note that subject to the boundary conditions at $x = \pm 1$ the solutions u_k, w_k of (2.3) are given by

$$(3.6) \quad \begin{aligned} u_k(x) &= a_k(s) \sinh(p_k(s)(x+1)) + U_{k,s}(x), & x \in (-1, 0), \\ w_k(x) &= b_k(s) \sinh(q_k(s)(1-x)) + W_{k,s}(x), & x \in (0, 1), \end{aligned}$$

where $a_k(s), b_k(s) \in \mathbb{C}$, $k \in \mathbb{N}$, are constants and

$$\begin{aligned} U_{k,s}(x) &= -\frac{1}{p_k(s)} \int_{-1}^x \sinh(p_k(s)(x-r)) H_{k,s}(r) dr, & x \in (-1, 0), \\ W_{k,s}(x) &= -\frac{1}{q_k(s)} \int_x^1 \sinh(q_k(s)(r-x)) h_k(r) dr, & x \in (0, 1), \end{aligned}$$

with $H_{k,s}(x) = is f_k(x) + g_k(x)$, $x \in (-1, 0)$. If we denote the 2×2 matrix appearing in (2.6) by $M_k(s)$ when $\lambda = is$, then we may write the coupling conditions at $x = 0$ in the form

$$M_k(s) \begin{pmatrix} a_k(s) \\ b_k(s) \end{pmatrix} = \begin{pmatrix} f_k(0) - is U_{k,s}(0) + W_{k,s}(0) \\ W'_{k,s}(0) - U'_{k,s}(0) \end{pmatrix}, \quad k \in \mathbb{N}.$$

Hence the solutions u_k and w_k , $k \in \mathbb{N}$, may be written, after some elementary but tedious manipulations, in the form

$$\begin{aligned} u_k(x) = & \frac{is q_k(s) \cosh q_k(s)}{p_k(s) \det M_k(s)} \left(\frac{p_k(s)}{is} \sinh(p_k(s)(x+1)) f_k(0) \right. \\ & - \sinh(p_k(s)x) \int_{-1}^x \sinh(p_k(s)(1+r)) H_{k,s}(r) dr \\ & - \sinh(p_k(s)(1+x)) \int_x^0 \sinh(p_k(s)r) H_{k,s}(r) dr \Big) \\ & + \frac{\sinh q_k(s)}{\det M_k(s)} \left(\cosh(p_k(s)x) \int_{-1}^x \sinh(p_k(s)(1+r)) H_{k,s}(r) dr \right. \\ & + \sinh(p_k(s)(1+x)) \int_x^0 \cosh(p_k(s)r) H_{k,s}(r) dr \Big) \\ & + \frac{1}{\det M_k(s)} \sinh(p_k(s)(x+1)) \int_0^1 \sinh(q_k(s)(1-r)) h_k(r) dr \end{aligned}$$

for $x \in (-1, 0)$ and

$$\begin{aligned} w_k(x) = & \frac{p_k(s) \cosh p_k(s)}{q_k(s) \det M_k(s)} \left(-q_k(s) \sinh(q_k(s)(1-x)) f_k(0) \right. \\ & + \sinh(q_k(s)(1-x)) \int_0^x \sinh(q_k(s)r) h_k(r) dr \\ & + \sinh(q_k(s)x) \int_x^1 \sinh(q_k(s)(1-r)) h_k(r) dr \Big) \\ & + \frac{is \sinh p_k(s)}{\det M_k(s)} \left(\sinh(q_k(s)(1-x)) \int_0^x \cosh(q_k(s)r) h_k(r) dr \right. \\ & + \cosh(q_k(s)x) \int_x^1 \sinh(q_k(s)(1-r)) h_k(r) dr \Big) \\ & + \frac{is}{\det M_k(s)} \sinh(q_k(s)(1-x)) \int_{-1}^0 \sinh(p_k(s)(1+r)) H_{k,s}(r) dr \end{aligned}$$

for $x \in (0, 1)$. Though somewhat laborious to derive, these formulas can be readily verified simply by substituting them into (2.3). Since $s^2 > k^2 \pi^2 + 1$, we have $k \lesssim |s|$ and hence $1 \leq |p_k(s)| \lesssim |s|$ and $|s|^{1/2} \lesssim |q_k(s)| \lesssim |s|$. Moreover, $p_k(s)$ is purely imaginary. Hence if we differentiate the expression for u_k and estimate the L^2 -norms crudely, and in particular insert a factor of $|s|/k$ for later convenience, we obtain

$$\begin{aligned} (3.7) \quad k \|u_k\|, |s| \|u_k\|, \|u'_k\| & \lesssim \frac{s^4 e^{\operatorname{Re} q_k(s)}}{k |p_k(s)| |\det M_k(s)|} \|z_k\| \\ \text{and} \quad \|w_k\| & \lesssim \frac{s^2 e^{\operatorname{Re} q_k(s)}}{k |\det M_k(s)|} \|z_k\|. \end{aligned}$$

Note that $\operatorname{Re} q_k(s) > 0$. Next we seek to bound the term $e^{-\operatorname{Re} q_k(s)} |\det M_k(s)|$ from below. A straightforward calculation yields

$$(3.8) \quad 2 \frac{|\det M_k(s)|}{e^{\operatorname{Re} q_k(s)}} \geq |sq_k(s) \sin |p_k(s)| - i|p_k(s)| \cos |p_k(s)|| \\ - e^{-2\operatorname{Re} q_k(s)} |sq_k(s) \sin |p_k(s)| + i|p_k(s)| \cos |p_k(s)||,$$

and squaring the modulus of first term on the right-hand side we obtain

$$s^2 (\operatorname{Re} q_k(s))^2 \sin^2 |p_k(s)| + (s \operatorname{Im} q_k(s) \sin |p_k(s)| - |p_k(s)| \cos |p_k(s)|)^2.$$

If the first summand in this expression is less than $1/8$, then

$$|p_k(s)|^2 \cos^2 |p_k(s)| \geq 1 - \frac{1}{8s^2 (\operatorname{Re} q_k(s))^2} \geq \frac{1}{2}$$

because $|s| \operatorname{Re} q_k(s) \geq 1/2$, and since $|\operatorname{Im} q_k(s)| \leq \operatorname{Re} q_k(s)$ it follows easily that the second summand in the expression must be at least $1/8$. Hence the first term on the right-hand side of (3.8) is bounded from below by $1/2\sqrt{2}$. The second term on the right-hand side of (3.8) converges to zero uniformly in k as $|s| \rightarrow \infty$, so if we choose the lower bound $s_0 \geq 1$ for $|s|$ to be sufficiently large we see that $e^{-\operatorname{Re} q_k(s)} |\det M_k(s)|$ is uniformly bounded away from zero. Thus (3.7) becomes

$$k\|u_k\|, |s|\|u_k\|, \|u'_k\| \lesssim \frac{s^4}{k|p_k(s)|} \|z_k\| \quad \text{and} \quad \|w_k\| \lesssim \frac{s^2}{k} \|z_k\|.$$

Now if $s^2 \leq 2k^2\pi^2$, then $|s| \lesssim k$ and hence

$$k\|u_k\|, |s|\|u_k\|, \|u'_k\| \lesssim |s|^3 \|z_k\| \quad \text{and} \quad \|w_k\| \lesssim |s| \|z_k\|.$$

On the other hand, if $s^2 > 2k^2\pi^2$, then $|s| \lesssim |p_k(s)|$ and therefore

$$(3.9) \quad k\|u_k\|, |s|\|u_k\|, \|u'_k\| \lesssim |s|^3 \|z_k\| \quad \text{and} \quad \|w_k\| \lesssim s^2 \|z_k\|.$$

This completes the proof. \square

Remark 3.2. (a) The explicit solutions to (2.3) given in the proof work for all $s \in \mathbb{R} \setminus \{0\}$. Hence in proving Theorem 2.1, where we only need to show that the inverse of $is - A$ is a bounded operator for each fixed $s \in \mathbb{R} \setminus \{0\}$, we may use these explicit solutions to obtain crude bounds for all $k \in \mathbb{N}$ such that $k^2\pi^2 + 1 < s^2$ and then obtain (2.4) by using only Case 1 of the above proof, which crucially does not require the lower bound s_0 for $|s|$ to be large.

(b) In the above proof we have not attempted to obtain the best possible growth estimate in each of the subcases. For instance, a more careful analysis would show that for $s^2 > 2k^2\pi^2$ one can improve (3.9) to

$$k\|u_k\|, |s|\|u_k\|, \|u'_k\| \lesssim s^2 \|z_k\| \quad \text{and} \quad \|w_k\| \lesssim |s| \|z_k\|.$$

As we shall see now, however, the overall growth rate of $|s|^3$ is sharp. This suggests that the largest contributions come from terms for which $k \asymp |s|$; see Remark 3.4 below.

3.2. Optimality of the upper bound. We now show that the upper bound obtained in Theorem 3.1 is sharp by proving that A has eigenvalues which approach the imaginary axis at a suitable rate at infinity.

THEOREM 3.3. *There exist sequences (λ_k^\pm) of eigenvalues of A and a constant $C > 0$ such that $\operatorname{Im} \lambda_k^\pm \sim \pm k\pi$ as $k \rightarrow \infty$ and*

$$(3.10) \quad -\frac{C}{|\operatorname{Im} \lambda_k^\pm|^3} \leq \operatorname{Re} \lambda_k^\pm < 0, \quad k \in \mathbb{N}.$$

In particular,

$$\limsup_{|s| \rightarrow \infty} \frac{\|R(is, A)\|}{|s|^3} > 0.$$

Proof. Consider the functions $F_k(\lambda) = \tanh p_k(\lambda)$ and

$$G_k(\lambda) = \frac{p_k(\lambda)}{\lambda q_k(\lambda)} \tanh q_k(\lambda), \quad \lambda \in \mathbb{C},$$

where p_k, q_k are as in Theorem 2.1, and let $\mu_k^\pm = \pm i\pi\sqrt{k^2 + 1}$, $k \in \mathbb{N}$. Note that $p_k(\mu_k^\pm) = i\pi$ and let $q_k^\pm = q_k(\mu_k^\pm)$, $k \in \mathbb{N}$. Then $F_k(\mu_k^\pm) = 0$ and

$$G_k(\mu_k^\pm) = \pm \frac{\tanh q_k^\pm}{q_k^\pm \sqrt{k^2 + 1}}, \quad k \in \mathbb{N}.$$

In particular, since $|q_k^\pm| \sim k\pi$ and $|\tanh q_k^\pm| \rightarrow 1$ as $k \rightarrow \infty$ we have $|G_k(\mu_k^\pm)| \sim (k^2\pi)^{-1}$ as $k \rightarrow \infty$. In fact, if we let $\Omega_k^\pm = \{\lambda \in \mathbb{C} : |\lambda - \mu_k^\pm| < k^{-3}\}$, $k \in \mathbb{N}$, a more careful argument using Taylor expansions shows that

$$(3.11) \quad \sup \{|G_k(\lambda)| : \lambda \in \partial\Omega_k^\pm\} \sim \frac{1}{k^2\pi}, \quad k \rightarrow \infty.$$

For $\lambda = \mu_k^\pm + z$ we have

$$F_k(\lambda) = i \tan \left(\pi \left(1 - \frac{z^2 \pm 2zi\pi\sqrt{k^2 + 1}}{\pi^2} \right)^{1/2} \right)$$

and hence another Taylor expansion shows that $|F_k(\lambda)| \geq (2k^2)^{-1}$ for $\lambda \in \partial\Omega_k^\pm$ provided k is sufficiently large. Thus by (3.11) we have $|G_k(\lambda)| < |F_k(\lambda)|$ for such values of λ and k , so Rouché's theorem implies that the function $F_k + G_k$ has roots $\lambda_k^\pm \in \Omega_k^\pm$ when k is sufficiently large. By Theorem 2.1 any root of $F_k + G_k$ is an eigenvalue of A , so we obtain (3.10). The final claim follows easily since $\|R(is, A)\| \geq \operatorname{dist}(is, \sigma(A))^{-1}$, $s \in \mathbb{R}$. \square

Remark 3.4. An alternative approach to proving optimality of the resolvent bound in Theorem 3.1 is to show directly, by estimating $\|R(is, A)z_0\|$ from below for suitable $s \in \mathbb{R}$ and $z_0 \in Z$, that the upper bounds in the latter part of the proof of that result cannot be improved. Note also that the points on the imaginary axis at which the resolvent norm is shown to be large are of the form $\pm is_k$, $k \in \mathbb{N}$, where $s_k \sim k\pi$ as $k \rightarrow \infty$; see Remark 3.2.

4. Energy decay. We now turn to the rate of energy decay for classical solutions of the abstract Cauchy problem (2.1) corresponding to the wave-heat system (1.1). Our main result, Theorem 4.2 below, is a consequence of the following abstract result on rates of decay for semigroups on Hilbert space due to Borichev and Tomilov [10]; see also [7, 17].

THEOREM 4.1. *Let Z be a Hilbert space and let $(T(t))_{t \geq 0}$ be a bounded C_0 -semigroup on Z with generator A . Suppose that $\sigma(A) \subseteq \mathbb{C}_-$. Then for any constant $\alpha > 0$ the following conditions are equivalent:*

- (i) $\|R(is, A)\| = O(|s|^\alpha)$ as $|s| \rightarrow \infty$;
- (ii) $\|T(t)A^{-1}\| = O(t^{-1/\alpha})$ as $t \rightarrow \infty$;
- (iii) $\|T(t)z_0\| = o(t^{-1/\alpha})$ as $t \rightarrow \infty$ for all $z_0 \in D(A)$.

We now come to the main result of this paper.

THEOREM 4.2. *For $z_0 \in D(A)$ the energy of the classical solution of (2.1) satisfies $E_{z_0}(t) = o(t^{-2/3})$ as $t \rightarrow \infty$.*

Proof. Note that for any $z_0 \in Z$ we have $E_{z_0}(t) = \frac{1}{2}\|T(t)z_0\|^2$, $t \geq 0$. Since $\|R(is, A)\| = O(|s|^3)$ as $|s| \rightarrow \infty$ by Theorem 3.1 it follows from Theorem 4.1 that $E_{z_0}(t) = o(t^{-2/3})$ as $t \rightarrow \infty$ for $z_0 \in D(A)$. \square

- Remark 4.3.* (a) The rate $t^{-2/3}$ in Theorem 4.2 is optimal in the sense that, given any positive function r satisfying $r(t) = o(t^{-2/3})$ as $t \rightarrow \infty$, there exists $z_0 \in D(A)$ such that $E_{z_0}(t) \neq o(r(t))$ as $t \rightarrow \infty$. This follows from Theorem 3.3 and the uniform boundedness principle together with [8, Proposition 1.3]; see also [4, Theorem 4.4.14].
- (b) It follows from Theorem 3.3 and standard C_0 -semigroup theory that there is no hope of finding a rate of energy decay which is valid for *all* initial values $z_0 \in Z$; see also [15, Lemma 3.1.7]. On the other hand, if $z_0 \in D(A^k)$ for some $k \in \mathbb{N}$, then it follows easily from the semigroup property that $E_{z_0}(t) = o(t^{-2k/3})$ as $t \rightarrow \infty$, so more regular solutions have faster energy decay.

We conclude by mentioning that methods similar to the ones presented here can be used to obtain sharp estimates for the rate of energy decay in various related problems, such as system (1.1) but with the coupling condition $u_t(0, y, t) = w(0, y, t)$ replaced by $u(0, y, t) = w(0, y, t)$ for $y \in (0, 1)$, $t > 0$. Another example is the following wave equation which is damped on one half of its rectangular domain but not on the other:

$$\begin{cases} u_{tt}(x, y, t) + \mathbb{1}_{\Omega_+}(x, y)u_t(x, y, t) = \Delta u(x, y, t), & (x, y) \in \Omega, \ t > 0, \\ u(x, y, t) = 0, & (x, y) \in \partial\Omega, \ t > 0, \\ u(x, y, 0) = u_0(x, y), \ u_t(x, y, 0) = v_0(x, y), & (x, y) \in \Omega. \end{cases}$$

Here $\Omega = (-1, 1) \times (0, 1)$ and $\Omega_+ = (0, 1) \times (0, 1)$ as in section 1, and u_0, v_0 are suitable functions defined on Ω . In this case the energy

$$E_{z_0}(t) = \frac{1}{2} \int_{\Omega} (|\nabla u(x, y, t)|^2 + |u_t(x, y, t)|^2) \, d(x, y), \quad t \geq 0,$$

of any classical solution u , with corresponding initial data $z_0 = (u_0, v_0)$, can be shown to satisfy $E_{z_0}(t) = o(t^{-4/3})$ as $t \rightarrow \infty$, and furthermore this estimate is sharp; see [3, Part IV.B], [18], and also [11, 14]. Our methods can also be adapted to study

the following wave equation on the square $\Omega_- = (-1, 0) \times (0, 1)$ subject to Dirichlet boundary conditions along three of its edges but with the coupled heat equation in (1.1) replaced by a dissipative boundary condition along the fourth edge:

$$(4.1) \quad \begin{cases} u_{tt}(x, y, t) = \Delta u(x, y, t), & (x, y) \in \Omega_-, \ t > 0, \\ u(x, 0, t) = u(x, 1, t) = 0, & x \in (-1, 0), \ t > 0, \\ u(-1, y, t) = 0, & y \in (0, 1), \ t > 0, \\ u_x(0, y, t) = -\kappa u_t(0, y, t), & y \in (0, 1), \ t > 0, \\ u(x, y, 0) = u_0(x, y), & (x, y) \in \Omega_-, \\ u_t(x, y, 0) = v_0(x, y), & (x, y) \in \Omega_-, \end{cases}$$

for suitable initial data u_0, v_0 defined on Ω_- and any constant $\kappa > 0$. Models of this type are considered, for instance, in [1, 2, 13]. By formulating (4.1) as an abstract Cauchy problem and proceeding as in sections 2 and 3 one obtains the sharp estimate $\|R(is, A)\| = O(|s|^2)$, $|s| \rightarrow \infty$, for the resolvent of the generator A of the corresponding contraction semigroup. It follows as in Theorem 4.2 that the energy of any classical solution u of (4.1) decays like $o(t^{-1})$ as $t \rightarrow \infty$, and this estimate too is optimal.

Acknowledgments. This work was carried out while L. Paunonen visited Oxford from January to June 2017 and while D. Seifert visited Tampere University of Technology in September 2017. All three authors would like to express their gratitude to the two anonymous referees, whose comments improved the paper.

REFERENCES

- [1] Z. ABBAS AND S. NICAISE, *The multidimensional wave equation with generalized acoustic boundary conditions I: Strong stability*, SIAM J. Control Optim., 53 (2015), pp. 2558–2581.
- [2] Z. ABBAS AND S. NICAISE, *The multidimensional wave equation with generalized acoustic boundary conditions II: Polynomial stability*, SIAM J. Control Optim., 53 (2015), pp. 2582–2607.
- [3] N. ANANTHARAMAN AND M. LÉAUTAUD, *Sharp polynomial decay rates for the damped wave equation on the torus*, Anal. PDE, 7 (2014), pp. 159–214.
- [4] W. ARENDT, C. J. K. BATTY, M. HIEBER, AND F. NEUBRANDER, *Vector-Valued Laplace Transforms and Cauchy Problems*, 2nd ed., Birkhäuser, Basel, 2011.
- [5] G. AVALOS, I. LASIECKA, AND R. TRIGGIANI, *Heat-wave interaction in 2–3 dimensions: optimal rational decay rate*, J. Math. Anal. Appl., 437 (2016), pp. 782–815.
- [6] G. AVALOS AND R. TRIGGIANI, *Rational decay rates for a PDE heat-structure interaction: A frequency domain approach*, Evol. Equ. Control Theory, 2 (2013), pp. 233–253.
- [7] C. J. K. BATTY, R. CHILL, AND Y. TOMILOV, *Fine scales of decay of operator semigroups*, J. Eur. Math. Soc., (JEMS), 18 (2016), pp. 853–929.
- [8] C. J. K. BATTY AND T. DUYCKAERTS, *Non-uniform stability for bounded semi-groups in Banach spaces*, J. Evol. Equ., 8 (2008), pp. 765–780.
- [9] C. J. K. BATTY, L. PAUNONEN, AND D. SEIFERT, *Optimal energy decay in a one-dimensional coupled wave-heat system*, J. Evol. Equ., 16 (2016), pp. 649–664.
- [10] A. BORICHEV AND Y. TOMILOV, *Optimal polynomial decay of functions and operator semi-groups*, Math. Ann., 347 (2010), pp. 455–478.
- [11] N. BURQ AND M. HITRIK, *Energy decay for damped wave equations on partially rectangular domains*, Math. Res. Lett., 14 (2007), pp. 35–47.
- [12] T. DUYCKAERTS, *Optimal decay rates of the energy of a hyperbolic-parabolic system coupled by an interface*, Asymptot. Anal., 51 (2007), pp. 17–45.
- [13] J. LAGNESE, *Decay of solutions of wave equations in a bounded region with boundary dissipation*, J. Differential Equations, 50 (1983), pp. 163–182.
- [14] Z. LIU AND B. RAO, *Characterization of polynomial decay rate for the solution of linear evolution equation*, Z. Angew. Math. Phys., 56 (2005), pp. 630–644.

- [15] J. M. A. M. VAN NEERVEN, *The Asymptotic Behaviour of Semigroups of Linear Operators*, Birkhäuser, Basel, 1996.
- [16] J. RAUCH, X. ZHANG, AND E. ZUAZUA, *Polynomial decay for a hyperbolic-parabolic coupled system*, J. Math. Pures Appl., 84 (2005), pp. 407–470.
- [17] J. ROZENDAAL, D. SEIFERT, AND R. STAHN, *Optimal rates of decay for operator semigroups on Hilbert space*, Adv. Math., 346 (2019), pp. 359–388.
- [18] R. STAHN, *Optimal decay rate for the wave equation on a square with constant damping on a strip*, Z. Angew. Math. Phys., 68(36), 2017.
- [19] X. ZHANG AND E. ZUAZUA, *Long-time behavior of a coupled heat-wave system arising in fluid-structure interaction*, Arch. Ration. Mech. Anal., 184 (2007), pp. 49–120.