

Advancing research paradigms and pathophysiological pathways in psoriatic arthritis and ankylosing spondylitis: Proceedings of the 2017 Platform for the Exchange of Expertise and Research (PEER) meeting

Sonya Abraham^{a,*}, Anne Barton^b, Lihi Eder^c, Adrian Lim^d, Dennis McGonagle^e, Neil McHugh^f, Stephen Pennington^g, Raj Sengupta^h, Stefan Siebertⁱ, Paul Bowness^j, Peter H. Schafer^k, Eva Cullen^l, Oliver FitzGerald^g

^a NIHR/Wellcome Trust Clinical Research Facility, Imperial Centre for Translational and Experimental Medicine, Imperial College Healthcare NHS Trust, Hammer-smith Hospital, Du Cane Road, London W12 0HS, UK

^b Centre for Musculoskeletal Research, University of Manchester and Central Manchester NHS Foundation Trust, Manchester Academic Health Sciences Centre, Manchester, UK

^c Women's College Research Institute, Women's College Hospital, University of Toronto, Toronto, ON, Canada

^d Charing Cross Hospital, Imperial College London, UK

^e Leeds Institute of Rheumatic and Musculoskeletal Medicine, University of Leeds, Chapel Allerton Hospital, Leeds, UK

^f Royal National Hospital for Rheumatic Diseases and Department of Pharmacy and Pharmacology, University of Bath, Bath, UK

^g St. Vincent's University Hospital and Conway Institute for Biomolecular Research, University College Dublin School of Medicine, Dublin, Ireland

^h Royal National Hospital for Rheumatic Diseases, Bath, UK

ⁱ Institute of Infection, Immunity and Inflammation, University of Glasgow, Glasgow, UK

^j Nuffield Department of Orthopaedics Rheumatology and Musculoskeletal Science, Oxford OX3 7LD, UK

^k Celgene Corporation, Summit, NJ, USA

^l Celgene Ltd, Uxbridge, UK

Keywords:

Ankylosing spondylitis

Immunology

Inflammation

Psoriatic arthritis

Rheumatic diseases

Introduction – Dr. Sonya Abraham

The seronegative spondyloarthropathies, including psoriatic arthritis (PsA) and ankylosing spondylitis (AS), are characterized by varied clinical symptoms, severity, and disease course [1,2]. Diagnosis and monitoring can be challenging because there is no definitive laboratory biomarker for reliably measuring inflammation or other disease processes associated with spondyloarthropathies. Over time, many patients with PsA and AS eventually experience significant disability and impaired quality of life [1,2]. This may be partially

accounted for by delays in diagnosis and subsequent treatment [3], as well as the presence of comorbidities.

In recent years, research efforts aimed at identifying risk factors for PsA, including clinical, imaging, genetic, and laboratory assessments, have yielded major advances. The Platform for the Exchange of Expertise and Research (PEER) was formed to facilitate the exchange of research insights, sharing of expertise, and discussion of unmet needs in rheumatology research.

The objective of the current report is to provide an overview of the 2017 PEER meeting, which was held on May 19–20, 2017, in London, UK, and highlighted the most up-to-date research findings regarding PsA and AS pathophysiology, early detection, comorbidities, and treatment.

Development of psoriatic arthritis in patients with psoriasis: insights from a prospective cohort – Dr. Lihi Eder

The majority of patients with PsA initially develop skin disease, with joint disease following approximately 7–13 years later [4,5]. A delayed diagnosis of PsA is associated with worse outcomes than when the diagnosis is made earlier in the disease course [3]. In order to identify patients at risk for PsA, and increase early diagnosis and

Abbreviations: AS, ankylosing spondylitis; EPIC, Epidemiology of Psoriatic Arthritis; HLA, human leukocyte antigen; IL, interleukin; MHC, major histocompatibility complex; IMAPA, Immune Metabolic Associations in Psoriatic Arthritis; MRI, magnetic resonance imaging; MRM, multiple reaction monitoring; NIHR, National Institute for Health Research; PDUS, power Doppler ultrasound; PEER, Platform for the Exchange of Expertise and Research; PROMISE, PROgnostic Markers In Spondyloarthritis; PROMPT, "early detection to improve Outcome in people with undiagnosed Psoriatic arthritis"; PsA, psoriatic arthritis; RA, rheumatoid arthritis; SMI, superb microvascular imaging; SpA, spondyloarthritis; SPADE, Spondyloarthritis Diagnosis Evaluation; TNF, tumor necrosis factor; TUDOR, Total bUrDen of psoriasis.

* Corresponding author.

E-mail address: s.abraham@imperial.ac.uk (S. Abraham).

<https://doi.org/10.1016/j.semarthrit.2018.10.008>

0049-0172/© 2018 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license. (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

treatment, the ongoing Toronto Psoriasis Cohort Study aims to evaluate the genetic and environmental risk factors associated with PsA development; enrolled patients with psoriasis are followed on a yearly basis until they develop PsA (Fig. 1) [6].

The study has suggested that the incidence of PsA is higher than previously estimated [6]. Based on the data collected, a prediction model estimated that over 20 years of follow-up, approximately 35% of the cohort patients were predicted to develop PsA [6]; this contrasts with only 5% cumulative prevalence among psoriasis patients described in a previous report of a Rochester, Minnesota, cohort [7]. In the Toronto Psoriasis Cohort Study, clinical factors (including psoriasis severity), presence of nail lesions or uveitis, lower level of patient education, and use of retinoids to treat psoriasis were predictors of PsA development [6,8]. A related body of research has also identified trauma as a significant predictor of PsA development [9–11]. By contrast, trauma is not associated with onset of rheumatoid arthritis (RA) [9].

Other factors associated with PsA development in the Toronto cohort as well as a UK cohort include higher body mass index; presence of non-specific musculoskeletal symptoms, including pain, fatigue, and stiffness, among psoriasis patients, even in the absence of distinct arthritis on physical examination; and certain biomarkers [6,12]. C-X-C motif chemokine 10 (CXCL10) was found at higher levels in the serum of patients with psoriasis who subsequently developed PsA and may be a potential biomarker for predicting PsA [13,14]. Subclinical enthesitis, as detected by magnetic resonance imaging (MRI) or ultrasound, is often present in patients with psoriasis and observed at the nail and joints; the clinical significance of this finding requires further investigation [8,15].

Overall, the findings derived from the Toronto cohort and other related investigations indicate that it may be possible to identify a subset of psoriasis patients who may be at higher risk for the development of PsA than other psoriasis patients. Based on the Toronto cohort analyses discussed above, PsA risk factors include trauma, higher body mass index, severe skin psoriasis, nail lesions, uveitis, and nonspecific musculoskeletal symptoms, as well as higher serum levels of CXCL10. Use of these findings in the clinic may give rise to a new model of care in which certain psoriasis patients flagged as at high risk for PsA could obtain treatment at a substantially earlier stage of PsA disease development. Prompt identification of risk factors for PsA and earlier treatment initiation could potentially prevent the development of full-blown PsA.

PROMPT research program: an overview and update – Professor Neil Mchugh

The ongoing UK National Institute for Health Research (NIHR)-funded program grant for applied research, “early detection to

imPROve OutCoMe in people with undiagnosed Psoriatic arthriTis” (PROMPT), encompasses a series of research activities aimed at facilitating accurate and early detection of PsA in UK primary care. An additional aim of the study is to develop patient-completed outcome measures of early disease that are relevant to patients with PsA and not adequately captured by currently available assessments.

The PROMPT research program includes focus groups examining what patients perceive to be important patient-reported outcome measures to enhance early screening, detection, and diagnosis in the primary care setting and in clinical treatment of PsA [16,17]. PROMPT draws upon findings from the ongoing Epidemiology of Psoriatic Arthritis (EPIC) study, which is based on analysis of the Clinical Practice Research Datalink database for epidemiological factors associated with PsA. It also encompasses ongoing research into cost-effectiveness, screening questionnaires, and barriers to PsA diagnosis.

To date, focus group studies have demonstrated that pain and fatigue are rated as most important by patients, yet neither of these conditions is adequately captured in current composite measures for PsA [16,17]. Moreover, a substantial proportion of patients are diagnosed with PsA within 13 years of the onset of cutaneous psoriasis [5]. The interval between onset of psoriasis and PsA appears to have a genetic basis, with human leukocyte antigen (HLA)-C*06 being associated with a prolonged delay in PsA development and HLA-B*27 being associated with very short interval or contemporaneous onset of psoriasis and PsA [18,19]. The PROMPT study further indicated that uveitis and Crohn's disease, but not ulcerative colitis, are more common in patients with PsA than in those with psoriasis alone [20]. The Total bUrDen of psORiasis (TUDOR), an ongoing randomized, controlled study, is central to PROMPT in evaluating enhanced surveillance of PsA compared with standard primary care. TUDOR is the first study to prospectively examine the impact of early diagnosis on long-term outcomes in PsA (Fig. 2); with a planned 5-year duration, outcomes are expected to become available in 2021.

The ongoing quest for early diagnosis and biomarker discovery in ankylosing spondylitis – Dr. Raj Sengupta

Goals for improving outcomes in AS include earlier detection and treatment, which represents a major challenge and unmet need. In 1999, the time between symptom onset and diagnosis was a median of 5 years, which had not improved in 2013, based on analysis of clinical data at 2 large UK secondary care centers [21].

As with PsA, delay in diagnosis is associated with worse outcomes, including poor function and mobility and increased risk of work disability [22,23]. When AS is diagnosed and treated earlier, the Bath Ankylosing Spondylitis Metrology Index score tends to be




Fig. 1. The pre-clinical phases of PsA [6]. MSK, musculoskeletal psoriatic arthritis. Risk factors shown are based on data from the Toronto Psoriasis Cohort Study, a long-term prospective observational study assessing the development of psoriatic arthritis in patients with psoriasis.




Fig. 2. Overview of the PROMPT investigative plan. CPRD, UK Clinical Practice Research Datalink; PROMPT, “early detection to imPROve OutcoMe in people with undiagnosed Psoriatic arthriTis”; PROMS, patient-reported outcome measure; RCT, randomized controlled trial.

maintained, indicating that patient mobility remains stable and does not deteriorate [24]. In 1 study, a higher proportion of patients with early disease (< 5 years) responded to adalimumab treatment compared with patients with later disease (≥ 5 years) [25]. Another study demonstrated that following initiation of tumor necrosis factor (TNF) inhibitor treatment, radiographic progression was significantly slower in early disease patients than in later disease patients [26].

Among the emerging biomarkers for early detection of AS are approximately 41 genes, including those that encode major histocompatibility complex (MHC) class 1 proteins; related genes such as *ERAP1*, *UBE2E3*, *UBE2L3*, and those involved in CD8 + T-cell function; microbial sensing genes; and those involved in other immune functions that are not clearly understood [27,28]. However, the usefulness of these biomarkers in the clinical setting is not yet well developed.

Other inflammatory markers, such as C-reactive protein, erythrocyte sedimentation rate, and vascular endothelial growth factor, are not sufficiently specific and sensitive to differentiate patients with AS from either healthy controls or those with other arthropathy. Some patients with AS may exhibit normal values, and elevations of these markers are seen in other arthropathies [29,30]. IL-17 and IL-23, however, have been found to be more consistently elevated in patients with AS compared with healthy controls [31]. Markers of bone turnover, such as Wnt-3a, sclerostin, monocyte chemoattractant protein-1, and calprotectin, are promising because they tend to be higher in patients with AS compared with healthy controls [30,32,33]. Among these, only monocyte chemoattractant-1 has demonstrated the capability to distinguish AS patients from healthy controls or those with mechanical low back pain [33]. Likewise, while anti-flagellin antibodies and diagnostic biomarkers (e.g., autoantibodies against protein phosphatase magnesium-dependent 1A) are also elevated in patients with AS, only autoantibodies against protein phosphatase magnesium-dependent 1A have shown promise in helping to distinguish between patients with AS vs. healthy controls or patients with RA or PsA, given that they correlate with baseline disease severity and vary according to changes in disease severity related to clinical response to TNF- α inhibitor treatment [34,35]. These findings, however, require further validation in an independent patient cohort.

Major challenges in the use of biomarkers for AS diagnosis include a lack of consistent results across studies, large variation among patients due to the heterogeneous nature of AS, and questions around the reliability of the assays. Large studies in well-defined patient

cohorts are needed to bring use of these biomarkers to the bedside. To address some of these challenges, the PROgnostic Markers In Spondyloarthritis (PROMISE) study is under way, with a planned enrollment to include 250 patients with axial SpA or AS, with assessments to identify prognostic markers in SpA, including patient-reported outcomes and radiograph, MRI, biomarker, and genetic analyses. In the meantime, Web-based tools aimed at healthcare providers and patients with the intention of improving early AS detection are available, including the Spondyloarthritis Diagnosis Evaluation (SPADE) tool for the global provider community (<http://www.spadetool.co.uk/>), Back in Focus website for primary general practitioners (<https://www.axialspabackinfocus.co.uk>, sponsored by AbbVie), and Don't Turn Your Back on It website for patients (<http://www.dontturnyourbackonit.co.uk>).

Subclinical enthesopathy and development of psoriatic arthritis – Professor Dennis McGonagle

Enthesitis is inflammation occurring in tissue that connects ligament to bone, driving cytokine-mediated immunobiologic response at sites in the body that otherwise lack an immune response [36]. It is a common hallmark of the spondyloarthropathies, including PsA and AS [36]. Ongoing work has revealed how both mechanical stress and the genes responsible for the interleukin (IL)-12 and IL-23 inflammatory pathway have been associated with the development of enthesitis [37].

In murine models of spondyloarthritis (SpA), enthesal-resident IL-23 receptor-positive T cells have been shown to be critical to SpA pathogenesis [37]. In human enthesal tissue, IL-23R-positive innate type 3 lymphoid cells are present and show elevated levels of transforming growth factor- β 1 and TNF- α transcripts [38]. Moreover, patients with SpA and PsA exhibit microdamage at ligament and tendon attachment sites associated with MHC type 1-linked signals, including HLA-B*27 [39,40]. Clinically asymptomatic patients often have enthesopathy observed on ultrasound [41,42] and preliminary studies have shown that the presence of subclinical enthesitis may predict the development of PsA [43]. Enthesopathy is also linked to nail disease, which in turn is linked to PsA development [7,44]. In patients with psoriasis, barrier dysfunction may contribute to adaptive MHC-1-associated immune responses that drive enthesopathy and PsA development (Fig. 3) [45].




Fig. 3. Enthesopathy in psoriasis: Adaptive immune priming [45]. DAMP, damage-associated molecular pattern; DC, dendritic cell; ERAP1, endoplasmic reticulum aminopeptidase 1; NK, natural killer cell; PAMP, pathogen-associated molecular pattern.

Systemic treatment aimed at psoriasis, including methotrexate, the anti-TNFs, and ustekinumab [46], as well as anti-IL-17s and apremilast, may help to resolve subclinical enthesitis because these agents also have licensed indications for treating clinical arthritis. In a study using methotrexate (with or without a concomitant TNF inhibitor) in patients with psoriasis or PsA, the number of entheses with inflammatory abnormalities based on ultrasound was significantly decreased with 6 months of treatment [46]. The enthesis may be useful for ultrasound screening for early or preclinical PsA, as well as a treatment target; likewise, the systemic treatment of psoriasis has the potential to prevent development of arthritic lesions [43,46].

Superb microvascular imaging: novel Doppler ultrasound technology and its application in rheumatic diseases – Professor Adrian Lim

With the availability of new imaging techniques, the potential exists to diagnose PsA earlier and to better understand disease progression. Well-established imaging techniques that have been used to assess vascular flow include color/power Doppler ultrasound (PDUS) and some advanced dynamic flow ultrasound technologies. PDUS is cost-effective compared with contrast-enhanced MRI and useful for assessing inflammation in joints or tendons [47,48]. It is the current accepted gold standard to denote active inflammation in joints and tendons.

Superb microvascular imaging (SMI) is an emerging and novel ultrasound technology that allows visualization of the microvasculature

without the need for ultrasound contrast. SMI uses filters and a unique algorithm to selectively detect and remove random clutter without compromising the clinical data of interest. SMI, utilizing 24 MHz power, can distinguish slower flowing vessels within the microvasculature, which may help to identify sites of active inflammation (Fig. 4). In a study directly comparing the ability of SMI to detect inflammation in tendons and joints vs. PDUS, SMI revealed vascularity that was not detected by PDUS in a significant number of cases ($P=0.007$); in cases where vascularity was detected by both PDUS and SMI, vascularity scores were significantly greater with SMI ($P < 0.001$) [49].

Three-dimensional ultrasound imaging is also on the horizon. While the technology is available, its use requires training on how to properly manipulate the images. Images are currently of low resolution; however, algorithms are improving. Three-dimensional ultrasound has potential uses in measuring and quantifying vascularity in combination with SMI, as well as imaging cartilaginous surfaces. In addition to its potential usefulness in screening and diagnosis, future studies may include SMI to measure changes in vascularity during treatment.

Psoriatic arthritis and immune-metabolic interactions – Dr. Stefan Siebert

PsA is associated with comorbid diabetes, hypertension, and cardiovascular disease, as well as hyperuricemia [50,51]. Accumulating evidence demonstrates that metabolic dysfunction underlying these comorbid conditions is a key component of the PsA disease process




Fig. 4. Superb microvascular imaging of the index finger in a patient with psoriatic arthritis.




Fig. 5. Targeting PDE4 suppresses inflammation and decreases adiposity. Reprinted with permission from Wu C and Rajagopalan S. Phosphodiesterase-4 inhibition as a therapeutic strategy for metabolic disorders. *Obes Rev* 2016;17:429–41 [58], © John Wiley and Sons. cAMP, cyclic adenosine monophosphate; PDE4, phosphodiesterase 4.

itself and is not fully accounted for by lifestyle factors. Patients with PsA are more likely to have metabolic syndrome, as well as a greater number of metabolic syndrome components, than patients with RA or other, non-inflammatory conditions [52–54].

PsA patients also are more likely to be obese than the general population and to have a higher body mass index than patients with RA [52,55]. Additionally, psoriasis patients who are overweight or obese are more likely to develop PsA [55,56]. Obesity may play a role in the development of PsA and may influence treatment response. The release of adipokines from fat cells can drive metabolic dysfunction as well as pro-inflammatory pathways (Fig. 5), potentially leading to increased PsA risk [57,58]. Obesity is also associated with decreasing likelihood of optimal clinical response to therapy in PsA [59,60], while weight loss is associated with increased likelihood of achieving good outcomes with TNF inhibitor therapy [61].

The metabolic component of PsA may be influenced by the treatment prescribed. One study showed that PsA and RA patients who were receiving TNF inhibitors or hydroxychloroquine were at lower risk for developing diabetes than patients who were receiving methotrexate [62]. However, certain disease-modifying anti-rheumatic drugs, including TNF inhibitors, are associated with weight gain or an increase in percentage of body fat [63,64]. Phosphodiesterase 4 inhibitors have been associated with weight loss as well as reduction in hemoglobin A1c and may decrease insulin resistance [58,65–67] (Fig. 5). The ongoing Immune Metabolic Associations in Psoriatic Arthritis (IMAPA) study in Glasgow, UK, is investigating the effects of phosphodiesterase 4 inhibition on immune-metabolic interactions in PsA using the phosphodiesterase 4 inhibitor apremilast (ClinicalTrials.gov identifier: NCT03399708).

New insights into the genetics of psoriatic arthritis – Professor Anne Barton

PsA is a complex disease that has environmental and strong genetic components that contribute to susceptibility [68]. Identifying PsA susceptibility genes can aid in predicting disease development,

helping to indicate suitable preventive strategies, as well as in the design of more targeted drug therapies. In a genome-wide association study, variants of the *TRAF3IP2* gene were associated with PsA and psoriasis [69]. In a more recent case-control association analysis, while PsA-associated loci exhibited extensive overlap with known psoriasis-susceptibility loci, the 5q31 locus was specific to PsA [70]. Within the 5q31 locus, fine mapping analysis of single nucleotide polymorphisms revealed *SLC22A5* to be a likely candidate causal gene associated with PsA, with rs715285 or rs10065787 being the possible causal variant [70]. This association is independent of the IL-13 association with psoriasis.

Another gene known to be very important in the development of psoriasis and PsA encodes the IL-23 receptor (*IL-23R*) [70]. However, distinctly different variants on the *IL-23R* gene are associated with psoriasis vs. PsA; the gene associated with psoriasis encodes a non-synonymous variant and may impact gene function, while the PsA-associated variant is located in the *IL-23R* promoter region and may impact gene expression, although that has yet to be experimentally confirmed (Fig. 6) [71].

Variants linked to PsA appear more often to co-localize with epigenetic markers of open chromatin in CD8+ memory T cells. Genes encoding HLA subtypes have been strongly associated with psoriasis and PsA [70]. *HLA-C*0602* is strongly associated with both conditions and yet it counterintuitively has been reported to be protective for PsA compared with psoriasis [70,72]. However, after controlling for age at onset of psoriasis, *HLA-C*0602* has been shown to be primarily associated with skin (psoriasis) rather than joint (PsA) disease [73]. By contrast, variants at *HLA-B* (amino acid position 97) are associated with PsA and AS. The specific amino acid residue at position 97 may mediate which disease develops, and individuals with an associated serine residue are at risk for PsA but not for AS; those with an associated asparagine residue are at risk for both PsA and AS [73,74].

While an extensive overlap has been seen among the genes associated with psoriasis and PsA, a distinct set of genes uniquely contribute to the risk of PsA. Ongoing work encompassing genome-wide association studies, analysis of rare gene variants, and analyses of




Fig. 6. Different IL-23R gene variants are associated with PsA vs. psoriasis [71]. PsA, psoriatic arthritis; SNP, single nucleotide polymorphism.

genes that may contribute to PsA-associated risk factors, such as nail pitting, uveitis, and severe psoriasis, are required to fully understand the genetic background to PsA.

Arthropathies under a proteomic spotlight – Professor Stephen Pennington

The entire set of proteins expressed by an organism or an organ system may be evaluated using mass spectrometry-based proteomics [75]. In an era of increased understanding of disease complexity, such a proteomic approach forms the cornerstone of studies into the identification and measurement of panels of protein biomarkers that correlate with presence or severity of a certain condition [76,77]. Such protein biomarker panels or signatures may be used to improve disease diagnosis or treatment and to better support personalized medicine. This approach and process (using antibody-based assays) has yielded some success in the setting of RA, with approval and clinical adoption of the Vectra® DA test (Crescendo Bioscience, Inc., South San Francisco, CA, USA), which examines 12 biomarkers associated with RA disease activity [78].

For rheumatic diseases such as PsA and AS, a wide variety of treatments are available; however, only a small proportion of patients respond to any single treatment. Because rheumatic diseases are so physiologically heterogeneous, systems-driven proteomics may be a valuable approach toward predicting patient response to treatment by allowing personalized treatment decisions based on a unique profile or signature of proteins detected in a patient's tissue, synovial fluid, or blood [79–81]. Ongoing proteomic projects in PsA include examination of the pathophysiologic role of the IL-17 receptor and its adaptor protein ACT-1, biomarkers that predict response to TNF inhibitor treatment, biomarkers that predict joint damage, and biomarkers that aid PsA diagnosis [82–86].

The proteomic strategy begins with unbiased protein discovery, followed by assay development. The use of multiple reaction monitoring (MRM) mass spectrometry for multi-protein assays supports rapid assay development and analytical validation. Following clinical evaluation, the assays are then subjected to regulatory approval and adoption in the clinical setting (Fig. 7).

With regard to developing a similar test for PsA, several key questions remain unanswered. We do not fully understand the biologic mechanisms underlying bony changes that occur in PsA. Moreover, no biomarker test clearly differentiates PsA from RA, and no known

predictors have been linked to response to a given treatment. To address these challenges, the ongoing Health Research Board (Ireland)-funded study (PAPRICA) aims to evaluate approximately 140 target candidate proteins considered to be involved in PsA pathophysiology using MRM assays on blood samples obtained from patients enrolled in a number of clinical studies. It is hoped that the impending findings may be calibrated to address a variety of clinical questions and issues, including differential PsA/RA diagnosis, predicting disease progression or prognosis, and predicting treatment response.

Genetics and immunology of ankylosing spondylitis – Professor Paul Bowness

AS and the related spondyloarthritides are a group of diseases that share common genetic and clinical features, including inflammation that involves IL-23- and IL-17-driven signaling, enthesitis, and new bone formation [87]. Phenotypically, SpA and AS are considered immune-mediated inflammatory diseases, somewhere on a spectrum between autoimmune diseases and autoinflammatory diseases. AS is associated with a major hereditary component; the presence of HLA-B*27 accounts for approximately 30% of the hereditary variation, while *ERAP1*, *IL-23R*, *IL-12B*, and other genes linked to Th17 responses play lesser roles (> 100 loci have been linked to AS in large genetic analyses) [87,88].

HLA-B*27 is an MHC class I molecule that is present on most cells of the body. It functions by presenting peptide antigens to CD8⁺ T cells [87]. Each HLA-B*27 molecule carries a specific peptide antigen and is capable of initiating a corresponding immune response. The role of HLA-B*27 in AS pathogenesis is not completely understood; however, several hypotheses (i.e., arthritogenic, unfolded protein response, and free heavy chain) are being studied (Fig. 8). According to the arthritogenic hypothesis, HLA-B*27 might present “arthritogenic” peptides (e.g., from joints) to CD8⁺ T cells, hence triggering inflammation [87]. Alternatively, the unfolded protein response hypothesis proposes that inappropriate HLA-B*27 folding or trafficking might also occur due to malfunctioning *ERAP1*, which normally acts to trim the peptides bound to HLA-B*27 molecules and therefore facilitates cell surface egress [87]. The free heavy chain hypothesis posits that HLA-B*27 might malfunction due to the presence of an unpaired cysteine residue at position 67, which can cause it to form homodimers by pairing with another HLA-B*27 residue. HLA-B*27




Fig. 7. Multiple reaction monitoring development. LC-MS/MS, liquid chromatography–mass spectrometry.

homodimers can form inside cells but also can be expressed on the cell surface, where they can then bind to several immune receptors, triggering aberrant immune responses, including Th17 responses [87]. Th17 cells in turn produce a number of ILs, including IL-17, IL-21, IL-22, and IL-23. CD4 cells, including Th17 cells that express the KIR3DL2 receptor, which can bind strongly to HLA-B*27 homodimers, are increased in patients with AS and are increased in the joints of patients with SpA [89]. Currently, immunotherapy for AS includes TNF inhibitors and biologic therapies targeting IL-17/23. Future treatments may include small molecule inhibitors that target components of the Th17 inflammatory responses, such as HLA-B*27 signaling, transcription factors, and ERAP1.

Acknowledgments

The Platform for the Exchange of Expertise and Research (PEER) meeting was held on May 19–20, 2017, in London, UK.

Conflicts of interest: Dr. Abraham has received grant/research support from AbbVie, Celgene Corporation, Merck Sharp & Dohme, and

UCB. Dr. Barton has received grant/research support, speaker fees, and consultant fees from Celgene Corporation, Eli Lilly, Pfizer, and Roche-Chugai. Dr. Eder has received research and educational grants from and served as a consultant for AbbVie, Amgen, Celgene Corporation, Janssen, and Novartis. Professor Lim has served as a luminary for Toshiba Medical Systems. Professor McGonagle has received grant/research support from AbbVie, Celgene Corporation, Janssen, Merck Sharp & Dohme, Novartis, Pfizer, and UCB; and served on the speakers bureau for AbbVie, Celgene Corporation, Janssen, Merck Sharp & Dohme, Novartis, and Pfizer. Professor McHugh has received grant/research support and honoraria from AbbVie, Celgene Corporation, Eli Lilly, Pfizer, and UCB. Professor Pennington is the founder of and chief scientific officer for Atturo. Dr. Sengupta has received honoraria/expenses for attendance at advisory board meetings from AbbVie, Celgene Corporation, Merck Sharp & Dohme, Novartis, Pfizer, and UCB. Dr. Siebert has received grant/research support from Bristol-Myers Squibb, Boehringer Ingelheim, Celgene Corporation, Janssen,




Fig. 8. Overview of hypotheses explaining the role of HLA-B*27 in spondyloarthritis pathogenesis. HLA, human leukocyte antigen.

Pfizer, and UCB, and has received consultation and speaker fees from AbbVie, Boehringer Ingelheim, Janssen, Novartis, Pfizer, Roche, and UCB. Professor Bowness has received research funding support from Celgene Corporation and Merck. Dr. Schafer is an employee of Celgene Corporation. Dr. Cullen is an employee of Celgene Ltd. Dr. Fitzgerald has received honoraria and grant/research support from and has been a member of advisory boards for AbbVie, Celgene Corporation, Janssen, Merck Sharp & Dohme, Pfizer, Roche, and UCB.

Funding sources: The medical education meeting was organized and sponsored by Celgene UK and Ireland. Technical assistance with editing, figure preparation, and styling of the manuscript for submission was provided by Peloton Advantage, LLC, and was funded by Celgene Corporation. The PROMPT portion of this report contains independent research funded by the National Institute for Health Research (Programme Grants for Applied Research, Early detection to improve outcome in patients with undiagnosed psoriatic arthritis, [RP-PG-1212-20007](#)); the views in each portion of the publication are those of the author and not necessarily those of the National Health Service, the National Institute for Health Research, or the Department of Health. The authors were fully responsible for all content and editorial decisions and received no financial support or other form of compensation related to the development of this manuscript.

References

- [1] Coates LC, Kavanaugh A, Mease PJ, Soriano ER, Laura Acosta FM, Armstrong AW, et al. Group for Research and Assessment of Psoriasis and Psoriatic Arthritis: treatment recommendations for psoriatic arthritis 2015. *Arthr Rheumatol* 2016;68:1060–71.
- [2] Braun J, van den Berg R, Baraliakos X, Boehm H, Burgos-Vargas R, Collantes-Estevéz E, et al. 2010 update of the ASAS/EULAR recommendations for the management of ankylosing spondylitis. *Ann Rheum Dis* 2011;70:896–904.
- [3] Haroon M, Gallagher P, Fitzgerald O. Diagnostic delay of more than 6 months contributes to poor radiographic and functional outcome in psoriatic arthritis. *Ann Rheum Dis* 2015;74:1045–50.
- [4] Ritchlin CT, Colbert RA, Gladman DD. Psoriatic arthritis. *N Engl J Med* 2017;376:957–70.
- [5] Tillett W, Charlton R, Nightingale A, Snowball J, Green A, Smith C, et al. Interval between onset of psoriasis and psoriatic arthritis comparing the UK Clinical Practice Research Datalink with a hospital-based cohort. *Rheumatology* 2017;56:2109–13. (Oxford).
- [6] Eder L, Haddad A, Rosen CF, Lee KA, Chandran V, Cook R, et al. The incidence and risk factors for psoriatic arthritis in patients with psoriasis: a prospective cohort study. *Arthr Rheumatol* 2016;68:915–23.
- [7] Wilson FC, Icen M, Crowson CS, McEvoy MT, Gabriel SE, Kremers HM. Incidence and clinical predictors of psoriatic arthritis in patients with psoriasis: a population-based study. *Arthr Rheum* 2009;61:233–9.
- [8] Tan AL, Benjamin M, Toumi H, Grainger AJ, Tanner SF, Emery P, et al. The relationship between the extensor tendon entheses and the nail in distal interphalangeal joint disease in psoriatic arthritis—a high-resolution MRI and histological study. *Rheumatology* 2007;46:253–6. (Oxford).
- [9] Thorarensen SM, Lu N, Ogdie A, Gelfand JM, Choi HK, Love TJ. Physical trauma recorded in primary care is associated with the onset of psoriatic arthritis among patients with psoriasis. *Ann Rheum Dis* 2017;76:521–5.
- [10] Eder L, Law T, Chandran V, Shanmugarajah S, Shen H, Rosen CF, et al. Association between environmental factors and onset of psoriatic arthritis in patients with psoriasis. *Arthr Care Res* 2011;63:1091–7. (Hoboken).
- [11] Pattison E, Harrison BJ, Griffiths CE, Silman AJ, Bruce IN. Environmental risk factors for the development of psoriatic arthritis: results from a case-control study. *Ann Rheum Dis* 2008;67:672–6.
- [12] Love TJ, Gudjonsson JE, Valdimarsson H, Gudbjornsson B. Psoriatic arthritis and onycholysis – results from the cross-sectional Reykjavik psoriatic arthritis study. *J Rheumatol* 2012;39:1441–4.
- [13] Pollock RA, Abji F, Liang K, Chandran V, Pellett FJ, Virtanen C, et al. Gene expression differences between psoriasis patients with and without inflammatory arthritis. *J Invest Dermatol* 2015;135:620–3.
- [14] Abji F, Pollock RA, Liang K, Chandran V, Gladman DD. Brief report: CXCL10 is a possible biomarker for the development of psoriatic arthritis among patients with psoriasis. *Arthr Rheumatol* 2016;68:2911–6.
- [15] Eder L, Jayakar J, Thavaneswaran A, Haddad A, Chandran V, Salonen D, et al. Is the MADRID Sonographic Enthesitis Index useful for differentiating psoriatic arthritis from psoriasis alone and healthy controls? *J Rheumatol* 2014;41:466–72.
- [16] Dures E, Hewlett S, Lord J, Bowen C, McHugh N, Tillett W. Important treatment outcomes for patients with psoriatic arthritis: a multisite qualitative study. *The Patient* 2017;10:455–62.
- [17] Tillett W, Dures E, Hewlett S, Helliwell PS, Fitzgerald O, Brooke M, et al. A multicenter nominal group study to rank outcomes important to patients, and their representation in existing composite outcome measures for psoriatic arthritis. *J Rheumatol* 2017;44:1445–52.
- [18] Queiro R, Tejon P, Alonso S, Coto P. Age at disease onset: a key factor for understanding psoriatic disease. *Rheumatology* 2014;53:1178–85. (Oxford).
- [19] Haroon M, Winchester R, Giles JT, Heffernan E, Fitzgerald O. Certain class I HLA alleles and haplotypes implicated in susceptibility play a role in determining specific features of the psoriatic arthritis phenotype. *Ann Rheum Dis* 2016;75:155–62.
- [20] Charlton R, Green A, Shaddick G, Snowball J, Nightingale A, Tillett W, et al. Risk of uveitis and inflammatory bowel disease in people with psoriatic arthritis: a population-based cohort study. *Ann Rheum Dis* 2018;77:277–80.
- [21] Sykes MP, Doll H, Sengupta R, Gaffney K. Delay to diagnosis in axial spondyloarthritis: are we improving in the UK? *Rheumatology* 2015;54:2283–4. (Oxford).
- [22] Aggarwal R, Malaviya AN. Diagnosis delay in patients with ankylosing spondylitis: factors and outcomes—an Indian perspective. *Clin Rheumatol* 2009;28:327–31.
- [23] Gunasekera W, Shaddick G, Jobling A, Smith A, Sengupta R. Diagnostic delay worsens mobility and work disability in ankylosing spondylitis [abstract AB0735]. *Ann Rheum Dis* 2014;73(Suppl 2):1046.
- [24] Poddubnyy D, Fedorova A, Listing J, Haibel H, Baraliakos X, Braun J, et al. Physical function and spinal mobility remain stable despite radiographic spinal progression in patients with ankylosing spondylitis treated with TNF-alpha inhibitors for up to 10 years. *J Rheumatol* 2016;43:2142–8.
- [25] Sieper J, van der Heijde D, Dougados M, Mease PJ, Maksymowych WP, Brown MA, et al. Efficacy and safety of adalimumab in patients with non-radiographic axial spondyloarthritis: results of a randomised placebo-controlled trial (ABILITY-1). *Ann Rheum Dis* 2013;72:815–22.
- [26] Haroon N, Inman RD, Learch TJ, Weisman MH, Lee M, Rahbar MH, et al. The impact of tumor necrosis factor alpha inhibitors on radiographic progression in ankylosing spondylitis. *Arthr Rheum* 2013;65:2645–54.
- [27] Reveille JD. An update on the contribution of the MHC to AS susceptibility. *Clin Rheumatol* 2014;33:749–57.
- [28] Cortes A, Hadler J, Pointon JP, Robinson PC, Karaderi T, Leo P, et al. Identification of multiple risk variants for ankylosing spondylitis through high-density genotyping of immune-related loci. *Nat Genet* 2013;45:730–8.
- [29] Dougados M, Gueguen A, Nakache JP, Velicitat P, Zeidler H, Veys E, et al. Clinical relevance of C-reactive protein in axial involvement of ankylosing spondylitis. *J Rheumatol* 1999;26:971–4.
- [30] Turina MC, Yeremenko N, van Gaalen F, van Oosterhout M, Berg JJ, Ramonda R, et al. Serum inflammatory biomarkers fail to identify early axial spondyloarthritis: results from the SPondyloArthritis Caught Early (SPACE) cohort. *RMD Open* 2017;3:e000319.
- [31] Mei Y, Pan F, Gao J, Ge R, Duan Z, Zeng Z, et al. Increased serum IL-17 and IL-23 in the patient with ankylosing spondylitis. *Clin Rheumatol* 2011;30:269–73.
- [32] Klingberg E, Nurkka M, Carlsten H, Forsblad-d'Elia H. Biomarkers of bone metabolism in ankylosing spondylitis in relation to osteoproliferation and osteoporosis. *J Rheumatol* 2014;41:1349–56.
- [33] Romero-Sanchez C, Tsou HK, Jan MS, Wong RH, Chang IC, Londono J, et al. Serum monocyte chemoattractant protein-1 concentrations distinguish patients with ankylosing spondylitis from patients with mechanical low back pain. *J Spinal Disord Tech* 2011;24:202–7.
- [34] Wallis D, Asaduzzaman A, Weisman M, Haroon N, Anton A, McGovern D, et al. Elevated serum anti-flagellin antibodies implicate subclinical bowel inflammation in ankylosing spondylitis: an observational study. *Arthr Res Ther* 2013;15:R166.
- [35] Kim YG, Sohn DH, Zhao X, Sokolove J, Lindstrom TM, Yoo B, et al. Role of protein phosphatase magnesium-dependent 1A and anti-protein phosphatase magnesium-dependent 1A autoantibodies in ankylosing spondylitis. *Arthr Rheumatol* 2014;66:2793–803.
- [36] McGonagle D, Conaghan PG, Emery P. Psoriatic arthritis: a unified concept twenty years on. *Arthr Rheum* 1999;42:1080–6.
- [37] Sherlock JP, Joyce-Shaikh B, Turner SP, Chao CC, Sathe M, Grein J, et al. IL-23 induces spondyloarthritis by acting on ROR-gamma+ CD3+CD4-CD8- enthesal resident T cells. *Nat Med* 2012;18:1069–76.
- [38] Cuthbert RJ, Fragkakis EM, Dunsmuir R, Li Z, Coles M, Marzo-Ortega H, et al. Brief report: group 3 innate lymphoid cells in human entheses. *Arthr Rheumatol* 2017;69:1816–22.
- [39] McGonagle D, Georgouli T. The importance of 'Mechnikov's thorn' for an improved understanding of 21st century medicine and immunology: a view from the eye. *Scand J Immunol* 2008;68:129–39.
- [40] Benjamin M, McGonagle D. Histopathologic changes at "synovio-enthesal complexes" suggesting a novel mechanism for synovitis in osteoarthritis and spondyloarthritis. *Arthr Rheum* 2007;56:3601–9.
- [41] Gisondi P, Tinazzi I, El-Dalati G, Gallo M, Biasi D, Barbara LM, et al. Lower limb enthesopathy in patients with psoriasis without clinical signs of arthropathy: a hospital-based case-control study. *Ann Rheum Dis* 2008;67:26–30.
- [42] Naredo E, Moller I, de Miguel E, Batlle-Gualda E, Acebes C, Brito E, et al. High prevalence of ultrasonographic synovitis and enthesopathy in patients with psoriasis without psoriatic arthritis: a prospective case-control study. *Rheumatology* 2011;50:1838–48. (Oxford).
- [43] Tinazzi I, McGonagle D, Biasi D, Confente S, Caimmi C, Girolomoni G, et al. Preliminary evidence that subclinical enthesopathy may predict psoriatic arthritis in patients with psoriasis. *J Rheumatol* 2011;38:2691–2.
- [44] Aydin SZ, Castillo-Gallego C, Ash ZR, Marzo-Ortega H, Emery P, Wakefield RJ, et al. Ultrasonographic assessment of nail in psoriatic disease shows a link between onychopathy and distal interphalangeal joint extensor tendon enthesopathy. *Dermatology* 2012;225:231–5.

- [45] McGonagle D, Aydin SZ, Gul A, Mahr A, Direskeneli H. 'MHC-I-opathy'-unified concept for spondyloarthritis and Behcet disease. *Nat Rev Rheumatol* 2015;11:731–40.
- [46] Acquacalda E, Albert C, Montaudie H, Fontas E, Danre A, Roux CH, et al. Ultrasound study of entheses in psoriasis patients with or without musculoskeletal symptoms: a prospective study. *Joint Bone Spine* 2015;82:267–71.
- [47] Backhaus M, Kamradt T, Sandrock D, Loreck D, Fritz J, Wolf KJ, et al. Arthritis of the finger joints: a comprehensive approach comparing conventional radiography, scintigraphy, ultrasound, and contrast-enhanced magnetic resonance imaging. *Arthr Rheum* 1999;42:1232–45.
- [48] Taljanovic MS, Melville DM, Gimber LH, Scalcione LR, Miller MD, Kwoh CK, et al. High-resolution US of rheumatologic diseases. *Radiographics* 2015;35:2026–48.
- [49] Lim AKP, Satchithananda K, Dick EA, Abraham S, Cosgrove DO. Microflow imaging: new doppler technology to detect low-grade inflammation in patients with arthritis. *Eur Radiol* 2018;28:1046–53.
- [50] Shah K, Mellars L, Changolkar A, Feldman SR. Real-world burden of comorbidities in US patients with psoriasis. *J Am Acad Dermatol* 2017;77:287–92.
- [51] Tsuruta N, Imafuku S, Narisawa Y. Hyperuricemia is an independent risk factor for psoriatic arthritis in psoriatic patients. *J Dermatol* 2017;44:1349–52.
- [52] Craciun L, Craciun P, Buicu F. Prevalence of metabolic syndrome in psoriatic arthritis and rheumatoid arthritis. *Acta Medica Marisensis* 2014;60:196–9.
- [53] Haroon M, Gallagher P, Heffernan E, FitzGerald O. High prevalence of metabolic syndrome and of insulin resistance in psoriatic arthritis is associated with the severity of underlying disease. *J Rheumatol* 2014;41:1357–65.
- [54] Haroon M, Rafiq Chaudhry AB, FitzGerald O. Higher prevalence of metabolic syndrome in patients with psoriatic arthritis: a comparison with a control group of noninflammatory rheumatologic conditions. *J Rheumatol* 2016;43:463–4.
- [55] Love TJ, Zhu Y, Zhang Y, Wall-Burns L, Ogdie A, Gelfand JM, et al. Obesity and the risk of psoriatic arthritis: a population-based study. *Ann Rheum Dis* 2012;71:1273–7.
- [56] Li W, Han J, Qureshi AA. Obesity and risk of incident psoriatic arthritis in US women. *Ann Rheum Dis* 2012;71:1267–72.
- [57] Davidovici BB, Sattar N, Prinz J, Puig L, Emery P, Barker JN, et al. Psoriasis and systemic inflammatory diseases: potential mechanistic links between skin disease and co-morbid conditions. *J Invest Dermatol* 2010;130:1785–96.
- [58] Wu C, Rajagopalan S. Phosphodiesterase-4 inhibition as a therapeutic strategy for metabolic disorders. *Obes Rev* 2016;17:429–41.
- [59] di Minno MN, Peluso R, Iervolino S, Lupoli R, Russolillo A, Scarpa R, et al. Obesity and the prediction of minimal disease activity: a prospective study in psoriatic arthritis. *Arthr Care Res* 2013;65:141–7. (Hoboken).
- [60] Eder L, Thavaneswaran A, Chandran V, Cook RJ, Gladman DD. Obesity is associated with a lower probability of achieving sustained minimal disease activity state among patients with psoriatic arthritis. *Ann Rheum Dis* 2015;74:813–7.
- [61] Di Minno MN, Peluso R, Iervolino S, Russolillo A, Lupoli R, Scarpa R. Weight loss and achievement of minimal disease activity in patients with psoriatic arthritis starting treatment with tumour necrosis factor alpha blockers. *Ann Rheum Dis* 2014;73:1157–62.
- [62] Solomon DH, Massarotti E, Garg R, Liu J, Canning C, Schneeweiss S. Association between disease-modifying antirheumatic drugs and diabetes risk in patients with rheumatoid arthritis and psoriasis. *JAMA* 2011;305:2525–31.
- [63] Di Renzo L, Saraceno R, Schipani C, Rizzo M, Bianchi A, Noce A, et al. Prospective assessment of body weight and body composition changes in patients with psoriasis receiving anti-TNF- α treatment. *Dermatol Ther* 2011;24:446–51.
- [64] Baker JF, Sauer BC, Cannon GW, Teng CC, Michaud K, Ibrahim S, et al. Changes in body mass related to the initiation of disease-modifying therapies in rheumatoid arthritis. *Arthr Rheumatol* 2016;68:1818–27.
- [65] Paul C, Cather J, Gooderham M, Poulin Y, Mrowietz U, Ferrandiz C, et al. Efficacy and safety of apremilast, an oral phosphodiesterase 4 inhibitor, in patients with moderate to severe plaque psoriasis over 52 weeks: a phase III, randomized, controlled trial (ESTEEM 2). *Br J Dermatol* 2015;173:1387–99.
- [66] Wouters EF, Bredenkroter D, Teichmann P, Brose M, Rabe KF, Fabbri LM, et al. Effect of the phosphodiesterase 4 inhibitor roflumilast on glucose metabolism in patients with treatment-naïve, newly diagnosed type 2 diabetes mellitus. *J Clin Endocrinol Metab* 2012;97:E1720–5.
- [67] Jensterle M, Salamon V, Kocjan T, Vrtacnik Bokal E, Janez A. Short term monotherapy with GLP-1 receptor agonist liraglutide or PDE 4 inhibitor roflumilast is superior to metformin in weight loss in obese PCOS women: a pilot randomized study. *J Ovarian Res* 2015;8:32.
- [68] Karason A, Love TJ, Gudbjornsson B. A strong heritability of psoriatic arthritis over four generations—the Reykjavik Psoriatic Arthritis Study. *Rheumatology* 2009;48:1424–8. (Oxford).
- [69] Huffmeier U, Uebe S, Ekici AB, Bowes J, Giardina E, Korendowycz E, et al. Common variants at TRAF3IP2 are associated with susceptibility to psoriatic arthritis and psoriasis. *Nat Genet* 2010;42:996–9.
- [70] Bowes J, Budu-Aggrey A, Huffmeier U, Uebe S, Steel K, Hebert HL, et al. Dense genotyping of immune-related susceptibility loci reveals new insights into the genetics of psoriatic arthritis. *Nat Commun* 2015;6:6046.
- [71] Budu-Aggrey A, Bowes J, Loehr S, Uebe S, Zervou MI, Helliwell P, et al. Replication of a distinct psoriatic arthritis risk variant at the IL23R locus. *Ann Rheum Dis* 2016;75:1417–8.
- [72] Okada Y, Han B, Tsoi LC, Stuart PE, Ellinghaus E, Tejasvi T, et al. Fine mapping major histocompatibility complex associations in psoriasis and its clinical subtypes. *Am J Hum Genet* 2014;95:162–72.
- [73] Bowes J, Ashcroft J, Dand N, Jalali-Najafabadi F, Bellou E, Ho P, et al. Cross-phenotype association mapping of the MHC identifies genetic variants that differentiate psoriatic arthritis from psoriasis. *Ann Rheum Dis* 2017;76:1774–9.
- [74] Cortes A, Pulit SL, Leo PJ, Pointon JJ, Robinson PC, Weisman MH, et al. Major histocompatibility complex associations of ankylosing spondylitis are complex and involve further epistasis with ERAP1. *Nat Commun* 2015;6:7146.
- [75] Aebersold R, Mann M. Mass-spectrometric exploration of proteome structure and function. *Nature* 2016;537:347–55.
- [76] Meyer JG, Schilling B. Clinical applications of quantitative proteomics using targeted and untargeted data-independent acquisition techniques. *Expert Rev Proteom* 2017;14:419–29.
- [77] Percy AJ, Byrns S, Pennington SR, Holmes DT, Anderson NL, Agreste TM, et al. Clinical translation of MS-based, quantitative plasma proteomics: status, challenges, requirements, and potential. *Expert Rev Proteom* 2016;13:673–84.
- [78] Centola M, Cavet G, Shen Y, Ramanujan S, Knowlton N, Swan KA, et al. Development of a multi-biomarker disease activity test for rheumatoid arthritis. *PLoS One* 2013;8:e60635.
- [79] Cretu D, Liang K, Saraon P, Batruch I, Diamandis EP, Chandran V. Quantitative tandem mass-spectrometry of skin tissue reveals putative psoriatic arthritis biomarkers. *Clin Proteom* 2015;12:1.
- [80] Mahendran SM, Oikonomopoulou K, Diamandis EP, Chandran V. Synovial fluid proteomics in the pursuit of arthritis mediators: an evolving field of novel biomarker discovery. *Crit Rev Clin Lab Sci* 2017;54:495–505.
- [81] McArdle A, Pennington S, FitzGerald O. Clinical features of psoriatic arthritis: a comprehensive review of unmet clinical needs. *Clin Rev Allergy Immunol* 2017, Jul 27. <https://doi.org/10.1007/s12016-017-8630-7>.
- [82] Doyle MS, Collins ES, FitzGerald OM, Pennington SR. New insight into the functions of the interleukin-17 receptor adaptor protein Act1 in psoriatic arthritis. *Arthr Res Ther* 2012;14:226.
- [83] Ademowo OS, Hernandez B, Collins E, Rooney C, Fearon U, van Kuijk AW, et al. Discovery and confirmation of a protein biomarker panel with potential to predict response to biological therapy in psoriatic arthritis. *Ann Rheum Dis* 2016;75:234–41.
- [84] Collins ES, Butt AQ, Gibson DS, Dunn MJ, Fearon U, van Kuijk AW, et al. A clinically based protein discovery strategy to identify potential biomarkers of response to anti-TNF-alpha treatment of psoriatic arthritis. *Proteom Clin Appl* 2016;10:645–62.
- [85] McArdle A, Flatley B, Pennington SR, FitzGerald O. Early biomarkers of joint damage in rheumatoid and psoriatic arthritis. *Arthr Res Ther* 2015;17:141.
- [86] Butt AQ, McArdle A, Gibson DS, FitzGerald O, Pennington SR. Psoriatic arthritis under a proteomic spotlight: application of novel technologies to advance diagnosis and management. *Curr Rheumatol Rep* 2015;17:35.
- [87] Tam LS, Gu J, Yu D. Pathogenesis of ankylosing spondylitis. *Nat Rev Rheumatol* 2010;6:399–405.
- [88] Ellinghaus D, Jostins L, Spain SL, Cortes A, Bethune J, Han B, et al. Analysis of five chronic inflammatory diseases identifies 27 new associations and highlights disease-specific patterns at shared loci. *Nat Genet* 2016;48:510–8.
- [89] Bowness P, Ridley A, Shaw J, Chan AT, Wong-Baeza I, Fleming M, et al. Th17 cells expressing KIR3DL2+ and responsive to HLA-B27 homodimers are increased in ankylosing spondylitis. *J Immunol* 2011;186:2672–80.