

## RESEARCH LETTER

10.1002/2016GL072202

## Key Points:

- Observations provide evidence against strongly stabilizing tropical cloud feedbacks
- Compensation between feedbacks in stable and deep convective regimes resulted in a small net feedback consistent with climate models
- High clouds decrease with warming in regions of deep convection but increase elsewhere, leading to smaller changes in high-cloud fraction

## Supporting Information:

- Supporting Information S1
- Figure S1
- Figure S2
- Figure S3
- Figure S4
- Figure S5
- Figure S6
- Figure S7
- Figure S8

## Correspondence to:

I. N. Williams,  
inwilliams@lbl.gov

## Citation:

Williams, I. N., and R. T. Pierrehumbert (2017), Observational evidence against strongly stabilizing tropical cloud feedbacks, *Geophys. Res. Lett.*, 44, 1503–1510, doi:10.1002/2016GL072202.

Received 1 DEC 2016

Accepted 16 JAN 2017

Accepted article online 19 JAN 2017

Published online 8 FEB 2017

Published 2017. This article is a US Government work and is in the public domain in the United States of America.

## Observational evidence against strongly stabilizing tropical cloud feedbacks

Ian N. Williams<sup>1</sup> and Raymond T. Pierrehumbert<sup>2</sup>
<sup>1</sup>Climate and Ecosystem Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, California, USA,

<sup>2</sup>Department of Physics, University of Oxford, Oxford, UK

**Abstract** We present a method to attribute cloud radiative feedbacks to convective processes, using subcloud layer buoyancy as a diagnostic of stable and deep convective regimes. Applying this approach to tropical remote sensing measurements over years 2000–2016 shows that an inferred negative short-term cloud feedback from deep convection was nearly offset by a positive cloud feedback from stable regimes. The net cloud feedback was within statistical uncertainty of the National Center for Atmospheric Research Community Atmosphere Model (CAM5) with historical forcings, with discrepancies in the partitioning of the cloud feedback into convective regimes. Compensation between high-cloud responses to tropics-wide warming in stable and unstable regimes resulted in smaller net changes in high-cloud fraction with warming. In addition, deep convection and associated high clouds set in at warmer temperatures in response to warming, as a consequence of nearly invariant subcloud buoyancy. This invariance further constrained the magnitude of cloud radiative feedbacks and is consistent with climate model projections.

## 1. Introduction

Tropical clouds contribute to much of the uncertainty in climate sensitivity to greenhouse gas radiative forcing [Wetherald and Manabe, 1988; Ramanathan et al., 1989]. Low-level clouds are the largest source of intermodel variance in climate projections [Bony and Dufresne, 2005; Webb et al., 2006; Soden and Vecchi, 2011; Sherwood et al., 2014]. However, there remains uncertainty in the radiative effects of deep convection and associated high clouds. It was recently estimated that observed climate sensitivity is near the lower end of the range of climate model predictions and that a modest reduction in the high climate sensitivities of some models could be achieved by adding a model temperature dependence to the precipitation efficiency of deep convection [Mauritsen and Stevens, 2015]. A related study found a decrease in tropical high-cloud amount with warming due to clustering or aggregation of deep convection in a climate model [Bony et al., 2016]. On the other hand, efforts to explicitly resolve deep convection at global scale suggest a large positive high-cloud feedback [Tsuchida et al., 2014], which may be sensitive to subgrid turbulence and microphysics parameterizations [Bretherton, 2015].

Observational estimates of deep convective cloud feedback are challenging in part because deep convection is coupled to large-scale climate dynamics in addition to surface temperature. Past estimates were based on local covariation of cloud radiative effect (CRE) and sea surface temperature (SST) [Ramanathan and Collins, 1991; Lindzen et al., 2001] and were confounded by the tendency for deep convection and warm SSTs to coincide with large-scale ascent [Fu et al., 1992; Wallace, 1992; Hartmann and Michelsen, 1993; Lau et al., 1994, 1997; Hartmann and Michelsen, 2002]. Moreover, high-cloud feedback may be governed by different processes in regions of deep convection (and large-scale ascent) compared to remote stable regions, since only half of all tropical cirrus clouds are formed by detrainment from deep convection [Luo and Rossow, 2004]. Understanding of deep convective cloud feedback could be improved by separating the intrinsic temperature response of clouds from the influence of large-scale circulation on cloud type and convective regimes. Additionally, a method to control for the influences of large-scale circulation would help eliminate one possible source of the time scale dependence of cloud feedbacks noted previously [Zhou et al., 2016] and thus would aid in evaluating climate predictions against short-term observations.

Here we use a measure of subcloud layer buoyancy ( $B$ ) to partition the cloud feedback into convective regimes. Deep convection and associated high clouds set in when the subcloud layer is buoyant with respect to the free troposphere ( $B > 0$ ). Moreover, when negative,  $B$  is a measure of inversion strength, which is highly correlated

with low clouds [Klein and Hartmann, 1993; Wood and Bretherton, 2006]. Definitions of deep convection based on  $B$  are largely invariant under tropics-wide warming, as shown here. A related method based on mid-tropospheric vertical velocity ( $\omega_{500}$ ) [Bony et al., 2004] is expected to yield similar results for deep convection, since  $\omega_{500}$  and  $B$  are dynamically coupled; i.e., where  $B$  is positive, deep convection contributes to large-scale ascent. Therefore,  $B$  reflects both the large-scale circulation and the thermodynamic environment to which it is coupled. Unlike  $\omega_{500}$ , stability-related metrics such as  $B$  can better distinguish between shallow and deep clouds [Medeiros and Stevens, 2011; Webb et al., 2015]. Furthermore,  $B$  can be calculated independently from thermodynamic soundings, whereas  $\omega_{500}$  must be derived from climate models.

Our objectives are to (1) demonstrate the utility of a buoyancy-based diagnostic of tropical cloud radiative effects, (2) diagnose observed short-term cloud feedbacks from clouds in deep convective and stable regimes, and (3) evaluate cloud feedbacks in a climate model.

## 2. Diagnosing Cloud Radiative Effects of Deep Convection

We used monthly TOA shortwave and longwave all-sky radiation measured from CERES (Clouds and Earth's Radiant Energy System) on the Terra satellite (CERES EBAF-TOA Ed2.8) Wielicki et al. [1996], and SST from the U.S. National Oceanic and Atmospheric Administration (NOAA) optimal interpolation data set [Reynolds and Smith, 1994; Reynolds et al., 2002]. CRE was calculated by subtracting the clear-sky from the all-sky radiative fluxes, using clear-sky fluxes from the European Centre for Medium-Range Weather Forecasts (ECMWF) interim reanalysis product at  $1^\circ \times 1^\circ$  resolution [Dee et al., 2011], as in Dessler [2010] and Dessler and Loeb [2013]. The resulting CRE was adjusted to account for noncloud effects using a radiative kernel approach [Shell et al., 2008]. We focused our analysis on the tropical oceans ( $30^\circ\text{S}$  to  $30^\circ\text{N}$ ), for years CERES data were available (March 2000 to May 2016). We compared CERES observations to a climate model simulation (1979–2006), from the Atmospheric Model Intercomparison Project (AMIP). We used a single run from the National Center for Atmospheric Research (NCAR) Community Atmosphere Model (CAM5.1) at  $1^\circ \times 1^\circ$  resolution with historical forcings (hereafter CAM5-AMIP). Years 1982–1983 and 1991–1992 were excluded due to volcanic aerosol.


To explore the relationship between CRE and SST, we averaged shortwave and longwave CRE within SST intervals of  $0.5^\circ\text{C}$  to obtain their SST distributions (Figure 1a). The threshold SST for deep convection is indicated by the sharp rise in shortwave and longwave CRE between  $26$  and  $29^\circ\text{C}$  (Figure 1a). The observed near cancellation between shortwave and longwave CRE [Kiehl, 1994] results in a smaller net cooling effect of clouds near the convective threshold SST. Note that CRE drops off at very high SSTs greater than  $30^\circ\text{C}$ , which has been interpreted in relation to large-scale circulations [Waliser, 1996] and wind evaporation feedback on SST “hot spots” [Sobel and Gildor, 2003].

To diagnose the convective processes influencing cloud type and CRE, we replaced SST with a measure of subcloud buoyancy ( $B$ ).  $B$  is proportional to the difference between the unsaturated moist static energy of the subcloud layer ( $h_0$ ), and the saturated moist static energy of the overlying free troposphere ( $h^*$ ). This follows from the temperature difference between cloudy air and the environment in the tropics. The free-tropospheric temperature profile can be approximated by a saturated moist adiabat, while the cloudy air originates near the surface and conserves moist static energy up to the cloud base [Randall, 2015]. This diagnostic is similar to the “entropy excess” introduced earlier [Williams et al., 2009]. However, moist static energy allows an approximate expression for buoyancy at cloud base to be written compactly as

$$B = \frac{h_0 - h^*}{c_p T (1 + \gamma)} \quad (1)$$

where  $B$  is the buoyancy nondimensionalized by the gravitational acceleration and  $c_p$  is the specific heat of dry air.  $T$  represents the temperature at cloud base, which for our purposes can be approximated by the near-surface air temperature (see supporting information S1 for details). Although variations in  $B$  are largely controlled by the numerator in equation (1), the denominator gives the expression convenient units of acceleration when  $B$  is multiplied by gravitational acceleration. The additional factor  $(1 + \gamma)$  arises from the temperature dependence of the saturation mixing ratio,

$$\gamma = \frac{L}{c_p} \frac{dq^*}{dT} \quad (2)$$



**Figure 1.** (a) Distribution of monthly cloud radiative effect (CRE) shown as a function of SST and (b) nondimensional subcloud layer buoyancy (i.e.,  $B$ ), over tropical oceans. Deep convection is indicated by the sharp increase in both (minus) shortwave (dashed blue lines) and longwave (solid red lines) CRE between 27 and 29°C. CRE declines over the warmest SSTs (SST > 30), whereas CRE generally increases with  $B > 0$ . Error bars indicate the standard error of the mean. Circles with error bars show the interquartile range for bins having less than 20 data points, and circles without error bars show individual data points.

where  $L$  is the latent heat of vaporization and the change in saturation mixing ratio ( $q^*$ ) with temperature is evaluated at constant pressure.


We calculated  $B$  using ECMWF interim reanalysis. Other reanalysis products gave similar results. We averaged  $h^*$  over the free troposphere between 300 and 925 hPa, and averaged  $h_0$  between 925 and 1000 hPa. There was little sensitivity to the exact choice of levels, although differences between modeled and observed  $B$  were larger when calculating  $h_0$  using surface values at 2 m. The buoyancy distribution of CRE was calculated by averaging CRE within  $B$  intervals of  $10^{-3}$ . The spatial distributions of stable and unstable regimes are shown in supporting information Figure S1.

Shortwave and longwave CRE sharply increase above  $B = 0$  (Figure 1b), indicating the presence of deep convection and high clouds. Cloud radiative effects are more clearly delineated by  $B$ , particularly in very stable and unstable regimes. The larger shortwave CRE under stable conditions is characteristic of low clouds and is consistent with previous work showing that low cloud amount is strongly correlated with lower tropospheric inversion strength [Klein and Hartmann, 1993] and subsidence [Bony and Dufresne, 2005]. Furthermore, the comparison of SST and  $B$  coordinates suggests that the hottest SSTs are not actually in areas of strong buoyancy. In fact, they are rather stable or neutrally stable regions (Figure S2). Instead, the maximum cloud radiative effect and maximum buoyancy are both occurring at somewhat cooler temperatures.

### 3. Response of Cloud Radiative Effect to Warming

We estimated distributions of CRE over SST and  $B$  during the 2015–2016 El Niño, and for the three coldest years in the CERES record as determined by averaging annual SSTs over the tropical oceans (30°S to 30°N). The averaging was performed from June to May, to maximize interannual variability associated with the El Niño–Southern Oscillation. The difference in average SST between the 2015–2016 El Niño year and the coldest years (2000, 2007, and 2011) was 0.60°C. The comparison (Figure 2a) shows that the convective threshold increases with tropics-wide warming, as seen by the shift in onset of strong CRE beyond 26°C.


The success of  $B$  in controlling for variations in the convective threshold is indicated by the approximate collapse of the warmest and coldest years CRE onto a single curve when plotted as a function of  $B$  (Figure 2b). A similar result was obtained for the Coupled Model Intercomparison Project (CMIP3), for the scenario in which equivalent  $\text{CO}_2$  concentration starts at preindustrial levels and increases  $1\% \text{ yr}^{-1}$  until doubling [Williams *et al.*, 2009, Figures 1b and 1d]. Compared to  $B$ , midtropospheric vertical velocity yielded similar results for deep convective clouds but was less successful in detecting low clouds (Figures S3 and S4).



**Figure 2.** Effects of tropics-wide SST variability on observed and modeled CRE, shown as functions of local SST (Figures 2a, 2c, 2e, and 2g) and  $B$  (Figures 2b, 2d, 2f, and 2h). (a) The SST distributions of (minus) shortwave (dashed lines) and longwave (solid lines) CRE shift almost uniformly to warmer SSTs during the 2015–2016 El Niño relative to the three coldest years observed. (b) The buoyancy distribution of CRE is approximately invariant with warming. (c) Relative frequency distributions show that warming is approximately uniform across SST. (d) Warming results in little change to the relative frequency distribution of buoyancy. (e–h) As in Figures 2a–2d but for CAM5-AMIP simulations, for the three warmest and coldest simulated years. Error bars indicate the range over all three years.

The entire SST relative frequency distribution shifts almost uniformly to warmer SSTs with tropics-wide warming (Figure 2c), demonstrating that the observed “cliff” in the relative frequency distribution is not indicative of an upper limit on SST imposed by deep convection. Unlike SST, the relative frequency distribution of  $B$  is almost invariant between the warmest and coldest years (Figure 2d), implying that the area covered by deep convection does not substantially increase in response to tropics-wide warming. Approximate invariance of the  $B$  distribution was also seen in climate model projections [Williams *et al.*, 2009].

Results from the CAM5-AMIP simulation (Figures 2e–2h) also show a shift in the SST threshold for deep convection with tropics-wide warming (Figures 2e and 2g), and a collapse of the CRE distributions onto a single curve in  $B$  coordinates for the three warmest (1987, 1997, and 2003) and coldest (1984, 1985, and 1988)



**Figure 3.** Change in CRE per kelvin tropical warming ( $\Delta\text{CRE}_{\text{adj}}/\Delta T$ ), estimated by linear regression of adjusted CRE against tropical average temperature anomalies within each  $0.25 \times 10^{-2}$  buoyancy interval. (a) Observed (CERES) shortwave (SW; dashed blue) and longwave (LW; red)  $\Delta\text{CRE}_{\text{adj}}/\Delta T$  are large in magnitude but of opposite sign in the deep convective regime ( $B > 0$ ). (b) Shortwave and longwave  $\Delta\text{CRE}_{\text{adj}}/\Delta T$  in CAM5 are smaller in magnitude than observed. (c) The net  $\Delta\text{CRE}_{\text{adj}}/\Delta T$  from deep convection is positive in CAM5 (red) but negative in CERES (black), while the net  $\Delta\text{CRE}_{\text{adj}}/\Delta T$  from stable cloud regimes ( $B < 0$ ) is less positive in CAM5 than in CERES. Error bars indicate the 95% confidence interval for regression slope.

simulated years (Figure 2f). Modeled CRE is stronger than observed for deep convective regimes ( $B > 0$ ) (Figures 2e and 2f), and modeled  $B$  (Figure 2h) is slightly higher than observed (Figure 2d), with a peak just above 0. However, the shift in SST threshold for deep convection is similar to that seen in global climate model simulations in response to  $\text{CO}_2$  doubling [Williams *et al.*, 2009, Figures 1a and 1c] and indicates that the models are reliably capturing the response of convective threshold SST to tropics-wide warming.


A limitation of reanalysis-derived  $B$  is that it may be affected by the atmospheric model used in the reanalysis. We compared ECMWF to twice daily radiosonde profiles from the Atmospheric Radiation Measurement (ARM) site on Nauru Island ( $0.5^\circ\text{S}$ ,  $166.9^\circ\text{E}$ ) using monthly temperature and relative humidity for years 1999–2012 [Ciesielski *et al.*, 2009]. Nauru is on the eastern edge of the western Pacific warm pool and sees a reduction in cloudiness (corresponding to negative  $B$ ) during La Niña [Long *et al.*, 2013]. Variations in the ARM-derived  $B$  closely tracked the ECMWF-derived  $B$  (Figure S5).

#### 4. Inferred Cloud Feedbacks in Deep Convective and Stable Regimes

We estimated short-term cloud radiative feedbacks by regressing monthly adjusted CRE anomalies (departures from monthly climatology) against surface temperature anomalies [Gregory *et al.*, 2004], using monthly tropical average surface air temperature [e.g., Zelinka and Hartmann, 2011]. These estimates, hereafter denoted by  $\Delta\text{CRE}_{\text{adj}}/\Delta T$ , were categorized into convective regimes by calculating the regression slopes conditional on  $B$ . Shortwave and longwave  $\Delta\text{CRE}_{\text{adj}}/\Delta T$  were substantially larger for deep convective clouds ( $B > 0$ ) than for stable clouds in both CERES observations (Figure 3a) and CAM5-AMIP (Figure 3b), consistent with previous studies (based on cloud top pressure (CTP)) that inferred large but nearly canceling shortwave (positive) and longwave (negative) cloud feedbacks from high clouds [Zelinka and Hartmann, 2011; Zhou *et al.*, 2013]. The positive shortwave  $\Delta\text{CRE}_{\text{adj}}/\Delta T$  here is slightly less effective at canceling the negative longwave  $\Delta\text{CRE}_{\text{adj}}/\Delta T$ , resulting in a net negative  $\Delta\text{CRE}_{\text{adj}}/\Delta T$  from deep convection (Figure 3c). This is not necessarily inconsistent with the small positive high-cloud feedback inferred previously [Zelinka and Hartmann, 2011; Zhou *et al.*, 2013], given uncertainties in both estimates, and because not all high clouds occur in deep convective regimes. Estimates of  $\Delta\text{CRE}_{\text{adj}}/\Delta T$  using  $\omega_{500}$  instead of  $B$  were qualitatively similar but of smaller magnitude (Figure S6), suggesting that  $B$  better characterizes the diversity of cloud responses to warming.

Relative to the observational estimate, CAM5-AMIP underestimated the negative longwave  $\Delta\text{CRE}_{\text{adj}}/\Delta T$  from deep convection (Figure 3b) and underestimated the positive shortwave  $\Delta\text{CRE}_{\text{adj}}/\Delta T$  from stable regimes. This result did not change by restricting the comparison to years 2000–2006 for which CAM5-AMIP and CERES observations were both available, although the shorter period increased the uncertainty (not shown). A similar result suggesting a positive cloud feedback in stable regimes can be seen in Figure 2b, in terms of a





**Figure 4.** MODIS cloud fraction (%) separated into cloud top pressure and optical thickness bins for (a) stable ( $-3 < B \times 10^2 < -2$ ), (b) marginally stable ( $-2 < B \times 10^2 < 0$ ), (c) deep convective ( $0 < B \times 10^2 < 1$ ), and (d) all regimes. (e–h) As in Figures 4a–4d but for the cloud fraction response to warming estimated by the slope of the linear regression of cloud fraction anomalies against tropical average temperature anomalies.

reduction in (negative) shortwave CRE during the 2015–2016 El Niño, particularly for  $B$  between  $-3$  and  $-0.5$  ( $\times 10^{-2}$ ). Most of the discrepancy at  $B = 0.5$  is due to the observed longwave response being more negative, despite the model's shortwave CRE having larger mean state biases for  $B > 0$  (cf. Figures 2b and 2f). This implies that mean state biases as a function of  $B$  are not trivially related to the biases in the response to warming. Note that the larger error bars for CAM5-AMIP in Figure 3b are due to larger scatter about the linear fit.

We estimated contributions of convective regimes to the net  $\Delta \text{CRE}_{\text{adj}}/\Delta T$  (supporting information Table S1) by multiplying  $\Delta \text{CRE}_{\text{adj}}/\Delta T$  in Figure 3 by the relative frequency of  $B$  bins. Feedbacks from stable and deep convective clouds were nearly equal and of opposite sign, contributing to a smaller net feedback in the observations ( $-0.06 \pm 0.73 \text{ W m}^{-2} \text{ K}^{-1}$ ). The degree of this cancelation and the negative sign of the net  $\Delta \text{CRE}_{\text{adj}}/\Delta T$  from deep convection was not captured in CAM5. This biased response may be related to the higher climate sensitivity seen in some models [Mauritsen and Stevens, 2015]. However, CAM5 also underestimated the positive  $\Delta \text{CRE}_{\text{adj}}/\Delta T$  from stable regimes, such that the net  $\Delta \text{CRE}_{\text{adj}}/\Delta T$  summed across all regimes ( $0.35 \pm 0.65 \text{ W m}^{-2} \text{ K}^{-1}$ ) was within statistical uncertainty of the observations. The per Kelvin change in the probability of  $B$  regimes (Figure S7) was relatively small, contributing  $-0.15 \text{ W m}^{-2} \text{ K}^{-1}$  and  $-0.01 \text{ W m}^{-2} \text{ K}^{-1}$  to the observed and modeled  $\Delta \text{CRE}_{\text{adj}}/\Delta T$ , respectively. For tropical ocean and land together, we inferred net short-term cloud feedbacks of  $-0.51 \pm 0.58 \text{ W m}^{-2} \text{ K}^{-1}$  in observations, and  $-0.60 \pm 0.45 \text{ W m}^{-2} \text{ K}^{-1}$  in CAM5-AMIP, similar to previous estimates [Trenberth et al., 2010].

Since stable regimes are not guaranteed to be free of high clouds, we used MODIS (Moderate Resolution Imaging Spectroradiometer) Collection 5 data to identify clouds by CTP and optical thickness (Figures 4a–4d) [King et al., 2003; Pincus et al., 2012]. The results confirm that some high clouds (CTP  $< 440$  hPa) occur in stable regimes, while most high clouds and almost all optically thick high clouds occur in deep convective regimes diagnosed by  $B > 0$ . It is noted that these histograms may change with the more reliable Aqua MODIS Collection 6 [Yue et al., 2016].

We estimated the cloud response to warming (Figures 4e–4h) using the slope of the linear regression of cloud fraction anomalies against tropical average temperature anomalies [e.g., Zelinka and Hartmann, 2011]. High-cloud fraction declined with warming in deep convective regimes ( $-6.4 \pm 2.8\% \text{ K}^{-1}$ ) but increased in stable and marginally stable regimes ( $1.1 \pm 1.2\% \text{ K}^{-1}$ ). Given their opposite sign, these differences are unlikely to

be explained by differences between local and tropical average warming. Warming occurred in all regimes, although the deep convective regime warmed more (Figure S8). Furthermore, there is a distinct upward shift in high-cloud fraction in the  $B < 0$  regimes, but not in the  $B > 0$  regimes. High-cloud CTP decreased by  $31 \pm 20$  hPa  $K^{-1}$  in stable regimes but did not significantly change ( $0 \pm 24$  hPa  $K^{-1}$ ) in deep convective regimes, as estimated by regression of CTP anomalies against tropical average surface temperature anomalies. This indicates that different processes govern temperature responses of high clouds in deep convective regimes compared to those that occur remotely from deep convection. The latter may be formed in situ or may be advected into stable regimes from deep convective regimes.

The compensating changes in high-cloud fraction between stable and deep convective regimes resulted in smaller net changes in high clouds with warming (Figure 4h), that are consistent with previous studies [Zelinka and Hartmann, 2011; Zhou et al., 2013]. The reduction in deep convective high-cloud fraction is consistent with the inferred negative longwave cloud feedback in deep convective regimes (Figure 3a). Note that cloud responses in stable regimes were not statistically significant, in part due to compensating responses in different regions (Figure S1). It is also noted that feedbacks inferred from tropical interannual variability are not necessarily indicative of global feedbacks [Trenberth et al., 2010], and short-term cloud feedbacks may differ from long-term global warming cloud feedbacks [Zelinka and Hartmann, 2011; Zhou et al., 2015, 2016]. The  $B$  coordinate may provide some remedy for the timescale dependence of cloud feedbacks, by attempting to control for short-term variations in the large-scale circulation and its coupling to convective instability.

## 5. Conclusions

The observations confirm two predictions of climate models: As a function of subcloud buoyancy ( $B$ ), the threshold for deep convection is nearly invariant with tropics-wide warming, such that cloud radiative effect (CRE) during the warmest and coldest years tends to collapse onto a single curve in  $B$  coordinates. Second, the frequency distribution of  $B$  is approximately invariant with warming, thus constraining the relative frequency of deep convective regimes. Applying this diagnostic to short-term observations, we inferred a negative cloud feedback in deep convective regimes that was not represented in CAM5. Our results suggest that high clouds respond differently to warming in deep convective compared to stable regimes. Compensation between these responses contributed to the small magnitude and uncertain sign of the observed net tropical cloud radiative feedback inferred from short-term changes in adjusted CRE. Understanding the different processes governing cloud feedbacks in stable and deep convective regimes will be important in refining climate model representation of cloud feedbacks.

## Acknowledgments

The CERES EBAF-TOA Ed2.8 data set was obtained from the NASA Langley Research Center Atmospheric Science Data Center, at <http://ceres.larc.nasa.gov>. The ECMWF interim reanalysis data were obtained from ECMWF public data sets web interface at <http://www.ecmwf.int/en/research/climate-reanalysis/era-interim>. CAM5-AMIP data were obtained from <https://www.earthsystemgrid.org/dataset/ucar.cgd.cesm4.cam5.1.amip.1d.002.html>. Nauru radiosonde data were obtained from <https://doi.org/10.5439/1095390>. MODIS data were obtained from CFMIP observations for model evaluation (<http://climserv.ipsl.polytechnique.fr/cfmip-obs/>) at [ftp://laadsweb.nascom.nasa.gov/NetCDF/L3\\_Monthly/V02/](ftp://laadsweb.nascom.nasa.gov/NetCDF/L3_Monthly/V02/). We thank Mark Zelinka and an anonymous reviewer for helpful comments and suggestions. This material is based upon work supported in part by the U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research, under contract DE-AC02-05CH11231.

## References

- Bony, S., and J. L. Dufresne (2005), Marine boundary layer clouds at the heart of tropical cloud feedback uncertainties in climate models, *Geophys. Res. Lett.*, 32, L20806, doi:10.1029/2005GL023851.
- Bony, S., J. L. Dufresne, H. Le Treut, J. J. Morcrette, and C. Senior (2004), On dynamic and thermodynamic components of cloud changes, *Clim. Dyn.*, 22(2–3), 71–86, doi:10.1007/s00382-003-0369-6.
- Bony, S., B. Stevens, D. Coppin, T. Becker, K. A. Reed, A. Voigt, and B. Medeiros (2016), Thermodynamic control of anvil cloud amount, *Proc. Natl. Acad. Sci. U.S.A.*, 113(32), 8927–8932, doi:10.1073/pnas.1601472113.
- Bretherton, C. S. (2015), Insights into low-latitude cloud feedbacks from high-resolution models, *Philos. Trans. R. Soc. A*, 373, 20140415, doi:10.1098/rsta.2014.0415.
- Ciesielski, P. E., R. H. Johnson, and J. Wang (2009), Correction of humidity biases in Vaisala RS80-H sondes during NAME, *J. Atmos. Ocean. Technol.*, 26(9), 1763–1780, doi:10.1175/2009JTECHA1222.1.
- Dee, D. P., et al. (2011), The ERA-Interim reanalysis: Configuration and performance of the data assimilation system, *Q. J. R. Meteorol. Soc.*, 137(656), 553–597, doi:10.1002/qj.828.
- Dessler, A. E. (2010), A determination of the cloud feedback from climate variations over the past decade, *Science*, 330(6010), 1523–1527, doi:10.1126/science.1192546.
- Dessler, A. E., and N. G. Loeb (2013), Impact of dataset choice on calculations of the short-term cloud feedback, *J. Geophys. Res. Atmos.*, 118, 2821–2826, doi:10.1002/jgrd.50199.
- Fu, R., A. D. Delgenio, W. B. Rossow, and W. T. Liu (1992), Cirrus-cloud thermostat for tropical sea-surface temperatures tested using satellite data, *Nature*, 358(6385), 394–397, doi:10.1038/358394a0.
- Gregory, J. M., W. J. Ingram, M. A. Palmer, G. S. Jones, P. A. Stott, R. B. Thorpe, J. A. Lowe, T. C. Johns, and K. D. Williams (2004), A new method for diagnosing radiative forcing and climate sensitivity, *Geophys. Res. Lett.*, 31, L03205, doi:10.1029/2003GL018747.
- Hartmann, D. L., and M. L. Michelsen (1993), Large-scale effects on the regulation of tropical sea-surface temperature, *J. Clim.*, 6(11), 2049–2062, doi:10.1175/1520-0442(1993)006<2049:LSEOTR>2.0.CO;2.
- Hartmann, D. L., and M. L. Michelsen (2002), No evidence for iris, *Bull. Am. Meteorol. Soc.*, 83(2), 249–254, doi:10.1175/1520-0477(2002)083<0249:NEFI>2.3.CO;2.
- Kiehl, J. (1994), On the observed near cancellation between longwave and shortwave cloud forcing in tropical regions, *J. Clim.*, 7(4), 559–565, doi:10.1175/1520-0442(1994)007<0559:OTONCB>2.0.CO;2.
- King, M. D., et al. (2003), Cloud and aerosol properties, precipitable water, and profiles of temperature and humidity from MODIS, *IEEE Trans. Geosci. Remote Sens.*, 41, 442–458.

- Klein, S. A., and D. L. Hartmann (1993), The seasonal cycle of low stratiform clouds, *J. Clim.*, 6(8), 1587–1606, doi:10.1175/1520-0442(1993)006<1587:tscols>2.0.co;2.
- Lau, K. M., C. H. Sui, M. D. Chou, and W. K. Tao (1994), An inquiry into the cirrus-cloud thermostat effect for tropical sea-surface temperature, *Geophys. Res. Lett.*, 21(12), 1157–1160, doi:10.1029/94GL00222.
- Lau, K. M., H. T. Wu, and S. Bony (1997), The role of large-scale atmospheric circulation in the relationship between tropical convection and sea surface temperature, *J. Clim.*, 10(3), 381–392, doi:10.1175/1520-0442(1997)010<0381:trolsa>2.0.co;2.
- Lindzen, R. S., M. D. Chou, and A. Y. Hou (2001), Does the Earth have an adaptive infrared iris?, *Bull. Am. Meteorol. Soc.*, 82(3), 417–432, doi:10.1175/1520-0477(2001)082<0417:dtehaa>2.3.co;2.
- Long, C. N., S. A. McFarlane, A. D. Genio, P. Minnis, T. P. Ackerman, J. Mather, J. Comstock, G. G. Mace, M. Jensen, and C. Jakob (2013), ARM research in the equatorial western Pacific: A decade and counting, *Bull. Am. Meteorol. Soc.*, 94(5), 695–708, doi:10.1175/BAMS-D-11-00137.1.
- Luo, Z., and W. B. Rossow (2004), Characterizing tropical cirrus life cycle, evolution, and interaction with upper-tropospheric water vapor using lagrangian trajectory analysis of satellite observations, *J. Clim.*, 17(23), 4541–4563, doi:10.1175/3222.1.
- Mauritsen, T., and B. Stevens (2015), Missing iris effect as a possible cause of muted hydrological change and high climate sensitivity in models, *Nat. Geosci.*, 8(5), 346–351, doi:10.1038/ngeo2414.
- Medeiros, B., and B. Stevens (2011), Revealing differences in GCM representations of low clouds, *Clim. Dyn.*, 36(1–2), 385–399, doi:10.1007/s00382-009-0694-5.
- Pincus, R., S. Platnick, S. A. Ackerman, R. S. Hemler, and R. J. P. Hofmann (2012), Reconciling simulated and observed views of clouds: MODIS, ISCCP, and the limits of instrument simulators, *J. Clim.*, 25(13), 4699–4720, doi:10.1175/JCLI-D-11-00267.1.
- Ramanathan, V., and W. Collins (1991), Thermodynamic regulation of ocean warming by cirrus clouds deduced from observations of the 1987 El Niño, *Nature*, 351(6321), 27–32, doi:10.1038/351027a0.
- Ramanathan, V., R. D. Cess, E. F. Harrison, P. Minnis, B. R. Barkstrom, E. Ahmad, and D. Hartmann (1989), Cloud-radiative forcing and climate: Results from the Earth radiation budget experiment, *Science*, 243(4887), 57–63, doi:10.1126/science.243.4887.57.
- Randall, D. A. (2015), *An Introduction to the Global Circulation of the Atmosphere*, 447 pp., Princeton Univ. Press, Princeton, N. J.
- Reynolds, R. W., and T. M. Smith (1994), Improved global sea-surface temperature analyses using optimum interpolation, *J. Clim.*, 7(6), 929–948, doi:10.1175/1520-0442(1994)007<0929:igssta>2.0.co;2.
- Reynolds, R. W., N. A. Rayner, T. M. Smith, D. C. Stokes, and W. Q. Wang (2002), An improved in situ and satellite SST analysis for climate, *J. Clim.*, 15(13), 1609–1625, doi:10.1175/1520-0442(2002)015<1609:aisas>2.0.co;2.
- Shell, K. M., J. T. Kiehl, and C. A. Shields (2008), Using the radiative kernel technique to calculate climate feedbacks in NCAR's Community Atmospheric Model, *J. Clim.*, 21(10), 2269–2282, doi:10.1175/2007JCLI2044.1.
- Sherwood, S. C., S. Bony, and J.-L. Dufresne (2014), Spread in model climate sensitivity traced to atmospheric convective mixing, *Nature*, 505(7481), 37–42, doi:10.1038/nature12829.
- Sobel, A. H., and H. Gildor (2003), A simple time-dependent model of SST hot spots, *J. Clim.*, 16(23), 3978–3992, doi:10.1175/1520-0442(2003)016<3978:astmos>2.0.co;2.
- Soden, B. J., and G. A. Vecchi (2011), The vertical distribution of cloud feedback in coupled ocean-atmosphere models, *Geophys. Res. Lett.*, 38, L12704, doi:10.1029/2011GL047632.
- Trenberth, K. E., J. T. Fasullo, C. O'Dell, and T. Wong (2010), Relationships between tropical sea surface temperature and top-of-atmosphere radiation, *Geophys. Res. Lett.*, 37, L03702, doi:10.1029/2009GL042314.
- Tsushima, Y., S.-I. Iga, H. Tomita, M. Satoh, A. T. Noda, and M. J. Webb (2014), High cloud increase in a perturbed SST experiment with a global nonhydrostatic model including explicit convective processes, *J. Adv. Model. Earth Syst.*, 6(3), 571–585, doi:10.1002/2013MS000301.
- Waliser, D. E. (1996), Formation and limiting mechanisms for very high sea surface temperature: Linking the dynamics and the thermodynamics, *J. Clim.*, 9(1), 161–188, doi:10.1175/1520-0442(1996)009<0161:falmfv>2.0.co;2.
- Wallace, J. M. (1992), Effect of deep convection on the regulation of tropical sea-surface temperature, *Nature*, 357(6375), 230–231, doi:10.1038/357230a0.
- Webb, M. J., et al. (2006), On the contribution of local feedback mechanisms to the range of climate sensitivity in two GCM ensembles, *Clim. Dyn.*, 27(1), 17–38, doi:10.1007/s00382-006-0111-2.
- Webb, M. J., et al. (2015), The impact of parametrized convection on cloud feedback, *Philos. Trans. A Math. Phys. Eng. Sci.*, 373, 2054, doi:10.1098/rsta.2014.0414.
- Wetherald, R. T., and S. Manabe (1988), Cloud feedback processes in a general circulation model, *J. Atmos. Sci.*, 45(8), 1397–1416, doi:10.1175/1520-0469(1988)045<1397:CFPIAG>2.0.CO;2.
- Wielicki, B. A., B. R. Barkstrom, E. F. Harrison, R. B. Lee, G. L. Smith, and J. E. Cooper (1996), Clouds and the Earth's radiant energy system (CERES): An Earth observing system experiment, *Bull. Am. Meteorol. Soc.*, 77(5), 853–868, doi:10.1175/1520-0477(1996)077<0853:catere>2.0.co;2.
- Williams, I. N., R. T. Pierrehumbert, and M. Huber (2009), Global warming, convective threshold and false thermostats, *Geophys. Res. Lett.*, 36, L21805, doi:10.1029/2009GL039849.
- Wood, R., and C. S. Bretherton (2006), On the relationship between stratiform low cloud cover and lower-tropospheric stability, *J. Clim.*, 19(24), 6425–6432, doi:10.1175/jcli3988.1.
- Yue, Q., B. H. Kahn, E. J. Fetzer, S. Wong, R. Frey, and K. G. Meyer (2016), On the response of MODIS cloud coverage to global mean surface air temperature, *J. Geophys. Res. Atmos.*, 121, doi:10.1002/2016JD025174.
- Zelinka, M. D., and D. L. Hartmann (2011), The observed sensitivity of high clouds to mean surface temperature anomalies in the tropics, *J. Geophys. Res.*, 116, D23103, doi:10.1029/2011JD016459.
- Zhou, C., M. D. Zelinka, A. E. Dessler, and P. Yang (2013), An analysis of the short-term cloud feedback using MODIS data, *J. Clim.*, 26(13), 4803–4815, doi:10.1175/JCLI-D-12-00547.1.
- Zhou, C., M. D. Zelinka, A. E. Dessler, and S. A. Klein (2015), The relationship between interannual and long-term cloud feedbacks, *Geophys. Res. Lett.*, 42, 10,463–10,469, doi:10.1002/2015GL066698.
- Zhou, C., M. D. Zelinka, and S. A. Klein (2016), Impact of decadal cloud variations on the Earth's energy budget, *Nat. Geosci.*, 9, 871–874, doi:10.1038/ngeo2828.