

## **Optimal Resource for Children’s Surgical Care: Executive Summary**

November 18<sup>th</sup>, 2018

**Authors:** Global Initiative for Children’s Surgery (GICS) Collaborators

### **Global Initiative for Children’s Surgery Collaborators List:**

David Grabski, Emmanuel Ameh, Doruk Ozgediz, Keith Oldham, Francis A. Abantanga, Mohamed Abdelmalak, Nurudeen Abdulraheem, Niyi Ade-Ajayi, Edna Adan Ismail, Adesoji Ademuyiwa, Eltayeb Ahmed, Sunday Ajike, Olugbemi Benedict Akintububo, Felix Alakaloko, Brendan Allen, Vanda Amado, Shanthi Anbuselvan, Jamie Anderson, Theophilus Teddy Kojo Anyomih, Leopold Asakpa, Gudeta Assegie, Jason Axt, Ruben Ayala, Frehun Ayele, Harshjeet Singh Bal, Rouma Bankole, Tahmina Banu, Tim Beacon, Zaitun Bokhari, Hiranya Kumar Borah, Eric Borgstein, Nick Boyd, Jason Brill, Britta Budde-Schwartzman, Fred Bulamba, Marilyn Butler, Bruce Bvulani, Sarah Cairo, Juan Francisco Campos Rodezno, Massimo Caputo, Milind Chitnis, Maija Cheung, Bruno Cigliano, Damian Clarke, Michael Cooper, Tessa Concepcion, Scott Corlew, David Cunningham, Sergio D’Agostino, Shukri Dahir, Bailey Deal, Miliard Derbew, Sushil Dhungel, David Drake, Elizabeth Drum, Bassey Edem, Stella Eguma, Olumide Elebute, Beda R. Espineda, Samuel Espinoza, Faye Evans, Omolara Faboya, Diana Farmer, Jacques Fadhili Bake, Tatiana Fazecas, Mohammad Rafi Fazli, Graham Fieggen, Anthony Figaji, Jean Louis Fils, Tamara Fitzgerald, Randall Flick, Gacelle Fossi, George Galiwango, Lily Saldaña Gallo, Mike Ganey, Zipporah Gathuya, Maryam Ghavami Adel, Vafa Ghorban Sabagh, Sridhar Gibikote, Hetal Gohil, Laura Goodman, Sarah Greenberg, Russell Gruen, Lars Hagander, Rahimullah Hamid, Erik Hansen, William Harkness, Mauricio Herrera, Intisar Hisham, Andrew Hodges, Sarah Hodges, Ai Xuan Holterman, Andrew Howard, Romeo Ignacio, Dawn Ireland, Enas Ismail, Rebecca Jacob, Anette Jacobsen, Zahra Jaffry, Deeptiman James, Ebor Jacob James, Adiyasuren Jamiyanjav, Kathy Jenkins, Guy Jensen, Maria Jimenez, Tarun John K Jacob, Walter Johnson, Anita Joselyn, Kokila Lakhoo, Nasser Kakembo, Patrick Kamalo, Neema Kaseje, Bertille Ki, Phyllis Kisa, Peter Kim, Krishna Kumar, Rashmi Kumar, Charlotte Kvasnovsky, Ananda Lamahewage, Monica Langer, Christopher Lavy, Taiwo Lawal, Colin Lazarus, Andrew Leather, Chelsea Lee, Basil Leodoro, Allison Linden, Katrine Lofberg, Jonathan Lord, Jerome Loveland, Leecarlo Millano Lumban Gaol, Vrisha Madhuri, Pavrette Magdala, Luc Kalisyia Malemo, Aesha Malik, John Mathai, Marcia Matias, Bothwell Mbuwayesango, Merrill McHoney, Liz McLeod, Ashish Minocha, Charles Mock, Mubarak Mohamed, Ivan Molina, Ashika Morar, Zahid Mukhtar, Mulewa Mulenga, Bhargava Mullapudi, Jack Mulu, Byambajav Munkhjargal, Arlene Muzira, Mary Nabukenya, Mark Newton, Jessica Ng, Karissa Nguyen, Laurence Isaaya Ntawunga, Peter M. Nthumba, Alp Numanoglu, Benedict Nwomeh, Kristin Ojomo, Maryrose Osazuwa, Emmanuel Owusu Abem, Shazia Peer, Norgrove Penny, Dan Poenaru, Robin Petroze, Vithya Priya, Ekta Rai, Lola Raji, Vinitha Paul Ravindran, Desigen Reddy, Henry Rice, Yona Ringo, Amezene Robelie, Jose Roberto Baratella, David Rothstein, Coleen Sabatini, Soumitra Saha, Saurabh Saluja, Lubna Samad, John Sekabira, Justina Seyi-Olajide, Bello B. Shehu, Sabina Siddiqui, Ritesh Shrestha, David Sigalet, Martin Situma, Emily Smith, Adrienne Soggi, David Spiegel, Peter Ssenyonga, Jacob Stephenson, Richard Stewart, Erin Stieber, Etienne St-Louis, Vinayak Shukla, Thomas Sims, Faustin Felicien Mouafo Tambo, Robert Tamburro, Mansi Tara, Ahmad Tariq, Reju Thomas, Leopold Torres Contreras, Stephen Ttendo, Benno Ure, Luca Vricella, Luis Vasquez, Vijayakumar Raju, Jorge Villacis, Gustavo Villanova, Catherine deVries, Amira Waheeb, Saber Waheeb, Albert Wandaogo, Anne Wesonga, Omolara Williams, Sigal Willner, Nyo Nyo Win, Hussein Wissanji, Paul Mwindekuma Wondoh, Garreth Wood, Naomi Wright, Benjamin Yapo, George Youngson, Yasmine Yousef, Denléwendé Sylvain Zabsonre, Luis Enrique Zea Salazar, Adiyasuren Zevee, Bistra Zheleva, Stephen Bickler

## Executive Summary

There is growing recognition that safe surgery and anesthesia should be an essential component of universal health care in low- and middle-income countries (LMICs). Landmark studies in 2015, by the Disease Control Priorities Project (DCP3) and Lancet Commission on Global Surgery (LCoGS) were important in defining the scale and scope of the surgical burden worldwide, priority surgical interventions, and economic impact of surgical care<sup>1,2</sup>. Shortly after the release of their two reports, the World Health Assembly passed resolution 68/15 designed to strengthen emergency and essential surgical and anesthesia care as a component of universal health coverage<sup>3</sup>. To facilitate implementation of the DCP3 and LCoGS recommendations a growing number of countries are developing national surgical, obstetric and anesthesia plans<sup>4</sup>. National surgical plans are an iterative process where stakeholders use country-level data to develop a contextually relevant and sustainable plan to ensure that surgical and obstetrical services are available for an entire country or region.

Of the 4.8 billion without access to surgical care, 1.7 billion are children and adolescents<sup>2,5</sup>. The vast majority of these children and adolescents live in LMICs, where the pediatric age group represents a disproportionately larger fraction of the population. In these settings, need for children's surgical care is high and poor care results in significant morbidity and mortality<sup>6</sup>. A child born in sub-Saharan Africa has an estimated 85% risk of requiring surgical care by age 15 years<sup>7</sup>. Improving surgical for children in LMICs can achieve several important goals, including early identification of correctable congenital anomalies and other surgical conditions, treatment of life-threatening injuries and burns, minimizing acute and long-term suffering, protecting

1  
2  
3 families from substantial economic losses, and increasing economic productivity. For  
4 many childhood diseases including congenital anomalies, cancer, injuries and other  
5 emergencies, surgery remains the only curative modality in low resource regions of the  
6 world. Surgical care for children also has a role in achieving health-related Sustainable  
7 Development Goals and targets, in particular 1) Ending preventable deaths in newborns  
8 and children under five, 2) Reduction in death and disability for road traffic injuries and  
9 non-communicable diseases, 3) Ensuring universal health coverage, and 4) Scaling up the  
10 health workforce <sup>8</sup>.

11  
12 The Global Initiative for Children's Surgery (GICS) is a consortium of providers,  
13 institutions, and allies from both resource-rich and poor settings, who envision a future  
14 where every child will have access to surgical care <sup>9</sup>. GICS includes representation from  
15 13 pediatric specialties: anesthesia, orthopedic surgery, neurosurgery, plastic surgery,  
16 urology, cardiac surgery, critical care, general pediatric surgery, otolaryngology,  
17 radiology, ophthalmology, oral and maxillofacial surgery and nursing. The goals of  
18 GICS are to: 1) promote the inclusion of children's surgical care within existing surgical  
19 and global child health initiatives; 2) provide strategic direction among the many  
20 children's surgery sub-specialties; and 3) identify high-priority needs and solutions from  
21 the perspective of local providers.

22  
23 At the inaugural GICS meeting in London, England (May 2016) providers from  
24 LMICs expressed the need for guidelines that could be adapted for use in their local  
25 environment, especially for the purpose of advocating for increased resources. The  
26 Optimal Resources for Surgical Care (OReCS) project thus evolved to provide guidelines  
27 for essential surgical care for children in LMICs. A document was drafted and refined at  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 the second and third GICS meetings (October 2016 and January 2018) with ongoing  
4 specialty expert input through multiple working groups. The OReCS document is divided  
5 into two parts: Part 1: Guidelines for Different Levels of Care, and Part II: Supplies,  
6 Equipment and Infrastructure. Both are included as Supplementary Material. The  
7  
8 OReCS document outlines the resources required to care for children with surgical  
9  
10 conditions at all levels of the health care system (Table 1). The resources needed for  
11  
12 different levels of children's surgical care are outlined in the form of templates, which are  
13  
14 designed to support basic, intermediate and complex/advanced surgical care for children  
15  
16 in low resource settings. The goal of these templates is to provide a guide for minimum  
17  
18 standards for those planning surgical services for children in low-resource settings. Thus,  
19  
20 the OReCS document provides pragmatic, yet comprehensive recommendations  
21  
22 regarding the resources, training and research priorities required for safe and high-quality  
23  
24 surgical services for children in LMICs.  
25  
26  
27  
28  
29  
30  
31  
32

33 In conclusion, GICS convened a meeting of key stakeholders, led by LIMIC  
34 surgical and anesthesia providers, to develop guidelines for emergency and essential  
35 surgical services for children. The resulting product was the OReCS document which  
36 describes the resources needed to care for children with surgical problems at all levels of  
37 the health care system in LMICs. The document provides a strategy for integrating  
38 children's surgical care into national surgery, obstetric and anesthesia plans, or for  
39 developing surgical services within a child health policy/plan. The OReCS document can  
40 also serve as an important starting point for gap analysis, quality improvement, hospital  
41 assessment, training curricula, and for advocacy for children's surgical care. The OReCS  
42 program provides and important opportunity to mobilize the diverse community of  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

international stakeholders to both implement and integrate children’s surgery into existing and future global health initiatives.

For Peer Review

## References

1. Mock CN, Donkor P, Gawande A, Jamison DT, Kruk ME, Debas HT. Essential surgery: key messages from Disease Control Priorities, 3rd edition. *Lancet* 2015; **385**(9983): 2209-19.
2. Meara JG, Leather AJ, Hagander L, et al. Global Surgery 2030: evidence and solutions for achieving health, welfare, and economic development. *Lancet* 2015.
3. Strengthening emergency and essential surgical care and anaesthesia as a component of universal health coverage. World Health Organization 2015. Available from: <http://apps.who.int/medicinedocs/documents/s21904en/s21904en.pdf> (Accessed August 25, 2018)
4. Surgical Care Systems Strengthening: Developing national surgical, obstetric and anaesthesia plans. World Health Organization 2017. Available from: <http://apps.who.int/iris/bitstream/handle/10665/255566/9789241512244-eng.pdf> (Accessed August 25, 2018).
5. Mullapudi B, Grabski D, Ameh E, Ozgediz D, Bickler SW. 1.7 billion children and adolescents lack access to surgical and anesthesia care worldwide. 2018. Unpublished Manuscript.
6. Bickler SW, Rode H. Surgical services for children in developing countries. *Bull World Health Organ* 2002; **80**(10): 829-35.
7. Bickler SW, Telfer ML, Sanno-Duanda B. Need for paediatric surgery care in an urban area of The Gambia. *Tropical doctor* 2003; **33**(2): 91-4.
8. Henry JA, Abdullah F. Global Surgical Care in the U.N. Post-2015 Sustainable Development Agenda. *World J Surg* 2016; **40**(1): 1-5.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

9. Goodman LF, St-Louis E, Yousef Y, et al. The Global Initiative for Children's Surgery: Optimal Resources for Improving Care. *Eur J Pediatr Surg* 2017.

For Peer Review

## Key Messages

- Surgical care has an incontrovertible, crosscutting role in achieving child health.
- Children have unique surgical conditions, present unique anesthetic challenges, and have special perioperative needs.
- The Optimal Resources for Children's Surgery document provides a strategy for integrating surgical care for children into National Surgical, Obstetric and Anesthesia Plans.
- There is an important opportunity to prevent death and reduce disability in children by scaling up surgical care in low- and middle-income countries.

For Peer Review

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

# OPTIMAL RESOURCES FOR CHILDREN'S SURGICAL CARE

## I. Guidelines for Different Levels of Care



**GICS**

**Global Initiative for Children's Surgery**

*Version 2.0 (September 24, 2018)*

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

GICS members contributing to this document (alphabetical order):

Francis Abantanga, Mohamed Abdelmalak, Nurudeen Toyin Abdulraheem, Niyi Ade-Ajayi, Sunday Ajike, Olugbemi Benedict Akintububo, Brendan Allen, Vanda Amado, Emmanuel Ameh, Shanthi Anbuselvan, Jamie Anderson, Theophilus Kojo Anyomih, Leopold Asakpa, Gudeta Assegie, Ruben Ayala, Frehun Ayele, Jacques Bake, Harshjeet Singh Bal, Roumanatou Bankole, Tahmina Banu, Tim Beacon, Stephen Bickler, Zaitun Bokhari, Hiranya Kumar Borah, Eric Borgstein, Nick Boyd, Jason Brill, Britta Budde-Schwartzman, Marilyn Butler, Sarah Cairo, Juan Francisco Campos-Rodezno, Massimo Caputo, Maija Cheung, Milind Chitnis, Bruno Cigliano, Lee Ann Clark, Tessa Concepcion, Michael Cooper, David Cunningham, Sergio D'Agostino, Shugri Dahir, Miliard Derbew, Catherine DeVries, Sushil Dhungel, Elizabeth Drum, Basse Edem, Stella Eguma, Olumide Elebute, Beda Espineda, Ivan Espinoza Tristan, Faye Evans, Diana Farmer, Tatiana Fazecas, Mohammad Rafi Fazli, Tony Figaji, Gacelle Fossi, George Galiwango, Lily Saldaña Gallo, Zipporah Gathuya, Maryam Ghavami Adel, Vafa Ghorban Sabagh, Sridhar Gibikote, Hetal Gohil, Laura Goodman, David Grabski, Sarah Greenberg, Russel Gruen, Rahimullah Hamid, Erik Hansen, William Harkness, Mauricio Herrera, Intisar Hisham, Andrew Hodges, Sarah Hodges, Aixuan Holterman, Andrew Howard, Romeo Ignacio, Dawn Ireland, Edna Aden Ismail, Enas Ismail, Rebecca Jacob, Anette Jacobsen, Deeptiman James, Ebor Jacob James, Adiyasuren Jamiyanjav, Kathy Jenkins, Guy Jensen, Tarun John Jacob, Walter Johnson, Anita Joselyn, Patrick Kamalo, Neema Kaseje, Bertille Ki, Krishna Kumar, Rashmi Kumar, Kokila Lakhoo, Ananda Lamahewage, Monica Langer, Chris Lavy, Colin Lazarus, Andrew Leather, Basil Leodoro, Allison Linden, Katrine Lofberg, Jonathan Lord, Jerome Loveland, Vrisha Madhuri, Luc Kalisya Malemo, Aeesha Malik, John Mathai, Marcia Matias, Bothwell Mbuwayesango, Elizabeth McLeod, Leecarlo Millano, Ashish Minocha, Mubarak Mohamed, Faustin Felicien Mouafo Tambo, Zahid Mukhtar, Mulewa Mulenga, Bhargava Mullapudi, Jack Mulu, Mary Nabukenya, Mark Newton, Jess Ng, Karissa Nguyen, Shehu Ntawunga, Peter Nthumba, Alp Numanoglu, Keith Oldham, Maryrose Osazuwa, Emmanuel Abem Owusu, Doruk Ozgediz, Vinitha Paul Ravindran, Shazia Peer, Norgrove Penny, Robin Petroze, Marina Petrulla, Dan Poenaru, Vithya Priya, Ekta Rai, Vijayakumar Raju, Desigen Reddy, Dean Rex, Henry Rice, Yona Ringo, Amezene Robelie, David Rothstein, Bankole Rouma, Coleen Sabatini, Soumitra Saha, Saurabh Saluja, Lubna Samad, John Sekabira, Bello Shehu, Ritesh Shrestha, Vinayak Shukla, Sabina Siddiqui, David Sigalet, Cicero Silva, Thomas Sims, Emily Smith, David Spiegel, Peter Ssenyonga, Jacob Stephenson, Richard Stewart, Erin Stieber, Etienne St-Louis, Robert Tamburro, Mansi Tara, Ahmad Tariq, Roy Thankchan, Reju Thomas, Stephen Ttendo, Benno Ure, C. Luis Vasquez, Jorge Villacis, Amira Waheeb, Saber Waheeb, Albert Wandaogo, Anne Wesonga, Sigal Willner, Nyo Nyo Win, Hussein Wissanji, Paul Mwindekuma Wondoh, Garreth Wood, Naomi Wright, Benjamin Yapo, George Youngson, Yasmine Yousef, Denléwendé Sylvain Zabsonre, Luis Enrique Zea Salazar, Adiyasuren Zevee, Bistra Zheleva

## Contents

1		
2		
3		
4	1. KEY MESSAGES.....	4
5	2. SUMMARY.....	5
6	3. BACKGROUND.....	6
7		
8	3.1 Introduction.....	6
9	3.2 Goals of the Optimal Resources for Children’s Project.....	7
10	3.3 Justification.....	8
11	4. SURGICAL NEEDS OF CHILDREN.....	9
12		
13	4.1 Epidemiology of childhood surgical conditions.....	9
14	4.2 Benefits of providing children’s surgical care.....	10
15	5. STRUCTURE OF HEALTH SYSTEMS IN LOW AND MIDDLE-INCOME COUNTRIES.....	11
16		
17	5.1. Overview.....	11
18	5.2 Types of facilities.....	11
19		
20	5.2.1 Health center.....	12
21	5.2.2 First-level hospital.....	12
22	5.2.3 Referral hospitals.....	13
23	6. LEVELS OF CHILDREN’S SURGICAL CARE.....	16
24		
25	6.1 Overview.....	16
26	6.2 Levels of surgical care.....	16
27		
28	6.2.1 Basic surgical care.....	16
29	6.2.2 Intermediate surgical care.....	17
30	6.2.3 Complex/advanced surgical care.....	17
31	6.3. Suggested levels of care by facility type.....	17
32	7. OPTIMAL RESOURCES FOR CHILDREN’S SURGICAL CARE BY LEVEL OF CARE.....	20
33		
34	7.1 Overview.....	20
35	7.2 Resource templates.....	20
36		
37	7.2.1 Optimal resources for basic surgical care (health center).....	21
38	7.2.2 Optimal resources for intermediate surgical care (first-level hospital).....	22
39	7.2.3 Optimal resources for advanced/complex surgical care (referral hospital).....	22
40	7.2.4 Optimal resources for Children’s Anesthesia Care.....	23
41	8. STRATEGIES FOR IMPROVING CHILDREN’S SURGICAL CARE.....	29
42		
43	8.1 Training.....	29
44	8.2 Quality improvement.....	31
45		
46	8.2.1 Structural improvement.....	32
47	8.2.2 Process improvement.....	32
48	8.2.3 Morbidity and mortality conferences.....	33
49	8.2.4 Preventable death studies.....	33
50	8.2.5 Complications.....	33
51	8.2.6 Risk-adjusted mortality.....	33
52		
53		
54		
55		
56		
57		
58		
59		
60		

1		
2		
3	8.2.7 Data collection and surveillance.....	34
4	8.3 Hospital assessment.....	34
5		
6	9. RESEARCH .....	36
7	9.1 Overview .....	36
8	9.2 Regional research hubs.....	36
9		
10	9.3 Children’s surgical database .....	37
11	10. INTEGRATION OF CHILDREN’S SURGERY INTO HEALTH SYSTEMS.....	39
12		
13	11. INTERACTION WITH AND COORDINATION OF STAKEHOLDERS .....	39
14	REFERENCES.....	41
15	APPENDIX 1: Childhood surgical procedures for different facility types classified by pediatric	
16	surgical subspecialty .....	44
17		
18	APPENDIX 2: Childhood surgical procedures for different facility types classified by Surgical	
19	Condition.....	46
20	APPENDIX 3. Human Resource for different types of facilities .....	48
21		
22		
23		
24		
25		
26		
27		
28		
29		
30		
31		
32		
33		
34		
35		
36		
37		
38		
39		
40		
41		
42		
43		
44		
45		
46		
47		
48		
49		
50		
51		
52		
53		
54		
55		
56		
57		
58		
59		
60		

## 1. KEY MESSAGES

---

- Surgical care has an incontrovertible, crosscutting role in achieving child health.
- Children have unique surgical conditions, present unique anesthetic challenges, and have special perioperative needs.
- The Optimal Resources for Children's Surgery document provides a strategy for integrating surgical care for children into National Surgical, Obstetric and Anesthesia Plans.
- There is an important opportunity to prevent death and reduce disability in children by scaling up surgical care in low- and middle-income countries.

For Peer Review

## 2. SUMMARY

Level of care	Type of facility (DCP3 classification)	Responsibilities	Age Treated	General Anesthesia	OPTIMAL RESOURCES FOR CHILDREN'S SURGICAL CARE				Quality and safety
					Human resources	Required skills	Infrastructure	Equipment & supplies	
Basic	Health center	Screening for surgical disease Resuscitation Referral to higher levels of care	All	No	Non-physician clinician Community Health Workers	Basic assessment and treatment skills	Infrastructure to support basic services	Wound care supplies	CME/CPD Periodic supervision and mentoring
Intermediate	First-level hospital	24/7 Emergency surgical care  Diagnosis and treatment of common surgical diseases	All	Yes, not including complex cases and minimal comorbidity (Limit of ASA I or II)	Physician Anesthesiologist +/- anesthesia provider and perioperative nursing	Skills to treat emergency and essential childhood surgical conditions Anesthesia +perioperative care	Children's ward Functional operating room and recovery area	Emergency and essential surgical equipment & supplies for children	M&M review
Complex/ Advanced	Referral hospital: Second- level	Selected more complex cases	All	Yes, including some complex cases with comorbidities (Limit of ASA III)	General surgeon often present Specialists in some areas of children's surgical care	Surgical, anesthesia, nursing, and ancillary staff with advanced skills in the majority of children's surgical care	Children's clinics High-dependency units	Equipment and supplies to support selected more complex services	Trauma conference  Tumor Board
	Referral hospital: Third-level	Comprehensive surgical care for children <sup>2</sup>	All	Yes, including complex cases with comorbidities (All ASA)	Specialists in all areas of children's surgical care; Advanced nursing/allied health & Pediatric anesthesiologist	Surgical, anesthesia, nursing, and ancillary staff with advanced skills in all areas of children's surgical care	Neonatal and Pediatric Intensive Care Units (NICU/PICU) Burn Unit	Equipment and supplies to support full spectrum of children's surgical care	As above
	Referral hospital: National Children's hospital	Multidisciplinary and chronic care  Training, education, research in all children's surgical specialties  Development of standards of care & Advocacy	All	As above	As above	As above Training in multidisciplinary and chronic care	As above	As above	As above  Multi-disciplinary program review

1. Levels of care defined as Basic (I), Intermediate (II) and Complex/Advanced (III). Higher levels assume presence of infrastructure/ services at lower levels
2. Type of complex/advanced care delivered at second and third level hospital and national children's hospital may vary considerably by resources available in a country
3. CME: Continuing medical education; CPD: Continuing professional development; M&M- Morbidity and Mortality; ASA- American Society of Anesthesia classification

### 3. BACKGROUND

---

#### 3.1 Introduction

Despite remarkable progress in improving the health of children worldwide during the past 25 years, there are areas in which progress has been much slower.<sup>1</sup> Surgical care of children is one of the most critical examples. In many low-resource settings, surgical care of children continues to be viewed as a non-essential component of maternal-child health programs and a small part of the health needs of the pediatric population. In the absence of surgical care, congenital anomalies go unrepaired, treatable injuries result in lifelong disabilities, and children die of easily correctable surgical problems, e.g., airway foreign bodies, incarcerated inguinal hernias, and abdominal emergencies.<sup>2</sup>

In the past several years, the Essential Surgery Volume of the Disease Control Priorities 3<sup>rd</sup> Edition (DCP3), Lancet Commission on Global Surgery and other stakeholders have highlighted the critical need for improving surgical care worldwide.<sup>3,4</sup> Subsequently, the World Health Assembly (WHO) passed resolution (A68/15) on strengthening emergency and essential surgical care and anesthesia as a component of universal health coverage (UHC).<sup>5,6</sup> None, however, have explicitly addressed the unique surgical needs of children and the resources required to meet these needs in low-resource settings, where children represent up to 40-50% of the population. Global initiatives directed at reducing the high rate of infant and under-five mortality rate of children has also overlooked the surgical needs of children.

The Global Initiative for Children's Surgery (GICS) is a consortium of providers, institutions, and allies from both resource-rich and low-resource settings, who envision a future where every child will have access to surgical care.<sup>7</sup> This optimal resources document has been developed by GICS and represents consensus of worldwide leaders and providers of children's surgery in low-resource settings targeting surgical care in low-resource settings. Importantly, the

1  
2  
3 recently updated WHO-WFSA (World Federation Of Societies Of Anesthesiologists) document  
4 detailing safe practice of anesthesia and perioperative care was incorporated into this work  
5 through the anesthesia working group of GICS.<sup>8</sup> This document provides standards for service  
6 delivery through the health system and defines types of anesthesia providers. GICS recognizes that  
7 surgery, anesthesia, and nursing disciplines must closely align to ensure safe service delivery.  
8 Deficiencies in any of these areas compromise care at all levels of the health system.  
9  
10  
11  
12  
13  
14  
15  
16

### 17 **3.2 Goals of the Optimal Resources for Children's Project**

18  
19 The goal of the Optimal Resources for Children's Surgery (OReCS) program is to identify and  
20 promote standards of care that will improve the surgical care of children worldwide. The OReCS  
21 Program seeks to accomplish this by more clearly defining what essential services should be made  
22 available to every child in every region of the world. The OReCS recommendations are also  
23 intended to expand upon the World Health Organization's recently developed standards for  
24 improving the quality of care for children and young adolescents in health facilities.<sup>9</sup>  
25  
26  
27  
28  
29  
30  
31

32 In multiple international GICS meetings, low-resource providers identified 1) training and  
33 staffing 2) physical resources 3) quality and safety, and 4) research as thematic areas that need to  
34 be addressed to improve children's surgery globally. Importantly, these thematic areas represent  
35 subsets of the WHO health systems building blocks (Workforce, Service Delivery, Medical Products,  
36 Vaccines & Technologies, Financing, Leadership & Governance, and Information Systems)<sup>10,11</sup>. The  
37 Optimal Resource Project addresses specific aspects of health systems that low resource providers  
38 believe must be addressed first.  
39  
40  
41  
42  
43  
44  
45  
46

47 The Optimal Resource Project also seeks to describe necessary resources by hospital level  
48 within a national health system. The resources for different levels of children's surgical care are  
49 outlined in the form of templates, which are designed to support basic, intermediate and  
50 complex/advanced surgical care for children in low resource settings. The goal of these templates is  
51 to provide a guide for minimum standards for those planning surgical services for children in low-  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 resource settings. Thus, OReCS is intended to provide pragmatic, yet comprehensive  
4 recommendations regarding the resources, training and research priorities required for safe and  
5 high-quality surgical services for children. These resources can be adapted to suit local contexts and  
6 settings.  
7  
8  
9  
10

11 These guidelines are intended for:

- 12 • Health care professionals: in delivery and providing children's surgical services, and  
13 development of general and subspecialty services
- 14 • Hospital managers and administrators: in organization and strengthening of children's  
15 surgical services
- 16 • Policy makers: to provide the necessary strategic planning support for children's surgery
- 17 • Health planners: to ensure specific services (i.e., intensive care, blood banking, transport  
18 and referral systems, availability of essential medicine) are delivered at appropriate levels  
19 of the health system
- 20 • Training institutions and colleges: in planning, education and scale up of training of human  
21 resources to support children's surgery
- 22 • Individuals and organizations: in advocating for children's surgery
- 23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38

### 39 **3.3 Justification**

40  
41 Approximately five billion people lack access to safe surgical care worldwide, with the  
42 majority living in low- and middle-income countries (LMICs).<sup>3,12</sup>. Given that children comprise 40%  
43 - 50% of LMICs populations, this means that worldwide in 2016 up to 2.0 billion children do not  
44 have access to safe surgical care. To attain universal health coverage (UHC) and achieve the specific  
45 goals and targets of the sustainable development goals (SDGs), access to safe and affordable  
46 surgical and anesthesia care must be provided *for all children*, regardless of geographic location or  
47 socioeconomic status. It is therefore crucial that children's surgery becomes a key component of  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 every country's National Surgical, Obstetric and Anesthesia Plans (NSOAPs) and National Maternal  
4 and Child Health Plans. Fully integrating children's surgery into health systems strategic planning  
5  
6 both capitalizes on the multiple specialty-specific collaborations necessary in children's surgery  
7  
8 and avoids vertical plans that may lead to fragmented health systems. This is particularly  
9  
10 important in poorest areas of the world where the surgical needs of children are the greatest.  
11  
12

13  
14 Published literature and previous work have emphasized that outcomes for children with  
15  
16 surgical problems are optimized when treatment is provided by those specifically trained to  
17  
18 provide care for this segment of the population<sup>13-15</sup>. Based on these findings, efforts have been made  
19  
20 to identify optimal resources for children's surgical care in high-income countries<sup>16-18</sup>, but these do  
21  
22 not address low-resource settings, where the needs of children often may be neglected.  
23

24  
25 Children develop different surgical diseases compared to adults, have unique physiologic  
26  
27 needs and require different operations. For example, they have specific anaesthetic risks,  
28  
29 difficulties maintaining their body temperature, metabolic differences compared to adults, risk for  
30  
31 medication errors, unique transfusion and resuscitation protocols, and challenges in vascular  
32  
33 access. Their care before, during and after operations also differs. Good outcomes depend on  
34  
35 having the right resources and recognizing that "children are not just small adults". For this reason,  
36  
37 children should be treated in child-friendly environments with appropriate resources and properly  
38  
39 trained professionals to adequately address and meet their unique needs.  
40  
41

## 42 43 **4. SURGICAL NEEDS OF CHILDREN**

---

### 44 45 46 **4.1 Epidemiology of childhood surgical conditions**

47  
48  
49 Surgical needs cut across the entire spectrum of children's health problems, with the most  
50  
51 common conditions being injuries, congenital anomalies, and infections requiring surgery.<sup>19,20</sup>  
52  
53 Further, surgical conditions are exceedingly common, with an estimated 85% of children requiring  
54  
55 surgical care at some time before reaching age 15 years.<sup>21</sup> Twenty per cent of children in a recent  
56  
57  
58  
59  
60

1  
2  
3 multi-country household survey had a surgical need, and 62% of these children had an unmet need,  
4  
5 corresponding to 3.7 million children in those countries alone.<sup>22</sup>  
6

7 While global efforts to improve neonatal survival have expanded, these programs have not  
8  
9 included neonatal surgery, and the mandate to ensure treatment of surgical congenital anomalies.<sup>23</sup>  
10  
11 Road traffic injuries and associated traumatic brain injuries also exact an extraordinary toll on  
12  
13 children, with more children in the 5-14 age group dying of injuries than HIV, tuberculosis, and  
14  
15 malaria combined.<sup>24</sup> Globally, 90% of burn injuries occur in low-income countries and children are  
16  
17 disproportionately affected.<sup>25</sup> Meanwhile, as global attention has shifted to non-communicable  
18  
19 diseases, cancer control efforts are growing, but generally have not emphasized resources required  
20  
21 to treat tumors in low- and middle-income countries.<sup>26</sup> For many childhood cancers in these  
22  
23 settings, surgery is the primary curative modality required to ensure survival, as chemotherapy and  
24  
25 radiation may not be available.  
26  
27  
28  
29

#### 30 **4.2 Benefits of providing children's surgical care**

31  
32 The provision of surgical care to children within a health system can achieve the following goals:  
33

- 34 • Identify and treat correctable congenital anomalies.
- 35
- 36 • Treat life-threatening injuries promptly to maximize the likelihood of survival.
- 37
- 38 • Ensure potentially disabling injuries are treated appropriately, in order to minimize  
39 functional impairment and to maximize return to independence and participation in  
40 community life.
- 41
- 42 • Minimize acute and long-term pain and suffering.
- 43
- 44 • Protect from substantial economic losses to entire families due to medical expenses, death,  
45 and disability.
- 46
- 47 • Increase economic productivity in families, communities, and countries
- 48
- 49 • Increased physical activities helping to prevent non-communicable diseases  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- Increase school attendance
- Improve equity in surgical care provision across all age groups, as children are often neglected

## 5. STRUCTURE OF HEALTH SYSTEMS IN LOW AND MIDDLE-INCOME COUNTRIES

---

### 5.1. Overview

Health care in most LMICs is delivered through a network of health care facilities; which vary by the type and complexity of care provided. The most common organizational scheme in LMICs is a national referral hospital supported by provincial general hospitals, district hospitals, health centers, and dispensaries. This network of facilities is often supplemented by privately owned and operated hospitals and clinics and care provided by faith-based organizations. It is important to appreciate that facility terminology may vary considerably worldwide, and that capabilities, and “platforms for service delivery” at each level vary significantly among different countries. It is also acknowledged that the division between different levels is somewhat arbitrary as many times patients seek care for less complicated surgical problems at higher-level facilities. Nevertheless, classification of health facilities is still extremely valuable for allocating resources to the different levels of the health system.

### 5.2 Types of facilities

The OReCS Program builds on the health care facility classification system used in the Disease Control Priorities, 3<sup>rd</sup> Edition (DCP3).<sup>27</sup> DCP3 classifies platforms of delivery into five categories: **1) population based, 2) community level, 3) health center, 4) first-level hospital, and 5) referral hospital.** The OReCS program focuses on the delivery of surgical care at the highest three levels, i.e., levels 3, 4 and 5. In addition, based on recommendations of children’s

1  
2  
3 surgery providers in LMICs, the OReCS Program further splits the referral hospital levels into  
4  
5 second, third level facilities and national children's hospitals. This distinction was important to  
6  
7 children's surgery providers in LMICs because of the substantially different complexity of care that  
8  
9 is typically provided at second and third-level hospitals. The children's hospital category was  
10  
11 added because of the tremendous impact this type of facility can have on training, research and  
12  
13 advocacy, both at the national and international level. It is important to appreciate nomenclature of  
14  
15 hospital levels can vary by country, or region of the world. As an important point of slightly  
16  
17 different nomenclature, the WHO descriptions of hospital levels include 3 levels, including primary  
18  
19 health care (first-level); second-level (generally the district hospital) and third-level. Table 1  
20  
21 reviews the classification system, as designation and nomenclature of hospital levels have  
22  
23 sometimes varied. As the DCP3 and WHO health care facility classification can be confusing, the  
24  
25 OReCS document includes both nomenclatures in most tables.  
26  
27  
28

### 29 **5.2.1 Health center**

30  
31 A health center is typically staffed by non-doctor providers, such as community health  
32  
33 workers, nurses, or medical assistants. It is the mainstay of health care throughout many of the  
34  
35 rural areas of low-income countries. Health centers serve an important role in the health care  
36  
37 system, as they represent the point of first contact for a child with any medical need, including  
38  
39 those with surgical problems. Programs exist at this level for recognition and treatment of life-  
40  
41 threatening problems at birth and in the neonatal period, but surgical problems have not been  
42  
43 targeted.  
44  
45

### 46 **5.2.2 First-level hospital**

47  
48 This level facility forms an integral part of the health system, as up to 90% of the global  
49  
50 population receives their emergency care at this level. How these facilities are staffed, and  
51  
52 importantly, the distance and ability to transfer to the next level of care can be highly variable.  
53  
54 First-level hospitals should have twenty-four-hour clinical services that can provide emergency  
55  
56  
57  
58  
59  
60

1  
2  
3 treatment for patients with life-threatening injuries, obstetrical emergencies and other child health,  
4 medical and surgical problems that require immediate intervention. Providers at these hospitals  
5 should be comfortable with basic resuscitation, initial care and management of injuries, burns, and  
6 infections, and be comfortable screening for congenital anomalies. Surgical providers should be  
7 familiar with basic pediatric surgical procedures.  
8  
9  
10  
11  
12

13  
14 As such, first-level hospitals should have a functional operating room with a provider  
15 capable of administering a safe general anesthetic as well as availability of oxygen, sterile  
16 instruments, anesthetic agents and essential medicines.<sup>8</sup> This is typically the first level of the  
17 health care system where a general anesthetic can be provided. A recovery area should be present  
18 and nurses should have perioperative care skills. This level of facility should also have the  
19 laboratory and diagnostic capabilities that are appropriate for the medical, surgical and outpatient  
20 care provided. In some areas, particularly in East Africa, non-physician surgeons have been trained  
21 to act in the capacity of general surgeons, performing Caesarean section and other general surgical  
22 procedures.  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

### 33 **5.2.3 Referral hospitals**

34  
35 Referral hospitals serve the important role of supporting the health care provided at lower  
36 levels of care, both in terms of provided care for more complicated patients and for training and  
37 education. The OReCS Program includes recommendation for three types of referral hospitals: 1)  
38 second-level hospital, 3) third-level hospital, and 4) national children's hospital.  
39  
40  
41  
42  
43

#### 44 **5.2.3.1 Second-level hospital**

45  
46 Second-level hospitals typically have a least one fully trained general surgeon and an  
47 anesthesiologist. The staff at such facilities may also include other specialists such as orthopedic  
48 surgeons. More advanced and specialized nurses may be present, along with some allied health  
49 personnel supporting surgical services (e.g. physical and occupational therapists). These facilities  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 are usually referred to as regional hospitals in parts of Africa, community health centers or district  
4 hospitals in India, or general hospitals in Latin America and parts of Africa.  
5  
6

### 7 **5.2.3.2 Third-level hospital**

9  
10 Third-level hospitals have the broadest range of subspecialties and represent the highest level  
11 of care within a country or geographic region. The capabilities of third-level hospitals vary widely  
12 worldwide. Some third-level hospitals have an extensive range of subspecialties, while others are  
13 more limited. Third-level hospitals serve the important additional function of being teaching  
14 hospitals. Stand-alone facilities or niche hospitals that focus on a specific specialty, or surgical  
15 conditions (e.g. orthopedic, craniofacial, neurosurgery) also belong to this category of facilities.  
16  
17 Human resources and skills should include complex care with a wide range of nursing and allied  
18 health personnel (See Appendix 3).  
19  
20  
21  
22  
23  
24  
25  
26

### 27 **5.2.3.3 National children's hospital**

28  
29 Children's hospitals focus exclusively on the health needs of children and adolescents. Such  
30 facilities may be part of a second or third level or may be an independent facility. Children's  
31 hospitals serve the important role of training, education, and research, and are typically better  
32 resourced (spectrum and numbers of available specialists, specialized support services, specialized  
33 equipment) for the care of children compared to the regular hospitals at those levels. Ideally, each  
34 country should have at least one designated national children's hospital.  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Table 1.** Definition and examples of different types of health care facilities in low- and middle-income countries. Type of facility is based on classification system used in DCP3 (World Bank) and the World Health Organization.

Facility classification		Description	Examples
DCP3 (2017)	WHO (2003)		
Health center	Level 1: Small hospital/health center	Health care facility usually located within a rural community  Provides basic health care services of very general nature.  Proper triage is a key expectation	Community health center Primary health center Comprehensive health center
First-level hospital	Level 2: District/provincial hospital	Services available are of a general nature and not specialized  First level at which general anesthesia is available  Proper triage is a key expectation.	General hospital District hospital Cottage hospital
Referral hospital	Second-level hospital	Level 2: District/provincial hospital  Clinical services are specialized by function, but not all specialties and limited specialized technical equipment are available	Regional hospital Provincial hospital General mission hospital
	Third-level hospital	Level 3: Referral hospital  Highly specialized hospital, with similarly specialized staff and technical equipment to support level of care provided  Clinical services are highly differentiated by function  Training is provided	Academic/teaching/university hospital National hospital Central hospital Niche / Specialized mission hospital
	National children's hospital	Level 3: Referral hospital  Highly specialized hospital, dedicated to the care of children  Highly specialized staff and technical equipment in all children's specialties are available  Complex multidisciplinary and chronic care Specialized teaching and research in all children's specialties provided	Children's hospital

## 6. LEVELS OF CHILDREN'S SURGICAL CARE

---

### 6.1 Overview

It is neither feasible, nor desirable to provide all types of surgical care at every level of the health care system. As such, it is important to decide which childhood surgical problems can be safely treated at different levels of the health care system. This approach is feasible because children's surgical conditions have variable complexity, with some childhood surgical conditions treatable with less skill and resources, while others require advanced training and greater resources. The majority of childhood surgical conditions are relatively simple to treat, making prevention of death and disability very cost-effective.

### 6.2 Levels of surgical care

The OReCS program defines three levels of surgical care for children: basic, intermediate and complex/advanced care (Table 2). These categories are based on the complexity of the surgical problem and the resources required to treat the problem.

#### 6.2.1 Basic surgical care

Many of the most common surgical problems of children are not difficult to treat and require only minimal resources. Examples of basic surgical care include repair of an uncomplicated laceration or splinting of a fracture. Nonetheless, basic surgical care does require a basic knowledge of childhood surgical conditions, such that simple problems can be treated and more complicated problems are recognized and referred to higher levels of care after appropriate stabilization and resuscitation. By definition basic surgical care are problems that do not require general anesthesia.

### 6.2.2 Intermediate surgical care

This level of care addresses the most common emergency and essential surgical conditions of childhood. Examples include uncomplicated appendicitis, management of the majority of childhood injuries and treatment of common surgical infections (e.g., empyema, uncomplicated osteomyelitis). More complicated surgical conditions, and those with comorbidities are referred to higher levels of care. Surgical problems that fall into this category may or may not require a general anesthetic. Perioperative nursing skills are required at this level.

### 6.2.3 Complex/advanced surgical care

Complex/advanced surgical care requires the greatest knowledge of childhood surgical conditions. Examples are neonatal surgery, repair of cleft lips and palates and pediatric neurosurgical and cardiac procedures. Multimodal pediatric oncology care including coordinated care with pediatric oncologists through a cancer care team would be found at this level. Treatment often involves highly specialized care and is often multidisciplinary including more specialized nursing skills and a wide range of allied health professionals.

## 6.3. Suggested levels of care by facility type

Countries vary markedly in their ability to pay for health care. Recognizing this, individual countries will need to make decisions regarding, where and how surgical resources will be allocated. The resources outlined are standards rather than estimates of current capacity, which may be below standards in some low-income countries. To facilitate this planning process the OReCS program has included the following criteria to help identify which level of care is most appropriate for the type of facility (Table 3). In a fashion similar to that used in *Guidelines for essential trauma care*, the priority for having each level of care was given a designation according to the following criteria:<sup>28</sup>

**“Essential” (E) care**

An “E” designation means that the stated level of care should be available in all cases. The E designation represents the minimal acceptable level of children’s surgical care common to all regions, including even those where access to resources is most severely restricted. These services could and should be provided to children with surgical problems at the health facility concerned, even in countries with the most limited financial resources (whose Ministries of Health have a total budget of only \$3–4 per capita per year). It is important to recognize that these essential items could be provided primarily through improvements in organization and planning, with a minimal increase in expenditures.

**“Desirable” (D) care**

The designated item represents a capability that increases the probability of a successful outcome for children with surgical conditions. Such items may not be possible to provide for all facilities in environments with the poorest access to resources. Hence, they are not listed as essential. However, for countries with greater resource availability, such items may ultimately be designated essential in their own national plans.

Finally, the OReCS program anticipates that in the process of developing national surgical plans, many countries, especially with greater resources, will appropriately decide to declare some of the items in the desirable category essential, or provide higher levels of care at lower levels of the health system. The converse is not true, items designated as essential should remain so designated, as a change may result in children’s surgical care below recommended standards.

**Table 2.** Levels of care based on the complexity of the surgical conditions

Level of care	Definition
I: Basic	Recognition and treatment of minor surgical conditions that do not require a general anesthetic. Referral of more complex surgical conditions and patients with important comorbidities to higher levels of care.
II: Intermediate	Recognition and treatment the most common emergency and essential childhood surgical conditions that may or may not require a general anesthetic. Referral of more complicated childhood surgical conditions and patients with important comorbidities to a higher level of care.
II: Complex/Advanced	Treatment of complex children's surgical conditions. Multidisciplinary and highly specialized care.

**Table 3.** Recommended level of surgical care for children by facility type/platform.

Type of facility (DCP3classification) <sup>a</sup>		Level of care		
		Basic	Intermediate	Complex/Advanced
Health center		E	-	-
First -level hospital		E	E	-
Referral hospital	Second level hospital	E	E	D*
	Third level hospital	E	E	D
	National Children's hospital	E	E	E

E = Essential; D = Desirable

D\*: depends of available resources and geographic location of facilities

<sup>a</sup> See Table 1 for equivalent health care facilities in the WHO classification system.

## 7. OPTIMAL RESOURCES FOR CHILDREN'S SURGICAL CARE BY LEVEL OF CARE

---

### 7.1 Overview

In most low resource settings, little consideration has been given to optimizing the resources available for treatment of children with surgical conditions. Many hospitals lack important equipment, some of which is inexpensive. GICS members have observed that the main reason for the absence of such vital equipment and supplies is usually a lack of organization and planning, rather than resource restriction. Programs to assure the supply and maintenance of children's surgical equipment, appropriate to specific country circumstances, could address deficiencies with relatively inexpensive but high-yield resources. Nevertheless, in 2018, these goals should be achievable for most children with surgical needs regardless of geographic location.

### 7.2 Resource templates

The OReCS program provides resource templates for each of three suggested levels of care (basic, intermediate, complex/advanced). These resource templates offer broad guidelines to how surgical care of children might be organized in a health system. Each template provides an overview of the scope of practice and provides specific resources for optimal children's surgical care. Scope of practice contains information on the function, age and comorbidities of patients, anesthesia required, and procedure type. The templates are organized into three sections: 1) training and staffing, 2) physical resources, and 3) quality and safety. Training implies that medical, nursing and other staff have the requisite skills to perform diagnostic and therapeutic activities safely and successfully. Proper training includes basic education (school and postgraduate training), and a process for continuing education to maintain their skills. Training issues are described in greater detail in Section 7.1.

1  
2  
3 Physical resources are defined as infrastructure, equipment and supplies. To be effective for  
4 a population these items must be available to all who need them without consideration of ability to  
5 pay. Equipment and supplies need to be readily available on an ongoing basis; where appropriate,  
6 24 hours a day, seven days a week. It also implies that supply chains within the health system are  
7 functional, and that organizational and administrative mechanisms are in place to replace needed  
8 items or to accomplish equipment repairs when required.  
9  
10  
11  
12  
13  
14  
15

16 The resource templates provide a list of childhood procedures that each respective level of  
17 the health system should be capable of performing safely. Our procedure recommendations align  
18 with and are an extension of the recommendations of the DCP3 essential surgery guidelines of 44  
19 procedures that are both cost effective and essential at each hospital level<sup>29</sup>. Some of the  
20 procedural recommendations by the DCP3 that are relevant to children and parallel our  
21 recommendations include incision and drainage at the health center level, appendectomy, trauma  
22 laparotomy, external fixation and emergency burr hole at the first level hospital, and operative  
23 repair of club foot, cleft lip/palate and anorectal malformations at the second and third level  
24 hospitals. For ease of reference, Appendix 1 includes a list of childhood procedures by specialty and  
25 hospital level. Appendix 2 provides recommended procedures by thematic areas: trauma/injury,  
26 congenital anomalies, infections, tumors and other.  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39

### 40 **7.2.1 Optimal resources for basic surgical care (health center)**

41  
42 The suggested optimal resources for basic surgical care are summarized in Table 4. The  
43 optimal resources for this level of care, which would be available at the health center, focus on  
44 recognition and triage of childhood surgical conditions. Treatment is limited to uncomplicated  
45 conditions that do not require a general anesthetic. The key resources required for this level of care  
46 are training, wound care supplies and education including community injury prevention strategies.  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

### 7.2.2 Optimal resources for intermediate surgical care (first-level hospital)

The suggested optimal resources for intermediate surgical care are summarized in Table 5. The optimal resources for this level of care should be available at the first level hospital and would focus on supporting emergency and essential surgical care of children. The particular resources required for this level of care are training and supplies needed to support the care, as well as monitoring and evaluation of care. This level of care assumes that a first-level hospital has a functional operating room with support services. Anesthesia and perioperative care providers including nursing is critical at this level. As previously noted, all surgical procedures in children < 1 year old should be referred to higher level of care except for emergent cases where no referral is possible. In the scenario that a child < 1 year old must undergo general anesthesia at this level, the most senior anesthesia provider should care for the child.

### 7.2.3 Optimal resources for advanced/complex surgical care (referral hospital)

The suggested optimal resources for advanced/complex surgical care are summarized in Tables 6. For this level of care, the key resources are training and the supplies needed to support a broad spectrum of specialized children's surgical care that is often multidisciplinary. Surgical, anesthesia, and nursing providers must have more advanced training, and a broader range of allied health personnel must be present to support specialized services. It must be emphasized that the quantity and type of advanced/complex surgical care delivered at different types of referral hospitals can vary considerably between countries, and depends on resources available as well as geographic location. Thus, individual countries must decide how best to distribute advanced/complex surgical care. To facilitate this decision-making the OReCS program has outlined some of the general differences that should distinguish the type of surgical care for child provided at second and third level facilities, and national children's hospitals (Table 8).

#### 7.2.4 Optimal resources for children's anesthesia care

In contrast to surgical subspecialties where procedures can be specifically defined by levels of care, anesthesia and peri-operative care must take into account the child's age and risk stratification as determined by the child's co-morbidities. This principle led to the following summary anesthesia recommendations which should be used in addition to the specific surgical recommendations by level of care. Of note, these recommendations include a specific recommendation for the first-level hospital in which all children < 1 year old that require general anesthesia should be transferred to a higher level of care. The only exception is infants requiring emergent surgical procedures in which transfer is not possible. This recommendation comes directly from observational data on adverse outcomes in infants. Children less than 1 month undergoing general anesthesia have an approximately 6 times higher risk of cardiac arrest than do infants between 1 month and 1 year of age and children between 1 month and 1 year have an approximately 4 times higher risk of anesthesia-related cardiac arrest than in children aged 1 to 18 years old<sup>30-32</sup>.

- At any level of care where children's surgery is performed under general anesthesia or regional anesthesia, there should be a trained anesthesiologist or non-physician anesthesia provider available to provide care during the procedure (exception is for minor surgery under local anesthesia)
- GICS and the WFSA view anesthesiology as a medical practice. Wherever and whenever possible, it is highly recommended that anesthesia delivered to a child should be provided, led or overseen by an anesthesiologist. When anesthesia is provided by non-anesthesiologist, these providers should be directed and supervised by anesthesiologists in accordance with their level of training. When no anesthesiologists are present at a specific hospital level, leadership should be provided by the most senior anesthesia provider available

- 1
- 2
- 3 • Alternatives to general anesthesia should be developed and explored (local and regional
- 4 anesthesia) at all levels of care whenever applicable
- 5
- 6
- 7
- 8 • Size specific anesthesia monitors and equipment should be available at all care levels and
- 9 facilities in which general anesthesia is provided to allow for children to be adequately
- 10 monitored
- 11
- 12
- 13
- 14 • Programs for post-operative pain management should be developed in at all levels where
- 15 children's surgery occurs
- 16
- 17
- 18 • Unit sufficient resources are available to ensure that physician anesthesiologists are
- 19 physically present at every level where children's surgery occurs, pediatric surgical
- 20 networks for telemedicine, telephone consultation or other communication should be
- 21 developed and made standard procedure that take into consideration specific anesthesia
- 22 perspectives in regard to caring for children
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60

**TABLE 4.** Optimal resources for basic surgical care. This type of care is designed to be delivered at the Health Center Level<sup>a</sup>.

<b>BASIC SURGICAL CARE</b>		
<b>SCOPE OF PRACTICE</b>		
<b>Function</b>	Screening for surgical disease Resuscitation and Triage Training of health care workers Referral to higher levels of care Community Health Education	
<b>Age of patients</b>	All ages	
<b>Anesthesia</b>	Local anesthesia for minor procedures.	
<b>Examples care provided (See Appendix 1 and 2 for a more detailed list)</b>	<b>Injuries</b>	Resuscitation with basic life support measures, suturing and dressing of simple wounds, splinting of closed, non-displaced fractures 1 <sup>st</sup> degree burns < 10% TBSA and not including face, hands, GCS Score
	<b>Congenital anomalies</b>	Screening for congenital anomalies (e.g., abdominal wall defects, hernias, anorectal and urogenital abnormalities, limb deformities, hip dysplasia, cardiac anomalies, neural tube defects and craniofacial anomalies (cleft lip/palate)
	<b>Infections</b>	Screening for surgical site infections, intra-abdominal infection, bone and joint infections. Treatment of superficial abscess with incision and drainage
	<b>Tumors</b>	Screening for tumors
	<b>Others</b>	Circumcision using a Plastibel Removal of visible foreign bodies in the ear and nose Hearing screening Vision screening
<b>OPTIMAL RESOURCES</b>		
<b>Training &amp; staffing</b>	<b>Human resources</b>	Community health workers, general practice nurses, and others
	<b>Required skills</b>	<u>Screening</u> : congenital anomalies, surgical diseases, surgical site infections <u>Resuscitation and stabilization</u> : basic life support measures including airway management, peripheral vascular access, insertion of nasogastric tube, initial burn management <u>Care of trauma wounds</u> : arrest of bleeding, cleaning, suturing, dressing, splinting <u>Care of postoperative wounds</u> : cleaning, dressing, removal of stitches <u>Pain management</u> : use of non-opioid analgesics
<b>Physical resources</b>	<b>Infrastructure</b>	Infrastructure to support basic services Availability of transportation to higher level of care
	<b>Equipment &amp; supplies</b>	See OReCS: Supplies, Equipment and Infrastructure Document
<b>Quality &amp; Safety</b>	Supervision and mentoring Data collection for Quality Control CME/CPD	

<sup>a</sup> Equivalent to small hospital/health center in the WHO health facility classification system (WHO Level 1).

**TABLE 5.** Optimal resources for intermediate surgical care. This type of care is designed to be delivered at First-Level Hospitals<sup>a</sup>.

INTERMEDIATE SURGICAL CARE		
SCOPE OF PRACTICE		
<b>Function</b>	24/7 emergency surgical care Diagnosis and treatment of the most common pediatric surgical conditions Training of health care workers Referral to next level of care for complex procedures	
<b>Age of patients</b>	All ages <sup>b</sup>	
<b>Anesthesia</b>	General anesthesia (ASA I+II) Referral ASA >2	
<b>Examples of care provided (see Appendix 1 and 2 for a more detailed list)</b>	<b>Injuries</b>	Resuscitation with advanced life support measures including fluid management and nutrition Trauma laparotomy Closed and open fractures Diagnosis and stabilization of neurological trauma (e.g., epidural hematoma, includes emergency burr hole if transfer not possible) Treatment of burns (<10% not involving face, hands and perineum)
	<b>Congenital anomalies</b>	Inguinal hernia repair in older children
	<b>Infections</b>	Incision and drainage of abscesses, pyomyositis and septic arthritis Appendicitis, Intestinal Perforation Thoracostomy tube for empyema Drainage and debridement of osteomyelitis
	<b>Tumors</b>	Excision of benign tumors
	<b>Others</b>	Intestinal obstruction (e.g., intussusception, enterolysis for adhesions) Foreign body removal from ear, nose, airway and esophagus
OPTIMAL RESOURCES		
<b>Training &amp; staffing</b>	<b>Human resources</b>	General doctor or non-physician surgical provider Anesthesiologist +/- anesthesia provider Perioperative and ward nursing, Physiotherapist
	<b>Required skills</b>	Basic knowledge of common childhood conditions Advanced Trauma Life Support (ATLS); Pediatric Advanced Life Support (PALS); SAFE course Antibiotic stewardship Pediatric pain management Pediatric Anesthesia experience
<b>Physical resources</b>	<b>Infrastructure</b>	Operating room, recovery area, and children's ward Availability of transportation to higher level of care
	<b>Equipment &amp; supplies</b>	See OReCS: Supplies, Equipment and Infrastructure Document
<b>Quality &amp; Safety</b>	Supervision and mentoring Data collection for Quality Control CME/CPD	

<sup>a</sup> Equivalent to a district or provincial hospital in the WHO health facility classification system (WHO Level 2).

<sup>b</sup> All babies < 1 year old should be referred to higher levels of care due to increased anesthetic risk. Only exception is life-threatening emergency when transportation is not possible.

**TABLE 6.** Optimal resources for complex/advanced surgical care. This type of care is designed to be delivered at the referral hospital level. Referral hospitals include Second- and Third levels facilities, and National Children's Hospitals<sup>a</sup>.

COMPLEX/ADVANCED SURGICAL		
SCOPE OF PRACTICE		
<b>Function</b>	Advanced diagnostic services and multidisciplinary specialized care Receive referrals from other hospital levels Training of health care workers Research and Advocacy Develop standards of care	
<b>Age of patients</b>	All ages	
<b>Anesthesia</b>	All types of anesthesia including neonatal, cardiac and neuro anesthesia	
<b>Examples of care provided (See Appendix 1 and 2 for a more detailed list)</b>	<b>Injuries</b>	All traumatic injuries referred from lower levels of care, including neurovascular injuries
	<b>Congenital anomalies</b>	All congenital anomalies referred from lower levels of care, including repair of anorectal malformations, urogenital anomalies, congenital heart disease and meningomyelocele, hydrocephalus
	<b>Infections</b>	All surgical infections referred from lower levels of care, including complex soft tissue infections, osteomyelitis, intracranial infections
	<b>Tumors</b>	All benign and malignant tumors, including abdominal and urological malignancies
	<b>Others</b>	
OPTIMAL REASOURCES		
<b>Training &amp; staffing</b>	<b>Human resources</b>	<u>Trainees:</u> Medical Students, Surgical Residents and Surgical Subspecialty Fellows <u>Physician Human Resources:</u> <u>Pediatric Surgical Specialists:</u> general pediatric surgery; cardiac surgery; ENT surgeon; neurosurgery, ophthalmology; oral and maxillofacial surgery; orthodontics; orthopedic surgery; plastic surgery; urology <u>General pediatricians and pediatric specialists:</u> Anesthesia, Critical Care, Cardiology, Oncology, Gastroenterology, Neurology, Nephrology, Radiology, and others <u>Non-Physician Human Resources:</u> All Support specialists and staff in relevant specialties including perfusionists, Speech pathologist, occupational/physical therapists
	<b>Required skills</b>	Appropriate skills and training necessary to provide such specialized pediatric surgical care including neonatal critical care Structured Education for Trainees (all levels) Management and administrative skills
<b>Physical resources</b>	<b>Infrastructure</b>	Dedicated space for pre-operative and post-operative evaluation Dedicated Children's inpatient ward, ICU and NICU Endoscopy capability- EGD, colonoscopy, bronchoscopy Complete radiology capability (CT/MRI/Ultrasound/Fluoroscopy) Cardiac catheterization lab, cardiopulmonary bypass
	<b>Equipment &amp; supplies</b>	See OReCS: Supplies, Equipment and Infrastructure Document
<b>Quality &amp; Safety</b>	Supervision and mentoring; Data collection for Quality Control CME/CPD; Morbidity and Mortality Conference Trauma Conference; Tumor Board	

<sup>a</sup> Equivalent to a district/provincial hospital or referral hospital in the WHO health facility classification system (WHO Level 2 or 3)

**TABLE 7.** Suggested distribution of advanced/complex care at different levels of referral hospitals. Higher levels of care assume the infrastructure/services/functions of lower levels of care are present.

Type of referral hospital	Responsibilities	General anesthesia	Human resources	Required skills	Infrastructure
<b>Second- level</b>	24/7 Emergency surgical care, Selected more complex cases  Health care worker training  Appropriate referral	Yes, including some complex cases with comorbidities (Limit of ASA III)	Specialists in some areas of children's surgical care provided	Advanced surgical, anesthesia, and nursing skills in the majority of children's surgical care	Children's wards, clinics, operating rooms, and recovery areas
<b>Third-level</b>	Comprehensive surgical care for children <sup>2</sup>	Yes, including complex cases with comorbidities (All ASA)	Specialists in all areas of children's surgical care provided Pediatric anesthesiologist	Advanced surgical, anesthesia, and nursing skills in all areas of children's surgical care	NICU, PICU, burn unit
<b>National Children's hospital</b>	Multidisciplinary and chronic care  Training, education and research in all children's surgical specialties  Development of standards of care Advocacy	As above	As above	As above  Training in multidisciplinary and chronic care	As above

## 8. STRATEGIES FOR IMPROVING CHILDREN'S SURGICAL CARE

---

### 8.1 Training

There is a critical need to improve the knowledge and skills of those who care for children with surgical conditions at all levels of the healthcare system, from the community health worker to the general doctor and pediatrician, and the surgical, anesthesia, and perioperative care specialists at the tertiary referral hospital. A variety of other professionals are also critical to the treatment of children with surgical conditions including, but not limited to nursing, laboratory and X-ray technicians, operating room personnel, and rehabilitation specialists. A table of recommended human resources at each health care level is provided in Appendix 3.

The scale of the global surgery workforce problem is perhaps best illustrated by estimation of the Lancet Commission on Global Surgery (LCoGS) that there will be a deficit of 2.3 million surgical providers by the year 2030. Krishnaswami *et al.* have recently reviewed pediatric surgical workforce needs with similarly dire conclusions.<sup>33</sup> Anesthesia providers and perioperative nursing staff are also severely limited.<sup>34</sup> Lack of funding, limited training opportunities and deficient political support, in combination with a critical lack of mentors, were identified as important limiting factors.

Greater attention is needed worldwide to define and optimize the training of doctors, non-physician providers, nurses, and allied health professionals in the surgical care of children. This includes developing means of improving knowledge and skills imparted during undergraduate medical education, as well as the knowledge and skills acquired during postgraduate training. There is a need to integrate the surgical care of children into basic medical school curriculum and house-officer training, as well as in specialist training, for example, in anesthesia, orthopedics, neurosurgery, among other disciplines. This expanded education would assure that all graduates assigned to work in rural hospitals would have the skills needed to assess and treat children with

1  
2  
3 the most common surgical conditions. Existing educational resources may be used and new  
4 integrated modules may be developed.<sup>35-41</sup> Standardization of curricula that focus on improving  
5 children's surgical care skills could be an important tool for raising the standard of such care  
6 worldwide. There is also a clear need to increase the number of fellowships that focus on children's  
7 surgical care after completion of formal surgical residency training.  
8  
9  
10  
11  
12

13  
14 Training should reflect the surgical needs of the pediatric population served. Rather than a  
15 vertical approach that focuses on specific diseases, a holistic approach based on packages of  
16 conditions is recommended. There is a need to define a specific set of diagnostic and therapeutic  
17 skills to care for children with surgical conditions at every level of the health care system (Table 8).  
18 Such training could be provided to all levels of the health system by professional associations in  
19 countries where this type of organization is possible. Alternatively, training packages could be  
20 delivered regionally when not possible in individual countries.  
21  
22  
23  
24  
25  
26  
27  
28  
29

30 Finally, there is need for continuing medical education (CME) for all practitioners caring for  
31 children with surgical conditions. CME is critical to prevent decay of knowledge and skills,  
32 especially among those who do not see large numbers of children with surgical problems daily. It  
33 also helps staff to maintain competence and learn about new and developing areas, technology, and  
34 innovations in their respective fields. Teamwork, staff and inter- and intra-departmental  
35 relationships are enhanced and thereby overall competencies and performance improves as well.  
36 In addition, CME courses offer the opportunity to better define the core essential elements of  
37 children's surgery for any environment. By so doing, they can influence the knowledge and skills  
38 imparted in undergraduate and postgraduate training. Adequate and appropriate resources should,  
39 therefore be committed to CME in a planned and ongoing manner within and between hospitals and  
40 levels of care.  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**TABLE 8.** Training objectives for different types of facilities

Type of facility (DCP3 classification) <sup>a</sup>	Level of Care <sup>b</sup>	Training Objectives
Health center	<b>I</b>	<p>Train birth attendants and community health workers to recognize the most common (often visible) congenital anomalies and provide basic counseling. This has been done successfully for conditions such as clubfoot, cleft lip, and cleft palate. Training could be expanded so that birth attendants screen for conditions such as hydrocephalus, spina bifida, hernia, anorectal malformation, hip dysplasia and undescended testes.</p> <p>Create community health worker manual focused on identifying and addressing surgical conditions and injury prevention</p>
First-level hospital	<b>I-II</b>	<p>Emphasis on training the workforce to identify and treat an essential package of children's surgical conditions. This would include some emergencies that must be treated at this first level of the health system where surgical services are provided.</p> <p>A major emphasis would be resuscitation and emergency care and transfer children to a higher level of the system where more definitive care can be provided.</p> <p>Safe anesthesia and perioperative care training and training of perioperative nursing</p> <p>Consider conducting training sessions at second or third level hospitals to aid in creation of links of referral.</p>
Referral hospital	Second- level hospital	<b>I, II, III</b> Essential children's surgical services and selected more complex conditions. Skills and procedural training to address a selection of more complex surgical needs
	Third level hospital	<b>I, II, III</b> Definitive treatment for comprehensive children's surgical services. Skills and procedural training to meet this need, across disciplines. May be met through separate specialists, or through surgical, anesthesia, and nursing providers trained across disciplines for specific procedures.
	National children's hospital	<b>I, II, III</b> Subspecialist training to meet comprehensive and more super-specialized surgical needs (e.g., cardiac or neurosurgery) Multi-disciplinary focused training integrating all members of the advanced health-care delivery team.

<sup>a</sup> See Table 1 for equivalent health care facilities in the WHO classification system.

<sup>b</sup> Levels of care is based on the complexity of surgical care (see Table 2 for details)

## 8.2 Quality improvement

A variety of strategies can be used to improve the quality of healthcare delivered to children.<sup>9</sup> Perhaps most important is medical audit, which improves medical care by routinely monitoring the elements of diagnosis, treatment and outcome. By always examining the outcomes of patients, it is possible to evaluate the performance of individual providers as well as the system in which they work. Quality improvement for children's surgical care should take several forms, including routine, systematic review of the structure, process and outcomes of care (Figure 1). For quality improvement activities to be effective they must be predictable and done an ongoing manner, with feedback provided to all stakeholders so that appropriate action can be taken. Reporting requirements are likely to be less intensive at basic and intermediate levels, and more extensive at advanced/complex levels.

### 8.2.1 Structural improvement

This refers to the physical and human resources available for care. By periodically assessing the available resources (e.g. number of trained specialists, procedure volume, availability of equipment and supplies) gaps and limitations can be identified and addressed to improve the quality of care. This also helps in planning and allocation of resources to ensure optimal utilization and cost effectiveness.

### 8.2.2 Process improvement

This relates to the care that patients receive. By assessing the process of care, tasks can be standardized and checklists and guidelines can be developed based on identified gaps and problems. This will help to improve the quality of surgical care. Process improvement includes monitoring and tracking of outcome measures and peer review; it helps to improve the quality of surgical care. This also helps to establish benchmarks against which proposed improvements can be compared as well as for future aspiration. Many outcomes can be monitored; some take little time while others are more complex.

### 8.2.3 Morbidity and mortality conferences

The field of surgery has a long tradition of discussing deaths and complications in search of identifying preventable factors. Morbidity and mortality conferences are based on open, non-punitive education focused on peer review with the goal of identifying correctable problems with the individual practitioners and systems, all designed to improve performance. Predetermined audit filters (e.g., anesthetic complications, inpatient deaths, surgical site infections) can help determine which cases need peer review. Actionable factors should be identified and acted upon to improve subsequent outcomes. This last process is referred to as “loop closure” and is a critical component of all quality improvement work.

### 8.2.4 Preventable death studies

Reviewing the deaths that occur at an individual hospital or within a health system is an essential tool for quality improvement. When a preventable death is identified, steps can be undertaken to prevent similar problems in the future.

### 8.2.5 Complications

Tracking complications is an important tool for assessing quality of care. This process attempts to identify complication rates that are higher than would normally be expected. Surgical site infections, for example, are gaining greater emphasis, but more work is needed to define optimal quality metrics in resource-limited settings.

### 8.2.6 Risk-adjusted mortality

This strategy allows hospitals to compare their outcomes based on the type of patients they treat. Comparisons can be made against other hospitals and predetermined national norms and benchmarks. Identifying hospitals with higher risk-adjusted death rates can be an important first step in the performance improvement process. Risk-adjusted mortality rates are commonly used to evaluate trauma programs. These can serve as the basis for specific quality improvement interventions.

### 8.2.7 Data collection and surveillance


Children's surgical services must incorporate ongoing collection of data at hospital and national levels. Data on surgical epidemiology, disease trends and patterns, treatments and outcome, as well as resources should be continuously evaluated. This will help in planning and modification of treatment strategies, resource allocation and training, as well as prevention and control measures. Such data collection and surveillance should include the deployment of electronic medical records system, disease specific registries (e.g. congenital anomalies, trauma, cancer, etc.) and trainee databases. Integrating this data collection with existing public health surveillance will be critical to linking with broader child health initiatives.

### 8.3 Hospital assessment

Hospital assessment with verification of resources has proven to be a necessary tool for improving quality of care in high-income countries. Most low-resource settings also have some mechanism for monitoring the function of their hospitals, but the completeness and effectiveness of this monitoring is variable.<sup>42</sup> Government-run hospitals may be more likely to have monitoring systems in place than private facilities. Hence, there is a precedent for the concept of assessment and verification as a mechanism for improving and monitoring children's surgical services in most low-resource settings.

The GICS program recommends assessment and evaluation of hospitals to be done by internal country accreditation mechanisms, where they exist (e.g. Ministry of Health, training colleges). In the absence of in-country accreditation programs, invited external reviewers could assist with assessment and evaluation.

**Figure 1:** Quality improvement for children’s surgical care



## 9. RESEARCH

---

### 9.1 Overview

The importance of research in improving surgical care of children in low-resource settings cannot be overstated. Its roles are multiple: providing relevant information for management in low-resource settings, quantifying global burden of surgical disease in children, identifying gaps in provision of care, and generating data for advocacy for equitable resource allocation for the surgical care of children.<sup>43</sup> While research may be through academically affiliated institutions, non-academic institutions providing children's surgical care may also be engaging in important scholarly activity, as the range of facilities providing children's surgical care is diverse. The key research needs identified by the GICS were (1) the creation of regional research hubs, and (2) the deployment of appropriate institutional patient databases.

### 9.2 Regional research hubs

These hubs would each cover and support several related countries in one region. Ideally, they would include one or several health care researchers. Their roles would be to provide research support to children's surgical providers, including seeking funding, project design, IRB requirements, research ethics, data management and analysis, and manuscript writing and publication. They would also be able to deploy and assist in the adoption and maintenance of the institutional databases. They would offer training for providers and trainees, primarily through workshops and hands-on, on-site research fellowships mentoring providers through all the stages of a research project completion. The governance and financing of these regional hubs would ideally develop through existing collaborations between countries or may be coordinated through GICS or similar international organization. The regional research hubs should also align with the regional offices of the WHO Emergency and Essential Surgical Care offices and contribute to their

1  
2  
3 Global Surgery Database <sup>44</sup>. At the national level this data should also be shared with the Ministry  
4  
5 of Health to help inform national health policy that is country specific.  
6  
7

### 8 **9.3 Children's surgical database**

9  
10 The database requested by GICS providers needs to be simple to use at the point-of-care (therefore  
11  
12 mobile-device based), limited in complexity to under 20 fields per case, able to function in both  
13  
14 online and offline mode, and stored remotely (cloud-based in the regional research hubs). The  
15  
16 target population would be pediatric surgical inpatients, though a similar tool could be used for  
17  
18 outpatient clinic patients. Data would be prospectively collected for both ongoing audit/quality  
19  
20 management and for actual research – thus, locally the data would contain patient identifiers, while  
21  
22 the remote upload would be anonymous. Consent for data collection and research would be  
23  
24 obtained on patient admission. The principal crosscutting areas identified for data collection and  
25  
26 subsequent research were congenital birth abnormalities, childhood cancer, and injury/critical  
27  
28 care. In order to foster long-term adoption of the database collection process and create an  
29  
30 institutional culture of continuous data collection, the database would be initially deployed for  
31  
32 short-term “sprint audits” and research projects (2-4 weeks), then gradually transitioned to longer-  
33  
34 term collection. More specific disease-based registries, may also be established. Detailed research  
35  
36 objectives by level of facility are provided in Table 9.  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**TABLE 9.** Research objective for different types of facilities

Type of facility <sup>a</sup> (DCP3 classification)		Level of Care <sup>b</sup>	Research Objectives
Health center		<b>I</b>	<p>Ensure basic birth and death registry</p> <p>Generate population-based data on the prevalence of congenital surgical conditions. The necessary data collection is better achieved by incorporating basic screening for congenital anomalies into the birth and 6-week routine infant vaccination campaigns and well-baby cards.</p> <p>A simple check of a box for suspected abnormality by body area may then engender referral to a regional institution</p> <p>Such population-based data may then be analysed by paediatric surgical providers at higher health care system levels with the assistance of the regional research hubs</p>
First level hospital		<b>I, II</b>	<p>Prevalence, burden of disease and access studies, and injury epidemiology studies.</p> <p>The regional research hubs would provide the technical and logistic support for such studies such as the development of surgical indicators/ data points.</p>
Referral hospital	Second and third-level hospital	<b>I, II, III</b>	<p>Database for surgical conditions in children for both epidemiological and outcome purposes. General surgeons typically provide the surgical care of children with several common conditions, and monitoring the quality of care provided at this level is important for identifying gaps in infrastructure, personnel, and training.</p> <p>Data generated through such databases would again be made available to children's surgical providers for research projects coordinated through the regional research hubs.</p>
	Children's Hospital	<b>I, II, III</b>	<p>At the national referral "centre of excellence", research would be conducted at multiple levels. Epidemiological and outcomes studies on a variety of children's surgical conditions at the national level would be possible using the standard inpatient database.</p> <p>Disease-based registries may be established</p> <p>Higher-level interventional studies would also be fostered by the regional research hubs in order to identify best practices for low-resource settings and contextualized management solutions</p> <p>Access to care, disparities, and financial risk protection and catastrophic health expenditure due to children's surgical conditions would be analysed across populations</p> <p>Basic and translational science programs to advance knowledge in local disease processes</p> <p>Hub for research administration and courses in ethical conduct of research</p>

<sup>a</sup> See Table 1 for equivalent health care facilities in the WHO classification system.

<sup>b</sup> Levels of care is based on the complexity of surgical care (see Table 2 for details).

## 10. INTEGRATION OF CHILDREN'S SURGERY INTO HEALTH SYSTEMS

---

Children's surgery should be integrated into an overall national health plan. This should include children's surgery being integrated into childhood healthcare plans as well as the National Surgical, Obstetric and Anesthesia Plans (NSOAPs). There are also opportunities to align with obstetric - maternal health initiatives focused around prenatal care and delivery. At the institutional level, children's surgery should exist as an integral component of child healthcare services to address the social, emotional and psychological needs of children. Such integration also ensures that available physical and human resources are adequately allocated for children's surgical care. At the national level, this involves political jurisdictions designating hospitals at varying levels of complexity for children's surgical care. Emergency child health services, transfer criteria and transfer arrangements between levels of care and between hospitals must be planned to ensure timeliness and safety. National children's surgical plans may also involve discussion of the role of "niche" hospitals, especially for non-emergent specialty care, as these hospitals provide essential care in many low-resource regions of the world. Governance and finance of these hospitals within the health system and the role of public-private partnerships are key areas for discussion.

## 11. INTERACTION WITH AND COORDINATION OF STAKEHOLDERS

---

Global health initiatives have often struggled to implement changes at scale. It is reasonable to expect that introducing the guidelines for the optimal surgical care of children will face similar challenges unless there is close collaboration with multiple stakeholders. This planning will require coordination through ministries of health, promotion through professional societies and networks as well as parent-community groups in grass-roots lobbying efforts. Furthermore, continued integration and coordination of specialties related to children's surgery is paramount to successful implementation of children's surgery health initiatives. Stakeholders at different levels of the

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

hospital system, including community health centers and district hospitals must similarly be represented in order to improve surgical capacity and outcomes for children in low-resource environments.

For Peer Review

**REFERENCES**

1. Farmer DL, Bickler SW. Preface. *Seminars in pediatric surgery* 2016; 25(1): 1-2.
2. Bickler SW, Sanno-Duanda B. Epidemiology of paediatric surgical admissions to a government referral hospital in the Gambia. *Bulletin of the World Health Organization* 2000; 78(11): 1330-6.
3. Meara JG, Leather AJ, Hagander L, et al. *Global Surgery 2030: evidence and solutions for achieving health, welfare, and economic development*. Lancet 2015.
4. Mock CN, Donkor P, Gawande A, et al. *Essential surgery: key messages from Disease Control Priorities, 3rd edition*. Lancet 2015; 385(9983): 2209-19.
5. Henry JA, Abdullah F. *Global Surgical Care in the U.N. Post-2015 Sustainable Development Agenda*. *World journal of surgery* 2016; 40(1): 1-5.
6. *Strengthening emergency and essential surgical care and anaesthesia as a component of universal health coverage*. Geneva: 68th World Health Assembly, 2015.
7. Goodman LF, St-Louis E, Yousef Y, et al. *The Global Initiative for Children's Surgery: Optimal Resources for Improving Care*. *Eur J Pediatr Surg* 2017.
8. Gelb A, Morriss W, Johnson W, Merry A, al. e. *WHO-WFSA International Standards for a Safe Practice of Anesthesia Workgroup (epublication)*. Geneva: World Health Organization, 2017.
9. *Standards for improving the quality of care for children and young adolescents in health facilities*. Geneva: World Health Organization; 2018.  
[http://www.who.int/maternal\\_child\\_adolescent/documents/quality-standards-child-adolescent/en/](http://www.who.int/maternal_child_adolescent/documents/quality-standards-child-adolescent/en/) (Accessed September 23, 2018)
10. *Everybody's business - strengthening health systems to improve health outcomes: WHO's framework for action*. Geneva: World Health Organization; 2007.  
[http://www.who.int/healthsystems/strategy/everybodys\\_business.pdf](http://www.who.int/healthsystems/strategy/everybodys_business.pdf) (Accessed August 20, 2018)
11. *Monitoring the building blocks of health systems: a handbook of indicators and their measurement strategies*. Geneva, World Health Organization; 2010.  
[http://www.who.int/healthinfo/systems/WHO\\_MBHSS\\_2010\\_full\\_web.pdf](http://www.who.int/healthinfo/systems/WHO_MBHSS_2010_full_web.pdf) (Accessed August 20, 2018)
12. Alkire BC, Raykar NP, Shrimme MG, et al. Global access to surgical care: a modelling study. *The Lancet Global health* 2015; 3(6): e316-23.
13. Rhee D, Papandria D, Yang J, et al. Comparison of pediatric surgical outcomes by the surgeon's degree of specialization in children. *Journal of pediatric surgery* 2013; 48(8): 1657-63.
14. Emil SG, Taylor MB. Appendicitis in children treated by pediatric versus general surgeons. *Journal of the American College of Surgeons* 2007; 204(1): 34-9.
15. McAteer JP, Kwon S, LaRiviere CA, Oldham KT, Goldin AB. Pediatric specialist care is associated with a lower risk of bowel resection in children with intussusception: a population-based analysis. *Journal of the American College of Surgeons* 2013; 217(2): 226-32.e1-3.

16. Arca MJ, Goldin AB, Oldham KT. Optimization of care for the pediatric surgical patient: Why now? *Seminars in pediatric surgery* 2015; 24(6): 311-4.
17. Goldin AB, Dasgupta R, Chen LE, et al. Optimizing resources for the surgical care of children: an American Pediatric Surgical Association Outcomes and Clinical Trials Committee consensus statement. *Journal of pediatric surgery* 2014; 49(5): 818-22.
18. Oldham K. Optimal resources for children's surgical care in the United States. *Journal of the American College of Surgeons* 2014; 218(3): 479-87, 87 e1-4.
19. Bickler SW, Rode H. Surgical services for children in developing countries. *Bulletin of the World Health Organization* 2002; 80(10): 829-35.
20. Poenaru D. The burden of pediatric surgical disease in low-resource settings: Discovering it, measuring it, and addressing it. *Journal of pediatric surgery* 2016; 51(2): 216-20.
21. Bickler SW, Telfer ML, Sanno-Duanda B. Need for paediatric surgery care in an urban area of The Gambia. *Tropical doctor* 2003; 33(2): 91-4.
22. Butler EK, Tran TM, Nagarajan N, et al. Epidemiology of pediatric surgical needs in low-income countries. *PloS one* 2017; 12(3): e0170968.
23. Ozgediz D, Langer M, Kisa P, Poenaru D. Pediatric surgery as an essential component of global child health. *Seminars in pediatric surgery* 2016; 25(1): 3-9.
24. Ozgediz D, Poenaru D. The burden of pediatric surgical conditions in low and middle income countries: a call to action. *Journal of pediatric surgery* 2012; 47(12): 2305-11.
25. Sengoelge M, El-Khatib Z, Laflamme L. The global burden of child burn injuries in light of country level economic development and income inequality. *Preventive medicine reports* 2017; 6: 115-20.
26. Howard SC, Davidson A, Luna-Fineman S, et al. A framework to develop adapted treatment regimens to manage pediatric cancer in low- and middle-income countries: The Pediatric Oncology in Developing Countries (PODC) Committee of the International Pediatric Oncology Society (SIOP). *Pediatric blood & cancer* 2017; 64 Suppl 5.
27. Jamison DT, Alwan A, Mock CN, et al. Universal health coverage and intersectoral action for health: key messages from Disease Control Priorities, 3rd edition. *Lancet* 2018 Mar 17;391(10125):1108-1120.
28. Mock C, Lormand J, Goosen J, Joshipura M, Peden M. *Guidelines for Essential Trauma Care*. Geneva: World Health Organization, 2004.
29. Mock CN, Donkor P, Gawande A, Jamison DT, Kruk ME, Debas HT. Essential surgery: key messages from Disease Control Priorities, 3rd edition. *Lancet (London, England)* 2015; 385(9983): 2209-19.
30. Cohen MM, Cameron CB, Duncan PG. Pediatric anesthesia morbidity and mortality in the perioperative period. *Anesthesia and analgesia* 1990; 70(2): 160-7.

- 1
- 2
- 3 31. Gobbo Braz L, Braz JR, Modolo NS, do Nascimento P, Brushi BA, Raquel de Carvalho L.  
4 Perioperative cardiac arrest and its mortality in children. A 9-year survey in a Brazilian tertiary  
5 teaching hospital. *Paediatric anaesthesia* 2006; 16(8): 860-6.  
6
- 7 32. Flick RP, Sprung J, Harrison TE, et al. Perioperative cardiac arrests in children between 1988 and  
8 2005 at a tertiary referral center: a study of 92,881 patients. *Anesthesiology* 2007; 106(2): 226-  
9 37; quiz 413-4.  
10
- 11 33. Krishnaswami S, Nwomeh BC, Ameh EA. The pediatric surgery workforce in low- and middle-  
12 income countries: problems and priorities. *Seminars in pediatric surgery* 2016; 25(1): 32-42.  
13
- 14 34. Kempthorne P, Morriss WW, Mellin-Olsen J, Gore-Booth J. The WFSA Global Anesthesia  
15 Workforce Survey. *Anesthesia and analgesia* 2017; 125(3): 981-90.  
16
- 17 35. Bickler S, Ameh E. *Surgical Care for Children: Guidelines for the primary referral hospital*:  
18 Macmillan; 2011.  
19
- 20 36. *Pocketbook of hospital care for children*: World Health Organization, 2005.  
21
- 22 37. Gosselin R, Spiegel D, Foltz M. *Global Orthopedics*: Springer; 2014.  
23
- 24 38. Carter LL. *Principles of Plastic Surgery in Africa*: Pan African Academy of Christian Surgeons;  
25 2003.  
26
- 27 39. Semer N. *Practical Plastic Surgery for Nonsurgeons*. New York, Lincoln, and Shanghai: Authors  
28 Choice Press; 2007.  
29
- 30 40. *Surgical Care at the District Hospital*. Geneva, World Health Organization; 2003.  
31 <http://www.who.int/surgery/publications/en/SCDH.pdf?ua=1> (Accessed August 28, 2018).  
32  
33
- 34 41. e-SAFE: Safer Anesthesia Through Education. 2014. <http://www.e-safe-anaesthesia.org/>  
35 (Accessed August 20, 2018).  
36
- 37 42. Galukande M, Katamba A, Nakasujja N, et al. Developing hospital accreditation standards in  
38 Uganda. *The International journal of health planning and management* 2016; 31(3): e204-18.  
39
- 40 43. Greenberg SL, Ng-Kamstra JS, Ameh EA, Ozgediz DE, Poenaru D, Bickler SW. An investment  
41 in knowledge: Research in global pediatric surgery for the 21st century. *Seminars in pediatric*  
42 *surgery* 2016; 25(1): 51-60.  
43
- 44 44. *Emergency and Essential Surgical Care- Global Database*. Geneva, World Health Organization.  
45 [http://www.who.int/surgery/eesc\\_database/en/](http://www.who.int/surgery/eesc_database/en/) (Accessed on August 20, 2018).  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**APPENDIX 1: Childhood surgical procedures for different facility types classified by pediatric surgical subspecialty**

Specialty	Health Center	First-level hospital	Referral hospital		
			Second-level hospital	Third-level hospital	National children's hospital
<b>Anesthesia</b>	Local anesthesia	General Anesthesia without invasive monitoring or post-operative ventilation; Children < 3 months old and ASA > 3 referred	General Anesthesia for most Major surgeries; all ages	General Anesthesia for all specialized surgical procedures; all ages and co-morbidities	General Anesthesia for specialized surgical procedures, all ages and co-morbidities
<b>Cardiac Surgery</b>	Cardiovascular examination for congenital heart defects, ECG	Medical treatment (PE2) of PDA-dependent diseases, Basic Echocardiography	Same as First-level hospital	Surgical closure of PDA, Surgical correction of simple coarctation (non-neonatal)	All cardiac surgery procedures including Catheterization for ASD and pulmonary valve stenosis, Palliative treatment of TOF
<b>Critical Care</b>	Recognition of illness severity, Treatment of pneumonia, dehydration	Airway Management, Resuscitation and stabilization, Venous Cut-down, Treatment of sepsis, Pneumonia, Meningitis	Acute pediatric emergencies, Treatment of ARDS, Septic Shock, Coma	All NICU and PICU procedures, Treatment of traumatic brain injury, Meningoencephalitis, Cardiac failure	All NICU and PICU procedures, All co-morbid conditions
<b>General Surgery</b>	I&D of superficial abscess, Simple and minor lacerations	Trauma and Emergency laparotomy, Emergency colostomy, Hernia repair Trauma amputation, Thoracostomy, Anoplasty (cut-back)	Trauma Thoracotomy, ARM & Hirschsprung Disease	All General Surgery procedures including repair of gastroschisis and omphalocele, intraabdominal malignancy	All General Surgery procedures
<b>Neurosurgery</b>	Diagnosis and Triage of Neurotrauma	Exploratory Burr Holes, Trans-fontanelle Ventricular Tap	Trauma Craniotomy	ICP monitor, Intracranial infection, Hydrocephalus, Myelomeningocele, Neuroendoscopy	Vascular Malformations, Complex cranial tumors, Congenital Spinal Deformities, Cranial Malformations, Epilepsy
<b>Ophthalmology</b>	Antibiotic eye drops, Irrigation of eye after chemical injury	Foreign Body, Corneal Scratch, Hyphema	Lid Lacerations, Corneal laceration, Cataract operation (> 10 years old)	Cataract surgery (all ages), squint surgery, primary closure of trauma, Simple ptosis surgery	Glaucoma and corneal procedures, Retinopathy of premature procedures, Oculoplastic procedures

<b>Oral Maxillofacial Surgery</b>	Simple lacerations, Antibiotics	Dental extraction	Repair of cleft lip/palate Fabrication of feeding plates, Pre-surgical naso-alveolar molding,	Facial reconstruction, Distraction osteogenesis, Osteosynthesis	All Oral Maxillofacial Procedures
<b>Orthopedic Surgery</b>	Splinting for simple fractures	Wound care for open fractures, Drainage for orthopedic infections (Osteomyelitis), Ponseti technique for club foot	Closed Reduction for Hip Dysplasia, Pinning slipped capital femoral epiphysis, Fasciotomy, Syndactyly release	Definitive management of osteomyelitis/septic arthritis, All vascular injuries, Spina Bifida, Muscular Dystrophy	Excision/Reconstruction of boney tumors, Correction of brachial plexus injury, Tendon transfers, multiple congenital deformities
<b>Otolaryngology</b>	Otoscopy, Rhinoscopy, Removal of foreign body from ear, nose	Drain superficial head and neck abscess, Tonsillitis treatment, Swallowed foreign body, Torticollis	Excision of benign neck mass, tonsillectomy	Parapharyngeal & retropharyngeal abscess, Tympanoplasty, Ossiculoplasty, Tympano-Mastoidecomy, Bone anchored hearing aids	Laryngotracheal reconstruction, Slide trachoplasty, Complex head and neck mass excision
<b>Plastic Surgery</b>	Simple Laceration Repair, Management of 1 <sup>st</sup> degree burns	Management of 2 <sup>nd</sup> degree burns, Escharotomy, Stabilization of Vascular Trauma and Wound Sepsis	Complex laceration repair, Vascular shunt to preserve inflow in trauma, Revision amputation,	Soft tissue coverage (grafts, local flaps, free tissue transfer), Repair of cleft lip/palate, management of craniofacial & congenital hand anomalies	Complex reconstruction after tumor excision, Management of complex cranial vault and hand anomalies
<b>Urology</b>	Urinary Tract Infection, Local wound care after circumcision	Circumcision, Paraphimosis, Orchiopexy, preputial-plasty, Urethral catheterization	Cystostomy, Vesicostomy, Orchiopexy, Imperforate hymen	Hypospadias and Epispadias treatment, Urethral valve resection, complex repair of bladder exstrophy, nephrostomy, Genitoplasty, Urolithiasis management	Mitroffanoff intervention for spina bifida, Kidney Transplant

Abbreviations- ARDS- Acute Respiratory Distress Syndrome; ARM- Anorectal Malformation; ASA- American Society of Anesthesiologist physical status classification; I&D- Incision and Drainage; NICU- Neonatal Intensive Care Unit; PE2- Prostaglandin E2, PICU- Pediatric Intensive Care Unit; TOF- Tetralogy of Fallot

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

**APPENDIX 2: Childhood surgical procedures for different facility types classified by Surgical Condition**

Surgical Condition	Health Center	First-level hospital	Referral hospital		
			Second-level hospital	Third-level hospital	National children's hospital
<b>Trauma/ Injuries</b>	<u>Resuscitation:</u> BLS <u>Injuries:</u> Wound care for simple lacerations, <u>Fractures:</u> closed non-displaced <u>Burns:</u> 1 <sup>st</sup> degree	<u>Resuscitation:</u> ALS, ATLS <u>Injuries:</u> Thoracostomy Trauma Laparotomy Trauma-amputation <u>Fractures:</u> Closed reduction, Open I&D with external fixator placement <u>Burns:</u> 1 <sup>st</sup> , 2 <sup>nd</sup> degree < 10% TBSA, excluding face, hands, perineum Simple escharotomy	<u>Resuscitation:</u> All, NICU, PICU <u>Injuries:</u> Trauma Thoracotomy Renal/Urethral trauma Vascular shunt Fasciotomy Craniotomy <u>Fractures:</u> Complex nonunion & malunion, Pinning for slipped capital femoral epiphysis <u>Burns:</u> 1 <sup>st</sup> , 2 <sup>nd</sup> degree < 10% TBSA excluding face, hands, perineum	<u>Resuscitation:</u> All, NICU, PICU <u>Injuries:</u> Complex Neuro-trauma Complex neurovascular extremity injuries including limb sparing operations <u>Fractures:</u> All including Cranial fractures <u>Burns:</u> >10% TBSA Burns including face, hands, perineum	All
<b>Congenital anomalies</b>	<u>Screening:</u> Abdominal hernias, Urogenital abnormalities, Hip dysplasia & limb deformities, Congenital heart defects, Neural tube defects, Craniofacial anomalies	Inguinal hernia/hydrocele repair, Relief of urinary obstruction: catheterization & suprapubic cystostomy, Club foot: non-operative manipulation, Treatment of PDA dependent disease with IV prostaglandin E2	Neonatal intestinal obstruction, Meatal stenosis, Orchidopexy for testicular torsion & palpable undescended testes, Closed reduction and Pavlik harness for hip dysplasia, Club foot operative treatment, Hydrocephalus	Gastroschisis & Omphalocele Anoplasty and Pull through for ARM/Hirschsprung, Hypospadias/epispadias, Development dysplasia of Hip, Clubfoot, PDA, Coarctations, TOF, Spina bifida, myelomeningocele, meningocele, dysmorphic syndromes, craniofacial abnormalities, syndactyly	All

<b>Infections</b>	I&D of superficial abscess	Thoracostomy for empyema Laparotomy for Appendectomy, typhoid complications, I&D of superficial, pyomyositis, perianal, anorectal abscess, Septic arthritis, Osteomyelitis, Extraction, drainage of abscess, caries	Leprosy, Tonsillectomy I&D of neck abscess	Advanced osteomyelitis, tenosynovitis, sepsis and septic shock, I&D of retropharyngeal abscess, treatment for intracranial infections (hydrocephalus, subdural empyema)	All
<b>Tumors</b>	Screening for all abdominal tumors	Biopsy of palpable or U/S visualized mass Excision of benign tumors/cysts	Biopsy of all tumors, Excision of benign tumors/cysts Diagnosis of malignant tumors	Abdominal and urologic malignant tumors Malignant tumors of extremities Head and neck tumors Simple intracranial tumors	All
<b>Others</b>	Male circumcision using plastibel, Rectal prolapse reduction Foreign body removal from ears, nose, mouth	Central venous line Laparotomy for Intestinal obstruction	Functional and obstructive urinary pathology (pelvic-ureteral obstruction) Arthrotomy synovectomy Shunt Tap & Trans-fontanelle Ventricle Tap	Muscular dystrophy, limb reconstruction ureteral duplication anomalies, vesico-ureteral reflux Free Tissue Transfer, Local Flaps, Amniotic band release tympanoplasty, Ossiculoplasty,	All

Abbreviations-ALS- Advanced Life Support; ATLS- Advanced Trauma Life Support; BLS- Basic Life Support; ARM- Anorectal Malformation; I&D- Incision and Drainage; NICU- Neonatal Intensive Care Unit; PICU- Pediatric Intensive Care Unit; PDA- Patent Ductus Arteriosus, TBSA- Total Body Surface Area; TOF- Tetralogy of Fallot; U/S- Ultrasound

APPENDIX 3. Human resource for different types of facilities

Type of facility (DCP3 classification) <sup>a</sup>	Level of Care <sup>b</sup>	Human Resources
Health center	I	<u>Non-Physicians:</u> Trained Birth Attendants, Community Health Workers, General Practice Nurse <u>Physicians:</u> General Medical Physician, Clinical Officers
First-level hospital	I-II	<u>Non-Physicians:</u> Perioperative Nurses, Physical and Occupational Therapist, Orthopedic officer, X-ray Technician, Biomed and Laboratory Technician, Anesthesiologist +/- Anesthesia provider, Hospital Administrator <u>Physicians:</u> General Doctor or Non-Physician Surgical Provider
Referral hospital	Second- level hospital	<u>Non-Physicians:</u> In addition to First-level Hospital Requirements, Pediatric Nurse, Orthotics Personnel available (on call basis), Respiratory Care Specialists, Nutritionist, Dental Technologist <u>Physician Resources:</u> Physician Anesthesiologist, General Surgeon, Orthopedic Surgeon, Neonatologist, Pediatric Critical Care Physician, Radiologist (On-call, Pediatric Radiologist Consultation by email)
	Third level hospital	<u>Non-Physicians:</u> In addition to Second-level Hospital Requirements, Cardiac Perfusionists and Speech pathologist, Palliative care, Trainees- Nursing, Specialty Staff Students, Medical Students, Residents, Fellows <u>Physician Resources:</u> <u>Pediatric Disciplines:</u> Cardiology (with interventional capability), Oncology, Gastroenterology, Neurology, Nephrology, Radiology, Anesthesiologist; Intensivist; Neonatal Intensivist and Others <u>Pediatric Surgery Disciplines (trained in pediatric subspecialty):</u> General (Pediatric) Surgeon, Cardiac Surgeon; ENT; Oral and Maxillofacial Surgeon; Orthodontist; Orthopedic Surgeon; Plastic and Reconstructive Surgeon; Urologist, Neurosurgeon with Pediatric Experience, Ophthalmologist
	National children's hospital	<u>Non-Physicians:</u> All Specialists and Staff in First and Second-level Hospital including Cardiac Perfusionists and Speech pathologist, Palliative care, Trainees- Nursing, Specialty Staff, Medical Students, Residents, Fellows <u>Physician Resources:</u> <u>Pediatric Specific Disciplines:</u> Cardiology (with interventional capability), Oncology, Gastroenterology, Neurology, Nephrology, Radiology including neuroradiology, Anesthesiologist including neuro and cardiac; Pediatric Intensivist; Neonatal Intensivist and Others <u>Pediatric Surgery Disciplines (trained in pediatric subspecialty):</u> General (Pediatric) Surgeon, Cardiac Surgeon; ENT; Oral and Maxillofacial Surgeon; Orthodontist; Orthopedic Surgeon; Plastic and Reconstructive Surgeon; Urologist, Neurosurgeon, Ophthalmologist, Transplant Surgeon

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

- <sup>a</sup> See Table 1 for equivalent health care facilities in the WHO classification system.
- <sup>b</sup> Levels of care is based on the complexity of surgical care (see Table 2 for details).

For Peer Review

# OPTIMAL RESOURCES FOR CHILDREN'S SURGICAL CARE

## II. Supplies, Equipment and Infrastructure



**GICS**

**Global Initiative for Children's Surgery**

*Version 2.0 (September 24, 2018)*

## Contents

1.	Summary.....	3
2.	Supplies and Equipment for Basic and Intermediate Care.....	4
	Table 1: Overview of essential childhood surgical supplies for basic and intermediate surgical care classified by specialty.....	4
	Table 2: Surgical Equipment for First Level Hospital.....	5
	Table 3: Surgical Consumables for First Level Hospital.....	9
	Table 4: Surgical Instruments for Major Procedures at First Level Hospital.....	12
	Table 5: Surgical Instruments for Minor Procedures at First Level Hospital.....	16
	Table 6: Anesthesia Supplies at First Level Hospital.....	18
	Table 7: Pediatric specific surgical supplies for First Level Hospital.....	21
3.	Supplies, Equipment and Infrastructure for Advanced and Complex Care.....	23
	Table 8: Overview of essential childhood surgical supplies for referral hospitals by specialty.....	23
	Table 9: Select pediatric specific surgical supplies for advanced and complex surgical care.....	24
	REFERENCES.....	30

## 1. Summary

This document is intended as a supplement to the Optimal Resources for Children's Surgical Care: Guidelines for Different Levels of Care. It contains detailed information on the supplies, equipment and infrastructure necessary to perform essential children's surgery. The information is organized by hospital level as well as by surgical and anesthesia subspecialty.

The focus of this document is the essential supplies and equipment necessary to provide intermediate care for children at a first-level hospital. The main reference are Tables 2 – 6, which itemize the essential supplies and equipment necessary to provide safe surgery at first-level hospitals<sup>1</sup>. Table 7 is the additional supplies and equipment necessary to provide surgical care specifically to children at first level hospitals. This additional information was developed directly from the specialty working groups described in the OReCS: Guidelines for Different Levels of Care document.

Equipment recommendations by the WHO Essential Surgical Care (EESC) Compiled from the WHO Manual for Surgical Care at the District Hospital is contained in our itemized lists<sup>2</sup>. The WHO list appropriately recommends both a pediatric anesthesia system and pediatric resuscitator sets at the district level hospital. We include specific equipment recommendations including ultrasound, and neonatal and pediatric sizes for sphygmomanometers, central venous catheters, chest tubes, ventricular drains, endotracheal tubes and cling film for bowel coverage.

The additional sections of this document detail the essential supplies and equipment necessary to provide advance and complex surgical care for children. This would typically be delivered at secondary and tertiary hospitals, respectively. An overview table of supplies and equipment by specialty for advanced and complex care is provided (Table 8). This information is followed by an itemized list of select and necessary supplies and equipment for children's surgery.

## 2. Supplies and Equipment for Basic and Intermediate Care

**Table 1:** Overview of essential childhood surgical supplies for basic and intermediate surgical care classified by specialty

Subspecialty	Pediatric Specific Supplies for Health Center	Pediatric Specific Supplies for Intermediate Care
<b>Anesthesia</b>	Pulse Oximeters, Thermometers, Non-invasive blood pressure monitors, Stethoscope	Oxygen supply, CO2 monitor, Laryngoscopes and Endotracheal tubes, Regional anesthesia & IV infusion
<b>Cardiac Surgery</b>	Pulse Oximeter with neonatal probe, Electrocardiogram	Echocardiography equipment
<b>Critical Care</b>	Equipment for acute care resuscitation, Basic medicine and equipment store	Equipment for resuscitation and stabilization, Equipment and supplies for airway management
<b>General Surgery</b>	Patient transfer ability to higher levels of care, Incision and Drainage Kit	Basic laparotomy instrument set, X-ray
<b>Neurosurgery</b>	Basic equipment to assess neurotrauma	Hudson Brace and Gigli saw for Burr Holes, Equipment for Trans-Fontanelle Ventricular Tap
<b>Ophthalmology</b>	Headlamp, Ophthalmoscope, Eye pads, Visual acuity chart, Saline for eye irrigation	Same as Health Center
<b>Oral Maxillofacial Surgery</b>	Dental Equipment- basic examination and surgical hand instruments, Arch bars & wires, Surgical drill and burs	Dental casting machine
<b>Orthopedics Surgery</b>	Wound care supplies, Splints, Plaster Bandages, Crepe Bandages	Basic orthopedic instrument set, Plaster rolls, X-ray machine, Equipment for plaster removal
<b>Otolaryngology</b>	Headlight, Basic ear and nose instruments, Screening Audiometer, Otoscope, Rhinoscope	Same as Health Center
<b>Plastic Surgery</b>	Suture material, Basic wound dressing packs: sterile field, dressing tray, dressing towel, cotton balls/swabs, forceps	Basic laparotomy or cesarean section instrument set
<b>Radiology</b>	Ultrasound, X-ray, C-arm	Computed Tomography Scanner
<b>Urology</b>	Wound care supplies, Antiseptics, Antibiotics	Urinary catheters, Basic minor surgical instrument set, Basic wound dressings packs

**Table 2: Surgical Equipment for First Level Hospital**

<b>OPERATING TABLE</b>	
Hydraulic or Motor Driven	
Height: approximately 85 cm.	
Tabletop dimensions: Length ~200cm, Width ~80 cm	
Exposed metallic parts to be made from stainless, acid proof steel.	
Mobile table base with central braking system	
Flexible inclination angles for back rest and leg rest, kidney elevator, Trendelenburg and reverse-Trendelenburg positions	
Trendelenburg and Reverse Trendelenburg at least 25°	
Lateral tilt at least 18°	
Back rest inclination angle at least +55° to -25°	
Head rest inclination angle at least 45° up	
Head rest inclination angle at least 20° down	
Automatic stop in horizontal position	
<b>Accessories</b>	
Anesthetic Screen with clamp with telescopic tubes	
Body restraint Strap with clamp	
Padded Shoulder supports	
Padded Leg support with swivel type clamp	
Padded arm Rests 450 -500 mm long with two arm Clamps	
Padded Lateral support with universal attachment Clamp	
Padded rubber mattresses with Mackintosh covers	
Head rest, IV drip stands attachable to the table	
Gel cushions for head and neck support	
<b>ORTHOPAEDIC TABLE ACCESSORY</b>	
Attachments to operating table to allow positioning for specific procedures	
Radiolucent surface	
<b>ORTHOPAEDIC TRACTION SYSTEM</b>	
<b>MOBILE C-ARM X-RAY SYSTEM</b>	
Portable imaging system with digital Flat Panel Detector	
Motorized, with range of movements: vertical >400 mm, horizontal travel >200 mm	
Orbital movement -30 to +90 degrees	
Swing/panning movement: +/- 12 degrees or more	
Depth of C-arm: 650 mm or more	
Needs to be lightweight, flexible and portable	

1	
2	
3	Imaging possible in multiple positions: sitting, supine (and possibly standing?)
4	Capability of pulse fluoroscopy to reduce radiation exposure
5	Connectivity to EMR database
6	Display - real-time imaging with viewing system
7	
8	
9	<b>SURGICAL STOOL</b>
10	Ergonomic design
11	Fixed flexible height with stump feet and anti-static cushion
12	Seamless upholstery
13	
14	
15	<b>ANAESTHETIST STOOL</b>
16	Ergonomic design
17	Seamless upholstery
18	
19	
20	<b>MAYO TABLE</b>
21	Stand for instrument trays
22	Antistatic castors, Stainless steel tray, 500 x 375 x 254mm
23	
24	<b>INSTRUMENT TROLLEY</b>
25	Stainless steel shelves
26	Anti-static castors, 610mm X 460mm
27	
28	
29	<b>ECG MONITOR</b>
30	Recording ECG Leads: 12 standard Leads
31	Recording Channels: 3/1 User selectable
32	LCD display of ECG
33	Lead switching: manual and automatic
34	Sensitivity, mm/mV: 5, 10, 20.
35	Calibration signal: automatic and manual.
36	Diagnostic frequency range 0.67-150 Hz or greater
37	Filters for mains frequency, low frequency, muscle artefact, high frequency
38	Power requirements: 220 V, 50 Hz
39	
40	
41	<b>Accessories</b>
42	Patient cable
43	Chest electrodes
44	Limb electrodes
45	Strap electrodes
46	Bottle of ECG gel
47	
48	<b>OPERATING LIGHTS</b>
49	Cordless
50	
51	
52	
53	
54	
55	
56	
57	
58	
59	
60	

1	
2	
3	LED light source
4	Li-ion battery with cost-effective replacement
5	Ergonomic design
6	Adjustable light intensity
7	Light field size 15 - 30 cm.
8	Color temperature 4200° Kelvin
9	Working range without focusing approximately 70 - 140 cm
10	
11	
12	
13	
14	<b>ELECTROSURGICAL UNIT</b>
15	High-frequency electrosurgical unit for execution of monopolar and bipolar surgery
16	Maximum output 300 W for monopolar cut
17	Activation: Double pedal switch to toggle between monopolar and bipolar functions
18	Bipolar electrode with pedal switch or with automatic Start/Stop system for coagulation
19	Memory: User shall be able to use at least 4 working programs
20	Safety: Neutral plate safety circuit, notifications and alerts with automatic reductions in power for safety
21	
22	
23	
24	<b>Accessories</b>
25	Operative Foot-switch (usable as alternative to handle)
26	Reusable neutral plate
27	Kit of autoclavable Electrodes
28	Bipolar Forceps: 6
29	Straight Forceps - 18 cm (7")
30	
31	
32	
33	<b>SUCTION PUMP</b>
34	Suction unit for Major Surgery Procedures, should be mains-powered
35	Mobile on 4 anti-static castors
36	Oil-free pump, maximum suction of at least 500 mm Hg
37	Free flow rate at least 25 l/min
38	Pedal Action
39	Protective thermal cut-out relay
40	Motor-protection cap preventing aspirated liquids or secretions from reaching and damaging the vacuum pump
41	Suction command with continuous adjustment
42	2 X 2,000ml canisters with airproof screwing-cap and independent overflow devices
43	Fast connectors and silicone tubing
44	Power Supply: 220V AC 50Hz.
45	Ventilation Fan for overheating
46	
47	
48	
49	
50	
51	
52	
53	<b>Accessories</b>
54	Silicone Tubing, Sterilizable
55	
56	
57	
58	
59	
60	

1	
2	
3	Transparent Cannula Holder, Sterilizable
4	Anti-Bacterial Filters
5	
6	<b>ANAESTHESIA MACHINE W/ VENTILATOR AND MONITOR</b>
7	
8	Anesthesia Machine used for delivering anesthetic agents to patient during surgery and ventilate patient
9	Patient monitoring system: ECG, EtCO <sub>2</sub> , Pulse Oximeter and airway pressure, NIBP, Temperature
10	
11	Anesthesia gas delivery system, O <sub>2</sub> /N <sub>2</sub> O/Compressed Air
12	Circle absorber system.
13	
14	Precision vaporizer for halothane
15	Anesthesia ventilator
16	
17	<b>Accessories</b>
18	Resuscitators, Adult / Child
19	
20	Set of face masks, 3 Sizes
21	
22	Corrugated hoses and tubing set for Pipeline O <sub>2</sub>
23	Corrugated hoses and tubing set for Pipeline Air
24	
25	Hose assembly for piped Oxygen supply
26	Hose assembly for piped Nitrous Oxide supply
27	Hose assembly for piped Air-supply
28	
29	Regulator and manometer for the pipe pressure hose (60 psi)
30	Connection from ventilator to anesthetic apparatus, autoclavable/reusable
31	
32	Test bag, 1 liter
33	Peak Flow Meter
34	
35	Power cord with grounding wire
36	<b>OTHER FURNITURE</b>
37	
38	Kickabout with s/s bowl
39	Drip stands, Double-hooked, Telescopic
40	
41	Radiation Shielding Apron, 0.35 mm Pb equivalent, Frontal, S, M, L, XL
42	Radiation Shielding Apron, 0.35 mm Pb equivalent, Full, S, M, L, XL
43	
44	
45	
46	
47	
48	
49	
50	
51	
52	
53	
54	
55	
56	
57	
58	
59	
60	

**Table 3: Surgical Consumables for First Level Hospital**

<b>DRESSING SUPPLIES</b>	
Precision Blood Glucose Test Strips	
Glucometers	
Gloves, Surgical, Sterile – 7, 7.5, 8, 8.5	
Gloves, Examination, Latex, Non-sterile, S, M, L	
IV tubing	
Betadine Solution	
Betadine Scrub	
Syringe, 3 ml, 5 ml, 10 ml, 20 ml	
Syringe, Insulin, 100 IU/1 ml + Fixed Needle	
Cast Padding, 3 inch, 4 inch, 6 inch	
Cast, Fiberglass, 3 inch, 4 inch	
Resting ECG Electrodes	
Gauze, Compression, Paraffin, 10 cm x 10 cm, sterile	
Gauze, Compression, 10 cm, 12 plies, 17 threads, sterile	
Cotton wool, hydrophilic, roll, 500 g	
Tape, Adhesive, roll, 2 cm	
Tape, Adhesive, roll, extensible, nonwoven, 10 cm x 10 m	
Tape, Adhesive, roll, perforated, 10 cm	
Bandage, Adhesive, elastic, 10 cm x 3 m	
Bandage, Extensible, Non-adhesive, 6 to 7 cm x 4 m	
Bandage, Crepe (Velpeau), 10 cm x 4 m	
Bandage, Elastic Tubular Net, Arm/leg, roll 25 m	
Strip, Adhesive for skin closure (Steri-strip), 6 mm, per 3	
Dressing, Haemostatic (Quikclot ACS+), unit	
Mask, Respirator, N95 (Fluidshield N95)	
Mask, Surgical	
Thermometers - digital	
Thermometers – oral glass	
Tegaderm, 4.75 cm x 4.75 cm, Transparent	
Bag, Urine, 2 L, Drainable + Non-return valves, Sterile	
Catheter, Urinary, Foley, Balloon, CH 10, 12, 16, 20 Sterile	
Drawsheet, disposable, plastified, absorbable, 60 x 60 cm	
Yankeur and pool suction tips - disposable/ reusable	
Scrub dispensers	

1	
2	
3	Ratex sponges - 4x4
4	
5	Surgical Blades - Size 10, 11, 12, 15, 20
6	<b>LINEN</b>
7	
8	Clogs, Operation theatre, Polyuretane, Washable
9	
10	Drape, Surgical, woven, 1 x 1 m & 1.5 x 1.5 m
11	Drawsheet, Rubber, 100 x 180 cm, with rivets
12	Drawsheet, Disposable, plastified, absorbable, 60 x 60 cm
13	
14	Drape, Sterile, nonwoven, 45 x 75 cm
15	Drape, sterile, nonwoven, Cover for Mayo table
16	
17	Drape, sterile, nonwoven, Cover for Caesarian Section Set
18	Drape, sterile, nonwoven, Cover for Orthopaedics Set
19	
20	Drape, sterile, nonwoven, Cover for Universal/Laparotomy Set
21	Isolation gown, single use, non-sterile
22	
23	Gown, Surgical, woven M, L, XL, XXL
24	Shoe Cover, non woven, Single Use
25	
26	Scrub Suit, Surgical, woven, S, M, L
27	<b>SUTURES</b>
28	
29	Prolene- 2-0,3-0,4-0 cuticular
30	Prolene- 5-0 plastic
31	
32	Prolene- 3-0,4-0 cardiovascular
33	Prolene - 0.0,1-0,2-0(general closure)
34	
35	Silk 0.0,1-0(general closure)
36	Silk 2-0(gastro intestinal)
37	
38	Ethibon- 0.0,1-0,2-0,5-0(general closure)
39	Ethilon- 3-0,4-0 cuticular
40	
41	Ethilon- 5-0 plastic
42	Ethilon- 1-0(general closure)
43	
44	PDS - 1-0(general closure)
45	PDS - 2-0(gastrointestinal)
46	
47	PDS- 3-0, 4-0 cuticular
48	Vicryl - 0.0,1-0,2-0,3-0(general closure)
49	Vicryl - 2-0,3-0(gastrointestinal)
50	
51	Vicryl - 3-0,4-0 cuticular
52	
53	Chromic - 0.0,1-0,2-0,3-0(general closure)
54	Chromic - 2-0,3-0(gastro intestinal)
55	
56	
57	
58	
59	
60	

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Chromic - 1-0,2-0(blunt needle 3/4 circle)
Surgical absorbable hemostat 3" x 4"

For Peer Review

**Table 4:** Surgical Instruments for Major Procedures at First Level Hospital

ITEM	QUANTITY
<b>ABDOMINAL/LAPAROTOMY SET</b>	
Scalpel Handle (#4), Standard	1
Scalpel Handle (#4), Long	1
Forceps, Tissue, Standard, 14.5 cm/1x2 Teeth, Straight	1
Forceps, Tissue, Standard, 25 cm/1x2 Teeth, Straight	1
Forceps, Tissue-Grasping, Duval, 23 cm, Jaws 20 mm	2
Forceps, Tissue-Grasping, DeBakey, 15 cm, 2mm Atraumatic elastic jaw	1
Forceps, Peritoneal, Faure, 21 cm, Slightly curved	2
Forceps, Intestinal, Doyen, 23 cm, Straight, Atraumatic elastic jaw	2
Forceps, Intestinal, Doyen, 23 cm, Curved, Atraumatic elastic jaw	2
Forceps, Hemostatic, Kelly, 14 cm, Curved, Non-toothed	10
Forceps, Hemostatic, Bengolea, 20 cm, Curved, Serrated	4
Forceps, Hemostatic, Crafood, 24 cm, Curved	2
Forceps, Hemostatic, Halsted-Mosquito, 12.5 cm, Non-Toothed, Curved	6
Forceps, Hemostatic, Kocher, 14 cm/1x2 Teeth, Straight	2
Forceps, Hemostatic, Kocher, 14 cm/1x2 Teeth, Curved	2
Forceps, Hemostatic, Mixer, 14 cm, Very delicate	1
Forceps, Hemostatic, Mixer, 23 cm, Delicate	1
Forceps, Hemostatic, Heiss, 24 cm, Curved, Right-angled	1
Forceps, Dressing, Standard, 20 cm, Straight	1
Forceps, Towel-Clamp, Backhaus, 13 cm	8
Forceps, Sponge, Foerster, 24 cm, Straight, Serrated jaws	2
Needle-holder, Baby-Crile-Wood, 15 cm, Standard	1
Needle-holder, Mayo-Hegar, 23 cm, Standard	1
Scissors, Mayo, 17 cm, Curved	1
Scissors, Mayo, 23 cm, Curved	1
Scissors, Metzenbaum, 17 cm, Curved	1
Scissors, Metzenbaum, 23 cm, Curved	1
Retractor, General, Deaver, Standard, 30 cm, 38 mm Edge, Flexible	2
Retractor, General, Farabeuf, 15 cm, Double-ended	2
Retractor, General, Langenbeck, 24 cm, 160 x 45 mm	2
Retractor, Abdominal, Balfour, Standard, Complete Set	1
Spatula, Abdominal, 25 cm x 27 mm, Malleable	2

Suction Tube, Poole, 23 cm x 10 mm diameter, Straight	1
Kidney Dish, Medium, 250 x 140 x 40 mm, Stainless Steel	2
Galipot, Round, 500 ml, 12 cm diameter, Stainless Steel	2
Towel Clips	4
Towel Clips, Non-perforating	2
Sterile, Blade, No. 22 (for Scalpel #4)	1
Sterile, Suction Tip, Yankauer, 11.1" x 30F Tip	1
Basket, Instruments, wired 24 x 25 x 10 cm, for sterilization	1
<b>CAESAREAN SECTION SET</b>	
Scalpel Handle (#4), for 20 Blade	1
Forceps, Tissue, Standard, 14.5 cm/1x2 Teeth, Straight	1
Forceps, Tissue, Standard, 25 cm/1x2 Teeth, Straight	1
Forceps, Tissue-Grasping, Duval, 23 cm, Jaws 20 mm	4
Forceps, Hemostatic, Kelly, 14 cm, Curved, Non-toothed	2
Forceps, Hemostatic, Wertheim, 23 cm/1x2 teeth, Straight	2
Forceps, Hemostatic, Green-Armytage, 22 cm x 13 mm	2
Forceps, Dressing, Blank, 14.5 cm, Straight, Atraumatic serrations	1
Forceps, Towel-Clamp, Backhaus, 13 cm	4
Forceps, Sponge, Foerster, 24 cm, Straight, Serrated jaws	2
Needle-holder, Mayo-Hegar, 18 cm, Standard	1
Scissors, Mayo, 17 cm, Curved	1
Scissors, Metzemaum, 18 cm, Curved	1
Retractor, General, Farabeuf, 15 cm, Double-ended	2
Retractor, Abdominal, Gosset, Central + 2 x 28 mm Blades	1
Suction Tip, Poole, 23 cm x 10 mm diameter, Straight	1
Kidney Dish, Medium, 250 x 140 x 40 mm, Stainless Steel	2
Gallipot, Round, 100 ml, 80 x 35 mm, Stainless Steel	2
Towel Clips	4
Towel Clips, Non-perforating	2
Sterile, Blade, No. 22 (for Scalpel #4)	1
Sterile, Suction Tip, Yankauer, 11.1" x 30F Tip	1
Basket, Instruments, wired 24 x 25 x 10 cm, for sterilization	1
<b>BASIC ORTHOPAEDIC SET</b>	
Scalpel Handle (#4), for 20 Blade	1
Metal Ruler 15 cm	1

1		
2		
3	Forceps, Towel-Clamp, Backhaus, 13 cm	4
4	Forceps, Hemostatic, Halsted-Mosquito, 12.5 cm, Curved	6
5	Forceps, Hemostatic, Pean, 14 cm, Straight	1
6	Forceps, Hemostatic, Collin, 16 cm, Oval jaws	1
7	Forceps, Dressing, Standard, 14.5 cm, Straight	2
8	Forceps, Tissue, Standard, 16 cm/1x2 Teeth, Straight	2
9	Forceps, Tissue, Allis, 15 cm/2x3 Teeth, Standard	2
10	Forceps, Tissue, Allis, 19 cm/5x6 Teeth, Standard	2
11	Forceps, Tissue, Stand, 14.5 cm x 1x2 Teeth, Straight	1
12	Needle-holder, Mayo-Hegar, 18 cm, Serrated	1
13	Needle-holder, Baby-Crile-Wood, 15 cm, Delicate	1
14	Scissors, Dissecting, Mayo, 17 cm, Curved	2
15	Scissors, Dissecting, Metzenbaum, 18 cm, Curved	2
16	Scissors, Operating, 14 cm, Straight, Sharp/Sharp	1
17	Scissors, Operating, 14 cm, Straight, Sharp/Blunt	1
18	Scissors, Operating, 14 cm, Curved, Sharp/Blunt	1
19	Scissors, Bandage, Lister, 20 cm, One larger ring handle	1
20	Clamp, Bone-Holding, Verbrugge, 24 cm, prehensile	1
21	Forceps, Bone-Holding, Lewin, 17.8 cm, Serrated	1
22	Forceps, Bone-Cutting, Liston, 19.7 cm, Angled	1
23	Forceps, Bone-Cutting, Stille-Liston, 26.7 cm, Straight	1
24	Rongeur, Bone, Luer, 15 cm x 5 mm, Curved	1
25	Rongeur, Bone, Stille-Luer, 23 cm x 10 mm, Double-acting	1
26	Curette, Bone, Volkman, 17 cm x 3.6 mm	1
27	Curette, Bone, Volkman, 17 cm x 6.8 mm	1
28	Curette, Bone, Volkman, 17 cm x 8.5 mm	1
29	Curette, Bone, Volkman, 17 cm x 13.0 mm	1
30	Raspatory, Bone, Lambotte, 21 cm Sharp, 10 mm Curved	1
31	Raspatory, Bone, Lambotte, 21 cm Sharp, 20 mm Curved	1
32	Retractor, Bone, Hohmann, 24 cm, 18 mm	1
33	Retractor, Bone, Verbrugge-Hohmann, 25 cm, 65 mm	1
34	Forceps, Towel-Clamp, Backhaus, 9 cm	4
35	Forceps, Towel-Clamp, Backhaus, 13 cm	12
36	Kidney Dish, Medium, 250 x 140 x 40 mm, Stainless Steel	2
37	Gallipot, Round, 100 ml, 80 x 35 mm, Stainless Steel	2
38		
39		
40		
41		
42		
43		
44		
45		
46		
47		
48		
49		
50		
51		
52		
53		
54		
55		
56		
57		
58		
59		
60		

Towel Clips	4
Towel Clips, Non-perforating	2
Tourniquet for limb surgery with adult cuff 66 x 11cm (26 x 4")	1
<b>EXTERNAL FIXATION SUPPLEMENT</b>	
Brace, External fixator	1
Connectors, Ø 15/15 mm	12
Pin, self-drilling, Ø 4 mm, 100 mm	6
Pin, self-drilling, Ø 5 mm, 110 mm	6
Pin, self-drilling, Ø 5 mm, 140 mm	6
Pin, self-drilling, Ø 5 mm, 170 mm	6
Pin, self-drilling, Ø 5 mm, 210 mm	6
Pin, transfixing, Ø 5 mm, 250 mm	1
Sleeve, Reduction, 15/4 mm	6
Sleeve, Reduction, 15/5 mm	12
Sleeve, Reduction, 15/9 mm	2
Handle for Trocar/sleeve	1
Trocar/Sleeve without handle, 70 mm, Ø 5 mm	1
Tube, Carbon-Fibre, 100 mm, Ø 15 mm	2
Tube, Carbon-Fibre, 125 mm, Ø 15 mm	2
Tube, Carbon-Fibre, 150 mm, Ø 15 mm	2
Tube, Carbon-Fibre, 200 mm, Ø 15 mm	2
Tube, Carbon-Fibre, 250 mm, Ø 15 mm	2
<b>BONE PINNING &amp; WIRING SUPPLEMENT</b>	
Drill, Hand, Manual, with 3-jaw Chuck	1
Pliers, Wire-Cutting, 21 cm, Front-acting	1
Pliers, Flat Nose, 17 cm, Serrated jaws, 17 cm	1
Bone Suturing Wire, Soft, Coil 10 m, 0.5 mm	1
Bone Suturing Wire, Soft, Coil 10 m, 2.0 mm	1
Wire, Kirschner, 23 cm, Trocar point Ø 1.8 mm	10
Wire, Kirschner, 23 cm, Trocar point Ø 2.5 mm	20

**Table 5:** Surgical Instruments for Minor Procedures at First Level Hospital

ITEM	QUANTITY
<b>MINOR ABDOMINAL SURGERY</b>	
SCALPEL HANDLE for blades 10 -15 mm	1
SCALPEL HANDLE for blades 18 - 36 mm	1
SCISSORS straight 145 mm	1
DISSECTING SCISSORS curved 140 mm	1
ABDOMINAL SCISSORS heavy design, curved 160 mm	1
DISSECTING SCISSORS curved 230 mm	1
DISSECTING SCISSORS Curved, blunt 115 mm	1
TENDON SCISSORS curved, sharp 110 mm	1
DISSECTING FORCEPS straight, toothless 145 mm	2
DISSECTING FORCEPS 140 mm	1
TISSUE FORCEPS 1X2 teeth 145 mm	2
INTESTINAL GRASPING FORCEPS 160 mm	2
ATRAUMATIC GRASPING FORCEPS 7 mm 200 mm	2
DRESSING FORCEPS straight 220 mm	1
TOWEL CLAMP curved 85 mm	4
TOWEL CLAMP straight 160 mm	4
HOLDING RING HANDLE INSTRUM 105 mm	2
ARTERY FORCEPS straight, 1X2 teeth 145 mm	2
MOSQUITO FORCEPS curved 140 mm	2
NEEDLE CASE perforated 50X25X5 mm	1
NEEDLE HOLDER 125 mm	1
GUIDING NEEDLE char 16 190 mm	1
RETRACTOR 55 X 47 mm 205 mm	2
RETRACTOR medium, 55 mm deep 200X170 mm	1
RETRACTOR double ended, 10 mm set of 2, 125 mm	2
FISTULA HOOK 2 sprong 220 mm	2
UNIVERSAL TROCAR set of 4 tips 1 handle	1
STERILIZATION CONTAINER 300X274X247 Size ½ mm	1
ROUND BOWL 8.3 X 5.4 X 4.1 cm	1
<b>INCISION &amp; DRAINAGE</b>	
SCALPEL HANDLE for blades 10 -15 mm	1
SCALPEL HANDLE for blades 18 - 36 135 mm	1

1		
2		
3	SCISSORS straight 145 mm	1
4	DISSECTING SCISSORS curved 140 mm	1
5	DISSECTING SCISSORS Curved, blunt 115 mm	1
6	TISSUE FORCEPS 1X2 teeth 115 mm	1
7	TOWEL CLAMP curved 85 mm	4
8	MOSQUITO FORCEPS curved 140 mm	2
9	NEEDLE HOLDER 125 mm	1
10	STERILIZATION CONTAINER 250 x 110 x 40 mm	1
11	<b>DILATION &amp; CURETTAGE</b>	
12	DISSECTING FORCEPS straight, toothless 115 mm	2
13	TISSUE FORCEPS 1X2 teeth 145 mm	2
14	DRESSING FORCEPS straight 220 mm	1
15	TOWEL CLAMP curved 85 mm	6
16	STERILIZATION CONTAINER 470X274X135 Size 3/4	1
17	ROUND BOWL 8.3X5.4X4.1 cm	1
18	UTERINE DRESSING FORCEPS 260 mm	1
19	VAGINAL SPECULA 105/95X35 mm	1
20	VAGINAL SPECULA 60X35mm 240 mm	1
21	WEIGHTED VAGINAL SPECULA with detachable weight	1
22	UTERINE PROBE malleable, curved diameter: 4 mm	1
23	SET WITH 14 DILATORS IN METAL CASE	1
24	UTERINE CURETTE 7.5 mm, blunt, rigid max 255 mm	1
25	UTERINE CURETTE 12 mm, blunt, rigid max 255 mm	1
26	UTERINE TENACULUM FORCEPS 175 mm	1
27		
28		
29		
30		
31		
32		
33		
34		
35		
36		
37		
38		
39		
40		
41		
42		
43		
44		
45		
46		
47		
48		
49		
50		
51		
52		
53		
54		
55		
56		
57		
58		
59		
60		

**Table 6: Anesthesia Supplies at First Level Hospital**

<b>ANESTHESIA CONSUMABLES/SINGLE-USE</b>	
Airway, Guedel, Size 0, Pediatric, Re-usable	
Airway, Guedel, Size 00, Neonate, Re-usable	
Airway, Guedel, Size 01, Infant, Re-usable	
Airway, Guedel, Size 02, Toddler, Re-usable	
Airway, Guedel, Size 03, Adolescent, Re-usable	
Airway, Guedel, Size 04, Adult, Re-usable	
Airway, Guedel, Size 05, Large Adult, Re-usable	
Mask, Anesthesia, with Rim + Hook, Size 2: Child	
Mask, Anesthesia, with Rim + Hook, Size 4: Adolescent	
Mask, Anesthesia, with Rim + Hook, Size 5: Adult	
Mask, Anesthesia, Size 1: Neonate	
Self-inflating bag (Ambu), Adolescent/Child + Masks	
Self-inflating bag (Ambu), Child/Neonate + Masks	
Catheter Mount, 150 mm	
Laryngeal Mask Airway, Size 2, 3, 4, 5 Re-usable	
Bag, Urine, 2 L, Drainable + Non-return Valves, Sterile	
Filter, Breathing Circuit, Neonate	
Filter, Breathing Circuit, Adult	
Catheter, Urinary, Foley, Balloon, CH 10, 12, 16 Sterile	
Tube, Endotracheal, Cuffed, Balloon, Ø 5.5 mm, 6.0 mm, 6.5 mm, 7.0 mm, 7.5 mm, 8.0 mm, 8.5 mm,	
Tube, Endotracheal, Uncuffed, Balloon, Ø 3.0 mm, 4.0 mm, 5.0 mm	
Tube, Oxygen, Conical Tip, 40 cm, CH 10	
Tube, Suction, Conical Tip, 50 cm, CH 10, CH 12, CH 14, CH 16, Single use	
Gauze, Compression, 10 cm, 12 plies, 17 threads, sterile	
Tape, Adhesive, Roll, 2 cm	
IV Cannula, injection port, s.u. 16 G (1.7 x 55 mm), Grey	
IV Cannula, injection port, s.u. 18 G (1.2 x 45 mm), Green	
IV Cannula, injection port, s.u. 20 G (1.0 x 32 mm), Pink	
IV Cannula, injection port, s.u. 22 G (0.8 x 25 mm), Blue	
Needle, 19 G (1.1 x 40 mm), Cream, IV	
Needle, 21 G (0.8 x 40 mm), Green, IM	
Needle, Intra-Osseous, 16G	
Needle, Intra-Osseous, 18G	
Needle, 22 G (0.7 x 90 mm), Blue	

1	
2	
3	Needle, 25 G (0.5 x 90 mm), Orange
4	Set, Infusion, 'Y', Luer lock with air inlet
5	Set, Infusion, Pediatric
6	Syringe, 5 ml, 10 ml, 20 ml
7	Bougie, Gum Elastic, 50 cm, CH 5, CH 8, CH 14 Sterile
8	
9	
10	
11	<b>INTRAOPERATIVE EQUIPMENT</b>
12	Handle for Laryngoscope
13	Laryngoscope, with Bulb, Fiberoptic
14	Laryngoscope, Macintosh blades, Adult
15	Laryngoscope, Macintosh blades, Pediatric
16	Spare bulb for Laryngoscope (2.7V)
17	Stylet for Intubation
18	Spare batteries for Laryngoscope, Dry cell, Alkaline, 1.5V, C(LR14)
19	Self-inflating Bag (Ambu), Adult/Child + Masks
20	Self-inflating Bag (Ambu), Child/Neonate + Masks
21	Valve for Self-inflating bag Ambu, Adult, 1 inlet, Complete
22	Valve for Self-inflating bag Ambu, Pediatric, Complete
23	Tourniquet, Pneumatic, Arm Cuff + Manometer + Leg Cuff
24	Concentrator, Oxygen, 10 L, 230V, 50 Hz
25	Pump, Suction, Electrical, 100-240 V
26	Pump, Suction, Mechanical (Twin Pump) + Collection bottles
27	Bandage, 6 cm x 5 m
28	Pressure Cuff, for pouch 500/1000 ml
29	Sphygmomanometer, One-hand manometer, Adult, Velcro
30	Sphygmomanometer, One-hand manometer, Pediatric, Velcro
31	Stethoscope, Double Cup, Clinician
32	Tourniquet, 100 x 1.8 cm, Elastic
33	Forceps, Hemostatic, Pean, 14 cm, Straight
34	Scissors, Dressing, 14.5 cm, Straight, Blunt/Blunt
35	Forceps, Magill, 24 cm, Adult
36	Forceps, Magill, 15 cm, Child
37	Battery, Dry cell, Alkaline, 1.5 V, R14 (C)
38	Connector, Biconical, Symmetrical, Ext. Ø 7-11 mm, Autoclavable
39	Oxygen Mask, Adult size
40	Oxygen Mask, Pediatric size
41	
42	
43	
44	
45	
46	
47	
48	
49	
50	
51	
52	
53	
54	
55	
56	
57	
58	
59	
60	

1	
2	
3	Tube, Silicone, Internal Ø 5 mm, 25 m, Autoclavable
4	Tube, Silicone, Internal Ø 8 mm, 10 m, Autoclavable
5	
6	Thermometer, Electronic, Accuracy 0.1° C
7	
8	Anesthesia Ventilator + Scavenger + Accessories
9	Capnograph/Pulse Oximeter, Portable
10	
11	Monitor, Multiparameter + Accessories
12	Portable Anesthesia System + Scavenger
13	
14	<b>RECOVERY ROOM</b>
15	Airway, Guedel, Size 0, Paediatric, Re-usable
16	Airway, Guedel, Size 00, Neonate, Re-usable
17	Airway, Guedel, Size 01, Infant, Re-usable
18	Airway, Guedel, Size 02, Toddler, Re-usable
19	Airway, Guedel, Size 03, Adolescent, Re-usable
20	Airway, Guedel, Size 04, Adult, Re-usable
21	Airway, Guedel, Size 05, Large Adult, Re-usable
22	
23	Mask, Anesthesia, with Rim + Hook, Size 2: Child
24	Mask, Anesthesia, with Rim + Hook, Size 4: Adolescent
25	Mask, Anesthesia, with Rim + Hook, Size 5: Adult
26	
27	Self-inflating Bag (Ambu), Adult/Child + Masks
28	Self-inflating Bag (Ambu), Child/Neonate + Masks
29	
30	Concentrator, Oxygen, 10 L, 230V, 50 Hz
31	
32	Oxygen Mask, Adult size
33	Oxygen Mask, Paediatric size
34	
35	Oximeter, Pulse + Accessories
36	
37	Sphygmomanometer, One-hand manometer, Adult, Velcro
38	Sphygmomanometer, One-hand manometer, Pediatric, Velcro
39	
40	Stethoscope, Double Cup, Clinician
41	
42	Tourniquet, 100 x 1.8 cm, Elastic
43	
44	Torch, Pen Light, 2 x R6 batteries
45	
46	Pump, Suction, Electrical, 100-240 V
47	
48	Pump, Suction, Mechanical (Twin Pump) + Collection bottles
49	
50	
51	
52	
53	
54	
55	
56	
57	
58	
59	
60	

**Table 7:** Pediatric specific surgical supplies for intermediate surgical care (essential surgical package for first-level hospital)

Equipment	
Operating Room Positioning Supports for Children (Soft pads, Jelly pads, Pediatric OR straps)	
Temperature control- Bair Hugger; Radiant Warmer (Panda, Giraffe, Bassinet); Cotton Wool (Large Rolls)	
Flat weighing scales (infant)	
Post-operative crib with over-head safety constraints	
Ultrasound with Echocardiography capability	
Sphygmomanometers with pediatric cuffs (neonatal, small, medium, large)	
Otoscope and Screening audiometer	
Rhinoscope	
Ophthalmoscope	
Consumables	
Central Venous Catheters (Pediatric Sizes- 3- 7)	
Urethral Catheters (Pediatric Sizes- 6F, 8F, 10F, 12F)	
Nasogastric and Feeding Tubes (Pediatric Sizes- 6F, 8F, 10F, 12F)	
Plastic bags for eviscerated intestines or Preformed Silo	
Peritoneal dialysis catheters (Pediatric Sizes- 8F-10F)	
Chest tubes (Pediatric Sizes- 10F- 20F)	
Ventricular Drains (Pediatric Sizes)	
Cling film for bowel coverage	
Instruments (Major Procedures)	
No additional instruments necessary	
Instruments (Minor Procedures)	
Item	Quantity
Pediatric Cut Down Set	
Forceps, Artery, Halsted-Mosquito Curved 4 <sup>3/4</sup> "	3
Forceps, Artery, Halsted-Mosquito Straight 4 <sup>3/4</sup> "	3
Forceps, dissecting, Adson straight, 1/2 Teeth 4 <sup>3/4</sup> "	1
Forceps, dissecting, Adson straight, Non-Teeth 4 <sup>3/4</sup> "	1
Forceps, sponge holding 7"	1
Skin hook, Gillies, sharp	2
Skin hook, Blunt	1
Vessel hook 4 <sup>3/4</sup> "	2
Bard Parker handle for Surgical Blade No. 3	1
Needle holder, Mayo 5 <sup>3/4</sup> "	1

Scissors, ligature, Spencer 3.5"	1
Iris scissors Curved 3.5" & 4.5"	1
Iris scissors Straight 3.5"	1
Gallipot 30mls	1
Kidney dish 100mls	1
Anesthesia Supplies	
Tube, Endotracheal, Uncuffed, Balloon, Ø 2.5 mm for neonates and small children	
Tube, Endotracheal, Uncuffed, Balloon, Ø 3.0 mm for small children	
Laryngoscope, Miller blades, Neonates, Size 00 and 0	
Ambu Bags (bag valve mask, neonatal sizes)	

### 3. Supplies, Equipment and Infrastructure for Advanced and Complex Care

**Table 8:** Overview of essential childhood surgical supplies for referral hospitals by specialty (essential surgical package for advanced/complex surgical care)

Subspecialty	Pediatric Specific Supplies for Advanced Surgical Care	Pediatric Specific Supplies for Complex Surgical Care
<b>Anesthesia</b>	Multi-parameter intra-operative monitor	Automated Anesthesia Machine, Continuous and invasive intra-operative monitor
<b>Cardiac Surgery</b>	Same Supplies for First Level Hospital	Cardiopulmonary bypass machine, Catheterization laboratory with pediatric-sized arterial and venous catheters
<b>Critical Care</b>	Equipment for Neonatal Intensive Care Unit and Pediatric Intensive Care Unit, Equipment for full hematologic lab, Arterial Blood Gas Machine	Equipment for Burn Unit and Dialysis Unit
<b>General Surgery</b>	Infant Laparotomy Set, Pediatric Thoracotomy Set	Laparoscopic and Thoracoscopic Equipment, Endoscope, Colonoscope
<b>Neurosurgery</b>	Craniotomy instrument set	Intra-operative neurophysiology monitoring, Neuro-navigational system, Surgical microscope, Neuro-endoscopy
<b>Ophthalmology</b>	Slit lamp and lens, Direct ophthalmoscope	Optic Laser for ROP, Cataract surgical set, Orthoptic and Optometric equipment
<b>Oral Maxillofacial</b>	Dental laboratory casting machine	Dental surgical instrument set
<b>Orthopedics Surgery</b>	Pediatric Orthopedic Set (extra fine and fine), Antibiotic loaded cement, Prostheses, Surgical Drill and Saw, Functioning Arthroscope, Fracture Fixation Equipment	K-wire Kit, Broken Screw removal instrument kit, Spine instrument set, Hip & Pelvic osteotomy set, Limb reconstruction system, microdrill system
<b>Otolaryngology</b>	Same Supplies for First Level Hospital	Operative Microscope, Airway stents, Airway balloon dilators, Bronchoscopy, nerve monitoring system, FESS instruments
<b>Plastic Surgery</b>	Skin-graft mesher, Plastic surgery instrument set, Dermatome Knife	Vascular Doppler Machine, Electric dermatome, Operating microscope, Microsurgery instrument set
<b>Radiology</b>	Magnetic Resonance Imaging, Fluoroscopy Unit	Same as Advanced Surgical Care
<b>Urology</b>	Pediatric Urology Instrument Set	Urethroscope, Cystoscope, Ureteroscopy, Lithotripsy machine, dialysis catheters

**Table 9:** Select pediatric specific surgical supplies for advanced and complex surgical care (essential surgical package for second-level, third-level and children's national hospital)

Equipment
Orthopedic Protheses
Thompson prosthesis
Talwalkar nail
Swanson prosthesis
Steinmann pin
Seidel nail
Smith Peterson nail with McLaughlin's plate
Smith Peterson nail
Rush nail
Moore's pin
Kuntscher nail
Kirschner wire
Grosse-Kempf nail
Ender's nail
Condylar blade plate
Austin-Moore
Hegar's Dilator Set (full range of 8 dilators, double ended)
3mm/4mm
5mm/6mm
7mm/8mm
9mm/10mm
11mm/12mm
13mm/14mm
15mm/16mm
17mm/18mm
Radiology
X-ray Machine (fixed, conventional)
X-ray Machine (mobile, conventional)
X-ray Machine with Fluoroscopy
X-ray Machine (Low dose/Lodox®)
Ultrasound Machine (conventional)
Ultrasound Machine (Vascular Doppler/Duplex)

1		
2		
3	Computed Tomography Scanner	
4	Magnetic Resonance Imaging Machine	
5		
6	Equipment Necessary for Additional Specialty Units	
7		
8	Neonatal Intensive Care Unit	
9	Pediatric Intensive Care Unit	
10		
11	Burn Unit	
12	Dialysis Unit	
13		
14	Catheterization laboratory & Cardiopulmonary bypass machine	
15	Endoscopic Suit (Upper, colonoscope, Urethroscope, etc.)	
16		
17	Consumables	
18		
19	Ventricular Drains	
20	Instruments (Major Procedures)	
21		
22	Item	Quantity
23	Infant Laparotomy Instrument Set	
24	Bard Parker Scalpel Handle No. 3	2
25	Op Sciss Str Sh/bl 5 1/2"	1
26	Halsted Mosq Fcp Str 5"	6
27	Halsted Mosq Fcp Cvd 5"	6
28	Crile Fcp Str 5 1/2"	4
29	Crile Fcp Cvd 5 1/2"	3
30	Kocher Fcp Str Del 1x2 5 1/2"	3
31	Kocher Fcp Cvd Del 1x2 5 1/2"	3
32	Metz Lahey Del Crv Sciss 5 3/4	1
33	Metzenbaum Sciss Del Cvd 7"	1
34	Roch Pean Str Del 5 1/2"	3
35	Roch Pean Cvd Del 5 1/2"	3
36	Semkin Dress Fcp Nar Serr 5"	1
37	Semkin Tiss Fcp Nar 1x2 5"	1
38	Babcock Fcp 5 1/2"	3
39	Allis Tiss Fcp Del 4x5 6"	3
40	Yankauer Suct Tube Ped	1
41	Miller Senn Retr Shrp D/e 6"	1
42	Miller Senn Retr Blnt D/e 6"	1
43	Balfour Retr pediatric 1 1/4"	1
44	Adson Tiss Fcp 1x2 Del 4 3/4"	1
45	Adson Dress Fcp Serr Del 4 3/4	1
46		
47		
48		
49		
50		
51		
52		
53		
54		
55		
56		
57		
58		
59		
60		

Parker De Retr Set/2 5/8x5	1
Zalkind Ribbon Retr 1/2"x7.75"	1
Zalkind Ribbon Retr 3/4"x7.75"	1
Ballengr Spng Fcps Str Serr 7"	2
Doyen-Baby Intestinal Clamps/Forceps (Straight, Non-crushing/attraumatic) 5"	2
Doyen-Baby Intestinal Clamps/Forceps (Curved, Non-crushing/attraumatic) 5"	2
Payr Intestinal Clamp 5-3/4"	2
Benson Pylorus Separator (Spreader) 5-3/4"	1
Autoclavable Instrument Tray/Box	1
Pediatric Urology Instrument Set	
Hartmann Mosquito Straight Delicate Ser 3 -1/2"	3
Hartmann Mosquito Curved Delicate Ser 3 -1/2"	3
Halsted Mosquito Forceps Straight 5"	6
Halsted Mosquito Forceps Curved 5"	6
Bard Parker Handle No. 3	2
Mayo-Hegar NH 7"	1
Crile-Murray NH Serrated 6" T-C Grip	1
Ryder Micro NH Serrated 6" T-C	1
Slender Tip Scissors Curved 5 -1/2"	1
Ragnell Dissecting Scissors Curved 5"	1
Semkin Dressing Forceps Nar 5"	2
Semkin Tissue Forceps Narrow 1x2 5"	2
Miller-Senn Retractor Sh D/E 6"	1
Miller-Senn Retractor Bl D/E 6"	1
Aim Retractor 2-3/4"	1
Operating Scissors S/BSt 5-1/2"	2
Strabismus Scissors Straight 4 -1/2"	1
Mayo Dissecting Scissors Straight 5 -1/2"	1
Mayo Dissecting Scissors Curved 6 -3/4"	1
Scissors Blunt Curved 5 -3/4"	1
Crile Forceps Straight 5 -1/2"	12
Crile Forceps Curved 5-1/2"	12
Wagner Scissors Straight S/S 4 -3/4	2
Wagner Scissors Curved S/B 4 -3/4"	1
Brown Adson Tissue Forceps 4-3/4"	2
Jansen Retractor Sh 3x3 4"	1

1		
2		
3	Adson Forceps Straight 7 -1/4"	3
4	Adson Forceps Curved 7 -1/4"	3
5	Frazier Suction Tube 12 Fr	2
6	Adson Tissue Forceps 1x2 4 -3/4"	1
7	Adson Dressing Forceps Serrated 4 -3/4"	1
8	Gemini Clamp Delicate Ang 5 -1/2"	2
9	Crile Retractor D/E Small	2
10	Zalkind Ribbon Retractor - 3/4" x 7"	2
11	Autoclavable Instrument Box/Tray with Cover	1
12	Pediatric Thoracotomy Set	
13	Weinstein Instrument Holder 10	2
14	Halstead Mosq Fcps Cvd Satin 5"	2
15	Crile Fcps Cvd Satin 5.5"	4
16	Crile Fcps Str Satin 5.5"	2
17	R-Pean Fcps Cvd Satin 4.5"	2
18	R-Pean Fcps Cvd Satin 5.5"	2
19	R-Pean Fcps Cvd Satin 6 1/4"	2
20	Allis Tiss Fcps 4x5 teeth 5 1/4"	2
21	Allis Tiss Fcps 5x6 teeth 6 1/4"	2
22	Allis Tiss Fcps 5x6 teeth 7"	2
23	Babcock Tiss Fcps Atraumatic 6"	2
24	Babcock Tiss Fcps Atraumatic 7"	2
25	Ochsner Fcps Str Satin 6 1/4"	2
26	Schmidt Fcps Closed Ring 7 1/2"	2
27	Bridge Fcps Cvd Paediatric	2
28	Mixer Fcps Baby 5 1/4"	2
29	Mixer-O'Shaughnessy Fcps Horiz Serr 7 1/4"	2
30	TC Masson Needle holder 10 1/2 "	1
31	TC Mayo-Hegar Needle holder 5 3/4"	1
32	TC Mayo-Hegar Needle holder 6 1/4"	1
33	TC Mayo-Hegar Needle holder 7 1/4"	1
34	TC Mayo Scs Cvd 4 1/2"	1
35	TC Metz Scs Cvd 6 3/4"	1
36	TC Metz Scs Cvd 7 1/8"	1
37	TC Metz Scs Cvd 8 1/8"	1
38	TC Mayo Scs Str 6 3/4"	1
39		
40		
41		
42		
43		
44		
45		
46		
47		
48		
49		
50		
51		
52		
53		
54		
55		
56		
57		
58		
59		
60		

1		
2		
3	Debakey Tangent Vascular Clmp Pediatric 3 <sup>7/8</sup> "	2
4	Debakey Tangent Vascular Clmp Pediatric 4 <sup>7/8</sup> "	2
5	Debakey Tangent Vascular Clmp Pediatric 5 <sup>7/8</sup> "	2
6		
7	Duval Lung Fcps 7 <sup>3/4</sup> "	2
8		
9	Foster Sponge Fcp Str Satin 7 <sup>1/8</sup> "	1
10		
11	Edna Towel Fcps 3.5"	2
12	Backhaus Fcps Satin 4 <sup>3/8</sup> "	2
13		
14	Bard Parker Handle for Surgical Blade 3	2
15	Dressing Fcps Serrated 4.5"	2
16	Dressing Fcps Serrated 5 <sup>3/4</sup> "	2
17	Dressing Fcps Serrated 6 <sup>7/8</sup> "	2
18		
19	Tissue Fcps 1x2 teeth 4.5"	2
20	Tissue Fcps 1x2 teeth 6 <sup>7/8</sup> "	2
21		
22	Debakey Vasc Tiss Fcps 6 <sup>1/4</sup> "	2
23	Debakey Vasc Tiss Fcps 7 <sup>7/8</sup> "	2
24		
25	Potts-Smith Dress Fcps 7 <sup>1/8</sup> "	2
26	Russian Tiss Fcps 6"	2
27		
28	Adson Tiss Fcps 1x2 teeth 4 <sup>3/4</sup> "	2
29		
30	Us Army-Navy Retr Set/2 6"	1 (set)
31	Cushing Vein Retr 7"	1
32		
33	Richardson Retr Loop-Handle Small	2
34	Richardson Retr Loop-Handle Medium	2
35	Kelly Retr Loop-Handle Small/Medium	2
36		
37	Balfour-Baby Abd Retr 4 <sup>7/8</sup> "	1
38	Finochietto Rib Retr Infant	1
39	Finochietto Rib Retr Baby	1
40		
41	Finochietto Rib Retr 135 mm spread	1
42		
43	Bailey-Baby Rib Contractor 6 <sup>1/8</sup> "	1
44	Bailey Rib Contractor (short claws) 7 <sup>7/8</sup> "	1
45	Bailey Rib Contractor (long claws) 7 <sup>7/8</sup> "	1
46		
47	Killingbeck/Allison Lung Retr Infant	1
48	Killingbeck/Allison Lung Retr Pediatric	1
49		
50	Autoclavable Instrument Tray/Box with cover	1
51	Additional Surgical Instrument Sets and Kits	
52	Laparoscopic and Thoracoscopic Equipment	
53	Pediatric Plastic Surgery Set	
54		
55		
56		
57		
58		
59		
60		

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Functioning Arthroscope with related equipment	
K-wire Kit, Broken Screw Removal Instrument Kit	
Spine Instrument Set	
Hip/Pelvic Osteotomy Set	
Craniotomy Instrument Set	
Dental surgical instrument set	
Instruments (Minor Procedures)	
Instrument	Quantity
No additional Minor Procedure Sets Necessary	
Anesthesia	
Multi-parameter intra-operative monitor	
Automated Anesthesia Machine, Continuous and invasive intra-operative monitor	

For Peer Review

**REFERENCES**

1. Personal communication R. Gruen, Nanyang Technological University Singapore, easiOR Project (Essential Affordable Smart Infrastructure – Operating Room), December 2017.
2. Guide to infrastructure and supplies at various levels of health care facilities. Emergency and Essential Surgical Care (EESC). Compiled from the WHO Manual Surgical Care at the District Hospital 2003. Available at: <http://www.who.int/surgery/publications/s15983epdf?ua=1> (Accessed 8/20/18).

For Peer Review