Paper 1:

Qualitative research as evidence – expanding the paradigm for Evidence-Based Health Care

Veronika Williams, (corresponding author)

Nuffield Department of Primary Care Health Sciences, University of Oxford, Radcliffe Observatory Quarter, Woodstock Road, Oxford, OX2 6GG

veronika.williams@phc.ox.ac.uk

Anne-Marie Boylan

Nuffield Department of Primary Care Health Sciences, University of Oxford, Radcliffe Observatory Quarter, Woodstock Road, Oxford, OX2 6GG

David Nunan

Nuffield Department of Primary Care Health Sciences, University of Oxford, Radcliffe Observatory Quarter, Woodstock Road, Oxford, OX2 6GG

Keywords: Qualitative health research, critical appraisal, patient experience

Competing interests: Drs Williams, Boylan and Nunan have no competing interests to disclose.

Contributorship: All authors contributed to the conception of this article; VW wrote the first draft of the article with AMB and DN contributing and revising subsequent drafts. All authors approved the final submitted article.

Acknowledgements: We would like to thank Dr Kamal Mahtani for feedback on an earlier version of this paper.

Funding info: there are no funders to report for this submission

Ethical approval information: not applicable

Data sharing statement: not applicable

Abstract:

In this first of three articles in the EBM learning series exploring the role and nature of qualitative research and what it can add to evidence-based health care (EBHC), we discuss why qualitative research is still often on the 'margins' of EBHC. We outline the role qualitative research can play and ways in which it can contribute to improving patient care and the need to fully integrate qualitative research within EBHC. The second paper will focus on the ongoing tensions of appraising qualitative research as part of practising evidence-based health with the third paper offering a different approach to appraisal.

Writing in the BMJ, David Sackett and John Wennberg pointed out over 20 years ago we need to stop ‘squabbling over the best methods’ and focus on the research question, since it is the nature of the question that determines the best research design to answer it.[1] Yet, the majority of evidence hierarchies and ‘levels’ [2] still do not include qualitative methodologies, and EBHC still largely focuses on the positivist perspective of generating and applying knowledge in health care practice .[3] Despite its lack of prominence in EBHC, qualitative research in health care has increasingly been conducted and published since Sackett and Wennberg’s editorial; and the publication of qualitative evidence syntheses have doubled in the past decade. [4]

The argument for qualitative research within EBHC has been made for over 20 years, [5] including a series of educational articles in the BMJ [6-9] and all of which have citations in the 1000s. [NB the journal has since reversed their position on qualitative research to one where qualitative research is no longer a priority for the journal, to the dismay of prominent proponents. [10]

To understand why qualitative research has yet to achieve the same status as other forms of evidence within the evidence based health care movement, we need to consider wider contextual factors surrounding research and the underlying epistemological differences between the quantitative and qualitative paradigms.

Qualitative research can be of exploratory and /or explanatory nature. It seeks to answer ‘how’ and ‘why’ questions rather than trying to establish causation or correlation using statistical methods and aims to explore different perspectives and experiences of phenomena. [11] It falls within an interpretivist/ constructivist paradigm¹, which cannot, and indeed does not aim to make definitive statements about its research findings. [11] The underlying premise of methodologies that use qualitative methods does not assume *one* truth that can be verified and replicated, but rather several truths which are equally valid. Thus, research within this paradigm does not propose hypotheses *a priori*² and is therefore less likely to produce ‘headline grabbing’ results than findings from other study types (observational cohort studies for example).

Arguably, health care and health care systems as well as populations are becoming increasingly complex. Growing emphasis is placed on exploring and understanding the patient experience of healthcare services and ill-health as a way to provide health care that is more ‘patient-centred’ both on a policy [12] and research level. [13] Equally, people are living longer but with more chronic conditions than ever before and there is also a clear need to consider the increasing cultural and ethnic complexity of patient populations. Seminal work by qualitative researchers has provided a much better and in-depth understanding of the patient experience of long-term conditions and thus provided the evidence to improve health services for this patient population [14-16]. Advancements in medical treatment and technology result in more complex procedures and interventions, and polypharmacy is on the rise. Quantitative data alone are unlikely to provide the insight needed to tackle these challenges. Simply knowing ‘what works’ is not enough. For complex interventions in particular, we need to know not only what works, but *why* it may (or may not) work, and how significant research findings can be translated into front-line health care and health service delivery. Therefore, traditional RCTs on their own may not be the best way to assess ‘efficacy’ and implementation [17].

¹ Interpretivism assumes that the world does not exist independently of our knowledge of it. Knowledge and meaningful reality are constructed in and out of interaction between humans and their world and are developed and transmitted in a social context

² Although some methodologies such as Grounded Theory aim to develop hypothetical statements as one of its outcomes

Complex interventions and implementation research rely on qualitative alongside quantitative methods.[17, 18] Using a mixed method approach in health services research allows us to understand how best to place a complex intervention within its current clinical context, and to identify the opportunities and barriers to changing clinical care. [19] Yet we also know that qualitative studies within RCTs remain uncommon and are often poorly integrated with the wider trial findings. [17] Demonstrating the importance of improving this, frameworks such as the [MRC guidance](#) for conducting and reporting process evaluation studies have since been developed to advise researchers how to integrate qualitative methodology within RCTs of complex interventions. [20]

Qualitative methods have long played an important role in research into quality and safety in health care – to understand how medical errors occur, to consider how to minimise them and to identify ways to improve quality of care and support better teamwork. Lingard et al's [21] study on identifying communication errors and challenges in teamwork in operating theatres that may lead to medical errors is one of the most cited (over 1000) qualitative studies in health care research. Their ethnography of general and vascular surgery over a three-month period revealed different types of communication errors. This led to the development of a pre-operative communication checklist [21, 22] which has led to significant reductions in communication errors amongst surgical teams. [23] The initial data from the ethnographic study provided the crucial evidence on which the checklist was based. Another area within health services research, where qualitative research methodologies have contributed important knowledge and evidence in recent years, is the implementation and adoption (or lack thereof) of technological innovations, particularly around digital health interventions. [24-28]

There is undoubtedly a place for qualitative research within EBHC and a sound argument for why it should be included in hierarchies of evidence. However, given its different epistemological perspective to positivist, quantitative research, and the fact that it is not taught to the same extent (if at all) in medical schools (since the prevailing epistemological stance of medicine is that of positivism), it is perhaps unsurprising that EBHC practitioners may find it difficult to understand the contribution it can make to clinical care. However, as with any research we need to be able to distinguish poor research from high quality research in order to judge its relevance and appropriateness for health care services. Increasingly, research collaborations between different disciplines and methodologists are bridging some of this gap. Yet to ensure that healthcare practitioners have sound knowledge of several research designs across the epistemological divide, we may need to further incorporate interdisciplinary training across health care professions (nursing research for example tends to be situated predominantly within a constructivist/ interpretivist paradigm).

We argue that it is important that EBHC practitioners develop the knowledge and skills to appraise and review the methods and methodologies of qualitative research. However, as we explain in our next paper, appraising qualitative research is a complex endeavour. We set out the rationale for why we need to appraise qualitative health research, and introduce the current debates and challenges of using existing frameworks and tools for its appraisal.

References

1. Sackett, D.L. and J.E. Wennberg, *Choosing the best research design for each question*. BMJ: British Medical Journal, 1997. **315**(7123): p. 1636.
2. Sackett, D.L., et al., *Evidence based medicine: what it is and what it isn't*. 1996, British Medical Journal Publishing Group.
3. Goldenberg, M.J., *On evidence and evidence-based medicine: lessons from the philosophy of science*. Social science & medicine, 2006. **62**(11): p. 2621-2632.
4. Hannes, K. and K. Macaitis, *A move to more systematic and transparent approaches in qualitative evidence synthesis: update on a review of published papers*. Qualitative Research, 2012. **12**(4): p. 402-442.
5. Green, J. and N. Britten, *Qualitative research and evidence based medicine*. BMJ, 1998. **316**(7139): p. 1230-1232.
6. Mays, N. and C. Pope, *Qualitative research: rigour and qualitative research*. BMJ, 1995. **311**(6997): p. 109-112.
7. Mays, N. and C. Pope, *Assessing quality in qualitative research*. BMJ: British Medical Journal, 2000. **320**(7226): p. 50.
8. Britten, N., *Qualitative research: qualitative interviews in medical research*. BMJ, 1995. **311**(6999): p. 251-253.
9. Kitzinger, J., *Qualitative research. Introducing focus groups*. BMJ: British medical journal, 1995. **311**(7000): p. 299.
10. Greenhalgh, T., et al., *An open letter to The BMJ editors on qualitative research*. Bmj, 2016. **352**(i563).
11. Crotty, M., *The foundations of social research: Meaning and perspective in the research process*. 1998: Sage.
12. Improvement, N., *NHS Patient Experience Improvement Framework*. 2018: London.
13. Coulter, A., et al., *Collecting data on patient experience is not enough: they must be used to improve care*. BMJ, 2014. **348**(mar26 1): p. g2225-g2225.
14. Bury, M., *Chronic illness as biographical disruption*. Sociology of health & illness, 1982. **4**(2): p. 167-182.
15. Charmaz, K., *'Discovering' chronic illness: Using grounded theory*. Social Science & Medicine, 1990. **30**(11): p. 1161-1172.
16. Charmaz, K., *Good days, bad days: The self in chronic illness and time*. 1991: Rutgers University Press.
17. Lewin, S., C. Glenton, and A.D. Oxman, *Use of qualitative methods alongside randomised controlled trials of complex healthcare interventions: methodological study*. BMJ, 2009. **339**.
18. Campbell, M., et al., *Framework for design and evaluation of complex interventions to improve health*. BMJ : British Medical Journal, 2000. **321**(7262): p. 694-696.
19. Campbell, N.C., et al., *Designing and evaluating complex interventions to improve health care*. BMJ, 2007. **334**(7591): p. 455-459.
20. Moore, G.F., et al., *Process evaluation of complex interventions: Medical Research Council guidance*. bmj, 2015. **350**: p. h1258.
21. Lingard, L., et al., *Communication failures in the operating room: an observational classification of recurrent types and effects*. BMJ Quality & Safety, 2004. **13**(5): p. 330-334.
22. Lingard, L., et al., *Getting teams to talk: development and pilot implementation of a checklist to promote interprofessional communication in the OR*. BMJ Quality & Safety, 2005. **14**(5): p. 340-346.
23. Lingard, L., et al., *Evaluation of a preoperative checklist and team briefing among surgeons, nurses, and anesthesiologists to reduce failures in communication*. Archives of surgery, 2008. **143**(1): p. 12-17.
24. Greenhalgh, T., et al., *Adoption, non-adoption, and abandonment of a personal electronic health record: case study of HealthSpace*. Bmj, 2010. **341**: p. c5814.

25. Greenhalgh, T., et al., *The organising vision for telehealth and telecare: discourse analysis*. BMJ Open, 2012. **2**(4).
26. Greenhalgh, T., et al., *Storylines of research in diffusion of innovation: a meta-narrative approach to systematic review*. Soc Sci Med, 2005. **61**: p. 417 - 430.
27. Lupton, D., *The digitally engaged patient: Self-monitoring and self-care in the digital health era*. Social Theory & Health, 2013. **11**(3): p. 256-270.
28. Morton, K., et al., *Using digital interventions for self-management of chronic physical health conditions: A meta-ethnography review of published studies*. Patient Education and Counseling, 2016.