

Molecular Classification ~~with Integration of~~ Immunophenotypes Improves Risk Assessment in Adult *BCR-ABL1* negative B-ALL

Elisabeth Paietta,^{1,*} Kathryn G. Roberts,^{2,*} Victoria Wang,³ Zhaohui Gu,² Georgina A. N. Buck,⁴ Deqing Pei,⁵ Cheng Cheng,⁵ Ross L. Levine,⁶ Omar Abdel-Wahab,⁶ Zhongshan Cheng,⁷ Gang Wu,⁷ Chunxu Qu,² Lei Shi,⁵ Stanley Pounds,⁵ Cheryl L. Willman,⁸ Richard Harvey,⁸ Janis Racevskis,¹ Jan Barinka,² Yanming Zhang,⁹ Gordon W. Dewald,^{10†} Rhett P. Ketterling,¹⁰ David Alejos,¹ Hillard M. Lazarus,¹¹ Selina M. Luger,¹² Letizia Foroni,¹³ Bela Patel,¹⁴ Adele K. Fielding,¹⁵ Ari Melnick,¹⁶ David I. Marks,¹⁷ Anthony V. Moorman,¹⁸ Peter H. Wiernik,¹⁹ Jacob M. Rowe,²⁰ Martin S. Tallman,²¹ Anthony H. Goldstone,²² Charles G. Mullighan,^{2,†} Mark R. Litzow^{23,†}

Affiliations:

¹Department of Oncology, Montefiore Medical Center, Bronx, NY; ²Department of Pathology, St. Jude Children's Research Hospital, Memphis, TN; ³Department of Data Science, Dana-Farber Cancer Institute, Boston, MA; ⁴Clinical Trial Service Unit, Oxford, UK; ⁵Department of Biostatistics, St. Jude Children's Research Hospital, Memphis, TN; ⁶Human Oncology and Pathogenesis Program and Leukemia Service, Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY; ⁷Centre for Applied Bioinformatics, St. Jude Children's Research Hospital, Memphis, TN; ⁸University of New Mexico Comprehensive Cancer Center, Albuquerque, NM; ⁹Department of Pathology, Memorial Sloan Kettering Cancer Center, New York, NY; ¹⁰Department of Laboratory Medicine and Pathology, Mayo Clinic, Rochester, MN; ¹¹University Hospitals Cleveland Medical Center, Case Western Reserve University, Cleveland, OH; ¹²Abramson Cancer Center, University of Pennsylvania, Philadelphia, PA; ¹³Imperial College, London, UK; ¹⁴Centre for Haemato-Oncology, Barts Cancer Institute, Queen Mary University

of London, London, UK; ¹⁵UCL Cancer Institute, London, UK; ¹⁶Division of Hematology and Medical Oncology, Department of Medicine, Weill Medical College of Cornell University, New York, NY; ¹⁷Bristol Haematology and Oncology Centre, Bristol, UK; ¹⁸Leukaemia Research Cytogenetics Group, Newcastle University Translational and Clinical Research Institute, Newcastle-upon-Tyne, UK; ¹⁹Cancer Research Foundation, Bronx, NY; ²⁰Hematology, Shaare Zedek Medical Center, Jerusalem, Israel; ²¹Leukemia Service, Memorial Sloan Kettering Cancer Center, Weill Cornell Medical College, New York, NY; ²²University College London Hospitals, London, UK; ²³Division of Hematology, Department of Internal Medicine, Mayo Clinic, Rochester, MN;

†deceased

*E.P and K.G.R. are joint first authors; †C.G.M. and M.R.L. are joint senior authors

Short Title: Genotyping improves outcome in B-lineage ALL

Correspondence:

Mark R. Litzow, Division of Hematology, Mayo Clinic, 200 First Street, SW, Rochester, MN 55905; phone: 507-284-2511; fax: 507-266-4972; e-mail: litzow.mark@mayo.edu and Charles G. Mullighan, Department of Pathology, St. Jude Children's Research Hospital, 262 Danny Thomas Pl, MS342, Memphis, TN 38105; phone: 901-595-3387; fax: 901-595-5947; e-mail: charles.mullighan@stjude.org;

Wordcounts: 248 for abstract; 3986 for text;

References: 70

Tables: 4, Figures: 4

Key Points:

- Genomic testing leads to improved risk assignment in the majority of adult patients with *BCR-ABL* negative B-lineage ALL.
- ~~Simultaneous analysis of leukemia genotypes and immunophenotypes refines outcome prediction.~~ Charles, we need a new second point-reviewer did not like it

Abstract

While genomic classification of pediatric B-lineage acute lymphoblastic leukemia (B-ALL) has improved risk assignment, similar advances in adults are lacking. The international UKALLXII/ECOG-ACRIN E2993 (NCT00002514) trial accrued 1229 *BCR-ABL1* negative adolescent/adult B-ALL patients (aged 14-65 years). While 93% of patients achieved remission, 41% relapsed at a median of 13 months (range 28 days to 12 years). Five-year overall survival (5yr-OS) was 42% (95% CI, 39, 44). Transcriptome sequencing (n=238), gene expression profiling by LDA (n=210), cytogenetics (n=197) and PCR for known fusions (n=274) enabled genomic subtyping of 282 E2993 patient samples (264 of 409 eligible for trial). Among patients in the outcome analysis (n=217), 29.5% had standard-risk genotypes (*DUX4*-rearranged [9.2%], *ETV6-RUNX1*-like [2.3%], *TCF3-PBX1* [6.9%], *PAX5* P80R [4.1%], high-hyperdiploid [6.9%]) with 5yr-OS 65-80%; 50.2% had high-risk genotypes (Ph-like [21.2%], *KMT2A-AFF1* [12%], low-hypodiploid/near-haploid [14.3%], *BCL2/MYC*-rearranged [2.8%]) with 5yr-OS of 0-27%; and 20.3% had intermediate-risk genotypes (*PAX5alt* [12.4%], *ZNF384*-like [5.1%], *MEF2D*-rearranged [2.8%]) with 5yr-OS of 33-45%. *IKZF1*-alterations occurred in 87% of Ph-like cases and *TP53*-mutations occurred in low-hypodiploid (54%) and *BCL2/MYC*-rearranged patients (33%). Of patients considered high-risk for relapse based on presenting age and WBC count, 40% harbored subtype-defining genetic alterations associated with standard- or intermediate-risk outcomes. We revealed distinct immunophenotypic features for *DUX4*-rearranged (CD13^{high}, CD34^{high}, CD2⁺, CD20⁻), *PAX5* P80R (CD10⁻, CD2⁺, CD25⁺), and

Ph-like (CD38weak) genotypes. These data in a large adult B-ALL cohort treated in a non-risk-adapted approach on a single trial show the important prognostic value of genomic analyses which may translate into therapeutic benefits for future patients.

Introduction

Despite recent advances in therapeutic options for patients with B-lineage acute lymphoblastic leukemia (B-ALL), treatment of adults remains challenging because of a high incidence of adverse prognostic factors and poorer tolerance of intensive therapy. The Medical Research Council (MRC), now National Cancer Research Institute, UKALLXII/ECOG-ACRIN Cancer Research Group E2993 trial (ClinicalTrials.gov NCT00002514) accrued 1229 adult patients with *BCR-ABL1* negative B-ALL between 1993 and 2006.¹⁻³ Among these patients, 41% of cases with successful cytogenetics had recurrent genetic abnormalities: t(v;11q23) (rearrangement of *KMT2A/MLL*), t(12;21)(p13;q22) (*ETV6-RUNX1*), t(1;19)(q23;p13.3) (*TCF3-PBX1*), t(5;14)(q31;q32) (*IGH-IL3*), or aneuploidy.⁴ Thus, in adult *BCR-ABL1* negative B-ALL, the majority of patients historically fell into the category of B-ALL not otherwise specified.⁵

Incorporation of measurable residual disease (MRD) to improve risk stratification, use of pediatric-inspired regimens for young adults, and the advent of targeted therapies, including monoclonal antibodies and chimeric antigen-receptor T-cells, have redefined the therapeutic paradigm of ALL.⁶ The genomic revolution has also led to improved categorization of B-ALL.⁷⁻¹⁰ In particular, the Ph-like

(*BCR-ABL1*-like) phenotype, characterized by a gene-expression profile similar to *BCR-ABL1* positive ALL (*BCR-ABL1* ALL) but with non-*BCR-ABL1* genetic alterations activating kinase signaling or alterations of lymphoid transcription factor genes,¹¹⁻¹⁴ has poor prognosis in both children and adults.¹⁵⁻²⁰ Due to their prognostic impact, Ph-like and hypodiploid B-ALL have been added as a provisional entities to the WHO Classification of ALL.^{21,22}

In light of these advances, we explored the clinical outcomes of the largest cohort ever reported of adult *BCR-ABL1* negative B-ALL, who were treated without any risk-adapted modifications, on the UKALLXII/E2993 trial. In parallel, we performed genomic analysis of E2993 patients to gain insight into associations of novel genotypes with treatment response.

Methods

Patient identification and treatment design

The MRC enrolled 820 and ECOG-ACRIN 409 *BCR-ABL1* negative B-ALL patients. Immunophenotyping and PCR for known fusions were performed centrally for E2993 patients (supplemental materials), while the MRC centrally reviewed institutional immunophenotyping. Local cytogenetic data were reviewed centrally.⁴ Chemotherapeutic regimens of the trial have been described.^{1-3,23} Patients over 35 years of age or with a WBC count greater than $30 \times 10^9/L$ at presentation were designated high-risk. The Institutional Review Board of participating centers approved this study, and informed consent was obtained in accordance with the Declaration of Helsinki.

Genomic profiling

Transcriptome sequencing (RNA-seq) was performed on 238 samples using TruSeq library preparation and HiSeq 2000/2500 or NovaSeq 6000 sequencers (Illumina). All sequence reads were paired-end, and sequencing was performed using (1) total RNA and stranded RNA-seq [100 base-pair (bp) reads] or (2) polyA-selected mRNA (100bp reads). Methods for RNA-seq processing and analysis are detailed in the supplemental materials. In a complementary approach, a 15-gene Taqman® quantitative RT-PCR low density array (LDA)^{15,24,25} that identifies the Ph-like ALL gene signature, *P2RY8-CRLF2*, *BCR-ABL1*, *ETV6-RUNX1*, *TCF3-PBX1*, *KMT2A*-rearranged and *DUX4*-rearranged ALL was performed on 210 samples.

To determine copy number alterations (CNAs) including focal deletions, single nucleotide polymorphism (SNP) analysis was performed on 129 of 282 samples with available DNA using either SNP 6.0 microarrays (Affymetrix, n=47) or the Infinium Omni2.5Exome-8 BeadChip Kit (Illumina, n=82) as previously described.^{14,15} DNA and RNA-seq data was available for 102 samples using the FoundationOne Heme panel.²⁶ PCR and Sanger Sequencing for single nucleotide variants (SNVs) and insertion/deletions (indels) was available for 47 samples.¹⁴ Genomic data is publicly available and has been deposited in the European Genome Phenome Archive, Accessions EGAS00001000654, EGAS00001001952, EGAS00001003266 and EGAS00001004638.

Molecular subtyping

Genomic classification for 282 cases (supplemental Figure S1) was based on the use of multiple assays where available: RNA-seq (n=238), LDA (n=210), cytogenetics (n=197) and PCR for known fusions (*ETV6-RUNX1*, *KMT2A-AFF1* and *TCF3-PBX1*; n=274) (supplemental Figure S2A). The overall schema for molecular classification of B-ALL, as described previously,²⁷ is provided in supplemental Figure S3. For cases with RNA-seq (n=238), we identified 17 leukemia subtypes defined by CNA for aneuploid cases, recurring genetic alterations (subtype-defining rearrangements and SNVs), distinct clustering by two-dimensional t-Distributed Stochastic Neighbor Embedding (tSNE) plot, and gene expression profiles determined by Prediction Analysis of Microarrays (PAM)²⁸ (supplemental Figure S4). For the remaining cases with cytogenetic and/or LDA data (n=43), we identified subtypes based on aneuploidy (hyperdiploidy, hypodiploidy), the presence of known fusions (*ETV6-RUNX1*, *KMT2A-R*, *TCF3-PBX1*) and gene expression (*DUX4-R*, Ph-like). One remaining case with *KMT2A-AFF1* was identified by routine clinical PCR. Details for the 282 E2993 cases with molecular classification are provided in supplemental Table S1.

Copy number alteration (CNA) detection

RNA-seq data may be used to robustly identify chromosomal and arm level CNAs (<https://github.com/honzee/RNAseqCNV/>). Genes were ordered based on the median absolute deviation (MAD) of their expression level across the samples and a subset (1/4 to 1/3) of the genes with least MAD were picked as stably expressed

genes. To assist the CNA calling, mutant allele frequency of SNVs detected from RNA-seq data were plotted against the gene expression level of the stably expressed genes to double check if the CNAs were reliable.

Statistical methods

Patient characteristics and remission rates were compared using χ^2 tests for heterogeneity, Mantel-Haenszel test for trend or the Mann-Whitney U test. The donor-no donor comparison included patients <50 or 55 years (age limits for potential ASCT). Censoring was at the earlier of date of last contact, or October 31, 2010, the data cut-off date. Relapse-free survival (RFS) was calculated from CR to relapse or death without relapse. Time to relapse was measured from start of induction. The log-rank method was used to compare time-to-event distributions. Where the hazard was nonproportional and the long-term survival was to be compared, the χ^2 *P* value for the difference in the survival percentages at 5 years was quoted. Cox regression analysis was used to investigate the association between a time-to-event variable and baseline covariates. Transplantation was entered into multivariable Cox models as a time-varying covariate. All *P* values were two-sided.

Both percent of leukemic lymphoblasts stained by individual antibodies and the median fluorescence intensity (MFI) of antibody binding divided by MFI in the absence of antibody (MFI ratio) were compared across all genetic subtypes by Kruskal-Wallis test, and then between all pairs of genetic subtypes by the Mann-Whitney U test, yielding 1501 comparisons. The *P* value threshold of 0.0209 was

determined by the adaptive profile information criteria²⁹ which balance the levels of false discovery and false negative rates.

Results

Biologic features for 1229 B-ALL patients

The median age of the cohort was 30 years (range 14 to 65). The median WBC count was $9.2 \times 10^9/L$ (range 0.1-930) and was $<30 \times 10^9/L$ in 73% of the patients (supplemental Table S2).

By immunophenotyping, 18.5% were CD10- pro-B ALL, 63% were CD10+ early pre-B ALL, 15.7% had pre-B ALL with cytoplasmic IgM and 2.8% were mature B-ALLs. No survival difference was seen according to maturational stages, when excluding mature B-ALL (supplemental Figure S5).

Cytogenetic analysis was unavailable in 401 (32.6%), normal (≥ 20 normal bone marrow metaphases) in 199 (16.2%) and abnormal in 629 patients (51.2%), based on standard G-banding (supplemental Table S3). In multivariate Cox analysis, patients with low-hypodiploid (LH) or complex karyotype or t(8;14) had inferior survival.

Molecular characterization, demographics and antigen expression in E2993 patients

Nucleic acids were available for molecular analysis from 282 E2993 patients, of whom 264 were eligible for trial (supplemental Figure S1). With the exception of

older age (median 35 vs 29 years, $p < 0.001$), this cohort did not differ with respect to demographic or clinical variables from *BCR-ABL1* negative B-ALL UKALLXII/E2993 patients who were not tested (supplemental Table S4). Thus, our study cohort is representative of the entire trial population.

In addition to subtypes recognized by karyotyping or PCR for chimeric fusions, such as *ETV6-RUNX1*, *TCF3-PBX1*, *TCF3-HLF*, *KMT2A*-rearranged (*KMT2A-R*) and aneuploid ALL (high-hyperdiploid [HH], low-hypodiploid [LH], near-haploid [NH], intrachromosomal amplification of chromosome 21 [iAMP21]), recently described genotypes included *DUX4*-rearranged (*DUX4-R*),^{30,31} *ETV6-RUNX1*-like,³² *PAX5alt* and *PAX5 P80R*,^{27,33-35} Ph-like with rearrangements and/or overexpression of *CRLF2* (Ph-like *CRLF2-R*) and Ph-like with genetic alterations activating other kinases or cytokine receptor signaling pathways (Ph-like non-*CRLF2-R*),¹¹⁻¹⁴ *ZNF384*-rearranged (*ZNF384-R*) and *ZNF384-R*-like^{30,36-39}, *MEF2D*-rearranged (*MEF2D-R*),^{40,41} *BCL2/MYC*,²⁷ and the genotype with concomitant *ZEB2* and *CEBPE* alterations,³⁴ using established criteria (supplemental Table S5 and supplemental Figure S3). For outcome analysis, LH and NH patients were combined despite their distinct biologies,^{42,43} because of their comparably poor prognosis.⁴⁴ Cases without a subtype-defining driver genetic alteration were excluded from further analysis ("B-other") (11%). Demographics and frequencies of genomic subgroups are summarized in Table 1. Barriers to the enrollment of minorities to adult cancer clinical trials during E2993, before current NIH initiatives,⁴⁵ prevented associations between genotype and race or ethnicity.

In 47% of cases, cytogenetic analysis failed or yielded a normal result. While for recurrent gene fusions, corresponding cytogenetic abnormalities were frequently detected, the majority of genetic alterations were cryptic in nature (supplemental Table S1). Within the Ph-like cohort, 62.5% had Ph-like *CRLF2*-R versus 37.5% with Ph-like non-*CRLF2*-R, similar to other reports in adult ALL.¹⁵⁻¹⁸ Among Ph-like *CRLF2*-R cases, the ratio of *IGH-CRLF2* to *P2RY8-CRLF2* rearrangements was 3:1, supporting a prevalence of *IGH-CRLF2* in adults.^{15,16} Genetic alterations in Ph-like non-*CRLF2*-R cases included rearrangements and mutations driving JAK-STAT signaling (*JAK2*, *EPOR*, *IL7R*, *TYK2*), fusions involving ABL-class genes (*ABL1*, *ABL2*, *PDGFRA*), and one case with *ETV6-NTRK3* fusion. As in pediatric ALL,³² a fusion involving *IKZF1* was observed in one of our *ETV6-RUNX1*-like ALL patients. Among *DUX4*-R patients, *IGH-DUX4*³⁰⁻³² and in the *MEF2D*-R group, *MEF2D-BCL9* fusions were the most common.^{40,41,46} *ZNF384* was recurrently rearranged to the transcriptional regulator EP300,^{36,38} while other partners (*TAF15*, *TCF3*)³⁷ were less common. All of the *BCL2/MYC* cases harbored rearrangements of *BCL2*, *BCL6* and/or *MYC* (Table 2).

The presence of SNVs, insertion/deletions (indels) and focal deletions was determined for key genes including *IKZF1*, *TP53*, and those related to JAK-STAT signaling (*JAK1*, *JAK2*, *IL7R*, *SH2B3*, *CRLF2*) (supplemental Figure 2B and supplemental Table S6). We observed a higher frequency of *IKZF1* alterations (deletion or mutation) in Ph-like than non Ph-like cases (27 of 31, 87% vs. 14 of 100, 14%), confirming earlier data.¹⁴ Overall, alterations in the JAK-STAT signaling pathway were identified in 15 of 271 cases assessed (5.5%), predominantly in Ph-

like ALL. Among 30 Ph-like *CRLF2*-R cases, we observed mutation of *JAK1* (n=1), *JAK2* (n=8) and *IL7R* (n=1). Two Ph-like *CRLF2*-R cases harbored *CRLF2* p.Phe232Cys in addition to mutation of *JAK1* or *JAK2*. An additional *IL7R* indel was identified in a Ph-like non-*CRLF2*-R case. One PAX5 P80R case harbored two *JAK2* mutations (p.Arg683Gly and p.Asp873Asn) and one B-other case with *IGH-CRLF2* (lacking the Ph-like signature) harbored *CRLF2* p.Phe232Cys. *TP53* mutations were more common in LH (15 of 28, 54, as expected,⁴² and *BCL2/MYC*-R (4 of 12, 33%) compared to the remaining cohort (5 of 231, 2.2%).

Clustering of antigen expression for samples with defined genotypes is illustrated in Figure 1 and distinguishing immunophenotypic features are described in supplemental Table S7. Unique observations included overexpression of CD13, CD34 and CD38 by *DUX4*-R compared to other genotypes ($p < 0.0001$). The T-cell antigen CD2, present in 68% of *DUX4*-R, was also found in 50% of PAX5 P80R patients. PAX5 P80R blasts were arrested at the pro-B stage, though with higher frequency of CD10+ blasts ($p = 0.0011$) and stronger CD10 expression ($p = 0.0052$) compared to prototype pro-B *KMT2A*-R blasts. PAX5 P80R blasts expressed CD33, while lacking CD65s and CD15s, the two myeloid antigens typically⁴⁷ expressed jointly in *KMT2A*-R. Antigen expression was indistinguishable between *KMT2A-AFF1* and *KMT2A-non-AFF1* blasts. Ph-like blasts with or without *CRLF2*-R differed only by a higher frequency of CD13+ blasts in Ph-like non-*CRLF2*-R cases ($p = 0.011$). Compared with *BCR-ABL1*, Ph-like blasts showed comparably weak CD38 but less dual CD13-CD33 ($p = 0.0001$) and CD25 expression ($p = 0.0003$), both characteristic features of *BCR-ABL1* ALL.⁴⁸ *ZNF384*-R/-like blasts were CD10- in

73% of cases but compared to *KMT2A-R*, CD24+ ($p=0.0088$) and CD20+ ($p<0.0001$) blasts were more frequent. *ZNF384-R*-like blasts lacked CD65s and CD15s but expressed dual CD13-CD33 comparable to *BCR-ABL1* ALL and CD25 in one-third of cases. Myeloperoxidase was not detected. There was neither overexpression of surface CD135 nor evidence of FLT3-ITD, a potentially targetable protein kinase suggested for *ZNF384-R*.⁴⁹ Blasts with *BCL2/MYC-R* genotype showed features of mature B-ALL with surface immunoglobulin monoclonality and strong CD20 but variable TdT and CD10 expression. LH blasts were often CD10- and CD65s+ or CD15s+, though dual CD65s-CD15s expression was rarer in LH than *KMT2A-R* cases ($p<0.0001$). This data supports the notion that LH ALL arises from transformation of an immature B-cell progenitor.⁴²

Outcome in the entire UKALLXII/E2993 B-ALL cohort

Of 1229 patients with *BCR-ABL1*-negative B-ALL, 93% achieved CR. The CR rate was higher in younger patients ($p<0.0001$), but did not differ by sex, WBC count or state of CNS involvement. Relapse occurred in 507 patients (41%). The median time to relapse was 13 months (range 28 days to 12 years). Among relapsed patients, 116/373 without transplant in first CR had a transplant post relapse (12 autologous, 30 sibling-ASCT, 66 matched-unrelated-donor and 8 alternative-donor transplants). Only 109 relapsed patients had an ASCT post relapse and 42 survived for a median of 4.5 years (10 days to 14.9 years) from relapse.

The 5-year OS was 42% (95% CI, 39, 44). In multivariate Cox analysis, increasing age ($p<0.0001$), higher WBC count ($p<0.0001$) and CNS involvement at diagnosis ($p=0.02$) were associated with shorter OS. In 326 patients randomized between chemotherapy and autologous transplant, the 5-year OS was 47% (95% CI 39, 54) versus 35% (95% CI 28, 43) ($p=0.03$). Figure 2a shows OS in these two groups at 10 years. In the donor-no donor analysis, the 5-year OS was 52% (95% CI 47, 58) for patients with a donor and 45% (95% CI 41, 50) for patients without (log rank $p>0.1$; χ^2 test of difference at 5 years, $p=0.05$) (Figure 2b with results carried out to 10 years). The relapse risk at 5 years for those with a donor was 30% (95% CI 24, 35) versus 55% (95% CI 50, 60) in those without (log rank $p<0.0001$; χ^2 at 5 years, $p<0.0001$). The percentage of deaths in remission at 5 years for those with a donor was 32% (95% CI 26, 37) versus 10% (95% CI 7, 13) for those without (log rank $p<0.0001$; χ^2 at 5 years, $p<0.0001$).

Outcome among molecular subgroups

Of 282 patients genotyped, 264 were treated on study. Outcome analysis was limited to subgroups with ≥ 5 patients and was done for 217 patients with molecular characterization (see supplemental Figure S1 for patient selection). *KMT2A-non-AFF1* patients were excluded due to their molecular heterogeneity, though outcome did not differ from that in *KMT2A-AFF1* patients. Ph-like patients with and without *CRLF2*-R were combined as their outcomes were superimposable.

CR rates ranged from 50% in the *BCL2/MYC* to 100% in the *ETV6-RUNX1*-like, HH, PAX5 P80R and *MEF2D*-R groups (supplemental Table S8). Confidence intervals overlapped, possibly due to the small sizes of some of the cohorts.

With a median of 8 years and 9 months of follow-up for survivors, RFS (supplemental Table S9) and OS (supplemental Table S10) varied markedly among groups (Figure 3). In Cox proportional-hazards models, outcomes of all groups were compared to that of *DUX4*-R as reference group. In univariable analysis (Table 3), *DUX4*-R, HH, *ETV6-RUNX1*-like, PAX5 P80R and *TCF3-PBX1* patients had comparable RFS and were designated molecular standard-risk (mSR). *BCL2/MYC*-R, *KMT2A-AFF1*, LN/NH and Ph-like patients did significantly worse than *DUX4*-R ($p < 0.001$), and they were considered molecular high-risk (mHR). PAX5alt, *ZNF384*-R/-like and *MEF2D*-R patients had intermediate-risk (mIR) of relapse with RFS inferior to *DUX4*-R patients, though with lower levels of significance. OS among mSR groups did not vary significantly. OS of mHR groups remained inferior to that of *DUX4*-R ($p < 0.001$), while among mIR, only PAX5alt patients remained significantly inferior to *DUX4*-R ($p = 0.01$) but not *ZNF384*-R/-like ($p = 0.111$) or *MEF2D*-R ($p = 0.098$) (Table 3). Molecular analysis stratified 64 patients (29.5%) as mSR, 44 (20.3%) as mIR, and 109 (50.2%) as mHR. To consider an effect of antigens on outcome beyond their association with various genotypes, we adjusted multivariable Cox models for the presence of CD34, CD10, CD19, CD24, CD20, CD22 and CD38. Neither antigen expression nor the type of transplant received affected prognosis (not shown).

OS in IKZF1-altered versus IKZF1 wild-type in Ph-like ALL did not differ, in part due to the expected^{12,14,15} high frequency of IKZF1 alterations in this group (22 of 29 cases treated on trial, 76%). With respect to RFS, the two cases with IKZF1 wild-type appeared to fare much better than the 24 IKZF1-altered cases who had achieved a CR (supplemental Figure S6), though cohorts were too small to reach significance. When comparing outcome of IKZF1 altered (n=41) versus IKZF1 (n=90) wild-type across the whole cohort, again no difference was detected. As a caveat, the cohort of IKZF1-altered cohort contained a majority of poor prognostic genotypes.

When applying the protocol-specified risk classification (age >35 years or WBC count >30x10⁹/L) to 216 molecularly characterized patients, 137 were allocated to the protocol-defined high-risk group (pHR). Of pHR patients, 21.2% had mSR and 18.2% had mIR genotypes (supplemental Table S11). pHR patients with mSR or mIR genotypes did significantly better than mHR patients among the pHR cohort (for RFS, HR=0.36 [95% CI 0.23, 0.57], p< 0.001; for OS, HR=0.39 [95% CI 0.25, 0.60], p< 0.001) (supplemental Table S12).

Figure 4 shows the effect of protocol-defined low- (pLR) and high-risk (pHR) on RFS and OS in molecular risk groups. RFS was not different between protocol-defined risk groups within each molecular risk group, while OS was positively affected both for mSR (HR=0.24, p=0.007) and mHR patients (HR=0.47, p=0.007) when these patients fit the criteria for pLR. Superior 5-year RFS and OS was experienced by patients in the pLR/mSR cohort (73.3% [95% CI 59.8, 89.9] and 87.9% [95% CI 77.5, 99.8], respectively), while patients in the pHR/mHR cohort did

worst (11.9% [95% CI 6.23, 22.9] and 12.9% [95% CI 7.19, 23.1], respectively) (Table 4).

As patients with sibling donor in the entire B-ALL cohort did better than patients without a donor, a donor-no donor analysis was done for molecular risk groups. Only 125 patients with molecular characterization had donor information available (57.6%). Neither RFS nor OS differed by donor availability in the mSR and mIR risk groups (supplemental Figure S7). In mHR patients, the data suggest longer RFS ($p=0.08$) and OS ($p=0.2$) for patients with a donor. Due to the small sample size of subgroups, interpretations of this data with respect to transplant recommendations are limited.

Discussion

A unique feature of our study is the integration of data from genomic testing and immunophenotyping with outcome in a large subset of unselected adult *BCR-ABL1* negative B-ALL patients treated on a single trial. Only 11% of patients were not assigned to a well-described genotype, a percentage as low as that achieved in children.

A review of prognostic biologic features in the entire UKALLXII/E2993 *BCR-ABL1* negative B-lineage cohort revealed that cytogenetic anomalies, but not B-cell maturation stage of leukemic lymphoblasts, adversely affected outcome. However, standard chromosome analysis was non-informative in 44% of E2993 patients. Since most genotypes in E2993 were not associated with recurrent chromosomal

abnormalities (supplemental Table S1), cytogenetic analysis cannot be considered an adequate approach for their detection.

Among genotypes identified, Ph-like ALL constituted the biggest group (19.2%), comparable to most studies in adult ALL,^{15,17,18,27} but lower than reported by others.^{16,19} Although Ph-like and *BCR-ABL1* ALL have similar genetic profiles, the immunophenotype of Ph-like ALL has remained undefined.⁵⁰ We show here that their antigen expression is similar and distinctive from that in other genotypes with respect to weak CD38 expression, but also different with lower frequency of dual CD13-CD33 positive and CD25+ blasts in Ph-like ALL. We describe, for the first time, CD25+ CD10-/dull pro-B *ZNF384-R*-like blasts. While overexpression of CD25 and its gene, interleukin-2 receptor α , are negative prognostic factors in *BCR-ABL1* ALL and *BCR-ABL1* negative ALL,⁵¹⁻⁵³ CD25 did not affect outcome in *ZNF384-R*-like patients (not shown). This supports the notion that antigen profiles affect outcome only when analyzed in conjunction with their underlying molecular counterparts. We found expression of the T-cell antigen CD2 not only in *DUX4-R*^{54,55} but also in PAX5 P80R cases. CD2 expression in two novel genotypes with favorable outcome explains its proposed role as a surrogate for good prognosis in pediatric B-ALL in the early literature.⁵⁶⁻⁵⁸

As previously seen in adult ALL,^{15-19,33,34,59} outcome of Ph-like patients was extremely poor, though we did not see inferior outcome of Ph-like *CRLF2-R* compared with Ph-like non-*CRLF2-R* patients.¹⁶ Noteworthy, an analysis of UKALLXII patients with deregulated *CRLF2* expression showed reduced event-free survival and OS.⁶⁰ Among our patients with *PAX5*-altered genotypes, *PAX5* P80R

had a favorable outcome, similar⁶¹ or better than reported,^{27,34} and significantly better than PAX5alt patients.²⁷ While *ZNF384-R*-like patients had an intermediate molecular risk, as reported,^{33,34,37,38,62} so did *MEF2D-R* patients, who did better than previously suggested.^{33,34,40,41} Our large group of adult *DUX4-R* patients confirmed the excellent outcome seen in children and young adults.^{27,30-32}

Genomic analysis revealed that, despite a median age of 35 years, one-third of our patients had a favorable molecular risk. Importantly, E2993 patients deemed high-risk based on age and presenting WBC count who were molecularly standard- or intermediate-risk did significantly better than molecular high-risk patients within the protocol-defined high-risk cohort. However, younger age and lower WBC count still positively affected OS both in the standard- and high-risk molecular subgroups (Figure 4), highlighting the importance of implementing genomic analysis in addition to traditional prognostic factors for risk stratification.

In summary, our results have important clinical implications. We have demonstrated that molecular subtypes differ strikingly in their responses to treatment on a single trial. While MRD is another accepted powerful prognostic indicator in adult B-ALL,^{63,64} its impact in genetic subgroups has rarely been analyzed. Ph-like ALL patients tend to remain MRD-positive after induction chemotherapy.^{16,17,20,65} In children, MRD was prognostically important for Ph-like or hypodiploid ALL.^{65,66} Other trials of Ph-like ALL,^{16,67} however, suggested that relapse-risk remained high despite risk-adapted therapy and outcomes remained poor even if MRD-negativity was achieved. Response kinetics, clinically relevant cut-off levels and optimal timing of MRD testing vary between low- and high-risk

genotypes.^{68,69} When UKALLXII/E2993 was activated in 1993, MRD testing was not yet standard of care in adult B-ALL. As a result, submission of follow-up specimens was not mandated, limiting our potential for MRD testing. The MRC assessed MRD in 161/820 UKALLXII *BCR/ABL1*-negative B-ALL patients (19.6% of their accrual) and demonstrated that MRD was a predictor of relapse both in protocol-defined standard- and high-risk patients, though cohorts were small.⁷⁰ While we could not answer the interaction of MRD and molecular genotype in the present study, we will get clarity from the recently completed phase 3 trial E1910 (ClinicalTrials.gov NCT02003222), in which flow cytometric MRD was determined throughout the course of disease, samples underwent multimodal genetic analyses, and treatment with blinatumomab was assigned or randomized based on MRD status.

Acknowledgements

This study was conducted in part by the ECOG-ACRIN Cancer Research Group (Peter J. O'Dwyer, MD and Mitchell D. Schnall, MD, PhD, Group Co-Chairs) and supported by the National Cancer Institute of the National Institutes of Health under the following award numbers: U10CA180820, U10CA180794, UG1CA189859, UG1CA232760, UG1CA233234, and UG1CA233290. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. Mention of trade names, commercial products, or organizations does not imply endorsement by the U.S. government. This work was further supported by National Cancer Institute National Institutes of Health (R35 CA197695 and P30 CA021765 [C.G.M.], R01 CA198089 [A.M.M.], P30 CA008748

[R.L.L.], P30 CA118100-15 [C.L.W.], R50 CA211-542-04 [R.H.], UG1 CA233332 [O.A.-W.]); Leukemia Lymphoma Society (SCOR 7013-17 [A.M.M.]); the National Institute of General Medical Sciences National Institutes of Health (P50 GM115279 [C.G.M.], the American Lebanese Syrian Associated Charities of St. Jude Children's Research Hospital (C.G.M.); Blood Cancer UK (15036 [A.V.M.], 15009 [B.P.]); Cancer Research UK (C27995/A21019)[A.V.M. and A.K.F.]. The authors acknowledge data provided by Foundation Medicine, Cambridge, MA, which were previously published.²⁶

Authorship

Contribution: E.P., K.G.R., C.G.M. and M.R.L. prepared the manuscript; E.P., K.G.R., Z.G., J.B., J.R., Y.Z., R.L.L., O.A.-W., Z.C; G.W.; C.Q.; L.S.; S.P.; C.L.W., R.H., L.F., B.P., A.M. and D.A. performed experiments and data analyses and helped with data interpretation; V.W., G.A.N.B., D.P. and C.C. performed statistical analyses; Y.Z. G.W.D., and A.V.M. centrally reviewed institutional cytogenetic data; H.M.L., S.M.L., A.K.F., D.I.M., P.H.W., J.M.R., M.S.T. and A.H.G. organized the clinical trial; all authors reviewed and approved the manuscript.

Conflict-of-interest disclosure: E.P. does consulting work with Supertechs Inc., C.G.M. has received grant funding from Abbvie, Loxo Oncology and Pfizer, speaking fees for Amgen, and is on the Advisory Board for Illumina; R.L.L. is on the supervisory board of Qiagen and is a scientific advisor for Imago, Mission Bio, Zentalis, Ajax, Auron, Prelude, C4 Therapeutics and Isoplexis; he receives research

support from and consulted for Celgene and Roche and has consulted for Incyte, Janssen, Astellas, Morphosys and Novartis; he has received honoraria from Roche, Lilly and Amgen for invited lectures and from Gilead for grant reviews. O.A.-W. has served as a consultant for H3B Biomedicine, Foundation Medicine Inc, Merck, Prelude Therapeutics and Janssen and is on the scientific advisory board of Envisagenics Inc, Pfizer Boulder and AlChemys Inc; he has received prior research funding from Loxo Oncology and H3B Biomedicine. D.I.M. does consulting work for Pfizer, Amgen and Novartis; S.M.L. receives honoraria from Daiichi-Sankyo, Pfizer, Bristol-Myers Squibb, Acceleron, Agios, Loxo Oncology and has institutional research support from Onconova, Kura, Hoffman La Roche, Ariad and Biosight. M.S.T. receives research funding from Abbvie, Cellerant, Orsenix, ADC Therapeutics, Biosight, Glycomimetics, Rafael Pharmaceuticals and Amgen, is on the advisory boards of Abbvie, BioLineRx, Daiichi-Sankyo, Orsenix, KAHR, Rigel, Nohla, Delta Fly Pharma, Tetrphase, Oncolyze, Jazz Pharma, Roche, Biosight, Novartis, and receives royalties from UpToDate. A.M.M. receives research support from Janssen, Daiichi-Sankyo, Sanofi and does consulting work for Epizyme, Constellation and Jubilant. The remaining authors declare no competing financial interests.

References

1. Rowe JM, Buck G, Burnett AK, et al: Induction therapy for adults with acute lymphoblastic leukemia: results of more than 1500 patients from

- the international ALL trial: MRC UKALL XII/ECOG E2993. *Blood*. 2005;106(12):3760-3767.
2. Goldstone AH, Richards SM, Lazarus HM, et al: In adults with standard-risk acute lymphoblastic leukemia, the greatest benefit is achieved from a matched sibling allogeneic transplantation in first complete remission, and an autologous transplantation is less effective than conventional consolidation/maintenance chemotherapy in all patients: final results of the International ALL Trial (MRC UKALL XII/ECOG E2993). *Blood*. 2008;111(4):1827-1833.
 3. Rowe JM, Buck G, Moorman AV, et al. Standard consolidation/maintenance chemotherapy is consistently superior to a single autologous transplant for adult patients with acute lymphoblastic leukemia: Results of the international ALL trial (MRC UKALL XII/ECOG E2993). *Blood*. 2008;112(11):3314.
 4. Moorman AV, Harrison CJ, Buck GAN, et al: Karyotype is an independent prognostic factor in adult acute lymphoblastic leukemia (ALL): analysis of cytogenetic data from patients treated on the Medical Research Council (MRC) UKALLXII/Eastern Cooperative Oncology Group (ECOG) 2993 trial. *Blood*. 2007;109(8):3189-3197.
 5. Swerdlow SH, Campo E, Harris NL. WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues. Lyon, France: IARC; 2008.

6. Hefazi M, Litzow M. Recent advances in the biology and treatment of B-cell acute lymphoblastic leukemia. *Blood and Lymphatic Cancer: Targets and Therapy*. 2018;8:47-61.
7. Iacobucci I, Mullighan CG: Genetic basis of acute lymphoblastic leukemia. *J Clin Oncol*. 2017;35(9):975-983.
8. Lilljebörn H, Fioretos T. New oncogenic subtypes in pediatric B-cell precursor acute lymphoblastic leukemia. *Blood*. 2017;130(12):1395-1401.
9. Roberts KG, Mullighan CG. The Biology of B-Progenitor Acute Lymphoblastic Leukemia. Cold Spring Harbor perspectives in medicine. 2019.
10. Pui CH, Nichols KE, Yang JJ. Somatic and germline genomics in paediatric acute lymphoblastic leukemia. *Nat Rev Clin Oncol*. 2019;16(4):227-240.
11. Den Boer ML, van Slegtenhorst M, De Menezes RX, et al. A subtype of childhood acute lymphoblastic leukaemia with poor treatment outcome: a genome-wide classification study. *Lancet Oncol*. 2009;10(2):125-134.
12. Mullighan CG, Su X, Zhang J, et al. Deletion of IKZF1 and prognosis in acute lymphoblastic leukemia. *N Engl J Med*. 2009;360(5):470-480.
13. Roberts KG, Morin RD, Zhang J, et al. Genetic alterations activating kinase and cytokine receptor signaling in high-risk acute lymphoblastic leukemia. *Cancer Cell*. 2012;22(2):153-166.

14. Roberts KG, Li Y, Payne-Turner D, et al. Targetable kinase-activating lesions in Ph-like acute lymphoblastic leukemia. *N Engl J Med*. 2014;371(11):1005-1015.
15. Roberts KG, Gu Z, Payne-Turner D, McCastlain K, et al. High frequency and poor outcome of Ph-like acute lymphoblastic leukemia in adults. *J Clin Oncol*. 2017;35(4):394-401.
16. Jain N, Roberts KG, Jabbour E, et al. Ph-like acute lymphoblastic leukemia: a high-risk subtype in adults. *Blood*. 2017;129(5):572-581.
17. Herold T, Schneider S, Metzeler KH, et al. Adults with Philadelphia chromosome-like acute lymphoblastic leukemia frequently have *IGH-CRLF2* and *JAK2* mutations, persistence of minimal residual disease and poor prognosis. *Haematologica*. 2017;102(1):130-138.
18. Tasian SK, Hurtz C, Wertheim GB, et al. High incidence of Philadelphia chromosome-like acute lymphoblastic leukemia (Ph-like ALL) in older adults with B-ALL. *Leukemia*. 2017;31(4):981-984.
19. Harvey RD, Tasian SK. Clinical diagnostics and treatment strategies for Philadelphia chromosome-like acute lymphoblastic leukemia. *Blood Adv*. 2020;4(1):218-228.
20. Chiaretti S, Messina M, Della Starza I, et al. Philadelphia-like acute lymphoblastic leukemia is associated with minimal residual disease persistence and poor outcome. First report of the minimal residual disease-oriented GIMEMA LAL1913. *Haematologica*. 2020;Doi: 10.3324/haematol.2020.247973.

- 21.Arber DA, Orazi A, Hasserjian R, et al. The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia. *Blood*. 2016;127(20):2391-2405.
- 22.Wang S, He G. 2016 Revision to the WHO classification of acute lymphoblastic leukemia. *J Transl Intern Med*. 2016;4:147-149.
- 23.Lazarus HM, Richards SM, Chopra R, et al. Central nervous system involvement in adult acute lymphoblastic leukemia at diagnosis: results from the international ALL trial MRC UKALL XII/ECOG E2993. *Blood*. 2006;108(2):465-472.
- 24.Harvey RC, Kang H, Roberts KG, et al. Development and validation of a highly sensitive and specific gene expression classifier to prospectively screen and identify B-precursor acute lymphoblastic leukemia (ALL) patients with a Philadelphia chromosome-like signature (“Ph-like” or BCR-ABL-Like”) for therapeutic targeting and clinical intervention. *Blood*. 2013;122(21):826.
- 25.Reshmi SC, Harvey RC, Roberts KG. Targetable kinase gene fusions in high-risk B-ALL: a study from the Children’s Oncology Group. *Blood*. 2017;129(25):3352-3361.
- 26.He J, Abdel-Wahab O, Nahas MK, et al. Integrated genomic DNA/RNA profiling of hematologic malignancies in the clinical setting. *Blood*. 2016;127(24):3004-3014.
- 27.Gu Z, Churchman ML, Roberts KG, et al. PAX5-driven subtypes of B-progenitor acute lymphoblastic leukemia. *Nat Genet*. 2019;51(2):296-307.

28. Tibshirani R, Hastie T, Narasimhan B, Chu G. Diagnosis of multiple cancer types by shrunken centroids of gene expression. *Proc Natl Acad Sci USA*. 2002;99(10):6567-6572.
29. Cheng C. An adaptive significance threshold criterion for massive multiple hypothesis testing. In: *Optimality: The Second Erich L. Lehmann Symposium, Institute of Mathematical Statistics, Lecture Notes-Monograph Series*. 2006;49:51-76.
30. Yasuda T, Tsuzuki S, Kawazu M, et al. Recurrent DUX4 fusions in B cell acute lymphoblastic leukemia of adolescents and young adults. *Nat Genet*. 2016;48:569–574.
31. Zhang J, McCastlain K, Wu G, et al. Simultaneous deregulation of DUX4 and ERG in acute lymphoblastic leukemia. *Nat Genet*. 2016;48(12):1481-1489.
32. Lilljebörn H, Henningsson R, Hyrenius-Wittsten A, et al: Identification of *ETV6-RUNX1*-like and *DUX4*-rearranged subtypes in paediatric B-cell precursor acute lymphoblastic leukaemia. *Nat Commun*. 2016;7:11790.
33. Liu Y-F, Wang B-Y, Zhang W-N, et al. Genomic profiling of adult and pediatric B-cell acute lymphoblastic leukemia. *EBioMedicine*. 2016;8:173-183.
34. Li J-F, Dai Y-T, Lilljebörn H, et al. Transcriptional landscape of B cell precursor acute lymphoblastic leukemia based on an international study of 1,223 cases. *Proc Natl Acad Sci USA*. 2018;115(50):E11711-E11720.

35. Zaliouva M, Stuchly J, Winkowska L, et al. Genomic landscape of pediatric B-other acute lymphoblastic leukemia in a consecutive European cohort. *Haematologica*. 2019;104(7):1396-13406.
36. Gocho Y, Kiyokawa N, Ichikawa H, et al: A novel recurrent *EP300-ZNF384* gene fusion in B-cell precursor acute lymphoblastic leukemia. *Leukemia*. 2015;29(12):2445-2448.
37. Hirabayashi S, Ohki K, Nakabayashi K, et al. ZNF384-related fusion genes define a subgroup of childhood B-cell precursor acute lymphoblastic leukemia with a characteristic immunotype. *Haematologica*. 2017;102(1):118-129.
38. McClure BJ, Heatley SL, Kok CH, et al. Pre-B acute lymphoblastic leukaemia recurrent fusion, EP300-ZNF384, is associated with a distinct gene expression. *Br J Cancer*. 2018;118:1000-1004.
39. Alexander TB, Gu Z, Iacobucci I, et al. The genetic basis and cell of origin of mixed phenotype acute leukaemia. *Nature*. 2018;562(7727):373-379.
40. Gu Z, Churchman M, Roberts K, et al. Genomic analyses identify recurrent MEF2D fusions in acute lymphoblastic leukemia. *Nat Commun*. 2016;7:133331
41. Suzuki K, Okuno Y, Kawashima N, et al. *MEF2D-BCL9* fusion gene is associated with high-risk acute B-cell precursor lymphoblastic leukemia in adolescents. *J Clin Oncol*. 2016;34(28):3451-3459.

42. Holmfeldt L, Wei L, Diaz-Flores E, et al: The genomic landscape of hypodiploid acute lymphoblastic leukemia. *Nature Genetics*. 2013;45(3):242-252.
43. Safavi S, Olsson L, Biloglav A, et al. Genetic and epigenetic characterization of hypodiploid acute lymphoblastic leukemia. *Oncotarget*. 2015;6(40):42793-42802.
44. Safavi S, Paulsson K. Near-haploid and low-hypodiploid acute lymphoblastic leukemia: two distinct subtypes with consistently poor prognosis. *Blood*. 2017;129(4):420-423.
45. Hamel LM, Penner LA, Albrecht TL, et al. Barriers to clinical trial enrollment in racial and ethnic minority patients with cancer. *Cancer Control*. 2016;23(4):327-337.
46. Ohki K, Kiyokawa N, Saito Y, et al. Clinical and molecular characteristics of *MEF2D* fusion-positive B-cell precursor acute lymphoblastic leukemia in childhood, including a novel translocation resulting in *MEF2D-HNRNPH1* gene fusion. *Haematologica*. 2019;104(1):128-137.
47. Marks DI, Moorman AV, Chilton L, et al. The clinical characteristics, biology, therapy and outcome of 85 adults with acute lymphoblastic leukemia and t(4;11)(q21;q23)/*MLL-AFF1* uniformly treated on UKALLXII/ECOG2993. *Haematologica*. 2013;98(6):945-952.
48. Paietta E. Immunobiology of Acute Leukemia. In Neoplastic Diseases of the Blood, 6th Edition (Wiernik P, Dutcher J, Gertz MA, eds) Springer: Media, PA, 2018;pp237-279.

49. Griffith M, Griffith OL, Krysiak K, et al. Comprehensive genomic analysis reveals FLT3 activation and a therapeutic strategy for a patient with relapsed adult B-lymphoblastic leukemia. *Exp Hematol*. 44(7):603-613.
50. Siegele BJ, Nardi V. Laboratory testing in *BCR-ABL1*-like (Philadelphia-like) B-lymphoblastic leukemia/lymphoma. *Am J Hematol*. 2018;93:971-977.
51. Geng H, Brennan S, Milne TA, et al: Integrative epigenomic analysis of adult B-acute lymphoblastic leukemia identifies biomarkers and therapeutic targets. *Cancer Discovery*. 2012;2(11):1004-1023.
52. Chen P, Chu A, Zia H, et al. CD25 expression in B lymphoblastic leukemia/lymphoma predicts t(9;22)(q34;q11)/Philadelphia chromosome translocation (Ph) and is associated with residual disease in Ph-negative patients. *Am J Clin Pathol*. 2016;146(5):632-638.
53. Sadras T, Heatley SL, Kok CH, et al. Differential expression of *MUC4*, *GPR110*, and *IL2RA* defines two groups of *CRLF2*-rearranged acute lymphoblastic leukemia patients with distinct secondary lesions. *Cancer Letters*. 2017;408:92-101.
54. Zaliouva M, Zimmermanova O, Dörge P, et al. ERG deletion is associated with CD2 and attenuates the negative impact of IKZF1 deletion in childhood acute lymphoblastic leukemia. *Leukemia*. 2014;28(1):182-185.
55. Schumich A, Zaliouva M, Fortschegger K, et al. CD371 cell surface expression: a unique feature of *DUX4*-rearranged acute lymphoblastic leukemia. *Haematologica*. 2019;104(8):e352-e255.

56. Uckun FM, Muraguchi A, Ledbetter JA, et al. Biphenotypic leukemic lymphocyte precursors in CD2+Cd19+acute lymphoblastic leukemia and their putative normal counterpart in human fetal hematopoietic tissues. *Blood*. 1989;73(4):1000-1015.
57. Cantu-Rajnoldi A, Putti MC, Schiro R, et al. Biological and clinical features of B-precursor childhood acute lymphoblastic leukemia showing CD2 and/or E-Rosette co-expression. *Haematologica*. 1992;77(5):384-391.
58. Dunphy CH, Chu J. Aberrant CD2 expression in precursor-B acute lymphoblastic leukemia of childhood. *Am J Hematol*. 1996;52(3):224-226.
59. Boer JM, Koenders JE, van der Holt B, et al. Expression profiling of adult acute lymphoblastic leukemia identifies a *BCR-ABL1*-like subgroup characterized by high non-response and relapse rates. *Haematologica*. 2015;100(7):e261-264.
60. Moorman AV, Schwab C, Ensor HM, et al. *IGH@* translocations, *CRLF2* deregulation, and microdeletions in adolescents and adults with acute lymphoblastic leukemia. *J Clin Oncol*. 2012;30(25):3100-3108.
61. Passet M, Boissel N, Sigaux F, et al. *PAX5* P80R mutation identifies a novel subtype of B-cell precursor acute lymphoblastic leukemia with favorable outcome. *Blood*. 2019;133(3):280-284.
62. Hirabayashi S, Butler E, Ohki K, et al. Acute lymphoblastic leukemia with Zinc-Finger Protein 384 (ZNF384)-related rearrangements: a

- retrospective analysis from the Ponte Di Legno Childhood ALL Working Group. *Blood*. 2019;134(Suppl 1):652.
63. Berry DA, Zhou S, Higley H, et al. Association of minimal residual disease with clinical outcome in pediatric and adult acute lymphoblastic leukemia. *JAMA Oncol*. 2017;3(7):e170580.
64. Bassan R, Brüggemann M, Radcliffe HS, et al. A systematic literature review and meta-analysis of minimal residual disease as a prognostic indicator in adult B-cell acute lymphoblastic leukemia. *Haematologica*. 2019;104(10):2028-2039.
65. Roberts KG, Pei D, Campana D, et al. Outcomes of children with BCR-ABL1-like acute lymphoblastic leukemia treated with risk-adapted therapy based on the levels of minimal residual disease. *J Clin Oncol*. 2014;32(27):3012-3020.
66. Mullighan CG, Jeha S, Pei D, et al. Outcome of children with hypodiploid ALL treated with risk-directed therapy based on MRD levels. *Blood*. 2015;126(26):2896-2899.
67. Heatley SL, Sadras T, Kok CH, et al. High prevalence of relapse in children with Philadelphia-like acute lymphoblastic leukemia despite risk-adapted treatment. *Haematologica*. 2017;102(12):e490-e493.
68. Pui C-H, Pei D, Raimondi SC, et al. Clinical impact of minimal residual disease in children with different subtypes of acute lymphoblastic leukemia treated with response-adapted therapy. *Leukemia*. 2017;31(2):333-339.

69. O'Connor D, Enshaei A, Bartram J, et al. Genotype-specific minimal residual disease interpretation improves stratification in pediatric acute lymphoblastic leukemia. *J Clin Oncol*. 2018;36(1):34-43.
70. Patel B, Rai L, Buck G, et al. Minimal residual disease is a significant predictor of treatment failure in non T-lineage adult acute lymphoblastic leukaemia: final results of the international trial UKALLXII/ECOG2993. *Br J Haematol*. 2010;148(1):80-89.

Table 1: Baseline characteristics of 251 molecularly characterized E2993 patients (excluding the 31 B-other patients), including patients who were not eligible for trial

Molecular subgroup	Number of patients	%	Age (years) median (Q1,Q3)	WBC x 10⁹/L median (Q1,Q3)	Sex (%) F/M
<i>DUX4-R</i>	22	8.8	26.0 (22.5,40.3)	9.3 (3.5,23.6)	52.4/47.6
<i>ETV6-RUNX1/-like*</i>	5	2.0	22.0 (20.0,26.0)	6.7 (2.0,9.7)	60.0/40.0
<i>TCF3-PBX1</i>	15	6.0	29.0 (22.5,37.5)	29.8 (10.4,61.2)	46.7/53.3
<i>KMT2A-AFF1</i>	27	10.8	38.0 (32.0,45.0)	92.0 (43.4,229.2)	76.9/23.1
<i>KMT2A-non-AFF1</i>	10	4.0	35.5 (31.5,46.5)	84.4 (6.1,190.2)	70/30
PAX5alt	27	10.8	38.0 (20.5,50.0)	9.2 (3.5,23.5)	18.5/81.5
PAX5 P80R	10	4.0	26.5 (20.8,46.3)	11.7 (3.9,22.5)	30/70
Ph-like <i>CRLF2-R</i>	30	12.0	28.0 (19.3,35.8)	23.3 (11.0,70.8)	20/80
Ph-like non- <i>CRLF2-R</i>	18	7.2	38.5 (30.0,48.8)	14.3 (6.1,60.4)	61.1/38.9
<i>ZNF384-R/-like*</i>	11	4.4	27.0 (26.0,42.5)	13.2 (6.2,36.7)	18.2/81.8
<i>MEF2D-R</i>	6	2.4	22.0 (19.0,43.8)	25.0 (10.0,32.9)	50.0/50.0
<i>BCL2/MYC</i>	12	4.8	40.0 (26.5,48.8)	28.5 (16.7,57.8)	36.4/63.6
<i>ZEB2/CEBPE</i>	4	1.6	43.0 (37.8,45.8)	32.6 (2.8,86.9)	50.0/50.0
<i>iAMP21</i>	3	1.2	21.0 (20.0,30.0)	4.2 (3.9,6.0)	33.3/66.7
<i>TCF3-HLF</i>	1	0.4	44 (44,44)	6.1 (6.1,6.1)	100/0
High Hyperdiploid	16	6.4	20.0 (18.0,29.3)	2.5 (1.9,9.5)	25/75
Low Hypodiploid/ Near Haploid	34	13.5	44.0 (36.0,53.0)	3.7 (2.9,7.6)	37.1/62.9

*, indicates combination of *ETV6-RUNX1* and *ETV6-RUNX1*-like as well as *ZNF384* and *ZNF384*-like patients

Table 2: Gene rearrangements and gene mutations of 251 molecularly characterized E2993 patients (excluding 31 B-other patients)

Molecular Subgroup	Gene rearrangements and mutations (# of patients)
<i>DUX4-R</i>	IGH-DUX4 (19), MZF1-DUX4 (1), DUX4 overexpression (2)
<i>ETV6-RUNX1</i>	ETV6-RUNX1 (3)
<i>ETV6-RUNX1-like</i>	IKZF1-FIGNL1 (1), no fusion (1)
<i>TCF3-PBX1</i>	TCF3-PBX1 (15)
<i>TCF3-HLF</i>	TCF3-HLF (1)
<i>KMT2A-R</i>	KMT2A-AFF1 (27), KMT2A-MLLT1 (5), KMT2A-EPS15 (2), KMT2A-MLLT3 (1), KMT2A-AFF1 ^{NEG} by PCR (2)
Ph-like <i>CRLF2-R</i>	IGH-CRLF2 (19), P2RY8-CRLF2 (6), IGH-CRLF2 + P2RY8-CRLF2 (1), USP9X-RBM10 (1), high CRLF2 expression and Ph-like score (2), NA (1)
Ph-like non- <i>CRLF2-R</i>	PAX5-JAK2 (2), SSBP2-JAK2 (2), PCM1-JAK2 (1), ZNF340-JAK2 (1), ZNF340-TYK2 (1), RCSD1-ABL1 (1), RCSD1-ABL2 (1), ETV6-NTRK3 (1), IGH-EPOR (1), IGK-EPOR (1), FIP1L1-PDGFR (1), IL7R p.L242ins (1), no fusion (4)
PAX5alt	PAX5-ETV6 (4), PAX5-ZNF318 (1), IGH-CRLF2 (1), P2RY8-CRLF2 (1), PAX5 G334fs (1), PAX5 R140L (1), PAX5 R59W (1), no fusion or mutation (17)
PAX5 P80R	P2RY8-CRLF2 (1), PAX5 p.Pro80Arg (9)
<i>ZNF384-R</i>	EP300-ZNF384 (6), TCF3-ZNF384 (2), TAF15-ZNF384 (1)
<i>ZNF384-like</i>	ZBT47-ZNF652 (1), no fusion (1)
<i>MEF2D-R</i>	MEF2D-BCL9 (5), MEF2D-SS18 (1)
<i>BCL2/MYC</i>	IGH-MYC (3), IGH-BCL2 (2), IGH-BCL2 + BACH2-MYC (1), REPS1-BCL2 (1), IGH-BCL2 + IGH-MYC (2), SMC4-BCL6 (1), IGH-BCL2 + MBNL1-BCL6 (1), IGH-BCL6 (1),
<i>ZEB2/CEBPE</i>	IGH-CEBPE (3), no fusion (1)
<i>iAMP21</i>	IGH-CRLF2 (1), no fusion (2)
HH	no fusion (9), NA (7)
LH/NH	CREBBP-UTRN (1), TFAP4-TRAP1 (1), PAX5 R140L (1), no fusion (29), NA (3)

no fusion, inability to detect a fusion with the methodology used; NA, RNA-seq was not performed

Table 3: Univariable Cox-model of relapse-free (RFS) and overall survival (OS) for molecular subgroups compared to the *DUX4*-R patients as reference group

Molecular subgroup	RFS Hazard Ratio	P-value	OS Hazard Ratio	P-value
<i>BCL2/MYC</i>	36.4 (7.9, 167.2)	<0.001	11.3 (3.2, 39.3)	<0.001
<i>KMT2A-AFF1</i>	7.8 (2.6, 23.3)	<0.001	6.6 (2.5, 17.1)	<0.001
LH/NH	7.6 (2.6, 22.4)	<0.001	5.3 (2.0, 14.1)	<0.001
Ph-like	7.5 (2.7, 21.3)	<0.001	4.9 (1.9, 12.6)	<0.001
<i>MEF2D-R</i>	5.9 (1.6, 21.8)	0.009	3.0 (0.8, 11.3)	0.098
PAX5alt	4.6 (1.5, 13.7)	0.006	3.7 (1.4, 10.0)	0.010
<i>ZNF384-R/-like</i>	4.0 (1.1, 14.3)	0.031	2.6 (0.8, 8.6)	0.111
HH	2.4 (0.7, 8.4)	0.154	1.0 (0.3, 3.7)	0.995
<i>ETV6-RUNX1-like</i>	2.0 (0.4, 11.1)	0.411	0.8 (0.1, 6.7)	0.819
<i>PAX5 P80R</i>	1.5 (0.3, 6.8)	0.585	1.2 (0.3, 5.1)	0.795
<i>TCF3-PBX1</i>	1.2 (0.3, 5.5)	0.781	1.6 (0.5, 5.7)	0.429

Table 4: 5-year relapse-free (RFS) and overall survival (OS) of E2993 patients by combined protocol-defined and molecular-defined risk assignment.

Combined risk group	5-year RFS % (95% confidence interval)	5-year OS % (95% confidence interval)
pLR/mSR	73.3 (59.8, 89.9)	87.9 (77.5, 99.8)
pLR/mIR	29.4 (14.1, 61.4)	31.6 (16.3, 61.2)
pLR/mHR	28.9 (14.7, 56.6)	36.2 (21.5, 60.9)
pHR/mSR	62.0 (45.2, 85.1)	53.2 (37.5, 75.6)
pHR/mIR	27.7 (13.8, 55.3)	42.0 (25.5, 69.4)
pHR/mHR	11.9 (6.23, 22.9)	12.9 (7.19, 23.1)

pLR, protocol-defined low-risk; pHR, protocol-defined high-risk; mSR, molecular standard-risk; mIR, molecular intermediate-risk; mHR, molecular high-risk;

Figure 1: Antigen expression in major molecular subgroups.




Figure 2a Overall survival in *BCR/ABL1*-negative B-ALL patients on UKALLXII/E2993 by randomization allocation at 10 years.




Figure 2b Overall survival of *BCR/ABL1*-negative B-ALL patients on UKALLXII/E2993 by matched sibling donor availability at 10 years.




Figure 3 Outcome of E2993 Molecular Subgroups.





Figure 4: Outcome of E2993 patients by protocol-defined and molecular risk assignment.



LEGENDS

Figure 1. Antigen expression in major molecular subgroups. Unsupervised hierarchical clustering of antigen expression in each of 413 E2993 patients. The patient cohort consists of 244 samples from *BCR-ABL1* negative E2993 patients which were genotyped (excluding molecular subgroups with <5 patients and B-other) and 169 *BCR-ABL1* E2993 ALL patients. This approach identified characteristic antigen expression profiles within each of the 13 molecular subtypes, shown as black bars above the heatmap. Columns represent patients and rows are antigens. Antigens are listed to the right of the heatmap. Expression levels reflect the percentage of antibody-binding leukemic lymphoblasts, except for CD38_mfi, which reflects intensity of antigen expression. High- and low-expression of antigens in the heatmap are shown in shades of red and blue, respectively. The first row above the heatmap aligns genotypes with B-lineage maturation stages. ZNF384-R includes ZNF384-like; ETV6-RUNX1 includes ETV6-RUNX1-like; KMT2A-R includes both KMT2A-AFF1 and KMT2A-non-AFF1; low_hypo, low-hypodiploid and near-haploid patients; mcd22, membrane (surface) CD22;

Figure 2a. Overall survival in *BCR/ABL1*-negative B-ALL patients on UKALLXII/E2993 by randomization allocation at 10 years. Patients without an allogeneic donor were randomly assigned to standard consolidation/maintenance chemotherapy or to receive a single autologous transplantation (autograft).

Figure 2b. Overall survival of BCR/ABL1-negative B-ALL patients on UKALLXII/E2993 by matched sibling donor availability at 10 years.

Figure 3. Outcome of E2993 Molecular Subgroups. Kaplan-Meier estimates for relapse-free (RFS) and overall survival (OS) of molecular subgroups included in the outcome analysis. The RFS analysis included only patients who achieved a complete remission (*DUX4*-R, 18 patients; *TCF3-PBX1*, 12; *PAX5* P80R, 9; Hyperdiploid, 15; *ETV6-RUNX1*/-like, 5; *ZNF384*-R/-like, 10; *PAX5alt*, 24; *MEF2D*-R, 6; *KMT2A-AFF1*, 21; Ph-like, 41; Hypodiploid/Haploid, 24; *BCL2/MYC*, 3). The OS analysis included all patients with outcome information (*DUX4*-R, 20 patients; *TCF3-PBX1*, 15; *PAX5* P80R, 9; Hyperdiploid, 15; *ETV6-RUNX1*/-like, 5; *ZNF384*-R/-like, 11; *PAX5alt*, 27; *MEF2D*-R, 6; *KMT2A-AFF1*, 26; Ph-like, 46; Hypodiploid/Haploid, 31; *BCL2/MYC*, 6). Among *DUX4*-R, *ETV6-RUNX1*/-like, *TCF3-PBX1*, *PAX5* P80R and hyperdiploid patients, neither RFS nor OS varied significantly.

Figure 4. Outcome of E2993 patients by protocol-defined and molecular-defined risk assignment. Kaplan-Meier estimates of relapse-free (RFS) and overall survival (OS) for patients stratified by their protocol-defined risk: low-risk (pLR) and high-risk (pHR), and molecular risk: standard-risk (mSR), intermediate-risk (mIR), and high-risk (mHR).