

Research Article

Andrea Mondino* and Christian Scharrer

A strict inequality for the minimization of the Willmore functional under isoperimetric constraint

<https://doi.org/10.1515/acv-2021-0002>

Received January 4, 2021; revised April 30, 2021; accepted May 5, 2021

Abstract: Inspired by previous work of Kusner and Bauer–Kuwert, we prove a strict inequality between the Willmore energies of two surfaces and their connected sum in the context of isoperimetric constraints. Building on previous work by Keller, Mondino and Rivière, our strict inequality leads to existence of minimizers for the isoperimetric constrained Willmore problem in every genus, provided the minimal energy lies strictly below 8π . Besides the geometric interest, such a minimization problem has been studied in the literature as a simplified model in the theory of lipid bilayer cell membranes.

Keywords: Willmore functional, isoperimetric constraint, existence of minimizers, strict energy inequality

MSC 2010: Primary 53A05; secondary 53A30

Communicated by: Tristan Rivière

1 Introduction

The Willmore energy of an immersed closed surface $f : \Sigma \rightarrow \mathbb{R}^3$ is given by

$$\mathcal{W}(f) = \frac{1}{4} \int_{\Sigma} H^2 d\mu,$$

where the mean curvature H is defined as the sum of the principal curvatures, and μ is the Radon measure corresponding to the pull-back metric of the Euclidean metric along f . The isoperimetric ratio is defined by

$$\text{iso}(f) = \frac{\text{area}(f)}{\text{vol}(f)^{\frac{2}{3}}}, \quad (1.1)$$

where

$$\text{area}(f) = \int_{\Sigma} 1 d\mu, \quad \text{vol}(f) = \frac{1}{3} \int_{\Sigma} n \cdot f d\mu \quad (1.2)$$

are the area and enclosed volume, and $n : \Sigma \rightarrow \mathbb{S}^2$ is the Gauss map. One can find different definitions of isoperimetric ratio in the literature. Note that with the choice (1.1), $\text{iso}(f)$ is invariant under constant scaling of f , and it is minimized by any parametrization of the round sphere $\mathbb{S}^2 \subset \mathbb{R}^3$, as a consequence of the Euclidean isoperimetric inequality. Thus

$$\text{image}(\text{iso}) = [\sqrt[3]{36\pi}, \infty),$$

*Corresponding author: Andrea Mondino, Mathematical Institute, University of Oxford, Oxford, United Kingdom, e-mail: andrea.mondino@maths.ox.ac.uk. <https://orcid.org/0000-0002-1932-7148>

Christian Scharrer, Mathematics Institute, University of Warwick, Coventry, United Kingdom, e-mail: c.scharrer@warwick.ac.uk. <https://orcid.org/0000-0002-6651-6179>

where the image is taken over the class of immersed surfaces. Denote with \mathcal{S}_g the set of smooth immersions $f: \Sigma \rightarrow \mathbb{R}^3$ where Σ is a closed surface (i.e. compact without boundary) with genus $(\Sigma) = g$. We are interested in the following minimization problem.

Problem 1.1 (Isoperimetric constrained Willmore problem). Let g be a non-negative integer and fix $\sigma > \sqrt[3]{36\pi}$. Minimize the Willmore energy \mathcal{W} in the class of immersions $f \in \mathcal{S}_g$ subject to the constraint $\text{iso}(f) = \sigma$. That is, find $f_0 \in \{f \in \mathcal{S}_g : \text{iso}(f) = \sigma\}$ such that

$$\mathcal{W}(f_0) \leq \mathcal{W}(f) \quad \text{for any } f \in \mathcal{S}_g \text{ with } \text{iso}(f) = \sigma. \quad (1.3)$$

Such an immersion f_0 satisfying (1.3) is referred to as *solution* or *minimizer*.

Beyond the geometric interest, the minimization problem 1.1 is partially motivated by a model for closed lipid bilayer cell membranes proposed by Canham [3] and Helfrich [5]. Indeed, the Willmore energy is the main term in the Canham–Helfrich functional which describes the free energy of a closed lipid bilayer:

$$F_{\text{Can-Hel}} := \int_{\text{lipid bilayer}} \left(\frac{k_c}{2} (2H + c_0)^2 + \bar{k}K + \lambda \right) + p \cdot V,$$

where c_0 is the spontaneous curvature, k_c and \bar{k} are bending rigidities, λ is the surface tension, K is the Gauss curvature, p is the osmotic pressure and V the enclosed volume. According to such a model, the shapes of cell membranes observed in nature correspond to (local) minimizers of $F_{\text{Can-Hel}}$. Notice that, if $c_0 = \lambda = p = 0$, then one obtains the Willmore functional (up to a scaling factor and a topological term, by the Gauss–Bonnet theorem). If instead λ and p do not vanish, they can be seen as Lagrange multipliers for area and volume constraints. Thanks to the scaling invariance of the Willmore functional, such a constrained problem is thus strictly related to the isoperimetric constrained Willmore problem 1.1.

Even if spherical membranes are most common, also higher genus membranes have been observed in nature: for toroidal shapes see [18, 23] and for higher genus see [17, 19, 29]. The Canham–Helfrich and Willmore energies are commonly used in mathematical biology, for instance in modelling red blood cells [3, 20], crista junctions in mitochondria [24] and folds of endoplasmatic reticulum [30]. In particular, the isoperimetric constrained Willmore problem was studied in the axially symmetric case by numerical approximation of the corresponding ordinary differential equations in [28] (see also [15]). Without symmetry assumptions, the existence of spherical minimizers was achieved by Schygulla [27] and the higher genus case was investigated by Keller, Mondino and Rivière [7]. The goal of the paper is to establish the following theorem (for more details, see Corollary 1.6 below).

Theorem 1.2. Let g be a non-negative integer and fix $\sigma > \sqrt[3]{36\pi}$. Assume that

$$\beta_g(\sigma) := \inf\{\mathcal{W}(f) : f \in \mathcal{S}_g, \text{iso}(f) = \sigma\} < 8\pi. \quad (1.4)$$

Then $\beta_g(\sigma)$ is attained by a smoothly embedded minimizer $f_0 \in \mathcal{S}_g$, i.e. f_0 satisfies (1.3).

In the remaining part of the introduction, we set Theorem 1.2 in context of the existing literature about both the free (i.e. unconstrained) and the isoperimetric constrained Willmore problems. We will start by discussing how the minimizers for the free problem already provide partial solutions to Problem 1.1.

Problem 1.3 (Classical Willmore problem). Let g be a non-negative integer. Minimize the Willmore energy \mathcal{W} in the class \mathcal{S}_g . That is, find $f_0 \in \mathcal{S}_g$ such that $\mathcal{W}(f_0) \leq \mathcal{W}(f)$ for any $f \in \mathcal{S}_g$.

As a first result on the unconstrained problem, Willmore [32] showed that the energy now bearing his name is bounded below by 4π on the class of closed surfaces, with equality only for round spheres, which solves the genus $g = 0$ case. Later, Simon [31] proved existence of Willmore minimizers with prescribed genus g , provided

$$\beta_g := \inf\{\mathcal{W}(f) : f \in \mathcal{S}_g\} < \min\{8\pi, \omega_g\}, \quad (1.5)$$

where

$$\omega_g = \min\left\{4\pi + \sum_{i=1}^p (\beta_{g_i} - 4\pi) : g = \sum_{i=1}^p g_i, 1 \leq g_i < g\right\}. \quad (1.6)$$

This assumption is used to obtain compactness in the direct method of calculus of variations. It was already known since the work of Willmore that $\beta_1 \leq 2\pi^2$, and therefore $\beta_1 < 8\pi$. Moreover, since by definition $\omega_1 = \infty$, Simon in particular proved existence of Willmore tori (i.e. genus $g = 1$ minimizers). Kusner [9] showed that $\beta_g < 8\pi$ for all $g \geq 1$, by estimating the area of minimal surfaces in the 3-sphere \mathbb{S}^3 found by Lawson [14]. Hence, in order to prove existence of Willmore minimizers with prescribed genus $g \geq 2$, the missing step was to show that

$$\beta_g < \omega_g. \quad (1.7)$$

There were some suggestions on that inequality before it was finally proven. Namely Simon [31] conjectured that $\beta_g \geq 6\pi$ for all $g \geq 1$, which would imply $\omega_g > 8\pi$, reducing the compactness assumption in (1.5) to the 8π -bound proven by Kusner. Simon's conjecture is now known to be true, but we will discuss it later. Furthermore, he explained that the non-strict inequality

$$\beta_g \leq \omega_g \quad (1.8)$$

is indeed true. To see this, he suggested to choose p surfaces $f_1 \in \mathcal{S}_{g_1}, \dots, f_p \in \mathcal{S}_{g_p}$ with Willmore energies close to $\beta_{g_1}, \dots, \beta_{g_p}$, respectively. Then, to each surface f_i , apply a sphere inversion

$$I_{a_i} : \mathbb{R}^3 \setminus \{a_i\} \rightarrow \mathbb{R}^3, \quad I_{a_i}(x) = \frac{x - a_i}{|x - a_i|^2}, \quad (1.9)$$

for some point $a_i \in \text{image } f_i$ of multiplicity one, turning the surface f_i into an unbounded surface $I_{a_i} \circ f_i$ with a planar end and Willmore energy $\mathcal{W}(I_{a_i} \circ f_i) = \mathcal{W}(f_i) - 4\pi$ (in fact, he suggested to choose a_i close to the image of f_i which results in a surface that already looks like a round sphere; the final construction however will look the same). Then focus on the part of $I_{a_i} \circ f_i$ that carries the genus of f_i , cut away the planar end and glue it into a large round sphere. The gluing can be done at small cost in terms of Willmore energy in such a way that the resulting surface looks like a round sphere with a cap of g_i handles, having the same genus as f_i and Willmore energy close to the sum $\mathcal{W}(\mathbb{S}^2) + \mathcal{W}(I_{a_i} \circ f_i)$. Gluing suitable sphere inversions of the surfaces f_1, \dots, f_p all into the same large sphere, results in a surface f with

$$\text{genus}(f) = \text{genus}(f_1) + \dots + \text{genus}(f_p)$$

that looks like a round sphere with p caps and Willmore energy

$$\mathcal{W}(f) \approx \mathcal{W}(\mathbb{S}^2) + \sum_{i=1}^p \mathcal{W}(I_{a_i} \circ f_i) \approx 4\pi + \sum_{i=1}^p (\beta_{g_i} - 4\pi),$$

which indeed implies the non-strict inequality (1.8). In fact, in order to prove either of the inequalities (1.7) or (1.8), one might assume that $p = 2$ in the definition of ω_g (see equation (1.6)). The general case then follows by induction. Kusner [9] developed the *conformal connected-sum* $M \# N$ of two given immersed surfaces M and N in \mathbb{R}^3 satisfying

$$\mathcal{W}(M \# N) = \mathcal{W}(M) + \mathcal{W}(N) - 4\pi,$$

which also implies the non-strict inequality (1.8). This kind of equation can be found in many mathematical concepts. Notable for instance is that the same equation holds true for the Euler characteristic χ of the connected sum $M \# N$ of two n -manifolds M and N :

$$\chi(M \# N) = \chi(M) + \chi(N) - \chi(\mathbb{S}^n).$$

Later, Kusner [10] suggested to invert the two surfaces at nonumbilic points after which the planar end of each surface is asymptotic to the graph of a biharmonic function with higher order terms decaying at least as fast as $\frac{1}{r}$. Therefore, one can then weld together two such inverted surfaces along a line in their planar ends and estimate the saved energy in terms of the energy of a biharmonic graph. Inspired from this idea, Bauer and Kuwert [2] finally found a proof for the strict inequality (1.7) and thus completed the existence proof of Problem 1.3. Given two smoothly immersed surfaces $f_i : \Sigma_i \rightarrow \mathbb{R}^3$ with $i = 1, 2$ neither of which is a round sphere, they constructed an immersed surface $f : \Sigma \rightarrow \mathbb{R}^3$ with topological type of the connected sum $\Sigma_1 \# \Sigma_2$

by inverting the first surface f_1 at a nonumbilic point and gluing the inverted surface directly into a large copy of the second surface, again at a nonumbilic point. The gluing was done by the graph of a biharmonic function. Thereby they inferred

$$\mathcal{W}(f) < \mathcal{W}(f_1) + \mathcal{W}(f_2) - 4\pi, \quad (1.10)$$

which implies (1.7).

An alternative way to prove the strict inequality for the high genus case follows from [13]. They proved that $\lim_{g \rightarrow \infty} \beta_g = 8\pi$, which then implies $\lim_{g \rightarrow \infty} \omega_g > 8\pi$. Later, existence of Willmore minimizers was also proven by a parametric approach in independent works of Kuwert and Li [11] (building on top of previous work of Müller and Šverák [22]) and Rivière [25] (building on top of Hélein's moving frames technique [4]). In the parametric approach, the inequality in (1.5) is needed to obtain compactness in the moduli space of conformal structures over a Riemann surface.

We next discuss the connection between the classical Willmore problem 1.3 and the isoperimetric constrained Willmore problem 1.1. Given any smoothly embedded surface $f: \Sigma \rightarrow \mathbb{R}^3$ (such are Willmore minimizers), one can find a smooth curve $\gamma: (0, \infty) \rightarrow \mathbb{R}^3$ with $\text{image}(\gamma) \cap \text{image}(f) = \emptyset$ such that the sphere inversions $I_\gamma \circ f$ (see the definition in (1.9)) have the following properties. The isoperimetric ratio $\text{iso}(I_{\gamma(r)} \circ f)$ varies smoothly in r ,

$$\lim_{r \rightarrow 0^+} \text{distance}(\gamma(r), \text{image}(f)) = 0, \quad \lim_{r \rightarrow \infty} \text{distance}(\gamma(r), \text{image}(f)) = \infty$$

and, most importantly,

$$\lim_{r \rightarrow 0^+} \text{iso}(I_{\gamma(r)} \circ f) = \text{iso}(\mathbb{S}^2), \quad \lim_{r \rightarrow \infty} \text{iso}(I_{\gamma(r)} \circ f) = \text{iso}(f).$$

Given any integer $g \geq 1$ and any (unconstrained) Willmore minimizer $\Sigma_g \in \mathcal{S}_g$, it follows that for all isoperimetric ratios σ in the non-empty interval

$$(\text{iso}(\mathbb{S}^2), \text{iso}(\Sigma_g)], \quad (1.11)$$

the isoperimetric constrained Willmore problem 1.1 corresponding to g and σ has a smoothly embedded solution, given by a suitable sphere inversion of Σ_g . In particular, the function $\beta_g: (\sqrt[3]{36\pi}, \infty) \rightarrow \mathbb{R}$ (defined in (1.4)) is constant on the interval in (1.11) and non-decreasing on the whole domain.

It follows by the proof of the Willmore conjecture by Marques and Neves [16] that

$$\beta_g \geq 2\pi^2 \quad \text{for all } g \geq 1, \quad (1.12)$$

with equality attained only on the Clifford torus (or inversions of it). In particular, $\beta_g \geq 6\pi$ as conjectured by Simon, and thus $\omega_g > 8\pi$. This fact reduces the compactness assumption in (1.5) to the 8π -bound proven by Kusner. Thereby, Marques and Neves provide another proof for the strict inequality (1.7) used to show existence of solutions for the classical Willmore problem 1.3, alternatively to the one of Bauer and Kuwert. Thus the interval in (1.11) for $g = 1$ reads as

$$\left(\sqrt[3]{36\pi}, \sqrt[3]{16\sqrt{2}\pi^2} \right]. \quad (1.13)$$

Notice that recently, Rivière [26] gave a PDE based proof of the Willmore conjecture (i.e. equation (1.12)).

The classical Willmore problem only provides solutions for the isoperimetric constrained Willmore problem via sphere inversions for genus $g \geq 1$, as for $g = 0$ the minimizer of the Willmore energy coincides with the minimizer of the isoperimetric ratio giving thus an empty interval in (1.11). The genus $g = 0$ case was fully solved by Schygulla [27] in the ambient approach (and later generalized to the non-zero spontaneous curvature case by the authors [21] using the parametric approach). We call *Schygulla spheres* the minimizers of the genus $g = 0$ case with isoperimetric ratio σ , and denote them with $\mathcal{S}(\sigma)$. The first existence result for the genus $g \geq 1$ case of the isoperimetric constrained Willmore problem was given by Keller, Mondino and Rivière [7]. They proved existence of smoothly embedded minimizers for all isoperimetric ratios σ satisfying

$$\beta_g(\sigma) < \min\{8\pi, \omega_g, (\beta_g + \beta_0(\sigma) - 4\pi)\}. \quad (1.14)$$

Compare this inequality with Simon's compactness assumption (1.5). Schygulla also showed that

$$\beta_0(\sigma) = \mathcal{W}(\mathcal{S}(\sigma))$$

is continuous in σ . Hence the right-hand side in (1.14) is continuous in σ . Consequently, by the result of Keller, Mondino and Rivière, one can show (see [7, Theorem 1.4]) that the set of isoperimetric ratios for which there exist constrained minimizers, is an open interval containing the interval in (1.11).

By the result of Marques and Neves, the constant ω_g on the right-hand side of (1.14) is redundant. The main result of this paper will be that also the last constant can be neglected, reducing the compactness assumption (1.14) to

$$\beta_g(\sigma) < 8\pi.$$

It is yet unknown whether or not this inequality is always satisfied. Instead, it is clear that the non-strict inequality

$$\beta_g(\sigma) \leq 8\pi$$

holds true. This can be seen by taking two concentric spheres of nearly the same radii and connecting them with $(g + 1)$ catenoidal necks, resulting in a surface of genus g . This construction was carried out by Kühnel and Pinkall [8]. The isoperimetric ratio of these surfaces tends to infinity, while the Willmore energy approaches 8π from above.

We can now state the main result of the present paper.

Theorem 1.4. *Suppose Σ_1, Σ_2 are two closed surfaces, $f_1 : \Sigma_1 \rightarrow \mathbb{R}^3$ is a smooth embedding, $f_2 : \Sigma_2 \rightarrow \mathbb{R}^3$ is a smooth immersion, and neither f_1 nor f_2 parametrize a round sphere. Denote with Σ the connected sum $\Sigma_1 \# \Sigma_2$. Then there exists a smooth immersion $f : \Sigma \rightarrow \mathbb{R}^3$ such that*

$$\text{iso}(f) = \text{iso}(f_2) \quad (1.15)$$

and

$$\mathcal{W}(f) < \mathcal{W}(f_1) + \mathcal{W}(f_2) - 4\pi. \quad (1.16)$$

Moreover, if also f_2 is an embedding, then f is an embedding as well.

Recall that the inequality in (1.16) has been proven by Bauer and Kuwert [2] in order to solve the classical Willmore problem (see (1.10)). It is novel that the same inequality remains valid under the additional condition on the isoperimetric ratios (1.15). Indeed, in order to prove Theorem 1.4, we use the same connected sum construction developed by Bauer and Kuwert. It will be shown in Section 3 that the connected sum already satisfies Equation (1.15) asymptotically (see Lemma 3.2). We then adjust the isoperimetric ratio by applying a first variation of the surface f_2 supported away from the pasting region, inspired by Huisken's volume preserving mean curvature flow [6] (see Lemma 2.1 in Section 2). Using existence of smoothly embedded Schygulla spheres as well as existence of smoothly embedded Willmore minimizers, we infer the following corollary.

Corollary 1.5. *Given any integer $g \geq 1$, there holds*

$$\beta_g(\sigma) < \beta_g + \beta_0(\sigma) - 4\pi \quad \text{for all } \sigma > \sqrt[3]{36\pi}.$$

This corollary answers to a question raised by Keller, Mondino and Rivière in [7, Remark 1.7 (iii)]. Our corollary together with their results leads to the following statement, regarding Problem 1.1.

Corollary 1.6. *Given any integer $g \geq 1$, the function $\beta_g(\cdot)$ (defined as in (1.4)) is non-decreasing, continuous, and bounded: $4\pi < \beta_g(\cdot) \leq 8\pi$. Moreover, $\beta_g(\sigma)$ is attained by a smoothly embedded minimizer for all σ with $\beta_g(\sigma) < 8\pi$.*

Remark 1.7. The Willmore energy of a torus of revolution with radii $0 < r < R$ is given by

$$\mathcal{W}(\mathbb{T}_c) = \frac{\pi^2}{c\sqrt{1-c^2}},$$

where $c = \frac{r}{R}$. Its minimum is attained at $c = 1/\sqrt{2}$, which results in the Clifford torus. Moreover,

$$c_1 := \inf\{c > 0 : \mathcal{W}(\mathbb{T}_c) < 8\pi\} = \sqrt{\frac{1}{2} - \frac{\sqrt{16 - \pi^2}}{8}}.$$

The isoperimetric ratio of a torus of revolution \mathbb{T}_c can be computed as $\text{iso}(\mathbb{T}_c) = \sqrt[3]{16\pi^2/c}$. We thus obtain existence of solutions for all isoperimetric ratios in the interval

$$(\text{iso}(\mathbb{S}^2), \text{iso}(\mathbb{T}_{c_1})) = \left(\sqrt[3]{36\pi}, \sqrt[3]{16\pi^2 / \sqrt{\frac{1}{2} - \frac{\sqrt{16 - \pi^2}}{8}}} \right). \quad (1.17)$$

Notice that for the Clifford torus $\mathbb{T} = \mathbb{T}_{1/\sqrt{2}}$ there holds

$$\text{iso}(\mathbb{T}) = \sqrt[3]{16\sqrt{2}\pi^2} < \text{iso}(\mathbb{T}_{c_1}).$$

Hence, our solution interval (1.17) is strictly larger than the interval in (1.13).

Remark 1.8. As recalled above, [7] showed that the solution interval is open and contains the interval in (1.11). As a consequence of our main results, we obtain that an improved upper bound for the solution interval is given by $\inf\{\sigma : \beta_g(\sigma) = 8\pi\}$.

Remark 1.9. It is expected that if $\beta_g(\sigma) < 8\pi$ for all $\sigma > \sqrt[3]{36\pi}$, then

$$\lim_{\sigma \rightarrow \infty} \beta_g(\sigma) = 8\pi.$$

Indeed, this was proven for $g = 0$ by Schygulla [27] (see also [12] for a detailed blow-up analysis).

2 A suitable volume preserving first variation

In this section, we construct a variation of a given surface that initially linearly decreases the isoperimetric ratio, provided the surface is not a round sphere. The variation vector field can even be supported away from a given point. Note that the variation constructed below coincides at first order with the volume preserving mean curvature flow as developed by Huisken [6], multiplied by a suitable cut-off function.

Lemma 2.1. *Suppose $f : \Sigma \rightarrow \mathbb{R}^3$ is a smoothly immersed closed surface which is not a round sphere and $q \in \Sigma$ is any given point. Then there exists $q \neq p \in \Sigma$ with the following property: For each neighborhood U of p there exists a smooth normal vector field $\xi : \Sigma \rightarrow \mathbb{R}^3$ compactly supported in U such that for $f_t := f + t\xi$ with $t \in \mathbb{R}$ the function $t \mapsto \text{iso}(f_t)$ is differentiable at $t = 0$, and*

$$\left. \frac{d}{dt} \right|_{t=0} \text{iso}(f_t) \neq 0.$$

Moreover,

$$\mathcal{W}(f_t) = \mathcal{W}(f) + O(t) \quad \text{as } t \rightarrow 0.$$

Proof. First of all, recall that for any smooth vector field $\xi : \Sigma \rightarrow \mathbb{R}^3$ the family $f_t = f + t\xi$ defines a variation of the immersion f . In particular, for small $t \in \mathbb{R}$, the map $f_t : \Sigma \rightarrow \mathbb{R}^3$ is again a smooth immersion. Thus area, volume, and Willmore energy are defined for f_t with t small.

By a classical theorem of Alexandrov [1], since $f : \Sigma \rightarrow \mathbb{R}^3$ is not a round sphere, the mean curvature H cannot be constant. Therefore, we can choose a point p in the non-empty boundary of the set

$$\{x \in \Sigma : H(x) = \max \text{image } H\},$$

where $\text{image } H$ is the image of the mean curvature H . In fact, given any $c \in \text{image } H$, we might as well have chosen p in the non-empty boundary of the level set $\{H = c\}$. In particular, we can make sure that $p \neq q$. Now,

given any neighborhood U of p , we pick a smooth function $\varphi : \Sigma \rightarrow \mathbb{R}$ compactly supported in U such that $\varphi \geq 0$ and $\varphi(p) = 1$. Let $n : \Sigma \rightarrow \mathbb{S}^2$ be the Gauss map and define the constant h and the vector field ξ by

$$h = \int_{\Sigma} \varphi H \, d\mu \Big/ \int_{\Sigma} \varphi \, d\mu \quad \text{and} \quad \xi = \varphi(H - h)n.$$

Then $\xi : \Sigma \rightarrow \mathbb{R}^3$ is a smooth vector field compactly supported in U . Using the first variation formula of the volume, we compute for $f_t = f + t\xi$ that

$$\left. \frac{d}{dt} \right|_{t=0} \text{vol}(f_t) = - \int_{\Sigma} n \cdot \xi \, d\mu = - \int_{\Sigma} \varphi(H - h) \, d\mu = 0. \quad (2.1)$$

Moreover, using the first variation formula of the area, it follows

$$\left. \frac{d}{dt} \right|_{t=0} \text{area}(f_t) = - \int_{\Sigma} Hn \cdot \xi \, d\mu = - \int_{\Sigma} \varphi H(H - h) \, d\mu = - \int_{\Sigma} \varphi(H - h)^2 \, d\mu < 0. \quad (2.2)$$

The last expression is non-zero due to our choice of the point p and the function φ . Using (2.1) and (2.2), we infer that

$$\left. \frac{d}{dt} \right|_{t=0} \text{iso}(f_t) = - \int_{\Sigma} \varphi(H - h)^2 \, d\mu \Big/ \text{vol}(f)^{\frac{2}{3}} < 0.$$

Finally, using the first variation formula for the Willmore energy, we see that the function $t \mapsto \mathcal{W}(f_t)$ is differentiable at $t = 0$, which implies the conclusion. \square

3 Isoperimetric balance of the connected sum

In this section, we recall the connected sum construction developed by Bauer and Kuwert [2] and estimate its change of isoperimetric ratio (see Lemma 3.2). Then we prove Theorem 1.4 by applying the volume preserving variation constructed in Lemma 2.1 to the connected sum.

Let $f_i : \Sigma_i \rightarrow \mathbb{R}^3$ for $i = 1, 2$ be two smoothly immersed closed surfaces neither of which is a round sphere such that

$$f_i^{-1}\{0\} = \{p_i\} \quad \text{for some } p_i \in \Sigma_i, \quad \text{image } Df_i(p_i) = \mathbb{R}^2 \times \{0\}. \quad (3.1)$$

For some $\rho > 0$, one can then pick smooth local graph representations

$$f_1(z) = (z, u(z)), \quad f_2(z) = (z, v(z)) \quad \text{for } z \in D_\rho,$$

where D_ρ is the open disk $\{z \in \mathbb{R}^2 : |z| < \rho\}$. Letting $P, Q : \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow \mathbb{R}$ be the second fundamental forms at the origin of f_1 and f_2 , respectively, we define the error terms φ and ψ such that

$$\begin{aligned} u(z) &= p(z) + \varphi(z), \quad \text{where } p(z) = \frac{1}{2}P(z, z) \text{ for } z \in D_\rho, \\ v(z) &= q(z) + \psi(z), \quad \text{where } q(z) = \frac{1}{2}Q(z, z) \text{ for } z \in D_\rho. \end{aligned}$$

We denote the trace-free parts of the second fundamental forms with

$$P^\circ(w, z) = P(w, z) - \frac{(\text{tr } P)}{2}w \cdot z, \quad Q^\circ(w, z) = Q(w, z) - \frac{(\text{tr } Q)}{2}w \cdot z.$$

In view of [2, Lemma 4.5], we may assume that in addition to (3.1) there also holds

$$\langle P^\circ, Q^\circ \rangle > 0. \quad (3.2)$$

By [2, Lemma 2.3], the inverted and translated surface

$$f_1^\circ : \Sigma_1 \setminus \{p_1\} \rightarrow \mathbb{R}^3, \quad f_1^\circ(p) = \frac{f_1(p)}{|f_1(p)|^2} - \frac{(\text{tr } P)}{4}e_3,$$

where $e_3 = (0, 0, 1)$ is the third unit vector in \mathbb{R}^3 , has a graph representation at infinity. That is, outside of a large ball around zero, f_1° is given by the graph of a smooth function u° on $\mathbb{R}^2 \setminus D_R$ for some $R > 0$ with

$$u^\circ(z) = p^\circ(z) + \varphi^\circ(z), \quad \text{where } p^\circ(z) = \frac{1}{2}P^\circ\left(\frac{z}{|z|}, \frac{z}{|z|}\right) \quad (3.3)$$

such that the error term satisfies

$$|z||\varphi^\circ(z)| + |z|^2|D\varphi^\circ(z)| + |z|^3|D^2\varphi^\circ(z)| \leq C \quad \text{for } z \in \mathbb{R}^2 \setminus D_R. \quad (3.4)$$

Given any function $w : \Omega \rightarrow \mathbb{R}$ for $\Omega \subset \mathbb{R}^2$ and given any scalar $\lambda > 0$, we define the scaled function w_λ by

$$w_\lambda : \Omega_\lambda = \{z \in \mathbb{R}^2 : \lambda^{-1}z \in \Omega\} \rightarrow \mathbb{R}, \quad w_\lambda(z) = \lambda w(\lambda^{-1}z).$$

Hence, for small $\alpha, \beta > 0$, the graph representations of the scaled surfaces αf_1° and $(1/\beta)f_2$ are given by

$$\begin{aligned} u_\alpha^\circ(z) &= p_\alpha^\circ(z) + \varphi_\alpha^\circ(z) & \text{for } z \in \mathbb{R}^2 \setminus D_{\alpha R}, \\ v_{1/\beta}(z) &= q_{1/\beta}(z) + \psi_{1/\beta}(z) & \text{for } z \in D_{\rho/\beta}. \end{aligned}$$

Next, pick a smooth function $\eta : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$\eta(t) = \begin{cases} 0, & t \leq \frac{1}{4}\sqrt{\alpha}, \\ 1, & t \geq \frac{3}{4}\sqrt{\alpha}, \end{cases}$$

and such that $|\eta| + \sqrt{\alpha}|\eta'| + \alpha|\eta''| \leq C$ for some $0 < C < \infty$ independent of α . Then, for a third parameter γ with $0 < \alpha, \beta \ll \gamma \ll 1$, define for $r = |z|$,

$$w(z) = \begin{cases} p_\alpha^\circ(z) + \eta(\gamma - r)\varphi_\alpha^\circ(z), & \alpha R < r \leq \gamma, \\ q_{1/\beta}(z) + \eta(r - 1)\psi_{1/\beta}(z), & 1 \leq r < \frac{\rho}{\beta}, \end{cases}$$

and notice that $w = u_\alpha^\circ$ for $r \leq \gamma - \frac{3}{4}\sqrt{\alpha}$ as well as $w = v_{1/\beta}$ for $r \geq 1 + \frac{3}{4}\sqrt{\alpha}$. Moreover, on $D_1 \setminus D_\gamma$, let w be the unique solution of the bi-harmonic Dirichlet–Neumann problem (see [2, Lemma 3.1 and Lemma 3.2])

$$\begin{cases} \Delta^2 w = 0 & \text{in } D_1 \setminus D_\gamma, \\ w = p_\alpha^\circ, \quad \partial_r w = \partial_r p_\alpha^\circ & \text{on } |z| = \gamma, \\ w = q_{1/\beta}, \quad \partial_r w = \partial_r q_{1/\beta} & \text{on } |z| = 1. \end{cases} \quad (3.5)$$

To define the pasted surface, let U be the complement in Σ_1 of the preimage of the set

$$\{z \in \mathbb{R}^2 : \gamma - \sqrt{\alpha} < |z| < \infty\}$$

under the map $\alpha \cdot \pi \circ f_1^\circ$, where $\pi : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ denotes the orthogonal projection. Analogously, let V be the complement in Σ_2 of the preimage of the set $\{z \in \mathbb{R}^2 : |z| < 1 + \sqrt{\alpha}\}$ under the map $\frac{1}{\beta} \cdot \pi \circ f_2$. Moreover, let

$$W = \{z \in \mathbb{R}^2 : \gamma - \sqrt{\alpha} \leq |z| \leq 1 + \sqrt{\alpha}\}.$$

Then we can write the connected sum $\Sigma = \Sigma_1 \# \Sigma_2$ as $\Sigma = (U \cup V \cup W) / \sim$, where the identification \sim is given by

$$\begin{aligned} p \sim z &= \alpha \pi(f_1^\circ(p)) & \text{for } p \in U, z \in W, \\ q \sim z &= \frac{1}{\beta} \pi(f_2(q)) & \text{for } q \in V, z \in W. \end{aligned}$$

Now, the immersion of the patched surface can be defined by

$$f : \Sigma \rightarrow \mathbb{R}^3, \quad f(x) = \begin{cases} \alpha f_1^\circ(p), & x = p \in U \subset \Sigma_1, \\ \frac{1}{\beta} f_2(q), & x = q \in V \subset \Sigma_2, \\ (z, w(z)), & x = z \in W. \end{cases} \quad (3.6)$$

The connected sum satisfies the following energy saving proven by Bauer and Kuwert [2].

Lemma 3.1 (See [2, Lemma 4.4]). *Taking $\beta = t\alpha$ for any $t > 0$, and letting α tend to zero, there holds*

$$\mathcal{W}(f) - (\mathcal{W}(f_1) + \mathcal{W}(f_2) - 4\pi) = \pi\alpha^2(|P^\circ|^2 - t\langle P^\circ, Q^\circ \rangle + O_t(y^2 \log(y)^2) + O_{t,y}(\alpha^{1/2})), \quad (3.7)$$

where the constants in O_t (respectively in $O_{t,y}$) depend on t (respectively on t and y).

We will show that the isoperimetric ratio of the connected sum behaves as follows.

Lemma 3.2. *Taking $\beta = t\alpha$ for any $t > 0$, and letting α tend to zero, there holds*

$$\text{iso}(f) = \text{iso}(f_2) + O_{t,y}(\alpha^{2+\frac{1}{2}}), \quad (3.8)$$

where the constant in $O_{t,y}$ depends on t and y .

Proof. First, we will compute the area of the surface $f: \Sigma \rightarrow \mathbb{R}^3$. By definition of the connected sum in equation (3.6), we can split the area into

$$\text{area}(f) = \text{area}(f|_U) + \text{area}(f|_W) + \text{area}(f|_V). \quad (3.9)$$

Let

$$U_1 = \Sigma_1 \setminus (\pi \circ f_1^\circ)^{-1}\{z \in \mathbb{R}^2 : R < |z| < \infty\},$$

where again π denotes the orthogonal projection of \mathbb{R}^3 onto \mathbb{R}^2 . Then we can write

$$\text{area}(f|_U) = \alpha^2 \text{area}(f_1^\circ|_{U_1}) + \int_{D_{\gamma-\sqrt{\alpha}} \setminus D_{\alpha R}} \sqrt{1 + |Du_\alpha^\circ|^2} d\mathcal{L}^2, \quad (3.10)$$

where \mathcal{L}^2 denotes the 2-dimensional Lebesgue measure. For p° defined as in equation (3.3), we have

$$Dp^\circ(z) = P^\circ\left(\frac{z}{|z|}, D\left(\frac{z}{|z|}\right)\right), \quad D\left(\frac{z}{|z|}\right) = \frac{\text{Id}}{|z|} - \frac{\langle z, \cdot \rangle}{|z|^3} z.$$

Hence, $|p^\circ(z)| + |z||Dp^\circ(z)| \leq C$ for $z \in \mathbb{R}^2 \setminus D_R$ and after scaling,

$$|p_\alpha^\circ(z)| + |z||Dp_\alpha^\circ(z)| \leq C\alpha \quad \text{for } z \in \mathbb{R}^2 \setminus D_{\alpha R}.$$

Moreover, from the error estimation in equation (3.4),

$$|z||\varphi_\alpha^\circ(z)| + |z|^2|D\varphi_\alpha^\circ(z)| \leq C\alpha^2 \quad \text{for } z \in \mathbb{R}^2 \setminus D_{\alpha R}.$$

Using $u_\alpha^\circ = p_\alpha^\circ + \varphi_\alpha^\circ$, we thus infer

$$|u_\alpha^\circ(z)| + |Du_\alpha^\circ(z)| \leq C \quad \text{for } z \in D_{\sqrt{\alpha}R} \setminus D_{\alpha R}, \quad (3.11)$$

$$|u_\alpha^\circ(z)| + |Du_\alpha^\circ(z)| \leq C\sqrt{\alpha} \quad \text{for } z \in \mathbb{R}^2 \setminus D_{\sqrt{\alpha}R}. \quad (3.12)$$

Therefore, the area in equation (3.10) can be estimated by

$$\text{area}(f|_U) \leq C\alpha^2 + C\mathcal{L}^2(D_{R\sqrt{\alpha}} \setminus D_{\alpha R}) + (1 + C\sqrt{\alpha})\mathcal{L}^2(D_\gamma) = \mathcal{L}^2(D_\gamma) + O(\sqrt{\alpha})$$

as $\alpha \rightarrow 0$. On the other hand,

$$\text{area}(f|_U) \geq \mathcal{L}^2(D_{\gamma-\sqrt{\alpha}}) = \mathcal{L}^2(D_\gamma) - O(\sqrt{\alpha}) \quad \text{as } \alpha \rightarrow 0,$$

and thus

$$\text{area}(f|_U) = \mathcal{L}^2(D_\gamma) + O(\sqrt{\alpha}) \quad \text{as } \alpha \rightarrow 0. \quad (3.13)$$

From [2, Equation (4.13)] it follows that

$$|v_{1/\beta}(z)| + |Dv_{1/\beta}(z)| \leq C(t)\alpha \quad \text{for } z \in D_{1+\sqrt{\alpha}}, \quad (3.14)$$

and hence

$$\text{area}((1/\beta)f_2) - \text{area}(f|_V) = \int_{D_{1+\sqrt{\alpha}}} \sqrt{1 + |Dv_{1/\beta}|^2} d\mathcal{L}^2 = \mathcal{L}^2(D_1) + O_t(\sqrt{\alpha}) \quad (3.15)$$

as $\alpha \rightarrow 0$. Because of the homogeneity of p° and q (notice that $p_\alpha^\circ = \alpha p^\circ$, $q_{1/\beta} = \beta q$), the parameters α and β enter linearly into the boundary values of w on $D_1 \setminus D_\gamma$ and thus linearly into the solution (3.5) (see [2, (3.29), (3.30), and (3.35)]). Therefore, using $\beta = t\alpha$ as well as [2, (4.21), (4.22), and (4.25)], we infer

$$|w(z)| + |Dw(z)| \leq C(t, \gamma)\alpha \quad \text{for } z \in D_{1+\sqrt{\alpha}} \setminus D_{\gamma-\sqrt{\alpha}}. \quad (3.16)$$

It follows

$$\text{area}(f|_W) = \int_{D_{1+\sqrt{\alpha}} \setminus D_{\gamma-\sqrt{\alpha}}} \sqrt{1 + |Dw|^2} \, d\mathcal{L}^2 = \mathcal{L}^2(D_1 \setminus D_\gamma) + O_{t,\gamma}(\sqrt{\alpha}) \quad (3.17)$$

as $\alpha \rightarrow 0$. Putting (3.13), (3.15), and (3.17) into (3.9), leads to

$$\text{area}(f) = \text{area}((1/\beta)f_2) + O_{t,\gamma}(\sqrt{\alpha}) \quad \text{as } \alpha \rightarrow 0,$$

and thus

$$\text{area}(f) = (t\alpha)^{-2}(\text{area}(f_2) + O_{t,\gamma}(\alpha^{2+\frac{1}{2}})) \quad \text{as } \alpha \rightarrow 0. \quad (3.18)$$

Next, we will estimate the volume of the patched surface $f: \Sigma \rightarrow \mathbb{R}^3$. Using the definition of the volume (1.2) as well as the formula for the Gauss map of graphical surfaces, we estimate

$$\begin{aligned} |\text{vol}(f) - \text{vol}((1/\beta)f_2)| &\leq \alpha^3 \text{vol}(f_1^\circ|_{U_1}) + \int_{D_{\gamma-\sqrt{\alpha}} \setminus D_{\alpha R}} |z| |Du_\alpha^\circ| + |u_\alpha^\circ| \, d\mathcal{L}^2(z) \\ &\quad + \int_{D_{1+\sqrt{\alpha}} \setminus D_{\gamma-\sqrt{\alpha}}} |z| |Dw| + |w| \, d\mathcal{L}^2(z) + \int_{D_{1+\sqrt{\alpha}}} |z| |Dv_{1/\beta}| + |v_{1/\beta}| \, d\mathcal{L}^2(z). \end{aligned}$$

In view of (3.11), (3.12), (3.14), and (3.16), we can see that the right-hand side is uniformly bounded in α for $0 < \alpha \ll \gamma \ll 1$ and $\beta = t\alpha$. That means,

$$\text{vol}(f) = \text{vol}((1/\beta)f_2) + O_{t,\gamma}(1) \quad \text{as } \alpha \rightarrow 0,$$

and therefore

$$\text{vol}(f) = (t\alpha)^{-3}(\text{vol}(f_2) + O_{t,\gamma}(\alpha^3)) \quad \text{as } \alpha \rightarrow 0. \quad (3.19)$$

Notice that by differentiability of the function $s \mapsto (\text{vol}(f_2) + s)^{-2/3}$ at $s = 0$, there holds

$$\frac{1}{\text{vol}(f_2)^{\frac{2}{3}}} = \frac{1}{(\text{vol}(f_2) + s)^{\frac{2}{3}}} + O(s) \quad \text{as } s \rightarrow 0.$$

Thus, using $\beta = t\alpha$, (3.18), and (3.19), we infer

$$\text{iso}(f) = \frac{\text{area}(f_2)}{(\text{vol}(f_2) + O_{t,\gamma}(\alpha^3))^{\frac{2}{3}}} + \frac{O_{t,\gamma}(\alpha^{2+\frac{1}{2}})}{(\text{vol}(f_2) + O_{t,\gamma}(\alpha^3))^{\frac{2}{3}}} = \text{iso}(f_2) + O_{t,\gamma}(\alpha^{2+\frac{1}{2}})$$

as $\alpha \rightarrow 0$, which finishes the proof. \square

Now, we can prove Theorem 1.4.

Proof of Theorem 1.4. Let Σ_1, Σ_2 be two closed surfaces, let $f_1: \Sigma_1 \rightarrow \mathbb{R}^3$ be a smooth embedding, and let $f_2: \Sigma_2 \rightarrow \mathbb{R}^3$ be a smooth immersion such that neither f_1 nor f_2 parametrize a round sphere. Notice that the multiplicity of f_2 does not affect the construction of the connected sum. First, pick $p_i \in \Sigma_i$ for $i = 1, 2$ according to (3.1) and (3.2), with $p_2 \in f_2^{-1}\{0\}$ instead of $\{p_2\} = f_2^{-1}\{0\}$.

Apply Lemma 2.1 to the surface $f_2: \Sigma_2 \rightarrow \mathbb{R}^3$: denote $f_{2,s} = f_2 + s\xi$ with ξ compactly supported away from the point p_2 and

$$\mathcal{W}(f_{2,s}) = \mathcal{W}(f_2) + O(s) \quad \text{as } s \rightarrow 0, \quad (3.20)$$

$$\text{iso}(f_{2,s}) = \text{iso}(f_2) - c_2 s + o(s) \quad \text{as } s \rightarrow 0, \quad (3.21)$$

for some $c_2 > 0$. Now, apply the connected sum construction described in this section to the surfaces $f_1: \Sigma_1 \rightarrow \mathbb{R}^3$ and $f_{2,s}: \Sigma_2 \rightarrow \mathbb{R}^3$; in this way, we obtain the glued surface $f_{s,\alpha}: \Sigma \rightarrow \mathbb{R}^3$, where Σ is the connected sum of Σ_1 and Σ_2 . Notice that the right-hand side in equation (3.7) does not depend on s as

the vector field ξ is compactly supported away from the patching area. Therefore, we can first choose $t > 0$ large enough such that $|P^\circ|^2 - t\langle P^\circ, Q^\circ \rangle < 0$ and then choose $0 < \gamma < 1$ small enough such that still $|P^\circ|^2 - t\langle P^\circ, Q^\circ \rangle + O_t(\gamma^2 \log(\gamma)^2) < 0$, to obtain from Lemma 3.1 that

$$\mathcal{W}(f_{s,\alpha}) - (\mathcal{W}(f_1) + \mathcal{W}(f_{2,s}) - 4\pi) = -c\alpha^2 + O(\alpha^{2+\frac{1}{2}}) \quad (3.22)$$

as $\alpha \rightarrow 0$ for some $c > 0$. Putting (3.20) into (3.22) and (3.21) into (3.8), we infer

$$\mathcal{W}(f_{s,\alpha}) - (\mathcal{W}(f_1) + \mathcal{W}(f_2) - 4\pi) = -c\alpha^2 + O(\alpha^{2+\frac{1}{2}}) + O(s), \quad (3.23)$$

$$\text{iso}(f_{s,\alpha}) - \text{iso}(f_2) = O(\alpha^{2+\frac{1}{2}}) - c_2s + o(s) \quad (3.24)$$

as $s, \alpha \rightarrow 0$. Picking any $2 < m < 2 + \frac{1}{2}$, we see that for small $\alpha > 0$ and for $|s| \leq \alpha^m$ the right-hand side in (3.23) is strictly negative, while for $s = \alpha^m$ the right-hand side in (3.24) is strictly negative and for $s = -\alpha^m$ the right-hand side in (3.24) is strictly positive. Notice that once α is fixed, $\text{iso}(f_{s,\alpha})$ depends continuously on s . Therefore, there exists $\alpha > 0$ small and $-\alpha^m < s < \alpha^m$ such that the right-hand side in (3.23) is strictly negative, while the right-hand side in (3.24) is zero. In other words, $f_{s,\alpha}$ satisfies (1.15) and (1.16). Notice that the immersion $f_{s,\alpha}$ is smooth everywhere except on the boundary of $D_1 \setminus D_\gamma$, where the bi-harmonic function meets the second fundamental forms with Dirichlet and Neumann conditions; see (3.5). In general, $f_{s,\alpha}$ is only $C^{1,1}$ -regular. It remains to show that one can approximate $f_{s,\alpha}$ by a smooth immersion without losing the conditions (1.15) and (1.16). In view of its construction, we can choose a local graph representation of $f_{s,\alpha}$ given by a function u defined on an open subset of \mathbb{R}^2 that contains the boundary of $D_1 \setminus D_\gamma$. By multiplying with a cut-off function, one can write $u = u_s + u_r$ such that u_s is smooth and u_r is $C^{1,1}$ -regular as well as compactly supported. The standard mollification u_r^ε of u_r is smooth, compactly supported, and converges to u_r as $\varepsilon \rightarrow 0$ in the Sobolev space $W^{2,p}$ for all $1 \leq p < \infty$. The immersions f^ε corresponding to $u^\varepsilon := u_s + u_r^\varepsilon$ are smooth and differ from $f_{s,\alpha}$ only on a small neighborhood of the boundary of $D_1 \setminus D_\gamma$. Moreover, there holds

$$|\mathcal{W}(f^\varepsilon) - \mathcal{W}(f_{s,\alpha})| + |\text{iso}(f^\varepsilon) - \text{iso}(f_{s,\alpha})| \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0.$$

Hence there exists $\eta > 0$ such that, for $\varepsilon > 0$ small enough, f^ε satisfies the following quantified version of (1.16):

$$\mathcal{W}(f^\varepsilon) < \mathcal{W}(f_1) + \mathcal{W}(f_2) - 4\pi - \eta.$$

Finally, we once again apply Lemma 2.1 away from the support of u_r^ε to re-establish (1.15) still keeping the validity of (1.16). \square

Funding: The first author is supported by the European Research Council (ERC), under the European's Union Horizon 2020 research and innovation programme, via the ERC Starting Grant "CURVATURE", grant agreement no. 802689. The second author is supported by the EPSRC as part of the MASDOC DTC at the University of Warwick, grant no. EP/H023364/1.

References

- [1] A. D. Alexandrov, A characteristic property of spheres, *Ann. Mat. Pura Appl. (4)* **58** (1962), 303–315.
- [2] M. Bauer and E. Kuwert, Existence of minimizing Willmore surfaces of prescribed genus, *Int. Math. Res. Not. IMRN* **2003** (2003), no. 10, 553–576.
- [3] P. B. Canham, The minimum energy of bending as a possible explanation of the biconcave shape of the human red blood cell, *J. Theoret. Biol.* **26** (1970), no. 1, 61–81.
- [4] F. Hélein, *Harmonic Maps, Conservation Laws and Moving Frames*, 2nd ed., Cambridge Tracts in Math. 150, Cambridge University, Cambridge, 2002.
- [5] W. Helfrich, Elastic properties of lipid bilayers: Theory and possible experiments, *Z. Naturforschung C* **28** (1973), no. 11–12, 693–703.
- [6] G. Huisken, The volume preserving mean curvature flow, *J. Reine Angew. Math.* **382** (1987), 35–48.
- [7] L. G. A. Keller, A. Mondino and T. Rivière, Embedded surfaces of arbitrary genus minimizing the Willmore energy under isoperimetric constraint, *Arch. Ration. Mech. Anal.* **212** (2014), no. 2, 645–682.

- [8] W. Kühnel and U. Pinkall, On total mean curvatures, *Quart. J. Math. Oxford Ser. (2)* **37** (1986), no. 148, 437–447.
- [9] R. Kusner, Comparison surfaces for the Willmore problem, *Pacific J. Math.* **138** (1989), no. 2, 317–345.
- [10] R. Kusner, Estimates for the biharmonic energy on unbounded planar domains, and the existence of surfaces of every genus that minimize the squared-mean-curvature integral, in: *Elliptic and Parabolic Methods in Geometry* (Minneapolis 1994), A K Peters, Wellesley (1996), 67–72.
- [11] E. Kuwert and Y. Li, $W^{2,2}$ -conformal immersions of a closed Riemann surface into \mathbb{R}^n , *Comm. Anal. Geom.* **20** (2012), no. 2, 313–340.
- [12] E. Kuwert and Y. Li, Asymptotics of Willmore minimizers with prescribed small isoperimetric ratio, *SIAM J. Math. Anal.* **50** (2018), no. 4, 4407–4425.
- [13] E. Kuwert, Y. Li and R. Schätzle, The large genus limit of the infimum of the Willmore energy, *Amer. J. Math.* **132** (2010), no. 1, 37–51.
- [14] H. B. Lawson, Jr., Complete minimal surfaces in S^3 , *Ann. of Math. (2)* **92** (1970), 335–374.
- [15] R. Lipowsky, The conformation of membranes, *Nature* **349** (1991), 475–481.
- [16] F. C. Marques and A. Neves, Min-max theory and the Willmore conjecture, *Ann. of Math. (2)* **179** (2014), no. 2, 683–782.
- [17] X. Michalet and D. Bensimon, Observation of stable shapes and conformal diffusion in genus 2 vesicles, *Science* **269** (1995), no. 5224, 666–668.
- [18] X. Michalet and D. Bensimon, Vesicles of toroidal topology: Observed morphology and shape transformations, *J. Phys. II* **5** (1995), no. 2, 263–287.
- [19] X. Michalet, D. Bensimon and B. Fourcade, Fluctuating vesicles of nonspherical topology, *Phys. Rev. Lett.* **72** (1994), no. 1, 168–171.
- [20] T. Milcent, Shape derivative of the willmore functional and applications to equilibrium shapes of vesicles, Research Report RR-7539, INRIA, 2011.
- [21] A. Mondino and C. Scharrer, Existence and regularity of spheres minimising the Canham–Helfrich energy, *Arch. Ration. Mech. Anal.* **236** (2020), no. 3, 1455–1485.
- [22] S. Müller and V. Šverák, On surfaces of finite total curvature, *J. Differential Geom.* **42** (1995), no. 2, 229–258.
- [23] M. Mutz and D. Bensimon, Observation of toroidal vesicles, *Phys. Rev. A* **43** (1991), no. 8, 4525–4528.
- [24] C. Renken, G. Siragusa, G. Perkins, L. Washington, J. Nulton, P. Salamon and T. G. Frey, A thermodynamic model describing the nature of the crista junction: A structural motif in the mitochondrion, *J. Struct. Biol.* **138** (2002), 137–144.
- [25] T. Rivière, Lipschitz conformal immersions from degenerating Riemann surfaces with L^2 -bounded second fundamental forms, *Adv. Calc. Var.* **6** (2013), no. 1, 1–31.
- [26] T. Rivière, Minmax hierarchies, minimal fibrations and a PDE based proof of the Willmore conjecture, preprint (2020), <https://arxiv.org/abs/2007.05467>.
- [27] J. Schygulla, Willmore minimizers with prescribed isoperimetric ratio, *Arch. Ration. Mech. Anal.* **203** (2012), no. 3, 901–941.
- [28] U. Seifert, K. Berndt and R. Lipowsky, Shape transformations of vesicles: Phase diagram for spontaneous-curvature and bilayer-coupling models, *Phys. Rev. A* **44** (1991), no. 2, Article ID 1182.
- [29] U. Seifert and R. Lipowsky, Morphology of vesicles, in: *Handbook of Biological Physics*, Elsevier, Amsterdam (1995), 403–462.
- [30] Y. Shibata, T. Shemesh, W. A. Prinz, A. A. Palazzo, M. M. Kozlov and T. A. Rapoport, Mechanisms determining the morphology of the peripheral ER, *Cell* **143** (2010), 774–788.
- [31] L. Simon, Existence of surfaces minimizing the Willmore functional, *Comm. Anal. Geom.* **1** (1993), no. 2, 281–326.
- [32] T. J. Willmore, Note on embedded surfaces, *An. Şti. Univ. Al. I. Cuza Iaş Mat. (N. S.)* **11B** (1965), 493–496.