

# **Acute Kidney Injury and Long-Term Risk for Cardiovascular Events and Mortality.**

Ayodele Odutayo<sup>1,2</sup>, Christopher X Wong<sup>3</sup>, Michael Farkouh<sup>1</sup>, Douglas G Altman<sup>2</sup>, Sally Hopewell<sup>2</sup>, Connor A Emdin<sup>4</sup>, Benjamin H Hunn<sup>5,6</sup>

<sup>1</sup>Faculty of Medicine, University of Toronto, Toronto, Ontario.

<sup>2</sup>Centre for Statistics in Medicine, University of Oxford, Oxford, United Kingdom.

<sup>3</sup>Nuffield Department of Population Health, University of Oxford, Oxford, United Kingdom.

<sup>4</sup>St. John's College, University of Oxford, Oxford, United Kingdom.

<sup>5</sup>School of Medicine, University of Tasmania, Hobart, Australia.

<sup>6</sup>Department of Physiology, Anatomy and Genetics, University of Oxford, South Parks Road, Oxford, United Kingdom.

**Running Title:** AKI and Cardiovascular Disease

## **Correspondence to:**

Centre for Statistics in Medicine,  
Nuffield Department of Orthopedics, Rheumatology and Musculoskeletal Sciences,  
University of Oxford, Oxford,  
United Kingdom,  
OX3 7LD.

Telephone: 011-44-74-6211-3024

Email: ayodele.odutayo@bnc.ox.ac.uk

**Word Count (excluding methods):** 1930

**Abstract Count:** 203

**Keywords:** Acute Kidney Injury; Cardiovascular Mortality; Stroke; Acute Myocardial Infarction; Congestive Heart Failure

**ABSTRACT**

We conducted a systematic review and meta-analysis to assess whether acute kidney injury was associated with long-term cardiovascular disease. We included cohort studies that examined adults with and without acute kidney injury and reported a multivariable adjusted relative risk for the association between acute kidney injury and cardiovascular mortality, major cardiovascular events and disease specific events: congestive heart failure, acute myocardial infarction and stroke. Twenty-five studies involving 254,408 adults (55,150 with acute kidney injury) were included. Acute kidney injury was associated with an 86% and 38% increased risk of cardiovascular mortality and major cardiovascular events respectively (relative risk 1.86, 95% CI: 1.72-2.01 and 1.38, 1.22-1.55). For disease specific events, acute kidney injury was associated with a 58% increased risk of heart failure (1.58, 1.46-1.72) and a 40% increased risk of acute myocardial infarction (1.40, 1.23-1.59). The elevated risk of heart failure and acute myocardial infarction persisted in subgroup analyses based on acute kidney injury severity and the proportion of adults with baseline ischemic heart disease. Finally, acute kidney injury was associated with a 15% increased risk of stroke (1.15, 1.04-1.28). Acute kidney injury is associated with an elevated risk of cardiovascular mortality and major cardiovascular events, particularly heart failure and acute myocardial infarction.

## INTRODUCTION

Acute kidney injury (AKI) affects patients across multiple clinical settings and is a leading cause of morbidity and mortality.<sup>1-7</sup> AKI is also associated with increased hospital length of stay and excess healthcare spending.<sup>1</sup> Despite the established clinical importance of AKI, it is undiagnosed in up to 25% of cases in developed countries<sup>8</sup> and up to 75% of cases in developing countries.<sup>9</sup> Accordingly, guideline committees and public health authorities<sup>10,11</sup> have called for more research into the long-term prognostic implications of AKI as a key element of efforts to improve the diagnosis and management of AKI.

While the long-term risk of mortality, chronic kidney disease (CKD) and end-stage renal disease (ESRD) in adults with AKI are well established, it remains less clear whether AKI confers an additional long-term risk of cardiovascular disease. Existing studies have provided conflicting results<sup>12-14</sup> and prior systematic reviews have pooled unadjusted and adjusted relative risk estimates,<sup>15</sup> combined estimates for hard endpoints – congestive heart failure (CHF), acute myocardial infarction and stroke – and soft (revascularization) endpoints;<sup>15</sup> employed inadequate search strategies for cardiovascular outcomes;<sup>3</sup> or failed to identify more than two multivariable adjusted studies for acute myocardial infarction or stroke.<sup>3</sup> Furthermore, individual studies have been limited to specific patient populations – contrast induced AKI or peri-operative AKI – and it is unclear whether the prognostic importance of AKI varies by clinical factors such as baseline cardiovascular disease. A more comprehensive analysis is therefore needed.

Accordingly, we conducted a systematic review and meta-analysis to determine the long-term risk of cardiovascular mortality and cardiovascular disease associated with AKI. We also examined whether the risk conferred by AKI varied based on the severity of AKI, the recovery of renal function, the clinical setting and the presence of baseline cardiovascular disease or baseline CKD.

## **RESULTS**

In total, 1192 studies were reviewed and 1034 were excluded in the abstract screen largely because they did not report an association between AKI and an outcome of interest (Supplemental Figure 1). Among 158 full text articles that were reviewed, 133 were further excluded. Accordingly, 25 studies involving 254,408 adults were included in this meta-analysis. Of these individuals, 55,150 adults had AKI (Table 1). With respect to the risk of bias assessment, most studies examined a highly selected group of participants that were not representative of the general population. Accordingly, only one study obtained a full rating (four stars) for cohort selection. Fifteen studies received a full rating for comparability based on their multivariable adjustment and all studies received a full rating for outcome evaluation.

### **Cardiovascular Mortality and Major Cardiovascular Events**

Six studies,<sup>13,14,16-19</sup> involving 54,641 participants (18,284 with AKI) examined cardiovascular mortality as an outcome. The median follow-up was 2.6 years (inter-

quartile interval [IQI]: 2.0-3.4). The pooled relative risk for patients with AKI versus those without AKI was 1.86 (95% CI 1.72-2.01,  $I^2$ : 0%, Figure 1 and Supplementary Figure 2) and the relative risk did not vary by setting or proportion of adults with baseline CHF or CKD (Supplementary Table 2). There were insufficient studies to examine whether the risk varied by the remaining subgroups of interest.

Six studies,<sup>13,20-24</sup> involving 40,489 participants (11,824 with AKI) examined major cardiovascular events as an outcome. The median follow-up was 1.4 years (IQI 1.2-1.9). The pooled relative risk was 1.38 (95% CI 1.23-1.55,  $I^2$ : 47%, Figure 1 and Supplementary Figure 3) and it did not vary by the proportion of adults with baseline CHF, IHD or stroke (Supplementary Table 3). There were insufficient studies to examine whether the risk varied by the remaining subgroups of interest: AKI severity, recovery of renal function, and clinical setting.

### **Congestive Heart Failure**

Ten studies,<sup>12-14,25-31</sup> involving 90,251 participants (13,431 with AKI), examined CHF as an outcome. The median follow-up was 2.9 years (IQI 1.7-3.8). The pooled relative risk of CHF was 1.58 (95% CI 1.46-1.72, Figure 1 and Supplementary Figure 4). Heterogeneity was minimal ( $I^2$ : 10%). There was also an increase in the relative risk of long-term CHF with increasing severity of AKI ( $p=0.005$  for interaction, Table 2). In an exploratory analysis that was limited to studies that additionally adjusted for baseline left ventricular ejection fraction, the differential risk for CHF based on AKI severity was maintained (Table 2). Furthermore, the relative risk of CHF was modestly lower in

studies in which the majority of adults did not have baseline IHD (1.43, 95% CI 1.27-1.61) compared to studies where a large proportion of adults had baseline IHD (1.71, 95% CI 1.56-1.87,  $p=0.020$ ; Table 2). There were no differences based on the clinical setting, the recovery of renal function or the proportion of adults with baseline CHF, stroke or CKD (Table 2).

### **Acute Myocardial Infarction**

Ten studies,<sup>13,14,18,25,28,30,32-35</sup> involving 90,498 participants (12,091 with AKI), examined acute myocardial infarction as an outcome. The median follow-up was 2.3 years (IQI 1.7-2.9). The pooled relative risk of acute myocardial infarction was 1.40 (95% CI 1.23-1.59, Figure 1 and Supplementary Figure 5). Heterogeneity was moderate ( $I^2$ : 43%). The relative risk of long-term acute myocardial infarction did not vary by clinical setting (Table 3). Furthermore, there were no differences in the risk of long-term acute myocardial infarction based on AKI recovery nor based on the severity of AKI ( $p=0.12$  and  $0.64$  respectively, Table 3). Notably, there was no difference in the relative risk of acute myocardial infarction in subgroup analyses based on the proportion of adults with baseline cardiovascular disease or CKD (Table 3)

### **Stroke**

Five studies,<sup>13,25,28,36,37</sup> involving 72,347 participants (13,475 with AKI) examined stroke as an outcome. The median follow-up was 2.7 years (IQI 2.0-3.4). The pooled relative risk was 1.15 (95% CI 1.03-1.28,  $I^2$ : 0%, Figure 1 and Supplementary Figure 6). The relative risk of stroke did not vary between subgroups (Supplementary Table 4).

### **Sensitivity Analyses and Assessment of Heterogeneity**

There were minimal differences in the relative risk of long-term cardiovascular disease based on whether studies initiated follow up after discharge from hospital and thereby excluded inpatient events or based on the duration of follow up (data not shown). Heterogeneity was limited for all analyses and therefore not explored further. There was no funnel plot asymmetry (Supplementary Figures 7 & 8).

### **DISCUSSION**

This study is a comprehensive summary of the long-term risk of cardiovascular mortality and cardiovascular disease in adults with AKI. There were three major findings. First, we found that AKI was associated with an 86% increased risk of cardiovascular mortality and a 38% increased risk of major cardiovascular events. Second, AKI was associated with a 58% increased risk of CHF and a 40% increased risk of acute myocardial infarction and the association between AKI and these outcomes persisted robust in subgroup analyses. Third, AKI was associated with a 15% increased risk of stroke.

Our study is the first comprehensive summary of the long-term risk of cardiovascular mortality and cardiovascular disease in adults with AKI. We noted that

AKI was associated with cardiovascular mortality and major cardiovascular events and we also found robust associations for individual disease specific events.

In particular, we found that AKI was associated with a 58% increased risk of CHF. Furthermore, there was an association between the severity of AKI and the risk of CHF and this association persisted in studies that additionally adjusted for left ventricular ejection fraction. Although the risk of CHF was marginally lower in studies with a low proportion of adults with baseline IHD, the relative risk in both subgroups were well above the line of no effect; the threshold for which there may be no difference in the risk of CHF between adults with and without AKI. It is therefore unlikely that the association between AKI and CHF is due to residual confounding. Furthermore, we found that AKI was associated with a 40% increase in the relative risk of acute myocardial infarction. Although the risk of acute myocardial infarction did not increase with AKI severity, it is noteworthy that we combined the moderate and severe AKI categories, which may account for the absence of an association. Finally, the risk of acute myocardial infarction did not vary by the proportion of adults with baseline IHD.

To date, individual studies have focused on specific subgroups of patients (e.g. with established cardiovascular disease),<sup>13,28</sup> and it remained unclear whether their findings were generalizable to populations without cardiovascular disease. Our assessment of the consistency of relative risk estimates based on the proportion of adults with baseline cardiovascular disease adds credence to the prognostic importance of AKI



in populations with varying levels of cardiovascular risk. AKI may therefore be an important prognostic marker in patients with varying baseline cardiovascular risk.

Finally, with respect to stroke, we found that AKI was associated with a 15% relative risk increase. However, given the small effect size and the proximity of the lower confidence interval to the line of no effect, the relationship between AKI and stroke may also be due to residual confounding.

Taken together, the association between AKI, CHF and acute myocardial infarction should receive greater attention from clinicians, particularly as small changes in clinical practice can improve outcomes for adults with AKI. For instance, in a retrospective cohort study, investigators demonstrated that AKI was unrecognized in 24% of patients admitted to hospital in a one-month period.<sup>8</sup> The authors also identified important gaps in the management of these patients, including failure to withhold nephrotoxic drugs in 39% of patients.<sup>8</sup> With greater awareness and optimal care, the progression of AKI from mild to severe may be preventable, which could reduce long-term cardiovascular morbidity.

Our findings also highlight the need of longitudinal follow-up for adults with AKI. Prior studies have demonstrated decreased mortality in AKI survivors receiving nephrologist follow up.<sup>38</sup> However, it remains unclear what form of clinical care should be provided to these individuals, beyond the management of renal functional loss.<sup>38</sup> Our

findings highlight the need for screening and management of CHF and acute myocardial infarction in adults with AKI.

The following limitations should be considered when interpreting the findings of our study. First, we lacked individual patient data, which prevented us from using a common definition of AKI and from ensuring similar variables were adjusted for in regression models. This is particularly important as the definition of AKI varied between studies and there were insufficient studies to compare the risk estimates based on AKI definition. Thus, the data we report includes pooled populations of patients defined using the AKIN, RIFLE, and KDIGO and other grading systems and this should be considered when interpreting the findings of our study. Second, the majority of studies included in our analysis did not achieve the full rating for selection in our risk of bias assessment because they examined highly selected populations. Third, due to the small number of studies providing stratified data for our meta-analysis, our assessment of the association between the severity of AKI and cardiovascular outcomes was based on comparisons between mild AKI and the combination of moderate-to-severe AKI, as opposed to the familiar three AKI severity grades defined in the consensus criteria. This may account for the absence of any difference in the risk of cardiovascular disease based on the severity of AKI. Finally, our study can only demonstrate an association between AKI and cardiovascular disease and these associations cannot be deemed as causal in nature. Specifically, it could be that patients who develop AKI have conditions that put them at increased risk for cardiovascular events, rather than being at higher risk because of AKI itself.

Taken together, AKI is associated with an increased relative risk of long-term cardiovascular mortality, major cardiovascular events and particularly CHF and IHD. In contrast, the association with stroke may be due to residual confounding. Further research is required into interventions to improve long-term outcomes for adults with AKI.

## **METHODS**

This study was conducted in accordance with the Meta-analysis of Observational Studies in Epidemiology (MOOSE) guidelines and the PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) guidelines.<sup>39,40</sup>

### **Data Sources and Searchers**

We conducted a systematic search of MEDLINE and EMBASE (inception to March 2015). A qualified research librarian developed the search strategy and included but was not limited to search terms such as “acute kidney injury”, “acute renal failure”, “contrast induced nephropathy”, “congestive heart failure”, “myocardial infarction” and “stroke”. This was supplemented by a review of past meta-analyses,<sup>3-5,7,15</sup> a detailed review of references of included studies and citation tracking with Google Scholar.

Cohort studies that examined adults with and without AKI and reported a measure of relative risk for the association between AKI and cardiovascular disease (see below) were included. Studies were also required to include a minimum of 50 participants with

AKI and 50 participants without AKI with at least 6 months mean/median follow up.

Only studies reporting multivariable adjusted estimates were included and at a minimum, studies were required to adjust for age and sex. Unadjusted studies or studies that did not list variables included in their regression model were excluded. No language restrictions were applied.

### **Data Extraction and Quality Assessment**

Two reviewers independently reviewed titles and abstracts to assess studies for their inclusion. Four reviewers independently abstracted data using standardized forms. Where available, we abstracted information on general study characteristics (Study name or investigator's name; Recruitment Date (mid-point of the recruitment period); Mean follow-up duration; Year of publication of the primary findings); summary information about the studied population that was clinically relevant to the outcomes specified for our study (definition of AKI; number of participants with and without AKI; mean age; number of men; number of adults with baseline CHF, IHD and stroke). We also extracted information on the following outcomes: cardiovascular mortality, major cardiovascular events (a composite of cardiovascular death, stroke, acute myocardial infarction and CHF) and disease specific events: CHF, acute myocardial infarction and stroke.

Relative risk estimates and associated 95% confidence intervals (CI) for the association between AKI and the aforementioned study outcomes were abstracted. The list of variables included in the multivariable regression model was also abstracted.

A risk of bias assessment was performed using the Newcastle-Ottawa Scale, which assesses studies on three broad domains: the selection of participants for study groups; the comparability of study groups; and the ascertainment of the outcome. A star rating system is used to identify studies that are at low risk of bias and the maximum numbers of stars achievable are: Selection (4 stars), Comparability (2 stars), and Outcome (cohort studies only; 3 stars). We applied strict criteria to assess comparability (of individuals with or without AKI) based on which variables were included in the multivariable models. To receive one star for comparability, studies were required to have adjusted for age, sex and at least one cardiovascular risk factor (hypertension, diabetes, smoking, cholesterol and chronic kidney disease) and a baseline history of cardiovascular disease (stroke or IHD) if applicable. To receive two stars, studies were additionally required to have adjusted for baseline serum creatinine, estimated glomerular filtration rate (eGFR) or history of CKD.

### **Subgroup Analyses, Sensitivity Analyses**

We performed four subgroup analyses in which included studies were grouped by the severity of AKI (mild versus moderate-to-severe), the recovery of renal function (complete versus partial or no renal recovery), clinical setting and baseline cardiovascular disease and CKD. Subgroup analyses were only performed when at least 2 studies could be obtained for each stratum.

For the analysis of severity, mild AKI was defined as AKIN 1 according to the AKIN (Acute Kidney Injury Network) classification<sup>41</sup> or “Risk” according to the RIFLE (Risk, Injury, Failure, Loss of kidney function, and End-stage kidney disease) classification<sup>42</sup> or Stage 1 according to the KDIGO (Kidney Disease Improving Global Outcomes) classification. Moderate to severe AKI was defined as either AKIN 2 or 3, RIFLE “Injury” or “Failure”, KDIGO Stage 2 or 3 or AKI requiring dialysis.<sup>43</sup> For the analysis of recovery, complete recovery was defined as a return of renal function to below the study-defined threshold for AKI and partial or no recovery was defined as the absence of this level of improvement in kidney function. Finally, with respect to setting, we specifically examined contrast-induced AKI and AKI in the peri-operative or ICU setting. For the subgroup analysis by baseline cardiovascular disease, we calculated the proportion of adults with a past history of CHF, IHD, stroke and CKD at cohort entry and separated studies halves based on this proportion. For studies which were limited to adults with acute myocardial infarction or adults receiving percutaneous coronary interventions or coronary artery bypass grafting, all study participants were identified as having baseline IHD. We compared the prognostic importance of AKI in the two groups.

Finally, we performed two sensitivity analyses for the overall pooled estimates for each outcome. First, studies were grouped based on whether follow-up for cardiovascular events commenced immediately after entry into the cohort or after a period of time (e.g. after hospital discharge). Second, studies were divided into halves based on average length of follow up.

### **Data Synthesis, Analysis and Planned Assessment of Heterogeneity**

Overall summary estimates for each of the aforementioned outcomes were computed using a random effects meta-analysis. Where studies reported multiple relative risk estimates for specific subgroups without reporting an overall summary estimate, the subgroup estimates were pooled using fixed effects meta-analysis to ensure a single relative risk was obtained for each study prior to the overall analysis. Heterogeneity was assessed using the  $I^2$  statistic. Results are presented graphically using forest plots.

In accordance with Cochrane collaboration guidelines, when at least 10 studies were included in meta-analysis we performed assessment for publication bias by visual inspection of funnel plots, supported by Egger's test.<sup>44</sup> Where an asymmetrical funnel plot was identified, the Copas selection model was used to account for small study effect bias.<sup>45</sup>

When high heterogeneity was observed ( $I^2 > 60\%$ ), we planned to assess whether heterogeneity was reduced by subgroup analyses and in sensitivity analyses. If

heterogeneity was not reduced to acceptable levels ( $I^2 < 60\%$ ), we additionally assessed if age accounted for heterogeneity by performing meta-regression based on the average age (categorical variable divided into halves). All analyses were performed using R Statistical Software (Version 3.0).

**Ethics Approval and Funding Source**

Ethics approval was not required. This study was unfunded.



**ACKNOWLEDGEMENTS**

None

**CONFLICT OF INTEREST**

None

**FUNDING**

This study received no external funding.

## REFERENCES

1. Chertow GM, Burdick E, Honour M, Bonventre JV, Bates DW: Acute kidney injury, mortality, length of stay, and costs in hospitalized patients. *J Am Soc Nephrol* 16: 3365–3370, 2005
2. Coca SG, Bauling P, Schiffner T, Howard CS, Teitelbaum I, Parikh CR: Contribution of acute kidney injury toward morbidity and mortality in burns: a contemporary analysis. *Am J Kidney Dis* 49: 517–523, 2007
3. Coca SG, Yusuf B, Shlipak MG, Garg AX, Parikh CR: Long-term risk of mortality and other adverse outcomes after acute kidney injury: a systematic review and meta-analysis. *Am J Kidney Dis* 53: 961–973, 2009
4. Coca SG, Singanamala S, Parikh CR: Chronic kidney disease after acute kidney injury: a systematic review and meta-analysis. *Kidney Int* 81: 442–448, 2012
5. Corredor C, Thomson R, Al-Subaie N: Long Term Consequences of Acute Kidney Injury Following Cardiac Surgery: A Systematic Review and Meta-Analysis. *J. Cardiothorac. Vasc. Anesth.* 30: 69–75, 2016
6. Pickering JW, James MT, Palmer SC: Acute kidney injury and prognosis after cardiopulmonary bypass: a meta-analysis of cohort studies. *Am J Kidney Dis* 65: 283–293, 2015
7. Gargiulo G, Sannino A, Capodanno D, Perrino C, Capranzano P, Barbanti M, Stabile E, Trimarco B, Tamburino C, Esposito G: Impact of postoperative acute kidney injury on clinical outcomes after transcatheter aortic valve implantation: A meta-analysis of 5,971 patients. *Cathet. Cardiovasc. Intervent.* 2015
8. Aitken E, Carruthers C, Gall L, Kerr L, Geddes C, Kingsmore D: Acute kidney injury: outcomes and quality of care. *QJM* 106: 323–332, 2013
9. Yang L, Xing G, Wang L, Wu Y, Li S, Xu G, He Q, Chen J, Chen M, Liu X, Zhu Z, Yang L, Lian X, Ding F, Li Y, Wang H, Wang J, Wang R, Mei C, Xu J, Li R, Cao J, Zhang L, Wang Y, Xu J, Bao B, Liu B, Chen H, Li S, Zha Y, Luo Q, Chen D, Shen Y, Liao Y, Zhang Z, Wang X, Zhang K, Liu L, Mao P, Guo C, Li J, Wang Z, Bai S, Shi S, Wang Y, Wang J, Liu Z, Wang F, Huang D, Wang S, Ge S, Shen Q, Zhang P, Wu L, Pan M, Zou X, Zhu P, Zhao J, Zhou M, Yang L, Hu W, Wang J, Liu B, Zhang T, Han J, Wen T, Zhao M, Wang H, ISN AKF 0by25 China Consortiums: Acute kidney injury in China: a cross-sectional survey. *Lancet* 386: 1465–1471, 2015
10. Ftouh S, Thomas M, Acute Kidney Injury Guideline Development Group: Acute kidney injury: summary of NICE guidance. *BMJ*. 347: f4930, 2013
11. Mehta RL, Cerdá J, Burdmann EA, Tonelli M, García-García G, Jha V,

- Susantitaphong P, Rocco M, Vanholder R, Sever MS, Cruz D, Jaber B, Lameire NH, Lombardi R, Lewington A, Feehally J, Finkelstein F, Levin N, Pannu N, Thomas B, Aronoff-Spencer E, Remuzzi G: International Society of Nephrology's Oby25 initiative for acute kidney injury (zero preventable deaths by 2025): a human rights case for nephrology. *Lancet* 385: 2616–2643, 2015
12. Shirakabe A, Hata N, Kobayashi N, Shinada T, Tomita K, Tsurumi M, Matsushita M, Okazaki H, Yamamoto Y, Yokoyama S, Asai K, Mizuno K: Long-term prognostic impact after acute kidney injury in patients with acute heart failure. *Int Heart J* 53: 313–319, 2012
  13. Mielniczuk LM, Pfeffer MA, Lewis EF, Blazing MA, de Lemos JA, Mohanavelu S, Rouleau J, Fox K, Pedersen TR, Califf RM: Acute decline in renal function, inflammation, and cardiovascular risk after an acute coronary syndrome. *Clinical journal of the American Society of Nephrology : CJASN* 4: 1811–1817, 2009
  14. Jose P, Skali H, Anavekar N, Tomson C, Krumholz HM, Rouleau JL, Moya L, Pfeffer MA, Solomon SD: Increase in creatinine and cardiovascular risk in patients with systolic dysfunction after myocardial infarction. *J Am Soc Nephrol* 17: 2886–2891, 2006
  15. James MT, Samuel SM, Manning MA, Tonelli M, Ghali WA, Faris P, Knudtson ML, Pannu N, Hemmelgarn BR: Contrast-induced acute kidney injury and risk of adverse clinical outcomes after coronary angiography: a systematic review and meta-analysis. *Circ Cardiovasc Interv* 6: 37–43, 2013
  16. Ozrazgat-Baslanti T, Thottakkara P, Huber M, Berg K, Gravenstein N, Tighe P, Lipori G, Segal MS, Hobson C, Bihorac A: Acute and Chronic Kidney Disease and Cardiovascular Mortality After Major Surgery. *Annals of Surgery* 2016
  17. Uyarel H, Cam N, Ergelen M, Akkaya E, Ayhan E, Isik T, Cicek G, Gunaydin ZY, Osmonov D, Gul M, Demirci D, Guney MR, Ozturk R, Yekeler I: Contrast-induced nephropathy in patients undergoing primary angioplasty for acute myocardial infarction: incidence, a simple risk score, and prognosis. *Archives of Medical Sciences* 4: 550–558, 2009
  18. Kim J-H, Yang JH, Choi S-H, Song YB, Hahn J-Y, Choi J-H, Lee SH, Gwon H-C: Predictors of outcomes of contrast-induced acute kidney injury after percutaneous coronary intervention in patients with chronic kidney disease. *Am J Cardiol* 114: 1830–1835, 2014
  19. Narula A, Mehran R, Weisz G, Dangas GD, Yu J, G  n  reux P, Nikolsky E, Brener SJ, Witzenbichler B, Guagliumi G, Clark AE, Fahy M, Xu Ke, Brodie BR, Stone GW: Contrast-induced acute kidney injury after primary percutaneous coronary intervention: results from the HORIZONS-AMI substudy. *Eur Heart J* 35: 1533–1540, 2014
  20. Blair JEA, Pang PS, Schrier RW, Metra M, Traver B, Cook T, Campia U,

- Ambrosy A, Burnett JC, Grinfeld L, Maggioni AP, Swedberg K, Udelson JE, Zannad F, Konstam MA, Gheorghiade M, EVEREST Investigators: Changes in renal function during hospitalization and soon after discharge in patients admitted for worsening heart failure in the placebo group of the EVEREST trial. *Eur Heart J* 32: 2563–2572, 2011
21. Chawla LS, Amdur RL, Shaw AD, Faselis C, Palant CE, Kimmel PL: Association between AKI and long-term renal and cardiovascular outcomes in United States veterans. *Clinical journal of the American Society of Nephrology : CJASN* 9: 448–456, 2014
  22. Hansen MK, Gammelager H, Jacobsen C-J, Hjortdal VE, Layton JB, Rasmussen BS, Andreasen JJ, Johnsen SP, Christiansen CF: Acute Kidney Injury and Long-term Risk of Cardiovascular Events After Cardiac Surgery: A Population-Based Cohort Study. *J. Cardiothorac. Vasc. Anesth.* 29: 617–625, 2015
  23. Metra M, Nodari S, Parrinello G, Bordonali T, Bugatti S, Danesi R, Fontanella B, Lombardi C, Milani P, Verzura G, Cotter G, Dittrich H, Massie BM, Dei Cas L: Worsening renal function in patients hospitalised for acute heart failure: clinical implications and prognostic significance. *Eur. J. Heart Fail.* 10: 188–195, 2008
  24. Watabe H, Sato A, Hoshi T, Takeyasu N, Abe D, Akiyama D, Kakefuda Y, Nishina H, Noguchi Y, Aonuma K: Association of contrast-induced acute kidney injury with long-term cardiovascular events in acute coronary syndrome patients with chronic kidney disease undergoing emergent percutaneous coronary intervention. *Int J Cardiol* 174: 57–63, 2014
  25. Gammelager H, Christiansen CF, Johansen MB, Tønnesen E, Jespersen B, Sørensen HT: Three-year risk of cardiovascular disease among intensive care patients with acute kidney injury: a population-based cohort study. *Crit Care* 18: 492, 2014
  26. Damman K, Jaarsma T, Voors AA, Navis G, Hillege HL, van Veldhuisen DJ, COACH investigators: Both in- and out-hospital worsening of renal function predict outcome in patients with heart failure: results from the Coordinating Study Evaluating Outcome of Advising and Counseling in Heart Failure (COACH). *Eur. J. Heart Fail.* 11: 847–854, 2009
  27. Damman K, Perez AC, Anand IS, Komajda M, McKelvie RS, Zile MR, Massie B, Carson PE, McMurray JJV: Worsening renal function and outcome in heart failure patients with preserved ejection fraction and the impact of angiotensin receptor blocker treatment. *J Am Coll Cardiol* 64: 1106–1113, 2014
  28. James MT, Ghali WA, Knudtson ML, Ravani P, Tonelli M, Faris P, Pannu N, Manns BJ, Klarenbach SW, Hemmelgarn BR, Alberta Provincial Project for Outcome Assessment in Coronary Heart Disease (APPROACH) Investigators: Associations between acute kidney injury and cardiovascular and renal outcomes after coronary angiography. *Circulation* 123: 409–416, 2011

29. Choi AI, Li Y, Parikh C, Volberding PA, Shlipak MG: Long-term clinical consequences of acute kidney injury in the HIV-infected. *Kidney Int* 78: 478–485, 2010
30. Goldberg A, Kogan E, Hammerman H, Markiewicz W, Aronson D: The impact of transient and persistent acute kidney injury on long-term outcomes after acute myocardial infarction. *Kidney Int* 76: 900–906, 2009
31. Olsson D, Sartipy U, Braunschweig F, Holzmann MJ: Acute kidney injury following coronary artery bypass surgery and long-term risk of heart failure. *Circ Heart Fail* 6: 83–90, 2013
32. Tsai M-L, Mao C-T, Chen D-Y, Hsieh I-C, Wen M-S, Chen T-H: Short- and long-term major cardiovascular adverse events in carotid artery interventions: a nationwide population-based cohort study in Taiwan. *PLoS ONE* 10: e0121016, 2015
33. Giacoppo D, Madhavan MV, Baber U, Warren J, Bansilal S, Witzenbichler B, Dangas GD, Kirtane AJ, Xu Ke, Kornowski R, Brener SJ, G  n  reux P, Stone GW, Mehran R: Impact of Contrast-Induced Acute Kidney Injury After Percutaneous Coronary Intervention on Short- and Long-Term Outcomes: Pooled Analysis From the HORIZONS-AMI and ACUTITY Trials. *Circ Cardiovasc Interv* 8: e002475, 2015
34. Ryd  n L, Ahnve S, Bell M, Hammar N, Ivert T, Sartipy U, Holzmann MJ: Acute kidney injury after coronary artery bypass grafting and long-term risk of myocardial infarction and death. *Int J Cardiol* 172: 190–195, 2014
35. Lindsay J, Apple S, Pinnow EE, Gevorkian N, Gruberg L, Satler LF, Pichard AD, Kent KM, Suddath W, Waksman R: Percutaneous coronary intervention-associated nephropathy foreshadows increased risk of late adverse events in patients with normal baseline serum creatinine. *Catheter Cardiovasc Interv* 59: 338–343, 2003
36. Holzmann MJ, Ryd  n L, Sartipy U: Acute kidney injury and long-term risk of stroke after coronary artery bypass surgery. *Int J Cardiol* 168: 5405–5410, 2013
37. Wu V-C, Wu C-H, Huang T-M, Wang C-Y, Lai C-F, Shiao C-C, Chang C-H, Lin S-L, Chen Y-Y, Chen Y-M, Chu T-S, Chiang W-C, Wu K-D, Tsai P-R, Chen L, Ko W-J, NSARF Group: Long-term risk of coronary events after AKI. *J Am Soc Nephrol* 25: 595–605, 2014
38. Harel Z, Wald R, Bargman JM, Mamdani M, Etchells E, Garg AX, Ray JG, Luo J, Li P, Quinn RR, Forster A, Perl J, Bell CM: Nephrologist follow-up improves all-cause mortality of severe acute kidney injury survivors. *Kidney Int* 83: 901–908, 2013
39. Stroup DF, Berlin JA, Morton SC, Olkin I, Williamson GD, Rennie D, Moher D,

Becker BJ, Sipe TA, Thacker SB: Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. pp 2008–2012, 2000

40. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group: Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med.* 151: 264–9– W64, 2009
41. Mehta R, Kellum J, Shah S, Molitoris B, Ronco C, Warnock D, Levin A, Network TAKI: Acute Kidney Injury Network: report of an initiative to improve outcomes in acute kidney injury. *Critical Care* 2008 12:R47 11: R31, 2007
42. Bellomo R, Ronco C, Kellum JA, Mehta RL, Palevsky P, Acute Dialysis Quality Initiative workgroup: Acute renal failure - definition, outcome measures, animal models, fluid therapy and information technology needs: the Second International Consensus Conference of the Acute Dialysis Quality Initiative (ADQI) Group. *Crit Care* 8: R204–12, 2004
43. Kidney Disease: Improving Global Outcomes (KDIGO): KDIGO clinical practice guidelines for the prevention, diagnosis, evaluation, and treatment of hepatitis C in chronic kidney disease. *Kidney Int. Suppl.* S1–99, 2008
44. Egger M, Davey Smith G, Schneider M, Minder C: Bias in meta-analysis detected by a simple, graphical test. *BMJ* 315: 629–634, 1997
45. Schwarzer G, Carpenter J, Rücker G: Empirical evaluation suggests Copas selection model preferable to trim-and-fill method for selection bias in meta-analysis. *Journal of Clinical Epidemiology* 63: 282–288, 2010



## FIGURE LEGENDS

### **Figure 1: Association Between Acute Kidney Injury and Cardiovascular Mortality and Cardiovascular Events**

CVD is cardiovascular; MACE is major adverse cardiovascular events; CHF is congestive heart failure; and AMI is acute myocardial infarction. Horizontal lines indicate 95% confidence intervals. Number of events is missing for MACE because it is not reported in all studies. The solid vertical line represents the line of no effect ( $RR=1.0$ ), wherein there is no difference in the risk of the outcome of interest between adults with and without acute kidney injury.



**Table 1: List of Included Studies**

<b>Study Author, Year</b>	<b>Outcome Reported</b>	<b>AKI Definition</b>	<b>Setting</b>	<b>Subjects (AKI)</b>	<b>CHF (%)</b>	<b>IHD (%)</b>	<b>Stroke (%)</b>	<b>CKD (%)</b>
Blair, 2011 <sup>20</sup>	MACE	Absolute increase in SCr $\geq 0.3$ mg/dL	Other	2021 (279)	2021 (100)	1425 (71)	NA	553 (27)
Chawla, 2014 <sup>21</sup>	MACE	AKIN Criteria: Peak SCr during admission at least 1.5–1.9 times higher than baseline or an increase of $\geq 0.3$ mg/dl (stage 1), an increase of at least 2.0–2.9 times baseline (stage 2), or an increase of $\geq 3.0$ times baseline or $\geq 4.0$ mg/dl (stage 3)	Other	28554 (9633)	2829 (10)	28554 (100)	653 (2)	NA
Choi, 2010 <sup>29</sup>	CHF	AKIN Criteria: AKIN 1 - An SCr increase of $\geq 0.3$ mg/dl, or a relative increase of 150-200%; AKIN 2 - A relative increase of 200-300%; AKIN 3 - A relative increase $>300\%$ , or in individuals with an SCr of $\geq 4.0$ mg/dl, an SCr increase of 0.5 mg/dl from the baseline	Other	17325 (3060)	358 (2)	1186 (7)	NA	1281 (7)
Damman, 2009 <sup>26</sup>	CHF	Absolute increase in serum creatinine $>26.5$ mmol/L ( $>0.3$ mg/dL) in combination with $>25\%$ increase in serum creatinine between two time points	Other	869 (106)	869 (100)	364 (42)	80 (9)	NA
Damman, 2014 <sup>27</sup>	CHF	Absolute increase in SCr of $\geq 26.5$ mmol/l ( $\geq 0.3$ mg/dl), together with a relative increase in SCr of $\geq 25\%$ between baseline and 8 weeks	Other	3595 (229)	3595 (100)	884 (25)	321 (9)	NA
Gammelager, 2014 <sup>25</sup>	Stroke, IHD, CHF	KDIGO Criteria - Stage 1: 1.5-1.9 times increase versus baseline or $\geq 0.3$ mg/dl increase within 48 hours; Stage 2: 2.0-2.9 times baseline; $\geq 3.0$ times baseline or increase in serum creatinine to $\geq 4.0$ mg/dl or initiation of renal replacement therapy	Peri-Operative or ICU	21556 (4792)	0	2620 (12)	0	1903 (9)
Giacoppo, 2015 <sup>33</sup>	IHD	Absolute increase of SCr $\geq 0.5$ mg/dL or relative increase $\geq 25\%$ from baseline value within the first 72 hours of intervention	CIN	9512 (1212)	NA	9512 (100)	376 (4)	1702 (18)
Goldberg, 2009 <sup>30</sup>	IHD, CHF	Transient mild AK - In this group, peak SCr during hospital course was $\geq 0.3$ – $0.49$ mg/dl higher than admission SCr, but pre-discharge SCr returned to a level that was $<0.3$ mg/dl higher than admission creatinine.); Persistent mild AKI –	CIN	1957 (294)	368 (19)	1957 (100)	NA	338 (17)

		where SCr did not return to <0.3 mg/dl; Transient moderate/severe AKI - peak SCr $\geq$ 0.5mg/dl higher than admission creatinine, but pre-discharge returned <0.5 mg/dl higher than admission SCr; Moderate/severe AKI – where SCr did not return to <0.5 mg/dl						
Hansen, 2015 <sup>22</sup>	MACE	AKIN Criteria - Stage 1 - Increase in SCr of $\geq$ 0.3 mg/dL (26.5 $\mu$ mol/L) within 48 hours OR increase in SCr of 1.5-1.9 times baseline; Stage 2 - Increase in serum creatinine of 2.0-2.9 times baseline; Stage 3 - Increase in serum creatinine of 3.0 times baseline OR increase in serum creatinine to $\geq$ 4.0 mg/dL (353.6 $\mu$ mol/L) and satisfaction of at least stage 1 OR initiation of renal replacement therapy.	Peri-Operative or ICU	4742 (1457)	717 (15)	2651 (56)	282 (6)	823 (17)
Holzmann, 2013 <sup>36</sup>	Stroke	Postoperative SCr that was the highest value during the entire postoperative period. Absolute increases in SCr: stage 1, 0.3–0.5 mg/dL (26–44 $\mu$ mol/L); stage 2, 0.5–1 mg/dL (44–88 $\mu$ mol/L); and stage 3, $\geq$ 1 mg/dL ( $\geq$ 88 $\mu$ mol/L)	Peri-Operative or ICU	23584 (2764)	943 (4)	23584 (100)	0	4481 (19)
James, 2011 <sup>28</sup>	Stroke, IHD, CHF	AKIN Criteria: AKI stage 1, $\geq$ 0.3 mg/dL absolute or 1.5- to 2.0-fold relative increase in SCr; AKI stage 2, >2- to 3-fold increase in SCr; AKI stage 3, >3-fold increase in serum creatinine or SCr $\geq$ 4.0 mg/dL with an acute rise of >0.5 mg/dL	CIN	14782 (1420)	2253 (15)	10617 (72)	1096 (7)	3543 (24)
Jose, 2006 <sup>14</sup>	CVD Mortality, IHD, CHF	Increase in SCr >0.3 mg/dl from baseline	Other	1854 (223)	108 (6)	1854 (100)	NA	610 (33)
Kim, 2014 <sup>16</sup>	CVD Mortality, IHD	SCr $\geq$ 25% or $\geq$ 0.5 mg/dl from baseline within 72 hours after contrast exposure.	CIN	297 (55)	NA	297 (100)	31 (10)	297 (100)
Lindsay, 2003 <sup>35</sup>	IHD	An increase in the SCr of $\geq$ 50% from baseline after the intervention	CIN	5967 (208)	NA	5967 (100)	539 (9)	0
Metra, 2008 <sup>23</sup>	MACE	A $\geq$ 0.3 mg/dL and a $\geq$ 25% increase in SCr from admission	Other	318 (107)	318 (100)	173 (54)	22 (7)	NA
Mielniczuk, 2009 <sup>13</sup>	CVD Mortality, Stroke,	RIFLE Criteria: Decrease in eGFR of $\geq$ 25% compared with baseline	Other	3795 (184)	182 (5)	3795 (100)	223 (6)	NA

	IHD, CHF							
Narula, 2014 <sup>19</sup>	CVD Mortality	Increase in SCr of $\geq 0.5$ mg/dl or a 25% relative rise in creatinine, within 48 h after contrast exposure	CIN	676 (338)	75 (11)	676 (100)	NA	138 (20)
Olsson, 2013 <sup>31</sup>	CHF	Increase in postoperative SCr values: stage 1 - 0.3 to $<0.5$ mg/dL (26 to $<44$ $\mu\text{mol/L}$ ); stage 2 - 0.5 to $<1$ mg/dL (44 to $<88$ $\mu\text{mol/L}$ ); and stage 3 - $\geq 1$ mg/dL ( $\geq 88$ $\mu\text{mol/L}$ )	Peri-Operative or ICU	24018 (2779)	0	24018 (100)	1153 (5)	NA
Ozrazgat-Baslanti, 2016 <sup>16</sup>	CVD Mortality	KDIGO Criteria: $\geq 50\%$ or 0.3 mg/dL increase in serum creatinine relative to the reference value and stratified AKI severity based on the maximum change in serum creatinine during hospitalization	Peri-Operative or ICU	45498 (16854)	2915 (6)	2727 (6)	3731 (8)	0
Ryden, 2014 <sup>34</sup>	IHD	Postoperative increase in SCr: stage 1 - 0.3 to 0.5 mg/dL (26 to 44 $\mu\text{mol/L}$ ); stage 2 - $>0.5$ to 1 mg/dL (44 to 88 $\mu\text{mol/L}$ ) and stage 3 - $\geq 1.0$ mg/dL ( $\geq 88$ $\mu\text{mol/L}$ )	Peri-Operative or ICU	27929 (3617)	1117 (4)	27929 (100)	1396 (5)	5586 (20)
Shirakabe, 2012 <sup>12</sup>	CHF	RIFLE Criteria: Class R - Increase in serum creatinine $\geq 1.5 \times$ baseline; Class I - Increase in serum creatinine $\geq 2.0 \times$ baseline; Class F - Increase in serum creatinine $\geq 3.0 \times$ baseline or $\geq 4$ mg/dL with an acute rise $> 0.5$ mg/dL	Other	500 (344)	500 (100)	201 (40)	NA	NA
Tsai, 2015 <sup>32</sup>	IHD	ICD-9 Code	Peri-Operative or ICU	2849 (86)	275 (10)	281 (10)	2698 (95)	164 (6)
Uyarel, 2009 <sup>17</sup>	CVD Mortality	An increase in SCr level $\geq 0.5$ mg/dl or $\geq 25\%$ from baseline (admission) within 72 h of radiocontrast administration	CIN	2521 (630)	154 (6)	2521 (100)	NA	NA
Watabe, 2014 <sup>24</sup>	MACE	A $\geq 0.5$ mg/dL and/or $\geq 25\%$ absolute increase in SCr from baseline	CIN	1059 (164)	151 (14)	1059 (100)	90 (8)	1059 (100)
Wu, 2014 <sup>37</sup>	Stroke	RIFLE Criteria: AKI was defined according to the RIFLE (risk, injury, failure, loss, and end stage) criteria	Other	8630 (4315)	1011 (12)	1305 (15)	0	1029 (12)

SCr is serum creatinine.

**Table 2: Subgroup Analysis for the Association Between Acute Kidney Injury and Congestive Heart Failure**

Stratification	Number of Studies	Setting*	Relative Risk	95% CI	I <sup>2</sup> (%)	P-value
<b>CHF – Setting</b>						
CIN	2	-	1.69	1.44-1.99	0	0.43
Peri-operative	6	-	1.57	1.23-2.01	83	
Other	2	-	1.46	1.25-1.70	0	
<b>CHF – Recovery</b>						
Complete Recovery	3	CIN (1) Peri-op (1) Other (1)	1.41	1.22-1.62	0	0.21
Partial or No Recovery	3	CIN (1) Peri-op (1) Other (1)	1.65	1.35-2.03	0	
<b>CHF – Severity</b>						
Mild AKI	6	CIN (2) Peri-op (2) Other (2)	1.34	1.11-1.61	61	0.005
Moderate to Severe AKI	6	CIN (2) Peri-op (2) Other (2)	1.94	1.62-2.31	49	
<b>CHF – Severity (adjusted for LVEF)</b>						
Mild AKI	3	CIN (2) Peri-op (1) Other (0)	1.57	1.37-1.79	0	0.032
Moderate to Severe AKI	3	CIN (2) Peri-op (1) Other (0)	1.93	1.69-2.20	0	
<b>CHF – Median Proportion Baseline CHF (min-max)</b>						
[0.02, 0-0.06]	5	CIN (0) Peri-op (2) Other (3)	1.56	1.35-1.80	44	0.91
[1, 0.15-1]	5	CIN (2) Peri-op (0) Other (3)	1.58	1.38-1.80	0	
<b>CHF – Median Proportion Baseline IHD (min-max)</b>						
[0.25, 0.07-0.42]	5	CIN (0) Peri-op (1) Other (4)	1.43	1.27-1.61	0	0.020
[1, 0.72-1]	5	CIN (2)	1.71	1.56-1.87	0	

		Peri-op (1) Other (2)				
<b>CHF – Median Proportion Baseline Stroke (min-max)</b>						
[0.05, 0-0.06]	3	CIN (0) Peri-op (2) Other (1)	1.55	1.25-1.92	68	0.83
[0.09, 0.07-0.09]	3	CIN (1) Peri-op (0) Other (2)	1.60	1.35-1.88	0	
<b>CHF – Median Proportion Baseline CKD (min-max)</b>						
[0.09, 0.07-0.17]	3	CIN (1) Peri-op (1) Other (1)	1.54	1.32-1.79	30	0.80
[0.28, 0.24-0.33]	2	CIN (1) Peri-op (0) Other (1)	1.59	1.32-1.91	0	

\*number of studies is provided in brackets. CVD is cardiovascular, IHD is ischemic heart disease, CHF is congestive heart failure, Peri-op is peri-operative or ICU, LVEF is left ventricular ejection fraction and CKD is chronic kidney disease.

**Table 3: Subgroup Analysis for the Association Between Acute Kidney Injury and Acute Myocardial Infarction**

Stratification	Number of Studies	Setting*	Relative Risk	95% CI	I <sup>2</sup> (%)	P-value
<b>AMI – Setting</b>						
CIN	5	-	1.43	1.18-1.75	48	0.93
Peri-operative	3	-	1.39	1.08-1.77	47	
Other	2	-	1.31	0.82-2.09	55	
<b>AMI – Recovery</b>						
Complete Recovery	2	CIN (1) Peri-op (1) Other (0)	1.14	0.87-1.48	0	0.12
Partial or No Recovery	2	CIN (1) Peri-op (1) Other (0)	1.55	1.16-2.07	0	
<b>AMI – Severity</b>						
Mild AKI	4	CIN (2) Peri-op (2) Other (0)	1.45	1.28-1.65	18	0.64
Moderate to Severe AKI	4	CIN (2) Peri-op (2) Other (0)	1.53	1.27-1.84	0	
<b>AMI – Median Proportion Baseline CHF (min-max)</b>						
[0.04; 0-0.06]	4	CIN (0) Peri-op (2) Other (2)	1.42	1.19-1.69	38	0.91
[0.15; 0.10-0.17]	3	CIN (2) Peri-op (1) Other (0)	1.40	1.17-1.67	0	
<b>AMI – Median Proportion Baseline IHD (min-max)</b>						
[0.12; 0.10-0.72]	3	CIN (1) Peri-op (2) Other (0)	1.32	1.10-1.57	2	0.44
[1; 1-1]	7	CIN (4) Peri-op (1) Other (2)	1.45	1.23-1.71	51	
<b>AMI – Median Proportion Baseline Stroke (min-max)</b>						
[0.04; 0-0.06]	4	CIN (1) Peri-op (2) Other (1)	1.37	1.14-1.65	62	0.49
[0.10; 0.07-0.95]	4	CIN (3)	1.59	1.08-2.34	47	

		Peri-op (1) Other (0)				
<b>AMI – Median Proportion Baseline CKD (min-max)</b>						
[8.8; 0-0.18]	5	CIN (3) Peri-op (2) Other (0)	1.32	1.09-1.61	43	0.45
[0.28; 0.20-1]	4	CIN (2) Peri-op (1) Other (1)	1.47	1.23-1.75	37	

\*number of studies is provided in brackets. CVD is cardiovascular, AMI is acute myocardial infarction, IHD is ischemic heart disease, CHF is congestive heart failure, CKD is chronic kidney disease and Peri-op is peri-operative or ICU.