

CKM, mixing and CP violation results at LHCb

Tommaso Pajero^{a,1}

^a*University of Oxford,*

Denys Wilkinson Building, Keble Road, OX1 3RH, Oxford, United Kingdom

E-mail: tommaso.pajero@physics.ox.ac.uk

We review the most precise tests of the CKM model of flavour physics performed by the LHCb collaboration during 2021. These include world best measurements of both time-dependent and time-independent CP violation in charm decays, the first observation of a nonzero mass difference of the neutral charm-meson eigenstates, and the most precise determination to date of the mass difference between the B_s^0 eigenstates. A new simultaneous combination of measurements of the CKM angle γ and of charm decays allows to improve the precision on the decay-width difference of the neutral charm-meson eigenstates by a factor of 2 with respect to the combination of charm results only.

*** *Particles and Nuclei International Conference – PANIC2021* ***

*** *5–10 September, 2021* ***

*** *Online* ***

¹On behalf of the LHCb collaboration.

1. Introduction

The LHCb experiment has established itself as a main player in the study of beauty and charm hadrons, thanks to their large production cross-sections at the LHC and to its dedicated design [1]. In the following sections we review the most precise tests of the CKM model performed by the LHCb collaboration during 2021. Throughout the text, the first quoted uncertainties are statistical and the second are systematic.

2. Mixing and CP violation in charm decays

Two years after the first observation of CP violation in charm decays through the difference of the time-integrated CP asymmetries of $D^0 \rightarrow K^+K^-$ and $D^0 \rightarrow \pi^+\pi^-$ decays [2], agreement on the dynamical origin of this phenomenon has not been achieved yet [3–7]. Measurements of CP violation in further decay channels could provide insights on the size of the QCD nonperturbative effects at play. The LHCb collaboration has recently measured the CP asymmetries of $D^0 \rightarrow K_S^0 K_S^0$ [8] and $D_{(s)}^+ \rightarrow h^+ h^0$ decays [9], where h^+ stands for a π^+ or K^+ meson, and h^0 for a π^0 or η meson. The former has been proposed as a promising channel to measure CP violation in the decay, since only diagrams vanishing in the U -spin limit contribute to its leading-order decay amplitudes, while CP violating contributions do not cancel in this limit [10, 11]. The new measurement, $A_{CP}(D^0 \rightarrow K_S^0 K_S^0) = (-3.1 \pm 1.2 \pm 0.4 \pm 0.2)\%$, where the third uncertainty is due to the precision with which the CP asymmetry of the normalisation channel is known [12], is the most precise to date and is compatible with zero within 2.4 standard deviations. The second measurement is the first of its kind performed at hadron colliders, and relies on rare Dalitz decays, $h^0 \rightarrow e^+ e^- \gamma$, or on two-photon decays where one of the photons converts into an electron-positron pair within the vertex detector, to allow triggering on the displaced D -meson decay vertex, which would be impossible to reconstruct using bare two-photon decays. The large charm-hadron production cross-section with respect to B factories counterbalances the lower branching fraction of these final states, and allows to obtain results equally or more precise, all being consistent with zero within uncertainties ranging from 1 to 10%.

Crucial advances have been achieved also in time-dependent measurements. In the following we describe mixing and CP violation through the theoretical parametrisation [13], which parametrises the off-diagonal terms of the effective Hamiltonian governing the evolution of the D^0 – \bar{D}^0 system, $\mathbf{H} \equiv \mathbf{M} - \frac{i}{2}\mathbf{\Gamma}$, with the mixing parameters $x_{12} \equiv 2|M_{12}/\Gamma|$ and $y_{12} \equiv |\Gamma_{12}/\Gamma|$ and the weak phases ϕ_2^M and ϕ_2^Γ , defined as the phases of M_{12} and Γ_{12} relative to their dominant $\Delta U = 2$ contribution. The parameters x_{12} and ϕ_2^M correspond to dispersive off-shell transitions, and are sensitive to interactions with new heavy particles, while y_{12} and ϕ_2^Γ quantify absorptive on-shell transitions.

A new measurement of $D^0 \rightarrow K_S^0 \pi^+ \pi^-$ decays [14], based on the data sample collected during 2015–2018, increases tenfold the number of candidates with respect to the previous determination with 2011–2012 data thanks to improved trigger [15]. The variation of strong phases across the Dalitz plot ensures sensitivity to all mixing parameters and CP violation phases. The data are analysed with a model-independent method, in which the Dalitz plot is divided into 8 regions symmetric with respect to its bisector, each having an approximately constant strong-phase difference between the amplitudes in the two halves of the plane [16]. The ratios of the yields in the opposite halves of




Figure 1: (Top-left) Dalitz plot of background-subtracted $D^0 \rightarrow K_S^0 \pi^+ \pi^-$ candidates; m_{\pm}^2 stands for $m^2(K_S^0 \pi^{\pm})$ ($m^2(K_S^0 \pi^{\mp})$) for D^0 (\bar{D}^0) decays. (Bottom-left) Iso- $\Delta\delta$ binning of the Dalitz plane. (Right) CP -averaged yield ratios as a function of decay time for each Dalitz region, with fit projections overlaid.

the plane are measured for each Dalitz-plot region, labelled “ b ”, and in 13 intervals of decay time, labelled “ j ”; see fig. 1. The yields in the denominator mostly correspond to Cabibbo-favoured decays and are nearly constant as a function of decay time, while the fraction of Cabibbo-favoured decays following mixing in the numerator is as important as that of doubly Cabibbo-suppressed decays and increases as a function of decay time, thus providing sensitivity to the mixing parameters. The ratios are equal to

$$R_{bj}^{\pm} \approx \frac{r_b + \sqrt{r_b} \operatorname{Re}[X_b^*(z_{CP} \pm \Delta z)] \langle t \rangle_j + \frac{1}{4} [|z_{CP} \pm \Delta z|^2 + r_b \operatorname{Re}(z_{CP}^2 - \Delta z^2)] \langle t^2 \rangle_j}{1 + \sqrt{r_b} \operatorname{Re}[X_b(z_{CP} \pm \Delta z)] \langle t \rangle_j + \frac{1}{4} [\operatorname{Re}(z_{CP}^2 - \Delta z^2) + r_b |z_{CP} \pm \Delta z|^2] \langle t^2 \rangle_j}, \quad (1)$$

where r_b is the ratio at zero decay time, $X_b \equiv \langle e^{i(\Delta\delta)} \rangle_b$ is the average of the exponential of the strong-phase difference, as measured at charm factories [17, 18], and the two complex parameters $z_{CP} \equiv -(y_{CP} + i x_{CP})$ and $\Delta z \equiv -(\Delta y + i \Delta x)$ are equal to

$$\begin{aligned} x_{CP} &\equiv x_{12} \cos \phi_2^M = (3.97 \pm 0.46 \pm 0.29) \times 10^{-3}, & \Delta x &\equiv -y_{12} \sin \phi_2^{\Gamma} = (-0.27 \pm 0.18 \pm 0.01) \times 10^{-3}, \\ y_{CP} &\equiv y_{12} \cos \phi_2^{\Gamma} = (4.59 \pm 1.20 \pm 0.85) \times 10^{-3}, & \Delta y &\equiv x_{12} \sin \phi_2^M = (0.20 \pm 0.36 \pm 0.13) \times 10^{-3}. \end{aligned}$$

The results for x_{CP} and Δx improve the precision of their world average by a factor of 3, and the former constitutes the first observation of a nonzero mass difference between the neutral charm meson eigenstates, with a significance greater than 7 standard deviations.

The limits on the ϕ_2^M angle can be further improved by measuring the slope of the linear expansion of the time-dependent asymmetry of the decay rates of D^0 and \bar{D}^0 mesons into the CP -even final state f , where f stands for $= K^+ K^-$ or $\pi^+ \pi^-$,

$$A_{CP}(f, t) \equiv \frac{\Gamma(D^0 \rightarrow f, t) - \Gamma(\bar{D}^0 \rightarrow f, t)}{\Gamma(D^0 \rightarrow f, t) + \Gamma(\bar{D}^0 \rightarrow f, t)} \approx a_f^d - \Delta y \frac{t}{\tau_{D^0}}, \quad (2)$$

where a_f^d is the CP asymmetry in the decay and Δy has been defined above. A new measurement relying on the 2015–2018 data sample [19], $\Delta y = (1.0 \pm 1.1 \pm 0.3) \times 10^{-4}$, is the most precise

search for CP violation performed to date at a hadron collider and improves the precision of the world average by around a factor of 2.

3. Mixing and CP violation in beauty decays

Precise measurements of the mass difference between the B_s^0 eigenstates, Δm_s , are needed to minimise the systematic uncertainty of measurements of the CKM angle γ through time-dependent analyses [20] and, if combined with measurements of its analogue for B^0 mesons, Δm_d , provide constraints on the unitarity of the CKM matrix. The LHCb collaboration has recently measured Δm_s using $B_s^0 \rightarrow D_s^- \pi^+$ decays collected during 2015–2018, where the D_s^- meson decays into the $K^+ K^- \pi^-$ or $\pi^+ \pi^- \pi^-$ final states [21]; see fig. 2 left. The result is, in natural units,

$$\Delta m_s = 17.7683 \pm 0.0051 \pm 0.0032 \text{ ps}^{-1}.$$

Its relative precision, 3×10^{-4} , is lower than that of the previous world average by a factor of 2.

Further results include the first search for CP violation through an amplitude analysis of $\Xi_b^- \rightarrow p K^- K^-$ decays [22] and a measurement of the CKM angle γ with $\Lambda_b^0 \rightarrow D p K^-$ decays [23]. Both are statistically limited and do not observe CP asymmetries differing significantly from zero.

4. First simultaneous combination of charm and beauty results

A recent combination of LHCb measurements of the CKM angle γ yields $\gamma = (65.4^{+3.8}_{-4.2})^\circ$ [24]. Measurements base on time-independent and time-dependent analyses are compatible with each other at the level of 2 standard deviations, the precision of the latter being worse by a factor of 2; see fig. 2 centre. For the first time, measurements of charm decays are included in the combination and charm mixing parameters are fitted simultaneously to the angle γ and to the beauty hadronic observables. The precision on the strong-phase difference between $D^0 \rightarrow K^- \pi^+$ and $D^0 \rightarrow K^+ \pi^-$ decays, $\delta_D^K \pi$, is improved by around a factor of 2 with respect to the previous world average, thanks to the improved precision of the ADS measurement in Ref. [25]. This allows to improve also the precision on the charm mixing parameter y_{12} by a factor of 2, thanks to the constraints set on $-y_{12} \cos \delta_D^K \pi + x_{12} \sin \delta_D^K \pi$ by time-dependent $D^0 \rightarrow K^\pm \pi^\mp$ measurements [26]; see fig. 2 right.




Figure 2: (Left) Decay-time distribution of $B_s^0 \rightarrow D_s^- \pi^+$ signal decays. The oscillations frequency is equal to Δm_s ; fit projections are overlaid. (Centre) One-dimensional 1 – CL profiles for the CKM angle γ from combinations using inputs from different B -meson species. (Right) Two-dimensional profile likelihood contours for the charm mixing parameters. The brown contours, drawn out from 1 to 5 standard deviations, show the improvement of this combination with respect to the previous world average, in blue.

References

- [1] LHCb collaboration, *LHCb detector performance*, *Int. J. Mod. Phys. A* **30** (2015) 1530022 [1412.6352].
- [2] LHCb collaboration, *Observation of CP violation in charm decays*, *Phys. Rev. Lett.* **122** (2019) 211803 [1903.08726].
- [3] A. Khodjamirian and A.A. Petrov, *Direct CP asymmetry in $D \rightarrow \pi^- \pi^+$ and $D \rightarrow K^- K^+$ in QCD-based approach*, *Phys. Lett. B* **774** (2017) 235 [1706.07780].
- [4] M. Chala, A. Lenz, A.V. Rusov and J. Scholtz, *Δ_{ACP} within the Standard Model and beyond*, *JHEP* **07** (2019) 161 [1903.10490].
- [5] Y. Grossman and S. Schacht, *The emergence of the $\Delta U = 0$ rule in charm physics*, *JHEP* **07** (2019) 020 [1903.10952].
- [6] H.-Y. Cheng and C.-W. Chiang, *Revisiting CP violation in $D \rightarrow PP$ and VP decays*, *Phys. Rev. D* **100** (2019) 093002 [1909.03063].
- [7] S. Schacht and A. Soni, *Enhancement of charm CP violation due to nearby resonances*, [2110.07619].
- [8] LHCb collaboration, *Measurement of CP asymmetry in $D^0 \rightarrow K_S^0 K_S^0$ decays*, *Phys. Rev. D* **104** (2021) L031102 [2105.01565].
- [9] LHCb collaboration, *Search for CP violation in $D_{(s)}^+ \rightarrow h^+ \pi^0$ and $D_{(s)}^+ \rightarrow h^+ \eta$ decays*, *JHEP* **06** (2021) 019 [2103.11058].
- [10] J. Brod, A.L. Kagan and J. Zupan, *Size of direct CP violation in singly Cabibbo-suppressed D decays*, *Phys. Rev. D* **86** (2012) 014023 [1111.5000].
- [11] U. Nierste and S. Schacht, *CP violation in $D^0 \rightarrow K_S K_S$* , *Phys. Rev. D* **92** (2015) 054036 [1508.00074].
- [12] LHCb collaboration, *Measurement of CP asymmetry in $D^0 \rightarrow K^+ K^-$ decays*, *Phys. Lett. B* **767** (2017) 177 [1610.09476].
- [13] A.L. Kagan and L. Silvestrini, *Dispersive and absorptive CP violation in $D^0 - \bar{D}^0$ mixing*, *Phys. Rev. D* **103** (2021) 053008 [2001.07207].
- [14] LHCb collaboration, *Observation of the mass difference between neutral charm-meson eigenstates*, *Phys. Rev. Lett.* **127** (2021) 111801 [2106.03744].
- [15] R. Aaij et al., *Tesla: an application for real-time data analysis in High Energy Physics*, *Comput. Phys. Commun.* **208** (2016) 35 [1604.05596].
- [16] A. Di Canto, J. Garra Tico, T. Gershon, N. Jurik, M. Martinelli, T. Pilař et al., *Novel method for measuring charm-mixing parameters using multibody decays*, *Phys. Rev. D* **99** (2019) 012007 [1811.01032].
- [17] CLEO collaboration, *Model-independent determination of the strong-phase difference between D^0 and $\bar{D}^0 \rightarrow K_{S,L}^0 h^+ h^-$ ($h = \pi, K$) and its impact on the measurement of the CKM angle γ/ϕ_3* , *Phys. Rev. D* **82** (2010) 112006 [1010.2817].
- [18] BESIII collaboration, *Model-independent determination of the relative strong-phase difference between D^0 and $\bar{D}^0 \rightarrow K_{S,L}^0 \pi^+ \pi^-$ and its impact on the measurement of the CKM angle γ/ϕ_3* , *Phys. Rev. D* **101** (2020) 112002 [2003.00091].
- [19] LHCb collaboration, *Search for time-dependent CP violation in $D^0 \rightarrow K^+ K^-$ and $D^0 \rightarrow \pi^+ \pi^-$ decays*, *Phys. Rev. D* **104** (2021) 072010 [2105.09889].
- [20] LHCb collaboration, *Measurement of CP asymmetry in $B_S^0 \rightarrow D_S^\mp K^\pm$ decays*, *JHEP* **03** (2018) 059 [1712.07428].
- [21] LHCb collaboration, *Precise determination of the $B_S^0 - \bar{B}_S^0$ oscillation frequency*, to appear in Nature Physics [2104.04421].
- [22] LHCb collaboration, *Search for CP violation in $\Xi_b^- \rightarrow p K^- K^-$ decays*, *Phys. Rev. D* **104** (2021) 052010 [2104.15074].
- [23] LHCb collaboration, *Observation of the suppressed $\Lambda_b^0 \rightarrow D p K^-$ decay with $D \rightarrow K^+ \pi^-$ and measurement of its CP asymmetry*, submitted to *Phys. Rev. D* [2109.02621].
- [24] LHCb collaboration, *First simultaneous determination of CKM angle γ and charm mixing parameters from a combination of LHCb measurements*, submitted to JHEP [2110.02350].
- [25] LHCb collaboration, *Measurement of CP observables in $B^\pm \rightarrow D^{(*)} K^\pm$ and $B^\pm \rightarrow D^{(*)} \pi^\pm$ decays using two-body D final states*, *JHEP* **04** (2021) 081 [2012.09903].
- [26] LHCb collaboration, *Updated determination of $D^0 - \bar{D}^0$ mixing and CP violation parameters with $D^0 \rightarrow K^+ \pi^-$ decays*, *Phys. Rev. D* **97** (2018) 031101 [1712.03220].