

Drivers of Rift Valley fever epidemics, Madagascar

Renaud Lancelot^{a,b,1}, Marina Béal^{b,c,d}, Vincent Michel Rakotoharinome^e, Soa Fy Andriamandimby^f, Jean-Michel Héraud^f, Caroline Coste^{a,b}, Andrea Apolloni^{a,b}, Cécile Squarizoni-Diaw^g, Stéphane de La Rocque^{a,b,h,i}, Pierre Formenty^j, Jérémy Bouyer^{a,b}, G. R. William Wint^k, and Eric Cardinale^{b,c,d}

^aCirad, UMR Contrôle des Maladies Animales Exotiques et Emergentes (CMAEE), Campus International de Baillarguet, 34398 Montpellier, France; ^bInra, UMR CMAEE 1309, Campus International de Baillarguet, 34398 Montpellier, France; ^cCirad, UMR CMAEE, 97490 Sainte Clotilde, La Réunion, France; ^dCentre de recherche et de veille sur les maladies émergentes de l'Océan Indien (CRVOI), 97490 Sainte Clotilde, La Réunion, France; ^eMinistère de l'Agriculture, de l'Elevage et de la Pêche, Direction des Services Vétérinaires, Ambatofotsikely, BP 530 Antananarivo 101, Madagascar; ^fUnité de Virologie, Institut Pasteur de Madagascar, BP 1274 Antananarivo 101, Madagascar; ^gAnimal Production and Health Division, Food and Agriculture Organization of the United Nations (FAO), Viale delle Terme di Caracalla, 00153 Rome, Italy; ^hWorld Organization for Animal Health (OIE), 12 rue de Prony 75017 Paris, France; ⁱInternational Health Regulations (IHR) Monitoring Procedure and Information Team, Global Capacity and Response Department, World Health Organization (WHO), 20 Avenue Appia, 1211 Geneva 27, Switzerland; ^jEmerging and Epidemic Zoonotic Diseases, Pandemic and Epidemic Disease department, World Health Organization (WHO), 20 Avenue Appia, Geneva 27, Switzerland; ^kEnvironmental Research Group Oxford (ERGO), Department of Zoology, South Parks Road, Oxford OX1 3PS, United Kingdom

This manuscript was compiled on December 5, 2016

Rift Valley fever (RVF) is a vector-borne viral disease widespread in Africa. The primary cycle involves mosquitoes, and wild and domestic ruminant hosts. Humans are usually contaminated after contact with infected ruminants. As many environmental, agricultural, epidemiological and anthropogenic factors are implicated in RVF spread, the multidisciplinary One Health approach was needed to identify the drivers of RVF epidemics in Madagascar. We examined the environmental patterns associated with these epidemics, comparing human and ruminant serological data with environmental and cattle-trade data. In contrast to East Africa, environmental drivers did not trigger the epidemics: they only modulated local RVFV transmission in ruminants. Instead, RVFV was introduced through ruminant trade and subsequent movement of cattle between trade hubs caused its long-distance spread within the country. Contact with cattle brought in from infected districts was associated with higher infection risk in slaughterhouse workers. The finding that anthropogenic rather than environmental factors are the main drivers of RVF infection in humans can be used to design better prevention and early detection in case of RVF resurgence in the region.

Vector-borne infection | zoonosis | El Niño | cattle trade | One Health

Rift Valley fever (RVF) is a vector-borne infection caused by the RVF virus (RVFV). Mosquito species from the *Aedes*, *Culex*, and *Mansonia* genera are the main RVFV vectors [1]. The virus persists in the environment either by vertical transmission occurring in some *Aedes* species, or by an enzootic cycle involving ruminants and mosquitoes. When environmental conditions are favourable to mosquito proliferation, this cycle is amplified by *Culex* populations, leading to RVF epidemics [2]. In the Horn of Africa, these circumstances are met during the warm phases of El Niño southern oscillation (ENSO), and the Indian Ocean dipole zonal mode: warm sea-surface temperatures in the equatorial eastern-central Pacific Ocean and the western equatorial Indian Ocean result in heavy autumn rainfall, and subsequent greening of the vegetation [3–5]. The latter can be monitored by the remotely-sensed normalized difference vegetation index (NDVI). Besides these environmental conditions triggering epidemics, long-distance RVFV dissemination is often related to ruminant trade [6], as reported in Madagascar and Saudi Arabia [7–9]

In domestic ruminants, the RVFV causes mass abortions and high neonatal mortality [10]. Humans may get infected after the bites of infected mosquitoes, or the consumption of raw milk. However, most clinical cases occur after contact with blood, aborted fetuses, and placenta of viremic animals.

Farmers, veterinarians, slaughterhouse workers, and butchers are thus the most exposed to the risk of RVF, as well as any people attending the slaughtering of viremic animals [11]: blood aerosol produced on this occasion is highly infectious [12]. The virus is not transmitted from person to person.

In Madagascar, the first known RVF epidemic occurred in 1990–91 on the East coast and in the central highlands [7]. A second epidemic occurred in 2008–2009 with official reports suggesting at least 700 suspected cases and 26 RVFV laboratory confirmed fatalities. However due to under-reporting, an excess of 10,000 human cases were estimated [13–15].

Madagascar is one of the poorest countries in the World, with 90% of the population living with less than USD 1.5 inhab.^{−1} d^{−1} [16]. Because public health resources are limited in such setting, it is crucial to identify the most exposed populations, based on scientific evidences. To this end, we aimed at understanding the circumstances in which RVF epidemics occurred. We firstly assessed the risk of RVFV introduction to ruminants via livestock trade. Then, we described the environmental conditions associated with the start of 1990 and 2008 epidemics, and with local RVFV transmission. We then aimed to build risk indices of RVFV human infection related to the

Significance Statement

Rift Valley fever (RVF) is an emerging, mosquito-borne viral infection of ruminants, transmissible to people, and linked to rainfall. By investigating a wider range of possible drivers this study confirms the assumption that RVF occurrence can also be dependent non-environmental drivers. In Madagascar, human and ruminant infections were geographically distinct, with introduction and long distance disease spread linked to livestock trading. Human infections were highest in those involved with slaughtering and handling fresh meat. The identification of possible introduction routes, major cattle trade hubs, and areas of high risk to ruminants, has shown how multidisciplinary analyses are needed to properly understand disease dynamics and spread thereby improving early detection and prevention of RVF in humans.

R.L., V.M.R., S.F.A., C.S.D., S.D.L.R., P.F., G.R.W.W., and E.C. designed research; R.L., M.B., V.M.R., S.F.A., J.M.H., C.C., A.A., C.S.D., and E.C. performed research; R.L., M.B., C.C., A.A., and G.R.W.W. analysed data; and R.L., M.B., S.F.A., J.M.H., C.C., A.A., S.D.L.R., P.F., J.B., G.R.W.W., and E.C. wrote the paper.

The authors declare no conflict of interest.

¹To whom correspondence should be addressed. E-mail: renaud.lancelotcirad.fr

local environment (local risk) and cattle trade (remote risk), assess their relative importance, identify and map high-risk areas, and assess the consequences for human health.

Results

Risk of virus introduction by livestock trade. Queries in the UN ComTrade database did not reveal direct importations of live ruminants from mainland Africa to Madagascar. However, they highlighted several official importations from the Union of Comoros in 2005, 2006, and 2007 (Tab. 1). Though the figures are small, they confirmed the risk of RVFV introduction in Madagascar from an infected country, via livestock trade.

Table 1. Importation of live ruminants in Madagascar from the Union of Comoros between 2005 and 2007. Source: UN ComTrade.

Year	Species	Quantity (head)
2005	cattle	30 ^a
	goats	9 ^a
2006	goats	14 ^a
2007	cattle	7
	goats	88
	sheep	47

^a Estimated figure from the reported financial value. No imports of live-ruminants were reported from the Union of Comoros in 2003 and 2004.

In addition to this official trade, illegal cattle movements between the Comoros Archipelago and Madagascar were probably much more frequent. Informal surveys conducted in 2009 and 2010 in the main Comoros harbours, and in the North-West of Madagascar (Mahajanga and Antsiranana), revealed the frequent presence of cattle and small ruminants on board of freighters and *botry* (dhows) travelling from Comoros Islands to Madagascar, and from port to port. This coastal navigation is widespread in Madagascar, given the weakness of the terrestrial-road network (Fig. S1). Therefore, RVFV could have been introduced to Madagascar through ruminant trade from Comoros Islands - which were previously infected [17, 18] - and further disseminated through coastal navigation between Malagasy sea ports.

Environmental conditions and RVFV epidemics.

Triggering RVF epidemics. Conditions at the start of the two RVF epidemics in humans (March 1990 and January 2008) are shown on Fig. 1. For the three indicators (SOI, rainfall, and NDVI) and three biomes (Fig. S2A), the two epidemics did not occur in typical RVF conditions according to the eastern African standards [14]. In March 1990, a marked negative anomaly was observed for southern oscillation index (SOI), one of the main indicators of ENSO (Fig. 1A). Rainfall was lower than normal in all three biomes (Fig. 1B). March 1990 was also at the end of a period of positive NDVI anomalies, and before a short period of negative anomalies for the dry forest and xeric shrubland. The NDVI pattern was not clear for the moist forest where the epidemics started (Fig. 1C). Conditions during the 2008 epidemic were almost the reverse as it occurred during a cold (positive) anomaly of SOI, when rainfall was close to normal in the moist forest and somewhat higher than normal in the two other biomes.




Fig. 1. Environmental conditions (monthly average) in the main Malagasy biomes with respect to the 1990-1991 and 2008-2009 RVF epidemics. (A) Standardized southern oscillation index [19]. (B) Standardized rainfall anomalies. Source: NOAA's precipitation reconstruction over Land (PREC/L) provided by the NOAA/OAR/ESRL PSD, Boulder, Colorado, USA. (C) Standardized NDVI anomalies. Source: GIMMS AVHRR NDVI version 3 [20]. On each plot, favourable conditions for RVF were drawn in red.

These general impressions were corroborated by the results of partial triadic analysis and hierarchical clustering (Fig. S3): the environmental conditions of RVF epidemics fell into quite different clusters in 1990-1991, and in 2008-2009: with a three-class partition, the 1990 epidemics occurred in a category of rainy seasons with close-to-normal mean values for SOI (0.01), rainfall (0.16), and NDVI (-0.11). In 2008, the epidemic fell in a category with a high mean SOI (1.38), and close-to-normal mean values for NDVI (0.07) and rainfall (-0.16).

RVFV spread at the end of 2008 epidemic. We modelled the sero-prevalence rate of immunoglobulins of type G (IgG) in ruminant sera collected at the end of 2008 epidemic to assess the role of environmental conditions in RVF spread in livestock. The importance of environmental predictors with respect to the selected subset of plausible models is displayed in Tab. S2. Multi-model averaged coefficients are shown in Tab. S3. Higher rainfall, municipalities within 50 km from a sea port, and lower altitude were associated with higher sero-prevalence rate in ruminants.

The ROC curve for the averaged model had an area under the curve of 74%. The map of predicted sero-prevalence rate showed high-risk areas on the north-western and north-eastern coasts (lowlands). Southern regions and highlands were less affected at the exception of sea-port municipalities (Fig. 2). See SI Results for details.




Fig. 2. Sero-prevalence rate of anti-RVVF IgG in ruminants (Madagascar) predicted by the averaged beta-binomial logistic regression model: (A) predicted rate, (B) coefficient of variation.

The cattle trade network and RVFV dissemination. Cattle trade data were collected monthly from 2007 to 2011, with large variations in numbers across the years. The overall network activity is presented on Fig. S4A: each segment corresponds to a link between two nodes; a segment is drawn if at least one movement has been recorded along that link on a given month; its colour is related to its recorded frequency during the five-year survey. To assess the influence of the cattle trade network on the risk of RVFV in humans, we quantified the trade flows using the most comprehensive data set, collected in 2010. This network had the largest number of nodes (257 farms, markets, and slaughterhouses) and links (346). We could distinguish two sets of links: those active throughout the year (56% of the links) constituting the *backbone* of the network; those *occasional* links active one or two months. Despite their large number, only a small volume (3%) of animals was traded on these occasional links. Most of the backbone nodes and links active in 2010 were already present in the previous year.

To assess the role of the occasional links on RVFV spread, we considered separate monthly network snapshots and we evaluated the corresponding epidemic threshold q [21] which provides an estimate of the critical probability for the virus to spread through the network: the lower its value, the higher the risk of spread. The backbone network was the smallest in January (rainy season) and then increased till reaching its plateau size in April (early dry season): Fig. S4B. The number of occasional links peaked in May and October. Despite these large variations, q was slightly affected by the presence of these occasional links. Conversely, it was strongly influenced by the variations of trade volume in the backbone network. The whole network was prone to the diffusion of RVFV: the average epidemic threshold was $\bar{q} \simeq 2.1\%$, the period of maximum risk from March to May, with $\hat{q} \simeq 1.5\%$, and the minimum risk

($\hat{q} \simeq 3.9\%$) in January.

The distribution of in/out degrees (number of incoming/outgoing links for a given node) matched a power-law distribution. Only a few nodes (Ihosi, Tsiroanomandy, Ambalavao, and Mampikony) were highly connected hubs and also had the highest betweenness. The four network hubs belonged to three different giant strong components (GSC) - Ambalavao and Ihosi were in the same one. Nodes belonging to these GSC were strongly connected between themselves, thus increasing the risk of infection for other locations in their neighbourhood [22]. These hubs were also cut-points in the network: their removal would disrupt the network connectedness and thus limit the diffusion of an infectious agent like RVFV.

The pastoral areas of South-West and North-West of Madagascar were the major sources of the traded cattle (Fig. 3A) and were characterized by high cattle density and low human density. The great majority of markets had a negative balance in cattle flows: they were sources in the network, with low local consumption (Fig. 3B). On the other hand, two hubs (Ambalavao and Tsiroanomandy) had a large, positive balance. Further information is provided on Fig. 3C: cattle were collected in the markets of Tsiroanomandy and Ihosi and then sent to Ambalavao for slaughtering, Tsiroanomandy also being a consumption centre. In addition, a close examination of these four hubs showed that many small, outgoing flows offered numerous opportunities for long-distance RVFV spread.

Cattle trade and the risk of RVFV infection in humans. The coefficient for the local, environment-related index of RVFV transmission to humans was not significant, in contrast to the coefficient for the remote, cattle-trade related index (SI Tab. S3). The odds ratio (OR) for the latter was ORC=1.7 (95% confidence interval (CI): [1.1; 2.7]). The risk of humans getting infected with RVFV increased with the intensity of cattle trade from areas with infected livestock.

The map of predicted sero-prevalence rate in humans (Fig. 4), as well as the plot of random variations (Fig. S8) both highlighted large spatial variations in the infection rate, with a clear spatial pattern. The highest sero-prevalence rates were encountered in the densely populated areas of central highlands. This pattern is emphasized in Fig. S8B: a strong additional risk (with respect to cattle-trade related risk) was found in the regions of Antsirabe and Lake Alaotra. See SI Results for details.

Discussion

Cattle trade and the risk of RVFV introduction. Imports of live ruminants from Comoros were the main driver for RVFV introduction to Madagascar livestock and subsequent trade-related movements of cattle led to its spread to humans, at least in 2008. Phylogenetic studies showing that Malagasy RVFV were closely related to viruses previously circulating in mainland Africa [15] suggest that RVFV was probably introduced into the Comoros Islands through cattle trade with East Africa [23]. The existence of illegal livestock importation from Comoros Islands provided opportunities for the introduction of RVFV in Madagascar. Preventing such introductions is therefore essential to avoid further RVF epidemics in Madagascar. In practice, strengthening communication between African, Comoros, and Malagasy public-health and veterinary services would be important to share early detection in the event of




Fig. 3. Cattle flows in the Malagasy cattle-trade network, 2010. (A) Municipality of origin (number of flows); (B) Balance in cattle flows at the municipality level (number of heads); (C) Directed flows (number of heads).




Fig. 4. Sero-prevalence rate in humans predicted by a mixed-effect binomial logistic regression model: (A) predicted rate, (B) coefficient of variation. On panel (A), numbers were placed at the centroid of districts with the highest predicted rates (10th decile): 1 Ambohidratrimo; 2 Antananarivo Renivohitra; 3 Antananarivo-Sud; 4 Antsirabe Rural; 5 Antsirabe Urban; 6 Betafo; 7 Faratsiho; 8 Ambatondrazaka; 9 Amparafaravola; 10 Anosibe; 11 Moramanga.

new RVFV circulation. Also, quarantine measures should be reinforced for ruminants exported from continental Africa to Comoros Islands.

Environmental conditions and RVF epidemics. Anyamba et al. [14] already pointed out the contrasting behaviour of the

disease in Madagascar, East and South Africa. Prolonged heavy rainfall and positive NDVI anomalies occurred after the first known epidemics of 1990, e.g. in 1994 with the occurrence of Gerald cyclone [24]. More recently, torrential rainfall occurred in February-March 2015 in Madagascar, related to the strong El Niño conditions [25]. No RVF outbreak could be detected despite specific surveillance measures in 1994 (shortly after the 1990-91 RVF epidemic), and strengthened national and regional surveillance in 2015 [26, 27].

RVF epidemics usually start in the arid environment of *dambos* (East Africa) or *pans* (southern Africa) [1], which are similar to ecosystems in the Toliara region of South-West of Madagascar that are covered by xeric shrublands (Fig. S2A). Though this sea-port municipality was hit by RVF, the sero-prevalence rate in ruminants was low in the neighbouring municipalities (Fig. 2), probably because the virus was introduced in sub-optimal conditions for the vectors (no heavy rainfall, no flooding). Conversely, high sero-prevalence rates were observed in cattle in the North-West of the island, covered by the dry forest. Moreover, in East Africa, *Aedes* mosquitoes are the primary RVFV vectors [1]: their biology and ecology is well adapted to arid environments. Regarding Madagascar, 23 mosquito species might be considered as potential RVFV vectors, including no flood-water *Aedes* mosquito species [28–30]. A single species, *Cx. antennatus*, meets all criteria for formal classification as an RVFV vector [31]. This mosquito is widespread in Madagascar (except in the North), including in rice paddies which cover large areas of the island. The introduction of infected animals in conjunction with *Culex* hatching, during a standard rainy season profile, might have amplified the outbreak locally, in many places.

Two major differences are thus highlighted between Madagascar and East Africa: (i) the lack of connection between the start of 1990-1991 and 2008-2009 RVF epidemics and El Niño events - and more generally, with anomalous heavy rainfall, and (ii) no obvious role of *Aedes* mosquitoes in the primary RVFV transmission cycle, as well as the wide distribution of *Culex* (and other mosquito species).

Apparently, the climatic conditions observed during the two epidemics are common in Madagascar: the drivers triggering RVFV epidemics must therefore be sought elsewhere. Nevertheless, these climatic conditions remain important for the amplification of the primary epidemiological cycle between mosquitoes and ruminants (Tab. S2).

Cattle trade as a driver of RVF epidemics. The Malagasy population is growing fast - from 16 to 24 million between 2000 and 2015 [16], and is concentrating in Antananarivo and other large cities (Fig. S2B). Cattle is omnipresent in Malagasy agriculture, economy, and culture. This leads to an ever-growing demand for cattle meat and draught power for crops. Consequently, the increasing cattle trade provides more opportunities for RVF epidemics to spread [7, 8]. Our description of the national cattle-trade network strongly supports this assumption and further extends a previous analysis in the North of Madagascar [8]. The connectedness of the cattle-trade network and its low percolation threshold make the risk of seeding epidemics high.

Lake Alaotra and Antsirabe (Fig. S2A) are two major crop and livestock farming regions. The former is the largest rice-production basin in Madagascar, with many paddies and swamps favourable to mosquito proliferation. The two regions are densely populated, with cities harbouring big livestock markets, and slaughterhouses to match local red meat needs. Therefore, there are many opportunities for human exposure to the RVFV when viremic animals are slaughtered.

Additionally, using the network of markets, butcheries and *hotely* (cheap restaurants for travellers), farmers try to sell their sick animals at the first clinical sign of any disease, to mitigate economic losses. When they do not succeed, they slaughter them and eat their meat [32]. This practice was probably at the origin of many unreported human infections during the 2008-2009 RVF epidemics in Madagascar.

More collaboration with sociologists and anthropologists are needed to decipher farmers' perceptions of animal diseases, and to assess the social acceptability of prevention, surveillance, and control measures, such as cattle vaccination to protect people (should enough vaccine be available), or cattle movement restrictions to avoid RVFV spread through hubs in the cattle-trade network [33]. If RVFV introduction into Malagasy livestock continues, targeted (risk-based) vaccination campaigns of ruminants might be organized to protect human populations. Individual protection measures, such as vaccination when the human vaccine becomes available, or wearing personal protective equipments such as gowns, gloves, safety glasses, and masks when slaughtering ruminants, would also be important to implement in the most exposed categories of people [34, 35], together with dissemination and training programs. This implementation of coordinated actions between Public Health and Veterinary Services would represent an important advance in the so-called "One Health" joint approach of human and animal health [27, 36].

Materials and Methods

To assess the risk of RVFV introduction, we scanned national and international databases on sea trade from 2003 to 2008. We also implemented informal surveys in harbours of Comoros Islands and north-western Madagascar during and after this epidemic (up to 2010).

To describe the environmental conditions of the past RVF epidemics, we used global data sets available as long-term time series: (i) the southern oscillation index (SOI) as a main indicator of El Niño southern oscillation (ENSO) of major importance for the climatic conditions in East Africa and the South-West tropical Indian Ocean [3]; (ii) rainfall data, of crucial importance for the epidemiology of vector-borne diseases [37]; and (iii) the NDVI as an indicator of rainfall impact on the vegetation [38]: forage resources for ruminants and resting sites for mosquitoes. These indicators were averaged over the main biomes (SI Material and methods). Firstly, we plotted the time series for each biome. Secondly, we selected data from the dry-forest biome where the highest sero-prevalence rate was observed in cattle in 2008 [39]. We then used partial triadic analysis (PTA), a multi-table version of principal component analysis (PCA) [40] to identify common or contrasted patterns in rainy seasons (November to March) from 1983 to 2011 (28 years).

To assess the risks of RVFV spread within Madagascar, we used two types of data: (i) environmental data selected among the factors of interest for mosquito-borne infections, i.e., related to the availability of resting or breeding sites, or the seasonal changes in conditions favouring the development of their immature stage, etc. [37]: see the list in (SI Material and methods). They were obtained from international databases, and processed using standard methodology; (ii) cattle-trade data collected during repeated nationwide surveys in livestock markets and slaughterhouses, implemented from 2007 to 2011 for the purpose of this study.

Environmental data were used in beta-binomial logistic regression (BBLR) models to predict the RVFV sero-prevalence rate for cattle at the municipality level. The serological data were collected after the 2008 epidemics, and published [39]. Sera were tested for the presence of anti-RVFV IgG. We adopted a multi-model inference approach to select the best predictors. A set of plausible models was kept for model averaging and prediction of sero-prevalence rate in cattle at the municipality level.

Cattle-trade data were used (i) to assess the risk of RVFV spread through the network, and (ii) to assess the risk of human infection with RVFV. Farms, markets and slaughterhouses were the nodes of a directed cattle-trade network. A link between two nodes corresponded to a trade movement of animals. The direction indicated the origin and destination places, and the volume was the number of traded animals.

We assessed the risk of RVFV infection in humans using sera collected during by a nation-wide survey in slaughterhouse workers achieved in 2008 and 2009. The study was approved by the Malagasy National Ethical Committee. Participants were included if they gave written informed consent. Sera were tested for the presence of anti-RVFV IgM - signing a recent infection, and the results were published [13]. We defined two indices for the risk of RVFV infection in humans: (i) an index of local RVFV transmission risk, defined by the product of local predicted sero-prevalence rate in cattle and cattle density. This index was built to capture the risk associated with the primary RVFV epidemiological cycle involving mosquitoes and domestic ruminants, in the absence of known wild hosts for RVFV in Madagascar [41, 42]. Humans are not involved in this cycle. Therefore, their density was not considered in the index; (ii) an index of RVFV transmission risk related to cattle trade. It was defined as the product of cattle incoming flow (number of head), predicted sero-prevalence rate in cattle at the origin, and local human density. This latter variable was included because meat consumption is proportionally higher in urban than in rural areas [43] - with higher densities of slaughterhouses, butchers, and meat markets. Consequently, the risk of RVFV infection related to cattle trade should be positively correlated with human density.

The effect of these two indices on human sero-prevalence rate was assessed with a BBLR model. Finally, a mixed-effect binomial logistic regression model was used to study the local variations of sero-prevalence rate in humans, with the administrative district and

region as the nested random effects associated with the intercept. See SI Material and methods for detailed information on data sources and statistical methods, as well as links to download the data sets used in the analyses.

ACKNOWLEDGMENTS. This study was partially funded by the World Health Organization (WHO) and the Food and Agriculture Organization of the United Nations (FAO) through the Central Emergency Response Fund of the United Nations. It was also partially funded by EU grant FP7-613996 VMERGE and is catalogued by the VMERGE Steering Committee as VMERGExxx (<http://www.vmerge.eu>). The contents of this publication are the sole responsibility of the authors and don't necessarily reflect the views of the European Commission.

1. Linthicum KJ, Britch SC, Anyamba A (2016) Rift Valley fever: an emerging mosquito-borne disease. *Annu. Rev. Entomol.* 61:395–415.
2. de La Rocque S, Formenty P (2010) *Rift Valley fever: disease ecology and early warning*, Sustainable management of animal production and health, eds. Odongo N, Garcia M, Viljoen G. (Food and Agriculture Organization of the United Nations, Rome, Italy), pp. 327–333.
3. Linthicum K et al. (1999) Climate and satellite indicators to forecast Rift Valley fever epidemics in Kenya. *Science* 285(5426):397–400.
4. Saji NH, Goswami BN, Vinayachandran PN, Yamagata T (1999) A dipole mode in the tropical indian ocean. *Nature* 401(6751):360–363.
5. Anyamba A et al. (2009) Prediction of a Rift Valley fever outbreak. *Proc. Natl. Acad. Sci. U. S. A.* 106(3):955–959.
6. Chevalier V, Pépin M, Plée L, Lancelot R (2010) Rift Valley fever - A threat for Europe? *Euro-surveillance* 15(10):pii=19506.
7. Morvan J, Saluzzo JF, Fontenille D, Rollin PE, Coulanges P (1991) Rift Valley fever on the East coast of Madagascar. *Res. Virol.* 142(6):475–482.
8. Nicolas G, Durand B, Duboz R, Rakotondravao R, Chevalier V (2013) Description and analysis of the cattle trade network in the Madagascar highlands: potential role in the diffusion of Rift Valley fever virus. *Acta Trop.* 126(1):19–27.
9. Shoemaker T et al. (2002) Genetic analysis of viruses associated with emergence of Rift Valley fever in Saudi Arabia and Yemen, 2000-01. *Emerg. Infect. Dis.* 8(12):1415–1420.
10. Gerdes G (2004) Rift Valley fever. *Rev. Sci. Tech. OIE* 23(2):613–623.
11. Anyangu AS et al. (2010) Risk factors for severe rift valley fever infection in kenya, 2007. *Am J Trop Med Hyg* 83(2 Suppl):14–21.
12. Reed C et al. (2013) Aerosol exposure to Rift Valley fever virus causes earlier and more severe neuropathology in the murine model, which has important implications for therapeutic development. *PLoS Negl. Trop. Dis.* 7(4):e2156.
13. Andriamandimby SF et al. (2010) Rift Valley fever during rainy seasons, Madagascar, 2008 and 2009. *Emerg. Infect. Dis.* 16(6):963–970.
14. Anyamba A et al. (2010) Prediction, assessment of the Rift Valley fever activity in East and Southern Africa 2006-2008 and possible vector control strategies. *Am. J. Trop. Med. Hyg.* 83(2 Suppl):43–51.
15. Carroll SA et al. (2011) Genetic evidence for Rift Valley fever outbreaks in Madagascar resulting from virus introductions from the East African mainland rather than enzootic maintenance. *J. Virol.* 85(13):6162–6167.
16. The World Bank Group (2015) *Countries: Madagascar* (1818 H Street, NW Washington, DC 20433 USA). Accessed 28 December 2015.
17. Sissoko D et al. (2009) Rift Valley fever, Mayotte, 2007-2008. *Emerg. Infect. Dis.* 15(4):568–570.
18. Cêtre-Sossah C et al. (2012) Prevalence of Rift Valley fever among ruminants, Mayotte. *Emerg. Infect. Dis.* 18(6):972–975.
19. NOAA (2015) *Southern oscillation index*. Accessed 30 December 2015.
20. Pinzon JE, Tucker CJ (2014) A non-stationary 1981–2012 AVHRR NDVI3g time series. *Remote Sensing* 6(8):6929–6960.
21. Volkova VV, Howey R, Savill NJ, Woolhouse MEJ (2010) Sheep Movement Networks and the Transmission of Infectious Diseases. *PLoS ONE* 5(6):e11185.
22. Danon L et al. (2011) Networks and the epidemiology of infectious disease. *Interdisciplinary Perspectives on Infectious Diseases* Vol.2011:Article no. 284909.
23. Maquart M et al. (2016) Phylogeographic reconstructions of a Rift Valley fever virus strain reveals transboundary animal movements from eastern continental Africa to the Union of the Comoros. *Transbound. Emerg. Dis.* 63(2):e281–e285.
24. Fitchett JM, Grab SW (2014) A 66-year tropical cyclone record for south-east africa: temporal trends in a global context. *Int. J. Climatol.* 34(13):3604–3615.
25. Monastersky R (2016) Monster El Niño probed by meteorologists. *Nature* 529(7586):267–268.
26. Zeller H, Andriananana R, Quirin R (1998) Surveillance de la fièvre de la Vallée du Rift à Madagascar [Rift Valley fever surveillance in Madagascar], (Institut Pasteur de Madagascar, Antananarivo), Technical Report Projet IPM / PSE/ BM: Janvier 1996 - Mars 1998. 18 p.
27. Cardinale E, Rasamoelina-Andriananivo H, Razafimandimby A, Lepec R, Flachet L (2015) A concrete regional “One Health” surveillance system and management of epidemics: a success story in the South West Indian Ocean in *ISVEE 14, Yucatan 2015 - Veterinary Epidemiology & Economics: Planning Our Future*, Session 2: One Health. (Meridia, Mexico), p. 55.
28. Ratovonjato J et al. (2011) Detection, isolation, and genetic characterization of Rift Valley fever virus from *Anopheles* (*Anopheles*) *coustani*, *Anopheles* (*Anopheles*) *squamosus*, and *Culex* (*Culex*) *antennatus* of the Haute Matsiatra region, Madagascar. *Vector Borne Zoonotic Dis.* 11(6):753–759.