

Research Article

Cite this article: Mutiso V, Ndeti D, Jeremiah E, Nyamai P, Walusaka S, Onyango V, Musyimi C, Bhui K and Mamah D (2025). The moderating and mediating role of resilience in the relationship between adverse childhood experiences and depression, PTSD, and suicidality in Kenyan youth. *Cambridge Prisms: Global Mental Health*, **12**, e88, 1–16
<https://doi.org/10.1017/gmh.2025.10041>

Received: 21 February 2025

Revised: 24 June 2025

Accepted: 15 July 2025

Keywords:

ACEs; resilience; depression; suicidality; PTSD

Corresponding author:

David Ndeti;

Email: dmndeti@amhf.or.ke

The moderating and mediating role of resilience in the relationship between adverse childhood experiences and depression, PTSD, and suicidality in Kenyan youth

Victoria Mutiso^{1,2}, David Ndeti^{1,2,3} , Eric Jeremiah^{1,2}, Pascalyne Nyamai^{1,2}, Samuel Walusaka^{1,2}, Veronica Onyango^{1,2}, Christine Musyimi^{1,2}, Kamaldeep Bhui^{4,5}  and Daniel Mamah⁶

¹Africa Institute of Mental and Brain Health (formerly Africa Mental Health Research and Training Foundation), Nairobi, Kenya; ²World Psychiatric Association Collaborating Centre for Research and Training, Nairobi, Kenya; ³Department of Psychiatry, University of Nairobi, Nairobi, Kenya; ⁴Department of Psychiatry, Nuffield Department of Primary Care Health Sciences, Wadham College, University of Oxford, Oxford, UK; ⁵World Psychiatric Association Collaborating Centre, Oxford, UK and ⁶Department of Psychiatry, Washington University School of Medicine, St. Louis, MO, USA

Abstract

Adverse childhood experiences (ACEs) are widely associated with mental health disorders, such as depression, post-traumatic stress disorder (PTSD), and suicidality. Resilience plays a role in mediation and moderation of these associations, yet there is limited data from Kenya on this. This cross-sectional study examined the role of resilience in the relationship between ACEs and mental health outcomes among 1,972 participants aged 14–25 years in the Nairobi Metropolitan area. Participants completed the Trauma and Distress Scale (ACEs), Patient Health Questionnaire-9 (depression), Columbia-Suicide Severity Rating Scale (suicidality), Harvard Trauma Questionnaire (PTSD), and Adult Resilience Measure-Revised (resilience). Analyses of moderation and mediation using Hayes Process Macro indicated that resilience moderated the association between ACEs with PTSD and depression, with minimal effect on suicidality. It also moderated specific associations, including emotional/physical neglect on ideation, physical abuse on lifetime behavior ($p = 0.0479$), and total ACEs on recent behavior ($p = 0.0514$). Resilience also partially mediated the effects of ACEs on PTSD and depression, and fully mediated suicidality for specific ACE domains (emotional neglect, physical neglect, and physical abuse on suicidal ideation and all ACEs on recent suicidal behaviors). Building resilience mitigates the effects of ACEs on depression, PTSD, and suicidality among Kenyan youth.

Impact statement

This study highlights the role of resilience in determining the mental health outcomes of young Kenyans who have experienced adverse childhood experiences (ACEs). Resilience mediates and moderates the effects of ACEs on depression, post-traumatic stress disorder (PTSD), and suicidality. This calls for policies and initiatives that strengthen resilience in our study community population. Despite being a powerful protective factor against depression and PTSD, the impact of resilience on suicidality differs according to the kind of ACE, which emphasizes the necessity for targeted intervention.

Introduction

Adverse childhood experiences (ACEs) consists of a spectrum of traumatic events occurring before the age of 18 years and include various forms of abuse, neglect, and household dysfunction (Youssef et al., 2013; Schulz et al., 2014). ACEs are becoming more widely acknowledged as a serious public health issue with profound effects on mental health and overall well-being (Sasidharan and Talwar, 2023; White et al., 2024), and in particular resulting in depression, post-traumatic stress disorder (PTSD), and suicidality (Benarous et al., 2017; Burstein and Greenfield, 2020; Ashaba et al., 2021). The World Health Organization highlights ACEs as a critical public health issue and calls for continued efforts to develop interventions to reduce ACEs and their effects on mental health (Oral et al., 2016; Bethell et al., 2017; Srivastav et al., 2020). The impact of ACEs may be worse in low-resource settings like Kenya due to limited mental health infrastructure and compounding socioeconomic stressors (Arena et al., 2025). Studies have shown that over 50% of Kenyan children and youth reported having experienced at least one

ACE, with physical abuse, emotional neglect, and household dysfunction being the most prevalent. Additionally, children who were exposed to ACEs had a higher risk of developing mental health conditions like PTSD and depression (Miedema *et al.*, 2023; Ferrajão *et al.*, 2024).

Individuals exposed to multiple ACEs are substantially more likely to experience suicidal ideations and behaviors (Angelakis *et al.*, 2019; Baiden *et al.*, 2019; Burstein and Greenfield, 2020). In Kenya, young adults who experienced ACEs had significantly higher odds of poor mental health outcomes, including suicidal and self-harm behaviors, particularly among males (adjusted odds ratio = 6.7, 95% confidence interval [CI]: 2.8–16.0), highlighting ACEs as key contributors to youth suicidality in the country (Seya *et al.*, 2024). While resilience can mediate the relationship between ACEs and suicidality (Giovannelli *et al.*, 2020), its moderating function is still unclear (Zahradnik *et al.*, 2010).

Evidence also indicates that ACEs are a major predictor of PTSD (Bielas *et al.*, 2016), with 25–35% of children exposed to trauma and PTSD (John *et al.*, 2007; Wang *et al.*, 2012). Nearly 50% of Nairobi high school pupils have been found to have PTSD (Ndeti *et al.*, 2007), which is consistent with studies in Nigeria (Akinrogunde, 2023) and Uganda (McMullen *et al.*, 2012). High levels of ACE exposure are also experienced by South African women, which may have an impact on the development of PTSD (Jewkes *et al.*, 2010) and in Kenyan youth (Ferrajão *et al.*, 2024).

ACEs have also been linked to depression (Turner *et al.*, 2006; Bethell *et al.*, 2014), with research from the United States, the United Kingdom, and Uganda demonstrating a robust correlation (Bellis *et al.*, 2014; Blum *et al.*, 2019; Elmore and Crouch, 2020; Houtepen *et al.*, 2020; Debowska *et al.*, 2024). In Kenya, ACEs, especially sexual abuse, are linked to depressive symptoms in young people, highlighting the necessity of focused interventions (Mutamba, 2020).

The role of resilience

Resilience is the ability to adapt positively to adversity, influenced by personal characteristics, family relationships, and more general social and environmental factors, and is associated with better mental health outcomes (Stainton *et al.*, 2019; Ungar and Theron, 2020). Resilience is a key protective factor in mental health, with evidence suggesting that it can moderate and mediate the relationship between ACEs and mental health outcomes, and that resilient individuals exposed to ACEs exhibit lower rates of depression and suicidality (Poole *et al.*, 2017; Kelifa *et al.*, 2020; Freeny *et al.*, 2021). Resilience serves as both a mediator and moderator in the relationship between ACEs versus suicidality and depression among youth (Rytilä-Manninen *et al.*, 2018; Okwori, 2022). Ugandan studies found that resilience moderated the relationship between war-related trauma and suicidal ideation, indicating that fostering moderate resilience can reduce the likelihood of suicidal behaviors in this context (Haroz *et al.*, 2013; Okello, 2014). A study in Germany revealed that social support enhances resilience, leading to better mental health outcomes among individuals with a history of ACEs (Wolff and Caravaca Sánchez, 2019). Resilience has become a protective factor against the effects of trauma on mental health (Seery *et al.*, 2010; Poole *et al.*, 2017; Clements-Nolle and Wadlington, 2019). Moreover, this study is grounded in “ordinary magic” theory of resilience, which conceptualizes resilience as a common, adaptive process rooted in individual and contextual strengths (Michael Garrett, 2016). In low-resource settings,

resilience is often linked to familial and community factors rather than individual traits alone (Banyard *et al.*, 2025). In Kenya, despite high rates of depression (Lawley, 2023), suicidal thoughts and intent (Leone, 2024), and PTSD (Kaaria *et al.*, 2024), we have limited research investigating the mediating and moderating roles of resilience in the relationship between ACEs and mental health outcomes.

Mediation and moderation

Mediation and moderation analyses are necessary for discovering the relationship between ACEs, resilience, and mental health consequences. Mediation explains how resilience serves as a pathway through which ACEs influence mental health disorders, such as depression, PTSD, and suicidality (Panagou and MacBeth, 2022; Panagou, 2021), potentially lowering the risks by mitigating emotional dysregulation caused by trauma (Elrefaay and Elyzal, 2024). Moderation examines when and under what conditions resilience alters the strength or direction of the relationship between ACEs and these mental health outcomes (Zhang *et al.*, 2023), especially within high-risk environments (Okwori, 2022). Understanding and exploring these dynamics in Kenyan youth is crucial for finding potential cultural and contextual barriers and developing effective mental health interventions.

Knowledge gap

While previous studies have explored ACEs and the links between risk factors and their impact on mental health outcomes worldwide, research gaps still exist regarding how resilience mediates and moderates these relationships in the Kenyan context, and, in particular, no studies have compared how different categories of ACEs affect depression, PTSD, and suicidality.

General aim

The general aim is to investigate the association between the different categories of ACEs and suicidality, PTSD, and depression, and the moderating and mediating roles of resilience in the outcomes of the different categories of ACEs in the Nairobi Metropolitan Area.

Specific aims

1. To describe the distribution and correlations between ACEs, resilience, suicidal ideation and behavior, PTSD, and depression outcomes in Kenyan youth.
2. To assess the moderating effect of resilience on the relationship between ACEs and suicidality, PTSD, and depression outcomes.
3. To explore the mediating role of resilience in the relationship between ACEs, suicidality, PTSD, and depression outcomes.
4. To explore if our data can inform resilience interventions to mitigate the relationship between ACEs and the outcomes of suicidality, PTSD, and depression.

This study was guided by the following hypotheses

H1: Higher levels of ACEs will be significantly associated with increased depression, PTSD, and suicidal ideation and behavior in Kenyan youth.

H2: Resilience will moderate the relationship between ACEs and mental health outcomes (suicidality, PTSD, and depression).

H3: Resilience will mediate the relationship between ACEs and mental health outcomes (suicidality, PTSD, and depression).

H4: Patterns observed in the data will identify key resilience factors that can inform the development of interventions aimed at mitigating the effects of ACEs on suicidality, PTSD, and depression.

Methodology

Study design and setting

This was a cross-sectional study design conducted in Nairobi metropolitan capital city and Kiambu County (peri-urban). Participants from the community, high school level, and middle-level college located within the metropolitan area were recruited into the study.

Sampling

This study utilized a population-based convenience sampling technique. The inclusion criteria were all youths aged 14–25 years who were available, willing, and residing within the designated neighborhoods. While recruitment was not randomized, it was comprehensive and inclusive, leveraging local community leaders and school and college administrations to reach a broad cross-section of the population. The exclusion criteria were applied to individuals outside the designated age range and those unwilling to participate. Individuals incapable of comprehending the questionnaire due to factors such as intoxication or illiteracy were also excluded to minimize bias and incomplete data from self-report questionnaires.

Procedures

Research assistant (RA) selection and training

Twelve RAs selected through a competitive process underwent a 2-day comprehensive in-person training program that covered data collection techniques and procedures, including administering surveys. Additionally, they were trained on ethical procedures, such as obtaining informed consent and assent, ensuring confidentiality, and handling data accurately. The training also emphasized proper data handling, including real-time data verification, secure storage, and entry protocols to minimize errors. The program also included mock interviews to simulate real-world scenarios.

Recruitment and resulting sample

After receiving ethical approvals, we approached the County Commissioners in Kiambu and Nairobi Counties to explain to them the nature of the study and solicit their support. They then connected us with the chiefs and sub-chiefs who collaborated with community opinion leaders (elders) to sensitize and mobilize the youths in their areas. The chiefs, sub-chiefs, and elders were given a script that included information on the study, an invitation to the study at the designated time, date, and venue inside the study locations. They used it to actively engage through local meetings, community gatherings, and announcements at local centers, to organize and urge the youth to participate in the study. English literacy was a requirement for participation since all the questionnaires were administered in English in this context. While these criteria may limit the generalizability of findings to all youth, nobody met this exclusion criterion. However, chiefs' and sub-chiefs' ability to speak English was not a criterion for youth participation, as

communication with potential participants was handled directly by the RAs. We assigned a number (1–12) per RA, with each number forming a distinct group. Participants were then assigned to the group in a restricted randomization, where a printed voucher with numbers 1–12 on each page was used. This ensured balanced group sizes and facilitated smooth distribution of the questionnaires, as well as efficient and simultaneous data collection across groups. Each group comprised a maximum of 25 participants and was led by the respective RA assigned to that group. The RAs distributed questionnaires, verified ages, and collected written informed consent and assents. To ensure standardization, participants responded to the questions based on their understanding, thereby minimizing any potential influence from external assistance.

A total of 1,972 participants participated in the study. RAs explained the study's goals, methods, risks, rewards, and confidentiality. Participants had the right to withdraw without facing consequences before providing their consent. None of the participants declined participation. Adults affirmed their voluntary participation as per ethical guidelines by signing informed consent and assent for those below 18 years, which is the legal age in Kenya for minors. Minors were accompanied by a parent or a legal guardian for consent only, but completed the questionnaires independently, with guardians not permitted in the data collection area. Refunds were made for any travel expenses or lost income during their participation, and snacks were provided.

Data collection

All the data were collected during the day from September 21, 2022, to December 15, 2022. Data were collected on sociodemographics, ACEs, suicidality, depression, PTSD, and resilience. Participants completed self-administered questionnaires confidentially in group settings within social halls, similar to an exam environment where no communication occurs among participants. Additionally, to ensure privacy, each participant was provided with sufficient space to fill out the questionnaire without being able to see the responses of others to minimize social desirability bias. Students used school/college halls, while community participants used public halls in Nairobi and Kiambu that were approved for research purposes.

Instruments

Sociodemographic profile

- We collected data on age, gender, marital status, religion, birth position, level of education, employment status, primary source of income, place of abode, and whether they were sharing the living space(s).
- Wealth Index:** This was developed by the World Bank for use in low- and middle-income countries (LMICs) and uses household items to determine family socioeconomic status. It was constructed using principal component analysis based on the Demographic Health Survey model, assessing socioeconomic data on household assets, water sources, floor type, toilet facilities, and primary fuel sources for cooking (McLorg et al., 2021). The Wealth Index is classified from high to low on a 5-point Likert scale (Rutstein, 2015).

Adult resilience measure-revised (ARM-R)

The ARM-R is a self-report questionnaire designed to assess resilience from a socio-ecological lens (Liebenberg and Moore, 2018; Jefferies et al., 2019; Resilience Research Centre, 2022). The ARM-R instrument consists of 17 items capturing the protective factors at

the individual, relational, and community levels, scored on a 5-point Likert-type scale ranging from 1 = “Not at all” to 5 = “A lot”. The ARM-R measures the capacity to access resources in an individual’s environment (personal resilience) and the extent to which social entities in the environment (e.g., family, peers, and institutions) provide resources (relational resilience). All items in the questionnaire are positively worded, facilitating straightforward scoring by summing up the scores directly. Total resilience scores range from 17 to 85, with higher scores indicating greater resilience. Developed through a cross-cultural exploration involving 14 communities across 11 countries worldwide (Ungar et al., 2008), the ARM-R demonstrates good psychometric properties, including strong internal reliability/consistency, content, and face validity, and is used worldwide, that is, in Syria (Ipekci, 2021) and Brazil (Ferreira et al., 2022) and across cultures in diverse studies of resilience (Ungar et al., 2008; Jefferies et al., 2019). We examined scale reliability using Cronbach’s α for the 17 items of the ARM-R and found that the $\alpha > 0.80$, indicating good consistency (DeVellis, 1991). The Cronbach’s α for 16 years and above was 0.904, and that for <16 years was 0.898, indicating its reliability in this age group in our study (Zrihan-Weitzman et al., 2023).

The traumatic and distress scale (TADS)

The TADS, a self-report questionnaire, was used for the assessment of multiple types of ACEs (Salokangas et al., 2016). ACEs such as abuse and neglect were evaluated using the TADS items, which measure experiences across five main domains, namely emotional abuse, physical abuse, sexual abuse, emotional neglect, and physical neglect, each comprising five items. Each item is rated on a 5-point Likert scale from 0 to 4 (0 = *never*, 1 = *rarely*, 2 = *sometimes*, 3 = *often*, and 4 = *nearly always*). We explored the five TADS domains for ACEs. The TADS has excellent internal consistency within each domain and establishes its validity, reliability, and clinical utility in retrospectively assessing childhood traumatization (Salokangas et al., 2016). The TADS total score, indicative of the overall extent of childhood trauma, was calculated as the total score of all five domains (De Venter et al., 2020; Salokangas et al., 2020). The internal reliability using Cronbach’s α for all the 43 TADS items and 25 items for the ACEs domain was calculated and found to be good ($\alpha = 0.880$ and $\alpha = 0.850$, respectively).

The Columbia–suicide severity rating scale (C-SSRS)

We assessed suicidal ideations and behaviors based on self-reported C-SSRS (Posner et al., 2011). Indeed, C-SSRS does not introduce new criteria for suicidality and, at best, brings all of them into one tool. The C-SSRS has been validated in a study of the Lebanese population, indicating its applicability in a variety of LMIC contexts, despite the lack of research on suicidality assessment instruments in these contexts (Zakhour et al., 2021). It measures four constructs to distinguish between the domains of suicidal ideation and behavior. The “severity subscale” measures the severity of ideation using a 5-point ordinal scale ranging from 1 (*wish to be dead*) to 5 (*suicidal intent with a plan*). The “intensity subscale” consists of five items, each rated on a 5-point ordinal scale, assessing frequency, duration, controllability, deterrents, and reason for ideation. The “behavior subscale” includes categories such as actual, aborted, and interrupted attempts, preparatory behavior, and non-suicidal self-injurious behavior. The “lethality subscale” rates actual attempts on a 6-point ordinal scale, with potential lethality rated if actual lethality is 0. Psychometric evaluations have consistently demonstrated the scale’s reliability, internal consistency, and construct validity, with evidence supporting its ability to accurately

predict suicide risk across different populations (Yershova et al., 2016; Austria-Corrales et al., 2023; Schwartzman et al., 2023). We examined the internal consistency of the C-SSRS Cronbach’s α , and found the internal consistency for lifetime and recent versions was $\alpha = 0.821$ and $\alpha = 0.801$, respectively, which demonstrated good reliability.

Post-traumatic stress disorder

The Harvard Trauma Questionnaire (HTQ) is a four-part instrument that focuses on (i) experiences of torture and other traumas frequently experienced; (ii) a subjective description of the most severe traumatic event(s) experienced; (iii) events associated with head injuries; and (iv) symptoms of PTSD and expressions of functional distress. It was developed in the early 1990s as a cross-cultural screening instrument to document trauma exposure, head trauma, and trauma-related symptoms. While the original tool focused on refugee segments (Indo-Chinese refugee populations), where it exhibited strong psychometric properties, it is increasingly being used in other contexts, including low-prevalence community samples (Mollica et al., 1992). It has since been translated and validated in several languages for many regions and population samples (Halepota and Wasif, 2001; Lhewa et al., 2007; Oruc et al., 2008; de Fouchier et al., 2012). The first 16 items were derived from the Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV) criteria for PTSD. The DSM-IV PTSD score is calculated from the first 16 items. The higher the scores on the DSM-IV PTSD items, the more likely it is that the respondent will have a PTSD diagnosis. Response options were “yes” and “no.” Cumulative trauma exposure scores were calculated by a count of “yes” responses to all items.

Patient Health Questionnaire-9

The PHQ-9 is a nine-item self-report questionnaire designed to measure depression (Kroenke and Spitzer, 2002). Items are rated on a 4-point Likert scale, ranging from 0 (*not at all*) to 3 (*nearly every day*). The cumulative score can range from 0 to 27, with higher scores indicative of more pronounced depression. The questions address sleep, energy, appetite, and other possible symptoms of depression by asking respondents how often they have “been bothered by any of the following problems” in the past 2 weeks. Originally validated in the United States, the PHQ-9 demonstrates robust psychometric properties and has been extensively employed in culturally diverse settings (Kroenke et al., 2001). Interpretation of the total score categorizes depression severity into minimal (0–4), mild (5–9), moderate (10–14), moderately severe (15–19), or severe (20–27) categories. The nine items of the PHQ-9 demonstrated good internal consistency in this study, with Cronbach’s α of 0.814, indicating a reliable measure of depressive symptoms.

Ethics

All procedures involving human subjects/patients were approved by the Nairobi Hospital Ethics Research Committee (Approval no. TNH-ERC/DMSR/ERP/022/22). The study obtained licensing from the National Commission for Science, Technology, and Innovation (NACOSTI) license NACOSTI/P/22/18097. Administrative permissions were sought from the county-level offices in Kiambu and Nairobi counties to allow engagement with community participants. Additionally, institutional approval from the colleges was obtained, as data collection involved students, ensuring compliance with institutional policies on research involving students. Informed written consent/assent was obtained from the

participants before data collection commenced. For participants younger than the age of 18 years, informed written consent to participate was obtained from their parents or legal guardians.

Statistical analysis

Data were analyzed using the Statistical Package for the Social Sciences (IBM SPSS version 25). Descriptive statistics were done to summarize the mean scores of ACEs, PTSD, depression, resilience, and suicidality. We utilized Spearman's rank correlation to evaluate the correlation between ACEs, resilience, depression, PTSD, and suicidality because it is robust to deviations from normality and effectively captures both linear and monotonic relationships. In mediation and moderation analysis, Hayes' SPSS Process Macro (version 4.2 beta) was used with ACEs as predictors and depression, PTSD, and suicidal ideation and behavior as dependent variables. Resilience was incorporated as both a moderator and mediator because of its recognized theoretical function in mitigating the psychological effects of ACEs, as it is seen as a critical component in improving coping strategies and lowering emotional discomfort (Morgan et al., 2021).

Moderation: An interaction term (ACEs $\times$ Resilience) was included in a regression model to assess whether resilience alters the strength of the relationship between ACEs and psychological outcomes. **Mediation:** Regression-based path analysis was conducted, exploring both the direct and indirect effects of ACEs on depression, PTSD, and suicidal ideation and behavior through resilience. CIs based on bootstrapping were applied to enhance the accuracy of identifying indirect effects, ensuring that this method is both robust and reliable. A significance level of $p < 0.05$ was used for all analyses, and effect sizes with 95% CIs were reported to evaluate the strength and robustness of the relationships. Hayes' Process Macro was chosen for its ability to simultaneously evaluate moderation and mediation effects, offering a comprehensive framework for analyzing both direct and complex indirect relationships (Hayes, 2018). The statistical assumptions for regression-based analyses, including linearity, homoscedasticity, and multicollinearity, were assessed before conducting the analyses. The variance inflation factors and tolerance values for the predictor, moderator, and mediator variables were below 10 and above 0.1, respectively, indicating no multicollinearity concern. In all the analyses, scores were treated as continuous variables to retain variability and statistical power (Wu and Leung, 2017). Higher scores indicated higher experiences or symptoms.

Results

Social demographics

The study included 1,972 participants (55% female, mean age = 20.49 ± 2.63 , range = 14–25 years). Most participants were from urban settings (91.5%), were Christians (87.2%), and were unmarried (60.4%). The unemployment rate was high (77.9%) since most participants were in school. More than half (53.1%) were in the lowest quintiles (1 and 2) (Table 1).

Descriptive statistics and correlations for ACEs, resilience, depression, suicidal ideation and behavior, and PTSD

There were higher levels of negligence than abuse among the participants. All five ACEs are positively correlated to one another, as well as with PTSD ($r = 0.101$ – 0.205 , $p < 0.05$) and depression ($r = 0.101$ –

Table 1. Social demographics

Variables	Categories	Total (N = 1,972)
Age	Mean $\pm$ SD (min, max)	20.49 $\pm$ 2.631 (14,25)
Region	Peri-urban	168 (8.5%)
	Urban	1,804 (91.5%)
Gender	Female	1,069 (55.0%)
	Male	876 (45.0%)
Religion	Protestant	816 (43.9%)
	Catholic	805 (43.3%)
	Muslim	103 (5.5%)
	Other	135 (7.3%)
Level of education	Primary	179 (9.1%)
	Secondary	1,031 (52.3%)
	Tertiary	623 (31.7%)
	University	135 (6.9%)
Current marital status	Never married	1,182 (60.4%)
	In a relationship	428 (21.9%)
	Married/cohabiting	225 (11.5%)
	Separated (not divorced)	39 (2.0%)
	Divorced	16 (0.8%)
	Widowed	8 (0.4%)
	Other	59 (3.0%)
Wealth Index	1. Lowest quintile	284 (15.3%)
	2	703 (37.8%)
	3	681 (36.7%)
	4	166 (8.9%)
	5. Highest quintile	24 (1.3%)
Current employment status	Unemployed	1,531 (77.9%)
	Volunteering	225 (11.5%)
	Part-time employment	75 (3.8%)
	Full-time employment	134 (6.8%)
Living arrangement	Alone	431 (22.2%)
	Both parents	628 (32.3%)
	Single parent (father or mother)	479 (24.7%)
	Sibling(s)	72 (3.7%)
	Partner	151 (7.8%)
	Son(s)/daughter(s)	29 (1.5%)
	Non relative(s) (friends, housemate)	41 (2.1%)
	Extended family	67 (3.4%)
	Colleagues (shared quarter, dormitory)	34 (1.7%)
	Other	11 (0.6%)
Place of abode	Owned by me	61 (3.1%)

(Continued)

Table 1. (Continued)

Variables	Categories	Total (N = 1,972)
	Hosted by a relative	84 (4.3%)
	Hosted by a non-relative	33 (1.7%)
	Rented house	1,228 (62.6%)
	Family house	475 (24.2%)
	College hostel	41 (2.1%)
	Rented private students' hostel/ dormitory	27 (1.4%)
	Others	12 (0.6%)

Note: The Wealth Index was constructed using the DHS standard and utilized the socioeconomic data on assets and utilities.

0.389, $p < 0.05$). Resilience was negatively correlated with PTSD ($r = -0.141$, $p < 0.05$), depression ($r = -0.361$, $p < 0.05$), suicidal behavior (recent) ($r = -0.198$, $p < 0.05$), and suicidal ideation (recent) ($r = -0.187$, $p < 0.05$). Additionally, PTSD and depression were positively correlated ($r = 0.332$, $p < 0.05$), as were depression and suicidal ideation lifetime ($r = 0.302$, $p < 0.05$) (Table 2).

Moderation analysis

Resilience as a moderator between ACEs and PTSD, depression, and suicidality (ideation and behavior)

Resilience had a significant negative relationship with all three outcomes ($p < 0.05$), affirming its protective role. The associations of interactions show a complex dynamic (Table 3):

- **PTSD:** Resilience moderated the association of emotional and physical negligence and sexual abuse on PTSD. Higher resilience lowers the negative influence of ACEs on PTSD symptoms ($p < 0.05$).
- **Depression:** Resilience acted as a buffer against depression, showing significant negative associations across ACE domains, such as emotional and physical negligence ($p < 0.05$).
- **Suicidality:** Emotional negligence ($\beta = 0.0022$, $p = 0.0059$) and physical negligence ($\beta = 0.0026$, $p = 0.0179$) showed a significant interaction effect on lifetime suicidal ideation. Additionally, resilience showed a borderline significant moderation for physical abuse on lifetime suicidal behavior ($\beta = -0.0024$, $p = 0.0479$) and recent suicidal behavior ($\beta = -0.0017$, $p = 0.05$) and total ACEs on recent suicidal behavior ($\beta = -0.0004$, $p = 0.0514$).

Mediation analysis

Pathways from ACEs to resilience to depression, PTSD and suicidality ideation and behavior

We report standardized β -coefficients for key paths in the moderation and mediation models, which ranged from small to medium effect sizes (Table 4).

- **ACEs and PTSD through resilience:** Resilience partially mediated the relationship between most ACEs (direct and indirect effects; $p < 0.05$) and PTSD symptoms. However, for the total ACE score, the indirect effect on PTSD was nonsignificant ($p > 0.05$), indicating no mediation. Emotional negligence demonstrated

complete mediation, as resilience absorbs the entire effect (indirect effect significant and direct effect nonsignificant).

- **ACEs and depression through resilience:** While all ACE domains showed significant direct and indirect effects (partial mediation), emotional and physical abuse revealed the strongest direct effects, underscoring resilience's partial mediation across most domains.
- **ACEs and suicide ideation through resilience:** Resilience partially mediated the relationship between ACEs and lifetime/recent suicide ideation for most ACE domains. However, emotional and physical negligence demonstrated complete mediation for lifetime ideation, while for recent suicidal ideation, resilience fully mediated the effects of physical abuse and negligence, while partial mediation was observed for all other ACEs, suggesting resilience can entirely offset the adverse impacts of these ACEs on ideation.
- **ACEs and suicide behavior through resilience:** Resilience fully mediated the effects of ACEs on suicidal behaviors for most domains ($p < 0.001$, indirect effects, nonsignificant direct effects). Partial mediation is observed for ACE domains, such as emotional, physical, and sexual abuse, in relation to lifetime suicidal behavior.

Discussion

Our study presents a comprehensive investigation in Kenya of the link between different categories of ACEs and depression, suicidality, and PTSD, with consideration of the role of resilience as a mediator or moderator in youth living in the Nairobi Metropolitan Area. We found a strong correlation (ACEs and symptoms of depression, PTSD, and suicidality). The relationship between ACEs and depression and PTSD was significantly moderated by resilience and partially mediated by resilience, indicating a role in protecting the mental health of Kenyan youth. While findings may offer insights relevant to similar urban and peri-urban youth settings in Kenya and comparable low-resource contexts, the nonrandom sampling approach limits broader generalizability. We stress that resilience is crucial in mitigating these negative effects of ACEs on outcomes and advocate for its integration into interventions.

Social demographics

All the participants were in their youth (up to 25 years). The majority lived in the urban area where most of our research sites were located; there were more females than males, which reflects increased enrollment of females in schools, an emerging observation in the Kenyan education system. Most were Christians, which is the predominant religion in Kenya, and most were unmarried since they were students and came from a lower socioeconomic status.

Moderation and mediation interpretations

ACEs, mental health (PTSD, suicidality, and depression), and the role of resilience

Our findings align with previous global and regional studies, which showed a direct positive relationship between ACEs and adverse mental health outcomes (PTSD, suicidality, and depression) (Ndetei et al., 2007; Jewkes et al., 2010; McMullen et al., 2012; Bielas et al., 2016; Angelakis et al., 2019; Baiden et al., 2019; Blum et al., 2019; Houtepen et al., 2020; Debowska et al., 2024; White et al., 2024). Further, we found that Kenyan youth exposed to ACEs exhibited high levels of depressive symptoms, with resilience that negatively

Table 2. Descriptive statistics (mean and standard deviations) and correlations for ACEs, depression, resilience, PTSD, suicide ideations and behavior

Measures	Mean	SD	Correlations												
			1	2	3	4	5	6	7	8	9	10	11	12	13
ACEs															
1. Emotional negligence	5.64	4.519	1												
2. Emotional abuse	3.32	3.491	.345**	1											
3. Physical negligence	4.64	3.408	.546**	.413**	1										
4. Physical abuse	3.1	3.453	.267**	.525**	.409**	1									
5. Sexual abuse	1.71	3.165	.236**	.341**	.287**	.450**	1								
6. Total ACE scores	17.76	12.646	.719**	.733**	.745**	.726**	.617**	1							
7. Resilience	64.36	14.374	−.479**	−.352**	−.384**	−.311**	−.210**	−.502**	1						
8. PTSD	6.44	2.559	.101**	.184**	.116**	.189**	.147**	.205**	−.141**	1					
9. Depression (PHQ–9)	7.45	5.576	.221**	.389**	.228**	.330**	.209**	.379**	−.361**	.332**	1				
10. Suicide Ideation Lifetime Score	0.2	0.397	.158**	.242**	.099**	.173**	.165**	.241**	−.208**	.127**	.302**	1			
11. Suicide Ideation Recent Score	0.12	0.82	.169**	.178**	.113**	.142**	.136**	.203**	−.187**	.094**	.259**	.607**	1		
12. Suicide behavior lifetime	0.19	0.39	.158**	.208**	.129**	.178**	.180**	.223**	−.232**	.114**	.218**	.458**	.381**	1	
13. Suicidality behavior recent	0.12	0.322	.170**	.181**	.132**	.164**	.166**	.208**	−.198**	.090**	.178**	.378**	.434**	.692**	1

Note: SD, standard deviation. The correlations show the strength and direction of relationships between the variables. The correlation marked (**) indicates significance at $p < 0.05$.

Table 3. Resilience as a moderator between ACES and PTSD, depression (PHQ-9), suicidal ideations, and behavior

	PTSD		Depression (PHQ-9)		Lifetime suicidal ideation		Recent suicidal ideation		Lifetime suicidal behavior		Recent suicidal behavior	
	estimates (95% CI)	p-value	Estimates (95% CI)	p-value	Estimates (95% CI)	p-value	Estimates (95% CI)	p-value	Estimates (95% CI)	p-value	Estimates (95% CI)	p-value
1. Emotional negligence												
Emotional negligence	−0.1390 (−0.2804, 0.0023)	0.0539	−0.5081 (−0.757, −0.2591)	<0.0001	−0.1092 (−0.2096, −0.0089)	0.0329	−0.0076 (−0.0922, 0.0769)	0.8596	0.0692 (−0.0422, 0.1805)	0.2231	0.0840 (0.0041, 0.1640)	0.0394
Resilience	−0.0393 (−0.0578, −0.0208)	<0.0001	−0.1944 (−0.2278, −0.1611)	<0.0001	−0.0322 (−0.0465, −0.018)	<0.0001	−0.0141 (−0.0259, −0.0022)	0.0199	−0.0239 (−0.0382, −0.0097)	0.001	−0.0096 (−0.0198, −0.0005)	0.0632
EN × Resilience	0.0027 (0.0005, 0.0049)	0.0153	0.0094 (0.0056, 0.0133)	<0.0001	0.0022 (0.0006, 0.0038)	0.0059	0.0007 (−0.0007, 0.002)	0.3385	−0.0007 (−0.0024, 0.001)	0.408	−0.0011 (−0.0024, −0.0001)	0.0677
2. Emotional abuse												
Emotional abuse	−0.0176 (−0.1934, 0.1582)	0.8442	0.4569 (0.1421, 0.7716)	0.0045	0.1209 (−0.008, 0.2498)	0.0659	0.1177 (0.0113, 0.2242)	0.0303	0.1035 (−0.0421, 0.2492)	0.1633	0.1000 (−0.0073, 0.2074)	0.0677
Resilience	−0.0231 (−0.0373, −0.0088)	0.0015	−0.1027 (−0.127, −0.0784)	<0.0001	−0.0111 (−0.0217, −0.0005)	0.0396	−0.0063 (−0.0153, 0.0026)	0.165	−0.023 (−0.0339, −0.0121)	<0.0001	−0.0109 (−0.0188, −0.0029)	0.0073
EA × Resilience	0.0022 (−0.0006, 0.0050)	0.1271	0.0007 (−0.0043, 0.0058)	0.7834	−0.0002 (−0.0023, 0.0019)	0.8484	−0.0011 (−0.0028, 0.0006)	0.2173	−0.0005 (−0.0028, 0.0018)	0.6514	−0.0013 (−0.0029, 0.0004)	0.1398
3. Physical negligence												
Physical negligence	−0.1389 (−0.3210, 0.0432)	0.1349	−0.4678 (−0.7897, −0.1459)	0.0044	−0.1553 (−0.2852, −0.0255)	0.0191	−0.0261 (−0.1346, 0.0823)	0.6363	−0.0662 (−0.2155, 0.083)	0.3842	0.0162 (−0.0935, 0.1259)	0.7724
Resilience	−0.0370 (−0.0549, −0.0191)	0.0001	−0.1803 (−0.2128, −0.1479)	<0.0001	−0.0339 (−0.0477, −0.0202)	<0.0001	−0.0172 (−0.0288, −0.0055)	0.004	−0.0354 (−0.0494, −0.0214)	<0.0001	−0.0166 (−0.0269, −0.0064)	0.0015
PN × Resilience	0.0031 (0.0003, 0.0060)	0.0303	0.0102 (0.0051, 0.0152)	0.0001	0.0026 (0.0004, 0.0047)	0.0179	0.0007 (−0.0011, 0.0025)	0.466	0.0013 (−0.001, 0.0035)	0.2809	−0.0003 (−0.0020, 0.0014)	0.7053
4. Physical abuse												
Physical abuse	0.0062 (−0.1584, 0.1708)	0.9413	0.379 (0.069, 0.689)	0.0166	0.0696 (−0.0616, 0.2008)	0.298	0.1201 (0.0124, 0.2278)	0.0289	0.2005 (0.048, 0.3531)	0.010	0.1140 (0.0073, 0.2207)	0.0363
Resilience	−0.0228 (−0.0366, −0.0089)	0.0013	−0.1159 (−0.1404, −0.0914)	<0.0001	−0.0167 (−0.0275, −0.0059)	0.0025	−0.0084 (−0.0176, 0.0009)	0.0782	−0.02 (−0.0313, −0.0087)	0.0005	−0.0125 (−0.0204, −0.0047)	0.0018
PA × Resilience	0.0018 (−0.0008, 0.0045)	0.1787	0.0004 (−0.0046, 0.0054)	0.8706	−0.0002 (−0.0024, 0.002)	0.8536	−0.0016 (−0.0034, 0.0002)	0.0818	−0.0024 (−0.0048, 0)	0.0479	−0.0017 (−0.0034, −0.0000)	0.05
5. Sexual abuse												
Sexual abuse	−0.1963 (−0.3973, −0.0046)	0.0555	0.2274 (−0.1414, 0.5961)	0.2267	0.1396 (0.0028, 0.2764)	0.0455	0.0653 (−0.0464, 0.177)	0.2514	0.1839 (0.0223, 0.3456)	0.0258	0.0645 (−0.0451, 0.1742)	0.2484
Resilience	−0.0305 (−0.0420, −0.0190)	<0.0001	−0.1359 (−0.1566, −0.1152)	<0.0001	−0.0155 (−0.0239, −0.007)	0.0004	−0.0129 (−0.0201, −0.0058)	0.0004	−0.0271 (−0.0358, −0.0183)	<0.0001	−0.0154 (−0.0220, −0.0088)	<0.0001

(Continued)

Table 3. (Continued)

	PTSD		Depression (PHQ-9)		Lifetime suicidal ideation		Recent suicidal ideation		Lifetime suicidal behavior		Recent suicidal behavior	
	estimates (95% CI)	p-value	Estimates (95% CI)	p-value	Estimates (95% CI)	p-value	Estimates (95% CI)	p-value	Estimates (95% CI)	p-value	Estimates (95% CI)	p-value
SA × Resilience	0.0049 (0.0017, 0.0082)	0.0028	0.0004 (−0.0056, 0.0064)	0.9019	−0.001 (−0.0033, 0.0012)	0.3635	−0.0004 (−0.0023, 0.0014)	0.6476	−0.002 (−0.0046, 0.0006)	0.1257	−0.0007 (−0.0025, 0.0010)	0.4117
6. Total ACEs domain scores												
Total ACE scores	−0.0364 (−0.0878, 0.0150)	0.1654	0.0111 (−0.0836, 0.1058)	0.8186	0.0109 (−0.0285, 0.0503)	0.5877	0.0286 (−0.0027, 0.0598)	0.0731	0.0346 (−0.0084, 0.0775)	0.1149	0.0323 (0.0040, 0.0605)	0.0252
Resilience	−0.0329 (−0.0531, −0.0128)	0.0014	−0.1306 (−0.1669, −0.0942)	<0.0001	−0.0139 (−0.0301, 0.0024)	0.0937	−0.004 (−0.0171, 0.009)	0.5429	−0.0182 (−0.0341, −0.0023)	0.0248	−0.0076 (−0.0182, 0.0030)	0.1591
T ACEs × Resilience	0.0012 (0.0004, 0.0020)	0.0046	0.0018 (0.0003, 0.0033)	0.0199	0.0002 (−0.0004, 0.0009)	0.4566	−0.0002 (−0.0007, 0.0003)	0.3652	−0.0002 (−0.0009, 0.0004)	0.4637	−0.0004 (−0.0009, 0.0000)	0.0514

Note: EA, emotional abuse; EN, emotional negligence; PA, physical abuse; PN, physical negligence; SA, sexual abuse; TACEs, total ACE scores. **Dependent variables** are PTSD, depression (PHQ-9), and suicidal ideation and behavior assessed across lifetime and recent. **Predictor variables** include emotional/physical/sexual neglect or abuse, total ACEs domain scores, and resilience. **Estimates (95% CI)**: The regression coefficients representing the relationship between the predictor variable and the outcome variable, along with the 95% confidence interval (CI). In some models, ACEs appear to be negatively associated with outcomes due to variable centering and interaction inclusion. These should not be interpreted as protective effects. Rather, they reflect conditional associations at average resilience levels. The interaction term often captures the key variance when moderation is present and does not contradict the established positive link between ACEs and poor mental health. Interaction terms (ACEs × Resilience) should be interpreted in context. When the main effect of ACEs is positive, a negative interaction indicates that resilience buffers the risk, while a positive interaction suggests amplification. When the main effect is negative, a positive interaction weakens that effect (possibly protective), and a negative interaction strengthens it (possibly harmful). The significant p-values are bolded.

Table 4. Mediation analysis: Examining how ACEs associate with mental health outcomes (PTSD, depression, suicidal ideation and behavior) both directly and indirectly through resilience

Effect	PTSD		Depression (PHQ-9)		Lifetime suicidal ideations		Recent suicidal ideations		Lifetime suicidal behavior		Recent suicidal behavior	
	Coefficient (95% CI)	p-value	Coefficient (95% CI)	p-value	Coefficient (95% CI)	p-value	Coefficient (95% CI)	p-value	Coefficient (95% CI)	p-value	Coefficient (95% CI)	p-value
Emotional negligence												
Total effect	0.063 (0.03, 0.095)	<0.001	0.274 (0.213, 0.336)	<0.001	0.053 (0.028, 0.078)	<0.001	0.047 (0.027–0.068)	<0.001	0.066 (0.041–0.091)	<0.001	0.036 (0.018–0.055)	<0.001
Direct effect	0.03 (–0.007, 0.067)	0.1076	0.075 (0.008, 0.142)	0.028	0.026 (–0.002, 0.055)	0.072	0.032 (0.008–0.056)	0.008	0.024 (–0.004 to 0.051)	0.093	0.012 (–0.008 to 0.032)	0.248
Indirect effect	0.033 (0.014, 0.051)	<0.001	0.199 (0.162, 0.241)	<0.001	0.027 (0.012, 0.042)	<0.001	0.015 (0.002–0.029)	<0.001	0.042 (0.026–0.060)	<0.001	0.025 (0.012–0.038)	<0.001
Emotional abuse												
Total effect	0.138 (0.099, 0.177)	<0.001	0.644 (0.571, 0.717)	<0.001	0.124 (0.094, 0.153)	<0.001	0.067 (0.043–0.091)	<0.001	0.106 (0.076–0.136)	<0.001	0.043 (0.021–0.065)	<0.001
Direct effect	0.115 (0.074, 0.157)	<0.001	0.5 (0.424, 0.575)	<0.001	0.109 (0.078, 0.14)	<0.001	0.053 (0.027–0.078)	<0.001	0.071 (0.039–0.103)	<0.001	0.021 (–0.002 to 0.044)	0.072
Indirect effect	0.023 (0.007, 0.039)	<0.001	0.144 (0.112, 0.181)	<0.001	0.015 (0.006, 0.026)	<0.001	0.014 (0.005–0.024)	<0.001	0.035 (0.020–0.053)	<0.001	0.022 (0.010–0.036)	<0.001
Physical negligence												
Total effect	0.09 (0.048, 0.131)	<0.001	0.365 (0.285, 0.445)	<0.001	0.03 (–0.003, 0.063)	0.070	0.036 (0.009–0.064)	0.009	0.059 (0.027–0.091)	<0.001	0.022 (–0.002 to 0.046)	0.07
Direct effect	0.056 (0.012, 0.101)	0.0126	0.158 (0.076, 0.241)	<0.001	–0.004 (–0.039, 0.031)	0.815	0.013 (–0.017 to 0.042)	0.399	0.014 (–0.020 to 0.048)	0.43	–0.004 (–0.029 to 0.021)	0.727
Indirect effect	0.033 (0.016, 0.052)	<0.001	0.207 (0.166, 0.25)	<0.001	0.035 (0.021, 0.05)	<0.001	0.024 (0.011–0.038)	<0.001	0.045 (0.027–0.065)	<0.001	0.027 (0.014–0.041)	<0.001
Physical abuse												
Total effect	0.136 (0.096, 0.176)	<0.001	0.549 (0.472, 0.627)	<0.001	0.08 (0.048, 0.112)	<0.001	0.047 (0.020–0.073)	0.001	0.085 (0.053–0.118)	<0.001	0.033 (0.010–0.056)	0.005
Direct effect	0.115 (0.073, 0.157)	<0.001	0.404 (0.326, 0.482)	<0.001	0.058 (0.024, 0.091)	0.001	0.028 (–0.000 to 0.056)	0.051	0.050 (0.017–0.084)	0.003	0.010 (–0.014 to 0.033)	0.409
Indirect effect	0.021 (0.007, 0.036)	<0.001	0.145 (0.112, 0.182)	<0.001	0.022 (0.012, 0.035)	<0.001	0.019 (0.009–0.031)	<0.001	0.035 (0.021–0.051)	<0.001	0.023 (0.011–0.036)	<0.001
Sexual abuse												
Total effect	0.123 (0.08, 0.166)	<0.001	0.381(0.295,0.466)	<0.001	0.091(0.059,0.124)	<0.001	0.051 (0.024–0.078)	<0.001	0.088 (0.055–0.121)	<0.001	0.035 (0.012–0.059)	0.004
Direct effect	0.103 (0.059, 0.147)	<0.001	0.25 (0.167, 0.332)	<0.001	0.078 (0.045, 0.111)	<0.001	0.040 (0.013–0.067)	0.004	0.060 (0.028–0.093)	<0.001	0.020 (–0.004 to 0.044)	0.104

(Continued)

Table 4. (Continued)

Effect	PTSD		Depression (PHQ-9)		Lifetime suicidal ideations		Recent suicidal ideations		Lifetime suicidal behavior		Recent suicidal behavior	
	Coefficient (95% CI)	<i>p</i> -value	Coefficient (95% CI)	<i>p</i> -value	Coefficient (95% CI)	<i>p</i> -value	Coefficient (95% CI)	<i>p</i> -value	Coefficient (95% CI)	<i>p</i> -value	Coefficient (95% CI)	<i>p</i> -value
Indirect effect	0.02 (0.01, 0.032)	<0.001	0.131 (0.096, 0.17)	<0.001	0.013 (0.006, 0.023)	<0.001	0.011 (0.004–0.019)	<0.001	0.027 (0.017–0.040)	<0.001	0.016 (0.007–0.026)	<0.001
Total ACE score domain												
Total effect	0.041 (0.029, 0.052)	<0.001	0.174 (0.152, 0.196)	<0.001	0.03 (0.021, 0.039)	<0.001	0.020 (0.012–0.027)	<0.001	0.032 (0.023–0.041)	<0.001	0.014 (0.008–0.020)	<0.001
Direct effect	0.036 (0.022, 0.049)	<0.001	0.12 (0.095, 0.144)	<0.001	0.025 (0.015, 0.036)	<0.001	0.015 (0.006–0.023)	0.001	0.019 (0.009–0.029)	<0.001	0.005 (–0.002 to 0.012)	0.153
Indirect effect	0.005 (–0.002, 0.012)	<i>p</i> > 0.05	0.054 (0.04, 0.069)	<0.001	0.005 (0, 0.01)	<0.001	0.005 (0.000–0.010)	<0.001	0.013 (0.006–0.021)	<0.001	0.009 (0.003–0.015)	<0.001

Note: Dependent variables: PTSD, depression, lifetime and recent suicidal ideations and behavior. Mediator (resilience). Independent variables (ACEs). Total effect: The “total effect of ACEs on the dependent variable” represents the combined direct and indirect effects of ACE on the dependent variable. Direct effect: The “direct effect of ACEs on the dependent variable” is the effect of ACE on the dependent variable when resilience is not considered. Indirect effect: The “indirect effect of ACE on the dependent variable through resilience” represents the mediated effect of ACE on the dependent variable through the mediator “Resilience.” The significant *p*-values are bolded.

correlated with both ACEs and depression, as has been found in other countries (Abbott and Slack, 2021; Girma *et al.*, 2021).

In relation to the three mental health outcomes in our study, resilience serves as a mediator and a moderator, although to varying degrees, and all agree with the global literature summarized under the Introduction section:

- i. Moderating role: While resilience moderated the effects of ACEs on depression and PTSD, there are conflicting results in suicidality. Resilience moderated specific ACE-suicidality links, including neglect, abuse, and total ACEs, but was less protective in cases involving emotional or sexual abuse.
- ii. Mediating role: The connection between ACEs and the three mental health outcomes was largely mediated by resilience, except for emotional neglect, where it fully mediated suicidality and PTSD. This implies that emotional neglect may have a greater impact on mental health through pathways linked to resilience than through actual psychological discomfort.

These findings emphasize the necessity of adjusting resilience-building interventions to certain ACE types, acknowledging that certain adversities can call for more involved or specialized methods. Those in violent and aggressive families are prone to PTSD, coupled with cultural beliefs and practices that tolerate and accept family violence and physical punishment, as was found by Fulu *et al.* (2017). Emotional and physical neglect, the most frequent ACEs, impede stress buffering and the development of secure attachment, increasing PTSD, as found by Margolin and Vickerman (2007) and Spinazzola *et al.* (2014).

It is noteworthy that resilience negatively correlates with ACEs and PTSD, suggesting that individuals with higher resilience levels are less susceptible to PTSD symptoms. This concurs with similar findings (Reich *et al.*, 2010; Biggs *et al.*, 2024). Notably, resilience moderates the relationship between ACEs and PTSD, particularly in mitigating the association of emotional and physical negligence and sexual abuse. These results align with existing literature, which underscores the importance of resilience as a protective factor against PTSD (Seery *et al.*, 2010; Rutter, 2013). Additionally, resilience partially mediated the relationship between all ACEs and PTSD symptoms, except emotional neglect, where it fully mediated this relationship, which aligns with the findings of previous studies (Guo *et al.*, 2021).

Resilience moderated the relationship between emotional and physical negligence with lifetime and recent suicidal ideation in agreement with studies from China and the United States, which emphasize resilience as a key factor in reducing the risk of suicidality in individuals exposed to ACEs (Thompson *et al.*, 2019; Yu *et al.*, 2021). However, resilience did not significantly moderate the effects of emotional and sexual abuse on suicidal behaviors, highlighting the variability in its protective effects across different types of ACEs, as has been found in other studies (Brodsky and Stanley, 2008; Dahlby, 2022). Notably, resilience demonstrated minimal moderating effects on suicidality, particularly for emotional and sexual abuse, suggesting that suicidality may operate through distinct pathways less responsive to general resilience mechanisms. For example, acute life stressors, perceived burdensomeness, thwarted belongingness, or the lack of care and peer support may have more immediate effects on suicidal thoughts and behavior (Glenn *et al.*, 2022). In addition, development and psychosocial pathways to suicidality may differ from PTSD and depression and thus may require other protective mechanisms, such as emotional regulation, connectedness, or specific cognitive coping strategies (Sher, 2019). These distinctions imply that ACEs measured in this

study may not be sufficient to buffer the effect of ACEs on suicidality alone and suggest that there is a need for more specific interventions that address the suicide-specific risk and protective factors. Pathways showing full mediation, such as emotional neglect leading to PTSD or suicidality, may reflect resilience mechanisms like emotional regulation and perceived support, which are more directly disrupted by neglect. In contrast, partial mediation in abuse-related pathways suggests additional unmeasured factors – such as trauma severity or interpersonal betrayal – may also influence outcomes beyond what resilience alone can buffer. The non-significant moderation effect of resilience on suicidality may suggest that suicidal ideation in this context is less influenced by internal coping mechanisms and more shaped by external stressors or acute trauma. Future studies should explore additional moderators such as peer support or access to mental health care. The observed partial mediation suggests that while resilience explains part of the relationship between ACEs and mental health outcomes, other pathways are also at play. This is consistent with ecological systems theory, which posits that individual outcomes are shaped by multiple interacting influences. Furthermore, resilience acted as both a partial and complete mediator in the relationship between ACEs and suicidal ideation or behaviors, depending on the type of adversity, which agrees with the prior study (Giovannelli *et al.*, 2020). For instance, resilience fully mediated the effects of emotional negligence on recent suicidal ideation while partially mediating the effects of emotional abuse, physical abuse, and sexual abuse ($p < 0.001$).

Resilience significantly moderates the relationship between ACEs and the PHQ-9 assessment tool. This suggests that the relationship between these ACEs and depression was much more pronounced in people with lower resilience levels, which agrees with prior literature (Poole *et al.*, 2017). Moreover, our findings show that resilience partially mediated the relationship between depression and ACEs, indicating that while it reduces the adverse effects of childhood adversity, it does not eliminate them. This aligns with broader research highlighting resilience as a critical but insufficient factor in addressing mental health outcomes related to ACEs (Luthar and Zelazo, 2003; Clements-Nolle and Waddington, 2019).

Comparison with regional and global literature

Our findings are consistent with other international studies, which found that resilience-building interventions yield substantial mental health benefits for clients who experienced childhood adversity (Malhi *et al.*, 2019; Ungar and Theron, 2020). Moreover, our findings align with regional studies in sub-Saharan Africa, such as Kenya (Mwiti, 2024) and Rwanda (Satinsky *et al.*, 2021), which also found ACEs to be significantly associated with poor mental health outcomes. However, consistent with global research, resilience emerged as a protective factor across settings.

Implications and recommendations

The implications and recommendations from our findings are that developing early prevention interventions like school-based programs, community-driven approaches, and policy-level interventions could prevent or lessen PTSD, suicidality, and depression among Kenyan youth, especially in impoverished and high-crime regions, which are observed in the Nairobi metropolitan area (Ndeti *et al.*, 2007; Vundi, 2022). Moreover, our results point to the protective role of resilience, but it is important that interventions tackle not only the consequences of ACEs (resilience) but also

the root causes of mental health problems. Early identification of at-risk children, prevention through trauma-informed care, as well as community-based interventions aimed toward the perpetuating factors of ACEs, poverty, violence, and parental neglect need to be the focus of prevention efforts. However, resilience-building programs should not be used to replace the prevention of ACEs.

Limitations and future directions

Cross-sectional data limits causal inference when conducting mediation and moderation analysis. Without repeated measures over time, it is not possible to determine the directionality of the relationships examined. Also, moderation analysis detects interaction effects at one point in time, yet it fails to prove which event occurs first or second between the moderator and exposure or outcome. Future studies would benefit from longitudinal approaches to examine how ACEs affect and the protective role and resilience evolve over time. They could provide insight into causal pathways and aid in timely intervention decisions for longitudinal data. While the use of convenience sampling may limit the generalizability of the findings to broader youth populations beyond the study areas, the selected sites were nonetheless representative of Kenyan youth. The English literacy requirement may have introduced selection bias by excluding less educated or marginalized youth. This could have limited the generalizability and potential skewness of findings toward more advantaged, resilient participants.

Additionally, self-report measures may introduce recall and social desirability bias. Future research should employ longitudinal controlled studies and potentially bias-free objective measures to strengthen the validity of findings. Also, guardian presence may have caused underreporting of sensitive issues, introducing social desirability bias and underestimating ACEs' impact. Finally, the use of HTQ may not align with DSM-5 or capture culturally specific trauma symptoms among Kenyan youth in community settings. Future research should use culturally adapted, updated tools for trauma assessment.

Conclusions

We conclude that our findings underscore the critical role of resilience-building interventions in mitigating the impact of ACEs on suicidality, particularly in resource-constrained settings like Nairobi. A similar conclusion was reached by other studies (Al-Sabah et al., 2015; Hamby et al., 2021; Morgan et al., 2021). Our research sheds important light on the complex relationships that exist between ACEs, resilience, and the three studied mental health outcomes in young people in Kenya, thereby addressing the knowledge gap and fulfilling our general objective and specific aims, and in the process providing positive answers to all our research questions. There is a necessity for resilience-focused interventions that are customized to each of the three mental health outcomes and adversity. The interventions should build resilience via school mental health programs, trauma-informed teaching, peer support, and counseling. Engaging caregivers and elders while honoring Kenyan cultural values like collectivism and religious coping enhances impact. Kenya may significantly reduce the long-term effects of childhood trauma on mental health by incorporating resilience-building techniques frameworks in educational institutions in the community and supported by appropriate policy to achieve better mental outcomes in mental disorders in our study. These interventions can be implemented by trained nonmental

health specialists and, in the process, achieve critical reach. Hence, reducing the risk of mental health disorders in Kenyan youth involves preventing ACEs and promoting resilience. All interventions must be holistic and take into consideration adverse environments while also teaching youth psychological tools to cope with adversity.

Open peer review. To view the open peer review materials for this article, please visit <http://doi.org/10.1017/gmh.2025.10041>.

Data availability statement. Requests for the data may be sent to the corresponding author.

Author contribution. D.M.N.: conceptualization and drafting of the paper. V.M.: oversight of data collection. C.M.: oversight of ethics. E.J.: statistical analysis and literature review. P.N.: draft review. S.W.: statistical analysis. V.O.: field work during data collection and literature review. K.B.: critique of the manuscript. D.M.: conceptualization and critique of the manuscript. All authors read and approved the final manuscript.

Financial support. This study was funded by the National Institutes of Health (NIH), grant/award number: 5R01MH127571-02.

Competing interests. The authors declare none.

Ethics statement. The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008. All procedures involving human subjects/patients were approved by the Nairobi Hospital Ethics Research Committee (approval no. TNH-ERC/DMSR/ERP/022/22). The study obtained licensing from the National Commission for Science, Technology, and Innovation (NACOSTI) license NACOSTI/P/22/18097. Informed written consent/assent was obtained from participants before data collection commenced. For participants younger than the age of 18 years, informed written consent to participate was obtained from their parents or legal guardians.

References

- Abbott M and Slack KS (2021) Exploring the relationship between childhood adversity and adult depression: A risk versus strengths-oriented approach. *Child Abuse & Neglect* 120, 105207.
- Akinrogunde PO (2023) *Adverse Childhood Experiences in the Yoruba Tribe/ Ethnic Group of Nigerians Living in America: A Phenomenological Exploration*. Repository of Liberty University.
- Al-Sabah R, Legerski J-P, Layne CM, Isakson B, Katalinski R, Pasalic H, Bosankic N and Pynoos RS (2015) Adolescent adjustment, caregiver-adolescent relationships, and outlook towards the future in the long-term aftermath of the Bosnian war. *Journal of Child & Adolescent Trauma* 8, 45–60.
- Angelakis I, Gillespie EL and Panagioti M (2019) Childhood maltreatment and adult suicidality: A comprehensive systematic review with meta-analysis. *Psychological Medicine* 49(7), 1057–1078.
- Arena JF, Adriani B, Mancino S, Sarno F, Modesti MN, Del Casale A and Ferracuti S (2025) Gender diversity and mental health: A systematic review of the impact of childhood trauma. *Psychiatry International* 6(1), 13.
- Ashaba S, Cooper-Vince CE, Maling S, Satinsky EN, Baguma C, Akena D, Nansera D, Bajunirwe F and Tsai AC (2021) Childhood trauma, major depressive disorder, suicidality, and the modifying role of social support among adolescents living with HIV in rural Uganda. *Journal of Affective Disorders Reports* 4, 100094.
- Austria-Corrales F, Jiménez-Tapia A, Astudillo-García CI, Arenas-Landgrave P, Xochihua-Tlecuitl T, Cruz-Cruz C, Rivera-Rivera I, Gómez-García JA, Palacios-Hernández B and Pérez-Amezcuea B (2023) The Columbia-suicide severity rating scale: Validity and psychometric properties of an online Spanish-language version in a Mexican population sample. *Frontiers in Public Health* 11, 1157581.

- Baiden P, Antwi-Boasiako K and den Dunnen W (2019) Effects of suicidal ideation on unmet mental health needs among young adults in Canada. *Journal of Child and Family Studies* 28, 436–446.
- Banyard V, Rousseau D, Shockley McCarthy K, Stavola J, Xu Y and Hamby S (2025) Community-level characteristics associated with resilience after adversity: A scoping review of research in urban locales. *Trauma, Violence, & Abuse* 26(2), 356–372.
- Bellis MA, Lowey H, Leckenby N, Hughes K and Harrison D (2014) Adverse childhood experiences: Retrospective study to determine their impact on adult health behaviours and health outcomes in a UK population. *Journal of Public Health* 36(1), 81–91.
- Benarous X, Raffin M, Bodeau N, Dhossche D, Cohen D and Consoli A (2017) Adverse childhood experiences among inpatient youths with severe and early-onset psychiatric disorders: Prevalence and clinical correlates. *Child Psychiatry & Human Development* 48, 248–259.
- Bethell CD, Newacheck P, Hawes E and Halfon N (2014) Adverse childhood experiences: Assessing the impact on health and school engagement and the mitigating role of resilience. *Health Affairs* 33(12), 2106–2115.
- Bethell CD, Solloway MR, Guinosso S, Hassink S, Srivastav A, Ford D and Simpson LA (2017) Prioritizing possibilities for child and family health: An agenda to address adverse childhood experiences and foster the social and emotional roots of well-being in pediatrics. *Academic Pediatrics* 17(7), S36–S50.
- Bielas H, Barra S, Skrivaneck C, Aebi M, Steinhäuser H-C, Bessler C and Plattner B (2016) The associations of cumulative adverse childhood experiences and irritability with mental disorders in detained male adolescent offenders. *Child and Adolescent Psychiatry and Mental Health* 10(1), 1–10.
- Biggs AT, Seech TR, Johnston SL and Russell DW (2024) Psychological endurance: How grit, resilience, and related factors contribute to sustained effort despite adversity. *The Journal of General Psychology*, 151, 1–43.
- Blum RW, Li M and Naranjo-Rivera G (2019) Measuring adverse child experiences among young adolescents globally: Relationships with depressive symptoms and violence perpetration. *Journal of Adolescent Health* 65(1), 86–93.
- Brodsky BS and Stanley B (2008) Adverse childhood experiences and suicidal behavior. *Psychiatric Clinics of North America* 31(2), 223–235.
- Burstein B and Greenfield B (2020) Childhood maltreatment and suicidality: One link in a tangled chain toward risk reduction. *JAMA Network Open* 3(8), e2013095–e2013095.
- Clements-Nolle K and Waddington R (2019) Adverse childhood experiences and psychological distress in juvenile offenders: The protective influence of resilience and youth assets. *Journal of Adolescent Health* 64(1), 49–55.
- Dahlby L (2022) *Mediators of Childhood Maltreatment and Suicidal Ideation: A Systematic Review and Research Proposal*. Thesis from The University of Edinburgh.
- de Fouchier C, Blanchet A, Hopkins W, Bui E, Ait-Aoudia M and Jehel L (2012) Validation of a French adaptation of the Harvard trauma questionnaire among torture survivors from sub-Saharan African countries. *European Journal of Psychotraumatology* 3(1), 19225.
- De Venter M, Illegems J, Van Royen R, Sabbe BGC, Moorkens G and Van Den Eede F (2020) The relationship between childhood trauma and the response to group cognitive-behavioural therapy for chronic fatigue syndrome. *Frontiers in Psychiatry* 11, 536.
- Debowska A, Boduszek D, Fray-Aiken C, Ochen EA, Powell-Booth KT, Nanfuka Kalule E, Harvey R, Turyomurugendo F, Nelson K and Willmott D (2024) Child abuse and neglect and associated mental health outcomes: A large, population-based survey among children and adolescents from Jamaica and Uganda. *Mental Health and Social Inclusion* 28(1), 42–65.
- DeVellis R (1991) *Scale development*. Newbury Park, CA: Sage.
- Elmore AL and Crouch E (2020) The association of adverse childhood experiences with anxiety and depression for children and youth, 8 to 17 years of age. *Academic Pediatrics* 20(5), 600–608.
- Elrefaay SMM and Elyzal AS (2024) Adverse childhood experiences and depression: The mediating role of resilience and emotional regulation. *Journal of Psychosocial Nursing and Mental Health Services* 62(1), 45–54.
- Ferrajão PC, Tourais B, Faria I, Dias J and Elklit A (2024) A latent class analysis of adverse life events for Kenyan adolescents. *Journal of Child & Adolescent Trauma* 17(2), 261–273.
- Ferreira AL, Bezerra MA, de Lima LX, Jefferies P, Coimbra RM, da Cunha DP, Santos AS, da Silva MLF, Brasil TL and Ungar M (2022) Adaptation and psychometric evaluation of a Brazilian version of the CYRM-28. *Estudos de Psicologia (Natal)* 27(1), 12–22.
- Freney J, Peskin M, Schick V, Cuccaro P, Addy R, Morgan R, Lopez KK and Johnson-Baker K (2021) Adverse childhood experiences, depression, resilience, & spirituality in African-American adolescents. *Journal of Child & Adolescent Trauma* 14, 209–221.
- Fulu E, Miedema S, Roselli T, McCook S, Chan KL, Haardörfer R, Jewkes R, Warner X, Lang J and Naved RT (2017) Pathways between childhood trauma, intimate partner violence, and harsh parenting: Findings from the UN multi-country study on men and violence in Asia and the Pacific. *The Lancet Global Health* 5(5), e512–e522.
- Giovannelli A, Mondì CF, Reynolds AJ and Ou S-R (2020) Adverse childhood experiences: Mechanisms of risk and resilience in a longitudinal urban cohort. *Development and Psychopathology* 32(4), 1418–1439.
- Girma S, Tsehay M, Mamaru A and Abera M (2021) Depression and its determinants among adolescents in Jimma town, Southwest Ethiopia. *PLoS One* 16(5), e0250927.
- Glenn CR, Kleiman EM, Kandlur R, Esposito EC and Liu RT (2022) Thwarted belongingness mediates interpersonal stress and suicidal thoughts: An intensive longitudinal study with high-risk adolescents. *Journal of Clinical Child & Adolescent Psychology* 51(3), 295–311.
- Guo T, Huang L, Hall DL, Jiao C, Chen S-T, Yu Q, Yeung A, Chi X and Zou L (2021) The relationship between childhood adversities and complex post-traumatic stress symptoms: A multiple mediation model. *European Journal of Psychotraumatology* 12(1), 1936921.
- Halepota AA and Wasif SA (2001) Harvard trauma questionnaire Urdu translation: The only cross-culturally validated screening instrument for the assessment of trauma torture and their sequelae. *Journal-Pakistan Medical Association* 51(8), 285–289.
- Hamby S, Elm JHL, Howell KH and Merrick MT (2021) Recognizing the cumulative burden of childhood adversities transforms science and practice for trauma and resilience. *American Psychologist* 76(2), 230.
- Haroz EE, Murray LK, Bolton P, Betancourt T and Bass JK (2013) Adolescent resilience in northern Uganda: The role of social support and prosocial behavior in reducing mental health problems. *Journal of Research on Adolescence* 23(1), 138–148.
- Hayes AF (2018) Partial, conditional, and moderated mediation: Quantification, inference, and interpretation. *Communication Monographs* 85(1), 4–40.
- Houtepen LC, Heron J, Suderman MJ, Fraser A, Chittleborough CR and Howe LD (2020) Associations of adverse childhood experiences with educational attainment and adolescent health and the role of family and socioeconomic factors: A prospective cohort study in the UK. *PLoS Medicine* 17(3), e1003031. <https://doi.org/10.1371/journal.pmed.1003031>.
- Ipekci B (2021) *Posttraumatic Stress and Resilience among Iraqi and Syrian Refugees*. Boston, USA: University of Massachusetts Boston.
- Jefferies P, McGarrigle L and Ungar M (2019) The CYRM-R: A Rasch-validated revision of the child and youth resilience measure. *Journal of Evidence-Based Social Work* 16(1), 70–92.
- Jewkes RK, Dunkle K, Nduna M, Jama PN and Puren A (2010) Associations between childhood adversity and depression, substance abuse and HIV and HSV2 incident infections in rural south African youth. *Child Abuse & Neglect* 34(11), 833–841.
- John PB, Russell S and Russell PSS (2007) The prevalence of posttraumatic stress disorder among children and adolescents affected by tsunami disaster in Tamil Nadu. *Disaster Management & Response* 5(1), 3–7.
- Kaaria BK, Nyagah V and Ntarangwe M (2024) The prevalence of adverse childhood experiences among adolescents in secondary schools in Imenti south Sub-County in Meru County. *Journal of Sociology, Psychology & Religious Studies* 6(3), 27–36.
- Kelifa MO, Yang Y, Herbert C, He Q and Wang P (2020) Psychological resilience and current stressful events as potential mediators between adverse childhood experiences and depression among college students in Eritrea. *Child Abuse & Neglect* 106, 104480.
- Kroenke K and Spitzer RL (2002) The PHQ-9: A new depression diagnostic and severity measure. *Psychiatric Annals* 32(9), 509–515.
- Kroenke K, Spitzer RL and Williams JBW (2001) The PHQ-9: Validity of a brief depression severity measure. *Journal of General Internal Medicine* 16(9), 606–613.

- Lawley K** (2023) *Multiple Adverse Childhood Experiences and Maternal Mental Health Outcomes among Kenyan Mothers with and without HIV*. Washington, USA: University of Washington.
- Leone DW** (2024) The relationship between adverse childhood experiences and mental health issues among transgender individuals in Kenya. *Journal of Gay & Lesbian Mental Health* 28(4), 592–626.
- Lhewa D, Banu S, Rosenfeld B and Keller A** (2007) Validation of a Tibetan translation of the Hopkins symptom checklist–25 and the Harvard trauma questionnaire. *Assessment* 14(3), 223–230.
- Liebenberg I and Moore JC** (2018) A social ecological measure of resilience for adults: The RRC-ARM. *Social Indicators Research* 136, 1–19.
- Luthar SS and Zelazo LB** (2003) Research on resilience: An integrative review. *Resilience and Vulnerability: Adaptation in the Context of Childhood Adversities* 2, 510–549.
- Malhi GS, Das P, Bell E, Mattingly G and Mannie Z** (2019) Modelling resilience in adolescence and adversity: A novel framework to inform research and practice. *Translational Psychiatry* 9(1), 316.
- Margolin G and Vickerman KA** (2007) Posttraumatic stress in children and adolescents exposed to family violence: I. Overview and issues. *Professional Psychology: Research and Practice* 38(6), 613.
- McLorg A, Omolo K, Sifuna P, Shaw A, Walia B and Larsen DA** (2021) Examining wealth trends in Kombewa, Kenya. *Social Indicators Research* 157(2), 631–651.
- McMullen JD, O'Callaghan PS, Richards JA, Eakin JG and Rafferty H** (2012) Screening for traumatic exposure and psychological distress among war-affected adolescents in post-conflict northern Uganda. *Social Psychiatry and Psychiatric Epidemiology* 47, 1489–1498.
- Michael Garrett P** (2016) Questioning tales of 'ordinary magic': resilience and neo-liberal reasoning. *British Journal of Social Work* 46(7), 1909–1925.
- Miedema SS, Chiang L, Annor FB and Achia T** (2023) Cross-time comparison of adverse childhood experience patterns among Kenyan youth: Violence against children and youth surveys, 2010 and 2019. *Child Abuse & Neglect* 141, 106153.
- Mollica RF, Caspi-Yavin Y, Bollini P, Truong T, Tor S and Lavelle J** (1992) The Harvard trauma questionnaire: Validating a cross-cultural instrument for measuring torture, trauma, and posttraumatic stress disorder in Indochinese refugees. *The Journal of Nervous and Mental Disease* 180(2), 111–116.
- Morgan CA, Chang Y-H, Choy O, Tsai M-C and Hsieh S** (2021) Adverse childhood experiences are associated with reduced psychological resilience in youth: A systematic review and meta-analysis. *Children* 9(1), 27.
- Mutamba E** (2020) *Adverse Childhood Experiences and associated correlates among Adolescents at a Rural Private High School in Kenya*. Repository of Tangaza University.
- Mwiti GK** (2024) Counseling and psychotherapy—An African perspective. *Open Access Library Journal* 11(7), 1–16.
- Ndeti DM, Ongecha-Owuor FA, Khasakhala L, Mutiso V, Odhiambo G and Kokonya DA** (2007) Traumatic experiences of Kenyan secondary school students. *Journal of Child & Adolescent Mental Health* 19(2), 147–155.
- Okello J** (2014) *War Trauma, Attachment and Risky Behaviour in Adolescents in Northern Uganda*. Ghent, Belgium: Ghent University.
- Okwori G** (2022) Role of individual, family, and community resilience in moderating effects of adverse childhood experiences on mental health among children. *Journal of Developmental & Behavioral Pediatrics* 43(7), e452–e462.
- Oral R, Ramirez M, Coohy C, Nakada S, Walz A, Kuntz A, Benoit J and Peek-Asa C** (2016) Adverse childhood experiences and trauma informed care: The future of health care. *Pediatric Research* 79(1), 227–233.
- Oruc L, Kapetanovic A, Pojskic N, Miley K, Forstbauer S, Mollica RF and Henderson DC** (2008) Screening for PTSD and depression in Bosnia and Herzegovina: Validating the Harvard trauma questionnaire and the Hopkins symptom checklist. *International Journal of Culture and Mental Health* 1(2), 105–116.
- Panagou C and MacBeth A** (2022) Deconstructing pathways to resilience: A systematic review of associations between psychosocial mechanisms and trans-diagnostic adult mental health outcomes in the context of adverse childhood experiences. *Clinical Psychology & Psychotherapy* 29(5), 1626–1654.
- Panagou K** (2021) *Deconstructing pathways to resilience: a systematic review of associations between psychosocial mechanisms and adult mental health outcomes in the context of ACEs; and, trajectories of risk and resilience: the role of empathy and perceived social support*. Thesis from The University of Edinburgh.
- Poole JC, Dobson KS and Pusch D** (2017) Childhood adversity and adult depression: The protective role of psychological resilience. *Child Abuse & Neglect* 64, 89–100.
- Posner K, Brown GK, Stanley B, Brent DA, Yershova KV, Oquendo MA, Currier GW, Melvin GA, Greenhill L and Shen S** (2011) The Columbia–suicide severity rating scale: Initial validity and internal consistency findings from three multisite studies with adolescents and adults. *American Journal of Psychiatry* 168(12), 1266–1277.
- Reich JW, Zautra AJ and Hall JS** (2010) *Handbook of Adult Resilience*. New York: Guilford Press.
- Resilience Research Centre** (2022) *CYRM and ARM User Manual v2.5*. Halifax: Resilience Research Centre. Dalhousie University.
- Rutstein SO** (2015) *Steps to Constructing the New DHS Wealth Index*. Rockville, MD: ICF International, pp. 6.
- Rutter M** (2013) Annual research review: Resilience – clinical implications. *Journal of Child Psychology and Psychiatry* 54(4), 474–487. <https://doi.org/10.1111/j.1469-7610.2012.02615.x>.
- Ryttilä-Manninen M, Haravuori H, Fröjd S, Marttunen M and Lindberg N** (2018) Mediators between adverse childhood experiences and suicidality. *Child Abuse & Neglect* 77, 99–109.
- Salokangas RKR, Schultze-Lutter F, Patterson P, von Reventlow HG, Heinimaa M, From T, Luutonen S, Hanka J, Kotimäki M and Tuominen L** (2016) Psychometric properties of the trauma and distress scale, TADS, in an adult community sample in Finland. *European Journal of Psychotraumatology* 7(1). <https://doi.org/10.3402/ejpt.v7.30062>.
- Salokangas RKR, Schultze-Lutter F, Schmidt SJ, Pesonen H, Luutonen S, Patterson P, von Reventlow HG, Heinimaa M, From T and Hietala J** (2020) Childhood physical abuse and emotional neglect are specifically associated with adult mental disorders. *Journal of Mental Health*, 29, 376–384.
- Sasidharan D and Talwar P** (2023) The influence of adverse childhood experiences on mental health: A review study. *International Journal of Health Sciences and Pharmacy (IJHSP)* 7(2), 156–179.
- Satinsky EN, Kakuhiire B, Baguma C, Rasmussen JD, Ashaba S, Cooper-Vince CE, Perkins JM, Kiconco A, Namara EB and Bangsberg DR** (2021) Adverse childhood experiences, adult depression, and suicidal ideation in rural Uganda: A cross-sectional, population-based study. *PLoS Medicine* 18(5), e1003642.
- Schulz A, Becker M, Van der Auwera S, Barnow S, Appel K, Mahler J, Schmidt CO, John U, Freyberger HJ and Grabe HJ** (2014) The impact of childhood trauma on depression: Does resilience matter? Population-based results from the study of health in Pomerania. *Journal of Psychosomatic Research* 77(2), 97–103.
- Schwartzman JM, Muscatello RA and Corbett BA** (2023) Assessing suicidal thoughts and behaviors and nonsuicidal self-injury in autistic and non-autistic early adolescents using the Columbia suicide severity rating scale. *Autism* 27(8), 2310–2323.
- Seery MD, Holman EA and Silver RC** (2010) Whatever does not kill us: Cumulative lifetime adversity, vulnerability, and resilience. *Journal of Personality and Social Psychology* 99(6), 1025.
- Seya M-KS, Matthews S, Zhu L, Brown C, Lefevre A, Agathis N, Chiang IF, Annor FB, McOwen J and Augusto A** (2024) Parenting-related positive childhood experiences, adverse childhood experiences, and mental health—Four sub-Saharan African countries. *Child Abuse & Neglect* 150, 106493.
- Sher L** (2019) Resilience as a focus of suicide research and prevention. *Acta Psychiatrica Scandinavica* 140(2), 169–180.
- Spinazzola J, Hodgdon H, Liang L-J, Ford JD, Layne CM, Pynoos R, Briggs EC, Stolbach B and Kisiel C** (2014) Unseen wounds: The contribution of psychological maltreatment to child and adolescent mental health and risk outcomes. *Psychological Trauma: Theory, Research, Practice, and Policy* 6(S1), S18.
- Srivastav A, Stropolis M, Moseley A and Daniels K** (2020) The empower action model: A framework for preventing adverse childhood experiences by promoting health, equity, and well-being across the life span. *Health Promotion Practice* 21(4), 525–534.
- Stainton A, Chisholm K, Kaiser N, Rosen M, Upthegrove R, Ruhrmann S and Wood SJ** (2019) Resilience as a multimodal dynamic process. *Early Intervention in Psychiatry* 13(4), 725–732.

- Thompson MP, Kingree JB and Lamis D** (2019) Associations of adverse childhood experiences and suicidal behaviors in adulthood in a US nationally representative sample. *Child: Care, Health and Development* 45(1), 121–128.
- Turner HA, Finkelhor D and Ormrod R** (2006) The effect of lifetime victimization on the mental health of children and adolescents. *Social Science & Medicine* 62(1), 13–27.
- Ungar M, Liebenberg L, Boothroyd R, Kwong WM, Lee TY, Leblanc J, Duque L and Makhnatch A** (2008) The study of youth resilience across cultures: Lessons from a pilot study of measurement development. *Research in Human Development* 5(3), 166–180.
- Ungar M and Theron L** (2020) Resilience and mental health: How multi-systemic processes contribute to positive outcomes. *The Lancet Psychiatry* 7(5), 441–448.
- Vundi C** (2022) *Prevalence and Patterns of Mental Health Disorders among Children and Adolescents with Physical Illness at the Outpatient Department of Kenyatta National Hospital*. Kenya: University of Nairobi.
- Wang W, Fu W, Wu J, Ma X, Sun X, Huang Y, Hashimoto K and Gao C** (2012) Prevalence of PTSD and depression among junior middle school students in a rural town far from the epicenter of the Wenchuan earthquake in China. *PLoS One* 7(7), e41665.
- White BM, Prasad R, Ammar N, Yaun JA and Shaban-Nejad A** (2024) Digital health innovations for screening and mitigating mental health impacts of adverse childhood experiences: Narrative review. *JMIR Pediatrics and Parenting* 7(1), e58403.
- Wolff N and Caravaca Sánchez F** (2019) Associations among psychological distress, adverse childhood experiences, social support, and resilience in incarcerated men. *Criminal Justice and Behavior* 46(11), 1630–1649.
- Wu H and Leung S-O** (2017) Can Likert scales be treated as interval scales? – A simulation study. *Journal of Social Service Research* 43(4), 527–532.
- Yershova K, Lesser A, Logan K and Posner K** (2016) Asking about suicide as suicide prevention: The Columbia suicide severity rating scale (C-SSRS). In *Understanding Suicide: From Diagnosis to Personalized Treatment*. Berlin Germany: Springer Nature. pp. 29–41.
- Youssef NA, Green KT, Dedert EA, Hertzberg JS, Calhoun PS, Dennis MF, Research Education and Clinical Center Workgroup M-AMI and Beckham JC** (2013) Exploration of the influence of childhood trauma, combat exposure, and the resilience construct on depression and suicidal ideation among US Iraq/Afghanistan era military personnel and veterans. *Archives of Suicide Research* 17(2), 106–122.
- Yu Z, Wang L, Chen W, Perrin N and Gross D** (2021) Childhood adversity and mental health among Chinese young adults: The protective role of resilience. *Journal of Advanced Nursing* 77(12), 4793–4804.
- Zahradnik M, Stewart SH, O'Connor RM, Stevens D, Ungar M and Wekerle C** (2010) Resilience moderates the relationship between exposure to violence and posttraumatic reexperiencing in Mi'kmaq youth. *International Journal of Mental Health and Addiction* 8, 408–420.
- Zakhour M, Haddad C, Sacre H, Fares K, Akel M, Obeid S, Salameh P and Hallit S** (2021) Suicidal ideation among Lebanese adults: Scale validation and correlates. *BMC Psychiatry* 21, 1–11.
- Zhang Y, Li Y, Jiang T and Zhang Q** (2023) Role of body mass index in the relationship between adverse childhood experiences, resilience, and mental health: A multivariate analysis. *BMC Psychiatry* 23(1), 460.
- Zrihan-Weitzman A, Merhav I, Farchi M and Peled-Avram M** (2023) Resilience as Mediator Between Two Restriction Stages of COVID-19 and Mental Distress. *The International Journal of Community and Social Development*, 25166026231207984.