













Pollen sterols are associated with phylogeny and environment but not with pollinator guilds

Pengjuan Zu^{1,2} , Hauke Koch¹ , Orlando Schwery³ , Samuel Pironon⁴ , Charlotte Phillips^{4,5} , Ian Ondo⁴ , Iain W. Farrell¹ , W. David Nes⁶ , Elynor Moore⁷ , Geraldine A. Wright⁷ , Dudley I. Farman⁸  and Philip C. Stevenson^{1,8} 

¹Royal Botanic Gardens, Kew, Natural Capital and Plant Health Department, Richmond, Surrey, TW9 3AB, UK; ²Department Fish Ecology and Evolution, Swiss Federal Institute of Aquatic Science and Technology, Seestrasse 79, Kastanienbaum CH-6047, Switzerland; ³New Mexico Consortium, 4200 W. Jemez Rd, Suite 301, Los Alamos, NM 87544, USA; ⁴Royal Botanic Gardens, Kew, Biodiversity Informatics and Spatial Analysis Department, Richmond, Surrey, TW9 3AB, UK; ⁵Royal Botanic Gardens, Kew, Conservation Science Department, Wakehurst Place, Ardingly, West Sussex, RH17 6TN, UK; ⁶Department of Chemistry & Biochemistry, Texas Tech University, Lubbock, TX 79424, USA; ⁷Department of Zoology, University of Oxford, 11a Mansfield Road, Oxford, OX1 3SZ, UK; ⁸Natural Resources Institute, University of Greenwich, Chatham, Kent, ME4 4TB, UK

Summary

Author for correspondence:
Pengjuan Zu
Email: Pengjuan.zu@gmail.com

Received: 16 October 2020
Accepted: 14 January 2021

New Phytologist (2021) 230: 1169–1184
doi: 10.1111/nph.17227

Key words: chemical ecology, chemotaxonomy, environmental factors, phytosterol diversity, phylogeny, plant–insect interactions, pollen nutrients, pollinator guild.

- Phytosterols are primary plant metabolites that have fundamental structural and regulatory functions. They are also essential nutrients for phytophagous insects, including pollinators, that cannot synthesize sterols. Despite the well-described composition and diversity in vegetative plant tissues, few studies have examined phytosterol diversity in pollen.
- We quantified 25 pollen phytosterols in 122 plant species (105 genera, 51 families) to determine their composition and diversity across plant taxa. We searched literature and databases for plant phylogeny, environmental conditions, and pollinator guilds of the species to examine the relationships with pollen sterols.
- 24-methylencholesterol, sitosterol and isofucosterol were the most common and abundant pollen sterols. We found phylogenetic clustering of twelve individual sterols, total sterol content and sterol diversity, and of sterol groupings that reflect their underlying biosynthesis pathway (C-24 alkylation, ring B desaturation). Plants originating in tropical-like climates (higher mean annual temperature, lower temperature seasonality, higher precipitation in wettest quarter) were more likely to record higher pollen sterol content. However, pollen sterol composition and content showed no clear relationship with pollinator guilds.
- Our study is the first to show that pollen sterol diversity is phylogenetically clustered and that pollen sterol content may adapt to environmental conditions.

Introduction

Phytosterols are a class of lipids with key metabolic and ecological functions for plants (Nes & McKean, 1977; Vanderplanck *et al.*, 2020a). For example, they regulate membrane fluidity and permeability (Grunwald, 1971; Schuler *et al.*, 1991; Hartmann, 1998), act as precursors for metabolic signals such as brassinosteroid growth hormones (Grove *et al.*, 1979; Chung *et al.*, 2010) that promote cell division, mediate reproduction in plants, and protect them against environmental stresses (Khripach *et al.*, 2000). Phytosterols may also modulate plant defence against bacterial pathogens (Posé *et al.*, 2009; Wang *et al.*, 2012; Ferrer *et al.*, 2017), and pollen sterols accelerate germination and tube growth and protect against desiccation (Kumar *et al.*, 2015; Rotsch *et al.*, 2017).

Phytosterols show considerable diversity with > 250 structures reported (Nes, 2011) with notable variation at the methine substitution (double bond) in ring B and methyl or ethyl

substitutions at C-24 (Fig. 1). The structural variation and composition of sterols in plant tissues is important for phytophagous insects since they cannot synthesize sterols *de novo*, and therefore depend upon specific plants to obtain the required sterols from their diet to sustain their development (Behmer & Elias, 1999, 2000; Lang *et al.*, 2012). This may be especially important for pollen feeding insects that require specific sterols. Honeybees, for example, require 24-methylencholesterol (Herbert *et al.*, 1980; Chakrabati *et al.*, 2020), so they must collect pollen from plant species that produce this sterol to rear brood. Bee sterols are similar to those occurring in the pollen on which they feed (Vanderplanck *et al.*, 2020a) but differ across bee taxa, suggesting bees are what they eat with respect to sterols. Wild pollinators range from pollen generalists to specialists (Rasmussen *et al.*, 2020), and for some species this specialism may be mediated by pollen sterols. Therefore, a landscape of flowers that does not provide the sterols required for a specific bee may be nutritionally deficient for that species. In general, however, the relationships

(a) 24-Methylenecholesterol



(b) Double bond (Δ or D) position in ring B (c) Substitution at C-24

Fig. 1 Chemical structure of 24-methylenecholesterol as an illustration of phytosterols, showing (a) carbon numbering (b) different substitutions in ring B, and (c) different substitutions at C-24.

between pollen sterols and the nutritional needs of pollinating insects have not yet been evaluated.

Conversely, plant sterol composition may evolve with antagonists as well as mutualists since the pathways for the synthesis of sterols overlap with those for some defence compounds against herbivores (Qi *et al.*, 2006). A range of naturally occurring insect toxins occur in pollen (Arnold *et al.*, 2014; Rivest & Forrest, 2020), with the likely role of protecting the male gamete – some sterols can also act as defensive compounds against arthropod herbivores (Jing & Behmer, 2020). They could also be toxic to pollen feeding insects to reduce damage to or excessive loss of pollen grains.

Abiotic conditions may affect phytosterol structural variations at different levels. At individual plant level, an ethyl substitution at C-24 (e.g. sitosterol and stigmasterol), for example, reinforces membrane cohesion (Piironen *et al.*, 2000; Dufourc, 2008), and therefore sterol structures may be altered in response to temperature variations. At population level, from limited heritability studies on phytosterols in plant seeds (Amar *et al.*, 2008; Velasco *et al.*, 2013), environmental factors have also been shown to contribute to sterol phenotypic variation, although much less compared to the contribution from genetic factors (heritabilities > 0.8 have been documented). At species level, pollen sterol composition seems to be highly variable between different species (Villette *et al.*, 2015; Vanderplanck *et al.*, 2020a) and can differ from the sterol composition in vegetative tissues (Nes, 1990; Nes *et al.*, 1993). However, no study has investigated whether pollen sterol variations at species level can be a consequence of evolutionary adaptation to environmental conditions.

Moreover, due to the limited number of studies on pollen sterol profiles, we lack a comprehensive and fundamental

understanding of the patterns of pollen sterol diversity across plant taxa. The question of whether pollen phytosterols are phylogenetically structured remains controversial. For example, Standifer *et al.* (1968) suggested a lack of phylogenetic constraints on pollen sterol composition, based on the evidence of large variation in three Salicaceae species. By contrast, Vanderplanck *et al.* (2020a) found similar pollen sterol composition within the genus *Salix*, and our interpretation of the data published by Villette *et al.* (2015) suggested that the occurrence of some pollen sterols was phylogenetically constrained. Since most studies focused on a few plant species, they were insufficient to reach a general overview of the patterns and drivers of pollen sterol diversity across plant taxa.

In this study we analysed pollen sterols, including saturated stanols, in 122 angiosperms representing 51 plant families and 25 plant orders. We further compiled data from the literature and from databases on plant phylogeny, pollinators, and environmental conditions within native geographic regions for each plant species to examine relationships between these factors and pollen sterol composition and diversity. Specifically, we ask the following questions: First, are pollen sterols phylogenetically clustered? Second, are pollen sterols correlated with abiotic environments? And third, are pollen sterols associated with pollinator guilds?

Materials and Methods

Pollen collection

From March to November 2018, we collected pollen from fresh flowers growing in the Royal Botanic Gardens (RBG), Kew, UK and nearby areas (see Supporting Information Table S1 for

details of collection dates and locations for each species). RBG Kew supports a diverse collection of living plant species from across the world. Before pollen collection, we used a fine-meshed bag to cover flower buds whenever possible to prevent potential contamination or removal due to pollinator visitation. When flowers were fully open, we gently shook the flower and collected pollen into a weighed 2 ml microcentrifuge tube (Safe-Lock, Eppendorf, Stevenage, UK). For species for which pollen was more difficult to harvest, such as in the cases of *Lamium purpureum* L. and *Ulex europaeus* L., we used small forceps to help push the pollen out or trigger pollen ejection, respectively. Pilot studies carried out in our laboratory (with *Helleborus foetidus* L., *Prunus avium* L., *Prunus spinosa* L., *Salix cinerea* L. and *Symphytum officinale* L.) showed a conserved pattern of pollen sterol composition: within-species variation was significantly lower than between-species variation (all P -values < 0.001 under multi-variate distribution tests, e.g. Hotelling test, Pillai test, and Wilks' lambda distribution test), consistent with findings on within- vs between-species variation of other pollen metabolites (Palmer-Young *et al.*, 2019). Therefore, we collected 2 to 5 replicates per species (for details see Table S1) and used the average quantities across replicates of each species for analysis. In total, we collected 308 samples from 122 species, representing 105 genera, 51 families and 25 orders across the major groups of seed plants (Gymnosperms, Nymphaeales, Monocots, Ranunculids, Caryophyllales, Asterids and Rosids; Table S1). Our selection of species was guided by a combination of practical considerations (feasibility of collecting sufficient pollen for analysis, availability of species at Kew) while attempting to maximize phylogenetic and ecological diversity of plants (pollinator guilds, ecological niches). Pollen weight (to 0.1 mg accuracy) and collection date were recorded for each sample. Pollen samples were stored in a freezer (-20°C) before extracting sterols.

Sterol content analysis

To extract sterols and stanols (from here referred to as phytosterols or pollen sterols) from the pollen, we added 500 μl 10% KOH in MeOH to the microcentrifuge tubes containing a weighed pollen sample. Then, an internal standard (20 μl of 0.5 mg ml^{-1} epicoprostanol) was added before incubating the tube for 2 h at 80°C for saponification. Phytosterols were then recovered into 1 ml hexane. After complete evaporation of the hexane, the phytosterols remained in the tube. We derivatized these with 20 μl Tri-Sil (Sigma) and then briefly vortexed them and injected them directly into an Agilent Technologies (Palo Alto, CA, USA) 7890A gas chromatograph connected to an Agilent Technologies 5975C MSD mass spectrometer (for gas chromatography–mass spectrometry (GC–MS)) and eluted over an Agilent DB5 column using a splitless injection at 250°C with a standard GC program at 170°C for 1 min, ramped to 280°C at $20^{\circ}\text{C min}^{-1}$ and monitoring between 50 and 550 amu.

All 25 phytosterols were identified by comparison of their retention time relative to cholesterol and mass spectra from authentic standards (W. David Nes collection; see Fig. S1 for mass spectra of each sterol) either directly through co-analysis or

using existing data and confirmed where data was available with the NIST (National Institute of Standards and Technology) mass spectral library (Guo *et al.*, 1995; Heupel & Nes, 1984; Nes *et al.*, 1977, 2003, 2009; Xu *et al.*, 1988).

To quantify the amount of each phytosterol, we used its relative peak area by calculating the ratio of the peak area of the targeted sterol to that of the internal standard. Then, by multiplying the ratio with the mass of the internal standard, we obtained the mass of each sterol in the sample. Compound identification (using target ions) and quantification were carried out with CHEMSTATION ENHANCED DATA ANALYSIS (v.E.01.00). In total, we identified 25 phytosterols in pollen (Table S1).

For each plant species, we calculated phytosterol amount ($\mu\text{g mg}^{-1}$ sampled pollen), total sterol content ($\mu\text{g mg}^{-1}$ sampled pollen), and the percentage of each sterol in the total phytosterol content. In addition, we calculated the chemical diversity index using Shannon entropy: where S is the total number of phytosterols, p_i is the percentage of the i^{th} phytosterol. Note that we used the total phytosterol number S as the base of log (instead of the natural base e) to scale the range of diversity index values to $[0, 1]$, with 1 indicating the highest diversity. This equates to calculating Shannon's equitability. Finally, for each phytosterol, we calculated its commonness and abundance across all plant species. Commonness is given by the proportion of plant species that contained that specific phytosterol (i.e. present/absent). Relative abundance was given by the average proportion of a specific sterol across all species.

Additionally, to understand how different phytosterols in pollen co-varied, we performed a factor analysis using the R package STATS (R Core Team, 2020) on the data for the relative weight of phytosterols measured in pollen across the entire data set. We set a criterion of eigenvalue > 1 for inclusion of extracted factors. A varimax rotation was used to adjust the fit of the factor analysis to variance in the data.

Moreover, based on biosynthetic reasoning as discussed by Benveniste (2004), we arranged the phytosterols identified in our pollen samples into alternate hypothetical biosynthetic pathways to cholesterol and 24-alkyl phytosterols.

Phylogenetic tree construction and analyses

We used the R package ROTL (Michonneau *et al.*, 2016) to download the induced subtree of only our focal taxa from the Open Tree of Life (OTL) synthetic tree (Hinchliff *et al.*, 2015; Rees & Cranston, 2017). If only the genus was known, OTL used the root of the genus for the subtree wherever possible. Name synonyms and corrections suggested by OTL for genus and species were adopted in our analyses (see Table S2). Taxa with subspecies or other epithets beyond species level were reduced to genus and species only (*Amaryllis belladonna* L., *Campanula fragilis* Cirillo, *Campanula isophylla* Moretti, *Euphorbia milii* Des Moul., *Hieracium umbellatum* L.). Only one terminal was retained to represent the two differently coloured varieties of *Hymenocallis littoralis* (Jacq.) Salisb.

We estimated divergence times with penalised likelihood using nine secondary calibration points. Using the R package APE

(Paradis *et al.*, 2004), we randomly resolved polytomies and computed branch lengths using Grafen's method. We looked up the inferred ages of seven clades from the large phylogeny of spermatophytes documented by Zanne *et al.* (2014): Angiospermae (243 million years ago (Ma)), Monocotyledoneae (171 Ma), Eudicotyledoneae (137 Ma), Superrosidae (118 Ma), Rosidae (117 Ma), Superasteridae (117 Ma), and Asteridae (108 Ma). The age of Nymphaea (78.07 Ma) was obtained from DateLife (Sanchez-Reyes & O'Meara, 2019), and we took the estimated origin of Spermatophyta to be 327 Ma (Smith *et al.*, 2010) to calibrate the root age. We used those times as minimal age constraints for a penalized likelihood analysis using *chronopl* in APE (Paradis *et al.*, 2004). Monophyly of families was checked using MONOPHY (Schwery & O'Meara, 2016).

To determine whether there is phylogenetic structure in the pollen sterol data, we used the function *phyloSignal* from the R package PHYLOSIGNAL (Keck *et al.*, 2016) to calculate Pagel's λ (Pagel, 1999) and Blomberg's K (Blomberg *et al.*, 2003), each with 999 iterations for *P*-value estimation. Phylogenetic signal was estimated this way for each of the individual sterol compounds (based on their percentage value), for sums of compounds belonging to each C-24 substitution (C0, C1, C2 indicating substitution with no carbon, a methyl and an ethyl), position of the olefinic or methine moiety in ring B (Δ^0 , Δ^5 , Δ^7 , Δ^8), sterol diversity index H, and total phytosterol content (absolute sterol amount mg^{-1} pollen). The output of these analyses was visualized using the R packages PHYTOOLS (Revell, 2012) and PICANTE (Kembel *et al.*, 2010).

Plant occurrence records and abiotic environmental data

To investigate whether species-level variations in pollen sterols are partially consequences of evolutionary adaptation to environmental conditions, we retrieved environmental information regarding the native geographic ranges of each species. Note that here we focused on 'long-term' prevailing abiotic conditions (e.g. climate) capable of shaping evolutionary changes of sterol composition at species level, as opposed to 'short-term' abiotic variables (e.g. stresses, weather) affecting traits via phenotypic plasticity at the individual level. For each species, we extracted geographic occurrence records from several global and continental databases: GBIF (Global Biodiversity Information Facility; <https://www.gbif.org/>) using the RGBIF package in R, BIEN (Botanical Information and Ecology Network; <http://bien.nceas.ucsb.edu/bien/>) using the BIEN R package, BioTIME (Dornelas *et al.*, 2018) and Rainbio (Dauby *et al.*, 2016). Because raw occurrence data from these databases contain taxonomic, spatial and temporal inconsistencies (Meyer *et al.*, 2016), we applied different cleaning filters using the COORDINATECLEANER package in R (Zizka *et al.*, 2019). We discarded nongeoreferenced records, records with latitude and longitude given as zero and having equal longitude and latitude, and points recorded before 1950, as well as fossil data, records corresponding to centroids of countries, capitals, known botanical institutions and GBIF headquarters, occurrences falling in the sea, cultivated records, and points indicated as having high coordinate uncertainty (> 20 km). We used the World Geographical Scheme for

Recording Plant Distribution (WGSRPD) database (Brummitt, 2001) from the World Checklist of Vascular Plants (WCVP, 2020) to discard records from species reported outside of their native regions at level 2 (regional or subcontinental level). Finally, we removed duplicates and thinned each species' occurrence dataset by keeping only one record per 10×10 arcmin grid cell to limit spatial autocorrelation. In total, 355 912 occurrence records were retrieved across all species (Table S1).

We quantified species' environmental niches based on a set of 13 climate, soil, and topography variables. Eight of them were bioclimatic variables (BIO1, BIO4, BIO10, BIO11, BIO12, BIO15, BIO16 and BIO17) extracted from the CHELSA database (Karger *et al.*, 2017), representing annual mean, seasonality, minimum and maximum temperature and precipitation (for a full list of variables and descriptions see Table S3). Four soil variables were extracted from the SoilGrids database (ISRIC, 2013; <https://data.isric.org>) and averaged across a 0–60 cm depth gradient: depth to bedrock, mean soil organic carbon stock, pH, and water capacity. Land slope was calculated using the *Slope* function in the *Spatial Analyst* toolbox of ARC/INFO/ARCGIS based on the Global Multi-resolution Terrain Elevation Database (GMTED) (Danielson & Gesch, 2011). To match the resolution of the occurrence records, all environmental variables were upsampled to 10 arcmin (*c.* 20 km) using the *resample* function of the RASTER package in R.

We extracted each of the 13 environmental variables at each occurrence point of each species using the *extract* function of the RASTER package in R. Mean environmental conditions were then calculated for each of the 13 variables across all occurrences of each species (i.e. environmental niche position along individual environmental gradients). We also created an environmental space summarizing the variation in the 13 environmental variables across the world using a principal component analysis (PCA) and the function *princomp* in the STATS package in R. We kept the first three component axes that explained 74% of the variation in the 13 variables: PC1 was mainly positively correlated with mean temperatures and negatively correlated with temperature seasonality and soil carbon content, PC2 was positively correlated with soil pH and negatively correlated with precipitation, and PC3 was positively correlated with soil depth to bedrock and negatively correlated with land slope (see Table S3 for variable contributions to PCA axes). To quantify the niche breadth of each species, we first drew three-dimensional alpha shapes around each set of occurrence points for each species in the environmental space defined by the PCA with an alpha value of two using the *ashape3d* function in the ALPHASHAPE3D package in R (Capinha & Pateiro-López, 2014). The alpha-shape is a profile technique used to compute species environmental niche envelopes using a flexible envelope fitting procedure that makes no assumption about the shape of the niche (Capinha & Pateiro-López, 2014). We then calculated the volume of each species' alpha shape as a measure of their environmental niche breadth using the *volume_ashape3d* function from the latter package. We also calculated the mean position of each species' alpha shape on the three retained main axes of the PCA (i.e. niche position; for individual variable contributions see Table S3). Because three-

dimensional alpha shapes require at least five occurrence points to be drawn, species with fewer records were discarded. We also discarded those species lacking sufficient and reliable geographic data or those with taxonomic uncertainty (i.e. we did not extract occurrence records for genera, subspecies and hybrids). In the end, we quantified niche breadth for 90 species, while 32 taxa were discarded, and niche position for 100 species, with 22 taxa discarded (Table S1).

Pollinator data collection

To study whether there is a relationship between plants' pollen sterols and their pollinators, we categorized plants in two different ways. Firstly, they were categorized based on pollinator guilds, as (1) bee, (2) fly, (3) Lepidoptera, (4) thrips, (5) generalist insect, (6) bird, or (7) wind pollinated. Secondly, we grouped plants by whether or not pollen acts as a reward for bee pollinators. On the one hand, bees depend on pollen as larval food and require pollen sterols as essential nutrients. Plants could therefore hypothetically attract bee pollinators with pollen sterol profiles of high nutritional quality. On the other hand, if pollen does not play a role as a bee reward (i.e. in non-bee-pollinated plants, and/or where nectar is the sole reward), sterol profiles that are of low quality or even toxic to bees could be expected, to prevent pollen robbery (as shown for some other chemical compounds in pollen, Rivest & Forrest, 2020).

To classify plant pollinator guilds and groups, we conducted literature searches for each plant species on Google Scholar, using the scientific name (including common synonyms) and “pollinat*”, OR “pollen”, OR “flower” as search terms. We examined relevant cited or citing references of publications found in this way for additional records, and consulted Knuth (1908, 1909) and Westrich (2018), or personal observations. If no sources on pollination and flower visitation were available, the pollinator guild was classified as ‘unknown’ (10 species in data set). We included plant species in the ‘pollen as bee reward’ group that both receive pollination services by bees (including some plants in the ‘generalist insect pollination’ category) and have records of bees collecting pollen. Plants were classified as ‘not producing pollen as bee reward’ if they were either not pollinated by bees, or, in case of bee pollination, had clear evidence of pollen not being collected by bees (e.g. pollen contained in pollinia of bee-pollinated orchids). Plants for which data on pollinator guild and collection of pollen by bees was missing were classified as ‘unknown’ (34 species in data set). A full list of relevant references and the assigned pollinator guilds is provided in Table S1.

Analyses of relationships between phytosterols and (a)biotic factors

To assess the association of sterol composition with environmental variables and pollinator guilds, we first calculated a Bray–Curtis distance matrix for sterol profiles of pairwise plant species comparisons, using absolute weights (μg) of each sterol mg^{-1} pollen. We then related this distance matrix to environmental factors and pollinator guild to study their relationships. Specifically, for abiotic environmental factors (continuous values), we

ran MRM (multiple regression on distance matrices) analyses (Lichstein, 2007) using an additive linear model with the Bray–Curtis distance matrix of pollen sterol composition dissimilarity as response, and environmental niche distance matrices for PC1, PC2, PC3 (calculated from pairwise Euclidean distances for all plants for their position on each of the PCs; see ‘Plant occurrence records and abiotic environmental data’ in the Materials and Methods section, above), and a phylogenetic distance matrix (pairwise phylogenetic distance in Ma; for more details on phylogeny see above) as independent variables. We used Pearson correlations with 10 000 permutations to test for significant associations. Calculations of distance matrices and MRM analyses were performed using the R package ECODIST (Goslee & Urban, 2007). For pollinator modes (categorical variables), we conducted analyses of similarities (ANOSIMs; Clarke, 1993) to test for significant differences of pollen sterol profiles between different pollinator groups (excluding pollinator groups with only one representative, i.e. wind, thrips, and fly) or between plant groups where pollen is used as a reward by bees or not. ANOSIMs were conducted in PAST v.4.03 (Hammer *et al.*, 2001) with 10 000 permutations. We illustrated the relationship of these factors to sterol profile similarity with 2D nonmetric multidimensional scaling (NMDS) ordination plots in PAST v.4.03 based on Bray–Curtis dissimilarities.

We furthermore examined associations of environmental variables and niche breadth with total sterol content and diversity. We calculated phylogenetic independent contrasts (Felsenstein, 1985; implemented in R package APE (Paradis *et al.*, 2004)) with the constructed plant phylogeny (see ‘Phylogenetic tree construction and analyses’ section in Materials and Methods section) for sterol content, Shannon diversity index H , positions on environmental principal component axes (PC1, PC2, PC3), and environmental niche breadth. Associations between contrasts of sterol content or diversity (as dependent variable) with contrasts of environmental principal components, niche breadth, or the 13 separate environmental factors were then individually evaluated by linear models in R (fitting the regression through the origin).

Finally, as 24-methylenecholesterol is a key sterol nutrient for honey bee development (Herbert *et al.*, 1980; Svoboda *et al.*, 1980) and could therefore have been selected for as an attracting reward in bee pollinated plants, we tested for differences in 24-methylenecholesterol content for plants that offer pollen as a reward for bees or not (or for which this interaction was unknown) with a phylogenetic ANOVA (Garland *et al.*, 1993), implemented in the R package PHYTOOLS (Revell, 2012), with 1000 simulations, and ‘Holm’ *post-hoc* testing. The same test was also conducted for total sterol content. Only species with phylogenetic information were included (pollen as bee reward: $n = 54$; pollen not bee reward: $n = 22$; unknown: $n = 24$).

Results

Pollen sterol composition and diversity across taxa

We profiled 25 phytosterols in the pollen of 122 plant species from 51 families, including representatives of Gymnosperms,

Nymphaeales, Monocots, Ranunculids, Caryophyllales, Asterids, and Rosids (Fig. 2; Table S1). These phytosterols can be arranged into biosynthetic pathways with three main distinct branches (i.e. C-24 – no substitution, C-24 methyl and C-24 ethyl groups, Fig 3; see Fig. 1 for illustrations of the group structure). Pollen phytosterols varied qualitatively and quantitatively across taxa, with each species exhibiting a distinctive sterol profile (Fig. 2).

Across all the sampled species, the most common pollen sterols (see ‘commonness’ at the bottom of Fig. 2) were sitosterol (recorded in 97.5% of sampled species), campesterol (88.5%), isofucosterol (82.0%), cholesterol (82.0%), cycloartenol (81.1%), 24-methylenecholesterol (ostreasterol, 73.0%), and stigmasterol (59.0%). The most abundant sterol dominating pollen sterol profiles (see ‘abundance’ at the bottom of Fig. 2) was 24-methylenecholesterol (on average accounting for 23% of total sterol content), followed by isofucosterol (21.5%), sitosterol (20.7%), and cycloartenol (17.7%). The first three are all Δ^5 sterols, of which 24-methylenecholesterol belongs to the 24C-methyl group, whereas sitosterol and isofucosterol belong to the C-24 ethyl group. Cholesterol, the primary sterol in animals, only represented a small portion (< 1%) of pollen sterol content, despite being common.

The pollen sterol diversity of plants varied dramatically, with a mean of 9.98 ± 4.46 (mean \pm SD) different phytosterols. For example, the carnivorous plant *Drosera regia* Stephens had almost exclusively 24-methylenecholesterol in pollen, whereas pollen from ivy (*Hedera helix* L.) contained 23 different sterols, tea pollen (*Camellia sinensis* L.) had 22 sterols, and pollen from the bellflowers *C. fragilis* Cirillo and *C. isophylla* Moretti had 23 and 19 sterols respectively. However, in all these species, only one to two sterol compounds were typically major components (contributing > 50% of total sterol content). The variation in the total weights of sterols led to a Shannon diversity index for pollen sterol composition ranging from 0 in *D. regia* to 0.64 in *H. helix* (Fig. 2; Table S1), with a mean of 0.34 (note that we standardized the maximum value of the Shannon diversity index to be 1.0, see the Materials and Methods section).

Covariance of pollen sterols

The factor analysis reduced the data to 12 independent latent factors that explained 73% of sterol variation (Table 1). Overall, phytosterols that have close positions in their biosynthetic

pathways (Fig. 3) or use the same enzyme (e.g. reductase) for production tend to align together with the same factors. For example, iso-obtusifolol is the precursor of 24-methylenelophenol, which then branches to either epifungisterol or avenasterol via episterol (Fig. 3). These four sterol compounds (not including episterol) largely aligned together with factor 1, which accounted for c. 9% of the variance (Table 1). Similar patterns also applied to factor 3 and factor 4, whose main contributing sterols represented the early cyclopropyl pathway intermediates. Factor 5 represented a strong positive correlation among the stanols (saturated in ring B) campestanol and sitostanol. Factors 6 and 7 represent products of Δ^{24} reduction. In addition, we found one inverse relationship between four of the most common phytosterols (in factor 2, accounting for 8% of the variance), where 24-methylenecholesterol was aligned in the opposite direction to the presence of three other phytosterols: sitosterol, campesterol, and stigmasterol.

Phylogenetic patterns

We found significant phylogenetic signal in 12 out of 25 phytosterols (percentage values of individual compounds), of which 7 were significant for both Pagel's λ and Blomberg's K, and 5 for only one of the tests (Fig. 2; Table 2). When grouping phytosterols based on the substitution at C-24 (C-24 methyl-, C-24 ethyl-, or C-24 no substitution or C-24 0) or based on the position of methine in ring B (Δ^0 , Δ^5 , Δ^7 , Δ^8), we found a significant phylogenetic signal (both Pagel's λ and Blomberg's K) for all groups except the Δ^8 sterols (Fig. 2; Table 2). Additionally, we found a significant signal for the Shannon diversity index and total sterol content ($\mu\text{g sterol mg}^{-1}$ pollen; Fig. 2; Table 2). These results remain largely consistent when excluding all taxa which are only identified to genus level. Note that λ and K are largely in agreement on which phytosterols showed significant signal (Table 2), although the significant estimates for λ are relatively high (0.585 to 1, mean = 0.847 for individual compounds; 0.668 to 0.906, mean = 0.79 for categories), whereas those for K are comparatively low (0.183 to 0.505, mean = 0.332 for individual compounds; 0.158 to 0.201, mean = 0.182 for categories). Some phylogenetic clustering of plants by overall sterol compositional similarity was also apparent in the NMDS plot, with, for example, plants in the Asteraceae, Asparagaceae, or Cactaceae sharing similar sterol profiles (Fig. S2).

Fig. 2 Pollen sterol profiles of plant species. Phylogenetic relationships are given on the left; bold numbers indicate families. Relative contributions of individual sterols to each species' total sterol content are given in the centre; commonness (proportion of plants containing an individual sterol) and relative abundance (average proportion of individual sterol in each species) are given at the bottom. Deeper reds indicate values closer to 1. Shannon diversity index (H), C-24 groups (C0, C1, C2), Δ groups (D0, D5, D7, D8), and total sterol content are given on the right. Circle size represents sums of relative sterol contents in the respective groups (0 to 1), and log of $\mu\text{g mg}^{-1}$ pollen for total sterol content. Sterol names and groups are coloured in the same fashion as illustrated in Fig. 1. Families: 1, Pinaceae; 2, Nymphaeaceae; 3, Colchicaceae; 4, Cannaceae; 5, Strelitziaceae; 6, Iridaceae; 7, Asphodelaceae; 8, Asparagaceae; 9, Amaryllidaceae; 10, Ranunculaceae; 11, Papaveraceae; 12, Paeoniaceae; 13, Geraniaceae; 14, Myrtaceae; 15, Onagraceae; 16, Cistaceae; 17, Malvaceae; 18, Oxalidaceae; 19, Salicaceae; 20, Linaceae; 21, Euphorbiaceae; 22, Fagaceae; 23, Cucurbitaceae; 24, Rosaceae; 25, Fabaceae; 26, Droseraceae; 27, Caryophyllaceae; 28, Nyctaginaceae; 29, Cactaceae; 30, Hydrangeaceae; 31, Polemoniaceae; 32, Theaceae; 33, Ericaceae; 34, Primulaceae; 35, Araliaceae; 36, Apiaceae; 37, Adoxaceae; 38, Caprifoliaceae; 39, Campanulaceae; 40, Menyanthaceae; 41, Asteraceae; 42, Apocynaceae; 43, Convolvulaceae; 44, Solanaceae; 45, Boraginaceae; 46, Gesneriaceae; 47, Scrophulariaceae; 48, Plantaginaceae; 49, Bignoniaceae; 50, Phrymaceae; 51, Lamiaceae.






Fig. 3 Hypothetical biosynthetic pathways of phytosterols identified in this study (pathways based on Benveniste, 2004). Details of C-24 and delta groups see Fig. 1.

range, but in general the explained variance (r^2) was low (Fig. 4; Table S4). Specifically, total sterol content correlated with environmental PC1 (associated with high mean temperatures, low temperature seasonality, and low soil carbon content; $P=0.015$,

$r^2=0.060$; Fig. 4). For linear models of individual environmental variables, species with higher total pollen sterol content tended to occur in locations with higher annual mean temperature, higher temperatures in the coldest quarter, higher precipitation in the

Table 1 Factor analysis identifying the covariance of 25 sterols measured across all the plant species surveyed (the main contributor(s) for each component is (are) highlighted in grey).

Component	1	2	3	4	5	6	7	8	9	10	11	12
Eigenvalue	2.146	2.053	1.874	1.761	1.581	1.460	1.421	1.356	1.234	1.105	1.098	1.069
Proportion of variance explained	0.086	0.082	0.075	0.070	0.063	0.058	0.057	0.054	0.049	0.044	0.044	0.043
Cumulative proportion of variance explained	0.086	0.168	0.243	0.313	0.376	0.434	0.491	0.545	0.594	0.638	0.682	0.725
Sterol	Factor loading											
Cycloartenol	−0.013	0.113	0.054	−0.683	−0.061	0.042	0.073	0.163	0.338	−0.131	0.107	−0.217
31-Norcycloartenol	−0.234	0.131	−0.022	−0.677	0.028	−0.009	−0.199	−0.251	−0.168	−0.094	−0.149	0.156
24,25-Dehydropollinastanol	0.019	0.065	−0.919	0.027	−0.022	−0.028	−0.060	0.049	−0.072	0.035	0.018	0.066
Pollinastanol	0.112	0.278	0.139	0.102	−0.046	−0.309	−0.415	0.278	−0.168	−0.286	−0.143	−0.027
Lathosterol	−0.021	0.039	−0.157	0.026	−0.004	−0.018	0.114	0.032	0.086	−0.040	−0.205	−0.740
Cholesterol	0.068	0.048	0.072	0.113	−0.194	0.670	0.090	−0.150	−0.081	0.091	−0.222	−0.080
31-Norcycloartenol	0.033	0.070	−0.924	0.032	−0.021	−0.033	0.024	0.067	0.011	0.013	−0.029	−0.222
14-Methylcholest-8-enol	0.064	−0.018	−0.038	0.048	−0.036	0.014	−0.736	−0.095	0.233	0.018	−0.001	−0.009
Desmosterol	0.051	0.077	−0.082	0.125	−0.069	−0.086	0.272	0.170	0.320	−0.110	−0.418	0.571
24-Methylencholesterol	0.208	0.531	0.256	0.266	−0.157	−0.378	−0.036	0.282	−0.350	0.215	−0.027	−0.073
24-Methylenecycloartenol	0.035	−0.036	0.035	−0.794	−0.035	−0.066	0.009	0.130	−0.054	0.128	−0.022	0.016
Cycloeucalenol	0.001	0.076	0.037	−0.001	−0.035	−0.017	−0.013	0.053	−0.057	−0.925	0.004	−0.005
Obtusifolol	−0.022	0.115	0.067	−0.003	0.003	−0.082	−0.143	0.052	0.801	0.084	−0.014	−0.005
Iso-obtusifolol	−0.741	−0.138	−0.005	−0.049	0.017	−0.026	−0.243	0.255	−0.070	0.071	−0.006	0.001
24-Methylenelophenol	−0.823	0.085	0.018	−0.079	0.056	−0.049	0.114	−0.036	−0.086	0.027	−0.062	0.053
Episterol	0.014	0.072	−0.002	0.058	−0.043	−0.098	0.088	−0.033	0.013	−0.009	0.842	0.117
Epifungisterol	−0.414	−0.005	−0.042	−0.211	0.041	0.056	−0.641	0.016	−0.091	−0.018	−0.072	0.101
Campesterol	0.052	−0.756	0.101	0.113	−0.101	−0.131	0.079	−0.024	−0.121	0.107	0.096	0.009
Campestanol	0.013	−0.042	0.019	0.028	0.823	−0.172	0.048	−0.076	0.046	0.006	−0.064	−0.056
Avenasterol	−0.752	0.078	0.044	0.045	−0.066	0.126	−0.022	0.026	0.250	−0.090	0.090	−0.128
Schottenol	−0.101	0.133	0.032	−0.031	0.250	0.778	−0.143	0.297	−0.045	−0.069	0.077	0.064
Sitosterol	0.104	−0.746	0.085	0.028	0.143	0.027	−0.005	−0.263	−0.036	−0.024	−0.152	−0.044
Sitostanol	−0.030	0.059	0.021	0.031	0.838	0.229	−0.029	0.155	−0.053	0.038	0.032	0.044
Isofucosterol	0.162	−0.091	0.133	0.121	−0.092	−0.072	−0.090	−0.771	−0.086	0.084	0.048	−0.010
Stigmasterol	−0.076	−0.653	0.018	0.031	−0.099	−0.095	−0.110	0.411	−0.059	0.085	−0.068	0.052

wettest quarter, and lower temperature seasonality (P -values < 0.05 for linear models of phylogenetic independent contrasts, r^2 between 0.05 and 0.08, Table S4), as is the case in tropical conditions. For Shannon's H diversity of pollen sterol profiles, the only significant association with environmental variables was a weak negative correlation with temperature seasonality ($P = 0.014$; $r^2 = 0.06$) (Table S4). None of the other environmental variables or principal components were significantly correlated with sterol content or diversity, nor was the total environmental niche breadth (Fig. 4; Table S4).

Sterols and pollinator guilds

We found that overall pollen sterol profiles were largely overlapping between plant groups with different pollinator guilds (bee, Lepidoptera, generalist insect, bird, unknown; Fig. 5a; ANOSIM: among group $R = -0.0069$, $P = 0.57$; no significant difference for any pairwise group comparison), and between plants with or without pollen as reward for bee pollinators (Fig. 5b; ANOSIM: $R = 0.033$, $P = 0.15$). This suggests that pollinator guilds or the use of pollen as reward by bees do not explain differences in pollen sterol composition. We note that wind pollinated angiosperms were not part of this dataset.

Neither 24-methylencholesterol nor total sterol content differed between plants with/without pollen as reward for bees (phylogenetic ANOVA: $P = 0.46$ and 0.66 respectively; Figs S3, S4).

Discussion

Phytosterols are primary metabolites in plants and are also essential nutrients for phytophagous insects, making them an important functional trait that provides a mechanistic link between plants and insects. Our study focused on the pollen sterol patterns across plant taxa, aiming to provide a more comprehensive overview of pollen sterol diversity and its relationship with plant phylogeny, abiotic environmental conditions, and pollinator guilds. We analysed 25 phytosterols in the pollen of > 120 angiosperms representing 51 plant families and identified covariance patterns among these phytosterols. Our data are the first to show significant phylogenetic signal for pollen phytosterols. Although environmental factors and pollinator guilds showed either weak or no relationships with pollen sterol content, future studies with more stratified sampling based on more finely defined pollinator species and environmental conditions could bring more insights about the drivers and importance of pollen sterol diversity.

Table 2 Identity of phytosterols in pollen of 122 plant species showing those with phylogenetic signal across species.

Trivial name	Semi-systematic name	Δ	C-24	λ	$p\text{-}\lambda$	K	$p\text{-}K$
Cycloartenol	4,4,14-trimethyl 9 β ,19-cyclo-cholest-24-en-3b-ol	0	C0	<0.001	1.000	0.103	0.407
31-Norcycloartenol	4,14-dimethyl 9 β ,19-cyclo-cholest-24-en-3b-ol	0	C0	1.003	0.001	0.505	0.021
24,25-Dehydropollinastanol	14-methyl 9 β ,19-cyclo-cholest-24-en-3b-ol	0	C0	1.003	0.001	0.354	0.036
Pollinastanol	14-methyl 9 β ,19-cyclo-cholestan-3b-ol	0	C0	0.681	0.001	0.169	0.107
Lathosterol	cholest-7-en-3b-ol	7	C0	<0.001	1.000	0.077	0.754
Cholesterol	cholest-5-en-3b-ol	5	C0	0.076	1.000	0.039	0.938
31-Norcycloartanol	4,14-dimethyl 9 β ,19-cyclo-cholestan-3b-ol	0	C0	1.003	0.001	0.208	0.050
14-Methylcholest-8-enol	14-methyl cholest-8-en-3b-ol	8	C0	<0.001	1.000	0.093	0.615
Desmosterol	cholesta-5,24-dien-3b-ol	5	C0	0.263	1.000	0.196	0.072
24-methylencholesterol	24-methyl cholesta-5,24(28)-dien-3b-ol	5	C1	0.747	0.001	0.183	0.012
24-Methylenecycloartanol	4,4,14,24-tetramethyl 9 β ,19-cyclo-cholest-24(28)-en-3b-ol	0	C1	<0.001	1.000	0.162	0.227
Cycloeucalenol	4,14,24-trimethyl 9 β ,19-cyclo-cholest-24(28)-en-3b-ol	0	C1	<0.001	1.000	0.219	0.147
Obtusifolol	4,14,24-trimethyl cholesta-8,24(28)-dien-3b-ol	8	C1	<0.001	1.000	0.167	0.132
Iso-obtusifolol	4,14,24-trimethyl cholesta-7,24(28)-dien-3b-ol	7	C1	0.973	0.001	0.250	0.039
24-Methylenelophenol	4,24-dimethyl cholesta-7, 24(28)-dien-3b-ol	7	C1	<0.001	1.000	0.102	0.495
Episterol	24-methyl cholesta-7,24(28)-dien-3b-ol	7	C1	0.990	0.001	0.505	0.021
Epifungisterol	24 α -methyl cholest-7-en-3b-ol	7	C1	0.694	0.018	0.236	0.055
Campesterol	24 α -methyl cholest-5-en-3b-ol	5	C1	0.585	0.005	0.240	0.011
Campestanol	24 α -methyl cholestan-3b-ol	0	C1	<0.001	1.000	0.290	0.119
Avenasterol	24-ethyl cholesta-7,24(28) trans-dien-3b-ol	7	C2	<0.001	1.000	0.069	0.814
Schottenol	24 α -ethyl cholest-7-en-3b-ol	7	C2	1.003	0.001	0.440	0.008
Sitosterol	24 α -ethyl cholest-5-en-3b-ol	5	C2	0.275	1.000	0.183	0.012
Sitostanol	24 α -ethyl cholestan-3b-ol	0	C2	<0.001	1.000	0.093	0.606
Isofucosterol	24-ethyl cholesta-5,24(28) trans-dien-3b-ol	5	C2	0.634	0.002	0.141	0.053
Stigmasterol	24 α -ethyl cholesta-5,22 trans-dien-3b-ol	5	C2	<0.001	1.000	0.149	0.200
H-diversity	Shannon Index			0.834	0.001	0.177	0.004

Grouped sterols	Grouping description				
Sum_C0	No substitution at C-24	0.906	0.001	0.196	0.025
Sum_C1	CH _n substitution at C-24	0.689	0.001	0.158	0.012
Sum_C2	C ₂ H _n substitution at C-24	0.668	0.001	0.169	0.002
Sum_D0	Phytosterols (Saturated ring B)	0.880	0.004	0.177	0.032
Sum_D5	Δ^5 Sterols	0.777	0.001	0.201	0.007
Sum_D7	Δ^7 Sterols	0.825	0.001	0.188	0.035
Sum_D8	Δ^8 Sterols	0.022	0.114	0.167	0.152
Total sterol		1.002	0.001	0.293	0.026

All values presented are based on the percentage values of sterols except total sterol content ($\mu\text{g mg}^{-1}$ of sampled pollen). Δ and C-24 value indicates structure of ring B and on the C-24 (see Fig. 1 for details). Pagel's λ and Blomberg's K are used for testing phylogenetic signal. P -values for each test are given accordingly ($p\text{-}\lambda$ and $p\text{-}K$). Sterols with significant phylogenetic signals are in bold.

Table 3 Multiple regression on distance matrices (MRM) analysis results showing regression coefficients and P -values for the multiple regression of pairwise distances on the first three environmental principal components (PC1–PC3) and phylogenetic distances against the sterol profile Bray–Curtis distance matrix.

Variable	Regression coefficient	P -value
Intercept	7.89×10^{-1}	0.24
Phylogenetic distance	2.27×10^{-5}	0.82
PC1	1.87×10^{-2}	0.0587
PC2	-2.0×10^{-2}	0.0585
PC3	-1.07×10^{-2}	0.3635

Phylogenetic structure of pollen sterols

Similar pollen sterol profiles in related taxa are ultimately due to shared evolutionary history and proximately due to shared genes

for the enzymes involved in their biosynthesis. Indeed, we show that phylogenetic patterns in pollen sterols in part reflect their relations in the underlying biosynthesis pathway. For example, we observed significant phylogenetic clustering of plant species whose pollen sterol profiles are dominated by C-24 methyl (C1-group), C-24 ethyl (C2-group) or nonsubstituted (C-24 C0-group) phytosterols (Fig. 2; Table 2), reflecting the bifurcation of biosynthesis pathways (Fig. 3). Key enzymes (SMTs, sterol methyltransferases) that bifurcate the phytosterol pathways are SMT1, which methylates cycloartenol (unsubstituted at C-24) to 24-methylenecycloartanol, and SMT2, the key and effective enzyme for the methylation of C-24 methyl to C-24 ethyl sterols (Akihisa *et al.*, 1991; Nes, 2000; Schaeffer *et al.*, 2001; Neelakanandan *et al.*, 2009). Based on our findings, it would seem likely that the expression of these enzymes follows phylogenetically conserved patterns in different clades. Similarly, phylogenetic clusters of the main sterol groups based on the presence or absence of and




Fig. 4 Correlation plots of phylogenetically independent contrasts (PICs) of positions on the environmental principal component axes (PC1–PC3) and environmental niche breadth against PICs of total pollen sterol amounts (top row) and sterol profile Shannon diversity indices (H, bottom row). Blue dashed lines indicate regression lines of linear models (with intercept set to zero); r^2 and P -values for linear models have been inserted into their respective plots. For PC loadings from each environmental variable see Supporting Information Table S3.

position of the double bond in ring B (e.g. Δ^5 , Δ^7 , or saturated ring B) suggest conserved expressions of specific desaturases (e.g. STE1, sterol-C5-desaturase) which convert saturated carbon bonds to methines, and reductases (e.g. DWF5, sterol- Δ^7 -reductase) which reduce methines to saturated bonds (Benveniste, 2004; Villette *et al.*, 2015).

Our factor analysis (Table 1) further revealed an inverse relationship between the abundance of the major $\Delta^{24,28}$ sterol (24-methylenecholesterol) and the C-24,28-saturated sterols: campesterol, sitosterol and stigmasterol. This suggests an overall trade-off between these groups, the balance of which may be governed by DWF1 (sterol- Δ^{24} -reductase) activity. Data from previous studies (Villette *et al.*, 2015; Vanderplanck *et al.*, 2020a) suggests a similar inverse correlation between 24-methylenecholesterol and the C-24 ethyl sterols, although this has not been explicitly stated. A high ratio of 24-methylenecholesterol to C-24,28-saturated sterols is evident in Cactaceae, Droseraceae, Rosaceae, Onagraceae, and Paeoniaceae. Conversely, C-24,28-saturated sterols are more abundant than 24-methylenecholesterol in Ericaceae, Primulaceae, Salicaceae and Amaryllidaceae. These families are not closely related, suggesting convergent evolution of sterol composition. Overall, this indicates an interplay of environmental selection pressures for particular structural groups and phylogenetic constraints on sterol biosynthesis enzyme expression.

The composition of phytosterols appears to be tissue-dependent (Nes, 1990; Nes *et al.*, 1993). For example, 24-methyleneccholesterol has been identified as the main pollen sterol in many Cactaceae (Fig. 2; Table S1) but is not abundant in their photosynthetic tissue (Standifer *et al.*, 1968; Salt *et al.*, 1987; Lusby *et al.*, 1993; Li, 1996). The unique functional roles in pollen

development when compared to the sporophyte may contribute to the distinct sterol profiles in pollen. We observed strong correlations among early, cyclopropyl sterol intermediates of the sterol pathway, particularly 9 β ,19-cyclopropyl sterols (Table 1). Cycloartenol, 31-norcycloartenol, and 24-methylenecycloartenol are correlated with each other: 31-norcycloartenol and 24-methylenecycloartenol are both derived from cycloartenol. 24,25-Dehydropollinastanol and 31-norcycloartenol also show high correlation and both are derived from 31-norcycloartenol. Co-occurrence of cyclopropyl sterols suggests a reduction in CPI1 (cyclopropyl isomerase) activity and truncation of the sterol pathway, either within the pollen grain or in the surrounding tapetum cells from which pollen coat sterols are derived. 9 β ,19-Cyclopropyl sterols have been identified as key components of the pollen coat in *Brassica napus* L. (Wu *et al.*, 1999; Villette *et al.*, 2015). In addition, cycloeucalenol is the main sterol synthesised in the growing pollen tube of *Nicotiana tabacum* (Villette *et al.*, 2015; Rotsch *et al.*, 2017).

Correlations of phytosterols with abiotic factors

The presence of different phytosterols could be evolutionary adaptations to environmental conditions. We detected a positive relationship between sterol content and temperature (particularly mean annual temperature and mean temperature of the coldest quarter), and a negative correlation with temperature seasonality, even though the overall association strength was low (Table S4). This indicates that plants found in cool and temperate climatic conditions were likely to have less pollen sterol than those found in areas of the world with warm climates and little seasonal




Fig. 5 2D-nMDS plots of sterol profiles for plants (a) with different pollinator guilds, and (b) with/without pollen as bee reward. Distances correspond to sterol profile dissimilarity (Bray–Curtis distances). Stress of NMDS solution: 0.202.

fluctuation (e.g. tropical climates). The association between warmer climates and higher total amounts of pollen sterols may have evolved as protection against membrane heat stress – the

role of phytosterols in adaptation to high temperature stress is established (Dufourc, 2008; Narayanan *et al.*, 2016). Phytosterols including campesterol, sitosterol, and avenasterol degrade

in stored grain more rapidly at higher temperatures (Wawrzyniak *et al.*, 2019), so higher sterol content in warmer climates may avert the risk of their rapid breakdown and limited availability. Besides this, other pollen sterol characteristics (e.g. sterol diversity and the overall pollen sterol composition) were not notably associated with abiotic factors. Our sampling, however, was biased towards plants originating from temperate regions (the predominant species available to us for sampling). Limited sampling towards extremes of the environmental gradients may have reduced our scope for the detection of associations between abiotic factors and pollen sterol characteristics. Future work should therefore be targeted at sampling additional plant species of more extreme environments to fill this gap. Note that our species were sampled at glasshouses (e.g. tropical glasshouse, alpine glasshouse) or outdoors at Royal Botanic Gardens Kew and nearby areas (Table S1) to get a first estimate of pollen sterol diversity across a broad range of species. Future in-depth studies on how abiotic conditions affect pollen sterol variation within species deserve further attention to build a more complete overview of pollen sterol diversity at different taxonomic levels.

Impact of sterol diversity on pollinators

Pollen sterol amount and composition did not differ significantly between bee pollinated and non-bee-pollinated plant species. This could indicate that pollen sterols have generally not been under selection by bee pollinators, although we acknowledge that our analysis combined all bee pollinated plants into one group. Therefore, it remains possible that pollen sterols play a role in finer scale interactions between different bee species of varying levels of pollen specialization and their host plants. We also note that, although we based our assessment of pollinator guilds on the best available data in the literature, the quality of evidence for the effective pollinators of the plants in our data set varied. This calls for further in-depth studies of the relationships between pollen sterols and pollinators, also including wind-pollinated angiosperm taxa missing in this work as points of comparison to animal pollinated plants.

A major knowledge gap exists in understanding how important specific phytosterols are for bees, particularly wild bee species, since many of them are pollen specialists. Plants adapt nectar chemistry to suit the specific needs of pollinators (Vandelook *et al.*, 2019) and could similarly alter nutritional chemistry of pollen to optimize its nutritional suitability for flower visitors. Bee pollinators require a dietary source of sterols (Wright *et al.*, 2018), and for this they must use the phytosterols found in pollen. Therefore, determining how lipid components of pollen vary qualitatively and quantitatively across different plant taxa is important in understanding how nutritionally limiting landscapes might be for bees, especially where they are not botanically diverse. For example, honeybee colony growth benefits from 24-methylenecholesterol (Herbert *et al.*, 1980). Thus, honeybees may be nutritionally limited in landscapes where floral resources do not provide 24-methylenecholesterol. Our data suggested that many Asteraceae (e.g. *Achillea ptarmica* L., *Tanacetum vulgare* L., *Achillea millefolium* L., *Jacobaea vulgaris* Gaertn., *Centaurea nigra*

L. and *Cirsium vulgare* (Savi.) Ten) are rich in Δ^7 -sterols (Fig. 2; Table S1) and lack the common honeybee-favourable Δ^5 -sterols (e.g. 24-methylenecholesterol). Δ^7 -sterols are known to be toxic to nonspecialist herbivores and can only be utilized by some insect species (Behmer & Nes, 2003; Lang *et al.*, 2012). Thus, plant species that produce unusual phytosterols in pollen may produce these as a defence against pollen herbivory, but some specialist bee species may have developed mechanisms overcoming this defense. Indeed, pollen foraging bees on Asteraceae plants are mostly specialized oligolectic bees, while polylectic bee species avoid the pollen despite the ubiquitous distribution of Asteraceae species and their substantial amount of pollen provision; this is known as the Asteraceae paradox (Müller & Kuhlmann, 2008). While the reasons for the Asteraceae paradox remain unclear, the abundance of Δ^7 -sterols we found in the pollen of Asteraceae species could provide a potential explanation (see also Vanderplanck *et al.*, 2018, 2020b).






Acknowledgements







We thank Wenxu Zhou for his guidance in analysing sterols, Amy Kendal-Smith, Richard Moore, and James Woodward for their help with pollen collection, Geoffrey Kite for support with the GC-MS, and Ellen Baker for helping revise the manuscript. We also thank the Horticultural staff of the Royal Botanic Gardens, Kew for maintenance of the outstanding living collections used to provide pollen for this study. The project was funded by a BBSRC grant to GAW (BB/P007449/1) and PCS (BB/P005276/1) and a Peter Sowerby Grant to PCS and HK. PZ was supported to work with DN at Texas Tech on a BBSRC International Travel Award Scheme grant (BB/S004653/1) to PCS. EM was funded by a BBSRC iCASE grant (BB/M011224/1) to the University of Oxford.

Author contributions

PZ, HK, and PCS designed the research. PZ collected the pollen and extracted and analysed sterols. WDN, IWF, and DIF helped with sterol identification and quantification. OS collected phylogenetic information on studied species and conducted phylogenetic analyses. SP, CP, and IO collected species geographic and environmental information and performed abiotic niche analyses. HK and PZ collected pollinator records on studied species, and HK conducted analyses with environmental factors and pollinator guilds. EM and GAW conducted factor analysis. WDN generated sterol biosynthesis pathways. PZ and PCS drafted the manuscript. HK, OS, SP, GAW and all other authors contributed to writing and revising the manuscript.

ORCID

Dudley I. Farman  <https://orcid.org/0000-0003-3579-3672>
Iain W. Farrell  <https://orcid.org/0000-0002-2257-8853>
Hauke Koch  <https://orcid.org/0000-0002-2694-7775>
Elynor Moore  <https://orcid.org/0000-0003-1531-6393>
Ian Ondo  <https://orcid.org/0000-0001-7816-5882>

Charlotte Phillips  <https://orcid.org/0000-0002-2200-9677>
 Samuel Pironon  <https://orcid.org/0000-0002-8937-7626>
 Orlando Schwery  <https://orcid.org/0000-0001-7373-7407>
 Philip C. Stevenson  <https://orcid.org/0000-0002-0736-3619>
 Geraldine A. Wright  <https://orcid.org/0000-0002-2749-021X>
 Pengjuan Zu  <https://orcid.org/0000-0002-3222-598X>

Data availability

All the source data used for this publication can be found in the Supporting Information Tables. Related codes for analyses can be found at: <http://doi.org/10.5281/zenodo.4448216>.

References

- Akihisa T, Kokke WCMC, Tamura T. 1991. Naturally occurring sterols and related compounds from plants. In: Patterson GW, Nes WD, eds. *Physiology and biochemistry of sterols*. Champaign, IL, USA: AOCS Press, 172–228.
- Amar S, Becker HC, Möllers C. 2008. Genetic variation and genotype × environment interactions of phytosterol content in three doubled haploid populations of winter rapeseed. *Crop Science* 48: 1000–1006.
- Arnold SEJ, Peralta Idrovo ME, Lomas Arias LJ, Belmain SR, Stevenson PC. 2014. Herbivore defence compounds occur in pollen and reduce bumblebee colony fitness. *Journal of Chemical Ecology* 40: 878–881.
- Behmer ST, Elias DO. 1999. The nutritional significance of sterol metabolic constraints in the generalist grasshopper *Schistocerca americana*. *Journal of Insect Physiology* 45: 339–348.
- Behmer ST, Elias DO. 2000. Sterol metabolic constraints as a factor contributing to the maintenance of diet mixing in grasshoppers (Orthoptera: Acrididae). *Physiological and Biochemical Zoology* 73: 219–230.
- Behmer ST, Nes WD. 2003. Insect sterol nutrition and physiology: a global overview. *Advances in Insect Physiology* 31: 1–72.
- Benveniste P. 2004. Biosynthesis and accumulation of sterols. *Annual Review of Plant Biology* 55: 429–457.
- Blomberg SP, Garland T Jr, Ives AR, Crespi B. 2003. Testing for phylogenetic signal in comparative data: behavioral traits are more labile. *Evolution* 57: 717–745.
- Brummitt RK. 2001. *World geographical scheme for recording plant distributions*, 2nd edn. Geneva Switzerland: Biodiversity Information Standards (TDWG). [WWW document] URL <http://www.tdwg.org/standards/109> [accessed 18 March 2020].
- Capinha C, Pateiro-López B. 2014. Predicting species distributions in new areas or time periods with alpha-shapes. *Ecological Informatics* 24: 231–237.
- Chakrabarti P, Lucas HM, Sagili RR. 2019. Evaluating effects of a critical micronutrient (24-methylenecholesterol) on honey bee physiology. *Annals of the Entomological Society of America* 113: 176–182.
- Chung HY, Fujioka S, Choe S, Lee S, Lee YH, Baek NI, Chung IS. 2010. Simultaneous suppression of three genes related to brassinosteroid (BR) biosynthesis altered campesterol and BR contents, and led to a dwarf phenotype in *Arabidopsis thaliana*. *Plant Cell Reports* 29: 397–402.
- Clarke KR. 1993. Non-parametric multivariate analyses of changes in community structure. *Australian Journal of Ecology* 18: 117–143.
- Danielson JJ, Gesch DB. 2011. *Global multi-resolution terrain elevation data 2010 (gmtd2010)*. Technical report. Reston, VA, USA: US Geological Survey.
- Dauby G, Zaiss R, Blach-Overgaard A, Catarino L, Damen T, Deblauwe V, Dessein S, Dransfield J, Droissart V, Duarte MC *et al.* 2016. Rainbio: a mega-database of tropical African vascular plants distributions. *PhytoKeys* 74: 1–18.
- Dornelas M, Antao LH, Moyes F, Bates AE, Magurran AE, Adam D, Akhmetzhanova AA, Appeltans W, Arcos JM, Arnold H *et al.* 2018. Biotime: a database of biodiversity time series for the anthropocene. *Global Ecology and Biogeography* 27: 760–786.
- Dufourc EJ. 2008. The role of phytosterols in plant adaptation to temperature. *Plant Signalling & Behavior* 3: 133–134.
- Felsenstein J. 1985. Phylogenies and the comparative method. *American Naturalist* 125: 1–15.
- Ferrer A, Altabella T, Arró M, Boronat A. 2017. Emerging roles for conjugated sterols in plants. *Progress in Lipid Research* 67: 27–37.
- Garland T, Dickerman AW, Janis CM, Jones JA. 1993. Phylogenetic analysis of covariance by computer simulation. *Systematic Biology* 42: 265–292.
- Goslee SC, Urban DL. 2007. The ecodist package for dissimilarity-based analysis of ecological data. *Journal of Statistical Software* 22: 1–19.
- Grove MD, Spencer GF, Rohwedder WK, Mandava N, Worley JF, Warthen JD, Steffens GL, Flippen-Anderson JL, Cook JC. 1979. Brassinolide, a plant growth-promoting steroid isolated from *Brassica napus* pollen. *Nature* 281: 216–217.
- Grunwald C. 1971. Effects of free sterols, steryl ester, and steryl glycoside on membrane permeability. *Plant Physiology* 48: 653–655.
- Guo D, Venkatramesh M, Nes WD. 1995. Developmental regulation of sterol biosynthesis in *Zea mays*. *Lipids* 30: 203–219.
- Hammer Ø, Harper DAT, Ryan PD. 2001. PAST: Paleontological statistics software package for education and data analysis. *Palaeontologia Electronica* 4: 1–9.
- Hartmann M-A. 1998. Plant sterols and the membrane environment. *Trends in Plant Science* 3: 170–175.
- Herbert EW, Svoboda JA, Thompson MJ, Shimanuki H. 1980. Sterol utilization in honey bees fed a synthetic diet: effects on brood rearing. *Journal of Insect Physiology* 26: 287–289.
- Heupel RC, Nes WD. 1984. Evidences for differences in sterol biosynthesis and derivatization in Sorghum. *Journal of Natural Products* 47: 292–299.
- Hinchliff CE, Smith SA, Allman JF, Burleigh JG, Chaudhary R, Coghill LM, Crandall KA, Deng J, Drew BT, Gazis R *et al.* 2015. Synthesis of phylogeny and taxonomy into a comprehensive tree of life. *Proceedings of the National Academy of Sciences, USA* 112: 12764–12769.
- ISRIC. 2013. *Soilgrids: an automated system for global soil mapping*. Wageningen, the Netherlands: International Soil Reference and Information Centre [WWW document] URL <https://data.isric.org>. [accessed 15 May 2020].
- Jing X, Behmer ST. 2020. Insect sterol nutrition: physiological mechanisms, ecology, and applications. *Annual Review of Entomology* 65: 251–271.
- Karger DN, Conrad O, Böhner J, Kawohl T, Kreft H, Soria-Auza RW, Zimmermann NE, Linder HP, Kessler M. 2017. Climatologies at high resolution for the earth's land surface areas. *Scientific Data* 4: 170122.
- Keck F, Rimet F, Bouchez A, Franc A. 2016. phylosignal: an R package to measure, test, and explore the phylogenetic signal. *Ecology and Evolution* 6: 2774–2780.
- Kembel S, Cowan P, Helmus M, Cornwell W, Morlon H, Ackerly D, Blomberg S, Webb C. 2010. Picante: R tools for integrating phylogenies and ecology. *Bioinformatics* 26: 1463–1464.
- Khrupach V, Zhabinsk V, de Groot A. 2000. Twenty years of brassinosteroids: steroidal plant hormones warrant better crops for the XXI century. *Annals of Botany* 86: 441–447.
- Knuth P. 1908. *Handbook of flower pollination*, vol. 2. Oxford, UK: Clarendon Press.
- Knuth P. 1909. *Handbook of flower pollination*, vol. 3. Oxford, UK: Clarendon Press.
- Kumar MS, Ali K, Dahuja A, Tyagi A. 2015. Role of phytosterols in drought stress tolerance in rice. *Plant Physiology and Biochemistry* 96: 83–89.
- Lang M, Murat S, Clark AG, Goupil G, Blais C, Matzkin LM, Guittard E, Yoshiyama-Yanagawa T, Kataoka H, Niwa R *et al.* 2012. Mutations in the neverland gene turned *Drosophila pachea* into an obligate specialist species. *Science* 337: 1658–1661.
- Li Y. 1996. *Stereochemical studies on the metabolism of sterols by Saccharomyces cerevisiae strain GL7*. PhD thesis, Texas Tech University, TX, USA.
- Lichstein JW. 2007. Multiple regression on distance matrices: a multivariate spatial analysis tool. *Plant Ecology* 188: 117–131.
- Lusby WR, Buchmann SL, Feldlaufer MF. 1993. Pollen sterols from three species of Sonoran cacti. *Lipids* 28: 469–470.
- Meyer C, Weigelt P, Kreft H. 2016. Multidimensional biases, gaps and uncertainties in global plant occurrence information. *Ecology Letters* 19: 992–1006.
- Michonneau F, Brown JW, Winter DJ. 2016. rotl: an R package to interact with the open tree of life data. *Methods in Ecology and Evolution* 7: 1476–1481.

- Müller A, Kuhlmann M. 2008. Pollen hosts of western palaearctic bees of the genus *Colletes* (Hymenoptera: Colletidae): the Asteraceae paradox. *Biological Journal of the Linnean Society* 95: 719–733.
- Narayanan S, Tamura PJ, Roth MR, Prasad PV, Welti R. 2016. Wheat leaf lipids during heat stress: I. High day and night temperatures result in major lipid alterations. *Plant, Cell & Environment* 39: 787–803.
- Neelakandan AK, Song Z, Wang J, Richards MH, Wu X, Valliyodan B, Nguyen HT, Nes WD. 2009. Cloning, functional expression, and phylogenetic analysis of plant 24C-methyltransferases involved in sitosterol biosynthesis. *Phytochemistry* 70: 1982–1998.
- Nes WD. 1990. Control of sterol biosynthesis and its importance to developmental regulation and evolution. *Recent Advances in Phytochemistry* 24: 283–327.
- Nes WD. 2000. Sterol methyl transferase: enzymology and inhibition. *Biochimica Biophysica Acta* 1529: 63–88.
- Nes WD. 2011. Biosynthesis of cholesterol and other sterols. *Chemical Reviews* 111: 6423–6451.
- Nes WR, Krevitz K, Joseph J, Nes WD, Harris B, Gibbons GF, Patterson GW. 1977. The phylogenetic distribution of sterols in tracheophytes. *Lipids* 12: 511–527.
- Nes WR, McKean ML. 1977. *Biochemistry of steroids and other isopentenoids*. Baltimore, MD, USA: University Park Press.
- Nes WD, Parker SR, Crumley FG, Ross SA. 1993. Regulation of phytosterol biosynthesis. In: Moore TS, ed. *Lipid metabolism in plants*. Boca Raton, FL, USA: CRC Press, 389–426.
- Nes WD, Song Z, Dennis AL, Zhou W, Nam J, Miller M. 2003. Biosynthesis of phytosterols: kinetic mechanism for the enzymatic C-methylation of sterols. *Journal of Biological Chemistry* 278: 34505–34516.
- Nes WD, Zhou W, Ganapathy K, Liu J, Vatsyayan R, Chamala S, Hernandez K, Miranda M. 2009. Sterol 24-C-methyltransferase: an enzymatic target for the disruption of ergosterol biosynthesis and homeostasis in *Cryptococcus neoformans*. *Archives of Biochemistry and Biophysics* 481: 210–218.
- Pagel M. 1999. Inferring the historical patterns of biological evolution. *Nature* 401: 877–884.
- Palmer-Young EC, Farrell IW, Adler IW, Milano NJ, Egan PA, Junker RR, Irwin RE, Stevenson PC. 2019. Chemistry of floral rewards: intra- and interspecific variability of nectar and pollen secondary metabolites across taxa. *Ecological Monographs* 89: e01335.
- Paradis E, Claude J, Strimmer K. 2004. Ape: analyses of phylogenetics and evolution in R language. *Bioinformatics* 20: 289–290.
- Piironen V, Lindsay DG, Miettinen TA, Toivo J, Lampi AM. 2000. Plant sterols: biosynthesis, biological function and their importance to human nutrition. *Journal of the Science of Food and Agriculture* 80: 939–966.
- Posé D, Castanedo I, Borsani O, Nieto B, Rosado A, Taconnat L, Ferrer A, Dolan L, Valpuesta V, Botella MA. 2009. Identification of the Arabidopsis dry2/sql1-5 mutant reveals a central role for sterols in drought tolerance and regulation of reactive oxygen species. *The Plant Journal* 59: 63–76.
- Qi X, Bakht S, Qin B, Leggett M, Hemmings A, Mellon F, Eagles J, Werck-Reichhart D, Schaller H, Lesot A *et al.* 2006. A different function for a member of an ancient and highly conserved cytochrome p450 family: from essential sterols to plant defense. *Proceedings of the National Academy of Sciences, USA* 103: 18848–18853.
- R Core Team. 2020. *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. R v.4.0.3 [WWW document] URL <https://www.R-project.org/>.
- Rasmussen C, Engel MS, Verecken NJ. 2020. A primer of host-plant specialization in bees. *Emerging Topics in Life Sciences* 4: 7–17.
- Rees JA, Cranston K. 2017. Automated assembly of a reference taxonomy for phylogenetic data synthesis. *Biodiversity Data Journal* 5: e12581.
- Revell LJ. 2012. Phytools: an R package for phylogenetic comparative biology (and other things). *Methods in Ecology and Evolution* 3: 217–223.
- Rivest S, Forrest JRK. 2020. Defence compounds in pollen: why do they occur and how do they affect the ecology and evolution of bees? *New Phytologist* 225: 1053–1064.
- Rotsch AH, Kopka J, Feussner I, Ischebeck T. 2017. Central metabolite and sterol profiling divides tobacco male gametophyte development and pollen tube growth into eight metabolic phases. *The Plant Journal* 92: 129–146.
- Salt TA, Tocker JE, Adler JH. 1987. Dominance of Δ^5 -sterols in eight species of the Cactaceae. *Phytochemistry* 26: 731–733.
- Sanchez-Reyes LL, O'Meara BC. 2019. Datelife: leveraging databases and analytical tools to reveal the dated Tree of Life. *bioRxiv*. doi: <https://doi.org/10.1101/782094>.
- Schaeffer A, Bronner R, Benveniste P, Schaller H. 2001. The ratio of campesterol to sitosterol that modulates growth in *Arabidopsis* is controlled by *STEROL METHYLTRANSFERASE 2;1*. *The Plant Journal* 25: 605–615.
- Schuler I, Milon A, Nakatani Y, Ourisson G, Albrecht A-M, Benveniste P, Hartman M-A. 1991. Differential effects of plant sterols on water permeability and on acyl chain ordering of soybean phosphatidylcholine bilayers. *Proceedings of the National Academy of Sciences, USA* 88: 6926–6930.
- Schwery O, O'Meara BC. 2016. MonoPhy: a simple R package to find and visualize monophyly issues. *PeerJ Computer Science* 2: e56.
- Smith SA, Beaulieu JM, Donoghue MJ. 2010. An uncorrelated relaxed-clock analysis suggests an earlier origin for flowering plants. *Proceedings of the National Academy of Sciences, USA* 107: 5897–5902.
- Standifer LN, Devys M, Barbier M. 1968. Pollen sterols—a mass spectrographic survey. *Phytochemistry* 7: 1361–1365.
- Svoboda JA, Thompson MJ, Herbert EW, Shimanuki H. 1980. Sterol utilization in honey bees fed a synthetic diet: Analysis of prepupal sterols. *Journal of Insect Physiology* 26: 287–289.
- Vandelook F, Janssens SB, Gijbels P, Fischer E, Van den Ende W, Honnay O, Abrahamczyk S. 2019. Nectar traits differ between pollination syndromes in Balsaminaceae. *Annals of Botany* 124: 269–279.
- Vanderplanck M, Decleves S, Roger N, Decroo C, Caulier G, Glauser G, Michez D. 2018. Is non-host pollen suitable for generalist bumblebees? *Insect Science* 25: 259–272.
- Vanderplanck M, Gilles H, Nonclercq D, Duez P, Gerbaux P. 2020b. Asteraceae paradox: chemical and mechanical protection of *Taraxacum* pollen. *Insects* 11: 304.
- Vanderplanck M, Zerck P-L, Lognay G, Michez D. 2020a. Generalized host-plant feeding can hide sterol-specialized foraging behaviors in bee-plant interactions. *Ecology and Evolution* 10: 150–162.
- Velasco L, Fernández-Cuesta Á, García-Ruiz JR, Fernández-Martínez JM, Domínguez-Giménez J. 2013. Genetic variation and genotype × environment interactions for seed phytosterols in sunflower. *Crop Science* 53: 1589–1593.
- Villette C, Berna A, Compagnon V, Schaller H. 2015. Plant sterol diversity in pollen from angiosperm. *Lipids* 50: 749–760.
- Wang K, Senthil-Kumar M, Ryu C-M, Kang L, Mysore KS. 2012. Phytosterols play a key role in plant innate immunity against bacterial pathogens by regulating nutrient efflux into the apoplast. *Plant Physiology* 158: 1789–1802.
- Wawrzyniak J, Gawrysiak-Witulska M, Rudzińska M. 2019. Dynamics of phytosterol degradation in a bulk of rapeseed stored under different temperature and humidity conditions. *Journal of Stored Products Research* 83: 292–304.
- WCVP. 2020. *World checklist of vascular plants, version 2.0*. London, UK: Royal Botanic Gardens, Kew [WWW document] URL <http://wcvp.science.kew.org/> [accessed 18 March 2020].
- Westrich P. 2018. *Die Wildbienen Deutschlands*. Stuttgart, Germany: Verlag Eugen Ulmer.
- Wright GA, Nicolson SW, Shafir S. 2018. Nutritional physiology and ecology of honey bees. *Annual Review of Entomology* 63: 327–344.
- Wu SS, Moreau RA, Whitaker BD, Huang AH. 1999. Steryl esters in the elaioplasts of the tapetum in developing *Brassica* anthers and their recovery on the pollen surface. *Lipids* 34: 517–523.
- Xu S, Norton RA, Crumley FG, Nes WD. 1988. Comparison of the chromatographic properties of sterols, select additional steroids and triterpenoids: gravity-flow liquid chromatography, thin-layer chromatography, gas-liquid chromatography, and high performance liquid chromatography. *Journal of Chromatography* 4: 377–398.
- Zanne AE, Tank DC, Cornwell WK, Eastman JM, Smith SA, FitzJohn RG, McGlinn DJ, O'Meara BC, Moles AT, Reich PB *et al.* 2014. Three keys to the radiation of angiosperms into freezing environments. *Nature* 506: 89–92.
- Zizka A, Silvestro D, Andermann T, Azevedo J, Duarte Ritter C, Edler D, Farooq H, Herdean A, Ariza M, Scharn R *et al.* 2019. Coordinated cleaner: standardized cleaning of occurrence records from biological collection databases. *Methods in Ecology and Evolution* 10: 744–751.

Supporting Information

Additional Supporting Information may be found online in the Supporting Information section at the end of the article.

Fig. S1 GC-MS spectra of the 25 phytosterols identified in our study (after Tri-sil derivatisation, extraction details see the Materials and Methods section).

Fig. S2 2D-NMDS plot: pollen sterol profile similarities between species of different plant families.

Fig. S3 24-methylenecholesterol content of plants without pollen as bee or with pollen as reward for bees.

Fig. S4 Total sterol content of plants without pollen as bee reward or with pollen as reward for bees.

Table S1 Data table (plant species, scores for different environmental variables/principal components, pollinator guilds, sterol composition including both relative and absolute amounts).

Table S2 Scientific name and family for all sampled species, along with suggested OTL synonyms and taxon IDs.

Table S3 Variable contributions to axes of PCA of 13 environmental variables.

Table S4 Test results: linear models of phylogenetic independent contrasts (PICs) of total sterol amount/diversity against PICs of environmental variables and niche breadth.

Please note: Wiley Blackwell are not responsible for the content or functionality of any Supporting Information supplied by the authors. Any queries (other than missing material) should be directed to the *New Phytologist* Central Office.



About New Phytologist

- *New Phytologist* is an electronic (online-only) journal owned by the New Phytologist Foundation, a **not-for-profit organization** dedicated to the promotion of plant science, facilitating projects from symposia to free access for our Tansley reviews and Tansley insights.
- Regular papers, Letters, Viewpoints, Research reviews, Rapid reports and both Modelling/Theory and Methods papers are encouraged. We are committed to rapid processing, from online submission through to publication 'as ready' via *Early View* – our average time to decision is <26 days. There are **no page or colour charges** and a PDF version will be provided for each article.
- The journal is available online at Wiley Online Library. Visit **www.newphytologist.com** to search the articles and register for table of contents email alerts.
- If you have any questions, do get in touch with Central Office (np-centraloffice@lancaster.ac.uk) or, if it is more convenient, our USA Office (np-usaoffice@lancaster.ac.uk)
- For submission instructions, subscription and all the latest information visit **www.newphytologist.com**