

1 **Nutritional niches reveal fundamental domestication tradeoffs in**  
2 **fungus-farming ants**

3  
4 Jonathan Z. Shik<sup>1,2,3\*</sup>, Pepijn W. Kooij<sup>2,4,9</sup>, David A. Donoso<sup>5,6</sup>, Juan C. Santos<sup>7</sup>, Ernesto  
5 B. Gomez<sup>3</sup>, Mariana Franco<sup>3</sup>, Antonin J. J. Crumière<sup>1</sup>, Xavier Arnan<sup>8,10</sup>, Jack Howe<sup>2,11</sup>,  
6 William T. Wcislo<sup>3</sup>, Jacobus J. Boomsma<sup>1,2</sup>

7  
8 <sup>1</sup>Section of Ecology and Evolution, Department of Biology, University of Copenhagen,  
9 Universitetsparken 15, 2100 Copenhagen, Denmark

10  
11 <sup>2</sup>Centre for Social Evolution, Department of Biology, University of Copenhagen,  
12 Universitetsparken 15, 2100 Copenhagen, Denmark

13  
14 <sup>3</sup>Smithsonian Tropical Research Institute, Apartado 0843-03092, Balboa, Ancon,  
15 Republic of Panama

16  
17 <sup>4</sup>Comparative Fungal Biology, Department of Comparative Plant and Fungal Biology,  
18 Royal Botanic Gardens, Kew, Richmond, TW9 3DS, United Kingdom

19  
20 <sup>5</sup>Departamento de Biología, Escuela Politécnica Nacional, Av. Ladrón de Guevara E11-  
21 253, Apartado: 17-01-2759, Quito, Ecuador.

22  
23 <sup>6</sup>Centro de Investigación de la Biodiversidad y Cambio Climático, Universidad  
24 Tecnológica Indoamérica, Quito EC170103, Ecuador

25  
26 <sup>7</sup>Department of Biological Sciences, St. John's University, 8000 Utopia Parkway,  
27 Queens, NY, 11439, USA

28  
29 <sup>8</sup>CREAF, Cerdanyola del Vallès, ES-08193 Catalunya, Spain

30  
31 <sup>9</sup>*present address*: Center for the Study of Social Insects, São Paulo State University  
32 (UNESP), Rio Claro-SP, 13506-900, Brazil.

33  
34 <sup>10</sup>*present address*: Department of Biological Sciences, University of Pernambuco, Rua  
35 Capitão Pedro Rodrigues, 105, 55290-000 Garanhuns, Pernambuco, Brazil

36  
37 <sup>11</sup>*present address*: Department of Zoology, University of Oxford, Oxford, OX 3PS, UK

38  
39 \*Correspondence: jonathan.shik@bio.ku.dk  
40  
41  
42

43 **ABSTRACT**

44 During crop domestication, human farmers traded greater productivity for higher crop  
45 vulnerability outside specialized cultivation conditions. We found a similar domestication  
46 tradeoff across the major co-evolutionary transitions in farming systems of attine ants.  
47 First, the fundamental nutritional niches (FNNs) of cultivars narrowed during ~ 60  
48 million years of naturally selected domestication, and laboratory experiments showed that  
49 ant farmers representing subsequent domestication stages strictly regulate protein harvest  
50 relative to cultivar FNNs. Second, ants with different farming systems differed in their  
51 abilities to harvest the resources that best matched the nutritional needs of their fungal  
52 cultivars. This was assessed by quantifying realized nutritional niches (RNNs) from  
53 analyses of items collected from the mandibles of laden ant foragers in the field. Third,  
54 extensive field collections suggest that among-colony genetic diversity of cultivars in  
55 small-scale farms may offer population-wide resilience benefits that species with large-  
56 scale farming colonies achieve by more elaborate and demanding cultivation practices of  
57 less diverse crops. Our results underscore that naturally selected farming systems have  
58 potential to shed light on nutritional tradeoffs that shaped the course of culturally evolved  
59 human farming.

## 60 INTRODUCTION

61 Farming evolved by natural selection in several social insect lineages and as a  
62 cultural innovation in our human ancestors. Although analogies across these domains  
63 need to be phrased carefully, it is reasonable to assume that both types of farming became  
64 more sophisticated over time which seems obvious from a crop perspective because our  
65 domesticated food plants often only remotely resemble their free-living ancestors. Over  
66 thousands of years, humans have continuously selected plant cultivars with nutritionally  
67 enhanced or enlarged leaves, roots, fruits and seeds<sup>1,2</sup> which facilitated modern  
68 agriculture's ecological expansion<sup>3</sup>. Domestication has also exposed production tradeoffs  
69 as increasingly specialized crops came to rely on specific abiotic conditions that farmers  
70 needed to provide to match their cultivars' shrinking fundamental niches for moisture,  
71 temperature and nutrients<sup>4</sup>. Such tradeoffs were especially pronounced when artificially  
72 selected traits ran counter to the naturally selected life histories that previously  
73 maximized fitness in wild cultivar ancestors confronting more hostile and fluctuating  
74 natural environments<sup>5,6</sup>. As domestication proceeded, culturally informed human farmers  
75 managed to push these tradeoffs to extremes. For instance, crops in historically recent  
76 eras have been farmed across huge metapopulations even though their vulnerability to  
77 herbivores and pathogens has often increased<sup>7</sup>, although the strength of such tradeoffs can  
78 vary<sup>8</sup>. This geographic expansion has been enhanced by technological solutions that  
79 enable farmers to provide their crops with consistently optimized realized niches (e.g.  
80 higher doses of pesticides).

81 The farming systems of ants<sup>9,10</sup>, macrotermite termites<sup>11</sup>, and ambrosia beetles<sup>10</sup>  
82 required millions of years of natural selection to produce narrow mutualistic co-

83 dependencies<sup>12</sup>. Rather than balancing nutritional needs by farming diverse foods as in  
84 large-scale human societies, these insect farmers specialized on clonal monocultures of a  
85 limited suite of fungi, like the fungus-farming attine ants that adopted agaricaceous  
86 fungal cultivars (Order: Agaricales)<sup>13,14</sup>. Despite this overall difference, ant farmers of  
87 diverse attine genera came to rear specialized cultivars with varying degrees of  
88 polyploidy<sup>15</sup> and basal metabolic rates<sup>16</sup> that appear linked to transitions to increasingly  
89 complex farming strategies. Throughout their shared co-evolutionary histories, fungus-  
90 farming ants also evolved specialized fungal enzyme recycling habits to increase crop  
91 productivity<sup>17</sup> and powerful antibiotic defences to control crop disease<sup>18-21</sup>. These  
92 fungicultural innovations have likely been instrumental in the ecological diversification  
93 and geographic expansion among the 19 extant attine genera encompassing > 240 species  
94 that now inhabit most neotropical rainforests and many drier habitats from Argentina to  
95 the northeastern United States<sup>22-25</sup>

96         Given these historical domestication patterns, we hypothesized that: 1) ant  
97 farmers have solved a tradeoff between crop yield and cultivar vulnerability, and 2) this  
98 tradeoff hinges upon nutrient availability as the core commodity of crop provisioning and  
99 yield. We used the well-established nutritional geometry approach to test this hypothesis,  
100 capitalizing on its conceptual and empirical tools for visualizing and modelling tradeoffs  
101 that organisms navigate to maintain nutritional homeostasis<sup>26-29</sup>. Our present study builds  
102 on an earlier one in which we used nutritional geometry to quantify a related type of  
103 nutritional tradeoff in the ant *Mycocepurus smithii*, a representative of the paleoattine  
104 clade that is sister to the neoattines that evolved more organizationally-complex farming  
105 systems. Colonies of *M. smithii* farm clones of a weakly domesticated fungal cultivar that

106 can maximize growth of edible hyphae in proportion to carbohydrate intake, but with the  
107 downside of encouraging cultivars to produce inedible mushrooms that may benefit  
108 fungal reproduction to the detriment of farmer fitness<sup>30</sup>. We now expand this nutritional  
109 geometry approach across the key evolutionary transitions towards higher organizational  
110 complexity in attine farming systems (Fig. 1a) by integrating their nutritional economy  
111 with ecological niche theory.

112         We tested the domestication tradeoff between yield and vulnerability by  
113 performing a study with the following three objectives: 1) To quantify and visualize the  
114 breadth of fundamental nutrient niches (FNNs)<sup>29,31</sup> when fungal cultivars are grown *in*  
115 *vitro* across artificial nutritional landscapes varying in absolute and relative abundance of  
116 protein and carbohydrate macronutrients (Fig. 1b). We used these results to interpret  
117 laboratory feeding experiments with nutritionally-defined substrates that tested how ant  
118 farmers prioritize macronutrient intake within the performance constraints imposed by  
119 their cultivar's FNNs. 2) To examine whether and how ant farmers target their cultivar's  
120 nutritional niches while foraging across natural nutritional landscapes in the field (Fig.  
121 1c). We quantified realized nutritional niches (RNNs)<sup>29,31</sup> of naturally collected substrates  
122 and assessed their nestedness within the experimentally obtained FNN dimensions of  
123 cultivars (Fig. 1d). 3) To assess whether individual small-scale farming colonies within  
124 local populations can successfully maintain genetically different cultivars while foraging  
125 within a single nutritional landscape. We assess whether small-scale farmers manage  
126 cultivar vulnerabilities differently than large-scale farmers that achieve system resilience  
127 by cultivating uniform crops of low genetic diversity.

128

129 **RESULTS**

130 ***Cultivar FNNs narrowed with full domestication but growth rates did not increase***

131 We first compared cultivar FNNs across three evolutionary stages of crop  
132 domestication<sup>14</sup> where we predicted that ant farmers solved the yield-vulnerability  
133 tradeoff in different ways. We started with the paleoattine genera *Mycocephurus* and  
134 *Apterostigma*, and the genus *Cyphomyrmex* that represents an early branch of the ‘lower’  
135 neoattines that would later produce the ‘higher’ neoattine genera described below (Fig.  
136 1A). These three representatives likely resemble the ancestral fungus-farming attines that  
137 arose ~ 60 MYA because all retained small, low-productivity colonies of tens to some  
138 hundreds of monomorphic workers and cultivars with undifferentiated hyphae to feed the  
139 ant farmers<sup>14</sup> (*but see* reference 32 for a case of a differentiated cultivar). As expected  
140 prior to full crop domestication, these cultivars exhibited broad FNNs with hyphal growth  
141 maximized across a wide range of imbalanced protein:carbohydrate (P:C) ratios (Fig. 2,  
142 Supplementary Table 1). These cultivar FNNs also varied widely among (but not within,  
143 Extended Data Fig. 1) cultivar types (Fig. 2), which likely reflects that these farmers have  
144 exchanged loosely domesticated fungi with free-living varieties over millions of years<sup>33</sup>.  
145 Despite this variation, the agaricaceous cultivars of *M. smithii* and *C. costatus*, as well as  
146 the pterulaceous ‘coral fungus’ *Myrmecopterula velohortorum* reared by *A. dentigerum*<sup>34</sup>  
147 all showed continuing or accelerated growth on carbohydrate-biased substrates at high  
148 nutritional concentrations (Fig. 2) relative to what can apparently be provided in detrital  
149 substrates collected by their ant farmers (*see* discussion of Figure 4a below).

150 We next measured the FNNs of cultivars in the higher-neoattine genera  
151 *Sericomyrmex*, *Mycetomoellerius* and *Paratrachymyrmex* (Fig. 1A) that evolved ~ 27-31

152 MYA<sup>21,22,35</sup>. The adaptive radiation of this clade coincided with the irreversible  
153 domestication of a specialized fungal lineage capable of concentrating nutrients in  
154 specialized swollen hyphal tips (gongylidia) bundled into staphylae<sup>36-38</sup>. The extant  
155 descendants of these higher neoattines represent an intermediate domestication phase  
156 characterized by farming fully domesticated cultivars but without significant changes in  
157 foraging substrates<sup>39,40</sup> and colony sizes<sup>16,41</sup> relative to the phylogenetically basal lower  
158 neoattines and paleoattines. Consistent with their domesticated crops being derived from  
159 a single fungal ancestor, the cultivars of these higher-neoattine farmers had less variable  
160 FNNs than the paleoattine and lower-neoattine cultivars, typically maximizing growth on  
161 substrates with 8-15 g/L protein and at rather low (8 to 25 g/L) absolute concentrations of  
162 carbohydrates (Fig. 2). The narrowed FNNs of these cultivars were confirmed because  
163 they often exhibited mortality in petri dish culture when they were provided with  
164 suboptimal nutrient mixtures (Extended Data Fig. 2, Supplementary Table 2), consistent  
165 with expectations of the hypothesized domestication tradeoff of specialization versus  
166 vulnerability.

167         The leafcutter ants represent the major expansion in scale and organizational  
168 complexity of neoattine fungus farming that appeared ~ 18.5 MYA<sup>21,22</sup> (Fig. 1a). The  
169 emergence of the leafcutter ants coincided with the use of freshly-cut vegetation from  
170 hundreds of plant species<sup>42</sup> to provision massive gardens of the now obligately  
171 polyploid<sup>15</sup> staphylae-bearing cultivar *Leucoagaricus gongylophorus*<sup>34</sup> to support tens of  
172 thousands of ants in colonies of *Acromyrmex* and up to millions of individuals in colonies  
173 of *Atta*<sup>24</sup>. The performance responses of *L. gongylophorus* cultivars isolated from  
174 colonies of *Acromyrmex echinatio*r and *Atta colombica* resembled those of the other

175 higher-neoattine cultivars (Fig. 2, Extended Data Fig. 1) and also often included steep  
176 mortality increases when confined to nutritional blends outside the cultivar's FNN  
177 (Extended Data Fig. 2). Thus, the protein-carbohydrate FNN of *L. gongylophorus*  
178 cultivars did not appear to have changed in connection to the broader ecological niche of  
179 substrate provisioning displayed by the leafcutter ants<sup>42</sup>.

180         The *L. gongylophorus* cultivar of leafcutter ants also exhibited statistically similar  
181 *in vitro* growth rates relative to cultivars of the other higher neoattines (Extended Data  
182 Fig. 3), which is remarkable because *Atta* foragers in the field provision cultivars with  
183 orders of magnitude more substrate mass than any other attine genera<sup>42</sup>. In fact, cultivars  
184 of the smallest lower-attine colonies of *A. dentigerum* and *C. costatus* grew faster *in vitro*  
185 on a standard potato dextrose agar (PDA) medium known to generally promote high  
186 cultivar growth rates<sup>25,43</sup> than all other cultivars tested (Extended Data Fig. 3). This result  
187 appears to be analogous to the insignificant changes in growth rate of plant crops during  
188 and after human domestication<sup>7</sup> and suggests that accelerated intrinsic cultivar growth  
189 rates neither drove nor responded to later increases in scale and organizational  
190 complexity of fungus-farming in neoattine ants.

191

### 192 ***Foraging ants collect nutrients within the FNN dimensions of their cultivars***

193         Ant and human farmers nutritionally provision their cultivars in fundamentally  
194 different ways. While humans farm mostly autotrophic plants that need targeted blends of  
195 inorganic NPK fertilizers, attine ant farmers maintain heterotrophic fungi that need  
196 organic inputs in the form of freshly cut plant material, or scavenged insect frass and  
197 other detritus that foragers must gather from their environment (Fig. 1). Fungal cultivars

198 also have no independent resource acquisition via roots, so the ants need to provide crops  
199 with most all their nutrients as crude forage<sup>44</sup>. These chemically and physically complex  
200 substrates contain the required protein and carbohydrate macronutrients but also water,  
201 micronutrients and vitamins, toxins and recalcitrant cell wall fibres<sup>37,45</sup>. We used  
202 nutritional geometry tools to cut through this complexity and explore whether and how *in*  
203 *vitro* measures of cultivar FNN breadth constrain the nutritional content of substrates  
204 collected by the fungus-farming ants when they forage.

205         We performed laboratory feeding experiments to measure nutrient-regulation  
206 strategies in attine colonies confined to single nutritionally-defined substrates with  
207 varying P:C ratios (Fig. 3a). For these experiments, we focused on comparing three attine  
208 species representing the main domestication transitions described above: paleoattines (*M.*  
209 *smithii*), higher neoattines (*Paratrachymyrmex cornetzi*), and crown-group neoattine  
210 leafcutters (*A. colombica*). We found that the leafcutter ants targeted broader protein  
211 dimensions than the two other species (Fig. 3b). Specifically, *A. colombica* foragers  
212 collected more of the protein-rich 1:1 P:C substrate than both *M. smithii* and *P. cornetzi*  
213 that tightly regulated protein intake to remain at low levels across all P:C substrates (Fig.  
214 3b, Extended Data Fig. 4, Supplementary Table 4). Colonies of *P. cornetzi* and *M. smithii*  
215 also had an aversion to substrates with macronutrient ratios above 1:3 P:C, possibly  
216 because these blends are associated with increased risks of crop failure (Extended Data  
217 Fig. 5), worker mortality (Supplementary Fig. 2a) and reduced total colony biomass  
218 (Supplementary Fig. 3a) for both *P. cornetzi* (Supplementary Table 6) and *M. smithii*<sup>30</sup>.  
219 The tight regulation of protein intake by foragers of these two species matched the

220 narrow *in vitro* protein FNNs of their cultivars (i.e., the steep declines in growth beyond  
221 15 g/L protein; Fig. 2).

222         The leafcutter ants did not face a similarly extreme vulnerability-yield tradeoff  
223 between starving fungal cultivars of carbohydrates or over-harvesting protein when  
224 confined to nutritionally imbalanced substrates (Fig. 3b, Extended Data Fig. 4,  
225 Supplementary Table 4). In fact, *A. colombica* colonies showed greater stability across all  
226 P:C substrates tested, remaining stable or increasing in both worker number  
227 (Supplementary Fig. 2b, Supplementary Table 6) and colony mass (Supplementary Fig.  
228 3b, Supplementary Table 6). Yet, *A. colombica* colonies also appeared to walk a  
229 nutritional tightrope since their foraging levels also varied with substrate macronutrient  
230 ratios. Specifically, *A. colombica* workers collected significantly less carbohydrates when  
231 confined to the most protein-biased substrates (3:1 and 6:1 P:C; Extended Data Fig. 4).  
232 Thus, while *A. colombica* colonies maintained relatively high foraging levels under these  
233 protein-biased conditions (Fig. 3b), they still experienced carbohydrate shortfall relative  
234 to levels that could maximize cultivar growth on carbohydrate-biased diets. This may be  
235 related to the steep *in vitro* decline in growth (Fig. 2) and survival (Extended Data Fig. 2)  
236 exhibited by *L. gongylophorus* beyond substrate protein concentrations of 30 g/L, that  
237 resembled (but was slightly higher than) the cultivars of *P. cornetzi* that we tested.  
238

### 239 ***Quantifying the RNN dimensions of foraging ant farmers in the field***

240         Targeting FNNs that maximize hyphal growth (Fig. 2) while avoiding inedible  
241 mushroom production is a documented nutrient-mediated tradeoff before attine ants fully  
242 domesticated their cultivars<sup>30</sup>. However, mushrooms were never produced by plated

243 cultivars of neotattine ants, so we expected them to exhibit more subtle domestication  
244 tradeoffs. Analogous to human crops that no longer exchange genes with wild or feral  
245 relatives, higher neotattine ant farmers are expected to nutritionally provision their  
246 cultivars to optimize resource allocation between somatic hyphal growth and the fraction  
247 of this growth to be harvested as edible staphylae with gongylidia. We therefore tested  
248 whether and how ant farmers in the field resolved nutrient provisioning vulnerabilities in  
249 their crops by estimating colony-level RNNs that could be mapped on the specific FNN  
250 dimensions obtained by rearing cultivars *in vitro*.

251         As before, we compared three attine species representing the transitions from  
252 collection of mostly frass—leaf material processed through the guts of insect herbivores  
253 (secondary herbivory) in *M. smithii* and *P. cornetzi* and freshly cut vegetation in *A.*  
254 *colombica* (primary herbivory). To assess RNNs, we collected, weighed, and catalogued  
255 samples of substrates collected from the mandibles of laden foragers as they returned to  
256 their nests, representing 77 hours of total field observations from 22 colonies of *M.*  
257 *smithii* (Supplementary Table 7), 44 colonies of *P. cornetzi* (Supplementary Table 7), and  
258 1 colony of *A. colombica* (Supplementary Table 8). We then nutritionally analysed  
259 substrates collected by foragers from the same study populations representing 19 colonies  
260 of *M. smithii*, 33 colonies of *P. cornetzi* (Supplementary Table 9), and 1 colony of *A.*  
261 *colombica* (Supplementary Table 8).

262         We expected colonies of each ant species to forage for natural substrates within  
263 the range of the experimentally obtained FNNs of their cultivars, i.e. that their RNNs  
264 would not exceed their cultivar's upper physiological tolerances and would tend to  
265 converge on the performance maximizing dimensions (dark red) of the nutritional FNNs

266 (Fig. 1c,d). We first found that foragers of *M. smithii* harvested sufficient amounts of  
267 protein (horizontal axis) to be able to avoid the cultivar FNN region that would produce  
268 inedible mushrooms, matching previously results<sup>30</sup>. However, while these detrital  
269 substrates provided a minimal protein level, they were also below the higher carbohydrate  
270 concentrations (vertical axis) that could accelerate cultivar growth (Fig. 4a). Second,  
271 colonies of *P. cornetzi* harvested substrates that yielded an RNN whose protein and  
272 carbohydrate amounts fell somewhat below the FNN regions maximizing both growth  
273 and staphyla density (Fig. 4b), while avoiding the high-concentration nutritional blends  
274 that induced cultivar mortality *in vitro* (Extended Data Fig. 2). Third, the representative  
275 *Atta* leafcutter ant colony achieved a comparatively broader RNN with higher maximal  
276 concentrations of macronutrients (Fig. 4c). This enlarged leafcutter RNN spanned the  
277 distinct FNN peaks for hyphal growth and staphyla production.

278

### 279 ***Cultivars with variable FNNs co-exist in a single nutritional provisioning environment***

280 Many attine ant species maintain mutualistic associations with several fungal  
281 haplotypes<sup>46-48</sup>. This trait also applies to human farming but is puzzling in attine farms  
282 that are constrained to rear a single cultivar clone which is usually vertically acquired<sup>12,49</sup>.  
283 For instance, if neighbouring colonies of the same ant species farm genetically distinct  
284 cultivars with different FNNs, one should expect considerable farmer plasticity to target  
285 RNNs that match their cultivar's specific needs. To assess the extent of performance  
286 variation across cultivars, we compared two attine species with different levels of fully  
287 domesticated crop specificity: 1) the leafcutter ant *A. colombica* as a single ant species  
288 (Extended Data Fig. 6) that farms a single species of *L. gongylophorus* cultivar (Fig. 5)

289 and 2) the higher neoattine *P. cornetzi* (Fig. 5, Extended Data Fig. 7, Extended Data Fig.  
290 8) known to farm one of several possible species of fungal cultivar across our study area  
291 in Panama<sup>48</sup>.

292 We found that *P. cornetzi* farms five cultivar haplotypes (Fig. 5) with similar  
293 FNNs for hyphal growth that also roughly matched those of the *L. gongylophorus* cultivar  
294 of *A. colombica* (Fig. 6a) and were consistent with other fully domesticated higher-  
295 neoattine cultivars tested (Fig. 2). However, the FNNs for staphyla density among  
296 isolates of the single cultivar species of *A. colombica* were much more consistent than the  
297 FNNs of the different cultivar haplotypes reared by *P. cornetzi* (Fig. 6a). In particular, the  
298 staphyla density FNN of cultivar haplotype 1 was similar to the FNN of *L. gongylophorus*  
299 reared by *A. colombica* and protein-biased (from 3:1 to 9:1 P:C) at intermediate nutrient  
300 concentrations (20 g/L) (Fig. 6a), while the staphyla-density FNN for *P. cornetzi*'s  
301 cultivar haplotype 3 was carbohydrate-biased (1:9 P:C) at high nutrient concentrations  
302 (40 to 60 g/L). This cultivar variation extended to intrinsic growth rates viewed across the  
303 distinct cultivar haplotypes farmed by *P. cornetzi* (Fig. 6b) although further sampling will  
304 be needed to confirm this apparent phylogenetic variation.

305 Variable FNNs for maximal staphylae density raise questions about the plasticity  
306 required by small-scale *P. cornetzi* farmers that found colonies with one specific crop  
307 symbiont and are unable to predict what mixture of forage items their habitat patch will  
308 provide over time. Such stochasticity may help explain why colonies primarily balance  
309 their foraging between two main detrital resources, frass (variable in protein and low in  
310 carbohydrates) and wood chips (variable in carbohydrates and low in protein), while also  
311 targeting nutritionally variable, but higher total concentration bits of plant material (e.g.

312 flowers, seeds, detrital leaf bits) (Fig. 4b, Supplementary Table 7). Further research will  
313 be required to resolve whether *P. cornetzi* colonies are actually able to dynamically adjust  
314 the cumulative sum of these alternative resources so they would achieve their optimal  
315 RNN under most circumstances.

316 The need for plasticity in cultivar provisioning also became apparent after we  
317 mapped spatial colony distributions across higher neoattine species within our 20-m<sup>2</sup>  
318 leaf litter plots. This showed that *P. cornetzi* is part of a diverse assemblage of similarly  
319 sized higher-neoattine (non-leafcutter) farming species (*Paratrachymyrmex*,  
320 *Mycetomoellerius*, and *Sericomyrmex*) in Soberanía Park (Extended Data Fig. 7,  
321 Extended Data Fig. 8), with colonies of 4-7 species (mean  $\pm$  SD:  $5 \pm 1$ ) co-existing in  
322 close sympatry and at high densities of 21 to 66 colonies (mean  $\pm$  SD:  $43 \pm 17$ ) per 20 m<sup>2</sup>  
323 plot. *P. cornetzi* was the most abundant of these farming ant species with  $20 \pm 13$  (range  
324 4-35) colonies per 20 m<sup>2</sup> (Fig. 6c, Extended Data Fig. 10). At a community scale, most of  
325 these fungus-farming ant species share at least some symbionts<sup>48</sup> so that neighbouring  
326 colonies separated by  $< 1$  m are within each other's foraging range. Different fungal  
327 haplotypes thus coexist within a single local nutritional environment.

328

## 329 **DISCUSSION**

### 330 ***Towards an ecological explanation of the history and current impact of ant farming***

331 Quantifying nutritional niches of fungus-farming ants reveals a domestication  
332 tradeoff between vulnerability and yield. First, weakly domesticated paleoattine and  
333 lower-neoattine cultivars tend to express generalist traits reminiscent of free-living fungi.  
334 These included broad and variable FNNs that likely matched ecological conditions found

335 in nutritional landscapes available outside regulated nest environments. For instance,  
336 their high growth rates at high-nutrient concentrations appear to exceed nutrients  
337 available from ant detritus provisioning, and may instead reflect an ability to exploit  
338 ephemeral pulses of leaf-litter resources<sup>50</sup>. Second, fully domesticated cultivars of higher-  
339 neoattine and leafcutter ants expressed similarly narrowed FNNs, likely reflecting that  
340 they are derived from a single fully domesticated *Leucoagaricus* ancestor. The trends  
341 observed across attine species generally matched expectations of a domestication tradeoff  
342 where increased productivity through specialized nutritional yield benefits (i.e., staphylae  
343 with gongyliidia) in a narrowed suite of environmental conditions coincided with an  
344 increasingly remote resemblance to the cultivar traits that characterized the ancestral free-  
345 living fungi.

346 Our results revealed only minor expansions in FNN breadth of the specialized *L.*  
347 *gongylophorus* cultivar of leafcutter ants compared to the genetically more variable but  
348 likewise fully domesticated cultivars of *P. cornetzi*. This result seemed surprising at first  
349 because leafcutter ants are remarkably generalist foragers that collect fresh leaves and  
350 flowers from hundreds of plant species<sup>42</sup>. However, the broad RNN dimensions of *Atta*  
351 colonies that we recorded in the field (Fig. 4c) were consistent with higher degrees of  
352 symbiotic resilience when we exposed colonies to nutritionally imbalanced substrates in  
353 lab experiments (Fig. 3). Moreover, leafcutter ant workers have evolved many specialized  
354 traits enhancing farming performance<sup>18-20,51-53</sup>. Specialized gardener castes may thus help  
355 to secondarily adjust the harvested substrates relative to required RNN dimensions when  
356 they macerate leaf fragments into a leaf-pulp mixed with glandular and faecal fluids<sup>17-19</sup>.  
357 A suite of bacterial symbionts was also recently discovered in leafcutter ant digestive

358 tracts<sup>54</sup> and fungus gardens<sup>55</sup> that may help to maintain the wide RNN of *Atta* leafcutter  
359 ants by providing nitrogen fixation services<sup>55</sup>, by recycling excess amino acids such as  
360 arginine, or by converting plant-sap citrate into acetate to directly fuel ant metabolism<sup>40</sup>.

361

### 362 ***A range of nutritional solutions to a fundamental domestication tradeoff***

363 Our study focused on three representative attine ant model species varying in  
364 colony size and scale of farming to explore: 1) the unique cultivation challenges and  
365 opportunities each symbiosis faces, and 2) how each farming system maximizes cultivar  
366 performance by mixing nutritionally variable substrates collected from the environment.  
367 Our results highlight that there is no single answer to how fungus-farming ants navigate  
368 the complex nutritional landscape of a tropical rainforest to target RNNs within the FNNs  
369 of their specific cultivars to maximize edible yield. Since the cultivars of paleoattines  
370 deliver undifferentiated hyphae as food, *M. smithii* ants need only to mix in sufficient  
371 protein via frass provisioning to avoid wasteful mushroom production, which is  
372 achievable but likely precludes rapid garden growth<sup>30</sup>. The two more complex farming  
373 systems have a differentiated cultivar so that maximizing crop growth (hyphal mass) and  
374 edible harvest (staphyla density) have become separate optimization challenges. In the  
375 higher-neoattine *P. cornetzi*, opportunities for maximizing yield appear to be somewhat  
376 constrained by the nutritional FNN targets that may vary across cultivar haplotypes (Fig.  
377 6a). Some farmers may thus maximize edible yield by collecting more flower petals to  
378 acquire extra carbohydrates, while others may benefit from the additional protein  
379 available in frass. To detect such dynamic nutritional provisioning strategies, longitudinal  
380 sampling of individual colonies will be useful for describing how RNNs change over

381 time, and broader sampling of colonies across habitat types will help determine the  
382 resiliency of RNNs among colonies with access to different suites of nutritionally  
383 variable substrates.

384 Our results further showed that the detrital substrates collected by paleoattine and  
385 non-leafcutter neoattine ants tend to be nutrient-poor resources, so that farming  
386 productivity is likely to be ecologically constrained because their cultivar FNNs peak at  
387 higher nutrient concentrations than the ants can usually provide. The expanded RNN of  
388 *Atta* leafcutter ants provisions a highly specialized cultivar and appears to have partially  
389 overcome such constraints by both being able to handle more diverse macronutrient  
390 mixtures and higher absolute macronutrient quantities. The *Atta* cultivar also appeared to  
391 be special because it offers the farming ants two distinct macronutrient performance  
392 targets – one for hyphal growth (a carbohydrate-biased FNN) and one for staphyla  
393 density (a protein-biased FNN). Thus, our results suggest that the RNN expansion  
394 achieved by *Atta* leafcutter ants may have allowed a novel form of farming flexibility that  
395 could capitalize on high-quality (protein-rich) substrates when available without  
396 simultaneously enhancing cultivar growth rate *per se*.

397 We expect that future work will elaborate on the ecological pathways of fungal  
398 cultivar domestication by focusing on nutritional aspects of: 1) the earliest domestication  
399 tradeoff at the origin of attine fungus farming between the yield of incipient crops and the  
400 costs of abandoning hunter-gathering<sup>16,56</sup>, 2) the diversification of farming practices  
401 across habitats and biomes<sup>57</sup> as attine lineages diverged and adapted to very different  
402 plant substrates<sup>14,22</sup>, and 3) the challenges of maintaining farming homeostasis in  
403 response to stochastic environmental variation<sup>25</sup>, particularly in large-scale farming

404 systems that evolved quite differently in *Acromyrmex* and *Atta* leafcutter ants. Further  
405 progress in answering questions of this kind is feasible because the attine ants are the  
406 only insect farming clade where many farming systems varying widely in scale and  
407 organizational complexity co-exist within the same ecosystems. This contrasts with the  
408 large-scale agricultural systems of humans that have tended to competitively displace  
409 subsistence farms in many habitats<sup>58</sup>, with the fungus-farming termites where only more  
410 organizationally-complex forms of agriculture exist<sup>11</sup>, and with the ambrosia beetles that  
411 invariably remained small-family cooperative breeders in tunnels that colony foundresses  
412 need to excavate in host trees<sup>59</sup>.

413 **METHODS**

414 *Study populations*

415 All attine colonies were harvested from the lowland tropical rainforest at  
416 Soberanía National Park, Panama (N 9.13528, W 79.72141) from 28 October 2013 to 10  
417 June 2015, with additional fieldwork conducted from 1 May to 30 June 2019. We located  
418 nest entrances of paleoattines, lower neoattines and higher neoattines under leaf litter by  
419 placing polenta bait on the ground and following laden workers back to their nests.  
420 Leafcutter nests were visible as large established colonies and small dirt mounds of  
421 recently founded colonies. Back in the lab at the Smithsonian Tropical Research Institute  
422 in Gamboa, we established colonies (see Supplementary Table 5 for demography) in  
423 plastic containers with *ad lib* water and ground polenta (or leaves for *Atta*), and  
424 acclimated them to lab conditions at 24°C. Ant vouchers were stored at the Smithsonian  
425 Tropical Research Institute (STRI) in Panama and fungal vouchers used in barcoding  
426 analyses were stored at Kew Gardens in the UK.

427

428 *Estimating ant cultivar FNNs and their growth performance in vitro*

429 We isolated fungus from 51 colonies of 10 attine species, including *Mycocepurus*  
430 *smithii* (10 colonies), *Apterostigma dentigerum* (3), *Cyphomyrmex rimosus* (1), *C.*  
431 *costatus* (3), *C. longiscapus* (1), *Paratrachymyrmex cornetzi* (13), *Mycetomoellerius*  
432 *zeteki* (2), *Sericomyrmex amabilis* (2), *Acromyrmex echinator* (3), and *Atta colombica*  
433 (13) (Supplementary Table 3). We grew these fungal cultivars on sealed sterile petri  
434 dishes containing PDA media (potato dextrose agar, DIFCO) and used them to generate  
435 pure fungal stock cultures. We then used PDA as a standard medium to compare cultivar  
436 growth rates<sup>25,43</sup> placing 19.6 mm<sup>2</sup> cylindrical plugs of pure culture into separate 60 x 15

437 mm petri dishes containing 15 ml of PDA. To estimate these intrinsic growth rates, we  
438 photographed the plates every 10 days for periods of 25 to 76 days (depending on the  
439 overall growth rate of the cultivar; Supplementary Table 3), and then used ImageJ (NIH  
440 Image; V 1.49) to measure cumulative fungal growth after 30 days (area mm<sup>2</sup>).

441 We visualized the fundamental nutritional niches (FNNs) of cultivar fungi  
442 isolated from nests of 10 attine species (N = 22 colonies), including *M. smithii* (1), *A.*  
443 *dentigerum* (3), *C. rimosus* (1), *C. costatus* (3), *C. longiscapus* (1), *P. cornetzi* (3), *M.*  
444 *zeteki* (2), *S. amabilis* (2), *A. echinator* (3), and *A. colombica* (3) (Supplementary Table  
445 3). We inoculated fungi in petri dishes containing 12 mL of 36 sterile synthetic agar-  
446 based ‘substrate’ treatments varying in P:C (1:9, 1:6, 1:3, 1:2, 1:1, 2:1, 3:1, 6:1, 9:1) and  
447 P + C concentration (8g/L, 20g/L, 40g/L, 60g/L) (n = 8 plates per substrate x dilution  
448 treatment, N = 288 plates per colony, N = 6,336 plates). We used growth rates on  
449 standard PDA media to set the length of P:C growth experiments for each cultivar  
450 (Supplementary Table 3).

451 Nutritionally-defined media that were used to quantify cultivars FNNs included  
452 distilled water and bacteriological agar (1.6% w:v; Amresco, Inc.), carbohydrates as  
453 equal parts sucrose (Doradita ® cane sugar) and starch (puriss p.a., from potato, reag.  
454 ISO, Sigma-Aldrich), and protein as equal parts bacto-peptone (enzymatic digest of  
455 protein, Becton, Dickinson and Company), trypticase peptone (pancreatic digest of  
456 casein, BD), and bacto tryptone (pancreatic digest of casein, BD). We also included a  
457 crushed multivitamin mixture (Centrum ®) at a concentration of 2% of the mass of  
458 protein + carbohydrates. See reference 30 for recipes. These ingredients were mixed with

459 200 ml distilled water on a stirring plate for 5 minutes and then sterilized by autoclaving  
460 at 121°C, yielding a pH of 6.9.

461 We mapped fungal growth areas across the 36-substrate arrays using the *fields*  
462 package<sup>61</sup> in R (3.2.4)<sup>62</sup>. We plotted nutritional landscape contours using non-parametric  
463 thin-plate splines and set the topological resolution of response surfaces to  $\lambda = 0.001$  as a  
464 smoothing parameter. We then used least-square regressions to assess the underlying  
465 significance of both linear and quadratic terms (and their interactions) and to verify the  
466 interpretation of FNN heatmaps based on fungal growth areas across the 36 protein and  
467 carbohydrate substrate combinations. Unless otherwise noted, we present heatmaps based  
468 on means of cultivars farmed by each attine species, but also provide all additional  
469 heatmaps for individual cultivars sampled from each attine colony in Figure 6A (*P.*  
470 *cornetzi* and *A. colombica*) and in the Extended Data Fig. 1 (all other colonies), as well as  
471 the underlying statistical analyses (Supplementary Table 1, Supplementary Table 10).

472 Each of the higher-neoattine and leafcutter cultivar haplotypes that we tested  
473 exhibited mortality on a subset of P:C substrates outside the range of the FNN. Mortality  
474 was indicated by inoculation plugs being clear (empty of fungus) or by lack of growth  
475 from the inoculation plug onto the P:C substrate by the end of the experiment (Extended  
476 Data Fig. 2). We generated heatmaps as described before ( $\lambda = 0.01$  for the percentage  
477 data) with survivorship averaged across cultivars isolated from nests of a given attine  
478 species, standardizing the heatmap colour scale from 0% to 100% survival. We  
479 statistically assessed the significance of underlying regressions as described for the  
480 growth area heatmaps. This type of mortality response was never observed in the  
481 paleoattine or lower-neoattine cultivars tested.

482

483 ***Laboratory no-choice feeding experiments with ant colonies and their natural cultivars***

484 We analysed nutrient regulation strategies of lab-acclimated colonies using  
485 nutritional geometry feeding experiments based on agar-based mixtures of protein and  
486 carbohydrate macronutrients (1.6g agar/L) at P:C ratios of 1:6, 1:3, 1:1, 3:1, and 6:1, and  
487 protein + carbohydrates dilutions of 100 g/L (for *P. cornetzi* and *A. colombica*)  
488 (Supplementary Table 11) or 20 g/L (for *M. smithii*)<sup>30</sup>. To evaluate evolutionary trends  
489 across a broader dataset than the one collected for the present study, we included some  
490 previously published data from a parallel study we performed on *M. smithii*<sup>30</sup>. In our no-  
491 choice experiments we confined colonies to a single P:C substrate for 29 days (*M.*  
492 *smithii*), 39 days (*P. cornetzi*), or 15 days (*A. colombica*). We provide raw pre- and post-  
493 experiment colony demography data for colonies in Supplementary Table 5. For  
494 standardized comparison across these experiments with different attine species, intake  
495 data were analysed for 15 days across all colonies.

496 We calculated protein and carbohydrate use by the ants from the agar P:C ratios  
497 and substrate dry mass loss estimated from dry:wet mass ratios of control agar fragments  
498 for each P:C ratio<sup>66</sup>. We weighed colonies on the first (initial colony mass) and last day  
499 (final colony mass) of the experiments, and calculated worker mortality rates by  
500 collecting all dead workers each time we changed substrates. During the *M. smithii* and *P.*  
501 *cornetzi* experiments, a number of colonies showed signs of crop failure (*i.e.*, elevated  
502 worker mortality, living workers ceasing foraging and discarding fungus garden  
503 material). For our demographic analyses, we assessed crops as having failed on the day  
504 colonies no longer had any fungus garden biomass left.

505 We performed statistical analyses in R 3.2.4<sup>62</sup>, exploring P:C substrate treatment  
506 effects on colony behaviour and performance separately for the three ant species (the *M.*  
507 *smithii* data were already similarly analysed in reference 30). We first used general linear  
508 models to test for substrate treatment effects on the response variables total  
509 macronutrients intake, protein intake, and carbohydrate intake with initial worker number  
510 as a covariate. These data were log-transformed prior to analyses to meet the assumptions  
511 of normality and homoscedasticity. For both *A. colombica* and *P. cornetzi*, we analysed  
512 cumulative substrate use over 15 days. For *P. cornetzi*, we also analysed macronutrient  
513 use ‘per day’, with number of days ranging from the full 39 days to any earlier day a  
514 specific colony was removed due to crop failure. We next performed a survival analysis  
515 (survfit, ggsvplot) in R using a Cox proportional hazards model to test for substrate  
516 treatment effects on crop failure in *P. cornetzi*, including initial fungus garden mass as a  
517 covariate. This analysis was not used for *A. colombica* because no colonies exhibited crop  
518 failure. To further explore the links between foraging behaviour and colony demography  
519 in both *P. cornetzi* and *A. colombica*, we used general linear models to test for effects of  
520 P:C macronutrient mixes on changes in worker number and colony mass, analysing the  
521 slopes of linear regressions for each colony from values on day 1 to values on the final  
522 day of the experiment (day 39 in *P. cornetzi*, day 15 *A. colombica*) or on the day the  
523 colony was removed from the experiment due to crop failure (a subset of *P. cornetzi*  
524 colonies).

525

526 *RNNs of fungal cultivars estimated from field-collected substrates*

527 To quantify RNNs for *P. cornetzi* and *M. smithii*, we first focused on cataloguing  
528 substrates harvested per unit time (Supplementary Table 7) and then on sampling the  
529 specific substrate categories to obtain sufficient biomass for nutritional analyses  
530 (Supplementary Table 9). We collected substrate items carried by laden workers  
531 returning to nests of 44 mapped *P. cornetzi* colonies from 8AM to 4PM during ~1-hour  
532 observation periods (43.7 total hours) (Supplementary Table 7). We collected substrates  
533 from 22 colonies of *M. smithii* using the same protocols, with sampling observations  
534 spanning over 20 hours (Supplementary Table 7, also see reference 30). Part of the raw  
535 biomass data of collected substrates for *M. smithii* and *P. cornetzi* were previously  
536 published<sup>40</sup> but were here explicitly analysed as RNNs and compared to fungal FNNs  
537 (Fig. 4a). These collections occurred during the wet season, a period of high attine  
538 activity, from Nov 16 to Dec 31 in 2013 and Jan 2 to Jan 17 in 2014 (Supplementary  
539 Table 7). During observation periods, we lay on trash bags next to nest entrances using a  
540 headlamp to maximize visibility and carefully grabbed laden foragers with forceps just  
541 before they disappeared underground. We collected these substrate pieces in Eppendorf  
542 tubes and then dried them at 60°C for 24 hours before classifying them into six  
543 categories: insect frass, leaf fragment, wood, flower, seed, and other (miscellaneous plant,  
544 insect piece, unknown), and weighing them to the nearest 1 µg on a Sartorius CP2P  
545 microbalance. We then calculated the fraction of biomass of each substrate type relative  
546 to the total biomass of substrate collected.

547 For nutritional analyses, we first homogenized individual frass samples collected  
548 from 33 colonies of *P. cornetzi* and 19 colonies of *M. smithii* located in the same leaf  
549 litter habitats as described above (Supplementary Table 9). We then determined their

550 elemental carbon (%C) and nitrogen (%N) in the lab of Ben Turner (Smithsonian  
551 Tropical Research Institute, Panama) with a Flash EA1112 analyser (CE Elantech, New  
552 Jersey, USA). The small size of these substrate fragments precluded direct macronutrient-  
553 level analyses of their percent protein and carbohydrates. To acquire these macronutrient  
554 estimates, we used 30 1-m<sup>2</sup> tarps placed on leaf litter for 24-hour intervals in the same  
555 forest in June 2019 to collect a large pooled sample of frass (and other detrital substrates)  
556 that fell from the canopy. Back in Copenhagen, we created a 1 g pooled frass sample of  
557 each substrate type (previously freeze dried just after collection in Panama using a SP  
558 Scientific BenchTop Pro with Omnitronics for 24 hours). We then homogenized these  
559 and analysed them for %C (Eurovector CN analyser coupled to an Isoprime isotope ratio  
560 mass spectrometer in the lab of Anders Michelsen, University of Copenhagen), as well as  
561 % total non-structural carbohydrates (%TNC: water soluble carbohydrates + starch). For  
562 %TNC measurements, we analysed 25 mg of frass with a Total Carbohydrate Assay Kit  
563 (Sigma-Aldrich) to determine water soluble carbohydrates, and 50 mg of frass with a  
564 Total Starch Assay Kit (Megazyme)<sup>67</sup> to quantify starch. We used 5 subsampled  
565 replicates for each quantification to calculate a conversion factor (2.59%) to substitute  
566 %C with %TNC for the frass and tiny wood fragments in the existing dataset  
567 (Supplementary Table 9).

568 From the elemental N frass data, we estimated crude protein (i.e. including non-  
569 available protein bound up by tannins) by multiplying substrate N mass by the standard  
570 conversion factor 6.25, following the Felton et al. protocol<sup>67</sup>. We then analysed %TNC in  
571 leaf-litter samples of seeds (n = 5) and a flower (n = 1), as well as pooled samples of  
572 flowers (n = 5) and tiny wood bits (n = 1) (Supplementary Table 9). In these non-frass

573 samples, we also quantified proteins using a CBQCA Protein Quantification Kit  
574 (Molecular Probes) (Supplementary Table 9).

575         The much larger quantities of fresh vegetative substrates harvested by leafcutter  
576 ants required different field collection methods and also facilitated more detailed NIRS  
577 nutritional analyses that were not possible for the small amounts of tiny substrate  
578 particles harvested from mandibles of paleoattine and higher-neoattine (non-leafcutter)  
579 ants. In May 2019, we located an *A. colombica* colony at La Laguna (W 9.11672, N -  
580 79.69514) and laid down on trash bags next to the most active trail, close to the nest  
581 entrance. We hand collected a total of 6,868 fragments (40,026 mg dry mass) carried by  
582 laden foragers during 3, 1.5 hour periods by two observers (9 total collection hours)  
583 between 9AM and 12AM (Supplementary Table 8). These fragments (almost all fresh  
584 plant material) were stored in new Ziplock bags every 30 minutes that were promptly  
585 transferred to a cooler. Back in the lab, we catalogued the forage fragment samples under  
586 a microscope based on the morphology of the veins of leaf fragments, weighed the  
587 specific fractions and freeze-dried for 24 hours as described above.

588         We then stored the samples at -20°C in new Ziploc bags with silica gel until  
589 subsequent nutrient and barcoding analyses. We extracted DNA from these samples and  
590 identified them using the Internal Transcribed Spacer 1 (ITS1) genetic marker (details in  
591 Supplementary methods; NCBI GenBank Accession codes provided in Supplementary  
592 Table 8). We then homogenized a subset of dried plant substrates into powder and used  
593 near infrared reflectance spectroscopy (NIRS)<sup>67</sup> to estimate the concentrations of total  
594 nitrogen, total protein, total non-structural carbohydrate, and starch in all 17 plant  
595 substrates sampled (details provided in Supplementary methods, Supplementary Table

596 12). The pie charts of harvested forage categories reported in Fig. 4 represent relative  
597 contributions to the total dry mass of substrate items collected by foraging workers  
598 (Supplementary Table 7, Supplementary Table 8). Based on barcoding analyses, we  
599 pooled a subset of the 17 samples that were identified as coming from the same plant  
600 species which gave a total of 13 plant species in the dataset (Supplementary Table 8). We  
601 converted the *in vitro* landscapes used to estimate cultivar FNNs from g/L to percent  
602 protein and carbohydrates based on P:C diet recipes to directly compare these data with  
603 RNNs based on macronutrient content in the natural forage obtained in the field. For  
604 simplicity, only the extreme carbohydrate-biased (1:9 P:C) and protein-based (9:1 P:C)  
605 nutritional rails are plotted in Fig. 4 to delineate the range of *in vitro* P:C conditions in  
606 which cultivar FNNs were measured.

607

#### 608 ***Obtaining accurate estimates of species diversity of sympatric farming systems***

609 The single species status of *A. colombica* and its fungal cultivar *L. gongylophorus*  
610 in central Panama are not in question and the same applies for *M. smithii* except that this  
611 paleoattine ant is known to be associated with a series of fungal cultivars<sup>68</sup>, some of  
612 which were also recognized in our previous study of the nutritional geometry of this  
613 species<sup>30</sup>. However, this is much more ambiguous for *P. cornetzi* and their fungal  
614 symbionts, because a previous small-scale study<sup>48</sup> showed that there is substantial cryptic  
615 diversity not only among the cultivars, but possibly also among the ants<sup>48</sup>. To avoid  
616 misinterpretations of our present results due to overlooking (and unjustifiably pooling)  
617 cryptic species of ants and cultivars, we did an extensive survey across a large meta-  
618 population *P. cornetzi* (54 km<sup>2</sup> of Soberanía Park forest) and within the local populations

619 represented by our six 20-m<sup>2</sup> plots (Extended Data Fig. 10). Within these study plots, we  
620 mapped all higher-neoattine nests (*Paratrachymyrmex*, *Mycetomoellerius*, *Sericomyrmex*)  
621 by dispersing oat-polenta bait in the leaf litter and following laden workers back to their  
622 nests (N = 263 total higher-neoattine nests across all plots, n = 27 ± 3 searching hours per  
623 plot), and then collected 2-4 workers from each nest in vials with 95% ethanol. We also  
624 sampled workers from 85 additional *Paratrachymyrmex* and *Mycetomoellerius* colonies  
625 distributed across Soberanía Park yielding 297 colonies sampled across these two genera.  
626 We similarly sampled 30 *A. colombica* colonies distributed across the Gamboa area.

627 We then identified the ants using morphological analysis of vouchers, as well as  
628 analysis of DNA microsatellite markers and mtDNA barcoding (details for each of these  
629 analyses provided in Supplementary methods). Briefly, we used DNA microsatellite  
630 analyses to examine the population structure of workers from 28 *A. colombica* colonies  
631 (Extended Data Fig. 6) and 297 *Paratrachymyrmex* and *Mycetomoellerius* colonies  
632 (Extended Data Fig. 7) using a total of nine variable markers (Ant7680, Ant859,  
633 Ant1343, Ant2341, Ant3993, Ant11400, Ant3653, Ant4155, Ant8498)<sup>69</sup>. To supplement  
634 these nuclear microsatellite data, we also used DNA barcoding analyses of a  
635 mitochondrial marker (~1100 bp of the Cytochrome Oxidase 1 gene), which included ant  
636 workers from 147 *Paratrachymyrmex*, 38 *Mycetomoellerius* colonies and 29 *A.*  
637 *colombica* colonies (Supplementary Table 13). A subset of these ants was paired with  
638 their fungal cultivars to examine farmer-cultivar association patterns, as we excavated 69  
639 colonies of putative *P. cornetzi* and 29 colonies of *A. colombica* (Supplementary Table  
640 14). For each of these colonies, we collected a clean 2-cm<sup>3</sup> fraction of fungus garden with  
641 forceps into vials with 95% ethanol soon after excavation. We identified these samples by

642 analysing mitochondrial DNA (~820 bp of the nuclear large subunit rRNA (LSU) and  
643 [~550 bp of the Internal Transcribed Spacer unit (ITS). We used these sequences (plus  
644 sequences from an additional *M. smithii* ant and fungal sample) to generate phylogenetic  
645 trees (details about tree generation are provided in Supplementary methods).

646         The resulting sequence information and supporting dataset for the 215 ant  
647 specimens is named DS-ATTINENG “Nutritional niches reveal fundamental  
648 domestication tradeoffs in fungus-farming ants” and has been deposited in the Barcode of  
649 Life Data System<sup>70</sup> (BOLD; [dx.doi.org/10.5883/DS-ATTINENG](https://dx.doi.org/10.5883/DS-ATTINENG)). The sequence data is  
650 also deposited in GenBank (Accession codes provided in Supplementary Table 13). The  
651 sequence information for 99 fungal cultivar samples in Fig. 5 has been deposited in  
652 GenBank (Accession codes for LSU and ITS provided in Supplementary Table 14). To  
653 further interpret our molecular evidence confirming known species and detecting cryptic  
654 species among the collected *P. cornetzi*-like colonies within Soberanía Park, we also  
655 explored the biogeographic distribution of this species complex by comparing our  
656 sequence data to sequences of 8 Costa Rican and 1 Ecuadorian *P. cornetzi* ant specimen  
657 available from BOLD<sup>81</sup> (Extended Data Fig. 8).

658

### 659 ***Cultivar diversity across colonies, in vitro FNN variation and growth performance***

660         We visualized the FNNs of isolated cultivars in Fig. 6A from three colonies of *P.*  
661 *cornetzi* (177605, 177611, 177612) and three colonies of *A. colombica* (177627, 177628,  
662 177626) (Supplementary Table 3). Cultivars were selected prior to determining their  
663 haplotype identity, after which the *P. cornetzi* cultivars used in the experiment were  
664 identified as haplotype 1 (177605, 177612) and haplotype 3 (177612) (Fig. 6a). Due to

665 general differences in growth rate among cultivar haplotypes on standard PDA medium,  
666 these haplotypes were grown on P:C substrates for different lengths of time  
667 (Supplementary Table 3). A trained researcher (MF) also estimated staphyla production  
668 rate (staphyla number per number of growth days) by visually scanning the high-  
669 resolution photos and validating counts by opening a subset of plates and directly  
670 counting staphyla (clusters of gongylidia) under a dissecting microscope at 40X  
671 magnification.

672 **ACKNOWLEDGEMENTS**

673 The Smithsonian Tropical Research Institute (STRI) provided support and access to  
674 facilities in Gamboa. The Autoridad Nacional del Ambiente y el Mar (ANAM) gave  
675 permission to the JJB and JZS lab groups to sample attine ants in Panama and export  
676 them to Denmark. We thank students of the Tropical Behavioural Ecology and Evolution  
677 field course in 2015 for additional assistance collecting colonies. JZS was supported by a  
678 Postdoctoral Fellowship via a Smithsonian Institution Competitive Grant to WTW, JJB  
679 and JZS, an EU Marie Skłodowska-Curie International Incoming Fellowship (327940),  
680 the Centre for Social Evolution at the University of Copenhagen (Danish National  
681 Research Foundation: DNRF57), an ERC Advanced grant (ANTS: 323085) to JJB, and  
682 an ERC Starting Grant to JZS (ELEVATE: 757810). JCS was supported by SJU start-up  
683 funds and NSF-DEB # 2016372.

684

685 **AUTHOR CONTRIBUTIONS**

686 J.Z.S., J.J.B., and W.T.W. conceived and designed the study. J.Z.S., E.B.G., M.F., A.J.J.C.,  
687 D.A.D., and P.W.K. performed field work, collected colonies, isolated fungal cultivars,  
688 and performed *in vitro* experiments with fungal cultivars. D.A.D., J.Z.S., P.W.K., and  
689 A.J.J.C. extracted DNA and performed DNA barcoding analyses for ants and fungal  
690 cultivars. P.W.K., J.Z.S. and J.H. performed the microsatellite analyses. X.A. and J.Z.S.  
691 performed statistical analyses. J.C.S. performed phylogenetic analyses. J.Z.S. and J.J.B.  
692 wrote the initial draft of the manuscript. J.Z.S., P.W.K., D.A.D., J.C.S., A.J.J.C., X.A.,  
693 J.H., W.T.W., and J.J.B. contributed to the interpretation of the data and to the editing of  
694 subsequent drafts of the manuscript.

695 **COMPETING INTERESTS**

696 The authors declare no competing interests

697

698 **DATA AVAILABILITY**

699 The DNA sequences generated during this study are available at NCBI GenBank. This  
700 includes the data for: the 215 ant specimens (Accession codes for COI sequences  
701 provided in Supplementary Table 13), the 99 fungal cultivar samples (Accession codes  
702 for LSU and ITS sequences provided in Supplementary Table 14), and plant sample ITS1  
703 sequences harvested by colonies of *A. colombica* (Accession codes provided in  
704 Supplementary Table 8). Ant sequence datasets and supporting information are also  
705 deposited in the Barcode of Life Data System<sup>70</sup> (BOLD) under the project titled DS-  
706 ATTINENG “Nutritional niches reveal fundamental domestication tradeoffs in fungus-  
707 farming ants” ([dx.doi.org/10.5883/DS-ATTINENG](https://dx.doi.org/10.5883/DS-ATTINENG)).

708

709 **CODE AVAILABILITY**

710 There is no custom code generated for this study. Sequence alignment matrices and  
711 newick files used to generate phylogenies in Fig. 5 (both fungal and ant trees) and  
712 Extended Data Fig. 8 are available as text files in Supplementary Data 1-3. All code used  
713 to generate results will be made available upon request.

714 **REFERENCES**

- 715 1. Piperno, D. & Pearsall, D. M. *The Origins of Agriculture in the Lowland Neotropics*  
716 (Academic Press, 1998).
- 717 2. Newell-McGloughlin, M. Nutritionally improved agricultural crops. *Plant. Physiol.*  
718 **147**, 939-953 (2008).
- 719 3. Green, R. E., Cornell, S. J., Scharlemann, J. P. W. & Balmford, A. Farming and the  
720 fate of wild nature. *Science* **307**, 550-555 (2005).
- 721 4. Meyer, R. S., DuVal, A. E. & Jensen, H. R. Patterns and processes in crop  
722 domestication: an historical review and quantitative analysis of 203 global food crops.  
723 *New Phytol.* **196**, 29-48 (2012).
- 724 5. Purugganan, M. D. & Fuller, D. Q. The nature of selection during plant  
725 domestication. *Nature* **457**, 843-848 (2009).
- 726 6. Milla, R., Osborne, C. P., Turcotte, M. M. & Violle, C. Plant domestication through  
727 an ecological lens. *Trends Ecol. Evol.* **30**, 463-469 (2015).
- 728 7. Evans L. T. *Crop evolution, adaptation and yield*. (Cambridge University Press,  
729 1993)
- 730 8. Turcotte, M. M., Turley, N. E. & Johnson, M. T. J. The impact of domestication on  
731 resistance to two generalist herbivores across 29 independent domestication events.  
732 *New Phytol.* **204**, 671-681 (2014).
- 733 9. Chomicki, G. & Renner, S. S. Farming by ants remodels nutrient uptake in epiphytes.  
734 *New Phytol.* **223**, 2011-2023 (2019).

- 735 10. Mueller, U. G., Gerardo, N. M., Aanen, D. K., Six, D. L. & Schultz, T. R. The  
736 evolution of agriculture in insects. *Ann. Rev. Ecol., Evol. Systemat.* **36**, 563-595  
737 (2005).
- 738 11. Aanen, D. K. et al. The evolution of fungus-growing termites and their mutualistic  
739 fungal symbionts. *Proc. Natl. Acad. Sci. USA* **99**, 14887-14892 (2002).
- 740 12. Mehdiabadi, N. J. & Schultz, T. R. Natural history of the fungus-farming ants  
741 (Hymenoptera: Formicidae: Myrmicinae: Attini). *Myrmec. News* **13**, 37-55 (2009).
- 742 13. Mueller, U. G., Scott, J. J., Ishak, H. D., Cooper, M. & Rodrigues, A. Monoculture of  
743 leafcutter ant gardens. *PLoS One* **9**, e12668 (2010).
- 744 14. Schultz, T. R. & Brady, S. G. Major evolutionary transitions in ant agriculture. *Proc.*  
745 *Natl. Acad. Sci. USA* **105**, 5435-5440 (2008).
- 746 15. Kooij, P. W., Aanen, D. K., Schiøtt, M. & Boomsma, J. J. Evolutionary advanced ant  
747 farmers rear polyploid fungal crops. *J. Evol. Biol.* **28**, 1911-1924 (2015).
- 748 16. Shik, J. Z. et al. Metabolism and the rise of fungus cultivation by ants. *Am. Nat.* **184**,  
749 364-373 (2014).
- 750 17. de Fine Licht, H. H. et al. Laccase detoxification mediates the nutritional alliance  
751 between leaf-cutting ants and fungus-garden symbionts. *Proc. Natl. Acad. Sci. USA*  
752 **110**, 583-587 (2012).
- 753 18. Fernández-Marín, H. et al. Functional role of phenylacetic acid from metapleural  
754 gland secretions in controlling fungal pathogens in evolutionarily derived leaf-cutting  
755 ants. *Proc. R. Soc. B* **282**, 20150212 (2015).
- 756 19. Fernández-Marín, H. et al. T. Dynamic disease management in *Trachymyrmex*  
757 fungus-growing ants (Attini: Formicidae). *Am. Nat.* **181**, 571-582 (2013).

- 758 20. Currie, C. R., Mueller, U. G. & Malloch, D. The agricultural pathology of ant fungus  
759 gardens. *Proc. Natl. Acad. Sci. USA* **96**, 7998-8002 (1999).
- 760 21. Nygaard, S. et al. Reciprocal genomic evolution in the ant-fungus agricultural  
761 symbiosis. *Nat. Comm.* **7**, 12233 (2016).
- 762 22. Branstetter, M. G. et al. Dry habitats were crucibles of domestication in the evolution  
763 of agriculture in ants. *Proc. R. Soc. B* **284**, 20170095 (2017).
- 764 23. Li, H. et al. Convergent evolution of complex structures for ant–bacterial defensive  
765 symbiosis in fungus-farming ants. *Proc. Natl. Acad. Sci. USA* **115**, 10720-10725  
766 (2018).
- 767 24. Hölldobler, B. & Wilson, E. O. *The Leafcutter Ants: Civilization by Instinct* (W. W.  
768 Norton and Company, 2010).
- 769 25. Mueller, U. G. et al. Evolution of cold-tolerant fungal symbionts permits winter  
770 fungiculture by leafcutter ants at the northern frontier of a tropical ant–fungus  
771 symbiosis. *Proc. Natl. Acad. Sci. USA* **108**, 4053-4056 (2011).
- 772 26. Simpson, S. J. & Raubenheimer, D. *The Nature of Nutrition: A Unifying Framework*  
773 *from Animal Adaptation to Human Obesity*. (Princeton Univ. Press, 2012).
- 774 27. Raubenheimer, D. Toward a quantitative nutritional ecology: the right-angled mixture  
775 triangle. *Ecol. Monogr.* **81**, 407-427 (2011).
- 776 28. Sperfeld, E. et al. Bridging ecological stoichiometry and nutritional geometry with  
777 homeostasis concepts and integrative models of organism nutrition. *Funct. Ecol.* **31**,  
778 286-296 (2017).
- 779 29. Shik, J. Z. & Dussutour A. Nutritional dimensions of invasive success. *Trends Ecol.*  
780 *Evol.* **2676**, X-X (2020).

- 781 30. Shik, J. Z. et al. Nutrition mediates the expression of cultivar-farmer conflict in a  
782 fungus-growing ant. *Proc. Natl. Acad. Sci. USA* **113**, 10121-10126 (2016).
- 783 31. Machovsky-Capuska, G. E., Senior, A. M., Simpson, S. J., & Raubenheimer, D. The  
784 multidimensional nutritional niche. *Trends Ecol. Evol.* **31**, 355-365 (2016).
- 785 32. Masiulionis, V. E. et al. A Brazilian Population of the Asexual Fungus-Growing Ant  
786 *Mycocepurus smithii* (Formicidae, Myrmicinae, Attini) Cultivates Fungal Symbionts  
787 with Gongyldia-Like Structures. *PLoS One* **9**, e103800 (2014).
- 788 33. Vo, T. L., Mikheyev, A. S. & Mueller, U. G. Free-living fungal symbionts  
789 (Lepiotaceae) of fungus-growing ants (Attini: Formicidae). *Mycologia* **101**, 206–210  
790 (2009).
- 791 34. Schultz, T. R. et al. The most relictual fungus-farming ant species cultivates the most  
792 recently evolved and highly domesticated fungal symbionts species. *Am. Nat.* **185**,  
793 693-703 (2015).
- 794 35. Solomon, S. E. et al. The molecular phylogenetics of *Trachymyrmex* Forel ants and  
795 their fungal cultivars provide insights into the origin and coevolutionary history of  
796 ‘higher-attine’ ant agriculture. *Systemat. Entomol.* **44**, 939-956 (2019).
- 797 36. Quinlan, R. J. & Cherrett, J. M. The role of fungus in the diet of the leaf-cutting ant  
798 *Atta cephalotes* (L.) *Ecol. Entomol.* **4**, 151-160 (1979).
- 799 37. Schiøtt, M., de Fin Licht, H. H., Lange, L. & Boomsma, J. J. Towards a molecular  
800 understanding of symbiont function: identification of a fungal gene for the  
801 degradation of xylan in the fungus gardens of leaf cutting ants. *BMC Biol.* **8**, 40  
802 (2008).

- 803 38. de Fine Licht, H. H., Boomsma, J. J. & Tunlid, A. Symbiotic adaptations in the fungal  
804 cultivar of leaf-cutting ants. *Nat. Comm.* **5**, 5675 (2014).
- 805 39. de Fine Licht, H. H. & Boomsma, J. J. Forage collection, substrate preparation, and  
806 diet composition in fungus-growing ants. *Ecol. Ent.* **35**, 259-269 (2010).
- 807 40. Sapountzis, P., Zhukova, M., Shik, J. Z., Schiøtt, M. & Boomsma, J. J.  
808 Reconstructing the symbiotic functions of intestinal Mollicutes in fungus-growing  
809 ants. *eLife* **7**, e39209 (2018).
- 810 41. Seal, J. N. & Tschinkel, W. R. Colony productivity of the fungus-gardening ant  
811 *Trachymyrmex septentrionalis* (Hymenoptera: Formicidae) in a Florida pine forest.  
812 *Ann. Ent. Soc. Amer.* **99**, 673-682 (2006).
- 813 42. Wirth, R., Beyschlag, W., Ryel, R. J. & Hölldobler, B. Annual foraging of the leaf-  
814 cutting ant *Atta colombica* in a semideciduous rain forest in Panama. *J. Trop. Ecol.*  
815 **13**, 741-757 (1997).
- 816 43. Cazin Jr., J., Wiemer, D. F. & Howard, J. J. Isolation, growth characteristics, and  
817 long-term storage of fungi cultivated by attine ants. *Appl. Environ. Microbiol.* **55**,  
818 1346–1350 (1989).
- 819 44. Mueller, U. G., Schultz, T. R., Currie, C. R., Adams, R. M. M. & Malloch, D. The  
820 origin of the attine ant-fungus mutualism. *Quart. Rev. Biol.* **76**, 169-197 (2001).
- 821 45. de Fine Licht, H. H., Schiøtt, M., Mueller, U. G. & Boomsma, J. J. Evolutionary  
822 transitions in enzyme activity of ant fungus gardens. *Evolution* **64**, 2055-2069 (2010).
- 823 46. Chapela, I. H., Rehner, S. A., Schultz, T. R. & Mueller, U. G. Evolutionary history of  
824 the symbiosis between fungus-growing ants and their fungi. *Science* **266**, 1691-1694  
825 (1994).

- 826 47. Mikheyev, A. S., Mueller, U. G. & Boomsma, J. J. Population genetic signatures of  
827 diffuse co-evolution between leaf-cutting ants and their cultivar fungi. *Molec. Ecol.*  
828 **16**, 209-216 (2007).
- 829 48. de Fine Licht, H. H. & Boomsma, J. J. Variable interaction specificity and symbionts  
830 performance in Panamanian *Trachymyrmex* and *Sericomyrmex* fungus-growing ants.  
831 *BMC Evol. Biol.* **14**, 244 (2014).
- 832 49. Howe, J., Schiøtt, M. & Boomsma, J. J. Horizontal partner exchange does not  
833 preclude stable mutualism in fungus-growing ants. *Behav. Ecol.* **30**, 372-382 (2018).
- 834 50. Cornejo, F. H., Varela, A. & Wright, S. J. Tropical forest litter decomposition under  
835 seasonal drought: nutrient release, fungi and bacteria. *Oikos* **70**, 183-190 (1994).
- 836 51. Wilson, E. O. Caste and division of labor in leaf-cutter ants (Hymenoptera:  
837 Formicidae: *Atta*) II. The ergonomic optimization of leaf cutting. *Behav. Ecol.*  
838 *Sociobiol.* **7**, 157-165 (1980).
- 839 52. Roces, F. & Hölldobler, B. Use of stridulation in foraging leaf-cutting ants:  
840 mechanical support during cutting or short-range recruitment signal? *Behav. Ecol.*  
841 *Sociobiol.* **39**, 293-299 (1996).
- 842 53. Kleineidam, C., Romani, R., Tautz, J. & Isidoro, N. Ultrastructure and physiology of  
843 the CO<sub>2</sub> sensitive sensillum ampullaceum in the leaf-cutting ant *Atta sexdens*. *Arthr.*  
844 *Struct. Devel.* **29**, 43-55 (2000).
- 845 54. Sapountzis, P., Nash, D. R., Schiøtt, M. & Boomsma, J. J. The evolution of  
846 abdominal microbiomes in fungus-growing ants. *Molec. Ecol.* **28**, 879-899 (2019).
- 847 55. Pinto-Tomás, A. A. et al. Symbiotic nitrogen fixation in the fungus gardens of leaf-  
848 cutter ants. *Science* **326**, 1120-1123 (2009).

- 849 56. Mummert, A. E., Esche, E., Robinson, J. & Armelagos, G. J. Stature and robusticity  
850 during the agricultural transition: evidence from the bioarchaeological record. *Econ.*  
851 *Human Biol.* **9**, 284-301 (2011).
- 852 57. Fuller, D. Q. et al. The domestication process and domestication rate in rice: spikelet  
853 bases from the Lower Yangtze. *Science* **323**, 1607–1610 (2009).
- 854 58. Sauer, C. O. *Agricultural Origins and Dispersals* (American Geographical Society,  
855 1952).
- 856 59. Nuotclà, J. A., Biedermann, P. H. W. & Taborsky, M. Pathogen defence is a potential  
857 driver of social evolution in ambrosia beetles. *Proc. R. Soc. B*, **286**, 20192332 (2019).
- 858 60. Dussutour, A., Latty, T., Beekman, M. & Simpson, S. J. Amoeboid organism solves  
859 complex nutritional challenges. *Proc. Natl. Acad. Sci. USA* **107**, 4607-4611 (2010).
- 860 61. Nychka, D., Furrer, R. & Sain, S. Fields: Tools for Spatial Data. R package version  
861 8.2-1. Available at <https://cran.r-project.org/web/packages/fields/index.html> (2015).
- 862 62. R CORE TEAM. R: A Language and Environment for Statistical Computing.  
863 Available <https://www.R-project.org/> (2018).
- 864 63. Dussutour, A. & Simpson, S. J. Description of a simple synthetic diet for studying  
865 nutritional responses in ants. *Insects. Soc.* **55**, 329-333 (2008).
- 866 64. Dussutour, A. & Simpson, S. J. Communal nutrition in ants. *Curr. Biol.* **19**, 740-744  
867 (2009).
- 868 65. Warbrick-Smith, J., Raubenheimer, D., Simpson, S. J. & Behmer, S. T. Three  
869 hundred and fifty generations of extreme food specialisation: testing predictions of  
870 nutritional ecology. *Ent. Exp. Applic.* **132**, 65-75 (2009).

- 871 66. Kay, A. D., Shik, J. Z., Van Alst, A., Miller, K. A., Kaspari, M. Diet composition  
872 does not affect ant colony tempo. *Funct. Ecol.*, **26**, 317-323 (2011).
- 873 67. Felton, A. M. et al. Nutritional ecology of *Ateles chamek* in lowland Bolivia: How  
874 macronutrient balancing influences food choices. *Int. J. Primatol.* **30**, 675-696  
875 (2009).
- 876 68. Kellner, K., Fernández-Marín, H., Ishak, H. D., Linksvayer, T. A. & Mueller, U. G.  
877 Co-evolutionary patterns and diversification of ant-fungus associations in the asexual  
878 fungus-farming ant *Mycocepurus smithii* in Panama. *J. Evol. Biol.* **26**, 1353-1362  
879 (2013).
- 880 69. Butler, I. A., Siletti, K., Oxley, P. R. & Kronauer, D. J. C. Conserved microsatellites  
881 in ants enable population genetic and colony pedigree studies across a wide range of  
882 species. *PLoS One* **9**, e107334 (2014).
- 883 70. Ratnasingham, S. & Hebert, P. D. N. BOLD: The Barcode of Life Data System  
884 ([www.barcodinglife.org](http://www.barcodinglife.org)). *Molec. Ecol. Notes* **7**, 355-364 (2007).
- 885

886 **FIGURE LEGENDS**

887 **FIGURE 1 Assessing domestication tradeoffs across transitions in mutualistic**  
888 **farming practice of the fungus-farming ants. a)** Attine ants harvest nutritionally  
889 variable detritus fragments in phylogenetically basal genera to freshly cut vegetation in  
890 the leafcutter ant crown group (schematic phylogeny<sup>21-23,34,35</sup> restricted to ten  
891 representative species co-occurring in Panama). The ants convert this crude forage into  
892 nutritional substrates for their fungal cultivars. Farming transitions included: Paleoattines  
893 (*Mycocepurus*, *Apterostigma*) that have retained small colonies of 10-100 workers and  
894 include a clade within *Apterostigma* secondarily adopting a coral fungus cultivar;  
895 Neoattines (*Cyphomyrmex*) that cultivate fungi in a hyphal form and a lineage of yeast  
896 farmers; Higher Neoattines (*Sericomyrmex*, *Mycetomoellerius*, *Paratrachymyrmex*) that  
897 farm a fully domesticated single lineage of gongylidia-bearing cultivars; Leafcutters  
898 including *Acromyrmex* with up to 1,000's of workers per colony, and *Atta* with up to  
899 millions of morphologically more specialized workers per colony. **b) Defining**  
900 *fundamental nutritional niches*<sup>29,31</sup> (FNNs) of fungal cultivars by quantifying their  
901 intrinsic tolerances and nutrient requirements after isolating cultivars from ant colonies  
902 and rearing them *in vitro* across substrates varying in concentrations and ratios of protein  
903 and carbohydrates. **c) Defining realized nutritional niches**<sup>29,31</sup> (RNNs) as the  
904 macronutrients that ant farmers offer to their cultivars by sampling and nutritionally  
905 analysing substrates collected from the mandibles of returning attine foragers in the field.  
906 **d) Assessing nutrient-provisioning strategies of farmers** by overlaying substrate RNNs  
907 atop cultivar FNNs. Foraging ants can maximize cultivar performance with a substrate  
908 RNN nested inside their cultivar's FNN. In the example provided, the FNN (red bullseye)  
909 is derived from a nutritional landscape of 36 diets (small grey circles) spanning nine P:C  
910 ratios from 1:9 (carbohydrate biased) to 9:1 (protein biased) along which P:C ratios  
911 remain constant (grey lines extending from the origin), but total protein and  
912 carbohydrates concentrations increase. The RNN (green polygon) is bounded by  
913 nutritional blends of substrates collected by foraging ants in the field, shown here as  
914 coloured circles matching substrates in the panel c pie chart. In panel a, ant specimen  
915 images are © antweb.org and attine foraging images are © Alex Wild. Where possible,  
916 we used pictures of the same species represented in this study, but the *Mycetomoellerius*  
917 picture represents the closely related *M. tucumanus*<sup>35</sup>, the *Apterostigma* picture is of an  
918 unknown species of that genus, and the *Atta* picture is of *A. cephalotes*. Photograph in  
919 panel c is by Sean Mattson.

920 **FIGURE 2 Heatmaps of cultivar growth indicate that domestication by higher-**  
921 **neoattine ants coincided with narrower fungal FNNs than in lower-attine cultivars.**  
922 **This revealed a domestication tradeoff when combined with cultivar mortality data**  
923 **(shown in Extended Data Fig. 2).** Bottom row - *Diverse lower-attine cultivars with*  
924 *variable FNNs.* Top row – *Fully domesticated higher-neoattine cultivars with narrowed*  
925 *FNNs,* showing maximal growth (red heatmap colour) on similar blends of protein and  
926 carbohydrates. Whereas dark blue heatmap values in the lower-attine plots indicate  
927 slower cultivar growth, dark blue values in the upper higher neoattine and leafcutter plots  
928 often indicated cultivar mortality (*see* Extended Data Fig. 2). In each of the ten plots,  
929 cultivar growth rates (hyphal area, mm<sup>2</sup>) are visualized across 36 experimentally defined  
930 artificial media varying in absolute (g/L) amounts and relative (P:C ratio) amounts of  
931 protein and carbohydrates<sup>30,60</sup>. As illustrated in Fig. 1D, these substrates spanned nine  
932 P:C ratios (1:9, 1:6, 1:3, 1:2, 1:1, 2:1, 3:1, 6:1, 9:1) and four protein plus carbohydrate  
933 concentrations (8, 20, 40, 60 g/L). We used the *fields* package<sup>61</sup> in R<sup>62</sup> to visualize the  
934 response surfaces obtained from nonparametric thin-plate splines. Heatmaps are averages  
935 of cultivars from three colonies (*Apterostigma dentigerum*, *Cyphomyrmex costatus*, *P.*  
936 *cornetzi*, *Acromyrmex echinator*, *A. colombica*) or two colonies (*Mycetomoellerius*  
937 *zeteki*, *Sericomyrmex amabilis*), or were the values from one colony (*M. smithii*,  
938 *Cyphomyrmex rimosus*, *C. longiscapus*). FNNs for cultivar growth were consistent when  
939 measured across multiple colonies of an attine species, so we assume that the three single  
940 colony estimates are representative (Extended Data Fig. 1). We used least-square  
941 regressions to assess the underlying significance of linear and quadratic terms (and the  
942 linear interaction) across the 36 protein and carbohydrate substrate combinations and to  
943 validate the interpretation of FNN heatmaps of the dependent variables growth area  
944 (Supplementary Table 1) and percent survival (Supplementary Table 2). All response  
945 surface regressions producing the heatmap colour-gradients shown here were statistically  
946 significant (Supplementary Table 1).  
947

948 **FIGURE 3 Laboratory experiments with colonies of three representative attine ant**  
949 **species showed that farming ants conformed to their cultivars' FNN dimensions**  
950 **when allowed to forage on single nutritionally-defined substrates. a)** A representative  
951 image of an experimental colony of *P. cornetzi*. **b)** Intake levels show that colonies of *M.*  
952 *smithii* and *P. cornetzi* tightly regulated protein concentrations to remain at low levels  
953 while allowing carbohydrate levels to fluctuate widely across substrates. In contrast,  
954 colonies of *A. colombica* allowed for greater fluctuations of protein intake while  
955 sustaining high carbohydrate levels, even when restricted to the 1:1 P:C nutritional rail  
956 that *M. smithii* and *P. cornetzi* avoided (*see also* Extended Data Fig. 4). These no-choice  
957 feeding experiments were performed on lab-acclimated colonies confined to single  
958 nutritional rails of agar-based substrates<sup>63</sup> (Supplementary Table 5) that constrained their  
959 intake to specific P:C ratios (1:6, 1:3, 1:1, 3:1, 6:1)<sup>26,30,64</sup>. Dashed lines reflect decisions  
960 to over- or under-collect one macronutrient to avoid imbalanced intake of another  
961 limiting macronutrient. Mean substrate harvest values  $\pm$  SE are presented coincident to  
962 intake rails<sup>65</sup>. We analysed cumulative intake rates over 15 days of feeding on these  
963 substrates (Supplementary Table 4). We present 15-day data for *P. cornetzi* and *M.*  
964 *smithii* to facilitate direct comparison with the *A. colombica* results, but the *P. cornetzi*  
965 experiment continued for 39 days, yielding long-term results that were consistent with the  
966 15-day data (Supplementary Fig. 1, Supplementary Table 4). The *M. smithii* experiment  
967 extended for 29 days and also yielded consistent results<sup>30</sup>. Colonies of *M. smithii* and *P.*  
968 *cornetzi* with collapsed fungus gardens were removed from the experiment on the day  
969 they had no remaining cultivar biomass left (Extended Data Fig. 5). In panel b, ant  
970 specimen images are © antweb.org

971 **FIGURE 4 Fungus-farming ants representing different stages of cultivar**  
972 **domestication and organizational complexity navigate nutritional landscapes to**  
973 **harvest RNNs relative to their cultivar's FNNs.** Lying down on the forest floor next to  
974 nest entrances, we collected and catalogued tiny substrate bits from the mandibles of  
975 laden workers returning to their colonies. Individual substrate types are represented by  
976 coloured wedges in pie charts corresponding to coloured circles comprising RNN maps  
977 (see Fig. 1C,D). **a)** The paleoattine ant *M. smithii* collected 52% insect frass, 29% wood  
978 pieces, 14% small seeds, 2% flowers, and 3% other undefined bits of detritus in terms of  
979 the proportion of sampled substrate biomass<sup>30</sup> (Supplementary Table 7). This yielded an  
980 RNN enabling the ants to provision cultivars with percent total non-structural  
981 carbohydrates (%TNC) ranging from 0.6 to 25.8% while also enabling protein  
982 percentages ranging from 0.1 to 27.0% (Supplementary Table 9). **b)** The higher-neoattine  
983 ant *P. cornetzi* rears a fully domesticated gongylidia-bearing cultivar, but colonies  
984 continue to forage like lower-attine ants, collecting detritus (60.1% insect frass, 10.7%  
985 detrital leaf fragments, 9.2% seeds, 3.5% flower pieces, 3.3% wood bits, 13.1% other,  
986 Supplementary Table 7). These substrates yielded a RNN with similar dimensions as  
987 harvested by *M. smithii*, with %TNC ranging from 0.6 to 25.8%, and crude protein  
988 ranging from 0.1 to 30.84% (Supplementary Table 9). **c)** The leafcutter ant *A. colombica*  
989 has a different foraging strategy of primary herbivory focusing on fresh leaves that  
990 foragers cut from many plant species (here represented by 13 different vegetative  
991 substrates harvested by a single colony, Supplementary Table 8). These substrates range  
992 widely in both %TNC (from 7.47 to 33.76%) and in crude protein (from 4.64 to 34.83%)  
993 (Supplementary Table 8). Capturing the macronutrient range of different substrate types  
994 collected by *M. smithii* and *P. cornetzi* required larger pooled samples of similar  
995 substrates collected from leaf litter traps and flower petal samples collected from the  
996 mandibles of *A. colombica* that resembled those foraged by *M. smithii* and *P. cornetzi*.  
997 Details about sampling and nutritional analyses are provided in the methods and  
998 Supplementary Tables 7-9. The slightly larger white areas in the low-concentration  
999 nutrient space (lower-left corners) of these fungal FNN heatmaps relative to their  
1000 representations in Fig. 2 resulted from the conversion of macronutrient units of g/L to  
1001 percent of dry diet mass within *in vitro* diet recipes. This facilitated comparison of  
1002 cultivar FNNs and substrate RNNs and did not impact the interpretation of the results.  
1003 The regressions underlying variation in growth area and staphylo density across P:C  
1004 substrates for fungal FNN plots were significant (Supplementary Table 1, Supplementary  
1005 Table 10).

1006 **FIGURE 5 Comparing two fully domesticated farming systems with different**  
1007 **operational scales and ecological impacts.** Within Soberanía National Park, the higher-  
1008 neoattine ant *P. cornetzi* cultivated five haplotypes of related *Leucoagaricus* fungi with  
1009 bootstrap support indicating that each of the five coloured cultivar clades represents a  
1010 separate fungal haplotype. Each of these fungal haplotypes was also widely distributed  
1011 across Soberanía Park (Extended Data Fig. 9) and likely across the neotropics<sup>35</sup>. In  
1012 contrast, the leafcutter ant *A. colombica* farms only a single cultivar species *L.*  
1013 *gongylophorus*. The bootstrap majority consensus barcoding tree based on the  
1014 Cytochrome Oxidase 1 (CO1) gene for the ants indicated that Panamanian *P. cornetzi*  
1015 includes a morphologically cryptic species (*hereafter P. ADG3274*) that farms an  
1016 overlapping diversity encompassing at least three of the five *P. cornetzi* cultivars. The  
1017 status of *P. ADG3274* as a distinct cryptic species was supported by microsatellite  
1018 analyses (Extended Data Fig. 7) and additional barcoding showing that while *P.*  
1019 *ADG3274* is relatively uncommon in our study site it has a regional distribution  
1020 extending at least to Costa Rica (Extended Data Fig. 8). We do not further consider *P.*  
1021 *ADG3274* in the present study. The bootstrap majority consensus barcoding tree for ants  
1022 is based on a ~1100 bp section of the CO1 gene and includes 130 colonies of *P. cornetzi*,  
1023 15 colonies of *P. ADG3274*, and 29 colonies of *A. colombica* (Supplementary Table 13).  
1024 The bootstrap majority consensus barcoding tree for fungi is based on a ~820 bp section  
1025 of the LSU gene and a ~550 bp section of the ITS marker, and includes samples from 69  
1026 colonies of *P. cornetzi* and *P. ADG3274* (fungal haplotype 1 [n = 26], 2 [n = 4], 3 [n =  
1027 13], 4 [n = 5], 5 [n = 21]) and 29 colonies of *A. colombica* (Supplementary Table 14).  
1028 Both ant and cultivar trees were rooted with a *M. smithii* ant or fungus sample. Ant  
1029 images are © Antweb.org.

1030 **FIGURE 6. Colonies of *P. cornetzi* co-exist within nutritional foraging environments**  
1031 **despite farming different cultivar haplotypes with variable properties of nutrient**  
1032 **yield and growth. a)** Fungal cultivars of *P. cornetzi* and *A. colombica* exhibited similar  
1033 FNNs for maximal hyphal growth (top row). However, *P. cornetzi* (fungal haplotypes 1  
1034 and 3) had more variable FNNs for staphyla density (bottom row) while differences  
1035 across these performance variables were much less variable for the *L. gongylophorus*  
1036 cultivars of *A. colombica* towards the right. All response surface regressions for fungal  
1037 growth area (Supplementary Table 1) and staphyla density (Supplementary Table 10)  
1038 were significant, supporting the interpretation of their response contours. **b)** Testing three  
1039 of the cultivar haplotypes isolated from 13 *P. cornetzi* colonies against *L. gongylophorus*  
1040 cultivars from 12 *Atta* colonies showed that the former grows slower and with greater  
1041 variation (means + SE of mycelial area per day of growth) on *in vitro* standardized PDA  
1042 media than the *A. colombica* cultivars ( $F_{1,22} = 12.76$ ,  $p = 0.002$ ). Visual inspection of the  
1043 growth results plotted on the phylogeny, further suggest strain-specific variation in mean  
1044 cultivar growth rate, although additional sampling will be needed to provide sufficient  
1045 statistical power for phylogenetically explicit analyses. **c)** *P. cornetzi* ants farm diverse  
1046 fungal cultivar haplotypes in sympatry, with four of the five identified cultivars co-  
1047 occurring within a single 20 m<sup>2</sup> monitoring plot within Soberanía park (Extended Data  
1048 Fig. 9). Thus, colonies separated by < 1 m often farmed different cultivar haplotypes. Our  
1049 estimates of local cultivar richness in this example plot are conservative as we sequenced  
1050 fungal cultivars from a subset of 18 (larger purple semi-circles) of the 35 *P. cornetzi*  
1051 colonies that we mapped over 23 searching hours. The small light-purple circles indicate  
1052 *P. cornetzi* colonies where only the ants were genotyped) and the stylized tree symbols  
1053 indicate trees > 1 m dbh.  
1054

## Extended Data Figure Legends

**Extended Data Figure 1 Cultivars exhibited consistent FNNs for hyphal growth when isolated from different colonies of each attine species**, supporting that the heatmaps based on species means (Fig. 2, Fig. 4B,C) accurately represent each cultivar's FNN. Additional details about how these heatmaps were generated and how they were interpreted are provided in Fig. 2. Heatmaps are provided here for attine species where multiple colonies were sampled. Additional colony-level heatmaps for *P. cornetzi* and *A. colombica* are provided in Fig. 6A. Collection IDs corresponding to experiment IDs were: Ae\_3 [177625], Ae\_2 [177624], Ae\_1 [177609], Tz\_30 [177632], Tz\_32 [177634], Sa\_23 [177623], Sa\_21 [177614], Ad\_5 [177629], Ad\_6 [177630], Ad\_7 [177631], Cc\_12 [37861], Cc\_13 [37862], and Cc14 [37864] (Supplementary Table 3). As in Figure 2, least-square regressions showed that each of the response surface regressions producing the heatmap colour-gradients was significant (Supplementary Table 1).

**Extended Data Figure 2 Cultivars of the higher-neoattine genera *Sericomyrmex*, *Mycetomoellerius*, *Paratrachymyrmex* (purple) and the leafcutter genera *Acromyrmex* (light green) and *Atta* (dark green) often exhibited mortality when confined to media with macronutrient mixtures outside their FNNs**. As shown in the figure legend, dark red indicates 100% survival and dark blue indicates 0% survival (a clear inoculation plug or a failure to colonize the agar plate from the inoculation plug). Percent mortality data were averaged across cultivars isolated from colonies of the same attine species (n = 2, *M. zeteki*; n = 3, *P. cornetzi*; n = 1, *S. amabilis*; n = 3, *A. echinator*; n = 3, *A. colombica*, Supplementary Table 2). Mortality data were not recorded for the *S. amabilis* colony Sa\_23. Small white areas on some of the plots indicate 100% survival.

**Extended Data Figure 3 The emergence of large-scale fungus farming by *Atta* leafcutter ants cannot be explained solely by an increase in intrinsic cultivar growth rate**. Coloured bars (corresponding to the farming transitions in the schematic tree of Fig. 1a) represent *in vitro* growth rates of cultivars during 30 days on a standard PDA medium. Vertical lines separate the major farming transitions that distinguish ten attine species sympatrically inhabiting the Panamanian rainforest of Soberanía park (see caption Fig. 1A). ANOVA showed that attine species farmed cultivars with significantly different growth rates after 30-days ( $F_{7,41} = 24.42$ ,  $p < 0.0001$ ) and letters indicate pairwise differences that were significant at  $P < 0.05$  in post-hoc Tukey tests. Numbers inside bars indicate numbers of colonies from which cultivars were isolated and used to calculate mean growth rates (+ SE) (Supplementary Table 3). The cultivars of *C. rimosus* and *C. longiscapus* were not included in the statistical analyses because we had no replicate colonies for these two ant species.

**Extended Data Figure 4 Cumulative amounts of protein and carbohydrates collected over 15 days by whole colonies of a) *P. cornetzi* and b) *A. colombica***. While colonies of *P. cornetzi* and *M. smithii*<sup>30</sup> avoided protein-biased substrates with

1101 ratios above 1:3 P:C, colonies of *A. colombica* collected large amounts of the 1:1 P:C  
1102 agar that *P. cornetzi* and *M. smithii* avoided. Colonies of *A. colombica* further  
1103 collected statistically similar levels of carbohydrates on 1:1, 1:3, and 1:6 P:C  
1104 substrates (Supplementary Table 4). This tolerance of higher protein levels enables *A.*  
1105 *colombica* to sustain higher carbohydrate intake levels on 1:1 P:C diets. Letters  
1106 indicate pairwise differences that were significant at  $P < 0.05$  (post-hoc Tukey tests)  
1107 (Supplementary Table 4). These intake plots (means + SE) provide an alternative  
1108 single-nutrient presentation of the bi-variate intake data presented in Fig. 3b.  
1109

1110 **Extended Data Figure 5 Colonies of *P. cornetzi* increasingly faced crop failure**  
1111 **when confined to P:C macronutrient mixtures with excess protein relative to**  
1112 **carbohydrates ( $\chi^2_4 = 17.9$ ;  $p = 0.001$ ).** While similar results were observed for *M.*  
1113 *smithii*<sup>30</sup>, no colonies of *A. colombica* experienced crop failure, even when confined  
1114 to the same protein-biased substrates. Cultivar survival probabilities were estimated  
1115 with a Cox proportional hazards model where substrate treatment was the explanatory  
1116 variable, initial garden mass was a covariate, and days remaining in the feeding  
1117 experiment was the response variable.  
1118

1119 **Extended Data Figure 6 STRUCTURE analyses of 9 microsatellite loci indicated**  
1120 **that workers from 28 *Atta colombica* colonies in the Gamboa area of Soberanía**  
1121 **park represent a single interbreeding population.** Specifically, the mean log-  
1122 likelihood was highest at  $K=1$  (although  $\Delta K$  cannot be assessed at  $K = 1$ ). When  $K$   
1123 was set to two, no individuals could be assigned to a single cluster. This Structure  
1124 plot included a burn-in of 250,000, MCMC reps of 500,000 and 25 iterations per  $K$ .  
1125

1126 **Extended Data Figure 7 STRUCTURE analyses of nuclear genetic marker data**  
1127 **from nine microsatellite loci from 297 *Paratrachymyrmex* and *Mycetomoellerius***  
1128 **workers,** from 194 and 103 colonies respectively, indicate that *P. cornetzi* and *P.*  
1129 *ADG3274* represent distinct ant species within the total community of eight species  
1130 belonging to these two ant genera in Soberanía Park. **a)** A STRUCTURE analysis  
1131 across all samples ( $N=297$ ) indicated that the most likely population subdivision is  
1132 two species because  $K = 2$  reached the highest ( $\Delta K$ ) resolution (burn-in 200000,  
1133 MCMC reps = 5000000, 25 iterations per  $K$ ). This analysis separated samples that  
1134 were all previously identified as *P. cornetzi* using morphological characters. It also  
1135 grouped three Barcode of Life Data System<sup>70</sup> ([www.boldsystems.org](http://www.boldsystems.org)) haplotypes  
1136 (AAP2324, ABY9919, ADG3341) separately from the newly recognized cryptic  
1137 species *P. ADG3274*, that otherwise clustered with the remaining species (light blue).  
1138 Although  $\Delta K$  cannot be calculated for  $K=1$ , the log-likelihood at  $K=1$  (-11121.864)  
1139 was substantially lower than the one at  $K=2$  (-9397.616), suggesting that ‘no  
1140 population subdivision’ is unlikely, consistent with all other species being  
1141 morphologically distinct. **b)** A similar STRUCTURE analysis using those individuals  
1142 assigned to cluster 1 in panel a ( $N=156$ ) indicated that three barcoded haplotypes  
1143 (AAP2324, ABY9919, ADG3341) located in the BOLD database represent a single  
1144 interbreeding population within Soberanía Park called *P. cornetzi*. Specifically,  $\Delta K$   
1145 was highest at 2 (1305.79) but the mean log likelihood was similar for  $K=1$  (-  
1146 3849.092) and  $K=2$  (-3698.860), suggesting that any existing population structure

1147 here is weak. The haplotypes also do not separate the dark blue and light blue bars  
1148 across our sampling sites consistent with failure to reliably assign individual workers  
1149 to clusters 1 or 2 as expected under high admixture rates. One sample was identified  
1150 as *P. ADG3274* by barcoding analysis (177340) (Fig. 5, Extended Data Fig. 8), but  
1151 could not be reliably distinguished from *P. cornetzi* by this STRUCTURE analysis. c)  
1152 A STRUCTURE plot generated for the remaining (non-*P. cornetzi*) individuals (N =  
1153 156), indicated that  $K=8$  was the most likely subdivision, with little to no admixture  
1154 between these species. The cryptic species *P. ADG3274* forms a distinct cluster in  
1155 this analysis. These genetic species assignments correspond well to species identities  
1156 obtained by morphological criteria and barcoding (Extended Data Fig. 8) and  
1157 included four currently named species (*P. bugnioni*, *M. isthmicus*, *M. opulentus*, and  
1158 *M. zeteki*), and three unnamed species (*M. PAN004* and *M. PAN002*, and *P.*  
1159 *ADG3274*). The ants *M. PAN004* and *M. PAN002* were placed in the genus  
1160 *Mycetomoellerius*<sup>35</sup> since they grouped with *M. isthmicus*, *M. opulentus*, and *M.*  
1161 *zeteki* in the CO1 barcoding tree (Extended Data Fig. 8, Supplementary Table 13) and  
1162 because they exhibited the required morphological characters to justify this decision.  
1163 This STRUCTURE analysis also suggested there are potentially more than one  
1164 species within the *M. zeteki* and *M. PAN002* clusters, and that six stray samples  
1165 (yellow bars) may also be distinct despite being morphologically grouped similar to  
1166 *P. bugnioni* (n = 2 samples), *M. isthmicus* (n = 1), *T. PAN004* (n = 1), and *M. zeteki*  
1167 (n = 2). Overall, the results of this analysis confirm that perhaps only ~ half of the  
1168 species are known even in a well-studied insect group in the tropics. The potential  
1169 taxonomic implications of the *P. cornetzi* and *P. ADG3274* species complex are  
1170 beyond the scope of the present study. Samples labelled with “no barcode” were ants  
1171 that were identified as belonging to *Paratrachymyrmex* and *Mycetomoellerius* using  
1172 morphological characters but were not included in barcoding analyses.

1173





1174 **Extended Data Figure 8 A majority-rule consensus tree based on COI sequences**  
1175 **of 185 *Paratrachymyrmex* and *Mycetomoellerius* ants supported our**  
1176 **microsatellite analyses as the same eight species were recognized as co-occurring**  
1177 **across Soberanía National Park.** Three of these species belonged to the genus  
1178 *Paratrachymyrmex* (*P. cornetzi* [n = 130], *P. ADG3274* [n = 15], *P. bugnioni* [n = 2])  
1179 and five belonged to the genus *Mycetomoellerius* (*M. zeteki* [n = 8], *M. opulentus* [n =  
1180 2], *M. isthmicus* [n = 7], *M. PAN002* [n = 10], *M. PAN004* [n = 11]; Supplementary  
1181 Table 13)<sup>22,35</sup> which is sister to the genus *Sericomyrmex*. This tree contains 9  
1182 additional public *P. cornetzi* COI sequences deposited at the Barcode of Life Data  
1183 System database<sup>70</sup> (BOLD, [www.boldsystems.org](http://www.boldsystems.org)) that supported that the main *P.*  
1184 *cornetzi* haplotype occurs at least from Ecuador to Costa Rica (N = 3 specimens), and  
1185 that the cryptic *P. cornetzi* haplotype [ADG3274] is distributed at least from the  
1186 Canal Zone in Panama to La Selva forest in Costa Rica (N = 6 specimens). The tree  
1187 was rooted by a worker of *M. smithii*.

1188


1189 **Extended Data Figure 9 Each of the five fungal haplotypes cultivated by *P.***  
1190 ***cornetzi* was widely distributed across Soberanía National Park.** For a map of  
1191 these sampling locations, see Extended Data Fig. 10, Supplementary Table 14. We  
1192 combined cultivars from La Seda plots 1 and 2 for this figure since these plots were

1193 within 200 m of each other. Four of the five fungal haplotypes that we obtained from  
1194 these Panama study plots matched cultivar haplotypes sampled from Brazilian  
1195 colonies of *Paratrachymyrmex* and *Sericomyrmex* by Solomon et al.<sup>35</sup>, suggesting  
1196 broad geographic distributions extending across Central and South America.  
1197 Additional Soberanía Park sampling localities where no specific plots were assigned  
1198 (so no labels in Extended Data Fig. 10) included La Laguna (N 9.1196, W -79.6942),  
1199 Horse Patch (N 9.11990, W -79.70730), and Barro Colorado Island (BCI: N 9.15744,  
1200 W -79.83523).

1201  
1202 **Extended Data Figure 10 We mapped local-scale higher neotattine colony**  
1203 **distributions within the six 20-m<sup>2</sup> study plots distributed across Soberanía**  
1204 **National Park.** This totalled 263 colonies (9 species) that were mapped by tracking  
1205 foraging workers back to their nests after which pinned specimens were screened in  
1206 microsatellite (Extended Data Fig. 7) DNA barcoding analyses (Extended Data Fig. 8,  
1207 Supplementary Table 13), as well as with morphological identification. We located  
1208 colonies by placing bait (polenta) in the leaf litter and observing foragers for  $27 \pm 3$   
1209 searching hours per plot (N = 163 total searching hours). The ant *M. opulentus* was  
1210 recorded within the forest, but not within any of the 20 m<sup>2</sup> plots. Nests were marked  
1211 by flags at nest entrances, which are inconspicuous holes under the leaf litter the  
1212 circumference of a pencil. Within 20-m<sup>2</sup> plots, we observed  $43 \pm 17$  (range: 21-66)  
1213 higher-neotattine colonies representing  $5 \pm 1$  (range: 4-7) higher-neotattine species.  
1214 Dashed lines indicate boundaries of 20m<sup>2</sup> plots, and an additional 5 m is provided  
1215 because some colonies were marked despite occurring just beyond the 20-m<sup>2</sup> plot  
1216 border lines. Trees > 1 m circumference dbh were also mapped. Plot abbreviations  
1217 were as follows: BP1: Bird Plot [N 9.16324, W -79.74553], GP1: Gamboa Plot [N  
1218 9.11489, W -79.69784], JG1: Juan Grande [N 9.13528, W -79.72141], LS1: La Seda  
1219 [N 9.15451, W -79.73583], LS2: La Seda [N 9.15624, W -79.73472], TB1: Thomas  
1220 Barbour [N 9.15744; W -79.83523]). The satellite image of the Panama Canal Zone is  
1221 from Google Maps and the country outline map is modified from Vemaps.com.




## Higher Neotattines




## Leafcutter farming

*A. echinator*


*A. colombica*



## Lower Attines



## Yeast farming




Paleoattine


Neoattine




a

## Lab feeding experiment





b











Leafcutter

*A. echinator*




Higher neotattine

*M. zeteki*




*S. amabilis*




Lower attine

*Apt. dentigerum*

Coral fungus farming










*C. costatus*





Paleoattine

Neoattine








a) *Paratrachymyrmex cornetzi*



b) *Atta colombica*



Diet P:C — 1:6 — 1:3 — 1:1 — 3:1 — 6:1



*P. cornetzi*

healthy fungus garden




Collapsed fungus garden



Cluster1 Cluster2




K=1




K=2


a) Testing whether *P. cornetzi* is isolated from populations of other *Paratrachymyrmex* and *Mycetomoellerius* species





b) Testing for population structure within *P. cornetzi*



c) Testing whether *P. ADG3274* is a reproductively isolated cryptic species







Higher-attine species

- *P. cornetzi*
- *P. ADG3274*
- *P. bugnioni*
- *M. isthmicus*
- *M. PAN002*
- *M. PAN004*
- *M. zeteki*
- *S. amabilis*
- not sequenced

