

The Nature of High $[\text{OIII}]_{88\mu\text{m}}/[\text{CII}]_{158\mu\text{m}}$ Galaxies in the Epoch of Reionization: Low Carbon Abundance and a Top-Heavy IMF?

Harley Katz^{1*}, Joakim Rosdahl², Taysun Kimm³, Thibault Garel⁴, Jérémy Blaizot², Martin G. Haehnelt⁵, Léo Michel-Dansac², Sergio Martin-Alvarez⁵, Julien Devriendt¹, Adrienne Slyz¹, Romain Teyssier⁶, Pierre Ocvirk⁷, Nicolas Laporte⁵, and Richard Ellis⁸

¹Sub-department of Astrophysics, University of Oxford, Keble Road, Oxford OX1 3RH, UK

²Univ Lyon, Univ Lyon1, Ens de Lyon, CNRS, Centre de Recherche Astrophysique de Lyon UMR5574, F-69230, Saint-Genis-Laval, France

³Department of Astronomy, Yonsei University, 50 Yonsei-ro, Seodaemun-gu, Seoul 03722, Republic of Korea

⁴Observatoire de Geneve, Universite de Geneve, 51 Ch. des Maillettes, 1290 Versoix, Switzerland

⁵Kavli Institute for Cosmology and Institute of Astronomy, Madingley Road, Cambridge CB3 0HA, UK

⁶Institute for Computational Science, University of Zurich, Winterthurerstrasse 190, CH-8057 Zurich, Switzerland

⁷Observatoire Astronomique de Strasbourg, Université de Strasbourg, CNRS UMR 7550, 11 rue de l'Université, 67000 Strasbourg, France

⁸Department of Physics and Astronomy, University College London, Gower Street, London WC1E 6BT, UK

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

ALMA observations of $z > 6$ galaxies have revealed abnormally high $[\text{OIII}]_{88\mu\text{m}}/[\text{CII}]_{158\mu\text{m}}$ ratios and $[\text{CII}]_{158\mu\text{m}}$ deficits compared to local galaxies. The origin of this behaviour is unknown. Numerous solutions have been proposed including differences in C and O abundance ratios, observational bias, and differences in ISM properties, including ionisation parameter, gas density, or photodissociation region (PDR) covering fraction. In order to elucidate the underlying physics that drives this high-redshift phenomenon, we employ SPHINX²⁰, a state-of-the-art, cosmological radiation-hydrodynamics simulation, that resolves detailed ISM properties of thousands of galaxies in the epoch of reionization. We find that the observed $z > 6$ $[\text{OIII}]_{88\mu\text{m}}$ -SFR and $[\text{CII}]_{158\mu\text{m}}$ -SFR relations can only be reproduced when the C/O abundance ratio is $\sim 8\times$ lower than Solar and the total metal production is $\sim 5.7\times$ higher than that of a Kroupa IMF. This implies that high-redshift galaxies are potentially primarily enriched by low-metallicity core-collapse supernovae with a more top-heavy IMF. As AGB stars and type-Ia supernova begin to contribute to the galaxy metallicity, both the $[\text{CII}]_{158\mu\text{m}}$ -SFR and $[\text{CII}]_{158\mu\text{m}}$ luminosity functions are predicted to converge to observed values at $z \sim 4.5$. While we demonstrate that ionisation parameter, LyC escape fraction, ISM gas density, and CMB attenuation all drive galaxies towards higher $[\text{OIII}]_{88\mu\text{m}}/[\text{CII}]_{158\mu\text{m}}$, observed values at $z > 6$ can only be reproduced with substantially lower C/O abundances compared to Solar. The combination of $[\text{CII}]_{158\mu\text{m}}$ and $[\text{OIII}]_{88\mu\text{m}}$ can be used to predict the values of ionisation parameter, ISM gas density, and LyC escape fraction and we provide estimates of these quantities for nine observed $z > 6$ galaxies. Finally, we demonstrate that $[\text{OI}]_{63\mu\text{m}}$ can be used as a replacement for $[\text{CII}]_{158\mu\text{m}}$ in high-redshift galaxies where $[\text{CII}]_{158\mu\text{m}}$ is unobserved and argue that more observation time should be used to target $[\text{OI}]_{63\mu\text{m}}$ at $z > 6$.

Key words: galaxies: high-redshift, galaxies: ISM, dark ages, reionization, first stars, ISM: lines and bands, ISM: kinematics and dynamics, galaxies: star formation

1 INTRODUCTION

Characterising the interstellar medium (ISM) of high-redshift galaxies deep into the epoch of reionization (EoR) represents one of the most interesting frontiers in the observational study of galaxy formation. Deep ALMA observations of forbidden infrared (IR) emission lines have now been confirmed out to $z > 9$ (e.g. Hashimoto et al. 2018; Laporte et al. 2019), providing insight into both the ISM properties and star formation histories of these primordial galaxies. As the sample of high-redshift systems with confirmed emission line de-

tections increases, certain differences have been discovered between reionization-epoch galaxies and those in the local Universe.

The $[\text{CII}]_{158\mu\text{m}}$ line (hereafter $[\text{CII}]$), is one of the dominant coolants of the ISM and hence one of the brightest IR emission lines in the local Universe (e.g. Spitzer 1978; Stacey et al. 1991). With an ionisation potential of 11.3eV, $[\text{CII}]$ emission can be stimulated in both ionised and neutral gas, often complicating its interpretation. Despite the wide range of ISM conditions that can emit $[\text{CII}]$, it has been empirically shown to strongly correlate with star formation in local galaxies (e.g. Boselli et al. 2002; De Looze et al. 2011, 2014; Herrera-Camus et al. 2015). Furthermore, observations have shown that the $[\text{CII}]$ line is especially bright in low-mass metal-poor dwarf

* E-mail: harley.katz@physics.ox.ac.uk

galaxies (e.g. [Poglitsch et al. 1995](#); [Cormier et al. 2010](#)), making it a particularly interesting target for studying the high-redshift Universe.

Earlier IR observations of $z > 6$ galaxies focused primarily on this line due to its predicted high luminosity and observability with ALMA. There are now more than 20 galaxies at $z > 6$ with [CII] detections as well as many more with non-detections. At slightly lower redshift (e.g. $4 \lesssim z \lesssim 6$), even larger samples of more than 100 galaxies have been surveyed ([Le Fèvre et al. 2020](#); [Béthermin et al. 2020](#)), of which 75 have been detected in [CII] emission ([Schaerer et al. 2020](#)). The non-detections are of particular interest because this places many of the EoR galaxies far below the local relation between [CII] and the star formation rate (SFR) (e.g. [Ota et al. 2014](#); [Maiolino et al. 2015](#); [Schaerer et al. 2015](#); [Pentericci et al. 2016](#); [Knudsen et al. 2016](#); [Bradač et al. 2017](#); [Laporte et al. 2019](#)). The exact origin of these deficits is currently unknown; however recent observations of extended, low surface brightness [CII] emission in the early Universe ([Fujimoto et al. 2019](#)) has helped relieve some of the tension between the high-redshift galaxies and the low-redshift relations ([Carniani et al. 2020](#)).

The [OIII] $_{88\mu\text{m}}$ line (hereafter [OIII]) is another strong IR emission line that has also been empirically established to correlate with star formation (e.g. [De Looze et al. 2014](#)). With an ionisation potential of 35.1 eV, [OIII] only traces ionised gas, in contrast to [CII]. Interestingly, in metal-poor local Universe dwarf galaxies, [OIII] often outshines [CII] (e.g. [Cormier et al. 2010](#); [Madden et al. 2013](#)), representing up to 2% of the total IR luminosity of the galaxy ([Cormier et al. 2015](#)). Since low-redshift metal poor dwarf galaxies are often considered analogues of high-redshift systems, [OIII] emission has been a particularly interesting target for the high-redshift Universe (e.g. [Inoue et al. 2016](#)) and has already been detected at $z > 9$ ([Hashimoto et al. 2018](#)).

While sample sizes of [OIII] emitting galaxies remain small at $z > 6$, there have now been ~ 10 detections, nearly all of which result in exceptionally high ratios of [OIII]/[CII], even compared to local metal-poor dwarf galaxies (e.g. [Laporte et al. 2019](#); [Harikane et al. 2020](#); [Carniani et al. 2020](#)). Even when correcting for extended [CII] emission ([Carniani et al. 2020](#)), the [OIII]/[CII] ratios can reach as high as 8, which represents the upper tail of the distribution of low-redshift metal-poor galaxies ([Madden et al. 2013](#)). Similar to the [CII] deficits observed in high-redshift galaxies, the exact origin of the high [OIII]/[CII] ratios is still debated as the ionisation parameter, photodissociation region (PDR) covering fraction, enrichment patterns, metallicity, ISM density, and many other physical properties of a galaxy can impact the [OIII]/[CII] ratio ([Harikane et al. 2020](#)).

Theoretical models, either analytic or numerical, are an invaluable tool for interpreting the origin of emission lines and physical properties of high-redshift galaxies. While historically, the understanding of emission lines, emission line ratios, and various strong-line diagnostics has relied upon idealised, 1D photoionization models (e.g. [Kewley & Dopita 2002](#); [Byler et al. 2017](#)), using codes such as CLOUDY ([Ferland et al. 2017](#)) or MAPPINGS ([Sutherland et al. 2018](#)), these models often lack some of the complex physical processes (e.g. SN feedback, magnetic fields, shocks, density inhomogeneities, etc.) that affect the physics of the ISM. Although they are significantly more computationally expensive and it is difficult to run “controlled” experiments as one can do with 1D photoionisation models, cosmological simulations provide the best insight into the complex physical nature of the high-redshift ISM.

Due to their computational expense, most recent studies of IR emission lines in cosmological simulations have focused on small samples or individual galaxies (e.g. [Vallini et al. 2015](#); [Olsen et al. 2017](#); [Katz et al. 2017, 2019b](#); [Pallottini et al. 2017, 2019a](#); [Arata](#)

[et al. 2020](#); [Lupi et al. 2019](#)), while only a few have attempted to model the emission in larger cosmological volumes (e.g. [Moriwaki et al. 2018](#); [Leung et al. 2020](#)). Obtaining larger samples of galaxies from simulations often relies on semi-analytic models (e.g. [Popping et al. 2016](#); [Lagache et al. 2018](#); [Popping et al. 2019](#)). It is not clear whether the results from small sample sizes can be generalised to the larger galaxy population while semi-analytic models do not resolve much of the physics that dictates ISM properties. Furthermore, previous attempts at large volumes containing many galaxies often compromise on important physics such as radiation transfer or sacrifice resolution due to computational expense.

Not all results from simulations agree on the expected trends between [OIII] and [CII] emission and various galaxy properties, likely due to differences in spatial and mass resolution, included physics, sub-grid models, and the method for modelling emission lines. For example, the [OIII]-SFR relations found in [Moriwaki et al. \(2018\)](#); [Arata et al. \(2020\)](#) point towards larger [OIII] luminosities per unit SFR compared to [Olsen et al. \(2017\)](#); [Katz et al. \(2019b\)](#). Likewise, the few galaxies simulated in [Katz et al. \(2019b\)](#) seem to fall closer to the local [CII]-SFR relation than those in ([Pallottini et al. 2017, 2019a](#)). The origin of these discrepancies can in part be due to galaxy-to-galaxy variance, in addition to differences in modelling techniques, which are well established to be far from straightforward ([Olsen et al. 2018](#); [Lupi et al. 2020](#)). Nevertheless, there have been many situations where simulations have agreed on emission-line properties of high-redshift galaxies and provide deep insights into or predictions of the early Universe. For example, the necessary but insufficient condition that LyC leakers have high O32 ([Katz et al. 2020b](#); [Barrow et al. 2020](#)) as confirmed at $z \sim 2-3$ ([Nakajima et al. 2020](#)) or the strong [OI] $_{63\mu\text{m}}$ (hereafter [OI]) emission line ([Olsen et al. 2017](#); [Katz et al. 2019b](#)), that was recently observed at $z > 6$ ([Rybak et al. 2020](#)).

While systematic uncertainties due to differences in modelling will continue to persist, issues due to galaxy-to-galaxy variance can be solved by simulating larger cosmological volumes. Until now there has been no full-box cosmological radiation hydrodynamics simulation with high enough resolution to semi-resolve the detailed structure of the ISM where these IR emission lines originate and with a large enough computational volume so that both the statistical properties of high-redshift galaxies can be studied and massive enough galaxies are probed to compare directly with observations. The primary goal of this work is to simulate the [CII] and [OIII] emission from such a state-of-the-art simulation to better understand the nature of high [OIII]/[CII] ratios at high redshift.

This paper is organised as follows: In Section 2, we briefly describe the SPHINX²⁰ simulation ([Rosdahl et al. in prep.](#)) as well as our modelling technique for line emission. In Section 3, we present the general [OIII] and [CII] properties of high-redshift galaxies at $z > 4.68$ and discuss the physics that drives high [OIII]/[CII] ratios. Finally, in Section 5, we present our discussion and conclusions.

2 NUMERICAL METHODS

2.1 Cosmological Simulations

This work makes use of the SPHINX²⁰ cosmological radiation hydrodynamics simulation ([Rosdahl et al. in prep.](#)). SPHINX²⁰ is the largest of all simulations in the SPHINX suite of cosmological radiation ([Rosdahl et al. 2018](#); [Katz et al. 2020a](#)) and magneto-radiation ([Katz et al. 2021](#)) hydrodynamics simulations and is designed to model both the inhomogeneous process of reionization and the escape of LyC photons from the ISM. SPHINX²⁰ is ideal for studying the emission line

properties of high-redshift galaxies because for more than 20,000 galaxies at $z \geq 4.68$, spanning a wide range in halo mass, stellar mass, and star-formation rate (SFR) in each simulation snapshot, we can probe the detailed properties of the ISM. This includes the inhomogeneous radiation field, as well as the detailed density structure, temperature, and metal enrichment, which represent key underlying quantities that impact emission line luminosities.

The details of SPHINX²⁰ are very similar to those presented in Rosdahl et al. (2018). We briefly describe the simulations with emphasis on the differences between the simulations used in this work compared to that of Rosdahl et al. (2018). The simulations are run with RAMSES-RT (Rosdahl et al. 2013; Rosdahl & Teysier 2015), which is a radiation hydrodynamics version of the open-source, adaptive-mesh refinement RAMSES code (Teyssier 2002). Initial conditions are generated with MUSIC (Hahn & Abel 2011) in a cosmological box with a comoving side length of 20 Mpc assuming the following cosmology: $\Omega_\Lambda = 0.6825$, $\Omega_m = 0.3175$, $\Omega_b = 0.049$, $h = 0.6711$, and $\sigma_8 = 0.83$. Due to the small volume of the simulations, the initial conditions were selected from multiple realisations to produce an average DM halo mass function at $z = 6$. The initial composition of the gas is set to be 76% H and 24% He by mass and have an initial metallicity of $3.2 \times 10^{-4} Z_\odot$. The dark matter particle mass is $2.5 \times 10^5 M_\odot$ and we have decreased the stellar particle mass from $1000 M_\odot$ in SPHINX¹⁰ to $400 M_\odot$ in SPHINX²⁰. The AMR grid is adaptively refined to resolve the local Jeans length by at least 4 cells or if the gas or DM mass of a cell is more than 8 times its initial value, resulting in a maximum physical resolution of $7.3 \text{ pc}/h$ at $z = 6$.

Stars are allowed to form in the simulation at the maximum level of refinement following a thermo-turbulent prescription (Kimm et al. 2017; Trebitsch et al. 2017; Rosdahl et al. 2018). Stellar feedback is included in the form of supernova (SN) explosions and Lyman-continuum radiation. Star particles with $3 \text{ Myr} < \text{age} < 50 \text{ Myr}$ can explode via individual SNe by sampling from a delay-time distribution. When a SN event occurs, momentum is injected into the host and surrounding cells such that the amount of momentum injected is dependent on the phase of the SN that is resolved (Kimm et al. 2015). Star particles also continuously inject ionising photons into their host cells depending on their mass, age, and metallicity. In contrast to SPHINX¹⁰, we only include two radiation energy bins to reduce the memory requirements of the simulation, set at the ionisation potentials of HI and HeI. The spectral energy distribution (SED) model employed in the simulation is BPASS (Eldridge et al. 2008; Stanway et al. 2016). We have updated the SED from SPHINX¹⁰ which uses v2.0 to v2.2.1 (Stanway & Eldridge 2018). Non-equilibrium chemistry is evolved locally for H, H⁺, e⁻, He, He⁺, and He⁺⁺ as are the associated cooling processes (see the Appendix of Rosdahl et al. 2013). Cooling from metal lines is included at $T > 10^4 \text{ K}$ following tables computed with CLOUDY (Ferland 1996) as well as at $T \leq 10^4 \text{ K}$ using the analytic model from Rosen & Bregman (1995).

For all simulation snapshots, we compute halo properties using the ADAPTAHOP halo finder Aubert et al. (2004); Tweed et al. (2009) in the most massive submaxima mode as described in Rosdahl et al. (2018). Furthermore, LyC escape fractions are also computed for each halo following Rosdahl et al. (2018), using the open-source RASCAS code (Michel-Dansac et al. 2020).

2.2 Emission-Line Modelling

Because our goal is to measure the [CII] and [OIII] emission line properties of high-redshift galaxies, we must compute the [CII] and [OIII] emission lines for each cell in the simulation. It is currently computationally intractable with our available resources to follow

the full non-equilibrium chemistry and radiation coupling for C and O directly in the simulation so we resort to post-processing the simulations in order to measure the [CII] and [OIII] luminosities. We select seven simulation snapshots at $z = 10, 9, 8, 7, 6, 5, \& 4.68$, the final redshift of our simulation for post-processing. Our method is similar in spirit to that used in Katz et al. (2019b,a, 2020b).

For each cell in the simulation, we know the density, temperature, metallicity and ionisation parameter (the ratio of ionising photons to hydrogen atoms), as well as the cell size and redshift. To obtain the radiation field at energies below 13.6eV, we have post-processed the simulation snapshots with radiation transfer in the optically thin limit. This is done by restarting the simulation assuming that the gas, dark matter, and stars are static and using RAMSES-RT to evolve the radiation in two additional energy bins (5.6eV-11.2eV and 11.2eV-13.6eV) for at least the light-crossing time¹ of the largest halo in the volume at that redshift. We then use CLOUDY (Ferland et al. 2017) to compute the [OIII] and [CII] emission lines from the cells based on their density, temperature, metallicity, redshift, and radiation field in the four² energy bins, i.e. 8 local parameters.

Because there are billions of cells across the seven snapshots, it is impractical to compute a CLOUDY model for each cell. Furthermore, computing a grid across our eight parameters would mean that some parameters are only sparsely sampled in the grid which could result in errors in interpolation. Hence we adopt a different approach as follows:

(i) For each snapshot we randomly sample between 1.5-4.5 million simulation cells from galaxies across the full halo mass range.

(ii) A CLOUDY model is run for the selected cells assuming an open geometry gas slab that has the same density, temperature, and metallicity as the simulation cell. The width of the slab is set to be the cell length and the shape of the radiation field within each radiation bin is set to follow that of the Milky Way (i.e. the total energy within the wavelength range of the radiation bin matches that of the simulation but the shape within the bin is kept constant for all cells). The CLOUDY models are all run in a constant temperature mode and a CMB background at the relevant redshift is also included.

(iii) After the photoionisation models have been computed, for each snapshot, we randomly split the results into a training, cross-validation, and test set such that 25% of cells are reserved for testing, 22.5% are used for cross-validation, and 52.5% of cells are used for training. For each line, we train two machine learning models. The first is a classifier that is used to predict whether the emission from the cell is $> 10^{-3} L_\odot$ while the latter is used to predict the log of the total luminosity of the cell. In contrast to Katz et al. (2019b), we predict the log of the luminosity rather than luminosity to better the accuracy in the low luminosity regions which is important for extended emission. For this work, we use the CATBOOST³ (Prokhorenkova et al. 2017; Veronika Dorogush et al. 2018) implementation of gradient boosted decision trees, which we find to be an improvement in terms of accuracy and well as training and prediction speed compared to the Random Forests used in Katz et al. (2019b). Models are trained for a maximum of 10,000 iterations and are stopped early if the loss-function does not improve for a period of 500 consecutive epochs. For the classifier, we use a binary cross-entropy loss function while

¹ Note that because the simulations use the variable speed of light approximation, we use a conservative estimate for the light crossing time by using the lowest value of the speed of light used in the simulation.

² This includes the two bins computed in post-processing and the two ionising radiation bins computed on-the-fly in the simulation.

³ <https://github.com/catboost/catboost>




Figure 1. Hydrogen column density (N_{H}), (left), [CII] surface brightness (centre), and [OIII] surface brightness (right) maps of the three most massive galaxies in SPHINX²⁰. These haloes have masses of $1.31 \times 10^{11} M_{\odot}$, $1.29 \times 10^{11} M_{\odot}$, and $9.97 \times 10^{10} M_{\odot}$, and 10Myr averaged star formation rates (SFRs) of $17.6 M_{\odot} \text{yr}^{-1}$, $20.7 M_{\odot} \text{yr}^{-1}$, and $39.7 M_{\odot} \text{yr}^{-1}$, respectively.

for the regressor, we aim to maximise the R^2 score. For [CII], we find on the test set that the mean accuracy of the classifier is $99.75 \pm 0.08\%$, the mean R^2 score of the regressor is 0.996 ± 0.001 , and that the total luminosity is conserved to within $2.89\% \pm 2.84\%$. For [OIII], we find on the test set that the mean accuracy of the classifier is $99.91 \pm 0.06\%$, the mean R^2 score of the regressor is 0.997 ± 0.002 , and that the total luminosity is conserved to within $11.7\% \pm 6.53\%$. Our values of

total luminosity conversion are well within the systematic errors of modelling the emission lines.

(iv) Finally, we apply the models to every cell in each halo across all seven snapshots.

To run the CLOUDY models, we must also set the abundances of each metal. For our fiducial model, we use the GASS command that sets the abundance ratios to be consistent with those of the Sun




Figure 2. [CII]- M_{vir} (top) and [OIII]- M_{vir} (bottom) relations for galaxies in SPHINX²⁰ at $z = 4.68 - 10$. The solid and dashed lines represent the results for the Solar abundance and core collapse SN abundance models, respectively. The shaded regions represent the 1σ scatter about the relation. The dotted lines in the top panel represent the semi-analytic models from [Silva et al. \(2015\)](#). Little evolution is seen in these relations as a function of redshift whereas systematic differences occur due to choice of abundance model.

([Grevesse et al. 2010](#)). Such abundances may not be representative of the early Universe due to different chemical enrichment scenarios (e.g. [Maiolino & Mannucci 2019](#)). For example, high-redshift galaxies may be predominantly enriched by core-collapse SN compared to a mix of core-collapse SN, type-Ia SN, and winds from AGB stars. For this reason, we also run an additional set of models where the chemical abundance ratios are consistent with the low metallicity SN yields of [Nomoto et al. \(2006\)](#). This is particularly important for the [OIII]/[CII] ratio because the C/O abundance ratio from [Nomoto et al. \(2006\)](#) is eight times less than that of Solar abundance patterns. Finally, we also test models including molecules but find that our results are not substantially different when these are included and so for the results presented in this work, molecules have been excluded.

3 RESULTS

In this section, we present our results on [CII] and [OIII] emission from high-redshift galaxies, with a particular focus on the origin of




Figure 3. [CII] (top) and [OIII] (bottom) luminosity functions at different redshifts for galaxies in SPHINX²⁰ versus observations. The solid and dashed lines represent the results for Solar abundance and core collapse SN abundance models, respectively. The shaded region represents the 1σ scatter about the relation. Observational constraints for the [CII] luminosity function are shown for [Swinbank et al. \(2012\)](#); [Capak et al. \(2015\)](#); [Aravena et al. \(2016\)](#); [Cooke et al. \(2018\)](#); [Uzgil et al. \(2021\)](#) and the dotted and dashed lines represent results from the semi-analytic models of [Lagache et al. \(2018\)](#) and [Popping et al. \(2016\)](#), respectively. No observational constraints are available for the [OIII] luminosity function at high redshift.

high [OIII]/[CII] ratios at $z > 6$ and how these two emission lines can be used to constrain the properties of high-redshift galaxies.

We have calculated the [CII] and [OIII] emission for thousands of galaxies in each simulation snapshot between $z = 10$ and $z = 4.68$. In Figure 1 we show density, [CII], and [OIII] maps of the three most massive haloes at $z = 6$, with halo masses of $1.31 \times 10^{11} M_{\odot}$, $1.29 \times 10^{11} M_{\odot}$, and $9.97 \times 10^{10} M_{\odot}$, respectively. These galaxies have 10Myr averaged star formation rates (SFRs) of $17.6 M_{\odot} \text{yr}^{-1}$, $20.7 M_{\odot} \text{yr}^{-1}$, and $39.7 M_{\odot} \text{yr}^{-1}$, well within the range of galaxies at have been observed at $z > 6$ (e.g. [Maiolino et al. 2015](#)).

These galaxies exhibit very different morphologies. The first is a turbulent, and partially disrupted rotating disc, the second is a triple major merger, and the third, while more similar to the first, possesses two strongly star forming regions and clear evidence of dynamical disturbance. Such properties are rather characteristic of high-redshift

galaxies where strong dynamical interactions are more common than in lower-redshift galaxies.

Consistent with our earlier work (Katz et al. 2019b), we find that the [CII] emission follows the densest gas structures in the galaxy while [OIII] exhibits a different morphology, more coincident with the young star forming regions and mildly lower density gas (see also Pallottini et al. 2019b; Lupi et al. 2020). With a higher ionisation potential and lower critical density compared to C⁺, such behaviour is expected for O⁺⁺.

3.1 General Properties

With such a large sample of galaxies, we can begin to overcome issues related to sample variance and make predictions regarding the relations between [CII] and [OIII] luminosity and galaxy properties.

3.1.1 [CII] – M_{vir} and [OIII] – M_{vir} relations

In Figure 2 we show the relation between [CII] and [OIII] luminosity and halo virial mass as a function of redshift for the models with Solar abundance ratios and core collapse SN abundance ratios. As halo mass increases, so do the [CII] and [OIII] luminosities. Our predicted relation between $L_{\text{[CII]}}$ and M_{vir} using Solar abundance ratios is in good agreement with the semi-analytic model of Silva et al. (2015) at $M_{\text{vir}} \gtrsim 9.5 M_{\odot}$. The two models deviate from each other at the low mass end. Our models with the same abundance ratios exhibit almost no redshift dependence. Similarly, the relation between $L_{\text{[OIII]}}$ and M_{vir} shows no evidence of redshift dependence.

Constraining the [CII]- M_{vir} and [OIII]- M_{vir} relations is of particular importance for forecasting intensity mapping results (e.g. Lagache 2018; Yue & Ferrara 2019) and we have tables of these values available online⁴.

3.1.2 [CII] and [OIII] luminosity functions

Since M_{vir} is not an observed quantity, we do not have any direct observational constraints between [OIII], [CII], and M_{vir} . However, observational constraints are available for the [CII] luminosity function and in the top panel of Figure 3, we compare our predicted [CII] luminosity functions as a function of redshift to various observational constraints (Swinbank et al. 2012; Capak et al. 2015; Aravena et al. 2016; Cooke et al. 2018; Uzgil et al. 2021) as well as other semi-analytic models (Popping et al. 2016; Lagache et al. 2018). We observe a strong redshift dependence of the [CII] luminosity function such that as redshift decreases, the number of [CII] emitters at all luminosities increases. Our simulations have very few galaxies with [CII] luminosities $> 10^{42} \text{erg s}^{-1}$ which is approximately the lower limit of where observational constraints begin. Nevertheless, we find that the $z \sim 5.5$ constraint from Capak et al. (2015) falls in between our $z = 5$ and $z = 6$ predictions. Our models predict far fewer [CII] emitters than the estimates from Aravena et al. (2016) at $z = 6-8$, and are more in line with the $z = 6-8$ upper limit from Uzgil et al. (2021). Compared to results from commonly used semi-analytic models, we predict significantly more [CII] emitters at $L_{\text{[CII]}} > 10^8 L_{\odot}$ than both Popping et al. (2016); Lagache et al. (2018). The model that uses core collapse SN abundance ratios predicts a luminosity function much lower than observations.

Unfortunately, there are no observational constraints on the [OIII] luminosity function at high redshift and it is not often discussed

in the literature (although see Moriwaki et al. 2018). Once again, we expect less redshift evolution in the [OIII] luminosity function compared to [CII] due to the fact that O is predominantly produced in core collapse SN; hence, at very high redshifts (e.g. $z \gtrsim 9$), [OIII] may be the most likely FIR line to be observed.

3.1.3 [CII] – SFR and [OIII] – SFR relations

There has been a considerable amount of debate in the literature, both from an observational and theoretical perspective on how the high-redshift [CII]-SFR relation compares to that in the local Universe. Early observations suggested a large deficit of [CII] emission in $z > 6$ galaxies compared to those at $z = 0$ (e.g. Maiolino et al. 2015; Pentericci et al. 2016; Laporte et al. 2019). However, more recent observations that account for extended [CII] emission reduce some of this difference (Carniani et al. 2020) and by $z \sim 4.5$, the observed relation is nearly identical to that in the local Universe (Schaerer et al. 2020).

The origin of [CII] deficits is currently unknown as simulations also do not agree. For example, the zoom-in simulations of individual high-redshift galaxies in Katz et al. (2019b) show that haloes tend to fall very close to the observed low-redshift relation, and when [CII] deficits do occur in their simulations, it is after a burst of star formation, when the galaxy is also at its brightest, perhaps suggesting an observational bias. The simulations of Arata et al. (2020) produce a [CII]-SFR relation steeper than that observed in the local Universe, likely due to different enrichment channels as well as different ISM properties, although their highly star-forming galaxies are consistent with $z = 0$ observations. In contrast, Pallottini et al. (2017, 2019a) find that for similar halo masses to Katz et al. (2019b), their haloes have both higher star formation rates and significantly larger [CII] deficits.

This disagreement is also present between different simulations regarding the high-redshift [OIII]-SFR relation where Katz et al. (2019b); Olsen et al. (2017) find [OIII] deficits while Arata et al. (2020); Moriwaki et al. (2018), find an [OIII]-SFR relation consistent with $z = 0$. Interestingly, while Arata et al. (2020) cannot produce lower luminosity [OIII] emitters, Katz et al. (2019b) struggles to produce high [OIII] emitters. In summary, the current lack of agreement in the literature, both in observations and simulations, prohibits robust interpretations of observed high-redshift galaxies and deserves further study.

In the top row of Figure 4, we show the [CII]-SFR relations for SPHINX²⁰ galaxies at different redshifts compared with observations. We show models using Solar abundances, core collapse SN abundances, and core collapse SN abundances where we have artificially and uniformly enhanced the metallicities of the gas cells in each simulated galaxy by a factor of 5.72. Regardless of the abundance model, we see a strong evolution with redshift in that at fixed SFR, the [CII] luminosity increases with decreasing redshift. To first order, the line luminosity scales with the ISM metallicity which is why the Solar abundance model produces higher [CII] for fixed SFR compared to the core collapse abundance model. In Figure 5, we plot the mean ISM metallicity as a function of SFR for various redshifts and see that as redshift decreases, the ISM metallicity at fixed SFR increases. Hence we would expect an increase in [CII] luminosity. The [CII] luminosity is also sensitive to the amount of low temperature neutral gas. We expect more of this gas is present in higher halo masses. In Figure 6 we show the evolution of the M_{vir} -SFR relation as a function of redshift and we also see here that at fixed SFR, halo mass increases with decreasing redshift. Hence this effect should also drive higher

⁴ <https://github.com/HarleyKatz/SPHINX-CII-OIII>




Figure 4. Mean [CII]-SFR relations (top) and [OIII]-SFR relations (bottom) for galaxies in the SPHINX²⁰ simulation at $z = 5 - 9$ compared to observations. The left, centre, and right rows represent models where we assume Solar abundances, core collapse SN abundances, or core collapse SN abundances with 5.72 times enhanced metallicity, respectively. The shaded regions represent the 1σ scatter about the relation. Black data points with errors represent observed galaxies at $z = 5 - 9$ whereas coloured data points represent the brightest [CII] and [OIII] emitters at each redshift. We show high-redshift fits from [Schaerer et al. \(2020\)](#); [Harikane et al. \(2020\)](#); [Carniani et al. \(2020\)](#) as well as $z = 0$ relations for dwarfs and starbursts (SBs) from [De Looze et al. \(2014\)](#).

[CII] luminosities both due to the increased amount of gas and the decreased susceptibility of the galaxy to SN feedback.

Since [OIII] is related to the efficiency of the feedback in the galaxy, we would expect the opposite effect compared to [CII]. The larger halo mass at fixed SFR would reduce the amount of [OIII] emission and counteract the increase in metallicity with decreasing redshift. The bottom row of Figure 4 shows the [OIII]-SFR relations as a function of redshift compared with observations. Regardless of the choice of abundance ratios, there is almost no redshift evolution. While it is expected that this relation does not evolve as strongly with redshift as the [CII]-SFR relation, it is an interesting coincidence that there is no redshift evolution at all.

The redshift evolution in the [CII]-SFR relation may be even more pronounced than shown in Figure 4 due to the expected evolution in C abundance. While O is predominantly formed in core collapse SNe soon after the onset of star formation, the formation timescale for C is significantly longer due to the fact that it is expelled into galaxies via both core collapse SNe and type-Ia SNe as well as in winds from asymptotic giant branch (AGB) stars (e.g. [Maiolino & Mannucci 2019](#)). Hence the amount of C per unit metallicity is expected to increase with decreasing redshift and therefore enhance

the redshift evolution in the relation. The top left and top centre panels of Figure 4 show the extreme cases of Solar abundances and core collapse SN abundances, respectively. Since we do not explicitly model the varying timescales of enrichment of different elements in the simulation, we have modulated the abundance ratios in post-processing to explore how this physics may affect the [CII]-SFR relation. Interestingly, when we assume Solar abundance ratios, we find that the simulated galaxies are in much better agreement with observations than when we assume core collapse SN abundance ratios. At $z \geq 6$, our Solar abundance relations are very consistent with the fitted relation from [Harikane et al. \(2020\)](#), and by $z = 4.68$, our relation has converged to that of [Schaerer et al. \(2015\)](#). At $\text{SFR} < 1 M_{\odot} \text{yr}^{-1}$, our relations seem to converge more towards those of [Carniani et al. \(2020\)](#) than that of [Harikane et al. \(2020\)](#), although at higher SFR, these two relations are relatively indistinguishable.

Compared to the local relation (shown as the yellow and brown lines in Figure 4), our $z \geq 6$ galaxies tend to produce small [CII] deficits even for Solar abundance ratios possibly due to the lower metallicity and lower halo mass per fixed SFR. The most [CII] bright galaxies in SPHINX²⁰ are however consistent with the local relation when assuming Solar abundance ratios. This is consistent with the




Figure 5. SFR- Z relation as a function of redshift. Z represents the mass-weighted gas-phase metallicity. The shaded region represents the 1σ standard deviation about the relation. As redshift decreases, galaxies at fixed SFR have higher metallicities.




Figure 6. SFR- M_{vir} relation as a function of redshift. The shaded region represents the 1σ standard deviation about the relation. As redshift decreases, galaxies at fixed SFR have higher halo masses or alternatively, galaxies at fixed halo mass have lower SFR.

results found in [Katz et al. \(2019b\)](#) which used similar subgrid physics and spatial/mass resolution and demonstrates the need for larger samples of galaxies.

When assuming core collapse SN abundance ratios, we find very strong [CII] deficits that appear inconsistent with the high-redshift galaxy population due to the drastically reduced C abundance. Taken at face-value, it would appear that our simulations prefer galaxies to exhibit abundance ratios closer to Solar values. However, we argue below that this is not necessarily the case.

One can see in the bottom row of Figure 4 that regardless whether we assume Solar abundances or core collapse SN abundances, our [OIII]-SFR relation underpredicts individual high-redshift observations as well as the fitted relation from [Harikane et al. \(2020\)](#). There is little difference between the [OIII]-SFR relation for the two abundance assumptions because the primary O enrichment channel is via core collapse SN and thus the O abundance is very similar be-

tween the two models. Hence while a Solar abundance ratios model can reproduce the [CII]-SFR relation at high redshift, it struggles with the [OIII]-SFR relation, as was the case in our earlier work [Katz et al. \(2019b\)](#). How then can we reconcile both relations with observations?

Changing the ISM properties of our high redshift galaxies is unlikely to relieve the tension. Because [OIII] and [CII] originate in different regions of temperature-density phase-space, attempting to create more [OIII]-emitting gas would reduce the amount of [CII]-emitting gas. Hence for the Solar abundance model, the reduction in [CII] would disrupt the current agreement in favour of improving the [OIII]-SFR relation. For this reason, this is not a preferred solution. If the galaxies were significantly gas richer at fixed halo mass in reality compared to what is predicted by the simulation, then this solution could be an option. However, [Katz et al. \(2020b\)](#) showed that for the SPHINX suite of simulations, these high-mass galaxies accrete gas relatively efficiently. The gas mass could only increase by a factor of about two as to not supersede the cosmic baryon fraction, which is not nearly enough to overcome the near order of magnitude deficit in the [OIII]-SFR relation.

Since when assuming Solar abundance ratios, the [CII]-SFR relation is in good agreement with observations, we argue that it is important that the total carbon mass in galaxies is not significantly different from our fiducial model. One solution to our problem could be to normalise the [Nomoto et al. \(2006\)](#) abundance patterns to have the same amount of C as the Solar abundance model. This implicitly assumes that the total metal mass in each cell is 5.72 times higher than what is currently predicted by the simulation. To explore this option, in the right panel of Figure 4, we plot the relations where we have simply scaled the [CII] and [OIII] luminosities of each galaxy for the core collapse SN abundance model by a factor of 5.72. Here we can see that the good agreement between the [CII]-SFR relation is once again recovered while the [OIII]-SFR relation is in significantly better agreement with the observed high-redshift galaxies. We emphasise that in this model, the C/O relation is the same as in the core collapse SN abundance model, but the total amount of metals is increased.

While this solution may seem *ad hoc*, there is precedent for this in high-redshift galaxies. Simulations show that low metallicity galaxies may produce a more top-heavy stellar initial mass function (IMF) compared to what is observed in the local Universe (e.g. [Safrańek-Shrader et al. 2014](#); [Chon et al. 2021](#)). More top-heavy IMFs result in an increased amount of SNe which leads to an increase in metal production as well as more high-energy photons which can form more O^{++} . In SPHINX²⁰, we already artificially boost the energy from SNe by a factor of 4 in order to match the high-redshift UV luminosity function ([Rosdahl et al. 2018](#)). This extra energy is thought to be needed in order to overcome numerical over-cooling; however it also may be reflecting the necessity for a more top-heavy IMF. Although the energy from SN in the simulation is artificially boosted, the metal yields are still consistent with a standard ([Kroupa 2001](#)) IMF. A more top-heavy IMF would produce an ISM that is similar to what we find in this work already due to the boosted feedback. Therefore, we argue that our results are potentially indicative of a top-heavy IMF in the early Universe. Future simulations that track both the evolving metal enrichment channels as well as sample different stellar IMFs can confirm this prediction, as also discussed by [Arata et al. \(2020\)](#).

3.1.4 [OIII]/[CII] ratios

In Figure 7 we plot mean [OIII]/[CII] versus SFR relations for SPHINX²⁰ galaxies compared with observations. We show both the




Figure 7. $[\text{OIII}]/[\text{CII}]$ versus SFR for galaxies in our simulation at different redshifts compared to observations for the Solar abundance model (top) and the core collapse SN abundance model (bottom). The shaded regions represent the 1σ scatter about the relation. Black data points represent $z = 5 - 9$ galaxies from Harikane et al. (2020) while red data points represent those same galaxies accounting for surface brightness dimming (Carniani et al. 2020). Orange points represent individual low redshift dwarf galaxies from the DGS survey (Madden et al. 2013). The red dotted line is the high-redshift fitted relation from Harikane et al. (2020) while the maroon and orange lines are the $z = 0$ relations for dwarf and starburst galaxies, respectively, from De Looze et al. (2014).

Solar abundance model (top) as well as the core collapse SN abundance model (bottom) and note that the model with the $5.72\times$ boosted metallicity that reproduces both the high-redshift $[\text{CII}]/\text{SFR}$ and $[\text{OII}]/\text{SFR}$ relations also predicts the same $[\text{OIII}]/[\text{CII}]$ ratios as the core collapse SN abundance model. Thus any future discussion of $[\text{OIII}]/[\text{CII}]$ ratios in the core collapse SN abundance model also applies to that with boosted metallicity.

In contrast to the observed relations, we find that the trend between $[\text{OIII}]/[\text{CII}]$ and SFR is not monotonic. Between $-2 < \log_{10}(\text{SFR}/M_{\odot}\text{yr}^{-1}) < 0$, we find decreasing $[\text{OIII}]/[\text{CII}]$ ratios as SFR decreases however, at $\log_{10}(\text{SFR}/M_{\odot}\text{yr}^{-1}) > 0$, the trend flattens, regardless of the assumption on abundance ratios. Redshift evolution pushes galaxies towards lower $[\text{OIII}]/[\text{CII}]$ with decreasing redshift, especially at $z < 6$.

As expected, compared to the observed high-redshift, $z = 5 - 9$ galaxies, the Solar abundance model produces $[\text{OIII}]/[\text{CII}]$ ratios that are far lower. At SFRs of $10M_{\odot}\text{yr}^{-1}$ the $[\text{OIII}]/[\text{CII}]$ deficit for the mean galaxy in our simulations is more than an order of magnitude compared to observations (see also Laporte et al. 2019). The $[\text{OIII}]/[\text{CII}]$ ratios are even lower than the metal poor dwarf galaxies in the local Universe and more consistent with local starburst galaxies. However, when we assume that the O and C abundances are similar to that of core collapse SNe, we find that the high $[\text{OIII}]/[\text{CII}]$ ratios at high redshift can be reproduced by our simulation. Still, we do not have many galaxies with $[\text{OIII}]/[\text{CII}] > 10$ as seen in Harikane et al. (2020). When accounting for $[\text{CII}]$ surface brightness dimming corrections (Carniani et al. 2020), we find very good agreement between $z = 5 - 9$ galaxies and the highly star-forming galaxies in SPHINX²⁰.

Because of the good agreement between the core collapse SN abundance model and observations, we argue that low C/O abundance ratios at high redshift could be the primary driver of high $[\text{OIII}]/[\text{CII}]$ ratios compared to the local Universe. Such an argument was proposed by Steidel et al. (2016) to explain the offsets of $z \sim 2 - 3$ galaxies on the BPT diagram. Similarly, Arata et al. (2020) demonstrated with cosmological simulations that explicitly track C and O abundances from different chemical enrichment channels that the C/O ratio is indeed significantly lower in high-redshift galaxies due to the delayed onset of AGB winds. However, Harikane et al. (2020) suggest that while lower C/O ratios may explain some of the enhanced $[\text{OIII}]/[\text{CII}]$ ratios observed at high redshift, the reduced C abundance alone is not enough to explain some of the non-detections of $[\text{CII}]$; hence, the ratio by itself cannot explain the galaxies with > 1 dex lower $[\text{CII}]/\text{SFR}$. Once again, these $[\text{CII}]$ deficits are partially relieved by surface brightness dimming corrections (Carniani et al. 2020); however, it is important to understand what other physical mechanisms can drive high $[\text{OIII}]/[\text{CII}]$ ratios as well as $[\text{CII}]$ deficits.

3.2 What else drives high $[\text{OIII}]/[\text{CII}]$ ratios?

Harikane et al. (2020) systematically discuss eight different effects that can impact the $[\text{OIII}]/[\text{CII}]$ ratio in high-redshift galaxies. They employ idealised models of the ISM using CLOUDY to better understand the impact of different ISM properties on the $[\text{OIII}]/[\text{CII}]$ ratio. While these simplified photoionization models are useful for isolating individual effects, real high-redshift galaxies exhibit a complicated, multi-phase ISM structure. In this section, we explore seven of the eight effects⁵ in the context of the SPHINX²⁰ simulations.

3.2.1 Lower PDR covering fractions

$[\text{OIII}]$ is generally emitted very close to stars from HII regions while $[\text{CII}]$ tends to originate further out in PDRs. Therefore if the HII region becomes large enough so that the gas cloud is density-bounded

⁵ The last effect is the lower C/O abundance ratios due to different enrichment channels in the early Universe which has already been discussed above.




Figure 8. 2D histograms of [CII]/SFR versus [OIII]/SFR for galaxies in the SPHINX²⁰ simulation between $6 \leq z \leq 10$ coloured by virial mass (top left), sSFR (top centre), 90th percentile ISM density (top right), ISM metallicity (bottom left), LyC escape fraction (bottom centre), and 90th percentile ISM ionisation parameter (bottom right). We show results for the core collapse SN abundance model with the $5.72\times$ enhanced metallicity. n_{H} , Z , and q have are calculated as the mass-weighted quantities in the [CII] and [OIII] emitting regions (Z , n_{H}) or HII regions (\bar{q}). Cyan points represent individual high-redshift galaxies from Carniani et al. (2020).

(i.e. the outer radius of the cloud is smaller than the Stromgren radius), we expect there to be very little [CII] emission. Molecular clouds exhibit complex density structures where some regions around the star are expected to remain shielded from the ionising radiation and exist in a neutral state. Harikane et al. (2020) argue that if the covering fraction of PDRs (C_{PDR}) surrounding stars is significantly less than unity, we would expect an increase in [OIII]/[CII].

Because the ISM structure in our simulations is highly complex, we cannot define a direct metric for C_{PDR} in our simulations that matches the idealised models of Harikane et al. (2020). Nevertheless, we have measured the LyC escape fraction (f_{esc}) for each galaxy in SPHINX²⁰ which provides a proxy for C_{PDR} on a galaxy scale. We expect that galaxies that have higher f_{esc} have lower PDR covering fractions and should thus exhibit higher ratios of [OIII]/[CII].

From simulations, it is well established that f_{esc} is a feedback regulated quantity (e.g. Trebitsch et al. 2017; Kimm et al. 2017). Hence it is the SN feedback that dictates the PDR covering fraction in our simulation more than the ionising photon production. Such an effect can only be realised in numerical simulations.

In the bottom-centre panel of Figure 8, we show a 2D histogram of [CII]/SFR versus [OIII]/SFR coloured by $\log_{10}(f_{\text{esc}})$. Galaxies in our simulation that exhibit the highest values of f_{esc} and, by deduction, lower PDR covering fractions also tend to exhibit weaker [CII]/SFR and higher [OIII]/SFR, or in other words, higher [OIII]/[CII] ratios.

As the Universe evolves and becomes more metal enriched, the typical mass of a galaxy also increases while the luminosity-weighted global LyC escape fractions tend to decrease (e.g. Rosdahl et al. 2018). Certain analytical models predict that f_{esc} evolves very strongly with redshift such that $f_{\text{esc}} \propto (1+z)^{3.4}$ (Haardt & Madau 2012). Harikane et al. (2020) show that for $C_{\text{PDR}} = 0$, [CII] luminosity is decreased by 99% and therefore, the evolving escape fraction can contribute to the high [OIII]/[CII] ratios seen at high redshift. Furthermore, given the correlation between [OIII]/[CII] and f_{esc} , Inoue et al. (2016); Katz et al. (2020b) argued that one can select LyC leakers from non-leakers based on this ratio alone. This metric was tentatively tested (Katz et al. 2020b) on the few low-redshift galaxies where observations of both lines exist and a common trend for the systems that were predicted to be leakers was that many appeared to have strong signatures of feedback, consistent with the results from our simulations.

To demonstrate this relation more explicitly, in Figure 9 we plot the mean value and 1σ scatter of f_{esc} versus $\log_{10}([\text{OIII}]/[\text{CII}])$ for each redshift interval for models where we assume either Solar abundance ratios or core collapse SN abundance ratios. Although there is considerable scatter in the relation, there is a very strong trend that the galaxies that have the highest values of [OIII]/[CII] also exhibit the highest f_{esc} . Assuming core collapse SN abundances systematically




Figure 9. [OIII]/[CII] versus f_{esc} for galaxies in our simulation versus observations. The coloured lines and shaded regions indicate the mean relation and 1σ scatter for different redshift bins. Solid and dashed lines represent the results when we assume Solar abundances and core collapse SN abundances, respectively. We only include galaxies in this plot that have an SFR $> 10^{-2}M_{\odot}\text{yr}^{-1}$.

shifts the relation towards higher values of [OIII]/[CII]; nevertheless, the trend persists.

Consistent with Harikane et al. (2020), we find that lower PDR covering fractions or equivalently increased LyC escape fractions can drive galaxies towards higher [OIII]/[CII].

3.2.2 Higher ISM densities

Galaxies that exhibit the highest ISM densities are expected to exhibit the lowest ratios of [CII]/SFR and [OIII]/SFR because once the density surpasses the critical density of the line, the emissivity increases as n rather than n^2 due to collisional de-excitation. In the top right panel of Figure 8 we show a 2D histogram of [CII]/SFR versus [OIII]/SFR, coloured by the 90th percentile density for gas that emits⁶ [CII] and [OIII]. There is a clear gradient indicating that the galaxies with the lowest values of [CII]/SFR and [OIII]/SFR also have the highest ISM densities.

The critical density of [CII] for interactions with hydrogen is $\sim 2800\text{cm}^{-3}$ while the critical density of [OIII] for interactions with electrons is $\sim 510\text{cm}^{-3}$ (Carilli & Walter 2013). Hence we expect that galaxies with higher ISM densities will exhibit lower ratios of [OIII]/[CII] because as density increases, we reach the critical density of [OIII] before [CII]. In Figure 10, we plot the 90th percentile ISM density of the [OIII] and [CII] emitting gas in the simulated galaxies versus [OIII]/[CII] for the cases where we assume both Solar abundances as well as core collapse SN abundances. For lower values of [OIII]/[CII], we see a decrease in ISM density with decreasing redshift. Consistent with expectations, galaxies with lower ISM densities exhibit higher ratios of [OIII]/[CII].

We do however stress that one should exercise caution when interpreting the trend between ISM density and [OIII]/[CII] for our simulated high-redshift galaxies. While such effects can be easily tested in controlled idealised models such as those presented in Harikane et al. (2020), in cosmological simulations, other parameters are not held




Figure 10. [OIII]/[CII] versus $\log_{10}(\bar{\rho}_{90}/\text{cm}^{-3})$ for galaxies in our simulation versus observations. ρ is defined to be the 90th percentile value from the [CII] and [OIII] emitting gas in the galaxies. The coloured lines indicate the median relation for different redshift bins. Solid and dashed lines represent the results when we assume Solar abundances and core collapse SN abundances, respectively. We only include galaxies in this plot that have an SFR $> 10^{-2}M_{\odot}\text{yr}^{-1}$.




Figure 11. Luminosity-weighted gas density of [CII] regions versus [OIII] regions for galaxies at $z = 6$ coloured by the ratio of [OIII]/[CII]. We only include galaxies that have SFR $> 10^{-2}M_{\odot}\text{yr}^{-1}$ as well as [CII] and [OIII] luminosities $> 10^3L_{\odot}$. [OIII]/[CII] ratios are shown for the Solar abundance model although the results are not fundamentally different for the core collapse SN abundance model.

fixed with the changing ISM density. For example, low density may indicate that the galaxy has been susceptible to strong SN feedback which can destroy the [CII] emitting regions of the galaxy, enhance the amount of O^{++} , and increase f_{esc} , all of which are important for setting the [OIII]/[CII] ratio. Furthermore, it is well established that [OIII] is emitted from different regions of temperature-density phase-space than [CII] and thus the gas that emits [OIII] may be characterised by a different density compared to that which emits [CII].

To demonstrate this, in Figure 11 we plot the luminosity-weighted mean gas density for [CII] against that of [OIII] coloured by the ratio of [OIII]/[CII]. While there is considerable scatter, in general, the

⁶ This gas is defined as having $T < 10^5\text{K}$ and $\rho > 1\text{cm}^{-3}$.

galaxy regions that emit strongly in [OIII] have significantly lower gas densities compared to the regions with high [CII] luminosity. It is often the case that the densities are more than an order of magnitude lower in the [OIII] emitting regions compared to [CII]. This suggests that idealised 1D photoionization models with a fixed ISM density poorly mimic the conditions of [OIII] and [CII] emitting regions in the ISM of high-redshift galaxies. Rather, our simulations are more akin to a constant pressure assumption rather than constant density for 1D CLOUDY models. This is why some in the literature have adopted the constant pressure assumption (see e.g. Nagao et al. 2011; Cormier et al. 2019; Harikane et al. 2020). Since, many of the galaxies with high [OIII]/[CII] ratios do not have ISM densities above the critical density for [OIII] and [CII], despite the large density contrasts between [CII] and [OIII] emitting regions, the higher ISM densities do not play a strong role in setting the [OIII]/[CII] ratio.

3.2.3 Higher ionisation parameters

The photoionization models of Harikane et al. (2020) show that models with higher ionisation parameters ($q = cn_{\gamma, \text{ion}}/n_{\text{H}}$) tend to exhibit higher ratios of [OIII]/[CII]. In our simulations, [CII] is predominantly emitted from higher density neutral regions whereas [OIII] originates in lower density, ionised gas with $T \gtrsim 10^4 \text{K}$. Additional high-energy photons can ionise some of the C^+ into C^{++} which also serves to reduce [CII] emission while additional hard ionising photons have the potential to create more O^{++} thereby increasing the ratio of [OIII]/[CII].

While it is trivial to set the ionisation parameter in a CLOUDY calculation, real galaxies exhibit huge ranges in q depending on location within the galaxy. In order to describe the galaxy by a single value, we select all HII regions defined as gas with $T < 10^{5.5} \text{K}$, $n_{\text{H}} > 1 \text{cm}^{-3}$ and $x_{\text{HII}} > 50\%$, and measure the 90th percentile mass-weighted ionisation parameter. In the bottom right panel of Figure 8 we show a 2D histogram of [CII]/SFR versus [OIII]/SFR coloured by $\log_{10}(\bar{q})$. As predicted by Harikane et al. (2020), we find the gradient in q evolves towards the upper left of the figure (i.e. high ionisation parameter results in lower [CII]/SFR and higher [OIII]/SFR). This can be seen more clearly in Figure 12 where we plot $\log_{10}(\bar{q})$ directly against [OIII]/[CII] for both the Solar abundance and core collapse SN abundance models. Ionisation parameters of galaxies with [OIII]/[CII] ratios of 10 can reach as high as $\log_{10}(\bar{q}) \gtrsim 9$. Therefore the higher [OIII]/[CII] ratios seen at high redshift could be partially due to higher ionisation parameters.

While measurements of the ionisation parameter at $z > 6$ are not yet possible, proxies from lower-redshift may give an indication as to whether reionization epoch systems truly exhibit higher values of q . The O32 diagnostic (i.e. $[\text{OIII}]_{4960\text{\AA}, 5007\text{\AA}}/[\text{OII}]_{3727\text{\AA}, 3729\text{\AA}}$) is often used as a proxy for ionisation parameter (e.g. Penston et al. 1990). Strom et al. (2017) demonstrated using the KBSS survey that star-forming galaxies at $z \sim 2-3$ exhibit considerably higher O32 compared to low-redshift SDSS galaxies, indicative of higher ionisation parameters. Similarly, Katz et al. (2019b) showed that massive high-redshift galaxies in the epoch of reionization have very similar strong-line diagnostics compared to these lower-redshift systems. Consistent results were also found for LyC leakers in the redshift range $z \sim 2-4$ (Nakajima et al. 2020).

In Figure 13, we compare SPHINX²⁰ galaxies with SFRs $> 1 M_{\odot}/\text{yr}$ to different classes of observed galaxies on the ionisation parameter-metallicity plane assuming the core-collapse SN abundance model with enhanced metallicity. The vast majority of star-forming SPHINX²⁰ galaxies exhibit ionisation parameters greater than that of SDSS galaxies, and often equal to or




Figure 12. [OIII]/[CII] versus $\log_{10}(\bar{q})$ for galaxies in our simulation. \bar{q} is defined to be the 90th percentile value from the HII regions in each galaxy. The coloured lines indicate the median relation for different redshift bins. Solid and dashed lines represent the results when we assume Solar abundances and core collapse SN abundances, respectively. We only include galaxies in this plot that have an SFR $> 10^{-2} M_{\odot} \text{yr}^{-1}$.




Figure 13. $\log_{10}(\bar{q})$ versus metallicity for galaxies in our simulation. \bar{q} is defined to be the 90th percentile value from the HII regions in each galaxy whereas metallicity is the mass-weighted mean gas-phase metallicity in the ISM. We only show galaxies with SFR $> 1 M_{\odot}/\text{yr}$ for the core-collapse SN abundance model with enhanced metallicity. Data points with error bars represent ranges in ionisation parameter and metallicity for SDSS galaxies, $z \sim 1$ galaxies, LyC leakers, Lyman-break analogues (LBAs), green pea (GP) galaxies, $z = 2-3$ Lyman-break galaxies (LBGs), and $z = 2-3$ Ly α emitters (LAEs) from Nakajima & Ouchi (2014).

greater than the ionisation parameters in $z = 2-3$ Ly α emitters or known LyC leakers. There is a clear trend that lower metallicity star-forming galaxies exhibit higher ionisation parameters. Furthermore, it is often the lower metallicity, higher ionisation parameter galaxies that exhibit the highest [OIII]/[CII] ratios. However, at fixed q , the [OIII]/[CII] seems to be relatively constant (this is explored further in the next section), hence we argue that q is more important for setting the [OIII]/[CII] ratio compared to Z .

In summary, observational evidence combined with theoretical models suggests that high-redshift galaxies exhibit higher ionisation




Figure 14. $[\text{OIII}]/[\text{CII}]$ versus $\log_{10}(Z/Z_{\odot})$ for galaxies in our simulation. The coloured lines indicate the mean relation for different redshift bins. Solid and dashed lines represent the results when we assume Solar abundances and core collapse SN abundances, respectively. We only include galaxies in this plot that have an $\text{SFR} > 10^{-2} M_{\odot} \text{yr}^{-1}$.

parameters which can contribute to the high ratios of $[\text{OIII}]/[\text{CII}]$. However, in the case of the solar abundance model, all of the observed high-redshift galaxies would need to exhibit ionisation parameters of order $\log(q) \sim 9$ in order to reach the $[\text{OIII}]/[\text{CII}]$ observed. The core collapse SN models can reach the observed ratios with a lower ionisation parameter.

3.2.4 Lower gas metallicities

To first order, the line luminosity of each simulation cell scales with the amount of metals present in the gas. [Harikane et al. \(2020\)](#) argue that for lower metallicity gas, one expects a drop in $[\text{OIII}]/\text{SFR}$ but a constant $[\text{CII}]/\text{SFR}$. Since the bulk of the $[\text{CII}]$ emission arises from PDRs, the PDR column density is proportional to $1/Z$ if dust dominates the self-shielding from UV photons ([Kaufman et al. 2006](#)). For dwarf galaxies with $\text{SFR} \geq 1 M_{\odot} \text{yr}^{-1}$ in the local Universe, [De Looze et al. \(2014\)](#) find an enhancement in $[\text{CII}]/\text{SFR}$ for metal-poor dwarf galaxies compared to other galaxy types.

Similar to ISM density, the $[\text{OIII}]/[\text{CII}]$ ratio is sensitive to numerous different effects that either affect or are impacted by metallicity. For example, lower metallicity stars have harder SEDs which could result in the production of more O^{++} , while the additional ionising photons may ionise some of the PDRs that would be present otherwise. The abundance ratios are expected to change with metallicity and [Arata et al. \(2020\)](#) predicted from simulations that lower metallicity galaxies exhibit higher ratios of $[\text{OIII}]/[\text{CII}]$. The final amount of momentum injected during the SN snowplow phase is weakly sensitive to the metallicity of the gas ([Blondin et al. 1998](#); [Thornton et al. 1998](#); [Karpov et al. 2020](#)) and similarly the ability for gas to cool is also sensitive to metallicity (e.g. [Sutherland & Dopita 1993](#)). Finally, there is a well established mass-metallicity relation in the local Universe ([Tremonti et al. 2004](#)) and total halo mass is partially responsible for setting the maximum densities achievable in our simulation.

In the bottom left panel of Figure 8, we show a 2D histogram of $[\text{CII}]/\text{SFR}$ versus $[\text{OIII}]/\text{SFR}$ for galaxies in our simulation coloured by average gas-phase metallicity in the regions of the galaxies that emit $[\text{OIII}]$ and $[\text{CII}]$. The higher metallicity galaxies evolve towards




Figure 15. Log difference between the $[\text{OIII}]/[\text{CII}]$ ratio for models that include or neglect the impact of the CMB on the $[\text{CII}]$ line as a function of SFR. We only include galaxies that have a $[\text{CII}]$ and $[\text{OIII}]$ luminosity greater than $10^3 L_{\odot}$. At $z = 6$, the $[\text{OIII}]/[\text{CII}]$ ratio only increases by ~ 0.05 dex indicating that it cannot be the dominant factor in the elevated $[\text{OIII}]/[\text{CII}]$ ratios observed at $z \geq 6$

the top right of the diagram, consistent with the idea that to first order, $[\text{OIII}]$ and $[\text{CII}]$ luminosity scale with total metallicity. Note that our CLOUDY calculations do not include dust, but this might be a reasonable approximation for high-redshift galaxies which tend to be metal-poor. Therefore, we do not recover the trend predicted by [Harikane et al. \(2020\)](#). Furthermore, as the abundance ratios in our simulation do not evolve with metallicity, we also do not recover the trend seen in [Arata et al. \(2020\)](#), although we emphasise that their trend is likely more realistic and the evolution is most likely bounded by our two different abundance ratio models. Rather, metallicity in our simulation, in the context of the $[\text{OIII}]/[\text{CII}]$ ratio, is a more sensitive indicator of the ISM physics and star formation history which helps set the emission line luminosities.

In Figure 14, we plot the mean relations of $[\text{OIII}]/[\text{CII}]$ versus $\log_{10}(Z/Z_{\odot})$ as a function of redshift. There is a strong evolution with decreasing redshift towards higher values of metallicity simply because as the simulation evolves, the galaxies become more metal enriched. The trend between $[\text{OIII}]/[\text{CII}]$ and metallicity is essentially flat indicating that for dust-poor galaxies, excluding the impact of evolving abundance ratios with metallicity, in general, the metallicity of the galaxy does not strongly impact the $[\text{OIII}]/[\text{CII}]$ ratio. We do however point out that for the galaxies with high SFRs, we indeed find a trend that lower metallicity galaxies tend to also exhibit higher ionisation parameters and higher $[\text{OIII}]/[\text{CII}]$ ratios (see Figure 13). However, at fixed q , the $[\text{OIII}]/[\text{CII}]$ ratio seems constant. While in general, metallicity may not impact the relation for the majority of galaxies in our simulation, at higher SFRs, since there is a trend between ionisation parameter and metallicity, it may appear that low metallicity correlates with high $[\text{OIII}]/[\text{CII}]$. This behaviour is in general different from what is suggested in [Harikane et al. \(2020\)](#), where it was predicted that the $[\text{OIII}]/[\text{CII}]$ ratio decreases with decreasing metallicity, if q is fixed. It is not yet clear however whether

the impact of dust can be completely excluded as some of the $z > 6$ objects have dust detections (e.g. [Harikane et al. 2020](#)).

3.2.5 Cosmic Microwave Background attenuation

The CMB can have multiple effects on high-redshift infrared line emission because the radiation temperature is significantly higher at $z > 6$ than it is in the local Universe. The CMB acts as both a background against which one has to observe the line and can also act as a heating term which is important for low temperature gas (e.g. [Lagache et al. 2018](#)). While the CMB can impact [OIII], it is particularly important in the context of [CII], especially for lower density, low-temperature gas (see e.g. Figure 1 of [Kohandel et al. 2019](#)). Strong CMB attenuation of the [CII] line can drive higher ratios of [OIII]/[CII] and partially explain the elevated ratio at high-redshift compared to the local Universe.

Both [Laporte et al. \(2019\)](#) and [Harikane et al. \(2020\)](#) considered the impact of CMB attenuation and concluded that the effect was not strong enough to be completely responsible for the elevated [OIII]/[CII] ratios. [Lagache et al. \(2018\)](#) predicted that the maximum increase in [OIII]/[CII] due to CMB attenuation is ~ 0.5 dex. This value is sensitive to both the temperature and density structure of the gas that emits [CII].

To measure the impact of CMB attenuation, we have run an additional set of CLOUDY models for our simulated galaxies at $z = 6$ where we have artificially set the CMB background to be consistent with the value at $z = 0$, thus removing the CMB attenuation effects. In Figure 15, we plot the ratio of [OIII]/[CII] with CMB attenuation to [OIII]/[CII] without CMB attenuation as a function of SFR. We have only considered galaxies that have [CII] and [OIII] luminosities of at least $10^3 L_{\odot}$. For most of the systems at $z = 6$, the enhancement in [OIII]/[CII] is only ~ 0.05 dex and thus we concur with [Laporte et al. \(2019\)](#) and [Harikane et al. \(2020\)](#) that while CMB attenuation enhances the [OIII]/[CII] ratio slightly, it cannot be a dominant factor.

3.2.6 Extended [CII] emission

Using stacks of $z = 5 - 7$ galaxies, [Fujimoto et al. \(2019\)](#) identified large haloes of [CII], extending out to ~ 10 kpc in galaxies with SFRs of $\sim 10 - 70 M_{\odot} \text{yr}^{-1}$. Taking into account this extended emission, [Carniani et al. \(2020\)](#) determined that earlier observations may underestimate the [CII] luminosity by up to a factor of ~ 2 . Simulations have struggled in reproducing large [CII] haloes (e.g. [Katz et al. 2019b](#); [Pallottini et al. 2019a](#); [Arata et al. 2020](#)) such as those observed in [Fujimoto et al. \(2019\)](#). Nevertheless, if the [CII] surface brightness profile is significantly more extended than that of [OIII], the underestimated values [CII] may lead to artificially high values of [OIII]/[CII].

In Figure 16, we plot the [OIII] and [CII] surface brightness profiles for a stack of the three most massive galaxies in our simulation at $z = 6$ for three different viewing angles. These are the galaxies shown in Figure 1 and represent the most similar galaxies in our simulation to those observed in [Fujimoto et al. \(2019\)](#). While the [CII] surface brightness profile is slightly more extended than that of [OIII], consistent with observations from [Carniani et al. \(2020\)](#), the difference is small.

Similar to [Arata et al. \(2020\)](#), there are regions at large radii that are either consistent with or exceed the surface brightness values seen in [Fujimoto et al. \(2019\)](#), although at many locations, we significantly underpredict the flux. Convolution with the ALMA PSF may help




Figure 16. Stacked [CII] and [OIII] surface brightness profiles at $z = 6$ for the three most massive galaxies in the simulation viewed along three different angles compared to an observed stack at $z = 5 - 7$ from [Fujimoto et al. \(2019\)](#). The [OIII] profile is only slightly more concentrated than that of [CII].

relieve some of the tension and smooth out the profile. Part of the extended profile that we see is driven by mergers. In the second row of Figure 1, we can see that the second most massive halo at $z = 6$ is a triple merger of three smaller galaxies. It is not particularly obvious how to choose a centre for stacking the images given the complex morphology and this can also play a role in the shape of the surface brightness profile. In Figure 16, we have opted to use the centre of the halo.

Because the [OIII] surface brightness profile is so similar to that of [CII], it is unlikely that galaxy [OIII] and [CII] morphologies are responsible for the high [OIII]/[CII] ratios observed at high redshift, consistent with the arguments posed in [Harikane et al. \(2020\)](#) because even when accounting for surface brightness dimming in observations ([Carniani et al. 2020](#)), high-redshift galaxies still exhibit high [OIII]/[CII] ratios.

3.2.7 Inclination effects

[Kohandel et al. \(2019\)](#) demonstrated using hydrodynamics simulations that because the shape of the [CII] spectrum changes with viewing angle of a galaxy, for orientations that lead to large velocity spreads (e.g. edge-on) the [CII] flux may not be high enough to be detected by ALMA. In contrast, when the galaxy is viewed face-on, it is more likely to be detected. [Harikane et al. \(2020\)](#) speculate that this effect is unlikely to change the [OIII]/[CII] ratio because if [CII] is undetected due to orientation, [OIII] is also likely to be impacted in a similar way. However, this argument implicitly assumes that [CII] and [OIII] have similarly shaped spectra or that [OIII] is not always more centrally concentrated. [Katz et al. \(2019b\)](#) showed that in certain cases, the first of these two assumptions is incorrect, especially during strong dynamical interactions or feedback events. This once again arises because [OIII] is emitted from a different gas phase than [CII].

In Figure 17 we show the [CII] and [OIII] spectra of the three most massive galaxies at $z = 6$ for three different viewing angles. We have normalised each spectrum so the full width at half maximum (FWHM) can be more easily compared between the lines that have different luminosities. Consistent with [Kohandel et al. \(2019\)](#), we also see strong variations in the spectra as a function of viewing




Figure 17. [CII] (cyan) and [OIII] (magenta) spectra for three different galaxies at $z = 6$ for three different viewing angles (solid, dashed, and dotted lines). The shape of the spectra change depending on the viewing angle and it is not necessarily the case that the [OIII] and [CII] spectra exhibit the same shape. We show the spectra for the Solar abundance model (although the shape will be similar for the core collapse SN abundance model) and the spectra have been smoothed over bins of 20 km/s. We have normalised each spectrum to its maximum value so that the FWHM can be easily compared.

angle as expected. While the [OIII] and [CII] spectra exhibit many similar features, the shapes are not identical. In some cases, the [CII] spectrum is more extended than that of [OIII] (see the solid lines in the top panel) which can lead to a [CII] deficit for this galaxy; however, this is not a generic feature. For example, in the second row, the dotted line peak for [OIII] is nearly 200 km/s offset from the [CII] peak towards negative velocity.

While galaxy orientation may explain [CII] deficits in specific high-redshift galaxies, it cannot explain the high [OIII]/[CII] ratios in all of the observed high-redshift systems, unless there is an observational bias that preferentially selects concentrated [OIII] emission or spatially extended [CII] emission.

3.3 Alternative Explanations for High [OIII]/[CII] Ratios

We have discussed eight different effects that can potentially drive high-redshift galaxies towards high [OIII]/[CII] ratios; however, there is still physics that is not considered in our model that may play a role in setting the [OIII]/[CII] ratios of high redshift galaxies.

At slightly lower redshifts than considered here, it has been shown that the [CII] line is optically thick in some of the brightest submillimeter galaxies (e.g. Neri et al. 2014). We can obtain a rough estimate of [CII] by measuring the optical depth to [CII] in every simulation cell. Following Goldsmith et al. (2012); Lagache et al. (2018), the optical depth (τ) can be written as

$$\tau = \tau_0 \frac{1 - e^{-T_*/T_{\text{ex}}}}{1 + \left(\frac{g_u}{g_l}\right) e^{-T_*/T_{\text{ex}}}}, \quad (1)$$

where $T_* = 91.25\text{K}$ is the equivalent temperature of the transition, $g_u = 4$ and $g_l = 2$ are the statistical weights of the upper and lower

levels, T_{ex} is the excitation temperature of the gas, and

$$\tau_0 = 7.49 \times 10^{-18} N_{\text{C}^+} / \delta v, \quad (2)$$

where N_{C^+} is the CII column density and δv is the line width. The line width is estimated based on the temperature of the cell and the microturbulence of the gas such that $\delta v = \sqrt{a_{\text{therm}}^2 + a_{\text{turb}}^2}$. Here,

$a_{\text{therm}} = \sqrt{\frac{2k_B T}{\mu m_p}}$ where T is the temperature of the cell, k_B is the Boltzmann constant, m_p is the proton mass and μ is the mean weight of $^{12}\text{C}^+$. To estimate the microturbulence, we follow Larson (1981); Franeck et al. (2018) and set $a_{\text{turb}} = 1.1 \times (l/\text{pc})^{0.38}$, where l is the physical length of a cell in the simulation. Based on our CLOUDY models, the fraction of C that is in the form of C^+ is well described as a function of temperature such that

$$f_{\text{C}^+} = \frac{1}{1 + e^{-(147.27 - 35.69 \log_{10}(T/K))}}. \quad (3)$$

N_{C^+} can then be calculated for every cell knowing the density, metallicity, f_{C^+} and cell size. We can measure the excitation temperature in each cell using the collisional deexcitation rate coefficients for neutral hydrogen and electrons as given in Goldsmith et al. (2012) and assuming a CMB background at $z = 6$. To calculate the true extinction, one must then ray trace through the simulation accounting also for the bulk velocities of every gas cell. Rather, to obtain a simple estimate for the effect of self-absorption, we assume that the emission can only be absorbed locally and that nearby cells exhibit similar τ . Under this simple approximation, for the most massive galaxy at $z = 6$, we find that $L_{[\text{CII}]}$ is reduced from 5.24×10^{41} ergs/s to 2.62×10^{41} ergs/s, which is equal to a 50% decrease in total luminosity. We have completed this exercise for the next two most massive galaxies at $z = 6$ and find similar reduction factors. Hence self-absorption can indeed be important for the most massive objects in the epoch of reionization; however, it is unlikely to be fully responsible for the high [OIII]/[CII] ratios seen, especially in the lower mass objects that are less metal enriched and when turbulence is high compared to the thermal velocity.

Another physical effect not considered here is different rates of depletion of C and O onto dust. While C and O are not expected to be heavily depleted like some other elements (e.g. Fe, Jenkins 2009), if for example C is more depleted than O, we would expect that the [OIII]/[CII] ratio would be enhanced. Because our simulated galaxies have low metallicity, we expect the effect of depletion of metals onto dust to be small.

We have hinted earlier that we also cannot rule out observational bias and the preferential selection of galaxies at high-redshift that either exhibit [CII] deficits or enhanced [OIII] emission. While observational bias does not explain why some high-redshift galaxies exhibit exceptionally high [OIII]/[CII] ratios, it may help explain why all of the nine observed high-redshift galaxies with both [OIII] and [CII] detections fall high compared to the local relation. Galaxies are often selected by means of UV dropout. Since UV photons are primarily emitted by young stars, at fixed halo mass, a galaxy is more likely than not to be observed during a burst of star formation. Since both ionising photons from young stars and SN feedback enhance [OIII] emission, we might expect that UV selection biases galaxies towards having high [OIII]/[CII]. Furthermore, galaxies that are more disrupted by feedback are likely to have less dust attenuation which favours brighter magnitudes at fixed SFR. This in turn suggests that galaxies not selected in the UV may not exhibit [CII] deficits. While the sample is still small, we point out that two galaxies at $z > 6$ that were selected based on blue, rest-frame optical colours for ALMA follow-up are more consistent with, or perhaps even above the local




Figure 18. 2D histograms of [CII] versus [OIII] coloured by the mean values of f_{esc} (left), 90th percentile ionisation parameter (centre), and 90th percentile ISM density (right) for galaxies in SPHINX²⁰ at $6 \leq z \leq 10$. The underlaid colour map shows the results from the fitted generalised linear model for each parameter. We have shown the results for the model that assumes the core collapse SN abundances with the metallicity scaled by a factor of 5.72 as this best reproduces the high-redshift [CII]-SFR and [OIII]-SFR relations. For comparison, we show nine $z > 6$ galaxies from Carniani et al. (2020).

[CII]-SFR relation (Smit et al. 2018). We do note that these galaxies may host AGN which have not been explicitly modelled in our work.

Finally, our models only consider stars as the sources of ionising radiation; however, if there are a substantial number of accreting black holes, either low-mass or obscured, they may evade observational detection while providing additional photons and feedback needed to increase [OIII]. Follow-up observations of additional lines, such as [NV] may help constrain this scenario.

4 DISCUSSION

4.1 Predicting Galaxy Properties with [OIII] and [CII]

As was shown in Section 3, the [OIII]/[CII] ratio is sensitive to many galaxy properties including the LyC escape fraction, ionisation parameter, ISM density, SFR, etc. [CII] and [OIII] are already used as SFR indicators (e.g. De Looze et al. 2014) whereas Katz et al. (2020b) showed that the combination of [OIII] and [CII] could be used to identify LyC leakers. Here, we aim to constrain the individual properties of high-redshift galaxies using the correlations found in SPHINX²⁰.

In Figure 18, we show 2D histograms of [CII] versus [OIII] where we have coloured each individual pixel by the mean value of f_{esc} (left), 90th percentile HII region ionisation parameter (centre), and 90th percentile HII region density (right) for simulated galaxies at $6 \leq z \leq 10$. The trends that were discussed in Section 3 are immediately visible. For example, the bias towards higher f_{esc} and q at high [OIII] for fixed [CII]. We show the model for the core collapse SN abundances with the 5.72 times enhanced metallicity as this is the model that best reproduces the [CII]-SFR and [OIII]-SFR relations at high redshift. We have fit relations between each parameter and [CII] and [OIII] using a generalised linear model such that:

$$\log_{10}(f_{\text{esc}}) = \text{Min}[\text{Max}[-0.91 + 0.95 \log_{10}\left(\frac{L_{[\text{OIII}]}}{L_{\odot}}\right) - 1.11 \log_{10}\left(\frac{L_{[\text{CII}]}}{L_{\odot}}\right), -10], 0], \quad (4)$$

$$\log_{10}\left(\frac{q}{\text{cm s}^{-1}}\right) = \text{Min}[\text{Max}[8.66 + 0.27 \log_{10}\left(\frac{L_{[\text{OIII}]}}{L_{\odot}}\right) - 0.29 \log_{10}\left(\frac{L_{[\text{CII}]}}{L_{\odot}}\right), 7.5], 9.5], \quad (5)$$

and

$$\log_{10}\left(\frac{n_{\text{H}}}{\text{cm}^{-3}}\right) = \text{Min}[\text{Max}[1.63 - 0.18 \log_{10}\left(\frac{L_{[\text{OIII}]}}{L_{\odot}}\right) + 0.21 \log_{10}\left(\frac{L_{[\text{CII}]}}{L_{\odot}}\right), 0], 4]. \quad (6)$$

We have defined these relations over fixed intervals for each parameter by the Min and Max functions in each equation. We have underlaid each of these relations in Figure 18 where one can see that the generalised linear models provide an adequate representation of the simulated galaxies.

Also shown in Figure 18 are nine $z > 6$ galaxies with [CII] and [OIII] detections from Carniani et al. (2020). Six of the nine galaxies have [CII] and [OIII] luminosities that fall either outside or very close to the end of the regions of the [CII]-[OIII] plane probed by SPHINX²⁰, while three galaxies, including MACS1149-JD1, the highest redshift galaxy in the sample, fall well within range of the simulation.

In Table 1, we list the values of f_{esc} , q , and n_{H} for each of the nine observed high-redshift galaxies as predicted by our model. Interestingly, these galaxies all have $f_{\text{esc}} \lesssim 5\%$, which is below the luminosity weighted values at all redshifts in SPHINX²⁰ and more consistent with the globally averaged value of f_{esc} (Rosdahl et al *in prep.*). Such values are typical of more massive galaxies at $z > 6$ in our simulations. These nine galaxies all have 90th percentile ionisation parameters with $\log_{10}(q/\text{cm s}^{-1}) > 8.5$ and 90th percentile HII region densities of $\log_{10}(n_{\text{H}}/\text{cm}^{-3}) > 1.5$. The LyC escape fractions scatter below that expected from the relation in Figure 9 due to the mass dependence of this quantity. The relations presented in Section 3 are weighted by number of galaxies rather than luminosity and are thus dominated by lower mass systems compared to what has been observed.

Table 1. Estimated ISM and HII region properties for nine $z > 6$ galaxies. Redshifts, [CII], and [OIII] luminosities are the surface brightness dimming corrected values from Carniani et al. (2020) while f_{esc} , ionisation parameter and density and predicted from our models. Ionisation parameter and density are estimated for HII regions.

Galaxy	Redshift	$L_{[\text{CII}]} / 10^8 L_{\odot}$	$L_{[\text{OIII}]} / 10^8 L_{\odot}$	f_{esc}	$\log_{10}(\bar{q} / \text{cm s}^{-1})$	$\log_{10}\left(\frac{n_{\text{H}}}{\text{cm}^{-3}}\right)$
MACS1149 – JD1	9.11	0.21 ± 0.05	0.88 ± 0.19	0.036 ± 0.016	8.69 ± 0.04	1.69 ± 0.03
A2744 – YD4	8.38	0.28 ± 0.09	0.88 ± 0.21	0.029 ± 0.035	8.66 ± 0.06	1.71 ± 0.04
MACS0416 – Y1	8.31	1.73 ± 0.25	14.0 ± 4.0	0.046 ± 0.015	8.75 ± 0.04	1.66 ± 0.03
SXDF – NB1006 – 2	7.21	3.0 ± 0.7	13.1 ± 2.9	0.024 ± 0.014	8.67 ± 0.04	1.72 ± 0.03
B14 – 65666	7.16	13.9 ± 1.8	39.0 ± 5.0	0.012 ± 0.002	8.61 ± 0.02	1.77 ± 0.02
BDF – 3299	7.11	1.16 ± 0.16	2.13 ± 0.24	0.012 ± 0.002	8.58 ± 0.02	1.77 ± 0.02
J0217 – 0208	6.2	15.6 ± 2.2	89.3 ± 2.1	0.023 ± 0.004	8.69 ± 0.02	1.71 ± 0.01
J0235 – 0532	6.09	5.1 ± 0.8	39.9 ± 3.1	0.038 ± 0.008	8.74 ± 0.02	1.67 ± 0.02
J1211 – 0118	6.03	15.1 ± 1.1	51.0 ± 7.0	0.014 ± 0.002	8.63 ± 0.02	1.75 ± 0.01

Compared to lower redshift galaxies, we find that our estimates for the ionisation parameters of the observed high-redshift galaxies tend to be more typical of $z \sim 2 - 3$ Ly α emitting galaxies rather than local galaxies from SDSS (see Table 3 of Sanders et al. 2015 and Figure 13). We emphasise that our values of q and n_{H} may not be directly comparable to those constrained by observations because of the different methods of measuring ionisation parameter. Our values represent the mass-weighted values of q and n_{H} in HII regions which is represented by gas with $n_{\text{H}} > 1 \text{ cm}^{-3}$, $T < 10^{5.5} \text{ K}$ and $x_{\text{HII}} > 50\%$.

We highlight that estimates of gas density in these high-redshift galaxies are not always consistent between different methods in the literature. For example for MACS1149_JD1, Yang & Lidz (2020) find an upper limit gas density of $\log_{10}(n_{\text{H}}/\text{cm}^3) = 2.4 - 2.8$ or $1.6 - 2.0$ depending on how scatter between HII regions is accounted for, whereas Vallini et al. (2021) estimate a value of $0.88^{+0.55}_{-0.19}$. We find a 90th percentile HII region density of 1.69 for the same galaxy. These density estimates are not all measuring the exact same quantity. The Vallini et al. (2021) values represent the average over the [OIII] and [CII] emitting regions while the Yang & Lidz (2020) values represent the upper limits in HII regions. Our 90th percentile values are perhaps more comparable to the upper limits quoted in Yang & Lidz (2020). Consistent with Vallini et al. (2021), we find that including both HII regions and PDRs in the calculation increases the density estimate.

4.2 The Case for [OI] Observations

Much of the complexity in interpreting ISM properties in real high-redshift galaxies using [OIII] and [CII] is that without knowing the star formation history and metallicity of the system, the different timescales for O and C enrichment can impact the determination of galaxy properties. Values of f_{esc} , q , and ρ are all sensitive to the C/O abundance ratio assumed. One potential way to circumvent this issue is to focus on line ratios that are not as impacted by the exact metal abundance ratios.

The [OI] 63 μm line has many similar properties to [CII] in that it is predominantly emitted by neutral gas since the ionisation energy of OI is 13.6eV (nearly exactly that of HI). [OI] has a very high critical density ($\sim 5 \times 10^5 \text{ cm}^{-3}$) and is known to strongly correlate with star formation in the local Universe (De Looze et al. 2014). Numerical simulations predict that [OI] can outshine [CII] at high redshift (e.g. Olsen et al. 2017; Katz et al. 2019b) and an [OI]/[CII] ratio > 1 was recently detected with ALMA at $z > 6$ (Rybak et al. 2020). [OI] is however impacted by self-absorption (Kraemer et al. 1996; Goldsmith 2019). The primary benefit of observing [OI] is that




Figure 19. 2D histogram of [OI] versus [OIII] coloured by the mean value of f_{esc} for galaxies in SPHINX²⁰ at $6 \leq z \leq 10$. The underlaid colour map shows results from the fitted generalised linear model. We have shown the results for the model that assumes core collapse SN abundances with the metallicity scaled by a factor of 5.72.

the comparison with [OIII] should not be as impacted by different enrichment channels.

In Figure 19 we show a 2D histogram of [OI] versus [OIII] coloured by the mean value of f_{esc} for galaxies in SPHINX²⁰ at $6 \leq z \leq 10$. Similar to [CII], galaxies with higher [OIII] at fixed [OI] exhibit higher escape fractions due to the fact that [OI] originates in predominantly neutral gas whereas [OIII] is emitted by ionised gas. The relation between [OIII], [OI], and f_{esc} is well defined by the generalised linear model:

$$\log_{10}(f_{\text{esc}}) = \text{Min}[\text{Max}[0.50 + 1.01 \log_{10}\left(\frac{L_{[\text{OIII}]}}{L_{\odot}}\right) - 1.16 \log_{10}\left(\frac{L_{[\text{OI}]}}{L_{\odot}}\right), -10], 0]. \quad (7)$$

The coefficients in this equation are nearly identical to the relation between [OIII], [CII], and f_{esc} , while the intercept has increased due to the fact that [OI] is brighter. Given the utility of [OI] as a

supplement or replacement for [CII], we argue that more observation time should be used to constrain [OI] emission from $z > 6$ galaxies.

4.3 Caveats

As with all numerical simulations, there are numerous caveats we must be aware of when interpreting our results. We have modelled the emission lines in post-processing with CLOUDY and our method for setting up the models can introduce systematic uncertainties (see e.g. [Olsen et al. 2018](#)). For example, we have assumed a gas slab being irradiated from one direction whereas one could assume a shell instead. Simulation cells that host star particles are more likely to act like shells whereas gas cells without stars are more similar to slabs. Furthermore, we employ a constant temperature assumption to account for shocks and SN feedback that are present in the simulation rather than allow CLOUDY to compute the photoionisation-equilibrium temperature of the gas. Such an assumption can lead to different line luminosities as [Lupi et al. \(2020\)](#) showed that a variable temperature CLOUDY model can increase the [OIII]/[CII] ratio by a factor of two. One could also choose a different shape for the input SED than used in this work which can impact both the ionisation states of individuals as well as the electron density. Rather than model a different SED for each gas cell, we have fixed the shape of the SED to be consistent with the Milky Way ISM and have normalised the photon count in each photon energy bin to match the simulation to conserve energy. The choice of SED within the cosmological simulation itself is also uncertain and it is well established that depending on the choice, the number of ionising photons as well as the escape fraction can change significantly (e.g. [Ma et al. 2016](#); [Rosdahl et al. 2018](#)). During post-processing, we also modelled the sub-ionising bands in the optically thin limit when in reality, molecules like H_2 or dust particles can absorb such photons and perhaps change the C^+ content of the galaxy. Finally, additional physics such as cosmic rays, magnetic fields, and subgrid turbulence are not included in the calculation, all of which can impact the emergent luminosities.

Our simulations are limited by both spatial and mass resolution which prevents us from modelling the detailed physics of the ISM on scales that we do not resolve. While our resolution is state-of-the-art in the context of full-box radiation hydrodynamics simulations, we do not resolve the formation of individual stars and their individual HII regions/PDRs where [OIII] and [CII] are emitted. Rather, we represent stellar clusters and therefore the structure of the ISM in the simulation may be different than in high-redshift galaxies. Furthermore, because we use a quasi-Lagrangian scheme for refining the AMR grid, most of our resolution tends to be concentrated on regions of high density. The origin of large [CII] haloes at high redshift is currently unknown and one suggestion is that there are small clumps of gas or small satellite galaxies in the outer regions of the haloes that emit [CII] (e.g. [Fujimoto et al. 2019](#)). Simulations that add additional resolution to the CGM (e.g. [Hummels et al. 2019](#); [van de Voort et al. 2019](#)) tend to resolve significantly more of these clumps.

Finally, we have assumed specific subgrid models for star formation and stellar feedback, all of which impact the stellar mass, metal enrichment, and ionising photon production. As we have argued, a more top-heavy stellar IMF which leads to enhanced SN feedback, ionising photon production, and metal enrichment may go a long way to reconciling the high-redshift [CII]-SFR, [OIII]-SFR, and [OIII]/[CII]-SFR relations. Future simulations will be needed to study these effects.

5 CONCLUSIONS

In this work, we have studied the possible origins of high [OIII]/[CII] ratios observed in $z > 6$ galaxies. We have used a new, state-of-the-art, full-box, cosmological radiation hydrodynamics simulation – SPHINX²⁰ – to model [CII] and [OIII] for more than 20,000 simulated star-forming galaxies at $z > 6$, allowing us to disentangle the physics that drives high [OIII]/[CII] ratios. Our main conclusions can be summarised as follows:

- We can only reproduce the observed [CII]-SFR, [OIII]-SFR, and [OIII]/[CII]-SFR relations at $z > 6$ if we assume that high-redshift galaxies are primarily enriched by low-metallicity core collapse SN and that the high-redshift stellar IMF is significantly more top heavy than the Kroupa IMF. This is consistent with expectations for low-metallicity star formation in the early Universe.
- The [CII]-SFR relation naturally converges to the observed values at $z \sim 4.5$ when Solar abundance ratios with a standard IMF are assumed, indicating that both the stellar IMF and enrichment channels evolve significantly as redshift decreases.
- While the reduced C/O ratios at $z > 6$ may be the primary reason why high-redshift galaxies exhibit high [OIII]/[CII] ratios, increased ionisation parameters, increased ISM density, increased LyC escape fractions, and to a small extent CMB attenuation also contribute to the high [OIII]/[CII] ratios. In contrast, lower gas metallicities, extended [CII] emission, and inclination effects are unlikely to explain the behaviour for all of the nine high-redshift galaxies with observations of both [CII] and [OIII].
- [OIII] emitting regions have significantly lower ISM densities than regions that emit [CII] and thus 1D models that consider a fixed gas density must be treated with caution when being used to study the ISM of the early Universe.
- [OIII] and [CII] can be used to constrain ISM properties and LyC escape fractions of high-redshift galaxies. We find that typical escape fractions of the observed systems are $\sim 1\% - 4\%$ while gas densities in the [OIII] and [CII] emitting regions can reach as high as $> 1000\text{cm}^{-3}$.
- [OI] can be used as both an alternative and supplement to [CII] observations in the high-redshift Universe because not only is it predicted to be brighter than [CII], it also originates from similar regions of the ISM meaning that it can be used to constrain similar galaxy characteristics as [CII]. We argue that more observation time should be spent on [OI] observations of $z > 6$ galaxies.

Our model for predicting emission lines from simulations is applicable to numerous other lines across the wavelength spectrum. In future work we aim to compare how the methods for predicting galaxy properties such as metallicity, ionisation parameter, and gas density from strong-line diagnostics translate to the early Universe.

ACKNOWLEDGEMENTS

This work was supported by the Programme National Cosmology et Galaxies (PNCG) of CNRS/INSU with INP and IN2P3, co-funded by CEA and CNES. MGH acknowledges support from the UKRI Science and Technology Facilities Council (grant numbers ST/N000927/1 and ST/S000623/1). Support by ERC Advanced Grant 320596 “The Emergence of Structure during the Epoch of reionization” is gratefully acknowledged. SMA acknowledges support by ERC Starting Grant 638707 “Black holes and their host galaxies: co-evolution across cosmic time”. TK was supported by the National Research Foundation of Korea (NRF-2019K2A9A1A0609137711 and

NRF-2020R1C1C100707911). TG acknowledges support from the European Research Council under grant agreement ERC-stg757258 (TRIPLE). The research of AS and JD is supported by Adrian Beecroft and STFC. Computing time for this work was provided by the Partnership for Advanced Computing in Europe (PRACE) as part of the “First luminous objects and reionization with SPHINX (cont.)” (2016153539, 2018184362, 2019215124) project. We thank Philipp Otte and Filipe Guimaraes for helpful support throughout the project and for the extra storage they provided us. We also thank GENCI for providing additional computing resources under GENCI grant A0070410560. RSE was supported by the European Research Council under the European Union Horizon 2020 research and innovation programme (grant agreement no. 669253).

DATA AVAILABILITY

The data underlying this article will be shared on reasonable request to the corresponding author.

REFERENCES

- Arata S., Yajima H., Nagamine K., Abe M., Khochfar S., 2020, *MNRAS*, **498**, 5541
- Aravena M., et al., 2016, *ApJ*, **833**, 71
- Aubert D., Pichon C., Colombi S., 2004, *MNRAS*, **352**, 376
- Barrow K. S. S., Robertson B. E., Ellis R. S., Nakajima K., Saxena A., Stark D. P., Tang M., 2020, *ApJ*, **902**, L39
- Béthermin M., et al., 2020, *A&A*, **643**, A2
- Blondin J. M., Wright E. B., Borkowski K. J., Reynolds S. P., 1998, *ApJ*, **500**, 342
- Boselli A., Gavazzi G., Lequeux J., Pierini D., 2002, *A&A*, **385**, 454
- Bradač M., et al., 2017, *ApJ*, **836**, L2
- Byler N., Dalcanton J. J., Conroy C., Johnson B. D., 2017, *ApJ*, **840**, 44
- Capak P. L., et al., 2015, *Nature*, **522**, 455
- Carilli C. L., Walter F., 2013, *ARA&A*, **51**, 105
- Carniani S., et al., 2020, *MNRAS*, **499**, 5136
- Chon S., Omukai K., Schneider R., 2021, arXiv e-prints, p. arXiv:2103.04997
- Cooke E. A., et al., 2018, *ApJ*, **861**, 100
- Cormier D., et al., 2010, *A&A*, **518**, L57
- Cormier D., et al., 2015, *A&A*, **578**, A53
- Cormier D., et al., 2019, *A&A*, **626**, A23
- De Looze I., Baes M., Bendo G. J., Cortese L., Fritz J., 2011, *MNRAS*, **416**, 2712
- De Looze I., et al., 2014, *A&A*, **568**, A62
- Eldridge J. J., Izzard R. G., Tout C. A., 2008, *MNRAS*, **384**, 1109
- Ferland G. J., 1996, Hazy, A Brief Introduction to Cloudy 90
- Ferland G. J., et al., 2017, *Rev. Mex. Astron. Astrofis.*, **53**, 385
- Franeck A., et al., 2018, *MNRAS*, **481**, 4277
- Fujimoto S., et al., 2019, *ApJ*, **887**, 107
- Goldsmith P. F., 2019, *ApJ*, **887**, 54
- Goldsmith P. F., Langer W. D., Pineda J. L., Velusamy T., 2012, *ApJS*, **203**, 13
- Grevesse N., Asplund M., Sauval A. J., Scott P., 2010, *Ap&SS*, **328**, 179
- Haardt F., Madau P., 2012, *ApJ*, **746**, 125
- Hahn O., Abel T., 2011, *MNRAS*, **415**, 2101
- Harikane Y., et al., 2020, *ApJ*, **896**, 93
- Hashimoto T., et al., 2018, *Nature*, **557**, 392
- Herrera-Camus R., et al., 2015, *ApJ*, **800**, 1
- Hummels C. B., et al., 2019, *ApJ*, **882**, 156
- Inoue A. K., et al., 2016, *Science*, **352**, 1559
- Jenkins E. B., 2009, *ApJ*, **700**, 1299
- Karpov P. I., Martizzi D., Macias P., Ramirez-Ruiz E., Kolborg A. N., Naiman J. P., 2020, *ApJ*, **896**, 66
- Katz H., Kimm T., Sijacki D., Haehnelt M. G., 2017, *MNRAS*, **468**, 4831
- Katz H., Laporte N., Ellis R. S., Devriendt J., Slyz A., 2019a, *MNRAS*, **484**, 4054
- Katz H., et al., 2019b, *MNRAS*, **487**, 5902
- Katz H., et al., 2020a, *MNRAS*, **494**, 2200
- Katz H., et al., 2020b, *MNRAS*, **498**, 164
- Katz H., et al., 2021, arXiv e-prints, p. arXiv:2101.11624
- Kaufman M. J., Wolfire M. G., Hollenbach D. J., 2006, *ApJ*, **644**, 283
- Kewley L. J., Dopita M. A., 2002, *ApJS*, **142**, 35
- Kimm T., Cen R., Devriendt J., Dubois Y., Slyz A., 2015, *MNRAS*, **451**, 2900
- Kimm T., Katz H., Haehnelt M., Rosdahl J., Devriendt J., Slyz A., 2017, *MNRAS*, **466**, 4826
- Knudsen K. K., Richard J., Kneib J.-P., Jauzac M., Clément B., Drouart G., Egami E., Lindroos L., 2016, *MNRAS*, **462**, L6
- Kohandel M., Pallottini A., Ferrara A., Zanella A., Behrens C., Carniani S., Gallerani S., Vallini L., 2019, *MNRAS*, **487**, 3007
- Kraemer K., Jackson J. M., Lane A. P., 1996, in American Astronomical Society Meeting Abstracts. p. 21.02
- Kroupa P., 2001, *MNRAS*, **322**, 231
- Lagache G., 2018, in Jelić V., van der Hulst T., eds, Vol. 333, Peering towards Cosmic Dawn. pp 228–233 (arXiv:1801.08054), doi:10.1017/S1743921318000558
- Lagache G., Cousin M., Chatzikos M., 2018, *A&A*, **609**, A130
- Laporte N., et al., 2019, *MNRAS*, **487**, L81
- Larson R. B., 1981, *MNRAS*, **194**, 809
- Le Fèvre O., et al., 2020, *A&A*, **643**, A1
- Leung T. K. D., Olsen K. P., Somerville R. S., Davé R., Greve T. R., Hayward C. C., Narayanan D., Popping G., 2020, *ApJ*, **905**, 102
- Lupi A., Volonteri M., Decarli R., Bovino S., Silk J., Bergeron J., 2019, *MNRAS*, **488**, 4004
- Lupi A., Pallottini A., Ferrara A., Bovino S., Carniani S., Vallini L., 2020, *MNRAS*, **496**, 5160
- Ma X., Hopkins P. F., Kasen D., Quataert E., Faucher-Giguère C.-A., Kereš D., Murray N., Strom A., 2016, *MNRAS*, **459**, 3614
- Madden S. C., et al., 2013, *PASP*, **125**, 600
- Maiolino R., Mannucci F., 2019, *A&ARv*, **27**, 3
- Maiolino R., et al., 2015, *MNRAS*, **452**, 54
- Michel-Dansac L., Blaizot J., Garel T., Verhamme A., Kimm T., Trebitsch M., 2020, *A&A*, **635**, A154
- Moriwaki K., et al., 2018, *MNRAS*, **481**, L84
- Nagao T., Maiolino R., Marconi A., Matsuhara H., 2011, *A&A*, **526**, A149
- Nakajima K., Ouchi M., 2014, *MNRAS*, **442**, 900
- Nakajima K., Ellis R. S., Robertson B. E., Tang M., Stark D. P., 2020, *ApJ*, **889**, 161
- Neri R., Downes D., Cox P., Walter F., 2014, *A&A*, **562**, A35
- Nomoto K., Tominaga N., Umeda H., Kobayashi C., Maeda K., 2006, *Nuclear Phys. A*, **777**, 424
- Olsen K., Greve T. R., Narayanan D., Thompson R., Davé R., Niebla Rios L., Stawinski S., 2017, *ApJ*, **846**, 105
- Olsen K., et al., 2018, *Galaxies*, **6**, 100
- Ota K., et al., 2014, *ApJ*, **792**, 34
- Pallottini A., Ferrara A., Gallerani S., Vallini L., Maiolino R., Salvadori S., 2017, *MNRAS*, **465**, 2540
- Pallottini A., et al., 2019a, *MNRAS*, **487**, 1689
- Pallottini A., et al., 2019b, *MNRAS*, **487**, 1689
- Penston M. V., et al., 1990, *A&A*, **236**, 53
- Pentericci L., et al., 2016, *ApJ*, **829**, L11
- Poglitsch A., Krabbe A., Madden S. C., Nikola T., Geis N., Johansson L. E. B., Stacey G. J., Sternberg A., 1995, *ApJ*, **454**, 293
- Popping G., van Kampen E., Decarli R., Spaans M., Somerville R. S., Trager S. C., 2016, *MNRAS*, **461**, 93
- Popping G., Narayanan D., Somerville R. S., Faisst A. L., Krumholz M. R., 2019, *MNRAS*, **482**, 4906
- Prokhorenkova L., Gusev G., Vorobev A., Veronika Dorogush A., Gulín A., 2017, arXiv e-prints, p. arXiv:1706.09516
- Rosdahl J., Teyssier R., 2015, *MNRAS*, **449**, 4380
- Rosdahl J., Blaizot J., Aubert D., Stranex T., Teyssier R., 2013, *MNRAS*, **436**, 2188
- Rosdahl J., et al., 2018, *MNRAS*, **479**, 994

- Rosen A., Bregman J. N., 1995, *ApJ*, **440**, 634
- Rybak M., Zavala J. A., Hodge J. A., Casey C. M., Werf P. v. d., 2020, *ApJ*, **889**, L11
- Safrank-Shrader C., Milosavljević M., Bromm V., 2014, *MNRAS*, **438**, 1669
- Sanders R. L., et al., 2015, *ApJ*, **799**, 138
- Schaerer D., Boone F., Zamojski M., Staguhn J., Dessauges-Zavadsky M., Finkelstein S., Combes F., 2015, *A&A*, **574**, A19
- Schaerer D., et al., 2020, *A&A*, **643**, A3
- Silva M., Santos M. G., Cooray A., Gong Y., 2015, *ApJ*, **806**, 209
- Smit R., et al., 2018, *Nature*, **553**, 178
- Spitzer L., 1978, Physical processes in the interstellar medium, doi:10.1002/9783527617722.
- Stacey G. J., Geis N., Genzel R., Lugten J. B., Poglitsch A., Sternberg A., Townes C. H., 1991, *ApJ*, **373**, 423
- Stanway E. R., Eldridge J. J., 2018, *MNRAS*, **479**, 75
- Stanway E. R., Eldridge J. J., Becker G. D., 2016, *MNRAS*, **456**, 485
- Steidel C. C., Strom A. L., Pettini M., Rudie G. C., Reddy N. A., Trainor R. F., 2016, *ApJ*, **826**, 159
- Strom A. L., Steidel C. C., Rudie G. C., Trainor R. F., Pettini M., Reddy N. A., 2017, *ApJ*, **836**, 164
- Sutherland R. S., Dopita M. A., 1993, *ApJS*, **88**, 253
- Sutherland R., Dopita M., Binette L., Groves B., 2018, MAPPINGS V: Astrophysical plasma modeling code (ascl:1807.005)
- Swinbank A. M., et al., 2012, *MNRAS*, **427**, 1066
- Teyssier R., 2002, *A&A*, **385**, 337
- Thornton K., Gaudlitz M., Janka H. T., Steinmetz M., 1998, *ApJ*, **500**, 95
- Trebtsch M., Blaizot J., Rosdahl J., Devriendt J., Slyz A., 2017, *MNRAS*, **470**, 224
- Tremonti C. A., et al., 2004, *ApJ*, **613**, 898
- Tweed D., Devriendt J., Blaizot J., Colombi S., Slyz A., 2009, *A&A*, **506**, 647
- Uzgil B. D., et al., 2021, arXiv e-prints, p. arXiv:2102.10706
- Vallini L., Gallerani S., Ferrara A., Pallottini A., Yue B., 2015, *ApJ*, **813**, 36
- Vallini L., Ferrara A., Pallottini A., Carniani S., Gallerani S., 2021, *MNRAS*, **505**, 5543
- Veronika Dorogush A., Ershov V., Gulin A., 2018, arXiv e-prints, p. arXiv:1810.11363
- Yang S., Lidz A., 2020, *MNRAS*, **499**, 3417
- Yue B., Ferrara A., 2019, *MNRAS*, **490**, 1928
- van de Voort F., Springel V., Mandelker N., van den Bosch F. C., Pakmor R., 2019, *MNRAS*, **482**, L85

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.