

INAPPROXIMABILITY OF THE INDEPENDENT SET POLYNOMIAL IN THE COMPLEX PLANE*

IVONA BEZÁKOVÁ[†], ANDREAS GALANIS[‡], LESLIE ANN GOLDBERG[‡], AND
DANIEL ŠTEFANKOVIČ[§]

Abstract. We study the complexity of approximating the value of the independent set polynomial $Z_G(\lambda)$ of a graph G with maximum degree Δ when the activity λ is a complex number. When λ is real, the complexity picture is well understood, and is captured by two real-valued thresholds λ^* and λ_c , which depend on Δ and satisfy $0 < \lambda^* < \lambda_c$. It is known that if λ is a real number in the interval $(-\lambda^*, \lambda_c)$ then there is a fully polynomial time approximation scheme (FPTAS) for approximating $Z_G(\lambda)$ on graphs G with maximum degree at most Δ . On the other hand, if λ is a real number outside of the (closed) interval, then approximation is NP-hard. The key to establishing this picture was the interpretation of the thresholds λ^* and λ_c on the Δ -regular tree. The “occupation ratio” of a Δ -regular tree T is the contribution to $Z_T(\lambda)$ from independent sets containing the root of the tree, divided by $Z_T(\lambda)$ itself. This occupation ratio converges to a limit, as the height of the tree grows, if and only if $\lambda \in [-\lambda^*, \lambda_c]$. Unsurprisingly, the case where λ is complex is more challenging. It is known that there is an FPTAS when λ is a complex number with norm at most λ^* and also when λ is in a small strip surrounding the real interval $[0, \lambda_c)$. However, neither of these results is believed to fully capture the truth about when approximation is possible. Peters and Regts identified the complex values of λ for which the occupation ratio of the Δ -regular tree converges. These values carve a cardioid-shaped region Λ_Δ in the complex plane, whose boundary includes the critical points $-\lambda^*$ and λ_c . Motivated by the picture in the real case, they asked whether Λ_Δ marks the true approximability threshold for general complex values λ . Our main result shows that for every λ outside of Λ_Δ , the problem of approximating $Z_G(\lambda)$ on graphs G with maximum degree at most Δ is indeed NP-hard. In fact, when λ is outside of Λ_Δ and is not a positive real number, we give the stronger result that approximating $Z_G(\lambda)$ is actually #P-hard. Further, on the negative real axis, when $\lambda < -\lambda^*$, we show that it is #P-hard to even decide whether $Z_G(\lambda) > 0$, resolving in the affirmative a conjecture of Harvey, Srivastava, and Vondrák. Our proof techniques are based around tools from complex analysis—specifically the study of iterative multivariate rational maps.

Key words. independent set polynomial, inapproximability, #P-hardness

AMS subject classifications. 68Q17, 68Q87

DOI. 10.1137/18M1184485

1. Introduction. The independent set polynomial is one of the most well studied graph polynomials, arising in combinatorics and in computer science. It is also known in statistical physics as the partition function of the hard-core model. This paper studies the computational complexity of evaluating the polynomial approximately

*Received by the editors May 2, 2018; accepted for publication (in revised form) June 29, 2020; published electronically September 24, 2020. A preliminary short version of the manuscript appeared in STOC, ACM, New York, 2018, pp. 1234–1240.

<https://doi.org/10.1137/18M1184485>

Funding: The first author was supported by NSF grant CCF-1319987. The second and third authors were supported by the European Research Council under the European Union’s Seventh Framework Programme (FP7/2007-2013) ERC grant agreement 334828. The paper reflects only the authors’ views and not the views of the ERC or the European Commission. The European Union is not liable for any use that may be made of the information contained therein. The fourth author was supported by NSF grant CCF-1563757.

[†]Department of Computer Science, Rochester Institute of Technology, Rochester, NY 14623 (ib@cs.rit.edu).

[‡]Department of Computer Science, University of Oxford, Wolfson Building, Parks Road, Oxford, OX1 3QD, UK (galanis.an@gmail.com, leslie.goldberg@cs.ox.ac.uk).

[§]Department of Computer Science, University of Rochester, Rochester, NY 14627 (stefanko@cs.rochester.edu).

when a parameter, called the *activity*, is complex. For properties of this polynomial in the complex plane, including connections to the Lovász local lemma, see the work of Scott and Sokal [14]. For $\lambda \in \mathbb{C}$ and a graph G the polynomial is defined as $Z_G(\lambda) := \sum_I \lambda^{|I|}$, where the sum ranges over all independent sets of G . We will be interested in the problem of approximating $Z_G(\lambda)$ when the maximum degree of G is bounded.

When λ is real, the complexity picture is well understood. For $\Delta \geq 3$, let \mathcal{G}_Δ be the set of graphs with maximum degree at most Δ . The complexity of approximating $Z_G(\lambda)$ for $G \in \mathcal{G}_\Delta$ is captured by two real-valued thresholds λ^* and λ_c which depend on Δ and satisfy $0 < \lambda^* < \lambda_c$. To be precise, $\lambda^* = \frac{(\Delta-1)^{\Delta-1}}{\Delta^\Delta}$ and $\lambda_c = \frac{(\Delta-1)^{\Delta-1}}{(\Delta-2)^\Delta}$. The known results are as follows.

1. If λ is in the interval $-\lambda^* < \lambda < \lambda_c$, there is a fully polynomial time approximation scheme (FPTAS) for approximating $Z_G(\lambda)$ on graphs $G \in \mathcal{G}_\Delta$. For $0 \leq \lambda < \lambda_c$, this follows from the work of Weitz [17], while for $-\lambda^* < \lambda < 0$ it follows from the works of Harvey, Srivastava, and Vondrák [8] and Patel and Regts [11].
2. If $\lambda < -\lambda^*$ or $\lambda > \lambda_c$, it is NP-hard to approximate $|Z_G(\lambda)|$ on graphs $G \in \mathcal{G}_\Delta$, even within an exponential factor. For $\lambda > \lambda_c$, this follows from the work of Sly and Sun [15], while for $\lambda < -\lambda^*$ it follows from the work of Galanis, Goldberg, and Štefankovič [6].

The key to establishing this complexity characterization was the following interpretation of the thresholds λ^* and λ_c . Given a Δ -regular tree T of height h with root ρ , let p_h denote the “occupation ratio” of the tree, which is given by $p_h = \frac{\sum_{I: \rho \in I} \lambda^{|I|}}{Z_T(\lambda)}$, where the sum ranges over the independent sets of T that include the root ρ . It turns out that the occupation ratio p_h converges to a limit as $h \rightarrow \infty$ if and only if the activity λ lies within the interval $[-\lambda^*, \lambda_c]$, so it turns out that the complexity of approximating $Z_G(\lambda)$ for $G \in \mathcal{G}_\Delta$ depends on whether this quantity converges.

Understanding the complexity picture in the case where $\lambda \in \mathbb{C}$ is more challenging. If λ is a complex number with norm at most λ^* then there is an FPTAS for approximating $Z_G(\lambda)$ on graphs $G \in \mathcal{G}_\Delta$. This is due to Harvey, Srivastava and Vondrák and to Patel and Regts [8, 11]. More recently, Peters and Regts [12] showed the existence of an FPTAS when λ is in a small strip surrounding the real interval $[0, \lambda_c)$. However, neither of these results is believed to fully capture the truth about when approximation is possible. Motivated by the real case, Peters and Regts [12] identified the values of λ for which the occupation ratio of the Δ -regular tree converges (for $\Delta \geq 3$). These values carve a cardioid-shaped region Λ_Δ in the complex plane, whose boundary includes the critical points $-\lambda^*$ and λ_c . The definition of Λ_Δ is as follows (see Figure 1):¹

$$(1.1) \quad \Lambda_\Delta = \left\{ \lambda \in \mathbb{C} \mid \exists z \in \mathbb{C} : |z| \leq 1/(\Delta-1), \lambda = \frac{z}{(1-z)^\Delta} \right\}.$$

Peters and Regts showed that, for every λ in the (strict) interior of Λ_Δ , the occupation ratio of the Δ -regular tree converges, and asked whether the region Λ_Δ marks the true approximability threshold for general complex values λ .

¹Technically, the word “cardioid” refers to a curve which can be obtained by a point on the perimeter of a circle which is rolling around a fixed circle of the same radius. The region (1.1) does not formally correspond to a cardioid in this sense, but its shape closely resembles a heart for all values of $\Delta \geq 3$, which justifies the (slight) abuse of terminology.




FIG. 1. The cardioid-shaped region Λ_Δ in the complex plane. We show that for all $\lambda \in \mathbb{C} \setminus (\Lambda_\Delta \cup \mathbb{R}_{\geq 0})$, approximating $Z_G(\lambda)$ is $\#P$ -hard. Previously, it was known that the problem is NP -hard on the real line in the intervals $\lambda < -\lambda^*$ and $\lambda > \lambda_c$. Note, we have that the thresholds $-\lambda^*, \lambda_c$ belong to Λ_Δ , by taking $z = \pm 1/(\Delta - 1)$ in (1).

Our main result shows that for every λ outside of the region Λ_Δ , the problem of approximating $Z_G(\lambda)$ on graphs $G \in \mathcal{G}_\Delta$ is indeed NP -hard, thus answering [12, Question 1]. In fact, when λ is outside of Λ_Δ and is not a positive real number, we establish the stronger result that approximating $Z_G(\lambda)$ is actually $\#P$ -hard. We do this by showing that an approximation algorithm for $Z_G(\lambda)$ can be converted into a polynomial-time algorithm for exactly counting independent sets. Further, on the negative real axis, when $\lambda < -\lambda^*$, we show that it is $\#P$ -hard to even decide whether $Z_G(\lambda) > 0$, resolving in the affirmative a conjecture of Harvey, Srivastava, and Vondrák [8, Conjecture 5.1].

We need the following notation to formally state our results. Given a complex number $x \in \mathbb{C}$, we use $|x|$ to denote its norm and $\text{Arg}(x)$ to denote the principal value of its argument in the range $[0, 2\pi)$. We also define $\arg(x) = \{\text{Arg}(x) + 2\pi j \mid j \in \mathbb{Z}\}$. For $y, z \in \mathbb{C}$, we use $d(y, z)$ to denote the Ziv distance between them [18], namely, $d(y, z) = \frac{|y-z|}{\max(|y|, |z|)}$. We denote by $\mathbb{C}_\mathbb{Q}$ the set of complex numbers whose real and imaginary parts are rational numbers (see Definition 3.4).

We consider the problems of multiplicatively approximating the norm of $Z_G(\lambda)$, additively approximating the argument of $Z_G(\lambda)$, and approximating $Z_G(\lambda)$ by producing a complex number \hat{Z} such that the Ziv distance $d(\hat{Z}, Z_G(\lambda))$ is small. We start with the following problem, which captures the approximation of the norm of $Z_G(\lambda)$.

Name $\#BipHardCoreNorm(\lambda, \Delta, K)$.

Instance A bipartite graph G with maximum degree at most Δ .

Output If $|Z_G(\lambda)| = 0$ then the algorithm may output any rational number. Otherwise, it must output a rational number \hat{N} such that $\hat{N}/K \leq |Z_G(\lambda)| \leq K\hat{N}$.

Our first theorem shows that it is $\#P$ -hard to approximate $|Z_G(\lambda)|$ on bipartite graphs of maximum degree Δ within a constant factor.

THEOREM 1.1. *Let $\Delta \geq 3$ and $\lambda \in \mathbb{C}_\mathbb{Q}$ be such that $\lambda \notin (\Lambda_\Delta \cup \mathbb{R}_{\geq 0})$. Then, $\#BipHardCoreNorm(\lambda, \Delta, 1.01)$ is $\#P$ -hard.*

Remark 1.2. The value “1.01” in the statement of Theorem 1.1 is not important. In fact, for any fixed $\epsilon > 0$ we can use the theorem, together with a standard powering argument, to show that it is $\#P$ -hard to approximate $|Z_G(\lambda)|$ within a factor of $2^{n^{1-\epsilon}}$.

The following problem captures the approximation of the argument of $Z_G(\lambda)$.

Name $\# \text{BipHardCoreArg}(\lambda, \Delta, \rho)$.

Instance A bipartite graph G with maximum degree at most Δ .

Output If $Z_G(\lambda) = 0$ then the algorithm may output any rational number. Otherwise, it must output a rational number \hat{A} such that, for some $a \in \arg(Z_G(\lambda))$, $|\hat{A} - a| \leq \rho$.

Our second theorem shows that it is $\#P$ -hard to approximate $\arg(Z_G(\lambda))$ on bipartite graphs of maximum degree Δ within an additive constant $\pi/3$.

THEOREM 1.3. *Let $\Delta \geq 3$ and $\lambda \in \mathbb{C}_{\mathbb{Q}}$ be such that $\lambda \notin (\Lambda_{\Delta} \cup \mathbb{R}_{\geq 0})$. Then, $\# \text{BipHardCoreArg}(\lambda, \Delta, \pi/3)$ is $\#P$ -hard.*

Theorem 1.3 also has the following immediate corollary for the case in which λ is a negative real number, resolving in the affirmative [8, Conjecture 5.1].

COROLLARY 1.4. *Let $\Delta \geq 3$ and $\lambda \in \mathbb{Q}$ be such that $\lambda < -\lambda^*$. Then, given as input a bipartite graph G with maximum degree Δ , it is $\#P$ -hard to decide whether $Z_G(\lambda) > 0$.*

Theorems 1.1 and 1.3 show as a corollary that it is $\#P$ -hard to approximate $Z_G(\lambda)$ within a small Ziv distance.

Name $\# \text{BipComplexHardCore}(\lambda, \Delta)$.

Instance A bipartite graph G with maximum degree at most Δ . A positive integer R , in unary.

Output If $Z_G(\lambda) = 0$ then the algorithm may output any complex number. Otherwise, it must output a complex number z such that $d(z, Z_G(\lambda)) \leq 1/R$.

COROLLARY 1.5. *Let $\Delta \geq 3$ and $\lambda \in \mathbb{C}_{\mathbb{Q}}$ be such that $\lambda \notin (\Lambda_{\Delta} \cup \mathbb{R}_{\geq 0})$. Then, the problem $\# \text{BipComplexHardCore}(\lambda, \Delta)$ is $\#P$ -hard.*

Corollary 1.5 follows immediately from Theorem 1.1 using the fact (see [7, Lemma 2.1]) that $d(z', z) \leq \epsilon$ implies $|z'|/|z| \leq 1/(1 - \epsilon)$. This fact implies (see [7, Lemma 2.2]) that there is a polynomial Turing reduction from $\# \text{BipHardCoreNorm}(\lambda, \Delta, K)$ to $\# \text{BipComplexHardCore}(\lambda, \Delta)$. Similarly, Corollary 1.5 can also be proved using Theorem 1.3. To see this, note that for $\epsilon \leq 1/3$, $d(z', z) \leq \epsilon$ implies that there are $a \in \arg(z)$ and $a' \in \arg(z')$ such that $|a - a'| \leq \sqrt{36\epsilon}/11$. This fact is proved in [7, Lemma 2.1] and it implies (see [7, Lemma 2.2]) that there is a polynomial Turing reduction from $\# \text{BipHardCoreArg}(\lambda, \Delta, \rho)$ to $\# \text{BipComplexHardCore}(\lambda, \Delta)$.

Note that our $\#P$ -hardness results for $\lambda \in \mathbb{C}_{\mathbb{Q}} \setminus (\Lambda_{\Delta} \cup \mathbb{R}_{\geq 0})$ highlight a difference in complexity between this case and the case where λ is a rational satisfying $\lambda > \lambda_c$. If λ is a positive rational then $Z_G(\lambda)$ can be efficiently approximated in polynomial time using an NP oracle, via the bisection technique of Valiant and Vazirani [16]. Thus, in that case approximation is NP-easy, and is unlikely to be $\#P$ -hard. The techniques for proving hardness also differ in the two cases.

1.1. Proof approach. To prove our inapproximability results, we construct graph gadgets which, when appended appropriately to a vertex, have the effect of altering the activity λ to any complex activity λ' that we wish, perhaps with some small error ϵ . In fact, it is essential for our $\#P$ -hardness results to be able to make the error ϵ exponentially small with respect to the number of the vertices in the graph (see the upcoming Proposition 2.2 for details).

Interestingly, our constructions are based on using tools from complex analysis for analyzing the iteration of rational maps. We start with the observation that $(\Delta - 1)$ -ary trees of height h can be used to “implement” activities λ' which correspond to the iterates of the complex rational map $f : x \mapsto \frac{1}{1 + \lambda x^{\Delta-1}}$. Crucially, we show that when $\lambda \notin \Lambda_\Delta$, all of the fixpoints of f are repelling, i.e., applying the map f at any point close to a fixpoint ω will push us away from the fixpoint. In the iteration of univariate complex rational maps, repelling fixpoints belong to the so-called Julia set of the map; a consequence of this is that iterating f in a neighborhood U of a repelling fixpoint gives rise to a chaotic behavior: after sufficiently many iterations, one ends up anywhere in the complex plane.

This sounds promising, but how can we get close to a *repelling* fixpoint of f in the first place? In fact, we need to be able to create arbitrary points in a neighborhood U of a repelling fixpoint and iterating f will not get us anywhere close (since the fixpoint is repelling). The key is to use a Fibonacci-type construction which requires analyzing a more intricate multivariate version of the map f . Surprisingly, we can show that the iterates of the multivariate version converge to the fixpoint ω of the univariate f with the smallest norm. Using convergence properties of the multivariate map around ω (and some extra work), we obtain a family of (univariate) contracting² maps Φ_1, \dots, Φ_t and a small neighborhood U around ω such that $U \subseteq \cup_{i=1}^t \Phi_i(U)$. The final step is to show that “contracting maps that cover yield exponential precision.” To do this we first show that, starting from any point in U , we can apply (some sequence of) Φ_1, \dots, Φ_t at most $\text{poly}(n)$ times to implement any point in U with precision $\exp(-\Omega(n))$. We then show that by iteratively applying the univariate map f and carefully tracking the distortion introduced, we can eventually implement any point in the complex plane with exponentially small error.

Section 2 gives a more detailed description of the proof approach, stating the key lemmas, and explaining how they relate to each other. Section 3 gives some preliminary definitions, which are not necessary for understanding the proof outline in section 2, but are necessary once we start with the proofs. Section 3 also imports some key facts from related works. Section 4 proves some key properties of the cardioid defined by (1.1). The main technical part of the paper is contained in sections 5 and 6. Section 5 shows how to implement activities with exponential precision. Section 6 shows how to obtain our inapproximability results, using these. Some of the technical proofs for section 5 are deferred to section 7.

1.2. New developments. After this paper was written, Bencs and Csikvári [2] discovered a new zero-free region inside the region Λ_Δ . Using the algorithm of Patel and Regts [11], this gives an FPTAS for approximating $Z_G(\lambda)$ on graphs $G \in \mathcal{G}_\Delta$ within this zero-free region. In addition to this, Rivera-Letelier [13] and Buys [4] showed the existence of zeros inside the region Λ_Δ , close to the boundary.

2. Proof outline. In this section, we give a more detailed outline of the proof of our results. We focus mainly on the case where $\lambda \in \mathbb{C} \setminus (\Lambda_\Delta \cup \mathbb{R})$. In section 2.7 we describe suitable modifications that will give us the ingredients needed for negative real values $\lambda \in \mathbb{Q} \setminus \Lambda_\Delta$.

Let $\lambda \in \mathbb{C}$ and $G = (V, E)$ be an arbitrary graph. We denote by \mathcal{I}_G the set of independent sets of G (including the empty independent set). For a vertex $v \in V$, we

²Let $\Phi : \mathbb{C} \rightarrow \mathbb{C}$ be a complex map. We say that Φ is contracting on a set $S \subseteq \mathbb{C}$ if there exists a real number $M < 1$ such that for all $x, y \in S$ it holds that $|\Phi(x) - \Phi(y)| \leq M|x - y|$.

will denote

$$Z_{G,v}^{\text{in}}(\lambda) := \sum_{I \in \mathcal{I}_G; v \in I} \lambda^{|I|}, \quad Z_{G,v}^{\text{out}}(\lambda) := \sum_{I \in \mathcal{I}_G; v \notin I} \lambda^{|I|}.$$

Thus, $Z_{G,v}^{\text{in}}(\lambda)$ is the contribution to the partition function $Z_G(\lambda)$ from those independent sets $I \in \mathcal{I}_G$ such that $v \in I$; similarly, $Z_{G,v}^{\text{out}}(\lambda)$ is the contribution to $Z_G(\lambda)$ from those $I \in \mathcal{I}_G$ such that $v \notin I$.

DEFINITION 2.1. *Fix a complex number λ that is not 0. Given λ , the graph G is said to implement the activity $\lambda' \in \mathbb{C}$ with accuracy $\epsilon > 0$ if there is a vertex v in G such that $Z_{G,v}^{\text{out}}(\lambda) \neq 0$ and*

1. *v has degree one in G , and*
2. *$\left| \frac{Z_{G,v}^{\text{in}}(\lambda)}{Z_{G,v}^{\text{out}}(\lambda)} - \lambda' \right| \leq \epsilon$.*

We call v the terminal of G . If item 2 holds with $\epsilon = 0$, then G is said to implement the activity λ' .

The key to obtaining our #P-hardness results is to show that, given *any* target activity $\lambda' \in \mathbb{C}$, we can construct in polynomial time a bipartite graph G that implements λ' with exponentially small accuracy, as a function of the size of λ' . More precisely, we use $\text{size}(\lambda', \epsilon)$ to denote the number of bits needed to represent the complex number $\lambda' \in \mathbb{C}_{\mathbb{Q}}$ and the rational ϵ (see Definition 3.4). The implementation that we need is captured by the following proposition.

PROPOSITION 2.2. *Let $\Delta \geq 3$ and $\lambda \in \mathbb{C}_{\mathbb{Q}}$ be such that $\lambda \notin \Lambda_{\Delta} \cup \mathbb{R}$. There is an algorithm which, on input $\lambda' \in \mathbb{C}_{\mathbb{Q}}$ and rational $\epsilon > 0$, outputs in $\text{poly}(\text{size}(\lambda', \epsilon))$ time a bipartite graph G of maximum degree at most Δ with terminal v that implements λ' with accuracy ϵ . Moreover, the algorithm outputs the values $Z_{G,v}^{\text{in}}(\lambda), Z_{G,v}^{\text{out}}(\lambda)$.*

Proposition 2.2 is extremely helpful in our reductions since it enables us to construct other gadgets very easily, e.g., equality gadgets that reduce the degree of a graph and gadgets that can turn it into a bipartite graph. The proofs of Theorems 1.1 and 1.3 show how to use these gadgets to obtain #P hardness. In this proof outline, we focus on the most difficult part which is the proof of Proposition 2.2.

To prove Proposition 2.2, we will make use of the following multivariate map:

$$(2.1) \quad (x_1, \dots, x_d) \mapsto \frac{1}{1 + \lambda x_1 \cdots x_d}, \text{ where } d := \Delta - 1.$$

If, starting from 1, there is a sequence of operations (2.1) which ends with the value x , for the purposes of this outline, we will loosely say that “we can generate the value x ” (the notion is formally defined in Definition 7.1). There is a simple correspondence between the values that we can generate and the activities that we can implement: in Lemma 7.2, we show that if we can generate a value x , we can also implement the activity λx using a tree of maximum degree Δ .³

To get some insight about the map (2.1), the first natural step is to look at the univariate case $x_1 = \cdots = x_d = x$, where the map (2.1) simplifies into

$$f : x \mapsto \frac{1}{1 + \lambda x^d}.$$

³Note the extra factor of λ when we pass to the implementation setting which is to ensure the degree requirement in item 1 of Definition 2.1; while the reader should not bother at this stage with this technical detail, the statements of our lemmas are usually about implementing activities and therefore have this extra factor λ incorporated.

Even analyzing the iterates of this map is a surprisingly intricate task; fortunately there is a rich theory concerning the iteration of complex rational maps which we can use (though much less is known in the multivariate setting!). In the next section, we review the basic ingredients of the theory that we need; see [1, 9] for detailed accounts on the subject.

2.1. Iteration of complex rational maps. We will use $\widehat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ to denote the Riemann sphere (complex numbers with infinity). To handle ∞ , it will be convenient to consider the *chordal* metric $d(\cdot, \cdot)$ on the Riemann sphere $\widehat{\mathbb{C}}$, which is given for $z, w \in \mathbb{C}$ by

$$d(z, w) = \frac{2|z - w|}{(1 + |z|^2)^{1/2}(1 + |w|^2)^{1/2}} \text{ and } d(z, \infty) = \lim_{w \rightarrow \infty} d(z, w) = \frac{2}{(1 + |z|^2)^{1/2}}.$$

Note that $d(z, w)$ is bounded by an absolute constant for all $z, w \in \widehat{\mathbb{C}}$.

Let $f : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$ be a complex rational map, i.e., $f(z) = P(z)/Q(z)$ for some coprime polynomials P, Q . We define $f(\infty)$ as the limit of $f(z)$ when $z \rightarrow \infty$. The *degree* of f is the maximum of the degrees of P, Q . A point $p \in \mathbb{C}$ is called a *pole* of f if $Q(p) = 0$; when $p = \infty$, p is a pole of f if 0 is a pole of $f(1/z)$.

Suppose that $z^* \in \mathbb{C}$ is a fixpoint of f , i.e., $f(z^*) = z^*$. The multiplier of f at z^* is given by $q = f'(z^*)$. If $z^* = \infty$, the multiplier of f at z^* is given by $1/f'(\infty)$. Depending on the value of $|q|$, the fixpoint z^* is classified as follows: (i) *attracting* if $|q| < 1$, (ii) *repelling* if $|q| > 1$, and (iii) *neutral* if $|q| = 1$.

For a nonnegative integer $n \geq 0$, we will denote by f^n the n -fold iterate of f (for $n = 0$, we let f^0 be the identity map). Given $z_0 \in \widehat{\mathbb{C}}$, the sequence of points $\{z_n\}$ defined by $z_n = f(z_{n-1}) = f^n(z_0)$ is called the *orbit* of z_0 .

Given a rational map $f : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$, we will be interested in the sensitivity of an orbit under small perturbations of the starting point. A point z_0 belongs to the *Fatou* set if, for every $\epsilon > 0$ there exists $\delta > 0$ such that, for any point z' with $d(z', z_0) \leq \delta$, it holds that $d(f^n(z'), f^n(z_0)) \leq \epsilon$ for all positive integers n (in other words, z_0 belongs to the Fatou set if the family of maps $\{f^n\}_{n \geq 1}$ is equicontinuous at z_0 under the chordal metric). A point z_0 belongs to the *Julia* set if z_0 does not belong to the Fatou set (i.e., the Julia set is the complement of the Fatou set).

LEMMA 2.3 (e.g., [9, Lemma 4.6]). *Every repelling fixpoint belongs to the Julia set.*

For $z \in \widehat{\mathbb{C}}$, the *grand orbit* $[z]$ is the set of points z' whose orbit intersects the orbit of z , i.e., for every $z' \in [z]$, there exist integers $m, n \geq 0$ such that $f^m(z) = f^n(z')$. The exceptional set of the map f is the set of points z whose grand orbit $[z]$ is finite. As we shall see in the upcoming Lemma 3.7, the exceptional set of a rational map f can have at most two points and, in our applications, it will in fact be empty.

For $z_0 \in \mathbb{C}$ and $r > 0$, we use $B(z_0, r)$ to denote the open ball of radius r around z_0 . A set U is a neighborhood of z_0 if U contains a ball $B(z_0, r)$ for some $r > 0$. We will use the following fact.

THEOREM 2.4 (see, e.g., [9, Theorem 4.10]). *Let $f : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$ be a complex rational map with exceptional set E_f . Let z_0 be a point in the Julia set and let U be an arbitrary neighborhood of z_0 . Then, the union of the forward images of U , i.e., the set $\bigcup_{n \geq 0} f^n(U)$, contains $\widehat{\mathbb{C}} \setminus E_f$.*

Peters and Regts [12] used a version of Theorem 2.4 to conclude the existence of trees T and λ 's close to the boundary of Λ_Δ such that $Z_T(\lambda) = 0$. We will use

Theorem 2.4 as a tool to get our #P-hardness results for any λ outside the cardioid Λ_Δ .

2.2. A characterization of the cardioid. To use the tools from the previous section, we will need to analyze the fixpoints of the map $f(z) = \frac{1}{1+\lambda z^{\Delta-1}}$. We denote by $\vartheta\Lambda_\Delta$ the following curve (which is actually the boundary of the region Λ_Δ defined in (1.1)):⁴

$$(2.2) \quad \vartheta\Lambda_\Delta = \left\{ \lambda \in \mathbb{C} \mid \exists z \in \mathbb{C} : |z| = 1/(\Delta - 1), \lambda = \frac{z}{(1-z)^\Delta} \right\}.$$

The following lemma is proved in section 4.

LEMMA 2.5. *Let $\Delta \geq 3$ and consider the map $f(z) = \frac{1}{1+\lambda z^{\Delta-1}}$ for $\lambda \in \mathbb{C}$. Then,*

1. *for all $\lambda \in \Lambda_\Delta \setminus \vartheta\Lambda_\Delta$, f has a unique attractive fixpoint; all other fixpoints are repelling;*
2. *for all $\lambda \in \vartheta\Lambda_\Delta$, f has a unique neutral fixpoint; all other fixpoints are repelling;*
3. *for all $\lambda \notin \Lambda_\Delta$, all of the fixpoints of f are repelling.*

2.3. Applying the theory. We are now in a position to discuss in detail how to apply the tools of section 2.1 and the result of section 2.2. Let $\lambda \in \mathbb{C}_\mathbb{Q} \setminus (\Lambda_\Delta \cup \mathbb{R})$. By Lemma 2.5, all of the fixpoints of the map $f(z) = \frac{1}{1+\lambda z^{\Delta-1}}$ are repelling. By Lemma 2.3, all of the repelling fixpoints belong to the Julia set of the map and, therefore, by applying Theorem 2.4, iteratively applying f to a neighborhood U of a repelling fixpoint gives the entire complex plane. Therefore, if we want to generate an arbitrary complex value $\lambda' \in \mathbb{C}$, it suffices to be able to generate values in a neighborhood U close to a repelling fixpoint of f . Of course, in our setting we will also need to do this efficiently, up to exponential precision. The following proposition is therefore the next important milestone. It formalizes exactly what we need to show in order to be able to prove Proposition 2.2.

PROPOSITION 2.6. *Let $\Delta \geq 3$ and $\lambda \in \mathbb{C}_\mathbb{Q} \setminus \mathbb{R}$, and set $d := \Delta - 1$. Let ω be the fixpoint of $f(x) = \frac{1}{1+\lambda x^d}$ with the smallest norm.⁵ There exists a rational $\rho > 0$ such that the following holds.*

There is a polynomial-time algorithm such that, on input $\lambda' \in B(\lambda\omega, \rho) \cap \mathbb{C}_\mathbb{Q}$ and rational $\epsilon > 0$, outputs a bipartite graph G of maximum degree at most Δ with terminal v that implements λ' with accuracy ϵ . Moreover, the algorithm outputs the values $Z_{G,v}^{\text{in}}(\lambda), Z_{G,v}^{\text{out}}(\lambda)$.

To briefly explain why Proposition 2.2 follows from Proposition 2.6, we first show how to use Proposition 2.6 to implement activities λx^* , where x^* is close to a pole p of f (i.e., a point p which satisfies $1 + \lambda p^d = 0$). For some $r > 0$, let U be the ball $B(\omega, r)$ of radius r around ω . Using Theorem 2.4, we find the first integer value of $N > 0$

⁴The fact that the curve $\vartheta\Lambda_\Delta$, as defined in (2.2), is the boundary of the region Λ_Δ (defined in (1.1)) follows from Lemma 4.1. Lemma 4.1 states that for every $\lambda \in \Lambda_\Delta$, there is a unique $z \in \mathbb{C}$ such that $|z| \leq 1/(\Delta - 1)$ and $\lambda = \frac{z}{(1-z)^\Delta}$. Thus, the function $g(z) = \frac{z}{(1-z)^\Delta}$ is holomorphic and injective on the open disc U given by $|z| < 1/(\Delta - 1)$. By the open mapping theorem, we have that $g(U)$ is an open set. Moreover, $|g(z)| \leq |z|/(1-|z|)^{\Delta-1}$, so $g(U)$ is bounded. Since g extends injectively and continuously to the closed disc $|z| \leq 1/(\Delta - 1)$, the boundary of $g(U)$ is given by the image of the circle $|z| = 1/(\Delta - 1)$. (Alternatively, one can also apply the domain invariance theorem to g .)

⁵Note, by Lemma 4.2, all the fixpoints of f have different norms for $\lambda \in \mathbb{C}_\mathbb{Q} \setminus \mathbb{R}$, so ω is well-defined.

such that a pole p^* belongs to $f^N(U)$; in fact, we can choose r (see Lemma 5.4) so that there exists a radius $r^* > 0$ such that $B(p^*, r^*) \subseteq f^N(U)$. The idea of “waiting till we hit the pole of f ” is that, up to this point, the iterates of f satisfy Lipschitz inequalities, i.e., it can be shown that there exists a real number $L > 0$ such that $|f^N(x_1) - f^N(x_2)| \leq L|x_1 - x_2|$ for all $x_1, x_2 \in U$. Therefore, for any desired target $x^* \in B(p^*, r^*)$ we can find $w^* \in U$ such that $f^N(w^*) = x^*$. We then implement λw^* using Proposition 2.6 with accuracy $\epsilon > 0$. Due to the Lipschitz inequality, this yields an implementation of λx^* with accuracy at most $\lambda L\epsilon$, i.e., just a constant factor distortion. Once we are able to create specified activities close to λp^* , where p^* is a pole we move from there using the recurrence $f(x) = \frac{1}{1+\lambda x^d}$ and this enables us to implement activities λ' with large norm. After that, we use the implementations that we have to implement activities λ' with small norm and, finally, λ' with moderate value of $|\lambda'|$ as well. See the proof of Proposition 2.2 in section 5.3 for more details.

2.4. Chasing repelling fixpoints. In this section, we focus on the proof of Proposition 2.6, whose proof (given in section 7) requires us to delve into the analysis of the multivariate map (restating (2.1))

$$(x_1, \dots, x_d) \mapsto \frac{1}{1 + \lambda x_1 \cdots x_d}, \text{ where } d := \Delta - 1.$$

Recall, in the scope of proving Proposition 2.6, our goal is to generate points close to a repelling fixpoint of the map $f : x \mapsto \frac{1}{1+\lambda x^d}$. Since λ is outside the cardioid region Λ_Δ , the fixpoints of the map f are repelling and therefore we cannot get close to any of them by just iterating f . Can the multivariate map make it easier to get to a fixpoint of f ? The answer to the question is yes, as the following lemma asserts.

LEMMA 2.7. *Let $\Delta \geq 3$ and $\lambda \in \mathbb{C} \setminus \mathbb{R}$, and set $d := \Delta - 1$. Let ω be the fixpoint of $f(x) = \frac{1}{1+\lambda x^d}$ with the smallest norm. For $k \geq 0$, let x_k be the sequence defined by*

$$(2.3) \quad x_0 = x_1 = \cdots = x_{d-1} = 1, \quad x_k = \frac{1}{1 + \lambda \prod_{i=1}^d x_{k-i}} \quad \text{for } k \geq d.$$

Then, the sequence x_k is well-defined (i.e., the denominator of (2.3) is nonzero for all $k \geq d$) and converges to the fixpoint ω as $k \rightarrow \infty$. Moreover, there exist infinitely many k such that $x_k \neq \omega$.

Note, Lemma 2.5 guarantees that the fixpoint ω in Lemma 2.7 is repelling when $\lambda \in \mathbb{C} \setminus (\Lambda_\Delta \cup \mathbb{R})$, so Lemma 2.7 indeed succeeds in getting us close to a repelling fixpoint in this case. It is instructive at this point to note that the sequence in (2.3) corresponds to a Fibonacci-type tree construction T_0, \dots, T_k , where for $k \geq d$ tree T_k consists of a root r with subtrees T_{k-d}, \dots, T_{k-1} rooted at the children of r . The trees T_{k-d}, \dots, T_{k-1} generate the values x_{k-d}, \dots, x_{k-1} , respectively, and the tree T_k generates the value x_k .

A few remarks about the proof of Lemma 2.7 are in order. Analyzing the behavior of multivariate recurrences such as the one in (2.3) is typically an extremely complicated task and the theory for understanding such recurrences appears to be still under development. Fortunately, the recurrence (2.3) can be understood in a surprisingly simple way by using the linear recurrence R_k defined by $R_0 = \cdots = R_d = 1$ and $R_{k+1} = R_k + \lambda R_{k-d}$ for all $k \geq d$, and observing that $x_k = R_k / R_{k+1}$ for all k . By interpreting R_k as the independent set polynomial of a claw-free graph evaluated at $\lambda \in \mathbb{C} \setminus \mathbb{R}$, we obtain using a result of Chudnovsky and Seymour [5] that $R_k \neq 0$. The detailed proof of Lemma 2.7 can be found in section 7.1.

2.5. Exponential precision via contracting maps that cover. Lemma 2.7 resolves the intriguing task of getting close to a repelling fixpoint ω of the univariate map when $\lambda \in \mathbb{C} \setminus (\Lambda_\Delta \cup \mathbb{R})$. But in the context of Proposition 2.6 we need to accomplish far more: we need to be able to generate any point which is in some (small) ball U around the fixpoint ω with exponentially small error ϵ .

To do this, we will focus on a small ball U around ω , i.e., $U = B(\omega, \delta)$ for some sufficiently small $\delta > 0$, and we will examine how the multivariate map (2.1) behaves when $x_1, \dots, x_d \in U$. In particular, we show in Lemma 7.9 that for any choice of $x_1, \dots, x_d \in U$ it holds that

$$(2.4) \quad \frac{1}{1 + \lambda x_1 \cdots x_d} \approx \omega + z((x_1 - \omega) + \cdots + (x_d - \omega)) \text{ for } z \in \mathbb{C} \setminus \mathbb{R} \text{ and } 0 < |z| < 1.$$

To establish (2.4) we choose $z = \omega - 1$ so it is important (see Lemma 7.4) that ω satisfies $0 < |\omega - 1| < 1$. Another important observation is that once we fix $x_1, \dots, x_{d-1} \in B(\omega, \delta)$, the resulting map Φ is contracting with respect to the remaining argument x_d (in the vicinity of ω); see Lemma 7.11 for a more detailed treatment of this contraction.

The observation that Φ is contracting will form the basis of our approach to iteratively reduce the accuracy with which we need to generate points (by going backwards): if we need to generate a desired $x \in U$ with error at most ϵ it suffices to be able to generate $\Phi^{-1}(x)$ with error at most $\epsilon/|z| > \epsilon$, i.e., to generate x with good accuracy, we only need to do the easier task of generating the point $\Phi^{-1}(x)$ with less restrictive accuracy. The only trouble is that, if we use a single map Φ , after a few iterations of the process the preimage $\Phi^{-1}(x)$ will eventually escape U . To address this, note that in the construction of the map Φ above, we had the freedom to choose arbitrary $x_1, \dots, x_{d-1} \in B(\omega, \delta)$. We will make use of this freedom and, in particular, we will use a family of contracting maps Φ_1, \dots, Φ_t (for some large constant t) instead of a single map Φ ; the large number of maps will allow us to guarantee that for all $x \in U$, at least one of the preimages $\Phi_1^{-1}(x), \dots, \Phi_t^{-1}(x)$ belongs in U , i.e., that the images $\Phi_1(U), \dots, \Phi_t(U)$ cover U . We will discuss in section 2.6 how to obtain the maps Φ_1, \dots, Φ_t , but first let us formalize the above into the following lemma, which is the basis of our technique for making the error exponentially small.

LEMMA 2.8. *Let $z_0 \in \mathbb{C}_\mathbb{Q}$, $r > 0$, be a rational, and U be the ball $B(z_0, r)$. Further, suppose that $\lambda'_1, \dots, \lambda'_t \in \mathbb{C}_\mathbb{Q}$ are such that the complex maps $\Phi_i : z \mapsto \frac{1}{1 + \lambda'_i z}$ with $i \in [t]$ satisfy the following:*

1. *for each $i \in [t]$, Φ_i is contracting on the ball U ;*
2. *$U \subseteq \bigcup_{i=1}^t \Phi_i(U)$.*

There is an algorithm which, on input of (i) a starting point $x_0 \in U \cap \mathbb{C}_\mathbb{Q}$, (ii) a target $x \in U \cap \mathbb{C}_\mathbb{Q}$, and (iii) a rational $\epsilon > 0$, outputs in $\text{poly}(\text{size}(x_0, x, \epsilon))$ time a number $\hat{x} \in U \cap \mathbb{C}_\mathbb{Q}$ and a sequence $i_1, i_2, \dots, i_k \in [t]$ such that

$$\hat{x} = \Phi_{i_k}(\Phi_{i_{k-1}}(\cdots \Phi_{i_1}(x_0) \cdots)) \text{ and } |\hat{x} - x| \leq \epsilon.$$

The proof of Lemma 2.8 can be carried out along the lines we sketched above; see section 5.1 for details. In that section, we also pair Lemma 2.8 with a path construction which, given the sequence of indices i_1, \dots, i_k , returns a path of length k that implements $\lambda \hat{x}$ (cf. footnote 3 for the extra factor of λ); see Lemma 5.2 for details.

2.6. Constructing the maps. We next turn to the last missing piece, which is to create the maps Φ_1, \dots, Φ_t which satisfy the hypotheses of Lemma 2.8 in a ball

$U = B(\omega, \delta)$ around the fixpoint ω for some small radius $\delta > 0$ (note, we are free to make δ as small as we wish). The following (standard) notions of “covering” and “density” will be relevant for this section.

DEFINITION 2.9. *Let $U \subseteq \mathbb{C}$. A set $F \subseteq U$ is called an ϵ -covering of U if for every $x \in U$ there exists $y \in F$ such that $|x - y| \leq \epsilon$. A set $F \subseteq U$ is called dense in U if F is an ϵ -covering of U for every $\epsilon > 0$.*

We have already seen in section 2.5 that, for arbitrary $x_1, \dots, x_d \in U$, we have (restating (2.4))

$$\frac{1}{1 + \lambda x_1 \cdots x_d} \approx \omega + z((x_1 - \omega) + \cdots + (x_d - \omega)) \text{ for } z \in \mathbb{C} \setminus \mathbb{R} \text{ and } 0 < |z| < 1.$$

We also discussed that, if we fix arbitrary $x_1, \dots, x_{d-1} \in U$, the resulting map $\Phi(x) = \frac{1}{1 + (\lambda x_1 \cdots x_{d-1})x}$ is contracting in U for all sufficiently small $\delta > 0$, and therefore we can easily take care of the contraction properties that we need (in the context of Lemma 2.8). The more difficult part is to control the preimage of the map Φ . We show in Lemma 7.10 that for $x, x_1, \dots, x_{d-1} \in U$, it holds that

$$\Phi^{-1}(x) = \frac{1}{\lambda x_1 \cdots x_{d-1}} \left(\frac{1}{x} - 1 \right) \approx \omega + \left(\frac{x - \omega}{z} - \sum_{j=1}^{d-1} (x_j - \omega) \right).$$

Therefore to ensure that $\Phi^{-1}(x)$ belongs to $U = B(\omega, \delta)$ we need to ensure that x_1, \dots, x_{d-1} are such that

$$(2.5) \quad \left| \frac{x - \omega}{z} - \sum_{j=1}^{d-1} (x_j - \omega) \right| < \delta/2.$$

Note that by Lemma 2.7 we can generate points arbitrarily close to ω and hence we can make each of $x_2 - \omega, \dots, x_{d-1} - \omega$ so small that they are effectively negligible in (2.5); then, to be able to satisfy (2.5), we need to be able to choose x_1 so that $|(x - \omega)/z - (x_1 - \omega)|$ is small, say less than $\delta/4$. Since $|(x - \omega)/z| \leq \delta/|z|$, the key will therefore be to produce a $(\delta/4)$ -covering of the slightly enlarged ball $B(\omega, \delta/|z|)$. Then, we can take x_1 to be one of the points in the $(\delta/4)$ -covering.

We will in fact show the following slightly more general lemma, which guarantees that we can indeed generate the required points around ω for any desired precision $\epsilon > 0$ provided that we choose δ small enough (and can therefore implement activities around $\lambda\omega$). Note that the lemma can be viewed as a “relaxed” version of Proposition 2.6 with much weaker guarantees.

LEMMA 2.10. *Let $\Delta \geq 3$ and $\lambda \in \mathbb{C}_{\mathbb{Q}} \setminus \mathbb{R}$, and set $d := \Delta - 1$. Let ω be the fixpoint of $f(x) = \frac{1}{1 + \lambda x^d}$ with the smallest norm. For any $\epsilon, \kappa > 0$ there exists a radius $\rho \in (0, \kappa)$ such that the following holds. For every $\lambda' \in B(\lambda\omega, \rho)$, there exists a tree G of maximum degree at most Δ that implements λ' with accuracy $\rho\epsilon$.*

But how can we “populate” the vicinity of ω , i.e., generate a covering of a ball $U = B(\omega, \delta)$? Lemma 2.7 only shows that we can generate points arbitrarily close to ω . The key once again is to use the multivariate map around ω and, in particular, the perturbation estimate in the right-hand side (r.h.s.) of (2.4). To focus on the displacement from ω , we will use the transformation $a_i = x_i - \omega$ so that (2.4) translates into the following operation

$$(a_1, \dots, a_d) \mapsto z(a_1 + \cdots + a_d),$$

i.e., if we have generated points which are displaced by a_1, \dots, a_d from ω , we can also generate a point which is roughly displaced by $z(a_1 + \dots + a_d)$ from ω ; we will only need to apply the operation a finite number of times, so the error coming from (2.4) will not matter critically and can be ignored in the following. We show in Lemma 7.8 that, using a sequence of such operations, we can generate points of the form $\omega + z^{N(p)}p(z)$, where p is an arbitrary polynomial with nonnegative integer coefficients and $N(p)$ is a positive integer which is determined by the number of operations we used to create p . We further show in Lemma 7.5, that for all $z \in \mathbb{C} \setminus \mathbb{R}$ with $|z| < 1$, the values $p(z)$, as p ranges over all polynomials with nonnegative integer coefficients, form a dense set of \mathbb{C} . Therefore, to obtain Lemma 2.10, we can choose an ϵ -covering F of the unit disc using a finite set of values $p(z)$ and set $\delta = z^N$, where $N = \max_{p \in F} N(p)$; then, we can generate the points $\omega + \delta p(z)$ for every $p \in F$, which form an $(\epsilon\delta)$ -covering of the ball $U = B(\omega, \delta)$, yielding Lemma 2.10. The full proof is in section 7.2.

2.7. Fitting the pieces together and proof for the real case. We briefly summarize the proof of Proposition 2.6. First, we get points close to a repelling fixpoint by showing Lemma 2.7 (discussed in section 2.4 and proved in section 7.1). Then, we bootstrap this into a moderately dense set of points around the fixpoint, yielding Lemma 2.10 (discussed in section 2.6 and proved in section 7.2). Further, we bootstrap this into exponential precision around the fixpoint using Lemma 2.8 (discussed in section 2.5 and proved in section 7.3). Finally, we propagate this exponential precision to the whole complex plane, therefore yielding Proposition 2.6 (discussed in section 2.3 and proved in section 5.3).

Finally, we mention the modifications needed for the real case when $\lambda < -\lambda^*$. The following proposition is the analogue of Proposition 2.2 and allows us to implement real activities with exponential precision.

PROPOSITION 2.11. *Let $\Delta \geq 3$ and $\lambda \in \mathbb{Q}$ be such that $\lambda < -\lambda^*$. There is an algorithm which, on input $\lambda', \epsilon \in \mathbb{Q}$ with $\epsilon > 0$, outputs in $\text{poly}(\text{size}(\lambda', \epsilon))$ time a bipartite graph G of maximum degree at most Δ with terminal v that implements λ' with accuracy ϵ . Moreover, the algorithm outputs the values $Z_{G,v}^{\text{in}}(\lambda), Z_{G,v}^{\text{out}}(\lambda)$.*

As in the complex case, we will need a moderately dense set of activities to get started, i.e., an analogue of Lemma 2.10; here, our job is somewhat simplified (relative to the case where $\lambda \in \mathbb{C} \setminus \mathbb{R}$) since we can use the following result of [6].

LEMMA 2.12 (see [6, Lemma 4]). *Let $\Delta \geq 3$ and $\lambda < -\lambda^*$. Then, for every $\lambda' \in \mathbb{R}$, for every $\epsilon > 0$, there exists a bipartite graph G of maximum degree at most Δ that implements λ' with accuracy ϵ .*

Note that Lemma 2.12 does not control the size of the graph G with respect to the accuracy ϵ , so it does not suffice to prove Proposition 2.11 on its own. In order to do this, we use the “contracting maps that cover” technique to get the exponential precision, i.e., the analogue of Lemma 2.8 restricted to the reals (see Lemma 5.1). The proof of Proposition 2.11 is completed in section 5.2.

Once the proofs of Propositions 2.2 and 2.11 are in place, we give the proofs of our #P-hardness results in section 6.

2.8. Dependencies between lemmas. The proofs of some theorems, propositions, and lemmas depend directly upon other theorems, propositions, and lemmas that are proved in this paper. To help the reader keep track of this, we provide Table 1. Note that Theorems 1.1 and 1.3 follow directly from Theorems 6.6 and 6.8.

TABLE 1
Dependencies between lemmas, etc.

Result	Depends directly on
Proposition 2.2	Proposition 2.6, Lemma 5.4
Lemma 2.5	Lemma 4.1
Proposition 2.6	Lemmas 2.8, 5.2, 7.12
Lemma 2.7	Lemmas 4.2, 7.3
Lemma 2.10	Lemmas 2.7, 7.2, 7.4, 7.5, 7.8, 7.9
Proposition 2.11	Lemmas 5.1, 5.2, 5.3
Lemma 5.4	Lemma 2.5
Lemma 6.3	Lemmas 6.1, 6.2
Theorem 6.4	Propositions 2.2, 2.11, Lemma 6.3
Theorem 6.6	Theorem 6.4
Theorem 6.8	Propositions 2.2, 2.11, Lemma 6.7
Lemma 7.4	Lemma 4.2
Lemma 7.11	Lemmas 7.4, 7.9
Lemma 7.12	Lemmas 2.10, 7.4, 7.9, 7.10, 7.11

3. Preliminaries.

3.1. Implementing activities. We recall the following definitions from [6], which we modify (slightly) here to account for complex activities.

DEFINITION 3.1. Let $\Delta \geq 2$ be an integer and $\lambda \in \mathbb{C}_{\neq 0}$. We say that (Δ, λ) implements the activity $\lambda' \in \mathbb{C}$ if there is a bipartite graph G of maximum degree at most Δ which implements the activity λ' . More generally, we say that (Δ, λ) implements a set of activities $S \subseteq \mathbb{C}$ if for every $\lambda' \in S$ it holds that (Δ, λ) implements λ' .

Implementing activities allows us to modify the activity at a particular vertex v . As in [6], it will therefore be useful to consider the hard-core model with nonuniform activities. Let $G = (V, E)$ be a graph and $\lambda = \{\lambda_v\}_{v \in V}$ be a complex vector, so that λ_v is the activity of the vertex $v \in V$. The hard-core partition function with activity vector λ is defined as

$$Z_G(\lambda) = \sum_{I \in \mathcal{I}_G} \prod_{v \in I} \lambda_v.$$

Note that by setting all vertex activities equal to λ we obtain the standard hard-core model with activity λ . For a vertex $v \in V$, we define $Z_G^{\text{in}}(\lambda)$ and $Z_G^{\text{out}}(\lambda)$ for the non-uniform model analogously to $Z_G^{\text{in}}(\lambda)$ and $Z_G^{\text{out}}(\lambda)$ for the uniform model, respectively.

The following lemma is proved in [6] for real values but the proof holds verbatim in the complex setting as well. The lemma connects the partition function $Z_G(\lambda)$ of a graph G with nonuniform activities to the partition function $Z_{G'}(\lambda)$ of an augmented graph G' with uniform activity λ (the augmented graph G' is obtained by sticking on each vertex v of G a graph G_v which implements the activity λ_v).

LEMMA 3.2 (see [6, Lemma 5]). Let $\lambda \in \mathbb{C}_{\neq 0}$, let $t \geq 1$ be an arbitrary integer, and let $\lambda'_1, \dots, \lambda'_t \in \mathbb{C}$. Suppose that, for $j \in [t]$, the graph G_j with terminal v_j implements the activity λ'_j , and let $C_j := Z_{G_j, v_j}^{\text{out}}(\lambda)$. Then, the following holds for every graph $G = (V, E)$ and every activity vector $\lambda = \{\lambda_v\}_{v \in V}$ such that $\lambda_v \in \{\lambda, \lambda'_1, \dots, \lambda'_t\}$ for every $v \in V$.

For $j \in [t]$, let $V_j := \{v \in V \mid \lambda_v = \lambda'_j\}$. Consider the graph G' obtained from G by attaching, for every $j \in [t]$ and every vertex $v \in V_j$, a copy of the graph G_j to the vertex v and identifying the terminal v_j with the vertex v . Then, for $C := \prod_{j=1}^t C_j^{|V_j|}$,

it holds that

$$(3.1) \quad Z_{G'}(\lambda) = C \cdot Z_G(\lambda),$$

$$(3.2) \quad \forall v \in V: Z_{G',v}^{\text{in}}(\lambda) = C \cdot Z_{G,v}^{\text{in}}(\lambda), \quad Z_{G',v}^{\text{out}}(\lambda) = C \cdot Z_{G,v}^{\text{out}}(\lambda).$$

Remark 3.3. As noted in [6, Remark 6], the construction of G' in Lemma 3.2 ensures that the degree of every vertex v in G with $\lambda_v = \lambda$ maintains its degree, while the degree of every other vertex v in G gets increased by one. Also, if the graph G is bipartite and the graphs G_j are bipartite for all $j = 1, \dots, t$, then G' is bipartite as well.

These observations will ensure in later applications of Lemma 3.2 that we do not blow up the degree and that we preserve the bipartiteness of the underlying graph G .

3.2. Finding roots of polynomials. To prove our $\#P$ -hardness results, we will sometimes need in our reductions to compute with accuracy ϵ roots of polynomial equations with coefficients in $\mathbb{C}_{\mathbb{Q}}$. We will therefore need some basic results that these procedures can be carried out in polynomial time. To formalize the running time, we will use the following definition for the size of a number in $\mathbb{C}_{\mathbb{Q}}$.

DEFINITION 3.4. Let $\mathbb{C}_{\mathbb{Q}}$ be the set of complex numbers whose real and imaginary parts are rationals. Let $\alpha \in \mathbb{C}_{\mathbb{Q}}$ and write $\alpha = \frac{a}{b} + i\frac{c}{d}$, where a, b, c, d are integers such that $\gcd(a, b) = 1$, $\gcd(c, d) = 1$. Then, the size of α , denoted by $\text{size}(\alpha)$, is given by $1 + \log(|a| + |b| + |c| + |d|)$.

For $\alpha_1, \dots, \alpha_t \in \mathbb{C}_{\mathbb{Q}}$, $\text{size}(\alpha_1, \dots, \alpha_t)$ denotes the total of the sizes of $\alpha_1, \dots, \alpha_t$.

We will need the following fact for finding roots of polynomials with complex coefficients.

LEMMA 3.5 (see, e.g., [10]). *There is an algorithm which, on input of a coefficient list $c_0, \dots, c_n \in \mathbb{C}_{\mathbb{Q}}$ with $c_n \neq 0$ and a rational $\epsilon > 0$, outputs in time $\text{poly}(n, \text{size}(c_0, \dots, c_n, \epsilon))$ numbers $\hat{\rho}_1, \dots, \hat{\rho}_n \in \mathbb{C}_{\mathbb{Q}}$ such that*

$$|\rho_1 - \hat{\rho}_1|, \dots, |\rho_n - \hat{\rho}_n| \leq \epsilon,$$

where ρ_1, \dots, ρ_n are the roots of the polynomial $P(x) = \sum_{i=0}^n c_i x^i$.

3.3. Lower bounds on polynomials evaluated at algebraic numbers. Let $P(x) = \sum_{i=0}^n a_i x^i$ be a polynomial with complex coefficients. The (naive) height of $P(x)$ is defined as $H(P) = \max_i \{|a_i|\}$. In transcendental number theory, the height of a polynomial in x is used to give a lower bound on its value when x is an algebraic number. The simple version of the lower bound that we use (Lemma 3.6 below) is from [7, Lemma 6.3] but the proof is entirely standard, and the proof given in [7] is taken from the proof of Theorem A.1 of Bugeaud's book [3].

Recall, that the minimal polynomial for an algebraic number is the monic polynomial of minimum degree and rational coefficients which has α as a root; the degree of an algebraic number is the degree of its minimal polynomial.

LEMMA 3.6 (Liouville's inequality). *Let $P(x)$ be an integer polynomial of degree n and $y \in \mathbb{C}$ be an algebraic number of degree d . Then either $p(y) = 0$ or $|P(y)| \geq c_y^{-n} (n+1) H(P)^{-d+1}$, where $c_y > 1$ is an effectively computable constant that only depends on y .*

3.4. Characterizing the exceptional set. We conclude this section with the following characterization of the exceptional set (cf. Theorem 2.4).

LEMMA 3.7 (see [9, Lemma 4.9] and [1, Theorem 4.1.2]). *Let $f : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$ be a complex rational map of degree ≥ 2 and let E_f denote its exceptional set. Then, $|E_f| \leq 2$. Moreover,*

- *if $E_f = \{\zeta\}$, then ζ is a fixpoint of f with multiplier 0.*
- *if $E_f = \{\zeta_1, \zeta_2\}$, then ζ_1, ζ_2 are fixpoints of f^2 with multiplier 0.*

4. Proving the properties of the cardioid. In this section, we will prove Lemma 2.5 which classifies the fixpoints of the map $f(z) = \frac{1}{1+\lambda z^{\Delta-1}}$ depending on the value of λ .

We start by proving the following property of the region Λ_Δ defined in (1.1).

LEMMA 4.1. *Let $\Delta \geq 3$ be an integer. Then, for every $\lambda \in \Lambda_\Delta$, there is a unique $z \in \mathbb{C}$ such that $|z| \leq 1/(\Delta-1)$ and $\lambda = \frac{z}{(1-z)^\Delta}$.*

Proof. Existence of $z \in \mathbb{C}$ with the required properties is immediate by the definition of Λ_Δ .

To show uniqueness, set $d := \Delta - 1$, and assume for the sake of contradiction that there exist $x, y \in \mathbb{C}$ with $x \neq y$ such that $|x|, |y| \leq 1/d$ and $\lambda = \frac{x}{(1-x)^{d+1}} = \frac{y}{(1-y)^{d+1}}$. This gives

$$x(1-y)^{d+1} - y(1-x)^{d+1} = 0.$$

By expanding terms we obtain that

$$x(1-y)^{d+1} - y(1-x)^{d+1} = (x-y) - \sum_{k=2}^{d+1} \binom{d+1}{k} (-1)^k xy (x^{k-1} - y^{k-1}).$$

Since $x \neq y$ and $x(1-y)^{d+1} - y(1-x)^{d+1} = 0$, we can factor out $(x-y)$ to obtain that

$$M = 1, \text{ where } M := \sum_{k=2}^{d+1} \binom{d+1}{k} (-1)^k xy \sum_{j=0}^{k-2} x^j y^{(k-2)-j}.$$

We will first show that $|M| = 1$ only if x, y are conjugate complex numbers satisfying $|x| = |y| = 1/d$. Then, we will bootstrap the argument to show that $|M| = 1$ further implies that $x, y \in \mathbb{R}$. Thus, we will get that $x = y$, contradicting that $x \neq y$.

The main observation is that since $|x|, |y| \leq 1/d$, by the triangle inequality we have that

$$\begin{aligned} 1 = |M| &= \left| \sum_{k=2}^{d+1} \binom{d+1}{k} (-1)^k xy \sum_{j=0}^{k-2} x^j y^{(k-2)-j} \right| \leq \sum_{k=2}^{d+1} \binom{d+1}{k} |x| |y| \sum_{j=0}^{k-2} |x|^j |y|^{k-2-j} \\ (4.1) \quad &\leq \sum_{k=2}^{d+1} \binom{d+1}{k} \frac{k-1}{d^k} = 1, \end{aligned}$$

where the last equality follows from subtracting the equalities

$$\begin{aligned} \sum_{k=2}^{d+1} \binom{d+1}{k} \frac{k}{d^k} &= \sum_{k=2}^{d+1} \binom{d}{k-1} \frac{d+1}{d^k} = \frac{d+1}{d} \sum_{k=1}^d \binom{d}{k} \frac{1}{d^k} = \left(1 + \frac{1}{d}\right)^{d+1} - \left(1 + \frac{1}{d}\right), \\ \sum_{k=2}^{d+1} \binom{d+1}{k} \frac{1}{d^k} &= \left(\frac{d+1}{d}\right)^{d+1} - 1 - \frac{d+1}{d}. \end{aligned}$$

It follows that the inequality in (4.1) must hold at equality, from where we obtain that $|x| = |y| = 1/d$. Further, since $\frac{x}{(1-x)^{d+1}} = \frac{y}{(1-y)^{d+1}}$, we obtain that $|1-x| = |1-y|$.

Now, since $x \neq y$, from $|x| = |y| = 1/d$ and $|1 - x| = |1 - y|$, we obtain that x and y are conjugate complex numbers with $|x| = |y| = 1/d$.

Since $x \neq y$ and x, y are conjugates, the imaginary parts of both x and y must be nonzero. Without loss of generality (w.l.o.g.), we may assume that the imaginary part of x is positive, so we can set $x = \frac{1}{d}e^{i\theta}$ and $y = \frac{1}{d}e^{-i\theta}$ for some $\theta \in (0, \pi)$. For each $k = 2, \dots, d+1$, let $M_k := xy \sum_{j=0}^{k-2} x^j y^{(k-2)-j}$. Observe that M_k is a real number (since it equals its conjugate) satisfying $|M_k| \leq (k-1)/d^k$. In fact, we have that

$$M_k = xy \frac{x^{k-1} - y^{k-1}}{x - y} = \frac{\sin((k-1)\theta)}{d^k \sin \theta}$$

for each $k = 2, \dots, d+1$ and in particular $M_2 = 1/d^2$. Using the triangle inequality again, we have that

$$1 = |M| = \left| \sum_{k=2}^{d+1} \binom{d+1}{k} (-1)^k M_k \right| \leq \sum_{k=2}^{d+1} \binom{d+1}{k} \frac{k-1}{d^k} = 1,$$

so the inequality must hold at equality. For this to happen, and since the M_k 's are real numbers, it must be the case that there exists $s \in \{0, 1\}$ so that $M_k = (-1)^{k+s}(k-1)/d^k$ for each $k = 2, \dots, d+1$. From $M_2 = 1/d^2$, we obtain that $s = 0$. Note that $d \geq 2$, so $M_3 = -2/d^3$ yields that $\sin(2\theta) = -2\sin(\theta)$. By the identity $\sin(2\theta) = 2\sin\theta\cos\theta$, we obtain that at least one of $\sin\theta = 0$ or $\cos\theta = -1$ must hold, which contradicts that $\theta \in (0, \pi)$.

Thus, it must be the case that $x = y$, concluding the proof of Lemma 4.1. \square

We are now ready to prove Lemma 2.5, which we restate here for convenience.

LEMMA 2.5. Let $\Delta \geq 3$ and consider the map $f(z) = \frac{1}{1+\lambda z^{\Delta-1}}$ for $\lambda \in \mathbb{C}$. Then,

1. for all $\lambda \in \Lambda_\Delta \setminus \vartheta\Lambda_\Delta$, f has a unique attractive fixpoint; all other fixpoints are repelling;
2. for all $\lambda \in \vartheta\Lambda_\Delta$, f has a unique neutral fixpoint; all other fixpoints are repelling;
3. for all $\lambda \notin \Lambda_\Delta$, all of the fixpoints of f are repelling.

Proof. Let $\omega \in \mathbb{C}$ be an arbitrary fixpoint of f so that

$$(4.2) \quad \omega = \frac{1}{1 + \lambda\omega^{\Delta-1}} \text{ or, equivalently, } \lambda\omega^\Delta = 1 - \omega.$$

Let $q := f'(\omega)$ be the multiplier of f at $z = \omega$. We have that

$$(4.3) \quad q = f'(\omega) = -\frac{(\Delta-1)\lambda\omega^{\Delta-2}}{(1 + \lambda\omega^{\Delta-1})^2} = -(\Delta-1)\lambda\omega^\Delta = -(\Delta-1)(1-\omega),$$

where in the latter two equalities we used (4.2). Let $\hat{\omega} := 1 - \omega$. Then, (4.2) and (4.3) give

$$(4.4) \quad \lambda = \frac{\hat{\omega}}{(1 - \hat{\omega})^\Delta}, \quad |q| = (\Delta-1)|\hat{\omega}|.$$

We are now ready to prove the lemma. Let $\omega_1, \dots, \omega_t$ denote the distinct fixpoints of f (note that $1 \leq t \leq \Delta$) and for $i \in [t]$, let q_i be the multiplier of f at ω_i . Further, let $\hat{\omega}_i = 1 - \omega_i$. Then, (4.4) gives that for all $i \in [t]$ it holds that

$$(4.5) \quad \lambda = \frac{\hat{\omega}_i}{(1 - \hat{\omega}_i)^\Delta}, \quad |q_i| = (\Delta-1)|\hat{\omega}_i|.$$

For $\lambda \in \Lambda_\Delta \setminus \vartheta\Lambda_\Delta$, there exists $j \in [t]$ such that $|\widehat{\omega}_j| < 1/(\Delta - 1)$ (by the definition of the regions Λ_Δ and $\vartheta\Lambda_\Delta$) and hence $|q_j| < 1$. By Lemma 4.1, for all $i \in [t]$ with $i \neq j$ it holds that $|\widehat{\omega}_i| > 1/(\Delta - 1)$ and hence $|q_i| > 1$. This proves item 1.

For $\lambda \in \vartheta\Lambda_\Delta$, there exists $j \in [t]$ such that $|\widehat{\omega}_j| = 1/(\Delta - 1)$ (by the definition of the region $\vartheta\Lambda_\Delta$) and hence $|q_j| = 1$. Again, by Lemma 4.1, for all $i \in [t]$ with $i \neq j$ it holds that $|\widehat{\omega}_i| > 1/(\Delta - 1)$ and hence $|q_i| > 1$. This proves item 2.

For $\lambda \notin \Lambda_\Delta$, we have that every $\widehat{\omega}_i$ satisfies $|\widehat{\omega}_i| > 1/(\Delta - 1)$ (by the definition of the region Λ_Δ) and hence $|q_i| > 1$. This proves item 3.

This concludes the proof of Lemma 2.5. \square

We close this section with the following lemma that applies to all $\lambda \in \mathbb{C} \setminus \mathbb{R}$.

LEMMA 4.2. *Let $\lambda \in \mathbb{C} \setminus \mathbb{R}$ and $\Delta \geq 3$. Then, the fixpoints of the map $f(z) = \frac{1}{1+\lambda z^{\Delta-1}}$ have (pairwise) distinct norms.*

Proof. For convenience, let $d := \Delta - 1$. The fixpoints are roots of $\lambda z^{d+1} + z - 1$. It will be convenient to reparameterize $z = 1/y$ and consider the roots of $\lambda + z^d - z^{d+1}$. (Note that for $y_1 = 1/z_1, y_2 = 1/z_2$, we have $|z_1| = |z_2|$ if and only if $|y_1| = |y_2|$.)

For the sake of contradiction, suppose that $\lambda = y^{d+1} - y^d$ has two roots y_1, y_2 of the same norm, that is $|y_1| = |y_2|$. We have

$$|\lambda| = |y_1|^d(|y_1 - 1|) = |y_2|^d(|y_2 - 1|),$$

and since $|y_1| = |y_2|$ we conclude $|y_1 - 1| = |y_2 - 1|$. Note that this means that y_1, y_2 lie on the intersection of two circles—one centered at 0 and one centered at 1. This means that y_1 and y_2 are conjugate (since the centers lie on the real line and if two circles intersect in 2 points then the points are symmetric about the segment connecting the centers) and thus

$$\bar{\lambda} = \bar{y}_1^{d+1} - \bar{y}_1^d = y_2^{d+1} - y_2^d = \lambda,$$

that is, $\lambda \in \mathbb{R}$. \square

5. Implementing activities with exponential precision. In this section, we prove our main implementation results, Propositions 2.2 and 2.11. We start in section 5.1 by proving Lemma 2.8 and its analogue for the real case that will help us obtain the exponential precision; we also give a path construction that will give us the desired implementations. Then, in section 5.2, we give the proof of Proposition 2.11 for the real case and, in section 5.3, the proof of Proposition 2.2 for the complex case.

5.1. Contracting maps that cover yield exponential precision. We first restate here Lemma 2.8 for convenience.

LEMMA 2.8. *Let $z_0 \in \mathbb{C}_\mathbb{Q}$, $r > 0$, be a rational, and U be the ball $B(z_0, r)$. Further, suppose that $\lambda'_1, \dots, \lambda'_t \in \mathbb{C}_\mathbb{Q}$ are such that the complex maps $\Phi_i : z \mapsto \frac{1}{1+\lambda'_i z}$ with $i \in [t]$ satisfy the following:*

1. *for each $i \in [t]$, Φ_i is contracting on the ball U ;*
2. *$U \subseteq \bigcup_{i=1}^t \Phi_i(U)$.*

There is an algorithm which, on input of (i) a starting point $x_0 \in U \cap \mathbb{C}_\mathbb{Q}$, (ii) a target $x \in U \cap \mathbb{C}_\mathbb{Q}$, and (iii) a rational $\epsilon > 0$, outputs in $\text{poly}(\text{size}(x_0, x, \epsilon))$ time a number $\hat{x} \in U \cap \mathbb{C}_\mathbb{Q}$ and a sequence $i_1, i_2, \dots, i_k \in [t]$ such that

$$\hat{x} = \Phi_{i_k}(\Phi_{i_{k-1}}(\dots \Phi_{i_1}(x_0) \dots)) \text{ and } |\hat{x} - x| \leq \epsilon.$$

The following is the exact analogue of Lemma 2.8 for the real case. For $z \in \mathbb{R}$ and $r > 0$, we use $I(z, r)$ to denote the interval of length $2r$ centered at z .

LEMMA 5.1. Let $z_0 \in \mathbb{Q}$ and $r > 0$, be a rational. Suppose further that $\lambda'_1, \dots, \lambda'_t \in \mathbb{Q}$ are such that the real maps $\Phi_i : z \mapsto \frac{1}{1+\lambda'_i z}$ with $i \in [t]$ satisfy the following:

1. for each $i \in [t]$, Φ_i is contracting on the interval $I(z_0, r)$;
2. $I(z_0, r) \subseteq \bigcup_{i=1}^t \Phi_i(I(z_0, r))$.

There is an algorithm which, on input of (i) a starting point $x_0 \in I(z_0, r) \cap \mathbb{Q}$, (ii) a target $x \in I(z_0, r) \cap \mathbb{Q}$, and (iii) a rational $\epsilon > 0$, outputs in $\text{poly}(\text{size}(x_0, x, \epsilon))$ time a number $\hat{x} \in I(z_0, r) \cap \mathbb{Q}$ and a sequence $i_1, i_2, \dots, i_k \in [t]$ such that

$$\hat{x} = \Phi_{i_k}(\Phi_{i_{k-1}}(\dots \Phi_{i_1}(x_0) \dots)) \text{ and } |\hat{x} - x| \leq \epsilon.$$

Proof of Lemmas 2.8 and 5.1. We focus on proving Lemma 2.8, since the proof of Lemma 5.1 is almost identical; one only needs to replace the ball $B(z_0, r)$ with the interval $I(z_0, r)$ in the following argument.

Since the maps Φ_i are contracting on the ball $B(z_0, r)$ for all $i \in [t]$, there exists a real number $M < 1$ such that for all $i \in [t]$ and all $x, y \in B(z_0, r)$, it holds that

$$(5.1) \quad |\Phi_i(x) - \Phi_i(y)| \leq M|x - y|.$$

W.l.o.g., we assume that $M \in \mathbb{Q}$.

Let $x_0, x \in B(z_0, r) \cap \mathbb{C}_{\mathbb{Q}}$ and $\epsilon \in (0, 1)$. We are now going to describe a procedure that produces a point that is at distance at most ϵ from x in time polynomial in $\text{size}(x_0, x, \epsilon)$.

```

GetPoint( $x, \epsilon$ )
if  $\epsilon \geq |x - x_0|$ , then return  $x_0$ 
else
  let  $i \in [t]$  be such that  $x \in \Phi_i(B(z_0, r))$ 
   $y \leftarrow \text{GetPoint}(\Phi_i^{-1}(x), \epsilon/M)$ 
  return  $\hat{x} = \Phi_i(y)$ 

```

Note that, in each recursive call of the procedure $\text{GetPoint}(\cdot, \cdot)$, the second parameter increases by a factor $1/M$ and hence the number N of recursive calls is bounded by $1 + \log_{1/M}(2r/\epsilon) = O(\text{size}(\epsilon))$. Moreover, we can find $i \in [t]$ such that $x \in \Phi_i(B(z_0, r))$ in time polynomial in $\text{size}(x, \epsilon)$ (since we can compute $\Phi_i^{-1}(x)$ for each $i \in [t]$ and check whether $|\Phi_i^{-1}(x) - z_0| \leq r$). Finally, note that, in each recursive call of the procedure $\text{GetPoint}(\cdot, \cdot)$, the first parameter is always from $B(z_0, r)$ since $x \in \Phi_i(B(z_0, r))$.

The correctness of the algorithm is proved by induction on the number N of recursive calls. In the base case $N = 0$, we have $\epsilon \geq |x - x_0|$ and the procedure returns x_0 which is at distance at most ϵ from x . For the inductive step we have $|y - \Phi_i^{-1}(x)| \leq \epsilon/M$ and hence by (5.1)

$$|\Phi_i(y) - x| \leq M|y - \Phi_i^{-1}(x)| \leq \epsilon.$$

It remains to observe that we can modify the procedure $\text{GetPoint}(x, \epsilon)$ so that it also returns the desired sequence i_1, i_2, \dots, i_k . \square

We will pair our applications of Lemmas 2.8 and 5.1 with the following path construction.

LEMMA 5.2. Fix $\lambda \in \mathbb{C}_{\mathbb{Q}}$. Let $x_0, \lambda'_1, \dots, \lambda'_t \in \mathbb{C}_{\mathbb{Q}} \setminus \{0\}$ and, for $i \in [t]$, consider the maps $\Phi_i(z) = \frac{1}{1+\lambda'_i z}$ for $z \neq -\frac{1}{\lambda'_i}$. Let $i_1, i_2, \dots, i_k \in [t]$ be a sequence such that $\Phi_{i_k}(\Phi_{i_{k-1}}(\dots \Phi_{i_1}(x_0) \dots)) = \hat{x}$ for some $\hat{x} \in \mathbb{C}$.

Let P be a path of length $k+1$ whose vertices are labeled as $v_0, v_1, \dots, v_k, v_{k+1}$. Let λ be the activity vector on P given by

$$\lambda_{v_0} = (1 - x_0)/x_0, \quad \lambda_{v_j} = \lambda'_{i_j} \text{ for } j \in [k], \quad \lambda_{v_{k+1}} = \lambda.$$

Then, it holds that $Z_{P,v_{k+1}}^{\text{out}}(\lambda) \neq 0$ and $\frac{Z_{P,v_{k+1}}^{\text{in}}(\lambda)}{Z_{P,v_{k+1}}^{\text{out}}(\lambda)} = \lambda \hat{x}$. Moreover, there is an algorithm that, on input $x_0, \lambda'_1, \dots, \lambda'_t, k$, computes the quantities $Z_{P,v_{k+1}}^{\text{in}}(\lambda)$ and $Z_{P,v_{k+1}}^{\text{out}}(\lambda)$ in time polynomial in k and $\text{size}(x_0, \lambda'_1, \dots, \lambda'_t)$.

Proof. For $j = 0, 1, \dots, k$, denote by P_j the path of length j induced by the vertices v_0, v_1, \dots, v_j ; the activity of a vertex v_i in P_j is equal to the activity of the vertex v_i in P . To simplify notation, we will drop the activity vector λ from notation, i.e., we will write $Z_{P_j,v_j}^{\text{in}}, Z_{P_j,v_j}^{\text{out}}, Z_{P_j}$ instead of $Z_{P_j,v_j}^{\text{in}}(\lambda), Z_{P_j,v_j}^{\text{out}}(\lambda), Z_{P_j}(\lambda)$.

We will show by induction on j that

$$(5.2) \quad \frac{Z_{P_j,v_j}^{\text{out}}}{Z_{P_j}} = x_j, \text{ where } x_j = \Phi_{i_j}(\Phi_{i_{j-1}}(\dots \Phi_{i_1}(x_0) \dots)).$$

For $j = 0$, we have

$$(5.3) \quad Z_{P_0,v_0}^{\text{out}} = 1, \quad Z_{P_0} = 1 + \lambda_0 = 1/x_0,$$

and therefore (5.2) holds. Assuming that (5.2) holds for some j in $0, \dots, k-1$, we show that it holds for $j+1$ as well. Note that $x_{j+1} = \Phi_{i_{j+1}}(x_j)$ and therefore $x_j \neq -1/\lambda'_{v_{j+1}}$, i.e., $Z_{P_{j+1},v_{j+1}}^{\text{out}}/Z_{P_{j+1}} \neq -1/\lambda'_{v_{j+1}}$. We have

$$(5.4) \quad \begin{aligned} Z_{P_{j+1},v_{j+1}}^{\text{out}} &= Z_{P_j}, \\ Z_{P_{j+1}} &= Z_{P_{j+1},v_{j+1}}^{\text{in}} + Z_{P_{j+1},v_{j+1}}^{\text{out}} = \lambda_{v_{j+1}} Z_{P_j,v_j}^{\text{out}} + Z_{P_j} \neq 0 \end{aligned}$$

and therefore

$$\frac{Z_{P_{j+1},v_{j+1}}^{\text{out}}}{Z_{P_{j+1}}} = \frac{1}{1 + \lambda_{v_{j+1}} \frac{Z_{P_j,v_j}^{\text{out}}}{Z_{P_j}}} = \Phi_{i_{j+1}}(x_j) = x_{j+1}.$$

This finishes the proof of (5.2) for all $j = 0, 1, \dots, k$. To conclude the proof, note that

$$(5.5) \quad \begin{aligned} Z_{P,v_{k+1}}^{\text{in}} &= \lambda_{v_{k+1}} Z_{P_k,v_k}^{\text{out}} = \lambda Z_{P_k,v_k}^{\text{out}}, \\ Z_{P,v_{k+1}}^{\text{out}} &= Z_{P_k} \neq 0, \end{aligned}$$

so that $\frac{Z_{P,v_{k+1}}^{\text{in}}}{Z_{P,v_{k+1}}^{\text{out}}} = \lambda \hat{x}$. Finally, note that using (5.3), (5.4), (5.5), we can also compute $Z_{P,v_{k+1}}^{\text{in}}(\lambda)$ and $Z_{P,v_{k+1}}^{\text{out}}(\lambda)$ in time polynomial in k and $\text{size}(x_0, \lambda'_1, \dots, \lambda'_t)$. This completes the proof of Lemma 5.2. \square

5.2. Proof of Proposition 2.11 (real case). To prove Proposition 2.11, we will bootstrap Lemma 2.12 to obtain implementations that control logarithmically the size of G in terms of the desired accuracy ϵ . The following technical lemma will allow us to use Lemma 5.1.

LEMMA 5.3. *Let $\lambda_1^* = -1/4$, $\lambda_2^* = -6/25$, and I be the interval $[7/4, 11/6]$. Let $\eta = 10^{-10}$, then the following holds for all $\lambda'_1 \in [\lambda_1^* - \eta, \lambda_1^* + \eta]$ and $\lambda'_2 \in [\lambda_2^* - \eta, \lambda_2^* + \eta]$.*

The maps $\Phi_1(x) = \frac{1}{1+\lambda'_1 x}$ and $\Phi_2(x) = \frac{1}{1+\lambda'_2 x}$ are contracting on the interval I and, moreover, $I \subseteq \Phi_1(I) \cup \Phi_2(I)$.

Proof. For all $\lambda'_1 \in [\lambda_1^* - \eta, \lambda_1^* + \eta]$ and $\lambda'_2 \in [\lambda_2^* - \eta, \lambda_2^* + \eta]$, we have that Φ_1, Φ_2 are increasing, while Φ'_1, Φ'_2 are decreasing. Therefore

$$\begin{aligned}\Phi_1(I) &= [\Phi_1(7/4), \Phi_1(11/6)], & \Phi_2(I) &= [\Phi_2(7/4), \Phi_2(11/6)], \\ \Phi'_1(I) &= [\Phi'_1(11/6), \Phi'_1(7/4)], & \Phi'_2(I) &= [\Phi'_2(11/6), \Phi'_2(7/4)].\end{aligned}$$

By calculating the relevant function values, we obtain that

$$[1.78, 1.84] \subseteq \Phi_1(I), \quad [1.73, 1.78] \subseteq \Phi_2(I), \quad \max_{x \in I} \{|\Phi'_1(x)|, |\Phi'_2(x)|\} \leq 9/10,$$

and the lemma follows. \square

We are now ready to give the proof of Proposition 2.11, which we restate here for convenience.

PROPOSITION 2.11. *Let $\Delta \geq 3$ and $\lambda \in \mathbb{Q}$ be such that $\lambda < -\lambda^*$. There is an algorithm which, on input $\lambda', \epsilon \in \mathbb{Q}$ with $\epsilon > 0$, outputs in $\text{poly}(\text{size}(\lambda', \epsilon))$ time a bipartite graph G of maximum degree at most Δ with terminal v that implements λ' with accuracy ϵ . Moreover, the algorithm outputs the values $Z_{G,v}^{\text{in}}(\lambda), Z_{G,v}^{\text{out}}(\lambda)$.*

Proof of Proposition 2.11. Let $I = [7/4, 11/6]$ be the interval of Lemma 5.3. The main idea of the proof is to use the maps in Lemma 5.3 in combination with Lemma 5.1 to get a subinterval of I where we can get exponentially accurate implementations. Then, we will propagate this exponential accuracy to the whole real line by using implementations from Lemma 2.12.

We next specify the activities that we will need to implement with constant precision via Lemma 5.3 (later, these activities will be combined with Lemma 5.1 to get the exponential precision). Let $\lambda_1^*, \lambda_2^*, \eta, I$ be as in Lemma 5.3. Moreover, let x_0^* be the midpoint of the interval I and let $\lambda_0^* = (1 - x_0^*)/x_0^*$, $\lambda_3^* = -1/x_0^*$. Finally, let $\lambda_4^* = \max\{10^{12}, 10^{12}|\lambda|\}$. (Note that all of these are rationals.)

By Lemma 2.12, for $i = 0, 1, \dots, 4$, there exists a bipartite graph G_i of maximum degree Δ which implements λ_i^* with accuracy η . Let v_i be the terminal of G_i and set $\lambda'_i = \frac{Z_{G_i, v_i}^{\text{in}}(\lambda)}{Z_{G_i, v_i}^{\text{out}}(\lambda)}$; we have that $\lambda'_i \in [\lambda_i^* - \eta, \lambda_i^* + \eta]$. Moreover, let $x_0 = 1/(1 + \lambda'_0)$ and note that $|x_0 - x_0^*| \leq 10^{-5}$, so that $x_0 \in I$. Also, let $x_3 = -1/\lambda'_3$ and note that $|x_3 - x_0^*| \leq 10^{-5}$, so that the interval $I^* = [x_3 - 10^{-3}, x_3 + 10^{-3}]$ is a subinterval of I . Finally, we have that

$$(5.6) \quad |\lambda'_4| \geq |\lambda_4^*| - \eta \geq |\lambda_4^*| - |\lambda_4^*|/2 \geq \max\{10^{11}, 10^{11}|\lambda|\}.$$

Suppose that we are given inputs $\lambda', \epsilon \in \mathbb{Q}$ with $\epsilon > 0$ and we want to output in $\text{poly}(\text{size}(\lambda', \epsilon))$ time a bipartite graph of maximum degree Δ that implements λ' with accuracy ϵ . Clearly, we may assume that $\epsilon \in (0, 1)$. The algorithm has three cases depending on the value of $|\lambda'|$.

Case I (large $|\lambda'|$): $|\lambda'| \geq \max\{10^4|\lambda|, 1\}$. Let x^* be a rational such that

$$(5.7) \quad \frac{\lambda}{1 + \lambda'_3 x^*} = \lambda', \text{ so that } x^* = \frac{1}{\lambda'_3} \left(\frac{\lambda}{\lambda'} - 1 \right).$$

Recall that $x_3 = -1/\lambda'_3$, so using the assumption $|\lambda'| \geq 10^4|\lambda|$ and that $|\lambda'_3| \geq |\lambda_3^*| - \eta \geq 1/10$, we have that

$$|x^* - x_3| = \frac{|\lambda|}{|\lambda' \lambda'_3|} \leq 10^{-3}.$$

It follows that x^* belongs to the interval I .

Note that Lemma 5.3 guarantees that the maps $\Phi_1(x) = \frac{1}{1+\lambda'_1 x}$ and $\Phi_2(x) = \frac{1}{1+\lambda'_2 x}$ satisfy the hypotheses of Lemma 5.1 with $z_0 = x_0^*$ and r equal to half the length of the interval I . Therefore, using the algorithm of Lemma 5.1, on input x_0, x^* and $\epsilon' = \epsilon \cdot \min \left\{ \frac{|\lambda|}{2|\lambda'\lambda'_3|}, \frac{|\lambda|}{2|(\lambda')^2\lambda'_3|}, 1 \right\}$, we obtain in time $\text{poly}(\text{size}(x_0, x^*, \epsilon')) = \text{poly}(\text{size}(\lambda', \epsilon))$ a number \hat{x} and a sequence $i_1, \dots, i_k \in \{1, 2\}$ such that

$$(5.8) \quad \hat{x} = \Phi_{i_k}(\Phi_{i_{k-1}}(\dots \Phi_{i_1}(x_0) \dots)) \text{ and } |\hat{x} - x^*| \leq \epsilon'.$$

Using (5.7) and (5.8), we have by the triangle inequality that

$$(5.9) \quad |1 + \lambda'_3 \hat{x}| \geq |1 + \lambda'_3 x^*| - |\lambda'_3(\hat{x} - x^*)| \geq \frac{|\lambda|}{|\lambda'|} - |\lambda'_3 \epsilon'| \geq \frac{|\lambda|}{2|\lambda'|}$$

and, therefore,

$$(5.10) \quad \begin{aligned} \left| \frac{\lambda}{1 + \lambda'_3 \hat{x}} - \lambda' \right| &= \left| \frac{\lambda}{1 + \lambda'_3 \hat{x}} - \frac{\lambda}{1 + \lambda'_3 x^*} \right| = \frac{|\lambda| |\lambda'_3| |\hat{x} - x^*|}{|1 + \lambda'_3 x^*| \cdot |1 + \lambda'_3 \hat{x}|} \\ &= \frac{|\lambda' \lambda'_3| |\hat{x} - x^*|}{|1 + \lambda'_3 \hat{x}|} \leq \frac{2|(\lambda')^2 \lambda'_3|}{|\lambda|} |\hat{x} - x^*| \leq \epsilon, \end{aligned}$$

where in the last equality we used (5.7), in the second to last inequality we used (5.9), and in the last inequality we used (5.8) and the choice of ϵ' .

Now, let P be a path of length $k+2$ with vertices labeled $v_0, v_1, \dots, v_{k+1}, v_{k+2}$. Let λ be the activity vector on P given by

$$\lambda_{v_0} = \lambda'_0, \quad \lambda_{v_j} = \lambda'_{i_j} \text{ for } j \in [k], \quad \lambda_{v_{k+1}} = \lambda'_3, \quad \lambda_{v_{k+2}} = \lambda.$$

Then, by Lemma 5.2, it holds that $Z_{P, v_{k+2}}^{\text{out}}(\lambda) \neq 0$ and $\frac{Z_{P, v_{k+2}}^{\text{in}}(\lambda)}{Z_{P, v_{k+2}}^{\text{out}}(\lambda)} = \frac{\lambda}{1 + \lambda'_3 \hat{x}}$; moreover, we can compute the values $Z_{P, v_{k+2}}^{\text{in}}(\lambda), Z_{P, v_{k+2}}^{\text{out}}(\lambda)$ in time $\text{poly}(k, \text{size}(\lambda'_0, \lambda'_1, \lambda'_2)) = \text{poly}(\text{size}(\lambda', \epsilon))$. Since the bipartite graphs G_0, G_1, G_2, G_3 implement the activities $\lambda'_0, \lambda'_1, \lambda'_2, \lambda'_3$, respectively, we obtain by applying Lemma 3.2 (to the path P with activity vector λ) a bipartite graph G' with maximum degree Δ and terminal v_{k+2} such that

$$(5.11) \quad Z_{G', v_{k+2}}^{\text{in}}(\lambda) = C \cdot Z_{P, v_{k+2}}^{\text{in}}(\lambda), \quad Z_{G', v_{k+2}}^{\text{out}}(\lambda) = C \cdot Z_{P, v_{k+2}}^{\text{out}}(\lambda),$$

where $C = \prod_{i=1}^3 (Z_{G_i, v_i}^{\text{out}}(\lambda))^{\{j \in \{0, \dots, k+2\} \mid \lambda_{v_j} = \lambda'_i\}}$. We conclude that

$$\frac{Z_{G', v_{k+2}}^{\text{in}}(\lambda)}{Z_{G', v_{k+2}}^{\text{out}}(\lambda)} = \frac{Z_{P, v_{k+2}}^{\text{in}}(\lambda)}{Z_{P, v_{k+2}}^{\text{out}}(\lambda)} = \frac{\lambda}{1 + \lambda'_3 \hat{x}}.$$

Combining this with (5.10), we obtain that G' with terminal v_{k+2} is a bipartite graph of maximum degree Δ which implements λ' with accuracy ϵ . Moreover, using (5.11), we can also compute the values $Z_{G', v_{k+2}}^{\text{in}}(\lambda), Z_{G', v_{k+2}}^{\text{out}}(\lambda)$.

Case II (small $|\lambda'|$): $|\lambda'| \leq \min\{10^{-5}, 10^{-5}|\lambda|\}$. We first assume that $\lambda' \neq 0$.

Let $\hat{\lambda}$ be such that

$$(5.12) \quad \frac{\lambda}{1 + \hat{\lambda}} = \lambda' \text{ so that } \hat{\lambda} = \frac{\lambda}{\lambda'} - 1.$$

Using the assumption $|\lambda'| \leq \min\{10^{-5}, 10^{-5}|\lambda|\}$ and the triangle inequality, we have that

$$|\hat{\lambda}| \geq \frac{|\lambda|}{|\lambda'|} - 1 \geq \frac{|\lambda|}{2|\lambda'|} \geq \max\{10^4|\lambda|, 1\}.$$

Therefore, by Case I, we can construct a bipartite graph G with terminal v that implements $\hat{\lambda}$ with accuracy $\epsilon' = \epsilon \cdot \min\{\frac{|\lambda|}{2|\lambda'|}, \frac{|\lambda|}{4|\lambda'|^2}, 1\}$ in time $\text{poly}(\text{size}(\hat{\lambda}, \epsilon')) = \text{poly}(\text{size}(\lambda', \epsilon))$. Moreover, we can compute the values $Z_{G,v}^{\text{in}}(\lambda)$ and $Z_{G,v}^{\text{out}}(\lambda)$ at the same time. Let $\lambda'' = \frac{Z_{G,v}^{\text{in}}(\lambda)}{Z_{G,v}^{\text{out}}(\lambda)}$ so that $|\lambda'' - \hat{\lambda}| \leq \epsilon'$. Using this and (5.12), we have

$$(5.13) \quad |1 + \lambda''| \geq |1 + \hat{\lambda}| - |(\lambda'' - \hat{\lambda})| \geq \frac{|\lambda|}{|\lambda'|} - \epsilon' \geq \frac{|\lambda|}{2|\lambda'|}$$

and, therefore,

$$(5.14) \quad \begin{aligned} \left| \frac{\lambda}{1 + \lambda''} - \lambda' \right| &= \left| \frac{\lambda}{1 + \lambda''} - \frac{\lambda}{1 + \hat{\lambda}} \right| = \frac{|\lambda| |\lambda'' - \hat{\lambda}|}{|1 + \hat{\lambda}| \cdot |1 + \lambda''|} \\ &= \frac{|\lambda'| |\lambda'' - \hat{\lambda}|}{|1 + \lambda''|} \leq \frac{2|\lambda'|^2}{|\lambda|} |\lambda'' - \hat{\lambda}| \leq \epsilon, \end{aligned}$$

where in the last equality we used (5.12), in the second to last inequality we used (5.13), and in the last inequality we used $|\lambda'' - \hat{\lambda}| \leq \epsilon'$ and the choice of ϵ' .

Now, let G' be the bipartite graph obtained from G by adding a new vertex u whose single neighbor is the terminal v of G . Then, we have that

$$(5.15) \quad Z_{G',u}^{\text{in}}(\lambda) = \lambda Z_{G,v}^{\text{out}}(\lambda), \quad Z_{G',u}^{\text{out}}(\lambda) = Z_G(\lambda) = Z_{G,v}^{\text{in}}(\lambda) + Z_{G,v}^{\text{out}}(\lambda).$$

We conclude that

$$\frac{Z_{G',u}^{\text{in}}(\lambda)}{Z_{G',u}^{\text{out}}(\lambda)} = \frac{\lambda Z_{G,v}^{\text{out}}(\lambda)}{Z_{G,v}^{\text{in}}(\lambda) + Z_{G,v}^{\text{out}}(\lambda)} = \frac{\lambda}{1 + \lambda''}.$$

Combining this with (5.14), we obtain that G' with terminal u is a bipartite graph of maximum degree Δ which implements λ' with accuracy ϵ . Moreover, using (5.15), we can also compute the values $Z_{G',u}^{\text{in}}(\lambda)$, $Z_{G',u}^{\text{out}}(\lambda)$.

To finish this case, it remains to argue for $\lambda' = 0$. In this case, for $\epsilon'' = \min\{\epsilon, 10^{-5}, 10^{-5}|\lambda|\}$, we can use the preceding method to implement the activity $\epsilon''/2 \neq 0$ with accuracy $\epsilon''/2$ in time $\text{poly}(\text{size}(\epsilon'')) = \text{poly}(\text{size}(\epsilon))$. The implemented activity λ'' satisfies by the triangle inequality $|\lambda''| \leq \epsilon$; hence, we have implemented the desired activity $\lambda' = 0$ with accuracy ϵ .

Case III (moderate $h|\lambda'|$): $\min\{10^{-5}, 10^{-5}|\lambda|\} < |\lambda'| < \max\{10^4|\lambda|, 1\}$. Let x^* be a rational such that

$$(5.16) \quad \frac{\lambda}{1 + \lambda'_4 x^*} = \lambda' \text{ so that } x^* = \frac{1}{\lambda'_4} \left(\frac{\lambda}{\lambda'} - 1 \right).$$

Using the assumption $10^{-5}|\lambda| < |\lambda'|$ and $|\lambda'_4| \geq \max\{10^{11}, 10^{11}|\lambda|\}$ (cf. (5.6)), we have that $|x^*| \leq \min\{10^{-5}/|\lambda|, 10^{-5}\}$. Let $\epsilon' = \epsilon \cdot \min\{\frac{1}{10^{10}|\lambda'_4|}, \frac{|\lambda|}{10^{10}|\lambda'_4|}\}$. Then, by the algorithm for Case II, we can implement the activity λx^* with precision ϵ' in time $\text{poly}(\text{size}(\lambda x^*, \epsilon')) = \text{poly}(\text{size}(\lambda', \epsilon))$. That is, we can construct a bipartite graph G

of maximum degree at most Δ with terminal v such that, for $\lambda'' := Z_{G,v}^{\text{in}}(\lambda)/Z_{G,v}^{\text{out}}(\lambda)$, it holds that

$$(5.17) \quad |\lambda'' - \lambda x^*| \leq \epsilon'.$$

Now, using (5.16) and (5.17), we have by the triangle inequality that

$$(5.18) \quad |1 + \lambda'_4(\lambda''/\lambda)| \geq |1 + \lambda'_4 x^*| - |\lambda'_4(x^* - \lambda''/\lambda)| \geq \frac{1}{10^4} - \frac{1}{10^{10}} \geq \frac{1}{10^5}$$

and, therefore,

$$(5.19) \quad \left| \frac{\lambda}{1 + \lambda'_4(\lambda''/\lambda)} - \lambda' \right| = \left| \frac{\lambda}{1 + \lambda'_4(\lambda''/\lambda)} - \frac{\lambda}{1 + \lambda'_4 x^*} \right| = \frac{|\lambda| |\lambda'_4| |x^* - (\lambda''/\lambda)|}{|1 + \lambda'_4 x^*| \cdot |1 + \lambda'_4(\lambda''/\lambda)|} \\ = \frac{|\lambda'| |\lambda'_4| |x^* - (\lambda''/\lambda)|}{|1 + \lambda'_4(\lambda''/\lambda)|} \leq 10^9 |\lambda'_4| |\lambda x^* - \lambda''| \leq \epsilon,$$

where in the last equality we used (5.16), in the second to last inequality we used (5.18) and $|\lambda'|/|\lambda| \leq 10^4$, and in the last inequality we used (5.17) and the choice of ϵ' .

Recall that G_4 is a bipartite graph of maximum degree Δ , with terminal v_4 , which implements the activity λ'_4 . Let G' be the bipartite graph obtained by taking a copy of G_4 and G and identifying the terminals v_4, v into a single vertex which we label u' (note that G' has maximum degree Δ as well since v_4 and v have degree one in G_4 and G , respectively). Then,

$$(5.20) \quad Z_{G',u'}^{\text{in}}(\lambda) = \frac{1}{\lambda} Z_{G_4,v_4}^{\text{in}}(\lambda) Z_{G,u}^{\text{in}}(\lambda), \quad Z_{G',u'}^{\text{out}}(\lambda) = Z_{G_4,v_4}^{\text{out}}(\lambda) Z_{G,v}^{\text{out}}(\lambda).$$

Consider the graph G'' obtained from G' by adding a new vertex u'' whose single neighbor is the vertex u' . Then, we have that

$$(5.21) \quad Z_{G'',u''}^{\text{in}}(\lambda) = \lambda Z_{G',u'}^{\text{out}}(\lambda), \quad Z_{G'',u''}^{\text{out}}(\lambda) = Z_{G'}(\lambda) = Z_{G',u'}^{\text{in}}(\lambda) + Z_{G',u'}^{\text{out}}(\lambda).$$

Using this in conjunction with (5.20), we conclude that

$$\frac{Z_{G'',u''}^{\text{in}}(\lambda)}{Z_{G'',u''}^{\text{out}}(\lambda)} = \frac{\lambda Z_{G',u'}^{\text{out}}(\lambda)}{Z_{G',u'}^{\text{in}}(\lambda) + Z_{G',u'}^{\text{out}}(\lambda)} = \frac{\lambda}{1 + \frac{Z_{G',u}^{\text{in}}(\lambda)}{Z_{G',u}^{\text{out}}(\lambda)}} = \frac{\lambda}{1 + \lambda'_4(\lambda''/\lambda)}.$$

Combining this with (5.19), we obtain that G'' is a bipartite graph of maximum degree at most Δ with terminal u'' which implements λ' with accuracy ϵ . Moreover, using (5.20) and (5.21), we can also compute the values $Z_{G'',u''}^{\text{in}}(\lambda), Z_{G'',u''}^{\text{out}}(\lambda)$.

This completes the three different cases of the algorithm, thus completing the proof of Proposition 2.11. \square

5.3. Proof of Proposition 2.2 (complex case). In this section, we prove Proposition 2.2 assuming Proposition 2.6 (the proof of the latter is given in section 7). Note that Proposition 2.6 applies for all $\lambda \in \mathbb{C}_{\mathbb{Q}} \setminus \mathbb{R}$. By restricting our attention to $\lambda \notin \Lambda_{\Delta} \cup \mathbb{R}$ and using the theory of section 2.1, we obtain the following.

LEMMA 5.4. *Let $\Delta \geq 3$ and $\lambda \in \mathbb{C}_{\mathbb{Q}} \setminus \mathbb{R}$ be such that $\lambda \notin \Lambda_{\Delta}$, and set $d := \Delta - 1$. Let ω be the fixpoint of $f(z) = \frac{1}{1+\lambda z^d}$ with the smallest norm, and p_1, \dots, p_d be the poles of f . Then, for any real number $\eta > 0$, there exist*

- (i) $N \in \mathbb{Z}_{\geq 1}$, (ii) a pole $p^* \in \{p_1, \dots, p_d\}$, (iii) rationals $L > 0$ and $r, r', r^* \in (0, \eta)$ such that $r' < r$ and all of the following hold:

1. $B(p^*, r^*) \subseteq f^N(B(\omega, r'))$;
2. $p_1, \dots, p_d \notin \bigcup_{n=0}^{N-1} f^n(B(\omega, r))$; and
3. for all $x_1, x_2 \in B(\omega, r)$, it holds that $|f^N(x_1) - f^N(x_2)| \leq L|x_1 - x_2|$.

Proof. Consider an arbitrary $\eta > 0$. Note that $\omega \neq p_1, \dots, p_d$, so there is no loss of generality in assuming that $\eta \leq |\omega - p_1|, \dots, |\omega - p_d|$. Let U' be the open ball $B(\omega, \eta/10)$ and note that our assumption on η ensures that $p_1, \dots, p_d \notin U'$.

Since $\lambda \notin \Lambda_\Delta \cup \mathbb{R}$, we have by Lemma 2.5 that ω is a repelling fixpoint of f and therefore ω belongs to the Julia set of f (by Lemma 2.3). Moreover, using Lemma 3.7, we have that the exceptional set of f is empty.⁶ Therefore, by Theorem 2.4, it holds that $\bigcup_{n=0}^{\infty} f^n(U') = \mathbb{C}$. Let N' be the smallest integer such that one of the poles p_1, \dots, p_d belongs to $f^{N'}(U')$, i.e., N' satisfies

$$(5.22) \quad \{p_1, \dots, p_d\} \cap \left(\bigcup_{n=0}^{N'-1} f^n(U') \right) = \emptyset \text{ and } \{p_1, \dots, p_d\} \cap f^{N'}(U') \neq \emptyset.$$

Note that $N' \geq 1$ since $p_1, \dots, p_d \notin U'$.

To prove the lemma, it will be important to ensure that the pole of f which belongs to $f^{N'}(U')$ does not sit on the boundary of any of the sets $f^0(U'), f^1(U'), \dots, f^{N'}(U')$. To achieve this, we will enlarge a little bit the ball U' as follows. Let \mathcal{P} be the union of the poles of the functions $f^1, \dots, f^{N'}$. Note that \mathcal{P} is a finite set, therefore, we can specify a radius $r \in (\eta/10, \eta)$ so that the boundary ∂U of the open ball $U = B(\omega, r)$ is disjoint from \mathcal{P} (i.e., $\partial U \cap \mathcal{P} = \emptyset$). Since $U' \subseteq U$ and $p_1, \dots, p_d \notin U$, we conclude from (5.22) that there exists a positive integer $N \leq N'$ such that

$$(5.23) \quad \{p_1, \dots, p_d\} \cap \left(\bigcup_{n=0}^{N-1} f^n(U) \right) = \emptyset \text{ and } \{p_1, \dots, p_d\} \cap f^N(U) \neq \emptyset,$$

i.e., $N \leq N'$ is the first integer such that a pole of f belongs to $f^N(U)$. Let $p^* \in \{p_1, \dots, p_d\}$ be an arbitrary pole of f such that $p^* \in f^N(U)$. We claim that

$$(5.24) \quad \text{for all } n \in \{0, \dots, N\}, p^* \text{ does not lie on the boundary of } f^n(U).$$

Indeed, observe that U is open and f^n is holomorphic on U for all $n = 0, \dots, N$ since $f^{n-1}(U)$ does not contain any pole of f . Therefore, by the open mapping theorem, we have that

$$(5.25) \quad f^n(U) \text{ is an open set for all } n = 0, \dots, N.$$

Since $p^* \in f^N(U)$, this already shows that p^* does not lie on the boundary of $f^N(U)$. For $n = 0, \dots, N-1$, we obtain from (5.25) and the open mapping theorem that p^* lies on the boundary of $f^n(U)$ only if p^* lies on the boundary of $f^n(\partial U)$, i.e., $f^n(\partial U)$ contains a pole of f and so ∂U contains a pole of f^{n+1} . In turn, this would imply that $\partial U \cap \mathcal{P} \neq \emptyset$, which is excluded by the choice of the radius r of U . This proves (5.24).

⁶To see this, let E_f denote the exceptional set of f . By Lemma 3.7, we have that a necessary condition for a point $x \in \mathbb{C}$ to be in E_f is that either $f'(x) = 0$ or $f'(f(x))f'(x) = 0$, which gives $x = 0$ as the only possible point. However, since $f(0) = 1$, we have that $x = 0$ cannot be a fixpoint of either f or f^2 and therefore, by Lemma 3.7, there is no point $x \in \mathbb{C}$ that belongs to E_f . Similarly, we have that $\infty \notin E_f$ since $x = \infty$ is not a fixpoint of either f or f^2 (by $f(\infty) = 0$ and $f(0) = 1$), proving that E_f is empty.

We are now ready to prove items 1 and 2 of the lemma. Namely, from (5.23), we have that $p_1, \dots, p_d \notin \bigcup_{n=0}^{N-1} f^n(B(\omega, r))$, which proves item 2. For item 1, note that from $p^* \in f^N(U)$ we obtain that there exists $x^* \in U = B(\omega, r)$ such that $f^N(x^*) = p^*$. Since U is an open ball, let r' be a rational such that $|x^* - \omega| < r' < r$ and consider the open ball $B(\omega, r')$. Then, we have that $p^* \in f^N(B(\omega, r'))$ (since $x^* \in B(\omega, r')$). We also have by the open mapping theorem that $f^N(B(\omega, r'))$ is open, therefore, there exists rational $r^* \in (0, \eta)$ such that $B(p^*, r^*) \subseteq f^N(B(\omega, r'))$, thus proving item 1.

It remains to prove item 3, which essentially follows from the fact $\bigcup_{n=0}^{N-1} f^n(U)$ does not contain any poles of f and (crude) Lipschitz arguments. In particular, note that, by (5.23) and (5.25), there exists a rational $\delta > 0$ such that

$$(5.26) \quad |x - p_1|, \dots, |x - p_d| \geq \delta \text{ for all } x \in \bigcup_{n=0}^{N-1} f^n(U).$$

Let $L_0 = 1$ and define $L_{n+1} = L_n \frac{d(|\omega| + L_n r)^{d-1}}{|\lambda| \delta^{2d}}$ for $n = 1, \dots, N-1$. For $n = 0, \dots, N$, we will show by induction that

$$(5.27) \quad \text{for all } x_1, x_2 \in U \text{ it holds that } |f^n(x_1) - f^n(x_2)| \leq L_n |x_1 - x_2|,$$

which clearly proves item 3 by taking L to be any rational $> L_N$. The base case $n = 0$ is trivial, so assume that (5.27) holds for some nonnegative integer $n \leq N-1$. Then, since ω is a fixpoint of f , we have that $f^n(\omega) = \omega$ and therefore $f^n(U) \subseteq B(\omega, L_n r)$, i.e.,

$$(5.28) \quad |x| \leq |\omega| + L_n r \text{ for all } x \in f^n(U).$$

Moreover, since $n \leq N-1$, we have by (5.26) that $|x - p_1|, \dots, |x - p_d| \geq \delta$ and hence by factoring $1 + \lambda x^d = \lambda(x - p_1) \cdots (x - p_d)$, we obtain

$$(5.29) \quad |1 + \lambda x^d| = |\lambda(x - p_1) \cdots (x - p_d)| \geq |\lambda| \delta^d \text{ for all } x \in f^n(U).$$

Let $x_1, x_2 \in U$ and set $z_1 = f^n(x_1)$, $z_2 = f^n(x_2)$, so that the inductive hypothesis translates into

$$(5.30) \quad |z_1 - z_2| \leq L_n |x_1 - x_2|.$$

We then obtain that

$$\begin{aligned} |f^{n+1}(x_1) - f^{n+1}(x_2)| &= |f(z_1) - f(z_2)| = \frac{|\lambda(z_1^d - z_2^d)|}{|(1 + \lambda z_1^d)(1 + \lambda z_2^d)|} \\ &= |z_1 - z_2| \cdot \frac{|\lambda| \cdot |z_1^{d-1} + z_1^{d-2} z_2 + \cdots + z_2^{d-1}|}{|(1 + \lambda z_1^d)(1 + \lambda z_2^d)|} \\ &\leq L_{n+1} |x_1 - x_2|, \end{aligned}$$

where in the last inequality we used (5.28), (5.29), and (5.30). This finishes the proof of (5.27) and, therefore, the proof of item 3. This concludes the proof of Lemma 5.4. \square

We are now ready to prove Proposition 2.2, which we restate here for convenience.

PROPOSITION 2.2. *Let $\Delta \geq 3$ and $\lambda \in \mathbb{C}_{\mathbb{Q}}$ be such that $\lambda \notin \Lambda_{\Delta} \cup \mathbb{R}$. There is an algorithm which, on input $\lambda' \in \mathbb{C}_{\mathbb{Q}}$ and rational $\epsilon > 0$, outputs in $\text{poly}(\text{size}(\lambda'), \epsilon)$ time a bipartite graph G of maximum degree at most Δ with terminal v that implements λ' with accuracy ϵ . Moreover, the algorithm outputs the values $Z_{G,v}^{\text{in}}(\lambda), Z_{G,v}^{\text{out}}(\lambda)$.*

Proof. For convenience, set $d := \Delta - 1$. Let ω be the fixpoint of $f(x) = \frac{1}{1+\lambda x^d}$ with the smallest norm, and p_1, \dots, p_d be the poles of f . Let $\rho > 0$ be the constant in Proposition 2.6. Then, by Lemma 5.4 (applied with $\eta = \rho/|\lambda|$), there exist a positive integer N , rationals $L > 0$, and $r, r', r^* \in (0, \eta)$ with $r' < r$, and a pole $p^* \in \{p_1, \dots, p_d\}$ such that

$$(5.31) \quad B(p^*, r^*) \subseteq f^N(B(\omega, r')), \quad p_1, \dots, p_d \notin \bigcup_{n=0}^{N-1} f^n(B(\omega, r))$$

$$(5.32) \quad \text{for all } x_1, x_2 \in B(\omega, r), \text{ it holds that } |f^N(x_1) - f^N(x_2)| \leq L|x_1 - x_2|.$$

We may assume that r^* is sufficiently small so that, for all poles $p \in \{p_1, \dots, p_d\}$ which are different than p^* it holds that

$$(5.33) \quad |x - p| \geq \delta \text{ for all } x \in B(p^*, r^*),$$

where $\delta > 0$ is a sufficiently small constant. Moreover, since p^* is a pole of f , we have that $1 + \lambda(p^*)^d = 0$, so there exists a unique integer $k \in \{0, 1, \dots, d-1\}$ so that $p^* = \frac{1}{|\lambda|^{1/d}} e^{i\theta + 2\pi i(k/d)}$, where $\theta = \frac{1}{d}(\pi - \text{Arg}(\lambda))$. Since k is an integer depending on λ but not on the inputs λ' or ϵ , the value of k which specifies p^* among the poles of f may be used by the algorithm.

Now, suppose that we are given inputs $\lambda' \in \mathbb{C}_{\mathbb{Q}}$ and rational $\epsilon > 0$. We want to output in time $\text{poly}(\text{size}(\lambda', \epsilon))$ a bipartite graph of maximum degree Δ that implements λ' with accuracy ϵ . Clearly, we may assume that $\epsilon \in (0, 1)$. Let $M > 0$ be a rational so that $M > 2/(r^*\delta^d)$. The algorithm has three cases depending on the value of $|\lambda'|$, namely,

$$\text{Case I: } |\lambda'| \geq M, \quad \text{Case II: } |\lambda'| \leq |\lambda|/(M+1), \quad \text{Case III: } |\lambda|/(M+1) < |\lambda'| < M.$$

Note that since $\lambda, \lambda' \in \mathbb{C}_{\mathbb{Q}}$ and $M \in \mathbb{Q}$, the algorithm can distinguish in time $\text{poly}(\text{size}(\lambda, \lambda', M)) = \text{poly}(\text{size}(\lambda'))$ which of the three cases applies.⁷

Case I (large $|\lambda'|$): $|\lambda'| \geq M$. The rough outline of the proof is to first specify and implement an activity λw for some appropriate w whose main property is that $\lambda/(1 + \lambda(f^N(w))^d)$ is ϵ -close to λ' ; then, we will show how to implement the activity $\lambda/(1 + \lambda(f^N(w))^d)$ by using an appropriate tree construction.

We begin by specifying w . We first claim that there exists a unique $x^* \in B(p^*, r^*/2)$ such that

$$(5.34) \quad \frac{\lambda}{1 + \lambda(x^*)^d} = \lambda'.$$

Indeed, for all x such that $|x - p^*| = r^*/2$ it holds that (using (5.33) and the choice of M)

$$(5.35) \quad |1 + \lambda x^d| = |\lambda| \cdot |x - p_1| \cdots |x - p_d| \geq |\lambda| r^* \delta^d / 2 > |\lambda|/|\lambda'|,$$

so by Rouché's theorem⁸ we have that the polynomial $1 + \lambda x^d - \frac{\lambda}{\lambda'}$ has the same number of roots as the polynomial $1 + \lambda x^d$ in the ball $B(p^*, r^*/2)$; the roots of the

⁷E.g., by squaring the inequalities, the radical of the norm goes away and the algorithm has just to compare rational numbers.

⁸Rouché's theorem says that for any functions f and g that are holomorphic inside a region B surrounded by a simple closed contour ∂B , if $|g(x)| < |f(x)|$ on ∂B , then f and $f + g$ have the same number of roots inside B .

latter polynomial are precisely the poles p_1, \dots, p_d and therefore, by (5.33), exactly one of those lies in the ball $B(p^*, r^*/2)$ (namely, p^*). This establishes the existence and uniqueness of $x^* \in B(p^*, r^*/2)$ satisfying (5.34).

By the first part of (5.31), we obtain that there exists $w^* \in B(\omega, r')$ such that $f^N(w^*) = x^*$, i.e.,

$$(5.36) \quad \frac{\lambda}{1 + \lambda(f^N(w^*))^d} = \lambda'.$$

A fact we will use later is that

$$(5.37) \quad |f^N(w^*) - p^*| = |x^* - p^*| > \tau, \text{ where } \tau > 0 \text{ is such that } d\tau(|p^*| + \tau)^{d-1} < 1/|\lambda'|.$$

To see this, note that for all x such that $|x - p^*| \leq \tau$ we have

$$\begin{aligned} |1 + \lambda x^d| &\leq |1 + \lambda(p^*)^d| + |\lambda| \cdot |x^d - (p^*)^d| = |\lambda| \cdot |x^d - (p^*)^d| \\ &\leq |\lambda| \cdot |x - p^*| \cdot \left| \sum_{j=0}^{d-1} x^j (p^*)^{d-1-j} \right| \leq d|\lambda|\tau(|p^*| + \tau)^{d-1} < |\lambda|/|\lambda'| \end{aligned}$$

and, therefore, by (5.34), it must be the case that $|x^* - p^*| > \tau$, thus proving (5.37). Note, we can compute $\tau \in \mathbb{Q}$ satisfying (5.37) in time $\text{poly}(\text{size}(\lambda'))$. Let

$$\hat{\epsilon} := \min \left\{ \frac{r - r'}{3}, \frac{r^*}{4L}, \frac{\tau}{4L}, \frac{1}{4dL|\lambda'|(|p^*| + r^*)^{d-1}}, \frac{\epsilon}{4dL(|\lambda'|^2)(|p^*| + r^*)^{d-1}}, 1 \right\},$$

and let $\epsilon' \in (0, \hat{\epsilon})$ be a rational with $\text{size}(\epsilon') = \text{poly}(\text{size}(\lambda'), \epsilon)$.

Note that $f^N(z)$ is a rational function of degree d^N and, in fact, we can write it as $f^N(z) = \frac{P(z)}{Q(z)}$, where $P(z), Q(z)$ are polynomials with coefficients in $\mathbb{C}_\mathbb{Q}$. Therefore, we can rewrite (5.36) as a polynomial equation in terms of w^* , whose degree is at most d^N (note that this is independent of λ' and ϵ) and whose coefficients have polynomial size in terms of $\text{size}(\lambda')$. Let w_1, \dots, w_t denote the roots of the polynomial. Using Lemma 3.5, we can compute $\hat{w}_1, \dots, \hat{w}_t \in \mathbb{C}_\mathbb{Q}$ such that $|w_i - \hat{w}_i| \leq \epsilon'/2$ for all $i \in [t]$. In particular, there exists $j \in [t]$ such that $|\hat{w}_j - w^*| \leq \epsilon'/2$ and therefore $\hat{w}_j \in B(\omega, r' + \epsilon'/2)$. Moreover, by applying (5.32) for $x_1 = \hat{w}_j$ and $x_2 = w^*$, we have

$$|f^N(\hat{w}_j) - x^*| = |f^N(\hat{w}_j) - f^N(w^*)| \leq L|\hat{w}_j - w^*| \leq L\epsilon'/2.$$

Therefore, by trying⁹ all of $\hat{w}_1, \dots, \hat{w}_t$, we can specify $\hat{w} \in \{\hat{w}_1, \dots, \hat{w}_t\}$ in time $\text{poly}(\text{size}(\lambda'), \epsilon)$ such that

$$(5.38) \quad \hat{w} \in B(\omega, r' + \epsilon'), \quad |f^N(\hat{w}) - x^*| \leq L\epsilon'.$$

Since $\epsilon' \leq (r - r')/3$ we have that $\hat{w} \in B(\omega, r)$. Recall that $r < \rho/|\lambda|$, so by the algorithm of Proposition 2.6, we can construct a bipartite graph G of maximum degree Δ with terminal v that implements $\lambda\hat{w}$ with accuracy $\lambda\epsilon'$ in time $\text{poly}(\text{size}(\lambda\hat{w}), \lambda\epsilon') =$

⁹Note that ω is the root of the polynomial $x + \lambda x^{d+1} - 1$ with the smallest norm and therefore, using Lemma 3.5, we can compute $\hat{\omega} \in \mathbb{C}_\mathbb{Q}$ such that $|\hat{\omega} - \omega| \leq \epsilon'/6$ in time $\text{poly}(\text{size}(\epsilon'))$. Similarly, x^* is the unique root of the polynomial $1 + \lambda x^d - \frac{\lambda}{\lambda'}$ in the ball $B(p^*, r^*/2)$ and therefore we can compute $\hat{x}^* \in \mathbb{C}_\mathbb{Q}$ such that $|\hat{x}^* - x^*| \leq L\epsilon'/6$ in time $\text{poly}(\text{size}(\lambda'), \epsilon')$. Then, for $i \in [t]$, we check whether $|\hat{w}_i - \hat{\omega}| \leq 2\epsilon'/3$ and $|f^N(\hat{w}_i) - \hat{x}^*| \leq 2L\epsilon'/3$; the check must pass for at least one $\hat{w} \in \{\hat{w}_1, \dots, \hat{w}_t\}$, and using the triangle inequality we obtain that \hat{w} satisfies (5.38).

$\text{poly}(\text{size}(\lambda', \epsilon))$. Moreover, we can compute the values $Z_{G,v}^{\text{in}}(\lambda)$ and $Z_{G,v}^{\text{out}}(\lambda)$ at the same time.

Let w be such that $\lambda w = \frac{Z_{G,v}^{\text{in}}(\lambda)}{Z_{G,v}^{\text{out}}(\lambda)}$, so that $|w - \hat{w}| \leq \epsilon'$. Since $r' + 2\epsilon' < r$, we obtain using (5.38) that $w \in B(\omega, r)$. Further, by applying (5.32) for $x_1 = w$ and $x_2 = \hat{w}$, we get that $|f^N(w) - f^N(\hat{w})| \leq L\epsilon'$ and therefore by the triangle inequality and (5.38) we have

$$(5.39) \quad |f^N(w) - f^N(w^*)| \leq 2L\epsilon'.$$

We will next show that

$$(5.40) \quad \left| \frac{\lambda}{1 + \lambda(f^N(w))^d} - \lambda' \right| \leq \epsilon.$$

Since $f^N(w^*) = x^*$ and $|x^*| \leq |p^*| + r^*/2$, we can conclude (using (5.39)) that $|f^N(w^*)|, |f^N(w)| \leq |p^*| + r^*$. In turn, this gives

$$(5.41) \quad \begin{aligned} |(f^N(w))^d - (f^N(w^*))^d| &= |f^N(w) - f^N(w^*)| \cdot \left| \sum_{j=0}^{d-1} (f^N(w))^j (f^N(w^*))^{d-1-j} \right| \\ &\leq 2dL(|p^*| + r^*)^{d-1}\epsilon' \leq 1/(2|\lambda'|), \end{aligned}$$

where in the last inequality we used that $\epsilon' \leq \frac{1}{4dL|\lambda'|(|p^*| + r^*)^{d-1}}$. From (5.36), (5.41), and the triangle inequality, we obtain that

$$(5.42) \quad \left| 1 + \lambda(f^N(w))^d \right| \geq \left| 1 + \lambda(f^N(w^*))^d \right| - \left| \lambda(f^N(w))^d - \lambda(f^N(w^*))^d \right| \geq \frac{|\lambda|}{2|\lambda'|}$$

and, therefore,

$$\begin{aligned} \left| \frac{\lambda}{1 + \lambda(f^N(w))^d} - \lambda' \right| &= \frac{|\lambda|^2 \cdot |(f^N(w))^d - (f^N(w^*))^d|}{|1 + \lambda(f^N(w))^d| \cdot |1 + \lambda(f^N(w^*))^d|} \\ &= \frac{|\lambda'\lambda| \cdot |(f^N(w))^d - (f^N(w^*))^d|}{|1 + \lambda(f^N(w))^d|} \leq 4dL|\lambda'|^2(|p^*| + r^*)^{d-1}\epsilon' \leq \epsilon, \end{aligned}$$

where in the last equality we used (5.36), in the second to last inequality we used (5.41) and (5.42), and in the last inequality we used the choice of ϵ' . This concludes the proof of (5.40). To ensure that certain partition functions are nonzero, we will need the following additional fact for w , namely, that

$$(5.43) \quad f^n(w) \neq p_1, \dots, p_d \text{ for all } n = 0, 1, \dots, N.$$

For $n = 0, 1, \dots, N-1$, this just follows from (5.31) and the fact that $w \in B(\omega, r)$. For $n = N$, we have from (5.39) and $f^N(w^*) = x^*$ that $f^N(w)$ is within distance $2L\epsilon' \leq r^*/2$ from $x^* \in B(p^*, r^*/2)$ and therefore $f^N(w) \in B(p^*, r^*)$. This implies that for all poles $p \neq p^*$ it holds that $f^N(w) \neq p$ (cf. (5.33)). For the pole p^* , we have from (5.39) that $f^N(w)$ is within distance $2L\epsilon' \leq \tau/2$ from $f^N(w^*)$ and therefore, using (5.37), we have that $|f^N(w) - p^*| \geq \tau/2 > 0$, i.e., $f^N(w) \neq p^*$. This finishes the proof of (5.43).

In light of (5.40), we next focus on implementing the activity $\lambda/(1 + \lambda(f^N(w))^d)$ using a bipartite graph of maximum degree Δ . For $h = 0, 1, \dots, N$, let T_h denote the d -ary tree of height h and denote the root of the tree by u_h . Let G_h be the bipartite graph of maximum degree Δ obtained from T_h by taking, for each leaf l of T_h , d distinct copies of the graph G (which implements λw) and identifying l with the d copies of the terminal v of G . Then, using that $\lambda w = Z_{G,v}^{\text{in}}(\lambda)/Z_{G,v}^{\text{out}}(\lambda)$, we have

$$(5.44) \quad \begin{aligned} Z_{G_0,u_0}^{\text{in}}(\lambda) &= \lambda(Z_{G,v}^{\text{in}}(\lambda)/\lambda)^d, & Z_{G_0,u_0}^{\text{out}}(\lambda) &= (Z_{G,v}^{\text{out}}(\lambda))^d, \\ Z_{G_0}(\lambda) &= Z_{G_0,u_0}^{\text{in}}(\lambda) + Z_{G_0,u_0}^{\text{out}}(\lambda) = (Z_{G,v}^{\text{out}}(\lambda))^d(1 + \lambda w^d). \end{aligned}$$

(5.43) ensures that $w \neq p_1, \dots, p_d$ so we have $1 + \lambda w^d \neq 0$. We therefore obtain from (5.44) that

$$(5.45) \quad Z_{G_0}(\lambda) \neq 0 \text{ and } \frac{Z_{G_0,u_0}^{\text{out}}(\lambda)}{Z_{G_0}(\lambda)} = \frac{(Z_{G,v}^{\text{out}}(\lambda))^d}{(Z_{G,v}^{\text{out}}(\lambda))^d(1 + \lambda w^d)} = f(w).$$

Further, for $h = 1, \dots, N$ it holds that

$$(5.46) \quad \begin{aligned} Z_{G_h,u_h}^{\text{in}}(\lambda) &= \lambda(Z_{G_{h-1},u_{h-1}}^{\text{out}}(\lambda))^d, & Z_{G_h,u_h}^{\text{out}}(\lambda) &= (Z_{G_{h-1}}(\lambda))^d, \\ Z_{G_h}(\lambda) &= Z_{G_h,u_h}^{\text{in}}(\lambda) + Z_{G_h,u_h}^{\text{out}}(\lambda) = (Z_{G_{h-1}}(\lambda))^d \left(1 + \lambda \frac{(Z_{G_{h-1},u_{h-1}}^{\text{out}}(\lambda))^d}{(Z_{G_{h-1}}(\lambda))^d} \right). \end{aligned}$$

We will show by induction that for all $h = 0, 1, \dots, N$ it holds that

$$(5.47) \quad Z_{G_h}(\lambda) \neq 0 \text{ and } \frac{Z_{G_h,u_h}^{\text{out}}(\lambda)}{Z_{G_h}(\lambda)} = f^{h+1}(w).$$

For $h = 0$, this is just (5.45). Assume that it holds for $h - 1$; we have by (5.46) and the induction hypothesis that

$$Z_{G_h}(\lambda) = (Z_{G_{h-1}}(\lambda))^d (1 + \lambda(f^h(w))^d) \neq 0,$$

where the disequality follows from $Z_{G_{h-1}}(\lambda) \neq 0$ and (5.43). We therefore obtain that

$$\frac{Z_{G_h,u_h}^{\text{out}}(\lambda)}{Z_{G_h}(\lambda)} = \frac{1}{1 + \lambda \left(\frac{Z_{G_{h-1},u_{h-1}}^{\text{out}}(\lambda)}{Z_{G_{h-1}}(\lambda)} \right)^d} = \frac{1}{1 + \lambda(f^h(w))^d} = f^{h+1}(w),$$

completing the induction and the proof of (5.47). For $h = N$, (5.47) gives that $Z_{G_N}(\lambda) \neq 0$ and

$$(5.48) \quad \frac{Z_{G_N,u_N}^{\text{out}}(\lambda)}{Z_{G_N}(\lambda)} = f^{N+1}(w) = f(f^N(w)) = \frac{1}{1 + \lambda(f^N(w))^d}.$$

Consider the graph G' obtained from G_N by adding a new vertex u' whose single neighbor is the vertex u_N . Then, we have that

$$(5.49) \quad Z_{G',u'}^{\text{in}}(\lambda) = \lambda Z_{G_N,u_N}^{\text{out}}(\lambda), \quad Z_{G',u'}^{\text{out}}(\lambda) = Z_{G_N}(\lambda) \neq 0.$$

Using this in conjunction with (5.48), we conclude that

$$\frac{Z_{G',u'}^{\text{in}}(\lambda)}{Z_{G',u'}^{\text{out}}(\lambda)} = \frac{\lambda Z_{G_N,u_N}^{\text{out}}(\lambda)}{Z_{G_N}(\lambda)} = \frac{\lambda}{1 + \lambda(f^N(w))^d}.$$

From this and (5.40), we obtain that G' is a bipartite graph of maximum degree Δ with terminal u' which implements λ' with accuracy ϵ . Moreover, using (5.44), (5.46), and (5.49), we can also compute the values $Z_{G',u'}^{\text{in}}(\lambda)$, $Z_{G',u'}^{\text{out}}(\lambda)$.

The remaining cases of the algorithm (Cases II and III below) are almost identical to Cases II and III of the algorithm in Proposition 2.11 for the real case, so we focus on the main differences (which mostly amount to modifying the upper and lower bounds for the relevant quantities). To align with the notation there, let G_4 be a bipartite graph of maximum degree at most Δ with terminal v_4 that implements a constant activity λ'_4 with

$$(5.50) \quad |\lambda'_4| > (M+1)(M+2), \text{ where } \lambda'_4 = \frac{Z_{G_4,v_4}^{\text{in}}(\lambda)}{Z_{G_4,v_4}^{\text{out}}(\lambda)}.$$

Note, this implementation can be done using the algorithm for Case I. We next give the details of the algorithm for the remaining cases.

Case II (small $|\lambda'|$): $|\lambda'| \leq |\lambda|/(M+1)$. We first assume that $\lambda' \neq 0$. Let $\hat{\lambda}$ be such that

$$(5.51) \quad \frac{\lambda}{1 + \hat{\lambda}} = \lambda' \text{ so that } \hat{\lambda} = \frac{\lambda}{\lambda'} - 1.$$

Let $\hat{\epsilon} = \epsilon \cdot \min\{\frac{|\lambda|}{2|\lambda'|}, \frac{|\lambda|}{2|\lambda'|^2}, 1\}$ and let ϵ' be a rational less than $\hat{\epsilon}$ so that $\text{size}(\epsilon') = \text{poly}(\text{size}(\lambda', \epsilon))$.

Using the assumption $|\lambda'| \leq |\lambda|/(M+1)$ and the triangle inequality, we have that $|\hat{\lambda}| \geq \frac{|\lambda|}{|\lambda'|} - 1 \geq M$. Therefore, by Case I, we can construct in time $\text{poly}(\text{size}(\lambda', \epsilon))$ a bipartite graph G of maximum degree Δ with terminal v that implements $\hat{\lambda}$ with accuracy ϵ' , and we can compute the values $Z_{G,v}^{\text{in}}(\lambda)$ and $Z_{G,v}^{\text{out}}(\lambda)$ at the same time. Let $\lambda'' = \frac{Z_{G,v}^{\text{in}}(\lambda)}{Z_{G,v}^{\text{out}}(\lambda)}$, so that $|\lambda'' - \hat{\lambda}| \leq \epsilon'$. Using this and (5.51), we have

$$(5.52) \quad |1 + \lambda''| \geq |1 + \hat{\lambda}| - |(\lambda'' - \hat{\lambda})| \geq \frac{|\lambda|}{|\lambda'|} - \epsilon' \geq \frac{|\lambda|}{2|\lambda'|}$$

and, therefore,

$$(5.53) \quad \left| \frac{\lambda}{1 + \lambda''} - \lambda' \right| = \frac{|\lambda'| |\lambda'' - \hat{\lambda}|}{|1 + \lambda''|} \leq \frac{2|\lambda'|^2}{|\lambda|} |\lambda'' - \hat{\lambda}| \leq \epsilon.$$

Now, let G' be the bipartite graph obtained from G by adding a new vertex u whose single neighbor is the terminal v of G . Then, just as in Case II of Proposition 2.11 we can conclude that G' with terminal u is a bipartite graph of maximum degree Δ which implements λ' with accuracy ϵ , and we can also compute the values $Z_{G',u}^{\text{in}}(\lambda)$, $Z_{G',u}^{\text{out}}(\lambda)$. The case $\lambda' = 0$ can be handled using the above technique by implementing the activity $\epsilon''/2 \neq 0$ with accuracy $\epsilon''/2$, where $\epsilon'' > 0$ is a rational less than $\min\{\epsilon, |\lambda|/(M+1)\}$ such that $\text{size}(\epsilon'') = \text{poly}(\text{size}(\epsilon))$; see Case II of the proof of Proposition 2.11 for more details.

Case III (moderate $|\lambda'|$): $|\lambda|/(M+1) < |\lambda'| < M$. Let x^* be such that

$$(5.54) \quad \frac{\lambda}{1 + \lambda'_4 x^*} = \lambda' \text{ so that } x^* = \frac{1}{\lambda'_4} \left(\frac{\lambda}{\lambda'} - 1 \right).$$

Let $\hat{\epsilon} = \epsilon \cdot \min\{\frac{|\lambda|^2}{2M|\lambda'_4|}, \frac{|\lambda|^2}{2M^2|\lambda'_4|}, 1\}$ and let ϵ' be a rational less than $\hat{\epsilon}$ so that $\text{size}(\epsilon') = \text{poly}(\text{size}(\lambda', \epsilon))$.

Using the assumption $|\lambda'| > |\lambda|/(M+1)$ and $|\lambda'_4| > (M+1)(M+2)$ (see (5.50)), we have that $|x^*| \leq 1/(M+1)$. By the algorithm for Case II, we can implement the activity λx^* with precision ϵ' in time $\text{poly}(\text{size}(\lambda x^*, \epsilon')) = \text{poly}(\text{size}(\lambda', \epsilon))$, i.e., we can construct a bipartite graph G of maximum degree at most Δ with terminal v such that, for $\lambda'' := Z_{G,v}^{\text{in}}(\lambda)/Z_{G,v}^{\text{out}}(\lambda)$, it holds that

$$(5.55) \quad |\lambda'' - \lambda x^*| \leq \epsilon'.$$

Now, using (5.54) and (5.55), we have by the triangle inequality that

$$(5.56) \quad |1 + \lambda'_4(\lambda''/\lambda)| \geq |1 + \lambda'_4 x^*| - |\lambda'_4(x^* - \lambda''/\lambda)| \geq \frac{|\lambda|}{M} - \frac{|\lambda'_4| \epsilon'}{|\lambda|} \geq \frac{|\lambda|}{2M}$$

and, therefore,

$$(5.57) \quad \left| \frac{\lambda}{1 + \lambda'_4(\lambda''/\lambda)} - \lambda' \right| = \left| \frac{\lambda}{1 + \lambda'_4(\lambda''/\lambda)} - \frac{\lambda}{1 + \lambda'_4 x^*} \right| = \frac{|\lambda| |\lambda'_4| |x^* - (\lambda''/\lambda)|}{|1 + \lambda'_4 x^*| \cdot |1 + \lambda'_4(\lambda''/\lambda)|} \\ = \frac{|\lambda'| |\lambda'_4| |x^* - (\lambda''/\lambda)|}{|1 + \lambda'_4(\lambda''/\lambda)|} \leq 2M^2 \frac{|\lambda'_4|}{|\lambda|^2} |\lambda x^* - \lambda''| \leq \epsilon,$$

where in the last equality we used (5.54), in the second to last inequality we used (5.56) and $|\lambda'| \leq M$, and in the last inequality we used (5.55) and the choice of ϵ' .

Recall from (5.50) that G_4 is a bipartite graph of maximum degree Δ with terminal v_4 which implements the activity λ'_4 . Let G' be the bipartite graph obtained by taking a copy of G_4 and G and identifying the terminals v_4, v into a single vertex which we label u' . Further, consider the graph G'' obtained from G' by adding a new vertex u'' whose single neighbor is the vertex u' . Then, just as in Case III of Proposition 2.11, using (5.57) we can conclude that G'' is a bipartite graph of maximum degree at most Δ with terminal u'' which implements λ' with accuracy ϵ , and we can also compute the values $Z_{G',u}^{\text{in}}(\lambda), Z_{G',u}^{\text{out}}(\lambda)$.

This completes the three different cases of the algorithm, thus completing the proof of Proposition 2.2. \square


6. #P-hardness. In order to prove Theorems 1.1 and 1.3 we first prove #P-hardness of multivariate versions of our problems. Instead of insisting that every vertex has activity λ , we allow the activities to be drawn from the set

$$(6.1) \quad \mathcal{L}(\lambda) = \{\lambda, -\lambda - 1, -1, 1\}.$$

Name #MVHardCoreNorm(λ, Δ, K).

Instance A graph G with maximum degree at most Δ . An activity vector $\lambda = \{\lambda_v\}_{v \in V}$, such that, for each $v \in V$, $\lambda_v \in \mathcal{L}(\lambda)$. For every vertex $v \in V$ with $\lambda_v \neq \lambda$, the degree of v in G must be at most 2.

Output If $|Z_G(\lambda)| = 0$ then the algorithm may output any rational number. Otherwise, it must output a rational number \hat{N} such that $\hat{N}/K \leq |Z_G(\lambda)| \leq K\hat{N}$.

FIG. 2. The binary equality gadgets B_i and B'_i used in the proof of Lemma 6.3.

Name $\#MVHardCoreArg(\lambda, \Delta, \rho)$.

Instance A graph $G = (V, E)$ with maximum degree at most Δ . An activity vector $\lambda = \{\lambda_v\}_{v \in V}$ such that, for each $v \in V$, $\lambda_v \in \mathcal{L}(\lambda)$. For every vertex $v \in V$ with $\lambda_v \neq \lambda$, the degree of v in G must be at most 2.

Output If $Z_G(\lambda) = 0$ then the algorithm may output any rational number. Otherwise, it must output a rational number \hat{A} such that, for some $a \in \arg(Z_G(\lambda))$, $|\hat{A} - a| \leq \rho$.

6.1. Reducing the degree using equality gadgets. Given a graph $B = (V, E)$ and two subsets T_{in} and T of the vertex set V satisfying $T_{\text{in}} \subseteq T$, let $\mathcal{I}_{B, T, T_{\text{in}}}$ denote the set of independent sets I of B such that $I \cap T = T_{\text{in}}$. Let $Z_{B, T, T_{\text{in}}}(\lambda) = \sum_{I \in \mathcal{I}_{B, T, T_{\text{in}}}} \lambda^I$. We use similar notation when activities are nonuniform. We start by introducing “equality” gadgets.

LEMMA 6.1. Let $\lambda \in \mathbb{C}$ be such that $\lambda \neq -1, 0$. Let B_i and B'_i be the graphs in Figure 2 with activity vectors λ and λ' , respectively, and set $T_i = \{u_i, v_i\}$, $T'_i = \{s_i, t_i\}$. Then,

$$Z_{B_i, T_i, \{u_i\}}(\lambda) = Z_{B_i, T_i, \{v_i\}}(\lambda) = 0, \quad Z_{B_i, T_i, \emptyset}(\lambda) = Z_{B_i, T_i, T_i}(\lambda) = C := \lambda(1 + \lambda) \neq 0,$$

and

$$Z_{B'_i, T'_i, \{s_i\}}(\lambda') = Z_{B'_i, T'_i, \{t_i\}}(\lambda') = 0, \quad Z_{B'_i, T'_i, \emptyset}(\lambda') = \frac{1}{\lambda} Z_{B'_i, T'_i, T'_i}(\lambda') = 1.$$

Proof. By enumerating the independent sets of B_i , we have

$$(6.2) \quad \begin{aligned} Z_{B_i, T_i, \{u_i\}}(\lambda) &= \lambda_{u_i}(1 + \lambda_{y_i} + \lambda_{z_i}), & Z_{B_i, T_i, \{v_i\}}(\lambda) &= \lambda_{v_i}(1 + \lambda_{x_i} + \lambda_{z_i}), \\ Z_{B_i, T_i, \emptyset}(\lambda) &= 1 + \lambda_{x_i} + \lambda_{y_i} + \lambda_{z_i} + \lambda_{x_i}\lambda_{y_i}, & Z_{B_i, T_i, T_i}(\lambda) &= \lambda_{u_i}\lambda_{v_i}(1 + \lambda_{z_i}). \end{aligned}$$

Observe that B'_i is obtained from B_i by “appending” the vertices s_i and t_i . Therefore,

$$(6.3) \quad \begin{aligned} Z_{B'_i, T'_i, \emptyset}(\lambda') &= Z_{B_i, T_i, \emptyset}(\lambda') + Z_{B_i, T_i, \{u_i\}}(\lambda') + Z_{B_i, T_i, \{v_i\}}(\lambda') + Z_{B_i, T_i, T_i}(\lambda'), \\ Z_{B'_i, T'_i, \{s_i\}}(\lambda') &= \lambda_{s_i}(Z_{B_i, T_i, \emptyset}(\lambda') + Z_{B_i, T_i, \{v_i\}}(\lambda')), \\ Z_{B'_i, T'_i, \{t_i\}}(\lambda') &= \lambda_{t_i}(Z_{B_i, T_i, \emptyset}(\lambda') + Z_{B_i, T_i, \{u_i\}}(\lambda')), \\ Z_{B'_i, T'_i, T'_i}(\lambda') &= \lambda_{s_i}\lambda_{t_i}Z_{B_i, T_i, \emptyset}(\lambda'). \end{aligned}$$

Using (6.2), we have that

$$Z_{B_i, T_i, \emptyset}(\lambda') = 1, \quad Z_{B_i, T_i, \{u_i\}}(\lambda') = Z_{B_i, T_i, \{v_i\}}(\lambda') = -1, \quad Z_{B_i, T_i, T_i}(\lambda') = 2.$$

Plugging this into (6.3) concludes the proof of the lemma. \square

The following lemma defines the gadgets for the case $\lambda = -1$.

LEMMA 6.2. *Let $\lambda = -1$. Let B_i be the path of length six with endpoints u_i, v_i , with activity vector λ , where every vertex has activity $\lambda = -1$, apart from the endpoint v_i which has activity $+1$. Let B'_i be the graph in Figure 2 with the activity vector λ' given there. Set $T_i = \{u_i, v_i\}$ and $T'_i = \{s_i, t_i\}$. Then,*

$$Z_{B_i, T_i, \{u_i\}}(\lambda) = Z_{B_i, T_i, \{v_i\}}(\lambda) = 0, \quad Z_{B_i, T_i, \emptyset}(\lambda) = Z_{B_i, T_i, T_i}(\lambda) = C := 1,$$

and

$$Z_{B'_i, T'_i, \{s_i\}}(\lambda') = Z_{B'_i, T'_i, \{t_i\}}(\lambda') = 0, \quad Z_{B'_i, T'_i, \emptyset}(\lambda') = \frac{1}{\lambda} Z_{B'_i, T'_i, T'_i}(\lambda') = 1.$$

Proof. Note that both graphs are paths of length six. We can therefore use the formula in (6.3). The lemma therefore follows by just making the substitutions. \square

The following lemma shows how to use the equality gadgets to replace high-degree vertices with equivalent subgraphs made up of low-degree vertices. For a graph G and a vertex v in G , we denote by $d_v(G)$ the number of neighbors of v .

LEMMA 6.3. *Let $\lambda \in \mathbb{C}_{\neq 0}$. If $\lambda = -1$, let $C = 1$; otherwise, let $C = \lambda(1 + \lambda) \neq 0$.*

Suppose that $G = (V, E)$ is a graph with an activity vector $\lambda = \{\lambda_v\}_{v \in V}$ such that, for every vertex $v \in V$, we have $\lambda_v \in \{1, \lambda\}$. Let $U_1 = \{v \in V \mid \lambda_v = 1\}$ and $U_\lambda = \{v \in V \mid \lambda_v = \lambda\}$. Consider an arbitrary set $S \subseteq V$. Then, there is a set S' of vertices (distinct from V) and a graph $G' = (V', E')$ with an activity vector $\lambda' = \{\lambda'_v\}_{v \in V'}$ such that

$$Z_G(\lambda) = \frac{Z_{G'}(\lambda')}{\prod_{v \in S \cap U_1} C^{d_v(G)} \prod_{v \in S \cap U_\lambda} C^{d_v(G)-1}}.$$

Furthermore,

- $V' = S' \cup (V \setminus S)$ and $|S'|$ is at most $8|V|^2$;
- every vertex $v \in S'$ has $d_v(G') \leq 3$ and $\lambda'_v \in \mathcal{L}(\lambda)$. If $\lambda'_v \neq \lambda$ then $d_v(G') \leq 2$;
- every vertex in $V(G) \setminus S$ has $d_v(G) = d_v(G')$ and $\lambda'_v = \lambda_v$.

Proof. To prove the lemma, we can assume that $|S| = 1$. (To prove the lemma for a larger set S we just repeatedly apply the singleton-set version to each of the vertices in S .) So let $S = \{v\}$ and, for convenience, let $d = d_v(G)$ be the degree of v in G . There are two cases.

Case 1: $v \in U_1$. Assume first that $d \neq 1$. In this case, S' will be the union of the vertices in the gadgets B_1, \dots, B_d by identifying vertex v_i of B_i with vertex u_{i+1} of B_{i+1} for each $i = 1, \dots, d$ (we will use the conventions that $u_{d+1} \equiv u_1$, $B_{d+1} \equiv B_1$, and $B_0 \equiv B_d$). To construct G' from G we replace v with the union of these gadgets. If a vertex w is in exactly one of these gadgets, then the activity λ'_w will be inherited from the gadget. Also, w will have no other neighbors in G' (other than the neighbors in its gadget). Now, for $i = 1, \dots, d$, the vertex u_i is in two gadgets, namely, B_i and B_{i-1} . In addition to having its gadget neighbors, this vertex u_i will be connected to the i th neighbor of v in G . Then we will set $\lambda'_{u_i} = \lambda$, since this is the product of the activities inherited from B_i and B_{i+1} . Note from Lemma 6.1 (and Lemma 6.2 in the case $\lambda = -1$) that, in the resulting graph G' , in any independent set of nonzero weight, either all v_i 's are in the independent set, or all v_i 's are missing. In each of these cases, we get a factor of C^d in the partition function. The construction for $d = 1$ is analogous, i.e., we replace v with the gadget B_1 , but now we do not do any

identification of vertices and we use u_1 to connect to the neighbor of v in G ; further, every vertex in B_1 retains its activity in G' . As before, we conclude that we get a factor of C in the partition function.

Case 2: $v \in U_\lambda$. Assume first that $d \neq 1$. In this case, S' will be the union of the gadgets $B_1, \dots, B_{d-1}, B'_d$. We will further identify vertex v_i of B_i with vertex u_{i+1} for each $i = 1, \dots, d-2$; for $i = d-1$, we will identify vertex v_{d-1} of B_{d-1} with vertex s_d of B'_d and vertex t_d of B'_d with vertex u_1 of B_1 . To construct G' from G we will replace v with the union of these gadgets. The assignment of activities is the same as in Case 1; the only difference is in the construction of the graph G' where now, for $i = 1, \dots, d-1$, the i th neighbor of v connects to the vertex u_i while the d th neighbor of v connects to the vertex s_d . From Lemma 6.1 (and Lemma 6.2 in the case $\lambda = -1$), we have that, in the resulting graph G' , in any independent set of nonzero weight, either all v_i 's are in the independent set, or all v_i 's are missing. If they are all in, then we get a factor of $C^{d-1}\lambda$ in the partition function; otherwise, we get a factor of C^{d-1} . The construction for $d = 1$ is analogous, i.e., we replace v with the gadget B'_1 , but now we do not do any identification of vertices and we use s_1 to connect to the neighbor of v in G ; further, every vertex in B'_1 retains its activity in G' . As before, we conclude that in any independent set of nonzero weight, either all v_i 's are in the independent set (contributing a factor of λ), or all v_i 's are missing (contributing a factor of 1).

This concludes the proof of Lemma 6.3. \square

6.2. #P-hardness of the multivariate problem.

THEOREM 6.4. *Let $\Delta \geq 3$ and $\lambda \in \mathbb{C}_\mathbb{Q}$ be a complex number such that $\lambda \notin (\Lambda_\Delta \cup \mathbb{R}_{\geq 0})$. Then, for $K = 1.02$, $\#\text{MVHardCoreNorm}(\lambda, \Delta, K)$ is #P-hard. Also, for $\rho = 9\pi/24$, $\#\text{MVHardCoreArg}(\lambda, \Delta, \rho)$ is #P-hard.*

Proof. Counting the number of independent sets of an input graph is a well known #P-hard problem. We will reduce this to both problems. To do this, let H be an n -vertex graph, and let $N = Z_H(1)$ denote the number of independent sets of H . Our goal is to show how to use an oracle for $\#\text{MVHardCoreNorm}(\lambda, \Delta, K)$ or $\#\text{MVHardCoreArg}(\lambda, \Delta, \rho)$ to compute N . Let $\epsilon = 1/50$ and $\eta = 1/35$.

Consider any rational number M in the range $0 \leq M \leq 2^n$. Let J_M be a graph with terminal v and maximum degree at most Δ that implements $-M$ with accuracy ϵ ; note, by applying Propositions 2.2 and 2.11 (for complex and real λ , respectively), there is an algorithm to construct J_M in time $\text{poly}(\text{size}(M, \epsilon)) = \text{poly}(\text{size}(M))$ that also outputs the exact values of $Z_{J_M, v}^{\text{in}}(\lambda), Z_{J_M, v}^{\text{out}}(\lambda)$. Let

$$(6.4) \quad \lambda_M := Z_{J_M, v}^{\text{in}}(\lambda) / Z_{J_M, v}^{\text{out}}(\lambda), \text{ and note that } Z_{J_M, v}^{\text{out}}(\lambda) \neq 0 \text{ and } |\lambda_M + M| \leq \epsilon.$$

Let G_M be the graph formed by taking the disjoint union of H and J_M and attaching the terminal v of J_M to every vertex in H . Let λ_M be the activity vector for G_M obtained by setting the activities of vertices originally belonging to H equal to 1 and the activities of vertices originally belonging to J_M equal to λ . Note that

$$(6.5) \quad Z_{G_M}(\lambda_M) = Z_{J_M, v}^{\text{in}}(\lambda) + Z_{J_M, v}^{\text{out}}(\lambda)Z_H(1) = Z_{J_M, v}^{\text{in}}(\lambda) + Z_{J_M, v}^{\text{out}}(\lambda)N.$$

Let G'_M and λ'_M be the graph and activity vector that are constructed by applying Lemma 6.3 to G_M with $S = V(H) \cup \{v\}$. Note that the size of G_M is at most a polynomial in $\text{size}(M)$ so (from Lemma 6.3) the size of G'_M is also at most a

polynomial in $\text{size}(M)$. From Lemma 6.3, we have that

$$(6.6) \quad Z_{G_M}(\lambda_M) = Z_{G'_M}(\lambda'_M)/W_H, \text{ where } W_H := C^{n-1} \prod_{v \in V(H)} C^{d_v(H)},$$

and C is the constant in Lemma 6.3. Furthermore, every vertex v of G'_M has degree at most Δ and every vertex v of G'_M with $\lambda'_v \neq \lambda$ has degree at most 2 in G'_M . Thus, G'_M is a valid input to $\#\text{MVHardCoreNorm}(\lambda, \Delta, K)$ and $\#\text{MVHardCoreArg}(\lambda, \Delta, \rho)$. Moreover, combining (6.5), (6.6) and dividing through by $Z_{J_M, v}^{\text{out}}(\lambda)$, we have

$$(6.7) \quad \lambda_M + N = \frac{Z_{G'_M}(\lambda'_M)}{W_H Z_{J_M, v}^{\text{out}}(\lambda)} =: f_M.$$

Part one: $\#\text{P}$ -hardness of $\#\text{MVHardCoreNorm}(\lambda, \Delta, 1.02)$.

By the triangle inequality (in the form $||a| - |b|| \leq |a + b|$), we have

$$||f_M| - |M - N|| \leq |\lambda_M + N + M - N| = |\lambda_M + M| \leq \epsilon,$$

where in the last inequality we used (6.4). Therefore, $|N - M| - \epsilon \leq |f_M| \leq |N - M| + \epsilon$.

Consider M so that $|N - M| \geq 1$. Then $|f_M|$ is not 0. From the definition of f_M in (6.7), this means that $|Z_{G'_M}(\lambda'_M)| \neq 0$ and hence we can use the oracle for $\#\text{MVHardCoreNorm}(\lambda, \Delta, 1.02)$ to produce an estimate for $|Z_{G'_M}(\lambda'_M)|$ within a factor of 1.02. We can also obtain an estimate of the value $|W_H Z_{J_M, v}^{\text{out}}(\lambda)|$ within a factor of $(1 + 7\eta/8)/1.02 = 1.025/1.02$, since $Z_{J_M, v}^{\text{out}}(\lambda)$ is output by the algorithm from Propositions 2.2 and 2.11. Combining these, we obtain an estimate \hat{f}_M for $|f_M|$ satisfying $(1 - \eta)|f_M| \leq \hat{f}_M \leq (1 + \eta)|f_M|$. We now use the binary search technique of [7].

The invariant that we will maintain is that we have an interval $[M_{\text{start}}, M_{\text{end}}]$ of real numbers with $M_{\text{start}} \leq N \leq M_{\text{end}}$. Initially, $M_{\text{start}} = 0$ and $M_{\text{end}} = 2^n$. Let $\ell = M_{\text{end}} - M_{\text{start}}$. If $\ell < 1$ then there is only one integer between M_{start} and M_{end} , so the value of N is known.

Suppose $\ell \geq 1$. For $i \in \{0, \dots, 8\}$, let $M_i = M_{\text{start}} + i\ell/8$. For $i \in \{0, \dots, 7\}$, let s_i be the sign (positive, negative, or zero) of $\hat{f}_{M_i} - \hat{f}_{M_{i+1}}$.

First, consider $i \in \{0, \dots, 7\}$ and suppose $N \geq M_{i+2}$. Then,

$$\begin{aligned} \hat{f}_{M_i} - \hat{f}_{M_{i+1}} &\geq (1 - \eta)|f_{M_i}| - (1 + \eta)|f_{M_{i+1}}| \\ &\geq (1 - \eta)(N - M_i - \epsilon) - (1 + \eta)(N - M_i - \ell/8 + \epsilon) \\ &= (1 + \eta)\ell/8 - 2\eta(N - M_i) - 2\epsilon \\ &\geq (1 + \eta)\ell/8 - 2\eta\ell - 2\epsilon. \end{aligned}$$

This is positive since $2\eta < (1 + \eta)/16$ and $\epsilon \leq \eta\ell$, so s_i is positive. Similarly, if $N \leq M_{i-1}$ then

$$\begin{aligned} \hat{f}_{M_i} - \hat{f}_{M_{i+1}} &\leq (1 + \eta)|f_{M_i}| - (1 - \eta)|f_{M_{i+1}}| \\ &\leq (1 + \eta)(M_i - N + \epsilon) - (1 - \eta)(M_i - N + \ell/8 - \epsilon) \\ &= -(1 - \eta)\ell/8 + 2\eta(M_i - N) + 2\epsilon \\ &\leq -(1 - \eta)\ell/8 + 2\eta\ell + 2\epsilon, \end{aligned}$$

so s_i is negative.

Now, consider i^* so that $M_{i^*} \leq N \leq M_{i^*+1}$. Then s_0, \dots, s_{i^*-2} are plus and s_{i^*+2}, \dots, s_7 are minus. So we have a (possibly nonempty) sequence of pluses followed by three arbitrary signs followed by a (possibly nonempty) sequence of minuses. If there are three minuses in a row at the end of the sequence, we must have $N \leq M_7$ (otherwise s_5 would have been a plus). So we can shrink the interval by redefining M_{end} to be M_7 . Otherwise, there are three pluses in a row at the beginning of the sequence. This means that $N \geq M_1$ (otherwise s_2 would have been negative). So, in this case, we can shrink the interval by redefining M_{start} to be M_1 . Either way, the interval shrinks to $7/8$ of its original size, so we can recurse on the new interval; after at most $\text{poly}(n)$ steps, we will have $M_{\text{end}} - M_{\text{start}} < 1$, which gives us the exact value of N .

Part two: #P-hardness of $\#\text{MVHardCoreArg}(\lambda, \Delta, 9\pi/24)$.

Consider any rational number M in the range $0 \leq M \leq 2^n$. We will show that, if $N > M + 1/7$ then there is an $a \in \arg(f_M)$ such that $-\pi/12 < a < \pi/12$. Also, if $N < M - 1/7$ then there is an $a \in \arg(f_M)$ such that $\pi - \pi/12 < a < \pi + \pi/12$.

To prove these claims, let $x_M = \lambda_M + M$ so by (6.4), $|x_M| \leq \epsilon$. Then by (6.7), $f_M = \lambda_M + N = x_M + N - M$. Suppose $N > M + 1/7$ and consider $\theta \in \arg(f_M) = \arg(x_M + N - M)$. For concreteness (by adding integer multiples of 2π if necessary), suppose that θ is in the range $[-\pi, \pi)$. Then $\tan(\theta) \leq \frac{\epsilon}{N-M} \leq 7\epsilon$. But $\tan(\pi/12) > 0.26 > 7\epsilon$. So $\theta \leq \pi/12$. Similarly, $\theta \geq -\pi/12$. The case $M > N + 1/7$ is similar (restricting θ to $[0, 2\pi)$).

Now suppose $|N - M| > 1/7$. Since $f_M = x_M + N - M$, we have $f_M \neq 0$. Since we can compute the value of $W_H Z_{J_M, v}^{\text{out}}(\lambda)$ exactly, we can also compute $\text{Arg}(W_H Z_{J_M, v}^{\text{out}}(\lambda))$ within $\pm\pi/48$. Using an oracle for $\#\text{MVHardCoreArg}(\lambda, \Delta, 9\pi/24)$ with input (G'_M, λ'_M) we thus obtain an estimate \hat{A}_M such that, for some $a \in \arg(f_M)$, $|\hat{A}_M - a| \leq 9\pi/24 + \pi/48 = 19\pi/48$.

As in Part one, we now do a binary search. Again, the invariant that we will maintain is that we have an interval $[M_{\text{start}}, M_{\text{end}}]$ of real numbers with $M_{\text{start}} \leq N \leq M_{\text{end}}$. Initially, $M_{\text{start}} = 0$ and $M_{\text{end}} = 2^n$. Let $\ell = M_{\text{end}} - M_{\text{start}}$. If $\ell < 1$ then there is only one integer between M_{start} and M_{end} , so the value of N is known.

Suppose $\ell \geq 1$. For $i \in \{0, \dots, 6\}$, let $M_i = M_{\text{start}} + i\ell/6$. Let s_i be minus if there is an integer j such that $\pi/2 < \hat{A}_{M_i} + 2\pi j < 3\pi/2$. Let s_i be plus if there is an integer j such that $-\pi/2 < \hat{A}_{M_i} + 2\pi j < \pi/2$. If neither of these occurs, then s_i is undefined.

If $N \leq M_{i-1}$ then s_i is minus. If $N \geq M_{i+1}$ then s_i is plus. So s_0, \dots, s_6 consists of a (possibly empty) sequence of pluses followed by one unknown value followed by a (possibly empty) sequence of minuses.

Suppose that s_0, s_1 , and s_2 are all plus. Then $N \geq M_1$ (otherwise s_2 would be minus). Otherwise, s_4, s_5 , and s_6 are all minus. In this case, $N \leq M_5$ (otherwise s_4 would be plus). Either way, we can shrink the interval to $5/6$ of its original length, so, as in Part one, we can discover the value of N . \square

6.3. Restricting to bipartite graphs. In this section, we reduce the problems $\#\text{MVHardCoreNorm}(\lambda, \Delta, K)$ and $\#\text{MVHardCoreArg}(\lambda, \Delta, \rho)$ to multivariate versions whose inputs are further restricted to bipartite graphs. To do this, we will need to enlarge slightly the set of activities that a vertex can have. Namely, let

$$(6.8) \quad \mathcal{L}_{\text{bip}}(\lambda) := \mathcal{L}(\lambda) \cup \{-2, -1/4\} = \{\lambda, -\lambda - 1, -1, 1, -2, -1/4\}.$$

We will consider the following problems.




FIG. 3. The gadget B''_i with activity vector λ'' used in the proof of Lemma 6.5.

Name #MVBipHardCoreNorm(λ, Δ, K).

Instance A bipartite graph G with maximum degree at most Δ . An activity vector $\lambda = \{\lambda_v\}_{v \in V}$, such that, for each $v \in V$, $\lambda_v \in \mathcal{L}_{\text{bip}}(\lambda)$. For every vertex $v \in V$ with $\lambda_v \neq \lambda$, the degree of v in G must be at most 2.

Output If $|Z_G(\lambda)| = 0$ then the algorithm may output any rational number. Otherwise, it must output a rational number \hat{N} such that $\hat{N}/K \leq |Z_G(\lambda)| \leq K\hat{N}$.

Name #MVBipHardCoreArg(λ, Δ, ρ).

Instance A bipartite graph $G = (V, E)$ with maximum degree at most Δ . An activity vector $\lambda = \{\lambda_v\}_{v \in V}$ such that, for each $v \in V$, $\lambda_v \in \mathcal{L}_{\text{bip}}(\lambda)$. For every vertex $v \in V$ with $\lambda_v \neq \lambda$, the degree of v in G must be at most 2.

Output If $Z_G(\lambda) = 0$ then the algorithm may output any rational number. Otherwise, it must output a rational number \hat{A} such that, for some $a \in \arg(Z_G(\lambda))$, $|\hat{A} - a| \leq \rho$.

LEMMA 6.5. Let B''_i be the graph in Figure 3 with activity vector λ , and set $T''_i = \{p_i, q_i\}$. Then, for any value of the activities $\lambda_{p_i}, \lambda_{q_i} \in \mathbb{C}$, it holds that

$$\begin{aligned} Z_{B''_i, T''_i, \emptyset}(\lambda'') &= 1, & Z_{B''_i, T''_i, \{p_i\}}(\lambda'') &= \lambda_{p_i}, \\ Z_{B''_i, T''_i, \{q_i\}}(\lambda'') &= \lambda_{q_i}, & Z_{B''_i, T''_i, T''_i}(\lambda'') &= 0. \end{aligned}$$

Proof. Note that B''_i is obtained from the path B'_i in Figure 2 by appending the vertices p_i, q_i , and in turn B'_i is obtained from the path B_i in Figure 2 by appending the vertices s_i, t_i . As in Lemma 6.1, we denote $T'_i = \{s_i, t_i\}$ and $T_i = \{u_i, v_i\}$. Completely analogously to (6.3), we have

$$\begin{aligned} (6.9) \quad Z_{B''_i, T''_i, \emptyset}(\lambda'') &= Z_{B'_i, T'_i, \emptyset}(\lambda'') + Z_{B'_i, T'_i, \{s_i\}}(\lambda'') + Z_{B'_i, T'_i, \{t_i\}}(\lambda'') + Z_{B'_i, T'_i, T'_i}(\lambda''), \\ Z_{B''_i, T''_i, \{p_i\}}(\lambda'') &= \lambda_{p_i} (Z_{B'_i, T'_i, \emptyset}(\lambda'') + Z_{B'_i, T'_i, \{t_i\}}(\lambda'')), \\ Z_{B''_i, T''_i, \{q_i\}}(\lambda'') &= \lambda_{q_i} (Z_{B'_i, T'_i, \emptyset}(\lambda'') + Z_{B'_i, T'_i, \{s_i\}}(\lambda'')), \\ Z_{B''_i, T''_i, T''_i}(\lambda'') &= \lambda_{p_i} \lambda_{q_i} Z_{B'_i, T'_i, \emptyset}(\lambda''). \end{aligned}$$

To compute the r.h.s. in (6.9) we will use (6.3) which expresses the relevant quantities

in terms of the gadget B_i in Figure 2. Using (6.2) and (6.3), we therefore obtain

$$Z_{B_i, T_i, \emptyset}(\lambda'') = -1/4, Z_{B_i, T_i, \{u_i\}}(\lambda'') = Z_{B_i, T_i, \{v_i\}}(\lambda'') = -1/4, Z_{B_i, T_i, T_i}(\lambda'') = 3/4, \\ Z_{B'_i, T'_i, \emptyset}(\lambda'') = 0, Z_{B'_i, T'_i, \{s_i\}}(\lambda') = Z_{B'_i, T'_i, \{t_i\}}(\lambda'') = 1, Z_{B'_i, T'_i, T'_i}(\lambda'') = -1.$$

Plugging this into (6.9) concludes the proof of the lemma. \square

THEOREM 6.6. *Let $\Delta \geq 3$ and $\lambda \in \mathbb{C}_{\mathbb{Q}}$ be a complex number such that $\lambda \notin (\Lambda_{\Delta} \cup \mathbb{R}_{\geq 0})$.*

Then, for $K = 1.02$, $\#\text{MVBipHardCoreNorm}(\lambda, \Delta, K)$ is $\#\text{P}$ -hard. Also, for $\rho = 9\pi/24$, $\#\text{MVBipHardCoreArg}(\lambda, \Delta, \rho)$ is $\#\text{P}$ -hard.

Proof. Let $G = (V, E)$ be an n -vertex graph with maximum degree at most Δ . Suppose that $\lambda = \{\lambda_v\}_{v \in V}$ is an activity vector for G such that, for each $v \in V$, $\lambda_v \in \mathcal{L}(\lambda)$. Let e_1, \dots, e_m be an arbitrary enumeration of the edges of G and suppose that $e_i = \{p_i, q_i\}$, where p_i and q_i are vertices of G . Let H be the bipartite graph constructed from G by replacing every edge e_i of G with the gadget B'_i from Figure 3 and denote by λ' the resulting activity vector on H (every vertex originally in G retains its activity in H). Observe that every vertex activity in H is from the set $\mathcal{L}_{\text{bip}}(\lambda)$. Moreover, $Z_G(\lambda) = Z_H(\lambda')$. The result therefore follows from Theorem 6.4. \square

We will need the following lower bound on $Z_G(\lambda)$, which follows from Lemma 3.6.

LEMMA 6.7. *Suppose that $\lambda \in \mathbb{C}_{\mathbb{Q}}$. Then, there exists a rational $C_{\lambda} > 1$ such that the following holds. For any n -vertex graph $G = (V, E)$ and any activity vector $\lambda = \{\lambda_v\}_{v \in V}$ such that $\lambda_v \in \mathcal{L}_{\text{bip}}(\lambda)$ for all $v \in V$, it holds that either $Z_G(\lambda) = 0$ or $|Z_G(\lambda)| > C_{\lambda}^{-n}$.*

Proof. Let $\lambda_1 = -\lambda - 1$, $\lambda_2 = -1$, $\lambda_3 = 1$, $\lambda_4 = -2$, and $\lambda_5 = -1/4$, so that $\mathcal{L}_{\text{bip}}(\lambda) = \{\lambda, \lambda_1, \dots, \lambda_5\}$.

Let $\{U_0, U_1, \dots, U_5\}$ be a partition of V such that λ_v is equal to λ if $v \in U_0$ and, for $i = 1, \dots, 5$, $\lambda_v = \lambda_i$ if $v \in U_i$. For an independent set $I \in \mathcal{I}_G$ and $i = 0, \dots, 5$, let $n_i(I) = |I \cap U_i|$ and $n_-(I) = n_1(I) + n_2(I) + n_4(I) + n_5(I)$. Then,

$$Z_G(\lambda) = \sum_{I \in \mathcal{I}_G} \lambda^{n_0(I)} \prod_{i=1}^5 \lambda_i^{n_i(I)} = \sum_{I \in \mathcal{I}_G} (-1)^{n_-(I)} 2^{n_4(I)} (1/4)^{n_5(I)} \lambda^{n_0(I)} (1 + \lambda)^{n_1(I)} \\ = \sum_{I \in \mathcal{I}_G} (-1)^{n_-(I)} 2^{n_4(I)} (1/4)^{n_5(I)} \lambda^{n_0(I)} \sum_{k=0}^{n_1(I)} \binom{n_1(I)}{k} \lambda^k.$$

Thus, we have that $4^n Z_G(\lambda)$ is an integer polynomial of λ . Moreover, observe that the absolute values of the coefficients of $4^n Z_G(\lambda)$ corresponding to an independent set $I \in \mathcal{I}_G$ sum to at most $4^n \cdot 2^n = 8^n$. Since $|\mathcal{I}_G| \leq 2^n$, we have that the absolute values of the coefficients of $4^n Z_G(\lambda)$ sum to at most $2^n \cdot 8^n = 16^n$. The result now follows by applying Liouville's inequality (cf. Lemma 3.6). \square

6.4. Reduction from the multivariate problem to the single-activity problem. The purpose of this section is to prove the following theorem.

THEOREM 6.8. *Let $\Delta \geq 3$ and $\lambda \in \mathbb{C}_{\mathbb{Q}}$ be a complex number such that $\lambda \notin (\Lambda_{\Delta} \cup \mathbb{R}_{\geq 0})$. Then there is a polynomial-time Turing reduction from the problem $\#\text{MVBipHardCoreNorm}(\lambda, \Delta, 1.02)$ to $\#\text{BipHardCoreNorm}(\lambda, \Delta, 1.01)$. There is also a polynomial-time Turing reduction from the problem $\#\text{MVBipHardCoreArg}(\lambda, \Delta, 9\pi/24)$ to the problem $\#\text{BipHardCoreArg}(\lambda, \Delta, \pi/3)$.*

Proof. Let $\lambda_1 = -\lambda - 1$, $\lambda_2 = -1$, $\lambda_3 = 1$, $\lambda_4 = -2$, $\lambda_5 = -1/4$ so that $\mathcal{L}_{\text{bip}}(\lambda) = \{\lambda, \lambda_1, \dots, \lambda_5\}$. Let $M > 1$ be a rational such that $M > \max\{|\lambda|, |\lambda_1|, \dots, |\lambda_5|\}$ and let $C_\lambda > 1$ be the rational in Lemma 6.7.

Let $G = (V, E)$ be an n -vertex bipartite graph with maximum degree at most Δ . Suppose that $\lambda = \{\lambda_v\}_{v \in V}$ is an activity vector for G such that, for each $v \in V$, $\lambda_v \in \mathcal{L}_{\text{bip}}(\lambda)$. Let $\{U_0, U_1, \dots, U_5\}$ be a partition of V such that λ_v is equal to λ if $v \in U_0$ and, for $i = 1, \dots, 5$, $\lambda_v = \lambda_i$ if $v \in U_i$. Suppose further that, for every vertex $v \in \bigcup_{i=1}^5 U_i$, the degree of v in G is at most 2.

Let $\epsilon := \frac{1}{10^4 n (4MC_\lambda)^n}$. For $i = 1, \dots, 5$, let J_i be a bipartite graph with maximum degree at most Δ with terminal v_i that implements λ_i with accuracy ϵ . Propositions 2.2 and 2.11 guarantee that J_i exists, and that it can be constructed in time $\text{poly}(\text{size}(\epsilon))$. The propositions also guarantee that $Z_{J_i, v_i}^{\text{out}}(\lambda) \neq 0$. Let $\lambda'_i = Z_{J_i, v_i}^{\text{in}}(\lambda) / Z_{J_i, v_i}^{\text{out}}(\lambda)$, so that $|\lambda'_i - \lambda_i| \leq \epsilon$. Note, we have the crude bound $|\lambda'_i| \leq 2M$ for all $i = 1, \dots, 5$.

Let λ' be the activity vector for G formed from λ by replacing every instance of λ_i with λ'_i for $i = 1, \dots, 5$. Let H be the bipartite graph constructed from G by replacing, for each $i \in [5]$, every vertex $v \in U_i$ with a (distinct) copy of J_i , relabeling the terminal v_i as v and attaching the terminal to the (at most two) neighbors of v in G (note that this is the same construction as the one in Lemma 3.2). Note that the maximum degree of H is at most Δ and, by Lemma 3.2,

$$(6.10) \quad Z_G(\lambda') = Z_H(\lambda) / C_H, \text{ where } C_H := \prod_{i=1}^5 \prod_{v \in U_i} Z_{J_i, v_i}^{\text{out}}(\lambda).$$

Note that using the output of the algorithm provided by Propositions 2.11 and 2.2, we can compute C_H exactly in time $\text{poly}(\text{size}(\epsilon))$.

We will show that, whenever $Z_G(\lambda) \neq 0$, it holds that

$$(6.11) \quad \frac{1.01}{1.015} |Z_G(\lambda')| \leq |Z_G(\lambda)| \leq \frac{1.015}{1.01} |Z_G(\lambda')|$$

and that

$$(6.12) \quad \text{there are } a \in \arg(Z_G(\lambda)) \text{ and } a' \in \arg(Z_G(\lambda')) \text{ such that } |a - a'| \leq \pi/30.$$

Before proving (6.11) and (6.12), we show that they give the desired reductions. To reduce $\#\text{MVBipHardCoreNorm}(\lambda, \Delta, 1.02)$ to $\#\text{BipHardCoreNorm}(\lambda, \Delta, 1.01)$, suppose that (G, λ) is an input to $\#\text{MVBipHardCoreNorm}(\lambda, \Delta, 1.02)$ and that $Z_G(\lambda) \neq 0$. By (6.11), we obtain that $Z_G(\lambda')$ is nonzero and, therefore, by (6.10), $|Z_H(\lambda)| \neq 0$ as well. Thus, an oracle for $\#\text{BipHardCoreNorm}(\lambda, \Delta, 1.01)$ with input H gives an approximation \hat{N} so that $\hat{N}/1.01 \leq |Z_H(\lambda)| \leq 1.01\hat{N}$. By (6.10) and (6.11), we therefore obtain that

$$\frac{\hat{N}}{1.015|C_H|} \leq |Z_G(\lambda)| \leq 1.015 \frac{\hat{N}}{|C_H|}.$$

As we noted earlier, the value C_H can be computed exactly in time $\text{poly}(\text{size}(\epsilon))$. Thus, it is easy, in time $\text{poly}(\text{size}(\epsilon))$, to compute a value \hat{C} such that

$$\frac{1.015}{1.02} \hat{C} \leq |C_H| \leq \frac{1.02}{1.015} \hat{C}.$$

Thus, the algorithm for $\# \text{MVBipHardCoreNorm}(\lambda, \Delta, 1.02)$ can return \hat{N}/\hat{C} as an output.

To reduce $\# \text{MVHardCoreArg}(\lambda, \Delta, 9\pi/24)$ to $\# \text{BipHardCoreArg}(\lambda, \Delta, \pi/3)$, suppose that (G, λ) is an input to $\# \text{MVHardCoreArg}(\lambda, \Delta, 9\pi/24)$. An oracle call to $\# \text{BipHardCoreArg}(\lambda, \Delta, \pi/3)$ with input H gives a value \hat{A} such that, for some $h \in \arg(Z_H(\lambda))$, $|\hat{A} - h| \leq \pi/3$. Consider a and a' from (6.12). By (6.10) there is a $c \in \arg(C_H)$ such that $a' = h - c$. Now, by the triangle inequality,

$$|a - (\hat{A} - c)| \leq |a - a'| + |a' - (h - c)| + |h - \hat{A}| \leq \pi/30 + 0 + \pi/3 = 11\pi/30.$$

Adding an integer multiple of 2π to both a and c on the left-hand side, we conclude that for every $\tilde{c} \in \arg(C_H)$ there exists an $\tilde{a} \in \arg(Z_G(\lambda))$ such that $|\tilde{a} - (\hat{A} - \tilde{c})| \leq 11\pi/30$. In particular, taking $\tilde{c} = \text{Arg}(C_H)$, there exists an $\tilde{a} \in \arg(Z_G(\lambda))$ such that $|\tilde{a} - (\hat{A} - \text{Arg}(C_H))| \leq 11\pi/30$. Thus, the algorithm for $\# \text{MVHardCoreArg}(\lambda, \Delta, 9\pi/24)$ can compute a value \hat{C} such that $|\hat{C} - \text{Arg}(C_H)| \leq 9\pi/24 - 11\pi/30$ and return the value $\hat{A} - \hat{C}$ as output.

So in the rest of the proof, we will establish (6.11) and (6.12). First, we show that, whenever $Z_G(\lambda) \neq 0$, it holds that

$$(6.13) \quad |Z_G(\lambda) - Z_G(\lambda')| \leq \frac{1}{10^4(C_\lambda)^n} \leq \frac{|Z_G(\lambda)|}{10^4}.$$

The rightmost inequality is immediate by Lemma 6.7. For the leftmost inequality, note that for all positive integers k and arbitrary complex numbers $x_1, y_1, \dots, x_k, y_k$ we have the telescopic expansion $\prod_{i=1}^k x_i - \prod_{i=1}^k y_i = \sum_{j=1}^k (x_j - y_j) \prod_{i=1}^{j-1} x_i \prod_{i=j+1}^k y_i$. Hence, for an arbitrary independent set $I \in \mathcal{I}_G$, we have that (using the crude bounds $|\lambda_i|, |\lambda'_i| < 2M$ and $|\lambda| < 2M$)

$$\left| \prod_{v \in I} \lambda_v - \prod_{v \in I} \lambda'_v \right| \leq \sum_{v \in I} (2M)^{|I|-1} |\lambda_v - \lambda'_v| \leq n(2M)^n \epsilon,$$

where in the last inequality we used that $|\lambda_v - \lambda'_v| \leq \epsilon$ for all $v \in V$. Since $|\mathcal{I}_G| \leq 2^n$ and $\epsilon = \frac{1}{10^4 n (4MC_\lambda)^n}$, we obtain (6.13).

Now we are ready to show (6.11) and (6.12) whenever $Z_G(\lambda) \neq 0$. In particular, by the triangle inequality and (6.13), we have that

$$\left| \frac{|Z_G(\lambda')|}{|Z_G(\lambda)|} - 1 \right| = \frac{||Z_G(\lambda')| - |Z_G(\lambda)||}{|Z_G(\lambda)|} \leq \frac{|Z_G(\lambda') - Z_G(\lambda)|}{|Z_G(\lambda)|} \leq 10^{-4},$$

which proves (6.11). In fact, using the inequality above and (6.11), it follows that the Ziv distance between $Z_G(\lambda)$ and $Z_G(\lambda')$ is at most 10^{-3} and, therefore, (6.12) follows from [7, Lemma 2.1]. \square

7. Proof of Proposition 2.6. In this section, we prove Proposition 2.6. To do this, we will focus on understanding the following type of sequences.

DEFINITION 7.1. Let $\Delta \geq 3$ and $\lambda \in \mathbb{C}_{\mathbb{Q}} \setminus \mathbb{R}$, and set $d := \Delta - 1$. A *hard-core program* is a sequence a_0, a_1, \dots , starting with $a_0 = 1$, and

$$(7.1) \quad a_k = \frac{1}{1 + \lambda a_{i_{k,1}} \cdots a_{i_{k,d}}} \quad \text{for } k \geq 1,$$

where $i_{k,1}, \dots, i_{k,d} \in \{0, \dots, k-1\}$. We say that the hard-core program generates $x \in \mathbb{C}$ if there exists an integer $k \geq 0$ such that $a_k = x$.

Our interest in hard-core programs is justified by the following lemma.

LEMMA 7.2. *Let $\lambda \in \mathbb{C}$ and $d \geq 2$. Suppose that a_0, a_1, \dots is a hard-core program. Then, for every $k \geq 1$, there exists a tree of maximum degree at most $\Delta = d + 1$ that implements the activity λa_k .*

Proof. The proof is by induction on k . For $k = 1$, we have that $i_{0,1} = \dots = i_{0,d} = 0$, so $a_1 = 1/(1 + \lambda)$. Let T be the single-edge tree $\{u, v\}$. Then, we have that

$$Z_{T,v}^{\text{in}}(\lambda) = \lambda, \quad Z_{T,v}^{\text{out}}(\lambda) = 1 + \lambda,$$

and therefore T with terminal v implements the activity $\frac{Z_{T,v}^{\text{in}}(\lambda)}{Z_{T,v}^{\text{out}}(\lambda)} = \lambda a_1$, as wanted.

Suppose that the statement is true for all values $\leq k$ and suppose that

$$a_{k+1} = \frac{1}{1 + \lambda a_{i_{k,1}} \cdots a_{i_{k,d}}}$$

for some $i_{k,1}, \dots, i_{k,d} \in \{0, \dots, k-1\}$. Let $J = \{j \in [d] \mid i_{k,j} \neq 0\}$ and note that for every $j \in [d] \setminus J$, it holds that $a_{i_{k,j}} = 1$. By the induction hypothesis, for every $j \in J$, there exists a tree T_j of maximum degree at most Δ , with terminal v_j , such that $\frac{Z_{T_j,v_j}^{\text{in}}(\lambda)}{Z_{T_j,v_j}^{\text{out}}(\lambda)} = \lambda a_{i_{k,j}}$. Let T be the tree obtained by taking the disjoint union of the trees T_j with $j \in J$ and identifying all the terminals v_j into a new vertex v . Then,

$$(7.2) \quad Z_{T,v}^{\text{in}}(\lambda) = \lambda \prod_{j \in J} \frac{Z_{T_j,v_j}^{\text{in}}(\lambda)}{\lambda}, \quad Z_{T,v}^{\text{out}}(\lambda) = \prod_{j \in J} Z_{T_j,v_j}^{\text{out}}(\lambda).$$

Now consider the tree T' obtained from T by adding a new vertex u whose single neighbor is the vertex v . Then,

$$Z_{T',u}^{\text{in}}(\lambda) = \lambda Z_{T,v}^{\text{out}}(\lambda), \quad Z_{T',u}^{\text{out}}(\lambda) = Z_T(\lambda).$$

Using this and (7.2), we therefore obtain that

$$\begin{aligned} \frac{Z_{T',u}^{\text{in}}(\lambda)}{Z_{T',u}^{\text{out}}(\lambda)} &= \lambda \frac{Z_{T,v}^{\text{out}}(\lambda)}{Z_T(\lambda)} = \lambda \frac{1}{1 + \frac{Z_{T,v}^{\text{in}}(\lambda)}{Z_{T,v}^{\text{out}}(\lambda)}} = \frac{\lambda}{1 + \lambda \prod_{j \in [J]} a_{i_{k,j}}} \\ &= \frac{\lambda}{1 + \lambda \prod_{j \in [d]} a_{i_{k,j}}} = \lambda a_{k+1}, \end{aligned}$$

where in the second to last equality we used that $a_{i_{k,j}} = 1$ for $j \in [d] \setminus J$. \square

7.1. Getting close to a repelling fixpoint. In this section, we will show how to generate points that are arbitrarily close to a fixpoint of the function $f(x) = \frac{1}{1 + \lambda x^{\Delta-1}}$ using a hard-core program. We start with the following lemma.

LEMMA 7.3. *Let $\Delta \geq 3$ and set $d := \Delta - 1$. Let p_k be a polynomial in λ defined by*

$$p_0 = p_1 = \dots = p_d = 1, \quad \text{and} \quad p_{k+1} = p_k + \lambda p_{k-d} \quad \text{for } k \geq d.$$

Then, for all $k \geq 0$, all roots of p_k are real.

Proof. Consider a graph $G_k = (V, E)$ with $V = \{d+1, \dots, k\}$ with i, j connected if $|i - j| \leq d$. We will show, by induction, that the independent set polynomial of G_k is p_k . The claim is true for $k = 0, \dots, d$ (since the graph is empty and $p_k = 1$). For $k+1 \geq d+1$ the claim follows by induction:

- the contribution of the independent sets with $k+1$ included is λ times the independence polynomial of G_{k-d} , which, by the inductive hypothesis, is λp_{k-d} ;
- the contribution of the independent sets with $k+1$ not included is the independent set polynomial of G_k , which, by the inductive hypothesis, is p_k .

Hence, the independent set polynomial of G_{k+1} is $p_k + \lambda p_{k-d} = p_{k+1}$.

A claw is a graph with 4 vertices a, b_1, b_2, b_3 and 3 edges ab_1, ab_2, ab_3 . A claw-free graph is a graph that does not contain a claw as an induced subgraph. We will show that G_k is claw-free. Suppose $a, b_1, b_2, b_3 \in \{d+1, \dots, k\}$ form a claw; w.l.o.g. $b_1 < b_2 < b_3$. Then $|a - b_i| \leq d$ for $i = 1, 2, 3$. If $b_2 < a$ then $b_2 - b_1 \leq a - b_1 \leq d$ and hence $b_1 b_2$ is an edge—a contradiction with the assumption that a, b_1, b_2, b_3 form a claw. If $b_2 > a$ then $b_3 - b_2 \leq b_3 - a \leq d$ and hence $b_2 b_3$ is an edge—a contradiction with the assumption that a, b_1, b_2, b_3 form a claw. Thus G_k is claw-free.

Now we use [5, Theorem 1.1] which states that the roots of the independent set polynomial of a claw-free graph are all real. \square

We will now show that we can get close to the fixpoint of f with the smallest norm.

LEMMA 2.7. *Let $\Delta \geq 3$ and $\lambda \in \mathbb{C} \setminus \mathbb{R}$, and set $d := \Delta - 1$. Let ω be the fixpoint of $f(x) = \frac{1}{1+\lambda x^d}$ with the smallest norm. For $k \geq 0$, let x_k be the sequence defined by*

$$(7.3) \quad x_0 = x_1 = \dots = x_{d-1} = 1, \quad x_k = \frac{1}{1 + \lambda \prod_{i=1}^d x_{k-i}} \quad \text{for } k \geq d.$$

Then, the sequence x_k is well-defined (i.e., the denominator of (7.3) is nonzero for all $k \geq d$) and converges to the fixpoint ω as $k \rightarrow \infty$. Moreover, there exist infinitely many k such that $x_k \neq \omega$.

Proof. For $k \geq 0$, let R_k be the sequence defined by

$$(7.4) \quad R_0 = R_1 = \dots = R_d = 1, \quad R_{k+1} = R_k + \lambda R_{k-d} \quad \text{for } k \geq d.$$

Observe that $R_k = p_k(\lambda)$, where p_k is the polynomial in Lemma 7.3. Since $\lambda \in \mathbb{C} \setminus \mathbb{R}$, Lemma 7.3 implies that $R_k \neq 0$ for all k and hence for $k \geq 0$ we can let

$$(7.5) \quad y_k = R_k / R_{k+1}.$$

Note that for $k \in \{0, \dots, d-1\}$ we have $y_k = x_k$. For $k \geq d$ we have

$$y_k = \frac{R_k}{R_{k+1}} = \frac{R_k}{R_k + \lambda R_{k-d}} = \frac{1}{1 + \lambda \frac{R_{k-d}}{R_k}} = \frac{1}{1 + \lambda \prod_{j=1}^d \frac{R_{k-j}}{R_{k-j+1}}} = \frac{1}{1 + \lambda \prod_{j=1}^d y_{k-j}}$$

and hence, by induction, $x_k = y_k$ for all k . It follows that the sequence x_k is well-defined.

Let $\omega_1, \dots, \omega_{d+1}$ be the fixpoints of $f(x) = \frac{1}{1+\lambda x^d}$ sorted in increasing order of their absolute value, so that $\omega_1 = \omega$. Note, since $\lambda \in \mathbb{C} \setminus \mathbb{R}$, by Lemma 4.2 we have that $|\omega_i| \neq |\omega_j|$ for different $i, j \in [d+1]$. Observe that the characteristic polynomial of the recurrence (7.4) is $z^{d+1} - z^d - \lambda$ and that the roots of the polynomial are

$1/\omega_1, \dots, 1/\omega_{d+1}$ (to verify this, use that $\lambda\omega_j^{d+1} + \omega_j - 1 = 0$ for $j \in [d+1]$). Therefore, from the theory of linear recurrences we have that there exist $\alpha_1, \dots, \alpha_{d+1} \in \mathbb{C}$ such that for all $k \geq 0$

$$R_k = \sum_{j=1}^{d+1} \alpha_j (1/\omega_j)^k.$$

Note that since $R_0 = R_1 = \dots = R_d = 1$ we have that $\alpha_1, \dots, \alpha_{d+1}$ is the solution of the following (Vandermonde) system

$$\sum_{j=1}^{d+1} (1/\omega_j)^k \alpha_j = 1 \quad \text{for } k \in \{0, \dots, d\}.$$

Suppose one of the $\alpha_1, \dots, \alpha_{d+1}$ is zero, say $\alpha_i = 0$ for some $i \in [d+1]$. Let $\alpha_{d+2} = -1$ and $\omega_{d+2} = 1$. For $j \in [d+1]$, note that $\omega_j \neq 1$ (since that would imply $\lambda = 0$) and therefore $\omega_j \neq \omega_{d+2}$. Then we have that $\{\alpha_j\}_{j \in [d+2] \setminus \{i\}}$ is a nonzero solution of the following (again Vandermonde) system

$$(7.6) \quad \sum_{j \in [d+2] \setminus \{i\}} (1/\omega_j)^k \alpha_j = 0 \quad \text{for } k \in \{0, \dots, d\}.$$

This is a contradiction, since the system only has a zero solution ($\omega_1, \dots, \omega_{d+1}, \omega_{d+2}$ are distinct). Thus none of the $\alpha_1, \dots, \alpha_{d+1}$ is zero and, in particular, $\alpha_1 \neq 0$. It follows that $x_k = R_k/R_{k+1}$ converges to $\omega_1 = \omega$ as $k \rightarrow \infty$.

To finish the proof, it remains to show that $x_k \neq \omega$ for infinitely many k . For the sake of contradiction, assume otherwise, and let k_0 be the largest integer such that $x_{k_0} \neq \omega$. From (7.3), we obtain

$$x_{k_0+d} = \frac{1}{1 + \lambda \prod_{j=0}^{d-1} x_{k_0+j}}.$$

By the choice of k_0 , we have $x_{k_0+1} = \dots = x_{k_0+d} = \omega$, which gives that $x_{k_0} = \omega$ (using that $\omega = \frac{1}{1+\lambda\omega^d}$), a contradiction.

This concludes the proof of Lemma 2.7. \square

Finally, we conclude this section with the following lemma, which will be useful later.

LEMMA 7.4. *For $\lambda \in \mathbb{C} \setminus \mathbb{R}$ and $d \geq 2$, let ω be the fixpoint of $f(x) = \frac{1}{1+\lambda x^d}$ with the smallest norm. Then, $\omega \in \mathbb{C} \setminus \mathbb{R}$ and $0 < |\omega - 1| < 1$.*

Proof. Since $1 - \omega = \lambda\omega^{d+1}$, we have that $|\omega - 1| > 0$ (otherwise, $\lambda = 0$) and $\omega \in \mathbb{C} \setminus \mathbb{R}$ (otherwise, $\lambda \in \mathbb{R}$). We therefore focus on showing that $|\omega - 1| < 1$.

Let $\omega_1, \dots, \omega_{d+1}$ be the fixpoints of f sorted in increasing order of their norm, so that $\omega_1 = \omega$. By Lemma 4.2, we have $|\omega_1| < \dots < |\omega_{d+1}|$, so by $1 - \omega_j = \lambda\omega_j^{d+1}$ ($j \in [d]$), we obtain that

$$(7.7) \quad |1 - \omega_1| < |1 - \omega_2| < \dots < |1 - \omega_{d+1}|.$$

Note that $\omega_1 - 1, \dots, \omega_{d+1} - 1$ are roots of $\lambda(y+1)^{d+1} + y = 0$; the coefficient of y^{d+1} and the coefficient of y^0 are both equal to λ , so by Vieta's formula,

$$(7.8) \quad \prod_{j=1}^{d+1} (1 - \omega_j) = 1.$$

Equations (7.7) and (7.8) imply $|1 - \omega_1| < 1$, as needed. \square

7.2. Bootstrapping a point next to a fixpoint to an ϵ -covering. Once we have the ability to obtain points close to the fixpoint ω , we proceed to the next step, which is creating a moderately dense set of points around ω (cf. Lemma 2.10).

The main idea of the proof of Lemma 2.10 is that close to the fixpoint the recurrence implements with a small error any polynomial with nonnegative integer coefficients (evaluated at $\omega - 1$). Then we use the fact that the values of these polynomials yield a dense set of points in \mathbb{C} . Before proceeding with the proof of Lemma 2.10 we state these ingredients formally.

LEMMA 7.5. *Let $z \in \mathbb{C} \setminus \mathbb{R}$ be such that $|z| < 1$. Let S be the set of values of polynomials with nonnegative integer coefficients evaluated at z (that is, $S = \{p(z) \mid p \in \mathbb{Z}_{\geq 0}[x]\}$). Then S is dense in \mathbb{C} .*

Proof. We can write $z = |z|e^{2\pi i x}$ for some $x \in [0, 1)$. Note that $x \neq 0$ and $x \neq 1/2$ (since we assumed $z \notin \mathbb{R}$).

First suppose that x is rational, that is, $x = p/q$ for integer coprime p, q , where $q \geq 3$. For any $k \in \{0, \dots, q-1\}$ we can obtain an arbitrarily small number on the ray with angle $2\pi k/q$ (by taking $(z^{p^{-1} \bmod q})^{k+q\ell}$ for large ℓ) and hence we have a dense set of points on the ray (taking integer multiples of the small number). Now we show how using the points on the rays we obtain a point arbitrarily close to any complex number $t \in \mathbb{C}$. First, we can write t as a convex combination of points on the rays, that is, $t = \alpha_0 r_0 + \dots + \alpha_{q-1} r_{q-1}$, where r_k is on the ray with angle $2\pi k/q$ (for $k \in \{0, \dots, q-1\}$), $\alpha_k \in [0, 1]$ (for $k \in \{0, \dots, q-1\}$), and $\sum_{k=0}^{q-1} \alpha_k = 1$. For $\epsilon > 0$, let $\hat{\alpha}_k$ be a rational such that $|\hat{\alpha}_k - \alpha_k| \leq \epsilon/q$ and let w be the product of the denominators of $\hat{\alpha}_0, \dots, \hat{\alpha}_{q-1}$. Since we have a dense set of points on each ray we can obtain \hat{r}_k on the ray with angle $2\pi k/q$ such that $|r_k/w - \hat{r}_k| \leq \epsilon/w$ (for $k \in \{0, \dots, q-1\}$). Now we argue that $\sum_{k=0}^{q-1} (w\hat{\alpha}_k)\hat{r}_k$ is close to t . We have

$$(7.9) \quad \left| t - \sum_{k=0}^{q-1} (w\hat{\alpha}_k)\hat{r}_k \right| = \left| \sum_{k=0}^{q-1} (\alpha_k - \hat{\alpha}_k)r_k + \sum_{k=0}^{q-1} \hat{\alpha}_k(r_k - w\hat{r}_k) \right| \leq \epsilon \sum_{k=0}^{q-1} |r_k| + (1 + \epsilon)\epsilon.$$

Taking ϵ sufficiently small we get a point arbitrarily close to t .

Now suppose x is irrational. Let $t = |t|e^{2\pi i y}$ be a complex number, where $y \in [0, 1)$. The fractional part $\{kx\}$ of kx for positive integers k is dense in $[0, 1)$ and, hence, for $\epsilon > 0$, there exists k such that $|\{kx\} - y| \leq \epsilon$ and $|z|^k \leq \epsilon$. Let $w = \lfloor |t|/|z|^k \rfloor$ so that $0 \leq |t| - w|z|^k \leq \epsilon$. Observe that $wz^k = w|z|^k e^{2\pi i \{kx\}}$, so by the triangle inequality

$$\begin{aligned} |t - wz^k| &\leq |t - |t|e^{2\pi i kx}| + ||t|e^{2\pi i kx} - wz^k| = |t| |e^{2\pi i(y - \{kx\})} - 1| + ||t| - w|z|^k| \\ &\leq \epsilon(1 + 2\pi|t|), \end{aligned}$$

where in the last inequality we used that for $\theta = 2\pi(y - \{kx\})$ it holds that

$$|e^{i\theta} - 1| = \sqrt{2 - 2\cos\theta} = 2|\sin(\theta/2)| = 2\sin|\theta/2| \leq |\theta| \leq 2\pi\epsilon.$$

Taking ϵ sufficiently small we get a point (wz^k) arbitrarily close to t . \square

Remark 7.6. Note that the assumption $|z| < 1$ is necessary—the lemma would be false for, e.g., $z = i$.

The following operation (as we will prove in Lemma 7.9 below) is a first-order approximation of operation (7.1) when applied to points around ω perturbed by

a_1, \dots, a_d :

$$(7.10) \quad (a_1, \dots, a_d) \mapsto z \sum_{i=1}^d a_i.$$

We will use a sequence of (7.10) to implement polynomials with nonnegative integer coefficients; the complexity of a polynomial will be the number of steps in the sequence.

DEFINITION 7.7. *A straight-line program with operation (7.10) is a sequence of assignments starting with $a_0 = 0$, $a_1 = 1$, and*

$$a_k = z(a_{i_{k,1}} + \dots + a_{i_{k,d}}) \quad \text{for } k = 2, 3, \dots,$$

where $i_{k,1}, \dots, i_{k,d} \in \{0, \dots, k-1\}$. We say that the straight-line program generates $x \in \mathbb{C}$ if there exists integer $k \geq 0$ such that $a_k = x$.

Using a finite sequence of (7.10), we can implement any polynomial with nonnegative integer coefficients, up to factors of z . More precisely, we have the following.

LEMMA 7.8. *Let $p \in \mathbb{Z}_{\geq 0}[z]$ be a polynomial with nonnegative integer coefficients. There exist nonnegative integers $k := k(p)$ and $n := n(p)$ and a straight-line program with operation (7.10) such that $a_k = z^n p(z)$.*

Proof. Let $p(z) = \sum_{j=0}^t c_j z^j$, where $c_t \neq 0$ or $t = 0$. We will prove the claim by induction on $t + \sum_{j=0}^t c_j$. The base case is $p(z) \equiv 0$; here we can take $k(p) = 0$ and $n(p) = 0$. Now assume $t + \sum_{j=0}^t c_j \geq 1$.

First assume that $c_0 \geq 1$. Let $q(z) = p(z) - 1$. By the induction hypothesis there exist n, k , and a straight-line program with operation (7.10) such that $a_k = z^n q(z)$. Let $a_{k+1} = za_1$ and $a_{k+j} = za_{k+j-1}$ for $j = 2, \dots, n$ (note that $a_{k+n} = z^n$). Finally, add $a_{k+n+1} = z(a_k + a_{k+n})$. Note that $a_{k+n+1} = z^{n+1}p(z)$.

Now assume $c_0 = 0$. Let $q(z) = p(z)/z$. By the induction hypothesis there exist n, k , and a straight-line program with operation (7.10) such that $a_k = z^n q(z)$. Let $a_{k+1} = za_k$. Note that $a_{k+1} = z^{n+1}p(z)$. \square

From the Taylor expansion we have that close to the fixpoint the multivariate hard-core recurrence implements operation (7.10) (with a small error).

LEMMA 7.9. *Suppose that $\lambda \in \mathbb{C} \setminus \mathbb{R}$ and $d \geq 2$. Let ω be the fixpoint of $f(x) = \frac{1}{1+\lambda x^d}$ with the smallest norm and set $z = \omega - 1$.*

There exist reals $C_0 := C_0(\omega, \lambda, d) > 1$ and $\delta_0 := \delta_0(\omega, \lambda, d) > 0$ such that for any $a_1, \dots, a_d \in \mathbb{C}$ with $|a_j| \leq \delta_0$ (for $j \in [d]$) we have

$$(7.11) \quad \frac{1}{1 + \lambda \prod_{j=1}^d (\omega + a_j)} = \omega + z \left(\sum_{j=1}^d a_j \right) + \tau,$$

where $|\tau| \leq C_0 \max_{j \in [d]} |a_j|^2$.

Proof. Let b_1, \dots, b_d be arbitrary complex numbers with $|b_1|, \dots, |b_d| \leq 1$. Let

$$F(t) = \frac{1}{G(t)}, \quad \text{where } G(t) = 1 + \lambda \prod_{j=1}^d (\omega + b_j t).$$

Then, using that ω is a fixpoint of $f(x) = \frac{1}{1+\lambda x^d}$, we have $G(0) = 1/\omega$ and

$$(7.12) \quad F'(0) = -\frac{\lambda \left(\sum_{j=1}^d b_j \right) (\omega)^{d-1}}{(1 + \lambda(\omega)^d)^2} = z \sum_{j=1}^d b_j.$$

Note that for all $t \in [0, 1]$

$$(7.13) \quad |G'(t)| = \left| \lambda \sum_{j=1}^d b_j \prod_{k \neq j} (\omega + b_k t) \right| \leq d|\lambda|(1 + |\omega|)^d.$$

Similarly, we have that for all $t \in [0, 1]$

$$|G''(t)| \leq d^2 |\lambda| (1 + |\omega|)^d.$$

Let $\delta_0 := \min \left\{ \frac{1}{2|\omega|d|\lambda|(1+|\omega|)^d}, 1 \right\}$. Note that (7.13) implies that for $t \in (0, \delta_0)$ we have

$$|G(t)| \geq |G(0)| - td|\lambda|(1 + |\omega|)^d \geq \frac{1}{|\omega|} - \delta_0 d|\lambda|(1 + |\omega|)^d \geq \frac{1}{2|\omega|},$$

and hence for $t \in (0, \delta_0)$

$$(7.14) \quad |F''(t)| = \left| \frac{2G'(t)^2}{G(t)^3} - \frac{G''(t)}{G(t)^2} \right| \leq 3(2|\omega|)^3 d^4 |\lambda|^2 (1 + |\omega|)^{2d} =: C,$$

which implies

$$(7.15) \quad |F(t) - F(0) - tF'(0)| \leq Ct^2.$$

Given a_1, \dots, a_d such that $|a_j| \leq \delta_0$, let $t = \max_j |a_j| \in (0, \delta_0)$ and $b_j = a_j/t$ (for $j \in [d]$); note,

$$\frac{1}{1 + \lambda \prod_{j=1}^d (\omega + a_j)} = F(t) \quad \text{and} \quad tF'(0) = z \sum_{j=1}^d a_j.$$

Let $C_0 := \max\{C, 2\} > 1$. The lemma now follows from (7.15), (7.12), and the fact that $F(0) = \omega$. \square

Finally we can prove Lemma 2.10, which we restate here for convenience.

LEMMA 2.10. *Let $\Delta \geq 3$ and $\lambda \in \mathbb{C}_{\mathbb{Q}} \setminus \mathbb{R}$, and set $d := \Delta - 1$. Let ω be the fixpoint of $f(x) = \frac{1}{1+\lambda x^d}$ with the smallest norm. For any $\epsilon, \kappa > 0$ there exists a radius $\rho \in (0, \kappa)$ such that the following holds. For every $\lambda' \in B(\lambda\omega, \rho)$, there exists a tree G of maximum degree at most Δ that implements λ' with accuracy $\rho\epsilon$.*

Proof. Consider arbitrary $\epsilon, \kappa > 0$ and let $z := \omega - 1$. By Lemma 7.4, we have that $z \in \mathbb{C} \setminus \mathbb{R}$ and $0 < |z| < 1$. Therefore, Lemma 7.5 gives that polynomials with nonnegative integer coefficients evaluated at z are dense in \mathbb{C} . Hence there exists a finite collection \mathcal{F} of polynomials with nonnegative integer coefficients whose values at z form an $\epsilon/2$ -covering of the unit disk (to obtain the collection take a finite $\epsilon/4$ -covering of the unit disk and for every point in the covering, using the density, get a polynomial whose value at z is at distance at most $\epsilon/4$ from the point).

By Lemma 7.8 every polynomial p with nonnegative integer coefficients can be generated, up to a factor $z^{n(p)}$, using $k(p)$ operations (7.10). Let

$$N := \max_{p \in \mathcal{F}} n(p) \quad \text{and} \quad K := N + \max_{p \in \mathcal{F}} k(p).$$

For every $p \in \mathcal{F}$, there is a straight-line program to generate $z^N p(z)$ using at most K applications of (7.10) (the extra N in the definition of K is to allow for extra operations (7.10) to move from $z^{n(p)} p(z)$ to $z^N p(z)$). Let $C_0 > 1$ and $\delta_0 > 0$ be the constants from Lemma 7.9. Let $Z(C_0, d, k) := C_0(2d)^{2k}$ for $k \geq 0$ and

$$(7.16) \quad \delta = \min \left\{ \delta_0/d^K, \frac{\epsilon|z|^N}{2Z(C_0, d, K)}, \frac{1}{Z(C_0, d, K)}, \frac{\kappa}{2|\lambda|} \right\}.$$

By Lemma 2.7, there is a hard-core program x_0, x_1, \dots such that for sufficiently large m, m' , it holds that $0 < |x_m - \omega| \leq \delta$ and $|x_{m'} - \omega| \leq |x_m - \omega|^2$. Let $y_1 = x_m$ and $y_0 = x_{m'}$, so that $0 < |y_1 - \omega| \leq \delta$ and $|y_0 - \omega| \leq |y_1 - \omega|^2$. Finally, let

$$h := y_1 - \omega \quad \text{and set} \quad r := |hz^N|.$$

Since $|z| < 1$, we have $r \leq |h| \leq \delta < \kappa/|\lambda|$.

We claim that for any u such that $(\omega + u) \in B(\omega, r)$, there exists a hard-core program that generates $\omega + u'$ with $|u - u'| \leq \epsilon r$. By Lemma 7.2, this implies that, for $\rho := |\lambda|r \in (0, \kappa)$, any activity in the ball $B(\lambda\omega, \rho)$ can be implemented with accuracy $\epsilon\rho$ by a tree of maximum degree Δ , as needed.

To obtain the desired hard-core program, observe first that u/r belongs to the unit disc, so there exists $p \in \mathcal{F}$ such that

$$(7.17) \quad |p(z) - u/r| \leq \epsilon/2.$$

Moreover, there is a straight-line program $\hat{y}_0, \hat{y}_1, \dots, \hat{y}_k$ with $k \leq K$ to generate $z^N p(z)$, i.e., $\hat{y}_0 = 0$, $\hat{y}_1 = 1$, and

$$(7.18) \quad \hat{y}_\ell = z(\hat{y}_{i_{\ell,1}} + \dots + \hat{y}_{i_{\ell,d}}) \quad \text{for } \ell = 2, 3, \dots, k,$$

where $i_{\ell,1}, \dots, i_{\ell,d} \in \{0, \dots, \ell-1\}$ and $\hat{y}_k = z^N p(z)$. Note that for all $\ell \in \{0, \dots, k\}$ we have by induction (using $|z| < 1$ from Lemma 7.4) that

$$(7.19) \quad |\hat{y}_\ell| \leq d^{\ell-1}.$$

We will next convert the straight-line program (7.18) into a hard-core program. For y_0, y_1 as above, let y_2, \dots, y_k be given by

$$(7.20) \quad y_\ell = \frac{1}{1 + \lambda \prod_{j=1}^d y_{i_{\ell,j}}} \quad \text{for } \ell = 2, 3, \dots, k.$$

We will prove that for all $\ell = 0, 1, \dots, k$ it holds that

$$(7.21) \quad y_\ell = \omega + h\hat{y}_\ell + \tau_\ell, \quad \text{where } |\tau_\ell| \leq |h|^2 Z(C_0, d, \ell).$$

Assuming (7.21) for the moment, let us conclude the proof of the claim by showing that y_k is at distance $\leq r\epsilon$ from $\omega + u$ and that it can be generated by a hard-core program.

In particular, for $\ell = k$, (7.21) gives (using that $\hat{y}_k = z^N p(z)$, $|h| = |y_1 - \omega| \leq \delta$, and (7.16))

$$y_k = \omega + h z^N p(z) + \tau_k, \text{ where } |\tau_k| \leq |h|^2 Z(C_0, d, K) \leq |h| |z|^N \epsilon / 2.$$

Combining this with (7.17) and recalling that $r = |h z^N|$, we obtain that

$$|y_k - (\omega + u)| = |h z^N p(z) + \tau_k - u| \leq r |p(z) - u/r| + |\tau_k| \leq r \epsilon.$$

To finish the proof of the claim, we only need to observe that, since y_0, y_1 can be generated using a hard-core program, we can also generate the sequence y_0, y_1, \dots, y_k using a hard-core program.

It remains to prove (7.21). We do this by induction. Note that $y_0 = \omega + 0 \cdot h + \tau_0$, where $|\tau_0| = |y_0 - \omega| \leq |h|^2$ and $y_1 = \omega + 1 \cdot h + 0$, covering the base cases of (7.21). For the induction step, assume that for all $0 \leq \ell' < \ell$ it holds that

$$(7.22) \quad y_{\ell'} = \omega + h \hat{y}_{\ell'} + \tau_{\ell'}, \text{ where } |\tau_{\ell'}| \leq |h|^2 Z(C_0, d, \ell').$$

For all $\ell \leq K$, we have by (7.16) that $Z(C_0, d, \ell) \leq Z(C_0, d, K) \leq 1/\delta \leq 1/h$ and $|h| d^{\ell-1} \leq \delta_0/d$. Therefore (7.19) and (7.22) give that, for all $0 \leq \ell' < \ell$,

$$(7.23) \quad |h \hat{y}_{\ell'} + \tau_{\ell'}| \leq |h| d^{\ell'-1} + |h|^2 Z(C_0, d, \ell') \leq |h| d^{\ell-1} + |h| \leq \delta_0.$$

Hence, we can apply Lemma 7.9 with a_j 's of the form $h \hat{y}_{\ell'} + \tau_{\ell'}$ ($\ell' \in \{0, 1, \dots, \ell-1\}$). From Lemma 7.9 and (7.23) we have

$$y_\ell = \frac{1}{1 + \lambda \prod_{j=1}^d (\omega + h \hat{y}_{i_{\ell,j}} + \tau_{i_{\ell,j}})} = \omega + h z \left(\sum_{j=1}^d \hat{y}_{i_{\ell,j}} \right) + \tau_\ell, \text{ where } \tau_\ell = z \sum_{j=1}^d \tau_{i_{\ell,j}} + \tau$$

and

$$|\tau| \leq C_0 \max_{\ell'=0, \dots, \ell-1} |h \hat{y}_{\ell'} + \tau_{\ell'}|^2 \leq C_0 |h|^2 (d^{\ell-1} + 1)^2 \leq C_0 |h|^2 d^{2\ell}.$$

Thus

$$\begin{aligned} |\tau_\ell| &\leq d |h|^2 Z(C_0, d, \ell-1) + C_0 |h|^2 d^{2\ell} = |h|^2 C_0 (d(2d)^{2\ell-2} + d^{2\ell}) \\ &\leq |h|^2 C_0 (2d)^{2\ell} = |h|^2 Z(C_0, d, \ell), \end{aligned}$$

completing the induction step. This finishes the proof of (7.21) and therefore the proof of Lemma 2.10. \square

7.3. Bootstrapping ϵ -covering to arbitrary density. Our next step is to use the “moderately dense” set of points in a small disk around ω to create a dense set of points. We first need a few technical results.

LEMMA 7.10. *Suppose that $\lambda \in \mathbb{C} \setminus \mathbb{R}$ and $d \geq 2$. Let ω be the fixpoint of $f(x) = \frac{1}{1 + \lambda x^d}$ with the smallest norm and let $z = \omega - 1$.*

There exist reals $C_1 := C_1(\omega, \lambda, d) > 0$ and $\delta_1 := \delta_1(\omega, \lambda, d) > 0$ such that for any $a_1, \dots, a_d \in \mathbb{C}$ with $|a_j| \leq \delta_1$ (for $j \in [d]$) we have

$$(7.24) \quad \frac{1}{\lambda} \left(\frac{1}{\omega + a_d} - 1 \right) \prod_{j=1}^{d-1} (\omega + a_k)^{-1} = \omega + \left(\frac{a_d}{z} - \sum_{j=1}^{d-1} a_j \right) + \tau,$$

where $|\tau| \leq C_1 \max_{k \in [d]} |a_k|^2$.

Proof. Let b_1, \dots, b_d be arbitrary complex numbers with $|b_1|, \dots, |b_d| \leq 1$. Let

$$F(t) = \frac{1}{\lambda} \left(\frac{1}{G_d(t)} - \frac{1}{G_{d-1}(t)} \right), \quad \text{where } G_k(t) := \prod_{j=1}^k (\omega + b_j t) \text{ for } k \in [d].$$

Then, $G_k(0) = \omega^k$ and $G'_k(0) = \omega^{k-1} \sum_{j=1}^k b_j$, so

$$(7.25) \quad F'(0) = -\frac{G'_d(0)}{\lambda(G_d(0))^2} + \frac{G'_{d-1}(0)}{\lambda(G_{d-1}(0))^2} = -\frac{\sum_{j=1}^d b_j}{\lambda\omega^{d+1}} + \frac{\sum_{j=1}^{d-1} b_j}{\lambda\omega^d} = \frac{b_d}{z} - \sum_{j=1}^{d-1} b_j,$$

where in the last equality we used that $\lambda\omega^{d+1} = 1 - \omega = -z$. As in (7.13), for $k \in [d]$ and all $t \in [0, 1]$, we have that

$$(7.26) \quad |G'_k(t)| \leq d(1 + |\omega|)^d.$$

Let $\delta_1 := \min \left\{ \frac{|\omega|^d}{2d(1+|\omega|)^d}, 1 \right\} > 0$. Note that (7.26) implies that for $k \in [d]$ and $t \in (0, \delta_1)$ we have

$$|G_k(t)| \geq |G_k(0)| - td(1 + |\omega|)^d \geq \frac{|\omega|^d}{2}.$$

Similarly to (7.26), for $k \in [d]$ and all $t \in [0, 1]$, we have

$$(7.27) \quad |G''_k(t)| \leq d^2(1 + |\omega|)^d,$$

and hence for $t \in (0, \delta_1)$, following the same argument as in (7.14),

$$|F''(t)| \leq \frac{6}{|\lambda|} (2/|\omega|^d)^3 d^4 (1 + |\omega|)^{2d} =: C_1,$$

which implies

$$(7.28) \quad |F(t) - F(0) - tF'(0)| \leq C_1 t^2.$$

Given a_1, \dots, a_d such that $|a_j| \leq \delta_1$, let $t = \max_j |a_j|$ and $b_j = a_j/t$ (for $j \in [d]$); then, from (7.25)

$$\frac{1}{\lambda} \left(\frac{1}{\omega + a_d} - 1 \right) \prod_{j=1}^{d-1} (\omega + a_j)^{-1} = F(t) \quad \text{and} \quad tF'(0) = \frac{a_d}{z} - \sum_{j=1}^{d-1} a_j.$$

The lemma now follows from (7.28) and the fact that $F(0) = \omega$. \square

LEMMA 7.11. *Suppose that $\lambda \in \mathbb{C} \setminus \mathbb{R}$ and $d \geq 2$. Let ω be the fixpoint of $f(x) = \frac{1}{1+\lambda x^d}$ with the smallest norm and let $z = \omega - 1$.*

There exist reals $C_2 := C_2(\omega, \lambda, d) > 0$ and $\delta_2 := \delta_2(\omega, \lambda, d) > 0$ such that for any $a_1, \dots, a_d \in \mathbb{C}$ with $|a_j| \leq \delta_2$ (for $j \in [d]$) we have

$$(7.29) \quad \frac{\partial}{\partial x} \frac{1}{1 + \lambda(\omega + x) \prod_{j=1}^{d-1} (\omega + a_j)} \Big|_{x=a_d} = z + \tau,$$

where $|\tau| \leq C_2 \max_{j \in [d]} |a_j|$.

Proof. Let C_0, δ_0 be the constants in Lemma 7.9 and let $\delta_2 := \min\{1, \delta_0\}$.

Suppose that $a_1, \dots, a_d \in \mathbb{C}$ with $|a_j| \leq \delta_2 \leq 1$. From Lemma 7.9 and the fact that $|z| < 1$ (cf. Lemma 7.4), we have

$$(7.30) \quad \left| \frac{1}{1 + \lambda \prod_{j=1}^d (\omega + a_j)} - \omega \right| \leq \left| z \sum_{j=1}^d a_j \right| + C_0 \max_{j \in [d]} |a_j|^2 \leq C \max_{j \in [d]} |a_j|, \text{ where } C := d + C_0.$$

From (7.13), we have

$$(7.31) \quad \left| \lambda \prod_{j=1}^{d-1} (\omega + a_j) - \lambda \omega^{d-1} \right| \leq C' \max_{j \in [d-1]} |a_j|, \text{ where } C' := d|\lambda|(1 + |\omega|)^d.$$

Note that $\frac{\partial}{\partial x} \frac{1}{1 + \lambda(\omega + x) \prod_{j=1}^{d-1} (\omega + a_j)} \Big|_{x=a_d}$ equals

$$- \frac{\lambda \prod_{j=1}^{d-1} (\omega + a_j)}{\left(1 + \lambda \prod_{j=1}^d (\omega + a_j)\right)^2} = -(\lambda \omega^{d-1} + \tau_1)(\omega + \tau_2)^2,$$

where $|\tau_1| \leq C' \max_{j \in [d]} |a_j|$ and $|\tau_2| \leq C \max_{j \in [d]} |a_j|$. Notice

$$-(\lambda \omega^{d-1} + \tau_1)(\omega + \tau_2)^2 = -\lambda \omega^{d+1} + \tau,$$

where $|\tau| \leq |\lambda| |\tau_2| |\omega|^{d-1} (2|\omega| + |\tau_2|) + |\tau_1| (|\omega| + |\tau_2|)^2$. Using that $\max_{j \in [d]} |a_j| \leq 1$, we obtain $|\tau| \leq C_2 \max_{j \in [d]} |a_j|$, where

$$C_2 := |\lambda| |\omega|^{d-1} C (2|\omega| + C) + C' (|\omega| + C) > 0.$$

This finishes the proof. \square

LEMMA 7.12. Let $\Delta \geq 3$ and $\lambda \in \mathbb{C}_{\mathbb{Q}} \setminus \mathbb{R}$, and set $d := \Delta - 1$. Let ω be the fixpoint of $f(x) = \frac{1}{1 + \lambda x^d}$ with the smallest norm. There is a set of activities $\{\lambda'_0, \lambda'_1, \dots, \lambda'_t\} \subseteq \mathbb{C}_{\mathbb{Q}} \setminus \{0\}$ and a real $r > 0$ such that the following hold for all $\hat{\omega} \in B(\omega, r)$:

1. for $i = 0$, $\frac{1}{1 + \lambda'_0} \in B(\hat{\omega}, 2r)$;
2. for $i = 1, \dots, t$, the map $\Phi_i : x \mapsto \frac{1}{1 + \lambda'_i x}$ is contracting on the ball $B(\hat{\omega}, 2r)$;
3. $B(\hat{\omega}, 2r) \subseteq \bigcup_{i=1}^t \Phi_i(B(\hat{\omega}, 2r))$.

Moreover, for $i = 0, 1, \dots, t$, there is a bipartite graph G'_i of degree Δ with a vertex w_i such that

$$\lambda'_i = \frac{Z_{G'_i, w_i}^{\text{in}}(\lambda)}{Z_{G'_i, w_i}^{\text{out}}(\lambda)} \quad \text{and} \quad \deg_{w_i}(G'_i) = \begin{cases} \Delta - 1 & \text{if } i = 0, \\ \Delta - 2 & \text{if } i = 1, \dots, t. \end{cases}$$

Proof. Take $C = \max\{C_0, C_1, C_2, 1\} > 0$ and $\delta = \min\{\delta_0, \delta_1, \delta_2, |\omega|, 1\} \in (0, 1]$, where $C_0, C_1, C_2, \delta_0, \delta_1, \delta_2$ are the constants from Lemmas 7.9, 7.10, and 7.11. Moreover, let $z = \omega - 1$ and recall by Lemma 7.4 that $0 < |z| < 1$. Let $\epsilon = \frac{1}{9}|z| > 0$. By Lemma 2.10, (Δ, λ) implements a set of activities $S = \{\lambda_1, \dots, \lambda_t\}$ which is an $(\epsilon\rho)$ -covering of $B(\lambda\omega, \rho)$ for some $\rho > 0$ satisfying

$$(7.32) \quad \rho < \frac{|\lambda|}{10C} \delta |z| (1 - |z|).$$

For convenience, for $i \in [t]$, define ζ_i by setting

$$\lambda_i = \lambda(\omega + \zeta_i) \text{ so that } \max_{i \in [t]} |\zeta_i| \leq \frac{\rho}{|\lambda|} < \frac{1}{5C} \delta |z|(1 - |z|) < \delta.$$

By Lemma 2.10, (Δ, λ) also implements an activity λ_0 such that $\lambda_0 = \lambda(\omega + \zeta_0)$ with $d|\zeta_0| < \epsilon\rho/|\lambda|$ (in particular, $|\zeta_0| < \delta$). Note that since $\lambda \in \mathbb{C}_{\mathbb{Q}}$ we have that $S \subseteq \mathbb{C}_{\mathbb{Q}}$.

Let $r := \rho \frac{|z|}{3|\lambda|}$. Moreover, let $\lambda'_0 := \lambda(\omega + \zeta_0)^d$, while for $i = 1, \dots, t$, let

$$\lambda'_i := \lambda(\omega + \zeta_i)(\omega + \zeta_0)^{d-2} \text{ and } \Phi_i \text{ be the map } x \mapsto \frac{1}{1 + \lambda'_i x}.$$

Note that $\{\lambda'_0, \dots, \lambda'_t\} \subseteq \mathbb{C}_{\mathbb{Q}} \setminus \{0\}$, since $\lambda \in \mathbb{C}_{\mathbb{Q}}$, $S \subseteq \mathbb{C}_{\mathbb{Q}}$, and for $i = 0, 1, \dots, t$ it holds that $|\zeta_i| < \delta \leq |\omega|$.

Consider an arbitrary $\hat{\omega} \in B(\omega, r)$. We first show that

$$(7.33) \quad \Phi_i \text{ is contracting on the ball } B(\hat{\omega}, 2r) \text{ for every } i \in [t].$$

Let $x \in B(\hat{\omega}, 2r)$. Since $|\hat{\omega} - \omega| \leq r$, we have $x \in B(\omega, 3r)$. Note that $3r < \delta$ and $\max_{i \in [t]} |\zeta_i| < \delta$, therefore from Lemma 7.11 (applied to $a_d = x - \omega$, $a_1 = \zeta_i$ and $a_2 = \dots = a_{d-1} = \zeta_0$), we have $\Phi'_i(x) = z + \tau$, where

$$|\tau| \leq C \max\{|\zeta_0|, |\zeta_i|, |x - \omega|\} \leq C \max\left\{\frac{\rho}{|\lambda|}, 3r\right\} = C \frac{\rho}{|\lambda|} \leq \frac{1 - |z|}{2},$$

and hence $|\Phi'_i(x)| \leq \frac{1 + |z|}{2}$, proving (7.33). We next show that

$$(7.34) \quad \text{for every } x \in B(\hat{\omega}, 2r), \text{ there exists } i \in [t] \text{ such that } \Phi_i^{-1}(x) \in B(\hat{\omega}, 2r).$$

We first calculate $\Phi_i^{-1}(x)$ for $i \in [t]$. By Lemma 7.10 (applied to $a_d = x - \omega$, $a_1 = \zeta_i$, and $a_2 = \dots = a_{d-1} = \zeta_0$), we have

$$\Phi_i^{-1}(x) = \frac{1}{\lambda} \left(\frac{1}{x} - 1 \right) \frac{1}{(\omega + \zeta_i)(\omega + \zeta_0)^{d-2}} = \omega + \left(\frac{x - \omega}{z} - \zeta_i - (d-2)\zeta_0 \right) + \tau,$$

where

$$|\tau| \leq C \max\{|\zeta_0|^2, |\zeta_i|^2, |x - \omega|^2\} \leq C \left(\frac{\rho}{|\lambda|} \right)^2 \leq \epsilon \frac{\rho}{|\lambda|} \quad \text{and} \quad (d-2)|\zeta_0| \leq d|\zeta_0| < \epsilon \frac{\rho}{|\lambda|}.$$

It follows that

$$(7.35) \quad \Phi_i^{-1}(x) - \omega = (\omega + \zeta) - (\omega + \zeta_i) + \tau', \text{ where } \zeta := \frac{x - \omega}{z} \text{ and } |\tau'| \leq 2\epsilon \frac{\rho}{|\lambda|}.$$

Note that

$$|\lambda\zeta| \leq |\lambda| \frac{|x - \omega|}{|z|} \leq |\lambda| \frac{3r}{|z|} = \rho,$$

so $\lambda(\omega + \zeta)$ belongs to the ball $B(\lambda\omega, \rho)$. In particular, we can choose λ_i from the $(\epsilon\rho)$ -covering such that

$$|\lambda(\omega + \zeta) - \lambda_i| \leq \epsilon\rho, \text{ which gives that } |(\omega + \zeta) - (\omega + \zeta_i)| \leq \epsilon \frac{\rho}{|\lambda|}.$$

Combining this with (7.35) and using that $\epsilon = \frac{1}{3}|z|$, we obtain that $|\Phi_i^{-1}(x) - \omega| \leq 3\epsilon \frac{\rho}{|\lambda|} = r$ and, therefore, by the triangle inequality $|\Phi_i^{-1}(x) - \hat{\omega}| \leq 2r$ (since $|\hat{\omega} - \omega| \leq r$). This proves (7.34). Finally, we show that

$$(7.36) \quad \frac{1}{1 + \lambda'_0} \in B(\hat{\omega}, 2r).$$

From Lemma 7.9 (applied to $a_1 = \dots = a_d = \zeta_0$), we obtain that

$$\frac{1}{1 + \lambda'_0} = \omega + d\zeta_0 + \tau, \text{ where } |\tau| \leq C|\zeta_0|^2 \leq \epsilon \frac{\rho}{|\lambda|} = r/3.$$

Since $d|\zeta_0| \leq \epsilon \frac{\rho}{|\lambda|} = r/3$, we obtain that $\frac{1}{1 + \lambda'_0} \in B(\omega, r)$ and therefore $\frac{1}{1 + \lambda'_0} \in B(\hat{\omega}, 2r)$ as well (since $|\hat{\omega} - \omega| \leq r$), finishing the proof of (7.36).

In light of (7.33), (7.34) and (7.36), to prove the lemma it remains to show the existence of the graphs G'_0, G'_1, \dots, G'_t claimed in the statement. Since (Δ, λ) implements the activities $\lambda_0, \lambda_1, \dots, \lambda_t$, for $i = 0, 1, \dots, t$ there exists a bipartite graph G_i of maximum degree at most Δ with terminal v_i such that $\lambda_i = \frac{Z_{G_i, v_i}^{\text{in}}(\lambda)}{Z_{G_i, v_i}^{\text{out}}(\lambda)}$. Consider first the case where we want to implement λ'_i for some $i \neq 0$. Construct the graph G'_i by taking $d - 2$ disjoint copies of the graph G_0 , one copy of the graph G_i , and identifying their terminals into a single vertex w_i . Then, the degree of w_i in G_i is $d - 1 = \Delta - 2$ and we have that

$$Z_{G'_i, w_i}^{\text{in}}(\lambda) = \lambda \left(\frac{Z_{G_0, v_0}^{\text{in}}(\lambda)}{\lambda} \right)^{d-2} \left(\frac{Z_{G_i, v_i}^{\text{in}}(\lambda)}{\lambda} \right), \quad Z_{G'_i, w_i}^{\text{out}}(\lambda) = (Z_{G_0, v_0}^{\text{out}}(\lambda))^{d-2} Z_{G_i, v_i}^{\text{out}}(\lambda)$$

and, therefore,

$$\frac{Z_{G'_i, w_i}^{\text{in}}(\lambda)}{Z_{G'_i, w_i}^{\text{out}}(\lambda)} = \lambda(\lambda_0/\lambda)^{d-2}(\lambda_i/\lambda) = \lambda(\omega + \zeta_i)(\omega + \zeta_0)^{d-2} = \lambda'_i,$$

as needed. The construction for the case $i = 0$ is analogous; to construct G'_0 , we take d disjoint copies of the graph G_0 and identify their terminals into a single vertex w_0 . Then, w_0 has degree $d = \Delta - 1$ and it holds that $\lambda_0 = Z_{G'_0, w_0}^{\text{in}}(\lambda)/Z_{G'_0, w_0}^{\text{out}}(\lambda)$. \square

Using Lemma 7.12, we can now prove Proposition 2.6 by applying Lemmas 2.8 and 5.2.

PROPOSITION 2.6. *Let $\Delta \geq 3$ and $\lambda \in \mathbb{C}_{\mathbb{Q}} \setminus \mathbb{R}$, and set $d := \Delta - 1$. Let ω be the fixpoint of $f(x) = \frac{1}{1 + \lambda x^d}$ with the smallest norm. There exists a rational $\rho > 0$ such that the following holds.*

There is a polynomial-time algorithm such that, on input $\lambda' \in B(\lambda, \rho) \cap \mathbb{C}_{\mathbb{Q}}$ and rational $\epsilon > 0$, outputs a bipartite graph G of maximum degree at most Δ with terminal v that implements λ' with accuracy ϵ . Moreover, the algorithm outputs the values $Z_{G, v}^{\text{in}}(\lambda), Z_{G, v}^{\text{out}}(\lambda)$.

Proof. Let $r > 0$, $\{\lambda'_0, \lambda'_1, \dots, \lambda'_t\} \subseteq \mathbb{C}_{\mathbb{Q}} \setminus \{0\}$, and let G'_0, \dots, G'_t (with vertices w_0, \dots, w_t , respectively) be as in Lemma 7.12. Let $\hat{\omega} \in \mathbb{C}_{\mathbb{Q}}$ be such that $|\omega - \hat{\omega}| < r$ and let $\rho > 0$ be a rational such that $|\omega - \hat{\omega}| < \rho/|\lambda| < r$. Note that the choice of $\hat{\omega}$ and ρ ensures that

$$(7.37) \quad B(\omega, \rho/|\lambda|) \subset B(\hat{\omega}, 2r)$$

since for all x with $|x - \omega| \leq \frac{\rho}{|\lambda|}$, we have by the triangle inequality $|x - \hat{\omega}| \leq |x - \omega| + |\omega - \hat{\omega}| < 2r$. Let

$$(7.38) \quad x_0 := \frac{1}{1 + \lambda'_0} \text{ so that, by Lemma 7.12, } x_0 \in B(\hat{\omega}, 2r).$$

Now, suppose that we are given inputs $\lambda' \in B(\lambda\omega, \rho) \cap \mathbb{C}_{\mathbb{Q}}$ and rational $\epsilon > 0$. We want to output in time $\text{poly}(\text{size}(\lambda', \epsilon))$ a bipartite graph of maximum degree Δ that implements λ' with accuracy ϵ . By Lemma 7.12, we have that the maps $\Phi_i(x) = \frac{1}{1 + \lambda'_i x}$ with $i \in [t]$ satisfy the hypotheses of Lemma 2.8 (with $z_0 = \hat{\omega}$ and radius $2r$). Moreover, by (7.37) and (7.38), we have that x_0 and $x^* = \frac{\lambda'}{\lambda}$ belong to the ball $B(\hat{\omega}, 2r)$. Let $\hat{\epsilon} = \epsilon/|\lambda|$ and $\epsilon' \in (0, \hat{\epsilon})$ be a rational such that $\text{size}(\epsilon') = \text{poly}(\text{size}(\hat{\epsilon})) = \text{poly}(\text{size}(\epsilon))$. Using the algorithm of Lemma 2.8 on input x_0, x^* , and ϵ' , we obtain in time $\text{poly}(\text{size}(x_0, x^*, \epsilon')) = \text{poly}(\text{size}(\lambda', \epsilon))$ a number $\hat{x} \in B(\hat{\omega}, 2r)$ and a sequence $i_1, \dots, i_k \in [t]$ such that

$$(7.39) \quad \hat{x} = \Phi_{i_k}(\Phi_{i_{k-1}}(\dots \Phi_{i_1}(x_0) \dots)) \text{ and } \left| \hat{x} - \frac{\lambda'}{\lambda} \right| \leq \epsilon' \leq \epsilon/|\lambda|.$$

For convenience, let $i_0 = 0$.

Now, let P be a path of length $k+1$ with vertices labeled $v_0, v_1, \dots, v_k, v_{k+1}$. Let λ be the activity vector on P given by

$$\lambda_{v_0} = \lambda'_0 = \frac{1 - x_0}{x_0}, \quad \lambda_{v_j} = \lambda'_{i_j} \text{ for } j \in [k], \quad \lambda_{v_{k+1}} = \lambda.$$

Then, by Lemma 5.2, it holds that $Z_{P, v_{k+1}}^{\text{out}}(\lambda) \neq 0$ and $\frac{Z_{P, v_{k+1}}^{\text{in}}(\lambda)}{Z_{P, v_{k+1}}^{\text{out}}(\lambda)} = \lambda \hat{x}$; moreover, we can also compute the values $Z_{P, v_{k+1}}^{\text{in}}(\lambda), Z_{P, v_{k+1}}^{\text{out}}(\lambda)$ in time polynomial in $k = \text{poly}(\text{size}(\lambda', \epsilon))$ and $\text{size}(x_0, \lambda'_1, \dots, \lambda'_t) = O(1)$.

Now, let G' be the bipartite graph obtained from the path P by taking for each $j = 0, 1, \dots, k$ a disjoint copy of the graph G'_{i_j} and identifying its vertex w_{i_j} with the vertex v_j of the path P . Using the degree specifications in Lemma 7.12, we have that G' has maximum degree Δ . Moreover, by Lemma 7.12 we have that $\lambda'_i = Z_{G'_i, w_i}^{\text{in}}(\lambda)/Z_{G'_i, w_i}^{\text{out}}(\lambda)$ for all $i = 0, 1, \dots, t$, so analogously to Lemma 3.2 we have that

$$(7.40) \quad Z_{G', v_{k+1}}^{\text{in}}(\lambda) = C \cdot Z_{P, v_{k+1}}^{\text{in}}(\lambda), \quad Z_{G', v_{k+1}}^{\text{out}}(\lambda) = C \cdot Z_{P, v_{k+1}}^{\text{out}}(\lambda),$$

where $C = \prod_{i=0}^t (Z_{G'_i, v_i}^{\text{out}}(\lambda))^{| \{j \in \{0, \dots, k\} \mid \lambda_{v_j} = \lambda'_i \} |}$. We conclude that

$$\frac{Z_{G', v_{k+1}}^{\text{in}}(\lambda)}{Z_{G', v_{k+1}}^{\text{out}}(\lambda)} = \frac{Z_{P, v_{k+1}}^{\text{in}}(\lambda)}{Z_{P, v_{k+1}}^{\text{out}}(\lambda)} = \lambda \hat{x}.$$

Combining this with (7.39), we obtain that G' with terminal v_{k+1} is a bipartite graph of maximum degree Δ which implements λ' with accuracy ϵ . Moreover, using (7.40), we can also compute the values $Z_{G', v_{k+1}}^{\text{in}}(\lambda), Z_{G', v_{k+1}}^{\text{out}}(\lambda)$. \square

REFERENCES

- [1] A. F. BEARDON, *Iteration of Rational Functions: Complex Analytic Dynamical Systems*, Grad. Texts in Math., Springer, New York, 1991.
- [2] F. BENCIS AND P. CSIKVÁRI, *Note on the Zero-Free Region of the Hard-Core Model*, preprint, <https://arxiv.org/abs/1812.07532> (2018).
- [3] Y. BUGEAUD, *Approximation by Algebraic Numbers*, Cambridge Tracts in Math., Cambridge University Press, Cambridge 2004.
- [4] P. BUYS, *On the Location of Roots of the Independence Polynomial of Bounded Degree Graphs*, preprint, <https://arxiv.org/abs/1903.05462> (2019).
- [5] M. CHUDNOVSKY AND P. D. SEYMOUR, *The roots of the independence polynomial of a clawfree graph*, J. Combin. Theory Ser. B, 97 (2007), pp. 350–357.
- [6] A. GALANIS, L. A. GOLDBERG, AND D. ŠTEFANKOVIČ, *Inapproximability of the independent set polynomial below the Shearer threshold*, in 44th International Colloquium on Automata, Languages, and Programming, ICALP 2017, Schloss Dagstuhl, Wadern, Germany, 2017, 28.
- [7] L. A. GOLDBERG AND H. GUO, *The complexity of approximating complex-valued Ising and Tutte partition functions*, Comput. Complexity, 26 (2017), pp. 765–833.
- [8] N. J. A. HARVEY, P. SRIVASTAVA, AND J. VONDRÁK, *Computing the independence polynomial: From the tree threshold down to the roots*, in Proceedings of the Twenty-Ninth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA '18, SIAM, Philadelphia, 2018, pp. 1557–1576.
- [9] J. W. MILNOR, *Dynamics in One Complex Variable*, Acad. Search Complete, Princeton University Press, Princeton, NJ, 2006.
- [10] C. A. NEFF AND J. H. REIF, *An efficient algorithm for the complex roots problem*, J. Complexity, 12 (1996), pp. 81–115.
- [11] V. PATEL AND G. REGTS, *Deterministic polynomial-time approximation algorithms for partition functions and graph polynomials*, SIAM J. Comput., 46 (2017), pp. 1893–1919.
- [12] H. PETERS AND G. REGTS, *On a conjecture of Sokal concerning roots of the independence polynomial*, Michigan Math. J., 68 (2019), pp. 33–55.
- [13] J. RIVERA-LETELIER, *private communication*, 2019.
- [14] A. D. SCOTT AND A. D. SOKAL, *The repulsive lattice gas, the independent-set polynomial, and the Lovász local lemma*, J. Stat. Phys., 118 (2005), pp. 1151–1261.
- [15] A. SLY AND N. SUN, *Counting in two-spin models on d -regular graphs*, Ann. Probab., 42 (2014), pp. 2383–2416.
- [16] L. G. VALIANT AND V. V. VAZIRANI, *NP is as easy as detecting unique solutions*, Theoret. Comput. Sci., 47 (1986), pp. 85–93.
- [17] D. WEITZ, *Counting independent sets up to the tree threshold*, in Proceedings of the Thirty-eighth Annual ACM Symposium on Theory of Computing, STOC '06, ACM, New York, 2006, pp. 140–149.
- [18] A. ZIV, *Relative distance—an error measure in round-off error analysis*, Math. Comp., 39 (1982), pp. 563–569.