

REVIEW

Contrasting Patterns in Crop Domestication and Domestication Rates: Recent Archaeobotanical Insights from the Old World

DORIAN Q FULLER*

Institute of Archaeology, University College London, 31–34 Gordon Square, London WC1H 0PY, UK

Received: 20 September 2006 Revision requested: 8 December 2006 Accepted: 31 January 2007 Published electronically: 10 May 2007

- **Background** Archaeobotany, the study of plant remains from sites of ancient human activity, provides data for studying the initial evolution of domesticated plants. An important background to this is defining the domestication syndrome, those traits by which domesticated plants differ from wild relatives. These traits include features that have been selected under the conditions of cultivation. From archaeological remains the easiest traits to study are seed size and in cereal crops the loss of natural seed dispersal.
- **Scope** The rate at which these features evolved and the ordering in which they evolved can now be documented for a few crops of Asia and Africa. This paper explores this in einkorn wheat (*Triticum monococcum*) and barley (*Hordeum vulgare*) from the Near East, rice (*Oryza sativa*) from China, mung (*Vigna radiata*) and urd (*Vigna mungo*) beans from India, and pearl millet (*Pennisetum glaucum*) from west Africa. Brief reference is made to similar data on lentils (*Lens culinaris*), peas (*Pisum sativum*), soybean (*Glycine max*) and adzuki bean (*Vigna angularis*). Available quantitative data from archaeological finds are compiled to explore changes with domestication. The disjunction in cereals between seed size increase and dispersal is explored, and rates at which these features evolved are estimated from archaeobotanical data. Contrasts between crops, especially between cereals and pulses, are examined.
- **Conclusions** These data suggest that in domesticated grasses, changes in grain size and shape evolved prior to non-shattering ears or panicles. Initial grain size increases may have evolved during the first centuries of cultivation, within perhaps 500–1000 years. Non-shattering infructescences were much slower, becoming fixed about 1000–2000 years later. This suggests a need to reconsider the role of sickle harvesting in domestication. Pulses, by contrast, do not show evidence for seed size increase in relation to the earliest cultivation, and seed size increase may be delayed by 2000–4000 years. This implies that conditions that were sufficient to select for larger seed size in Poaceae were not sufficient in Fabaceae. It is proposed that animal-drawn ploughs (or ards) provided the selection pressure for larger seeds in legumes. This implies different thresholds of selective pressure, for example in relation to differing seed ontogenetics and underlying genetic architecture in these families. Pearl millet (*Pennisetum glaucum*) may show some similarities to the pulses in terms of a lag-time before truly larger-grained forms evolved.

Key words: Domestication, cultivation, cereals, pulses, archaeobotany, *Triticum*, *Hordeum*, *Oryza*, *Vigna*, *Pennisetum*.

INTRODUCTION

This paper will consider archaeobotanical evidence for the evolutionary stages of domestication and the rates of evolution of the domestication syndrome for a select number of the best archaeologically documented crops in the Old World. The quantitative increase in archaeobotanical data in recent years indicates that the origins of crop cultivation was a dynamic and multi-stage evolutionary process that occurred independently in numerous world regions involving numerous crops. This evidence results from growth in archaeobotanical research in recent years. Systematic sampling for archaeobotanical remains, using the technique of flotation in the field, began slowly in the 1960s and 1970s, and led to the collection of much larger and more complete data sets (Pearsall, 2000; Fuller, 2002). In subsequent years, increasing numbers of sites were sampled and more practitioners of archaeobotany have joined the field. In the past two decades, refinements in identification,

analysis and modelling have occurred. Taken together these mean that just in the past few years it has become possible to offer new insights into plant domestication and the multiple origins of crops.

As it has been recognized that domestication is a multi-stage process (e.g. Ford, 1985; Harris, 1989, 1996), models have summarized the expected stages and their archaeological visibility. A classic model is that of Harris (1989), who distinguishes four general stages: (1) wild plant food procurement (true hunting and gathering), (2) wild plant food production (the very beginnings of cultivation), (3) systematic cultivation (of morphologically wild plants) and finally (4) agriculture based on domesticated plants. Domestication results from the earlier stages of wild plant food production and systematic cultivation, making crops more dependent on humans for survival but also more productive. Through all of these stages people put increasing labour effort into a single unit of land and a single field of crops, in other words this tracks an intensification of production. But the reward is increased

*For correspondence. E-mail d.fuller@ucl.ac.uk




FIG. 1. An evolutionary model from foraging to agriculture, with archaeobotanical expectations indicated at the bottom (modified from Harris, 1989). The stages of pre-domestication cultivation are shaded. In this version, domestication is represented as a process of gradual frequency change, with an earlier, more rapid 'semi-domestication' and a later, slower fixation of full domestication. The gap in time elapsed between these two can be taken as a minimal estimate of domestication rate (d.r.).

productivity of that land, and the ability to produce larger surpluses, to feed more people, or accumulate as wealth. An updated version of the Harris chart is given here (Fig. 1), in which I have modified domestication to indicate that it is a protracted quantitative process of change and that those rates may vary for different aspects of that change. Two points should be highlighted: first, that there is necessarily a stage of production (cultivation) that precedes morphological domestication, and archaeobotanists have been developing means to recognize this. Second, this transition represents an important change in human behaviour in terms of the nature of efficiency. While for foragers time is an important measure of efficiency (transport times, processing time, etc.) (see Kelly, 1995), for farmers, who have invested labour in smaller parcels of land for delayed returns, space becomes a crucial metric for efficiency, i.e. yield per unit area (Hillman and Davies, 1990: 69–70; Bar-Yosef, 1998). This cultural change is probably crucial to the change in selective pressures on plants that evolve into domesticates.

AGRICULTURE EVOLVING: THE INTERPLAY OF CULTURAL BEHAVIOUR AND PLANT GENETICS

The origins and evolution of agriculture involves considering how human food-procurement behaviours have changed strategically and how plant populations have evolved in response. The beginnings of food production represents a strategic shift in human behaviour, towards the manipulation of the soil environment (clearance, tillage) and through an influence on the composition of plant populations in that soil (via preferential seeding and tending of one or a few species). From the vantage point of present-day plant populations, it is possible to identify the closest wild relatives of crops, where and in what habitat they

occur, and how these modern domesticates differ from their presumed wild ancestors (e.g. Zohary, 1969; Harlan, 1992). The transition from the ancient wild forms to the modern domesticated forms was an evolutionary process, and it is necessary to ask whether this can be broken down into distinct stages, relating to different selective pressures and human behaviours, and at what rate this evolutionary process took place. Archaeobotany, the study of plant remains from sites of ancient human activity, which is directly correlated through its stratigraphic context with the archaeological record of human behaviour. Archaeobotany, however, has its limits imposed by preservation, and as most remains are preserved by being charred in a fire, we are usually limited to clear morphological distinctions and metrical shifts.

Before considering the archaeobotanical data, we must have in mind a framework for assessing changes. The 'domestication syndrome' provides such a framework by highlighting a set of characters that differ between domesticated crops and their wild ancestors (Harlan *et al.*, 1973; Hawkes, 1983; Zohary and Hopf, 2000; Gepts, 2004). These characters can be related to different aspects of cultivation in terms of what causes them to evolve. It should be noted that the domestication syndrome differs for different kinds of crop-plants. Thus, fruit trees, vines and tubers are not 'domesticated' in the same way as grain crops such as cereals and pulses, in as much as they tend to be reproduced vegetatively and to represent the human choice of favourable mutants (Zohary and Spiegel-Roy, 1975; Hather, 1996; Kislev *et al.*, 2006). This paper will consider only the domestication of seed-propagated herbaceous annuals.

There has been much discussion about how these domestications evolved. Important contributions come from field ecological studies of wild progenitors (e.g. Zohary, 1969; Hillman, 2000), experimental harvesting of wild progenitors using different methods (Harlan, 1967; Ladzinsky, 1987; Unger-Hamilton, 1989; Hillman and Davies, 1990; Anderson, 1992; Lu, 1998; Kislev *et al.*, 2004) and experimental cultivation of wild progenitors (e.g. Oka and Morishima, 1971; Zohary, 1989; Hillman and Davies, 1990; Willcox, 1992, 1999). Nevertheless there remain important areas of debate about the nature of the changes in human practices between collecting and cultivating, and what motivated those cultural shifts, and resulting adaptive changes in the plants. While these changes require sowing of seeds stored from a previous harvest, other selection pressures are involved which can be related to harvesting methods, soil conditions (created through tillage, etc.) and perhaps crop-processing methods. The domestication syndrome in grain crops usually includes the following six criteria.

1. Elimination/reduction of natural seed dispersal

For example, non-shattering rachis in cereals, non-dehiscent pod in pulses. This is often regarded as the single most important domestication trait ('domestication' *sensu stricto*). It makes a species dependent upon the human farmer for survival. It means that instead of shedding seeds when they are mature, a plant retains them.

Instead those seeds must be separated by the addition of human labour (threshing and winnowing), and then the seeds are dispersed by the farmer. Higher yields can be produced because the farmer can now wait until all, or most, of the grains on a plant have matured. This trait can only evolve under conditions of particular kinds of intensive harvesting that favour plants that retain their seeds, followed by sowing from the harvested seeds. All forms of cultivation may not select for this trait. While the evolution of this trait has often been attributed to the early use of sickle tools (e.g. Wilke *et al.*, 1972; Hillman and Davies, 1990; Bar-Yosef, 1998), current evidence suggests that this need not be the case, as discussed below.

2. Reduction in seed dispersal aids

This is connected to the last trait but is selected for in a different way. Plants often have a range of structures that aid seed dispersal, including hairs, barbs, awns and even the general shape of the spikelet in grasses. Thus, domesticated wheat spikelets are less hairy, have shorter or no awns and are plump, whereas in the wild they are heavily haired, barbed and aerodynamic in shape (see Hillman and Davies, 1990). All of these tend to be greatly reduced in the domesticated form. This can be considered to have come about by the removal of natural selection for effective dispersal, and once removed metabolic 'expenditure' on these structures is reduced. This trait might evolve under initial cultivation and be regarded as part of 'semi-domestication'.

3. Trends towards increasing seed/fruit size

This is likely to be selected for by open environments in which larger seedlings have advantages, surviving deeper burial within disturbed soils and thus this trait should be selected for by tillage and cultivation generally (Harlan *et al.*, 1973). Larger seeds are strongly correlated with larger seedlings in many species including cereal and legume crops (Baskin and Baskin, 2001: 214; Krishnasamy and Seshu, 1989). Comparative ecology indicates that larger seeds generally have competitive advantages over smaller seeds under certain kinds of competition including deeper burial (Maranon and Grubb, 1993; Westoby *et al.*, 1996). Although there may be some exceptions, this is demonstrated from cereal relatives (e.g. *Panicum*, *Lolium*, *Avena* and *Aegilops* among grasses; Baskin and Baskin, 2001: 212–213), and in Mediterranean habitats (Maranon and Grubb, 1993). Experimental cultivation of rice found some increase in average grain weight within just five generations (Oka and Morishima, 1971), suggesting this can indeed evolve quickly. This trait is the key archaeological indicator of 'semi-domestication' in cereals.

4. Loss of germination inhibition

In the wild many seeds will only germinate after certain conditions have passed, conditions of daylength, temperature, or after the seed coat is physically damaged. Crops tend to germinate as soon as they are wet and planted. This change is often signalled by changes in the seed,

such as thinner seed coats. This is also selected for simply by cultivation, and sowing from harvested yield, as those seeds that do not readily germinate will not contribute to the harvest. This has produced a particularly important character for the study of New World *Chenopodium* domestications (e.g. Smith, 1989; Bruno and Whitehead, 2003). So far it has been less useful for Old World crops. Although in principle it can be applied to pulses (Butler, 1989; Plitman and Kislev, 1989), seed coats are preserved rarely in charred archaeological specimens.

5. Synchronous tillering and ripening

This sometimes includes a shift from perennial to annual. Planting at one time and harvesting at one time will favour plants that grow in synchronization.

6. More compact growth habit

For example, reduction in branching, e.g. dense spikes or seed heads (the 'sunflower effect'; Harlan, 1995: 199), e.g. from climbing habit to self-standing. Harvesting methods, like those that select for non-shattering types (no. 1, above) will also favour plants with single and compact parts to be harvested.

Of particular importance to the archaeobotanist are those changes that can be identified in archaeological material. This is likely to include nos. 1–4, although no. 4 is only preserved in certain kinds of seeds, and no. 2 may be difficult to recognize because hairs are often destroyed by carbonization. For this reason, especially for most cereals, it is criteria 1 and 3 that archaeologists look at. Grain size (no. 3) is made complicated because of the potentially wide range of variation in modern populations, and the effects of charring (which causes shrinkage and sometimes distortion). If preserved, remains of the cereal ear rachis or spikelet base can provide clear evidence for the mode of shattering (no. 1). In wild types there should be a smooth scar, indicating normal abscission, whereas in domesticated (but also in very immature) plants the scar will be rough because the ear has been broken apart by threshing.

The contrast between shattering and grain size is significant. Tough (or non-shattering) rachis ears occur as a rare, deleterious genetic mutation in most wild grass populations, which has been demonstrated in wild barley (Kislev, 1997; Kislev *et al.*, 2004). If wild cereals were harvested simply by passing through stands and shaking or beating ears to knock seeds into a basket then the shattering, wild-type ears would be the ones to predominate in the next year's crop. Ethnographically this is the most common method documented for wild grass and forb seed gathering (e.g. Harris, 1984; Harlan, 1989). In addition, harvesting experiments in wild wheat and wild *Setaria* millet indicate that this is significantly more efficient in terms of return per unit of labour time (Hillman and Davies, 1990; Lu, 1998). By contrast, if people harvested with a sickle and cutting the entire ear, or plucking individual ears, or pulling plants up from the roots, this would tend to disperse shattering seeds and retain all non-shattering mutants. Therefore, these could be replanted the following year

and over time would come to dominate the population at the expense of wild, shattering types (Hillman and Davies, 1990). Under ideal circumstances of new soils being sown, without a seed bank of the wild form, and sickle harvesting of mature/near-mature plants, and self-pollination or sufficient distance from wild stands, the tough rachis genotype could evolve very rapidly, with estimates of 20–100 years suggested by Hillman and Davies (1990). As will be seen below, however, the archaeobotanical evidence appears to indicate a much slower process.

SCOPE

This paper will attempt to arrive at differing modal trajectories in crop domestication and rates of domestication based on dated archaeological evidence. Of particular significance is the distinction between different aspects of the domestication syndrome. First there is the evolution of grain size, or loss of dispersal appendages, such as hairs and awns, which can be called ‘semi-domestication’. Second there is the loss of wild seed dispersal (non-shattering), i.e. full ‘domestication’. The time between these two developments can be taken to define a domestication rate (‘d.r.’ in Fig. 1). This provides an archaeological estimate of the time it took for fully domesticated types to evolve in population genetics terms, i.e. to come to dominate cultivated populations and be genetically fixed. Archaeological evidence will be reviewed as it relates to the evolution of different aspects of the domestication syndrome. A small selection of crops from the Old World are explored which represent those with the best quantitative archaeobotanical evidence to date, and which are drawn from across four different centres of domestication in Asia and Africa.

This paper will begin by exploring a typical cereal model for domestication, reviewing current evidence for the evolution of the domestication syndrome in some of the main cereals of the Old World: wheat and barley in the Near East, and rice in southern China. Although the available archaeobotanical evidence is quite different, current indications are that similar evolutionary processes took place in each case, with evolution of grain shape and size preceding the loss of wild-type seed dispersal.

This paper will also explore a pulse model of domestication, as the trajectory followed in the evolution of the full domestication syndrome appears to have differed in legumes. In particular, there appears to be a long lag-time between early cultivation and domestication (in terms of seed dispersal and germination inhibition) and seed size increase. In pulses, the seed size increase must be considered a form of cultivar advancement, occurring much later after the initial development of agriculture (Fuller and Harvey, 2006: 257). This can be related to changes in agricultural techniques. This will be explored through the archaeobotanical evidence for two closely related pulses domesticated in India, the mungbean (*Vigna radiata*) and urdbean (*Vigna mungo*). A similar model seems to apply to Near Eastern pulses such as lentils (*Lens culinaris*) and peas (*Pisum sativum*), and possibly east Asian legumes, adzuki bean (*Vigna angularis*) and soybean (*Glycine max*). Finally, the case of West African pearl millet

(*Pennisetum glaucum*) is explored as a case in which a cereal shows some similarities to the pulse domestication model. A brief overview of the current state of archaeobotanical research in each case region is provided (the Near East, China, India and sub-Saharan Africa).

Archaeobotanical evidence is most often preserved carbonized, as the result of charring, and this process affects the size and shape of seeds. In general, carbonization leads to shrinkage. Based on a range of experimental studies (e.g. Lone *et al.*, 1993; Willcox, 2004; Braadbaart and van Bergen, 2005; Nesbitt, 2006: 21), most carbonized seeds appear to shrink between 10 and 20%, with a slight bias towards higher shrinkage in the longest dimension, i.e. a tendency to become more spherical. Therefore, in comparing carbonized seeds with modern seeds some correction factor is needed. In this paper I use a 10% shrinkage estimate for cereals and a higher 20% shrinkage estimate for pulses. The higher estimate for pulses should serve to over-emphasize potential seed enlargement in pulses. The most important evidence, however, remains comparisons between archaeological, carbonized assemblages of different sites and periods.

Note on ages

Throughout this paper ages are referred in calendar years BC or AD, based on the calibration of radiocarbon ages. For measured radiocarbon ages back to 11 000 BP calibration is made with measurements from tree-ring data, while older dates are calibrated by atmospheric ¹⁴C estimated from coral (Reimer *et al.*, 2004). This was done with OxCal v.3.9 software. For this process and software, see the Oxford University Radiocarbon Laboratory webpage: <http://c14.arch.ox.ac.uk/>.

THE CEREAL MODEL FOR DOMESTICATION: WHEAT AND BARLEY

Near Eastern agricultural origins: general background

The Near Eastern centre of crop domestication is often called the ‘fertile crescent’, and, as is well known, a number of crop progenitors can be found in a restricted set of adjacent vegetation zones from the Mediterranean oak woodlands, through the open grassland steppe (Fig. 2). The wild wheats (*Triticum* spp.) and barley (*Hordeum vulgare*) occur in the slightly drier, more open parkland steppe with dispersed shrubs, wild almond trees and oaks, while the wild pulses of south-west Asia (*Lens culinaris*, *Pisum sativum*, *Cicer arietinum*, *Lathyrus sativus*, *Vicia* spp.) occur in the clearings of nearby woodlands and rocky talus slopes (Zohary and Hopf, 2000). That hunter-gatherers had long been exploiting wild cereals when available without cultivation is established from finds at Ohlalo II (21 000–18 500 BC) of wild emmer (*Triticum diococcoides*) and barley (*Hordeum spontaneum*) together with several other small-seeded grasses, fruits and acorns (Kislev *et al.*, 1992; Weiss *et al.*, 2004). Climatic changes at the end of the Pleistocene are regarded as important for their impact on the availability of these wild progenitors and human subsistence and are the




FIG. 2. Map of south-west Asia, showing the locations of sites with archaeobotanical evidence that contributes to understanding the origins and spread of agriculture. Sites are differentiated on the basis of whether they provide evidence for pre-domestication cultivation, enlarged grains, mixed or predominantly domestic-type rachis data. Note that these sites represent a range of periods, and many sites have multiple phases of use, in which case the earliest phase with significant archaeobotanical data is represented. Shaded areas indicate the general distribution of wild progenitors (based on Zohary and Hopf, 2000, with some refinements from Willcox, 2005). It should be noted that wild emmer (*Triticum dicoccoides*) occurs over a sub-set of the wild barley zone, and mainly in the western part of the crescent.

favoured component underpinning explanations for why cultivation began. Of particular importance is the Younger Dryas dry (cold) episode from approx. 11 500 to 9800 BC (Bar-Yosef, 1998, 2003; Harris, 1998; Hillman *et al.*, 2001; Byrd, 2005). Prior to this event the climate was favourable and dense populations of hunter-gatherers (the Late Epipalaeolithic phase, 12 500–9700 BC) were settled in territories that included some possible year-round settlements. The Younger Dryas brought an end to this, with most sites being abandoned, and it is argued that a few groups may have resorted to cultivation during this period. Village populations reappeared throughout the area during the Pre-Pottery Neolithic A (PPNA) Period (9700–8700 BC), and many of them appear to have been cultivators. Finds of domesticated plants are generally widespread in the subsequent Pre-Pottery Neolithic B (PPNB; 8700–6200 BC), and it is by this period that they began to spread beyond the domestication zone into central Turkey, Cyprus, Crete and southern Greece. Domesticated animals first occur about 8200 BC at the start of the Middle PPNB (Garrard, 2000; Bar-Yosef, 2003; Colledge *et al.*, 2004; Byrd, 2005).

In recent years two important advances have occurred in archaeobotany: the recognition of pre-domestication cultivation and evidence for different sub-centres of crop domestication within this region. Evidence for pre-domestication cultivation has been recognized through the statistical composition of wild seed assemblages, for nearly 10 years (Colledge, 1998, 2001, 2002; Harris, 1998; Willcox, 1999, 2002; Hillman, 2000; Hillman *et al.*, 2001).

As is well known from later agricultural periods, archaeobotanical assemblages are made up predominately of crops and weeds, together with some gathered fruits and nuts (Jones, 1985). This pattern is already recognized in the PPNA and in some Late Epipalaeolithic sites (Fig. 2), by samples dominated by wild cereals together with seeds of herbaceous taxa that flourish in disturbed soils, which are best known today as arable weeds. This includes taxa such as small legumes (Trifoliaceae), Boraginaceae, small-seeded grasses (*Hordeum murinum*, *Bromus* sp., *Lolium* sp., *Lepturus pubescens*, *Lolium* sp.), Polygonaceae, *Silene* (Caryophyllaceae), *Galium* (Rubiaceae) and *Centaurea* (Asteraceae). Modelling of the impact of Younger Dryas climate change on species availability indicates that only cereals and these weeds show frequency increases against the grain of the expected climate trends through the sequence at Abu Hureyra, suggesting their persistence in a new habitat: the arable field (Hillman *et al.*, 2001). This is also suggested by the continued association of these same weeds with domesticated cereals in later periods (from the PPNB, 8700–6200 BC, and onwards).

A number of advances in recent years have made it increasingly clear that separate histories need to be traced for different sub-regions, which were separate in as much as the beginnings of agriculture relied on different combinations of species. Some suggestive evidence comes from genetics. For example, in barley (*Hordeum vulgare*) two variant genes control whether or not the ear shatters. A recessive mutation in either gene locus leads to the domesticated condition, while the dominant variant at either locus

confers wild-type shattering ears (Zohary and Hopf, 2000: 59–60). The existence of these two variants argues for two domestications for barley. Recent work on emmer wheat has identified two different lineages of a gluten gene which are so different that they are estimated to have evolved apart 100 000s of years ago, and thus amongst wild emmer wheat, long before domestication. Such evidence implies two separate domestications of emmer (Allaby *et al.*, 1999: 305; Brown, 1999; reviewed in Jones and Brown, 2000). Another source of evidence for multiple domestications of the ‘same’ (or similar) crops comes from refinements in archaeobotanical identification criteria. Thus, for example, it is possible on the basis of grain shape to distinguish einkorn wheat (*Triticum monococcum*) with single-grained spikelets (from wild *Triticum boeoticum* subsp. *aegilopoides*) from einkorn with two-grained spikelets (from wild *T. boeoticum* subsp. *thaudar* or *T. urartu*). Modern domesticated einkorn (*T. monococcum*) is normally only one-grained, but archaeobotanical evidence indicates the presence of one of these two grained forms as a wild cereal from the late Pleistocene in Syria (Hillman, 2000; Willcox, 2002, 2005), and later as a domesticated cereal in Syria, Turkey and into Neolithic Europe (Kreuz and Boenke, 2002). It persists in parts of Europe as late as the Iron Age, after 1000 BC (Kreuz and Boenke, 2002; Köhler-Schneider, 2003), and disappears in its Syrian homeland during the Chalcolithic, approx. 5000 BC (Van Zeist, 1999; Kreuz and Boenke, 2002). This implies an additional two-grained einkorn domestication but this crop went extinct in prehistory. Similarly, there is now evidence for an extinct emmer-like wheat, with distinctive glume architecture not yet paralleled in any studied modern wild or cultivated landrace but with some similarities to *Triticum timopheevi* (Jones *et al.*, 2000; Köhler-Schneider, 2003). It is known archaeologically from the eastern half of Europe, Turkey and Turkmenistan. It persists in parts of Europe as late as the Bronze Age (approx. 1000 BC) (Köhler-Schneider, 2003). In addition, early sites in Syria appear to have cultivated a local form of rye (*Secale* cf. *montanum*), but rye did not become a major crop of the Neolithic Near East despite occasional later finds (Hillman, 2000: 392), and was probably a different species from the later European rye (*Secale cereale*), domesticated from a field weed in Late Bronze Age or Iron Age times (approx. 1000 BC) (Küster, 2000). Taken together the archaeobotanical morphotypes and genetics suggest a minimum of seven domestications of wheat and barley in the Near Eastern Fertile Crescent region, and there is no reason to attribute them all to a single micro-region or a single process of agricultural origins, but at least two or perhaps three (Willcox, 2005).

Cereal grain size increase

There is a growing morphometric database for wheat and barley from the Near East (Colledge, 2001, 2004; Willcox, 2004). This indicates that wheat and barley grains increased in size starting in the PPNA and earliest PPNB. This is before clear and widespread evidence for tough rachises

and loss of natural seed dispersal. It is well known that wild and domesticated cereal grains differ in size and this has been used to infer the domesticated status of cereals, already in the PPNA and the earliest PPNB, including sites from the Jordan Valley (e.g. Fig. 3), the upper Euphrates in Syria, and the first settlements on Cyprus (Colledge, 2001, 2004).

This evolutionary shift can be illustrated from internal archaeobotanical evidence too, i.e. by charting grain metrics through time, as recently demonstrated with samples from the site of Jerf el Ahmar on the Upper Euphrates (Willcox, 2004). Fig. 4A shows the contrast between the barley grains from the early phase at Jerf el Ahmar (9500–8800 BC) and the much later Chalcolithic site of Kosak Shamali (approx. 5000 BC), in which all of the grains are larger and comparable with the domesticated size range inferred from modern material. If we look at the later phase at Jerf el Ahmar (approx. 8500 BC), however, it can be seen that many of the grains are of the larger size (Fig. 4B). This implies evolution towards larger grain size during the occupation of this site, but recovered rachis remains indicate that ears were still of the wild, shattering type. A similar pattern is found for the einkorn wheat grains (*Triticum monococcum*), which also include some mixture of rye (*Secale cereale*) (Fig. 4C, D). Given that occupation of Jerf el Ahmar lasted less than a millennium and perhaps only 500 years, we can suggest the rate of evolution of large grains, as occurring in a few centuries and certainly more rapidly than a millennium. This explains why large domestic-type grains are already widespread and predominant on most Near Eastern sites by the start of the PPNB (approx. 8700 BC). From the even earlier grain assemblage from Abu Hureyra, where cultivation has been inferred from the arable weed flora, a few large and plump rye grains were found, comparable with domesticated grains (Hillman, 2000; Hillman *et al.*, 2001). These might represent the earliest indications of selection for grain size increase under cultivation, but possibly a local dead-end trajectory.

This evidence raises the question of how large-grained varieties evolved. One possibility is that methods of




FIG. 3. Pre-Pottery Neolithic B wheat grain measurements from the Jordan Valley (after Colledge, 2001). This indicates the predominance of the larger domesticated-type grains. The gap between the two groups is comparable with that in modern reference material. These sites are dominated by wild-type barley rachis (see below).




FIG. 4. Scatter-plots of archaeological grain measurements showing the increase in grain size under early pre-domestication cultivation (after Willcox, 2004). (A) Barley grain measurements, comparing early Pre-Pottery Neolithic A Jefer el Ahmar with the much later domesticated material from Kosak Shamali. (B) Comparing early and late Jefer el Ahmar, indicating that shift towards larger grain size had already occurred. (C) Similar comparison of einkorn grains (probably including some rye grains) at early Jefer el Ahmar and Kosak Shamali. (D) Trend towards larger grain sizes over the course of Jefer el Ahmar occupation.

processing, such as using sieves after threshing and winnowing, served to bias stored cereals towards larger grains (on traditional crop-processing and the use of sieves, see Hillman, 1984). This, however, would tend to remove 'tail grain', i.e. the smaller grains on an ear, and would be expected to make very weak selection for genetically larger-grained plants, as the selected grains would be from the same genetic population as the discarded small grains. More likely seems to be the role of tillage. The practice of preparing land with tillage is likely to have been necessary to remove competing vegetation from favourable micro-environments for early cultivation, such as the breaks of slopes where ground-water tables are higher. Such areas would naturally be cloaked in herbaceous flora and undershrubs such as *Tamarix*, so tillage would be necessary (Hillman, 2000: 396; Hillman *et al.*, 2001: 387). Sowing grains into tilled soil with an underlying water table would give advantages to those best able to germinate successfully from greater burial, namely larger-seeded individuals.

Evolution of the tough rachis

The evolution of non-shattering ears was also a gradual process. Although theoretically it could have happened very quickly, as demonstrated under ideal experimental

conditions (Hillman and Davies, 1990), the archaeobotanical evidence indicates a much more gradual evolution of non-shattering ears. A recent quantitative assessment of the wheat and barley rachis remains from a few sites suggested a gradual increase in the proportion of the domesticated-type ears over the course of the PPNB (Tanno and Willcox, 2006a). A larger data set has been compiled here in which evidence for einkorn wheat spikelet bases and barley rachis segments are considered (Fig. 5A, B). In general, there is contrast between early sites which are largely or entirely of the wild-type chaff remains and later sites dominated by domesticated-type remains, with some intermediate proportions for sites chronologically in the middle. Although there are individual site exceptions, such as barley rachis from Jordan valley sites (e.g. Wadi Jilat), these may relate to particular local circumstances of higher reliance on wild plant foods (indicated by other data from this site, Colledge, 2001). In addition, wild cereals, especially barley, may persist as weeds of cultivation. Nevertheless, at a regional level the overall trend is clear.

These data can then be transformed to produce a regression of domestication rate over time (Fig. 6A, B). In these diagrams individual site proportions are plotted, both by the minimal figure, based on just the clear non-




FIG. 5. Proportions of wild and domesticated barley and einkorn rachis/spikelet remains on early Near Eastern sites, arranged chronologically from left to right, and grouped into broader phases. (A) Barley rachis types, including domesticated (tough), wild (shattering) and uncertain (but more likely wild). (B) Proportions of einkorn spikelet forks/glume bases, including domesticated (tough), wild (shattering) and uncertain (but more likely domesticated). Sites, approximate ages and data sources: Ohalo 2, 21 000–18 500 BC (Kislev *et al.*, 1992); Wadi Hammeh, approx. 12 000 BC (Colledge, 2001); Mureybit, 10 500–9500 BC (Van Zeist and Bakker Heeres, 1986); Iraq-ed-Dubb, approx. 9300 BC (Colledge, 2001); Jerf el Ahmar (early, with two-grained einkorn), 9700–9300 BC (Willcox, 1999, 2002); Wadi Jilat 7, 8800–8300 BC, Wadi Jilat 13, 7000–6500 BC (Colledge, 2001); Aswad, 8700–8000 BC (Van Zeist and Bakker Heeres, 1985; Tanno and Willcox, 2006a); Azraq 31, 7500–7000 BC (Colledge, 2001); Wadi Fidan A, 7500–7000 BC, Wadi Fidan C, 7000–6500 BC (Colledge, 2001); El Kowm, 7500–6800 BC (De Moulins, 1997); Catal Hoyuk, 7400–6800 BC (Fairbairn *et al.*, 2002); Ramad, 7500–6500 BC (Van Zeist and Bakker Heeres, 1985; Tanno and Willcox, 2006a); Magzaliyeh, 7100–6400 BC (Willcox, 2006); Tell el Kherkh, 8600–8300 BC (Tanno and Willcox 2006a, b); Nevali Cori, 8500–8000 BC (Tanno and Willcox, 2006a); Qaramel, approx. 10000 BC (Tanno and Willcox, 2006a); Netiv Hagdud, 9500–9000 BC (Kislev, 1997); Cafer Hoyuk, (IX–XIII) 8300–7700 BC, (III–VIII) 7500–7000 BC (De Moulins, 1997); Kosak Shamali, approx. 5000 BC (Tanno and Willcox, 2006a).

dehiscent type, and a larger figure which includes uncertain but possible domesticates. The true proportion of domesticates should fall within this range. Sites are plotted against a median age estimate for each site. In addition,

for barley the averages from each phase (e.g. PPNA, Early PPNB, Late PPNB) are plotted. The trend from wild dominance to domesticated dominance is clear but appears fairly gradual and slow. The rates of evolution do




FIG. 6. Domestication rates in barley and einkorn modelled from archaeobotanical data (based on Fig. 5). Proportion of domesticated type for each site is plotted by a box against a median estimate of site age. A margin of error is indicated by the line which connects the sum of domesticated and uncertain types (indicated by a cross or \times). Trend lines are shown based on the lower estimate. (A) Barley domestication rate model, on which period averages are also plotted for the PPNa, Early PPNB and Late PPNB, in which the diamond indicates the proportion of domesticated types and the circle the sum of domesticated and uncertain types. (B) Einkorn domestication rate model; the much later Kosak Shamali has been excluded.

not come anywhere near the 20–100 years estimated by Hillman and Davies (1990) on the assumption of sickle harvesting morphologically wild near-mature plants, or uprooting of whole plants. This vast difference in rate raises questions about how we explain the selection for domesticated non-shattering genotypes, an issue to which I return below.

From these data we can also suggest that domestication occurred somewhat more quickly in wheat, perhaps around 1500 years, as opposed to barley which shifts over a period of 2000 years or slightly more. An earlier domestication date for wheat has been previously noted (e.g. McCorrison, 2000). A number of factors might be suggested to account for this. Slightly higher average rates of cross-pollination are reported in barley, estimated at about 1–2% (Allard, 1988; Morrell *et al.*, 2005), as opposed to wheats, which are less than 1% (Hillman and Davies, 1990: 62), although this seems unlikely to be sufficient in itself. Of importance may be the fact that wild

barley (*Hordeum spontaneum*) is more prone to be a weed of cultivation, thereby maintaining introgression, whereas wheat cultivation, which is more water-demanding and may have been more carefully located, would have been cut off from the wild progenitor. In addition, the persistence of wild barley as a weed will affect the composition of archaeobotanical assemblages and rachis counts may therefore be an underestimate of the proportion of domesticated types among crop populations.

Most remarkable is the difference in evolutionary rate by comparison with grain size (semi-domestication), of 500–1000 years. The shift in rachises (full domestication) appears to start mainly after large grain size had already begun to evolve, and we might therefore suggest a minimum estimate of 2000 years for the evolution of both aspects of the domestication syndrome.

CHINESE RICE DOMESTICATION: THE IMPACT OF UNEVEN RIPENING

Chinese agricultural origins: background

Systematic archaeobotany in China has only recently begun and evidence for the beginnings of agriculture is still limited (Fig. 7). In northern China the millets *Setaria italica* and *Panicum miliaceum* were the initial crops. The earliest well-documented millets are from approx. 6000 BC at Xinglonggou, in Eastern Inner Mongolia, by which time plump-grained *Setaria* were already established but grains of *P. miliaceum* were near the wild-type in size and shape (Zhao, 2005). Subsequently, millet cultivation was established in much of the Yellow River basin by 5500 BC (the Peiligang, Cishan, Beixin and Dadiwan cultures) (Lu, 1999; Crawford, 2005; Crawford *et al.*, 2005). Rice from the south was added to this agricultural system only in the third millennium BC, with a few rice finds from Late Yangshao contexts (3000–2500 BC) and many more from the Longshan period (2500–2000 BC). In recent years, the orthodoxy has been that rice agriculture began early, perhaps at the start of the Holocene or late Pleistocene, in the Middle Yangtze, perhaps amongst seasonally inhabited cave-sites (e.g. Lu, 1999; Crawford, 2005). Critical re-assessment, however, suggests that rice may have been independently domesticated in both the middle and the lower Yangtze river areas, and that domestication was later than has been presumed (Fuller *et al.*, 2007).

Domestication as morphometric and maturity shift

It is now well established that *Oryza sativa* was domesticated more than once, with distinct origins of tropical monsoonal *indica* and marshland, sub-tropical *japonica*. Genetic studies are now numerous in support of these separate origins (e.g. Chen *et al.*, 1993; Cheng *et al.*, 2003; Londo *et al.*, 2006). Archaeological evidence also favours distinct centres of early rice cultivation in the Middle Ganges valley, India (Fuller, 2002, 2006), and the Middle or Lower Yangtze, China (Lu, 1999). This situation falsifies the older suggestions regarding Chinese Neolithic rice as




FIG. 7. Map of East Asia indicating early millet and rice sites, with inset of Yangtze region, showing archaeological sites mentioned in this article: 1, Hemudu; 2, Tianluoshan; 3, Kuahuqiao; 4, Shangshan; 5, Liangzhu; 6, Majiabang area, including Nanzhuangqiao, Luojiqiao and Pu'anqiao; 7, Nanhebang; 8, Maqiao; 9, Songze; 10, Xujiawan; 11, Chuodun; 12, Weidun; 13, Longnan and Caoxieshan; 14, Qiucheng; 15, Longqiuzhuang; 16, Sanxingcun; 17, Lingjiatan; 18, Jiahu; 19, Bashidang; 20, Pengtoushan.

predominantly *indica*, or an ancient 'intermediate type' that evolved into both *indica* and *japonica*. In light of this, I and collaborators have reconsidered published sources on early Chinese rice, and begun new archaeobotanical investigations (Fuller *et al.*, 2007). Similar research has been recently pursued on Indian rice (Harvey, 2006). Previous studies essentially began with the assumption that archaeological rice was a crop, and only asked instead whether it was *indica*, *japonica* or something 'intermediate' (e.g. Oka, 1988; Zhang, 2002). In fact, the morphometric data of early rice often fit well within the range of wild species, including *Oryza rufipogon* but also sometimes *Oryza* spp. not in the *sativa*-complex. On this basis we have suggested that another wild species might sometimes have been utilized as a hunter-gatherer resource (Fig. 8A, B). Based on modern geographical distributions *O. officinalis* might be suggested (cf. Vaughan, 1994), and this species produces substantial grain numbers per plant, although further consideration of feasibility must take into account potential stand sizes (generally small), as well as potential processing techniques that would be required for dehusking this thick-hulled species. Other *Oryza* species or a particularly small-grained race of *O. rufipogon* should also be considered.

Much of the grain, however, is notably long and skinny, which could also be a trait of immature rice (Fig. 8B). Consideration of rice development, with its extended period of uneven ripening (more than 2 weeks, Hoshikawa, 1993), and expectations from optimal foraging

theory and ethnographic expectations about harvesting lead us to expect early users of wild rices to have targeted immature plants to maximize their recovery of near-mature grains. Indeed in order to maximize grain yields wild rice should be targeted in the first 3 or 4 d after the very first grains mature, but at this time slightly more than 40% of the grains are still immature by more than 6 d. Within a week potential yields are vastly reduced by something on the order of 70–80% of mature grains (Fig. 9). As with ethnographically documented hunter-gatherers, we would therefore expect wild rice to be harvested when immature (e.g. the Bagundji wild *Panicum* gatherers of Australia: Allen, 1974; cf. Harris, 1984).

Archaeologically, the morphometric data available from Kuahuqiao and Longqiuzhuang (mainly of the earlier Majiabang period, which equates with the Hemudu period) suggests that grain assemblages are dominated by immature grains (Fig. 8B). It must be noted that this assumes the mature grains would have been in the range of modern domesticates, but they do appear somewhat plumper than expected for mature *O. rufipogon*. This in fact is plausible if we consider the likelihood that early pre-domestication cultivation had already begun to select for larger grains. It must be noted, however, that grain size evolution in rice has been complex and not always towards larger grain sizes. Many short-grained *japonica* races may have shorter, but often still plumper, grains than those in the wild progenitor. It is, nevertheless, clear that grain shape has changed under domestication and




FIG. 8. (A) Scatter plot of length and width of grains measured in modern populations (15 grains measured from 72 populations; Fuller *et al.*, 2007). (B) Grain measurements from selected Neolithic sites, showing maximum and minimum measured ranges with solid lines and statistical standard deviations with dashed lines (as reported). Note that grains from Kuahuqiao, Bashidang and the lower (Majiabang period) levels (8–6) at Longqiuzhuang fall largely or entirely in the expected immature grain proportions, while the latest grains from Longqiuzhuang, Songze period (level 4), indicate a clear shift towards longer and fatter grains that can be regarded as fully mature, and thus domesticated. Choudun (Late Majiabang) also indicates a shift towards mature *japonica*-type grains, but suggests local population differences from the domesticated rice at Longqiuzhuang. The small grains from Jiahu are suggestive of wild rice not in the *sativa* complex, such as *O. officinalis*. Sources: Kuahuqiao (Zheng *et al.*, 2004b), Longqiuzhuang (Huang and Zhang, 2000), Jiahu (Henan Provincial Institute of Archaeology, 1999), Chuodun (Tang, 2003) and Middle Yangzte Bashidang (Pei, 1998).

may also be affected by the proportions of mature and immature grains represented in a harvest.

In addition, the reduction in hairs on the awns of rice recovered from Hemudu (Sato, 2002) implies relaxation of natural selection for seed dispersal aids, which would be expected under cultivation. The Hemudu culture is rightly famous for its numerous wood and bone spades, tools which imply soil manipulation and probably planting. This is in contrast to Kuahuqiao, which produced just two poorly made proto-type spades. The slight difference in grain thickness between Kuahuqiao (6000–5400 BC) and the Majiabang period Longqiuzhuang (5000–4000 BC) could therefore suggest selection for larger grains under early cultivation, and we might therefore suggest the beginnings of cultivation between the end of the Kuahuqiao phase (approx. 5400 BC) and the early Hemudu/

Majiabang phase (from 5000 BC). This would imply a rate of evolution on a par with the grain-size increase in wheat and barley. Thus, the evidence from Hemudu and the earlier Majiabang period (from early to mid fifth millennium BC) both suggest pre-domestication cultivation, which may have started already at the end of or after the Kuahuqiao period. The rice at this stage can be regarded as semi-domesticated, like that of PPNA to early PPNB wheat and barley.

A clear contrast is seen with assemblages from around 4000 BC. The latest assemblage from Longqiuzhuang (early Songze period, just after approx. 4000 BC) has grains which are significantly longer, plumper (2.5–3 mm) and most likely fully mature. In addition, significantly plumper grains have been recovered from Chuodun (late Majiabang, just before approx. 4000 BC). This suggests




FIG. 9. (A) Graph indicating the expected relative frequency of grains reaching maturity during eight stages, of 2 d each, on an individual rice plant (based on anthesis data of a modern *japonica* cultivar, from Hoshikawa, 1993). (B) A graph converting this data into potential grain yields to a hunter-gatherer if this were a morphologically wild plant. This indicates the proportion of grains more than 6 d immature which would differ in grain proportions from mature grains. Although this represents an individual plant it must be assumed that a population of wild rice has individuals that begin this process at different times over a period of weeks. With cultivation there should be a tendency for plants to become synchronous.

an important morphological shift in archaeological rice occurred in the Lower Yangtze region during the later fifth millennium BC. This shift seems most likely to be due to a shift towards harvesting of mature panicles as opposed to immature panicles, rather than an evolutionary development in grain shape. Such a shift would imply that it became feasible to allow grains to mature on the plant without loss of the grains, or in other words that tough, domesticated-type rachises had evolved to dominate the rice populations being harvested. This hypothesis is currently being tested by myself and collaborators on large assemblages of preserved rice spikelet bases from the fifth millennium BC site of Tian Luo Shan. Additional evidence comes from measurements on bulliform phytoliths, silica bodies from the leaves of rice (Fig. 10). These bulliforms shift towards larger sizes between the Majiabang and Songze periods, i.e. before and after 4000 BC. Modern studies suggest that the size of these phytoliths relates to plant maturity and tillering stage (Zheng *et al.*, 2003).

These data suggest a rate of evolution from the beginning of cultivation to full domestication on the order of 1000–1500 years, similar in magnitude but slightly faster than that in the Near East. An important factor in this is that Late Majiabang period sites are known that preserve early paddy field systems. These consist of networks of shallow




FIG. 10. Size increase in Lower Yangzi rice phytoliths. (A) Measured horizontal length and vertical length of rice bulliform phytoliths from Majiabang period samples (M); (B) measurements from samples of the subsequent Songze (S) and Liangzhu (L) phases. The dashed oval represents the distribution of the Majiabang measurements. Data re-plotted from Zheng *et al.* (1994, 2004a, b) and Wang and Ding (2000).

pools, and ditches, as at Caioxieshan and Chuodun (Zou *et al.*, 2000; Gu, 2003). This development in cultivation techniques would imply greater separation of sown populations from wild stands, and a reduction therefore in cross-pollination with free-growing wild rice, which has been estimated at 10–50% (Oka and Morishima, 1967). This separation into distinct field systems may be highly significant, as, theoretically, out-breeding species should evolve more slowly towards adaptations to cultivation (Allard, 1988).

Also important is evidence for climatic change, which is likely to have been reducing populations of wild rice. Globally the mid Holocene, after 6000 BC, and especially between 5000 and 3000 BC, was a period of cooling, as

indicated for example by declining methane levels from Greenland ice cores (Blunier *et al.*, 1995). This is linked to declines in monsoon rainfall in East Africa (Gasse, 2000), South Asia (Madella and Fuller, 2006) and East Asia (Wang *et al.*, 1999). This process would be expected to have impacted South China, pushing more tropical taxa such as wild rice further south. Also significant is pollen core evidence from the Lower Yangtze region that indicates a marked decline in broad-leaved trees, including oaks, after 5200 BC (Yu *et al.*, 2000; Yi *et al.*, 2003; Tao *et al.*, 2006).

This is significant because evidence from Kuahuqiao, Hemudu and Tian Luo Shan suggests that it was acorns (*Quercus*, *Lithocarpus*, *Cyclobalanopsis*) that were the dietary staples of these cultures. These were stored in vast quantities in storage pits, along with water chestnuts (*Trapa bispinosa*) (Zhejiang Province Institute of Archaeology, 2003; 2004; my unpubl. data). As comparative cases indicate, e.g. ethnographic California or the archaeology of the Jomon culture of Japan, tree nuts, and particularly acorns, make ideal staple food sources (e.g. Heizer and Elsasser, 1980: 82–144; Takahashi and Hosoya, 2002; Barlow and Heck, 2002). Therefore, the decline in nut-bearing trees may have promoted increased reliance on, and cultivation of, rice (Fuller *et al.*, 2007).

These climatic factors may also explain the apparent difference in the domestication rate between Near Eastern cereals and East Asian rice. In the Early Holocene, the climate was warmer and wetter and therefore conducive to the expansion and persistence of wild wheat and barley stands in the Near East. By contrast, in Mid-Holocene China wild rice would have been in decline and on its way to local extinction. This may have promoted a somewhat faster rate of domestication.

THE PULSE DOMESTICATION MODEL: SEED SIZE AS EPIPHENOMENON

The domestication syndrome for pulses is essentially the same as that for cereals, with increases in seed size, reduced pod-shattering and importantly loss of germination inhibition (Plitman and Kislev, 1989; Smartt, 1990; Zohary and Hopf, 2000). This need not imply, however, that the rates of domestication and the order in which these traits evolved are necessarily the same. In fact, evidence suggests a difference from cereals. This can be illustrated with evidence for the early evolution of the Indian mungbean (*Vigna radiata*) and urdbean (*Vigna mungo*) under cultivation.

South Asia has received less attention from archaeologists studying early agriculture, although a large number of minor cereals (millets), pulses, cucurbitaceae crops and *indica*-type rices were domesticated there. Archaeology together with the modern distributions of wild progenitors suggest as least three, and perhaps five, distinct centres of plant domestication in India, including South India, Gujarat and the Ganges Plain (for a comprehensive review, see Fuller, 2006; also Fuller, 2002). These domestication events are associated with the mid-Holocene, between 4000 and 2000 BC, and occurred

amongst local populations of hunter-gatherers in monsoonal India, although some influences from the west due to the spread of livestock cannot be ruled out. In the north-west, early agriculture was based on the dispersal of wheat, barley and winter pulse cultivation from the Near East, together with domesticated animals. Despite the long delay before agriculture developed in other parts of India the earliest phases suggest local packages of crops into which wheat, barley or Near Eastern pulses were adopted.

The current distribution of wild mung and urd, taken together with their archaeological record, suggests three areas of domestication (Fig. 11). *Vigna radiata* was probably brought under cultivation in the north-west, perhaps in the Himalayan foothills in the Punjab region, and in the far south, as early finds in both regions are widely separated. Early finds of urdbean come from Gujarat and the northern Peninsula where wild populations of this species persist (*Vigna mungo* var. *sylvestris*) without associated wild mungbean (*Vigna radiata* var. *sublobata sensu stricto*) (Fuller and Harvey, 2006; see also Tomooka *et al.*, 2003). Early finds in South India of *V. radiata* occur in the driest savannah environments of the peninsula together with large quantities of horsegram (*Macrotyloma uniflorum*) and small millets (*Brachiaria ramosa*, *Setaria verticillata*), and in some cases associated with introduced domesticates (Fuller *et al.*, 2004). The association with other crops, their large quantity and the difference from the habitat of the wild progenitor all imply cultivation of these early mungbeans (on the past environmental conditions at these sites, see Fuller and Korisettar, 2004).

These early Neolithic pulses, however, do not appear to be different in size from their wild progenitor (Fig. 12A, B). Even assuming the upper end of 20% shrinkage, measured specimens from southern Neolithic sites fall almost entirely within the size range represented by modern wild populations (Fuller and Harvey, 2006). By contrast, those from Early Historic to Early Medieval periods fall in the range of modern domesticated populations (Fig. 13A). These data suggest on peninsular India selection for increasing pulse seed size was slight through the second millennium BC but had occurred by the first millennium AD. The evidence from late Chalcolithic Tuljapur Garhi (late second millennium BC to early first millennium BC) shows a wide size range from wild-type to domesticated-type, although the average falls in the latter range. This site might represent evidence for the actual process of size increase, suggesting that this occurred most markedly during the late second millennium BC through the Iron Age.

These data raise the question of what change in the environmental conditions, most likely in terms of agricultural practices, selected for seed enlargement in these pulses. Presumably domestication in terms of dispersal and germination was selected for early and rapidly, necessary to make any pulse a worthwhile crop (Ladzinsky, 1987, 1993; Zohary, 1989; Blumler, 1991). If it was being routinely harvested, either by pod-plucking or plant uprooting, and sown from stores we would expect domesticated




FIG. 11. A map of the wild progenitors of *Vigna radiata* and *V. mungo* in India in relation to the moist deciduous forests and the region with extensive wild rice populations (based on Tomooka *et al.*, 2003; Fuller and Harvey, 2006). Archaeobotanical finds of the *Vigna* pulses are indicated which include secure species-level identifications. Sites are numbered: 1, Semthan; 2, Hund; 3, Balu; 4, Kunal; 5, Burthana Tigrana; 6, Mitithal; 7, Hulas; 8, Hulaskhera; 9, Charda; 10, Imlidh-Kurd; 11, Narhan; 12, Khairadih; 13, Malhar; 14, Senuwar; 15, Tokwa; 16, Mahagara; 17, Koldihwa; 18, Balathal; 19, Babar Kot; 20, Rojdi (two phases); 21, Oriyo Timbo; 22, Kaothe; 23, Tuljapur Garhi; 24, Paithan; 25, Apegaon; 26, Bhokardan; 27, Nevasa; 28, Inamgaon; 29, Terr; 30, Golabai Sassan; 31, Piklihal; 32, Hallur; 33, Tekkalakota, Kurugodu, Sanganakallu and Hiregudda; 34, Hattibelagallu; 35, Sanyasula Cave; 36, Veerapuram; 37, Rupanagudi; 38, Ramapuram, Hanumantaraopeta and Peddamudiyam; 39, Kodumanal; 40, Perur. For details of primary sources, see Fuller and Harvey (2006).

forms to have evolved with less dehiscent pods and loss of dormancy. But size increase was not part of the initial domestication syndrome. What explains the delay? I would suggest that changes in agricultural techniques, such as deeper tillage (with ards), created a selective advantage for larger seeded genotypes (Fuller and Harvey, 2006: 257). In this regard the shift during the Iron Age (first millennium BC) is significant as this is the period when ard tillage began on the Indian peninsula, and perhaps in the latest second millennium BC (cf. Shinde, 1987; Fuller *et al.*, 2001).

Measurements from northern India suggest a similar pattern in the middle Ganges and Orissa, but suggest a

contrast with early pulses from the north-west (Fig. 13D). What can be seen in the plot is that Neolithic seeds fall largely in the expected primitive size range, while later sites such as late Chalcolithic Narhan and later Iron Age and Early Historic sites have large *Vigna* grains. Interestingly, two early sites, Balu and Kunal from the later third millennium BC, also have large *Vigna* seeds. Both of these sites, however, are within the Eastern Indus Civilization cultural zone, where we expect deep tillage with ards to have been the norm (Allchin and Allchin, 1997: 170; Miller, 2003). It might be suggested that in the Indus zone enlarged seeds have already been selected for by tillage in the third millennium, whereas further




FIG. 12. Metrical data for Indian *Vigna* pulses indicating no size increase with domestication. (A) Modern length–width measurements in *Vigna radiata*, *V. mungo* and their wild progenitors. The separation between them has been adjusted for a high estimate of 20% shrinkage with carbonization and shown on the other plots as a dashed line. (B) Neolithic *Vigna* finds from South India (before 1400 BC) (data from Fuller and Harvey, 2006).

east this process did not occur until the late second millennium BC. This of course raises the questions of whether the *Vigna* pulses introduced into the middle Ganges zone had diffused before emergence of large-seeded forms, or had diffused from the south rather than the Harappan zone, or whether the absence of positive selection through tillage could have led to a reversion to smaller seed size.

This model provides an explanation for the archaeological difficulty in identifying the beginnings of pulse domestication. Pods are not normally preserved, and germination characters are cryptic in rarely preserved seed coats (Butler, 1989). Only in peas does a clear distinction exist between smooth-testa domesticates (*Pisum sativum* spp. *sativum*) and tuberculate-testa wild peas (*Pisum sativum* ssp. *elatius*) (Butler, 1989; Zohary and Hopf, 2000). Early finds of pulses co-occur with cereal crops and weed assemblages, suggesting that they too were crops, but no measurable change in seed size is noted (Garrard, 2000; Zohary and Hopf, 2000; Hillman *et al.*, 2001; Tanno and Willcox, 2006b; Weiss *et al.*, 2006). Even millennia after the beginnings of cereal cultivation and the establishment of domesticated cereals, when agriculture had dispersed to Europe, finds of peas and lentils (e.g. from Neolithic and




FIG. 13. (A) Late Chalcolithic (1400–900 BC), Iron Age (900–200 BC) and Early Historic (200 BC – 400 AD) *Vigna* finds from South India, indicating size increase. (B) Archaeological *Vigna* measurements from northern India, including Ganges valley Neolithic, Iron Age and Harappan Bronze Age civilization finds, suggesting a correlation between larger size and plough agriculture (data from Fuller and Harvey, 2006).

Early Bronze Age Greece, Bulgaria, Germany and the Netherlands) are little different from their pre-agricultural Near Eastern ancestors in terms of average and minimum sizes, although there is a slight tendency for larger size maxima (e.g. Van Zeist and Bottema, 1971; Bakels, 1978; Hopf, 1973; Housely, 1981; Hansen, 1991). Perhaps the earliest sites with notably large lentils and peas are some Near Eastern pottery Neolithic sites (approx. 6000–5000 BC), such as Erbaa in Turkey and Tepe Sabz in Iran (Helbaek, 1969; Van Zeist and Buitenhuis, 1983), but such sites are the minority, with consistently enlarged pulse seeds encountered only in the Late Bronze Age, Iron Age and Roman periods. Taken together this suggests that there was an even longer delay in serious size increase in Near Eastern pulses than in Indian *Vigna*, something on the order 2000–4000 years, or even longer in some regions. It also raises the possibility that some size increases in particular regions and periods were contingent on local circumstances of selection. It may also be noted that limited data on early East Asian soybeans (*Glycine max*) and adzuki beans (*Vigna angularis*) indicate that as late as the third millennium BC they are still largely within the wild size range, with enlarged seed sizes from the later Bronze Age

or Iron Age (Crawford and Lee, 2003; Crawford *et al.*, 2005), periods in which ploughs were established.

A PULSE-LIKE CEREAL? THE CASE OF AFRICAN PEARL MILLET

A final interesting case is provided by pearl millet (*Pennisetum glaucum*), an important domesticate of West Africa (D'Andrea and Casey, 2002; Fuller, 2003; Zach and Klee, 2003). Archaeobotanical research in Africa remains very limited especially given the size of the continent. Most studies have focused on Egypt, but there is a growing database from across the Sahara and in sub-Saharan West Africa. As outlined by Harlan (1971) there are a wide range of crops with varying wild progenitor distributions in Africa (Fig. 14). In terms of major cereal domestications the northern savannas and the Sahel zone south of the Sahara are important. *Sorghum bicolor* ssp. *bicolor* is likely to have originated from the eastern part of the savannas, such as between Lake Chad and the Western Sudan, or north-west Ethiopia (Harlan and Stemler, 1976). By contrast, pearl millet has been linked to one, or more likely two, domestications in the Sahel zone west of Lake Chad (Tostain, 1992; Fuller, 2003; Neumann, 2003). An important factor in where and when agriculture developed in Africa is climate change. As

wetter conditions of the early (10 000–6200 BC) and mid Holocene (5900–2200 BC) caused a northward shift in vegetation zones and much of the Sahara was savanna-like. This environment supported populations of pottery-making hunter-gatherers, who utilized a range of wild millet-grasses, including wild *Sorghum bicolor* in the east (Egypt's Western Desert), and these groups adopted livestock early, certainly by 6000–5000 BC (Marshall and Hildebrand, 2002; Fuller, 2005). As conditions dried, after 3500 BC, and populations with their herds were forced south, some appear to have taken up cultivation. Evidence for the actual transition to agriculture remains elusive but in the early second millennium BC widely dispersed populations in West Africa were cultivating morphologically domesticated pearl millet (D'Andrea and Casey, 2002). That pearl millet had dispersed to India by approx. 1700 BC (Fuller, 2003) suggests that this process began by the third millennium BC and saw rapid dispersal of cropping across the northern savannas.

Pearl millet domestication is inferred from two sets of evidence. First, there was loss of natural seed shedding, which is linked to the shift from sessile involucre to development of a non-dehiscent peduncle (Poncet *et al.*, 2000). This shift is already evident from ceramic impressions of pearl millet chaff by 1700–1500 BC in Mauretania (Amblard and Pernes, 1989; MacDonald *et al.*, 2003), and




FIG. 14. A synoptic geography of early agricultural developments and precursors in Africa. Shown are the modern distributions of wild *Sorghum bicolor* and *Pennisetum glaucum* with genetic connections to the domesticates (after Harlan, 1971; Tostain, 1992). The previously wetter conditions imply a northward shift in the Sahara–Sahel transition (see Gasse, 2000; Marshall and Hildebrand, 2002). Early Holocene ceramic-using forager sites based on Jesse (2003), and mid-Holocene pastoral sites based on Jousse (2004). Early evidence for wild sorghum gathering is indicated (based on Stemler, 1990; Barakat and Fahmy, 1999; Wasylkova and Dahlberg, 1999). The spread of Near Eastern crops is indicated in the Nile Valley vis-à-vis the pre-ceramic Neolithic distribution in the Eastern Mediterranean. Sites with early pearl millet are numbered: 1, Dhar Tichitt sites (cited in D'Andrea and Casey, 2002); 2, Dhar Oualata sites (Amblard and Pernes, 1989); 3, Djiganyai (MacDonald *et al.*, 2003); 4, Winde Koroji (MacDonald, 1996); 5, Karkarichinkat (cited in D'Andrea and Casey, 2002); 6, Ti-n-Akof (cited in D'Andrea and Casey, 2002); 7, Oursi (cited in D'Andrea and Casey, 2002); 8, Birimi (D'Andrea *et al.*, 2001); 9, Ganjigana (Klee *et al.*, 2004); 10, Kursakata (Zach and Klee, 2003). Historical sites with pearl millet metrical data: 11, Arondo (cited in Zach and Klee, 2003); 12, Jarma (Pelling, 2005); 13, Qasr Ibrim (Steele and Bunting, 1982).

slightly later in Nigeria (Klee *et al.*, 2000, 2004). Pearl millet shows a subtle but clear change in grain shape, becoming apically thicker and more club-shaped than its wild counterpart, i.e. an increased thickness/breadth ratio (Brunken *et al.*, 1977; D'Andrea *et al.*, 2001; Zach and Klee, 2003). However, a major increase in seed size appears delayed (D'Andrea *et al.*, 2001: 346). Figure 15A shows the expected range of domesticated and wild forms based on reported modern populations, adjusted for 10% shrinkage. Figure 15B plots some archaeological populations.

Of note is that early West Africa populations, from the second and first millennia BC, have their averages firmly in the wild size range, although there are long tails of variation that extend into the larger size range (e.g. at Birimi). One of the earliest finds of pearl millet from India comes from Surkotada, Gujarat, approx. 1700 BC, which can be seen to fall with these early domesticated African populations. By contrast, a rather later, Gangetic population from Narhan (where arid tillage is established, and where larger *Vigna* pulses were present, see Fig. 13D) is

markedly larger, suggesting selection for larger-grained pearl millet. In Africa larger grained populations appear in the first millennium AD, represented by finds from Nubia and Libya, as well as Medieval Senegal. However, the continued small-grained populations in Early Historic South India (Nevasa) and apparent reversion in later Medieval Libya suggests that there may be factors that work against gigantism in pearl millet, and in the absence reinforcing selection populations may tend towards the smaller size ranges.

This raises questions about the selection pressures involved in large-grained *Pennisetum*. As both Libya and South India lack wild populations, this cannot be attributed to cross-pollination with wild types. There may be some constraints particular to this crop, as one experiment indicates that optimal germination occurred under higher temperatures that resulted in lower average grain weights (Mohamed *et al.*, 1985). In addition, pearl millet involucres are polymorphic in grain count with the vast majority producing two grains, a large minority with one larger grain, and a further minority producing 3–9 grains, which are




FIG. 15. Scatter plots of pearl millet (*Pennisetum glaucum*) grain width vs. thickness. (A) Modern population averages and minima of domesticated populations reduced by 10% to account for shrinkage, compared with modern wild population averages with maxima, reduced by 10%. Dashed line indicates expected separation between wild and domesticated forms. Sources: Brunken *et al.* (1977) and Zach and Klee (2003). (B) Plots of archaeological site averages and ranges. Early West Africa averages (Birimi, 1700–1500 BCE; Kursakata, 1500–800 BCE) fall in the wild zone although ranges extend into the larger domesticated zone. The earliest finds in India (Surkotada, approx. 1700 BCE) are close to these as are Early Historic (200 BCE–300 CE) Nevasa in southern India. North Indian Narhan (1400–800 BCE) shows a marked shift towards larger sizes comparable with modern domesticates, as does early medieval Qasr Ibrim (Egypt, approx. 450 CE: this find is preserved by dessication and has been reduced to be comparable with carbonized material). Jarma in south-west Libya may show an apparent shift towards somewhat larger grains during the early first millennium CE, comparable with the size found in medieval Senegal at Arundo. Later Medieval Jarma has shifted back towards to near wild size range. Sources: Kajale (1977), Steele and Bunting (1982), Chanchala (1995), Saraswat *et al.* (1994), D'Andrea *et al.* (2001) and Zach and Klee (2003) (Jarma data from Ruth Pelling, personal communication).

necessarily smaller (Godbole, 1925). Thus, selection for higher grain counts, and more reliable germination, might conflict with selection for larger seed sizes. Nevertheless as a working hypothesis, I would propose that, as with pulses, there is a deeper burial threshold that selected for gigantism in pearl millet at some times and in some locations. In that regard it might be noted that the larger grain populations in Libya and Nubia, like that in Gangetic India, are associated with more intensive plough cultures, whereas ards were not present in West Africa and may have declined in post-Garamantean Libya. Thus, we can hypothesize that large-grained varieties evolved under plough systems and then dispersed back to West Africa at a later date. If so, this would imply separate events of grain enlargement in India and north-eastern Africa. While initial cultivation must have selected for non-shattering, and slight changes in grain weight and shape (the club shape), serious gigantism may have required a stronger selection pressure and therefore evolved later: a millennium or more later in India, and two millennia later in Africa.

CONCLUSIONS

Archaeological evidence indicates that the entire domestication syndrome did not suddenly appear when people began to cultivate plants. Rather, different aspects of the syndrome evolved in response to the new ecological conditions of early cultivation. What these data suggest is that in domesticated grasses, changes in grain size and shape ('semi-domestication') evolved prior to non-shattering ears or panicles ('domestication' *sensu stricto*). While initial grain size increases may have evolved during the first centuries of cultivation, within perhaps 500–1000 years, non-shattering was much slower, becoming fixed about 1000–2000 years subsequently. Pulses by contrast do not show evidence for seed size increase in relation to the earliest cultivation, but seed size increase may be delayed by 2000–4000 years. This implies that conditions that were sufficient to select for larger seed-size in Poaceae were not sufficient in Fabaceae. This implies different thresholds of selective pressure in relation to differing seed ontogenetics and underlying genetic architecture in these families. Pearl millet (*Pennisetum glaucum*) may show some similarities to the pulses in terms of a lag-time before truly larger-grained forms evolved. These results may aid in predicting when and where certain crop domestications are likely to have occurred based on counting backwards from the earliest known domestic finds. Thus, for example, we would predict that pearl millet cultivation began by 3200–2700 BC. These results also raise questions about taxonomically linked differences in evolution under the selection forces of cultivation.

Reconsidering sickles and cereal domestication

There has been a tendency to assume that harvesting with a sickle was the selective force that led to domestication, i.e. non-shattering (as discussed above). The archaeological evidence, however, does not support this in any

documented case. In China, as discussed already, rice grains begin to plump and increase in size but domestication is indicated by the shift towards predominantly mature-grained harvests (and inferred non-shattering), during the fifth millennium BC, and by approx. 4000 BC. In this region there are no clear archaeological sickles until after 3500 BC, the Later Songze period (approx. 3500 BC), after which they become widespread in the Liangzhu culture (3300–2200 BC). These sickles may be a cultural borrowing from millet cultivators in central China, where such tools were in use since at least 5000 BC (cf. Chang, 1986). Even in central and northern China, the earliest sickles occur at sites that already have millet cultivation, and earliest documented domestic millets from Xinglonggou (near Chifeng, China), before 6000 BC (Zhao, 2005), come from a culture without sickles. In China, sickles consistently represent a technology development after domesticated plants are fully established.

In the Near East sickles were in use prior to agriculture and must now be argued to be transferred to agriculture relatively late, *after* domestication. Preserved sickles, and more commonly lithic sickle blades, are known from Natufian contexts (13 000–10 500 BC), in a period for which there is no evidence for domesticates, and non-shattering domesticates continued to be absent through the PPNA (through 8800 BC) (see Fig. 3). Microscopic studies of 'sickle gloss' have been used to suggest they were cereal-harvesting (Unger-Hamilton, 1989; Anderson, 1992), but we cannot rule out harvesting of sedges (Cyperaceae) and reeds (*Phragmites*) as materials for basketry or thatching. As suggested by Sauer (1958), the early Natufian sickles were prototype saws, designed for raw material gathering rather than seed collecting. As indicated by the archaeobotanical evidence reviewed above, the rate of evolution of tough rachis einkorn and barley is far too slow to be accounted for by a model of strong selective pressure that would be expected if sickling was carried out regularly, as modelled by Hillman and Davies (1990). Thus, it would appear that early cultivators continued to employ the time-efficient harvesting methods associated with hunter-gatherers. Once cultivated, and populations had noticeably large proportions (majorities) of non-shattering types, then the transfer of the sickle technology to agriculture may have been seen as an obvious enhancement. In evolutionary terms the sickle is thus an 'exaptation' (*sensu* Gould and Vrba, 1982), in that it developed for some other purpose, and was later transferred to crop-harvesting of *already domesticated* crops.

I would propose alternative explanations for the selection of domesticated-type crops that can account for the slow creep towards domestication. As others have noted, the harvesting of cereals when green, i.e. immature, regardless of technique, will not select for domesticated types (Hillman and Davies, 1990; Willcox, 1999). Harvesting green, however, may not provide full returns from a given stand of crops, as additional seeds (including late tillers) may form and approach maturation subsequent to the harvest. For the early farmers, who have invested significant labour into a restricted unit of land, it becomes important

to maximize returns from that unit of land (as noted by Hillman and Davies, 1990: 69; Bar-Yosef, 1998). This may encourage multiple episodes of harvest. Later harvestings, whether by plucking or beating, will encounter domesticated genotypes in a higher frequency than earlier harvests. If, as an aspect of random variation, some farming households choose to store the late harvest as seed for sowing the following year, those fields so sown will start an increase in the domesticated type. Other households, however, may store for sowing their earlier harvests. Therefore, taken at the level of a human community, or on a regional scale, there might be only a very small proportion of sown crop that had some selection for the domesticated type. Such a model might therefore account for significantly longer periods involved in the fixation of non-shattering types in cultivated populations. By contrast, every farmer and every sown population would be under selective pressure to germinate rapidly, leading to seed size increase and loss of germination inhibitors. Similarly, natural selection for dispersal aids such as awns will be uniformly reduced. Thus, we should expect these ‘semi-domestication’ traits to evolve more rapidly.

Domestication as an interdisciplinary study of evolution

Domestication in plants is not one thing, nor has it been one uniform process. While there are recurrent parallels, due to the same selective pressures of cultivation, different domestication traits have evolved at different rates and these have varied markedly across families, such as between cereals and legumes. Further archaeobotanical research will help to pin down the actual rates at which different domesticates evolved, and needs to be expanded to address a larger range of species. The archaeological record also provides insights into what people are doing during this evolutionary process in terms of their technologies and ecological adaptations. Understanding past domestications is an exciting area of interdisciplinary investigation, between archaeologists and plant scientists, which may offer insights relevant to future directions in the evolution of crops under human manipulation.

ACKNOWLEDGEMENTS

These ideas have benefited from discussions with Gordon Hillman, Sue Colledge and David Harris. I must thank Emma Harvey, Ling Qin, Mervyn Jupe, Ruth Pelling and Sue Colledge for assistance in compiling some of the data sets. I thank David Harris, Sue Colledge, Mary Ann Murray and Chris Stevens for their comments on a draft of this paper, and the input of the peer-reviewers. I thank the OECD for financial support to participate in the workshop on ‘Domestication, Super-Domestication and Gigantism’. Funding to pay the Open Access publication charges for this article was provided by the OECD.

LITERATURE CITED

Allaby RG, Banerjee M, Brown TA. 1999. Evolution of the high molecular weight glutenin loci of the A, B, D, and G genomes of wheat. *Genome* 42: 116–128.

- Allard RW. 1988. Genetic changes associated with the evolution of adaptability in cultivated plants and their wild progenitors. *The Journal of Heredity* 79: 225–238.
- Allchin FR, Allchin B. 1997. *Origins of a civilization. The prehistory and early archaeology of South Asia*. New Delhi: Penguin Books India.
- Allen H. 1974. The Bagundji of the Darling Basin: cereal gatherers in an uncertain environment. *World Archaeology* 5: 309–322.
- Amblard S, Pernes J. 1989. The identification of cultivated pearl millet (*Pennisetum*) amongst plant impressions on pottery from Oued Chebbi (Dhar Oualata, Mauritania). *African Archaeological Review* 7: 117–126.
- Anderson P. 1992. Experimental cultivation, harvest and threshing of wild cereals and their relevance for interpreting the use of Epipalaeolithic and Neolithic artefacts. In: Anderson P ed. *Prehistoire de L'agriculture: Nouvelles approches experimentales et ethnographiques*. Paris: CNRS, 179–209.
- Bakels C. 1978. *Four Linearbandkeramik settlements and their environments: a palynological study of Sittard, Stein, Esloo and Heinheim*. PhD dissertation, University of Leiden, Leiden.
- Barakat H, Fahmy AD. 1999. Wild grasses as ‘Neolithic’ food resources in the eastern Sahara: a review of evidence from Egypt. In: van der Veen M ed. *The exploitation of plant resources in Ancient Africa*. New York: Kluwer/Plenum, 33–46.
- Bar-Yosef O. 1998. The Natufian culture in the Levant. *Evolutionary Anthropology* 6: 159–177.
- Bar-Yosef O. 2003. The Natufian culture and the early Neolithic: social and economic trends in southwestern Asia. In: Bellwood P, Renfrew C eds. *Examining the farming/language dispersal hypothesis*. Cambridge: McDonald Institute for Archaeological Research, 113–126.
- Barlow KR, Heck M. 2002. More on acorn eating during the Natufian: expected patterning in diet and the archaeological record of subsistence. In: Mason SLR, Hather J eds. *Hunter-gatherer archaeobotany. perspectives from the northern temperate zone*. London: Institute of Archaeology, UCL, 128–145.
- Baskin CC, Baskin JM. 2001. *Seeds. Ecology, biogeography, and evolution of dormancy and germination*. New York: Academic Press.
- Blumler MA. 1991. Modelling the origins of legume domestication and cultivation. *Economic Botany* 45: 243–250.
- Blunier T, Chappellaz J, Schwander J, Stauffer B, Raynaud D. 1995. Variations in atmospheric methane concentration during the Holocene epoch. *Nature* 374: 46–49.
- Braadbaart F, van Bergen PF. 2005. Digital imaging analysis of size and shape of wheat and pea upon heating under anoxic conditions as a function of temperature. *Vegetation History and Archaeobotany* 14: 67–75.
- Brown TA. 1999. How ancient DNA may help in understanding the origin and spread of agriculture. *Philosophical Transactions of the Royal Society B: Biological Sciences* 354: 89–98.
- Brunken J, De Wet JM, Harlan JR. 1977. The morphology and domestication of pearl millet. *Economic Botany* 31: 163–174.
- Bruno M, Whitehead WT. 2003. *Chenopodium* cultivation and formative period agriculture at Ciripa, Bolivia. *Latin American Antiquity* 14: 339–355.
- Butler A. 1989. Cryptic anatomical characters as evidence of early cultivation in the grain legumes (pulses). In: Harris DR, Hillman GC eds. *Foraging and farming*. London: Unwin and Hyman, 390–407.
- Byrd BF. 2005. Reassessing the emergence of village life in the Near East. *Journal of Archaeological Research* 13: 231–290.
- Chanchala S. 1995. Harappan plant economy of Kutch, Gujarat. *Geophytology* 23: 227–233.
- Chang KC. 1986. *The archaeology of Ancient China*, 4th edn. New Haven: Yale University Press.
- Chen W-B, Nakamura I, Sato Y-I, Nakai H. 1993. Indica and Japonica differentiation in Chinese landraces. *Euphytica* 75: 195–201.
- Cheng C, Motohashi R, Tchuchimoto S, Fukuta Y, Ohtsubo H, Ohtsubo E. 2003. Polyphyletic origin of cultivated rice: based on the interspersed patterns of SINES. *Molecular Biology and Evolution* 20: 67–75.
- Colledge S. 1998. Identifying pre-domestication cultivation using multivariate analysis. In: Damania AB, Valkoun J, Willcox G, Qualset CO eds. *The origins of agriculture and crop domestication*. Aleppo, Syria: ICARDA, 121–131.

- Colledge S. 2001.** *Plant exploitation on epipalaeolithic and early neolithic sites in the Levant*. Oxford: British Archaeological Reports.
- Colledge S. 2002.** Identifying pre-domestication cultivation in the archaeological record using multivariate analysis: presenting the case for quantification. In: Cappers RTJ, Bottema S eds. *The dawn of farming in the Near East*. Berlin: ex Oriente, 141–151.
- Colledge S. 2004.** Reappraisal of the archaeobotanical evidence for the emergence and dispersal of the ‘founder crops’. In: Peltenberg E, Wasse A, eds. *Neolithic revolution. new perspectives on southwest Asia in light of recent discoveries on Cyprus*. Oxford: Oxbow Books, 49–60.
- Colledge S, Conolly J, Shennan S. 2004.** Archaeobotanical evidence for the spread of farming in the eastern Mediterranean. *Current Anthropology* **45**: S35–S58.
- Crawford G. 2005.** East Asian plant domestication. In: Stark MT, ed. *Archaeology of Asia*. Oxford: Blackwell Publishing, 77–95.
- Crawford G, Lee G-A. 2003.** Agricultural origins in the Korean Peninsula. *Antiquity* **77**: 87–95.
- Crawford G, Underhill A, Zhao Z, Lee G-A, Feinman G, Nicholas L et al. 2005.** Late Neolithic plant remains from northern China: preliminary results from Liangchengzhen, Shandong. *Current Anthropology* **46**: 309–317.
- D’Andrea AC, Casey J. 2002.** Pearl millet and Kintampo subsistence. *African Archaeological Review* **19**: 147–173.
- D’Andrea AC, Klee M, Casey J. 2001.** Archaeobotanical evidence for pearl millet (*Pennisetum glaucum*) in sub-Saharan West Africa. *Antiquity* **75**: 341–348.
- De Moulins D. 1997.** *Agricultural changes at Euphrates and steppe sites in the mid-8th to the 6th millennium B.C.* Oxford: British Archaeological Reports.
- Fairbairn A, Asouti E, Near J, Martinoli D. 2002.** Macro-botanical evidence for plant use at Neolithic Catalhoyuk, southcentral Anatolia, Turkey. *Vegetation History and Archaeobotany* **11**: 41–54.
- Ford R. 1985.** The process of plant food production in prehistoric North America. In: Ford R, ed. *Prehistoric food production in North America*. Ann Arbor: University of Michigan Museum of Anthropology, 1–18.
- Fuller DQ. 2002.** Fifty years of archaeobotanical studies in India: laying a solid foundation. In: Settar S, Korisettar R eds. *Indian Archaeology in retrospect, Volume III. Archaeology and interactive disciplines*. Delhi: Manohar, 247–363.
- Fuller DQ. 2003.** African crops in prehistoric South Asia: a critical review. In: Neumann K, Butler A, Kahlheber S eds. *Food, fuel and fields. Progress in African archaeobotany*. Köln: Heinrich-Barth Institut, 239–271.
- Fuller DQ. 2005.** Farming: Stone Age farmers of the savanna. In: Shillington K, ed. *Encyclopedia of African history*. New York: Fitzroy Dearborn, 521–522.
- Fuller DQ. 2006.** Agricultural origins and frontiers in South Asia: a working synthesis. *Journal of World Prehistory* **20**: 1–86.
- Fuller DQ, Harvey E. 2006.** The archaeobotany of Indian pulses: identification, processing and evidence for cultivation. *Environmental Archaeology* **11**: 219–246.
- Fuller DQ, Korisettar R. 2004.** The Vegetational context of early agriculture in South India. *Man and Environment* **29**: 7–27.
- Fuller DQ, Korisettar R, Venkatasubbaiah PC. 2001.** Southern Neolithic cultivation systems: a reconstruction based on archaeobotanical evidence. *South Asian Studies* **17**: 171–187.
- Fuller DQ, Korisettar R, Venkatasubbaiah PC, Jones MK. 2004.** Early plant domestications in southern India: some preliminary archaeobotanical results. *Vegetation History and Archaeobotany* **13**: 115–129.
- Fuller DQ, Harvey E, Qin L. 2007.** Presumed domestication? Evidence for Wild rice cultivation in the 5th Millennium BC of the Lower Yangtze region. *Antiquity* **81** (in press).
- Garrard A. 2000.** Charting the emergence of cereal and pulse domestication in South-West Asia. *Environmental Archaeology* **4**: 67–86.
- Gasse F. 2000.** Hydrological changes in the African tropics since the last glacial maximum. *Quaternary Science Review* **19**: 189–211.
- Gepts P. 2004.** Crop domestication as a long-term selection experiment. *Plant Breeding Reviews* **24**: 1–44.
- Godbole SV. 1925.** *Pennisetum typhoidum*. Studies on the Bajra Crop; I: The morphology of *Pennisetum typhoidum*. *Memoirs of the Department of Agriculture in India* (Agricultural Research Institute, Pusa) **14**: 247–268.
- Gould SJ, Vrba ES. 1982.** Exaptation—A missing term in the science of form. *Paleobiology* **8**: 4–15.
- Gu J. 2003.** The rice paddy land from Chuodun Site of Majiabang Period. *Dongnan Wenhua [southeast Culture]* **2003** (special issue 1): 42–45 [in Chinese].
- Hansen J. 1991.** *The palaeoethnobotany of Franchthi Cave*. Bloomington: Indiana University Press.
- Harlan JR. 1967.** A wild wheat harvest in Turkey. *Archaeology* **20**: 197–201.
- Harlan JR. 1971.** Agricultural origins: centers and noncenters. *Science* **174**: 468–474.
- Harlan JR. 1989.** Wild grass-seed harvesting in the Sahara and sub-Saharan of Africa. In: Harris DR, Hillman GC eds. *Foraging and farming: the exploitation of plant resources*. London: Unwin and Hyman, 79–98.
- Harlan JR. 1992.** *Crops and ancient man*. Madison: American Society for Agronomy.
- Harlan JR. 1995.** *The living fields*. Cambridge: Cambridge University Press.
- Harlan JR, Stemler AB. 1976.** The races of Sorghum in Africa. In: Harlan JR, De Wet MJM, Stemler AB, eds. *The origins of African plant domestication*. The Hague: Mouton, 465–478.
- Harlan JR, De Wet MJM, Price EG. 1973.** Comparative evolution of cereals. *Evolution* **27**: 311–325.
- Harris DR. 1984.** Ethnohistorical evidence for the exploitation of wild grasses and forbes: its scope and archaeological implications. In: Van Zeist W, Casparie WA, eds. *Plants and ancient man. Studies in paleoethnobotany*. Rotterdam: A. A. Balkema, 63–69.
- Harris DR. 1989.** An evolutionary continuum of people-plant interaction. In: Harris DR, Hillman GC eds. *Foraging and farming: the evolution of plant exploitation*. London: Routledge, 11–26.
- Harris DR. 1996.** Introduction: themes and concepts in the study of early agriculture. In: Harris DR, ed. *The origins and spread of agriculture and pastoralism in Eurasia*. London: UCL Press, 1–9.
- Harris DR. 1998.** The origins of agriculture in southwest Asia. *The Review of Archaeology* **19**: 5–11.
- Harvey EL. 2006.** *Early agricultural communities in Northern and Eastern India: an archaeobotanical investigation*. Unpublished PhD dissertation, University College London.
- Hather J. 1996.** The origins of tropical vegetation: Zingiberaceae, Araceae and Dioscoreaceae in Southeast Asia. In: Harris DR ed. *The Origins and spread of agriculture and pastoralism in Eurasia*. London: UCL Press, 538–550.
- Hawkes JG. 1983.** *The diversity of crop plants*. Cambridge, MA: Harvard University Press.
- Heizer RF, Elsasser AB. 1980.** *The natural world of the California Indians*. Berkeley: University of California Press.
- Helbaek H. 1969.** Plant collecting, dry-farming and irrigation agriculture in prehistoric Deh Luran. In: Hole F, Flannery KV, Neely JA eds. *Prehistory and human ecology of the Deh Luran Plain*. Ann Arbor: Museum of Anthropology, University of Michigan, 383–426.
- Henan Provincial Institute of Archaeology 1999.** *Wuyang jiahu (Jiahu Site in Wuyang)*. Beijing: Science Press [in Chinese].
- Hillman GC. 1984.** Interpretation of archaeological plant remains: the application of ethnographic models from Turkey. In: Van Zeist W, Casparie WA, eds. *Plants and ancient man. Studies in paleoethnobotany*. Rotterdam: A. A. Balkema, 1–41.
- Hillman GC. 2000.** Abu Hureyra 1: The Epipalaeolithic. In: Moore AMT, Hillman GC, Legge AJ, eds. *Village on the Euphrates. From foraging to farming at Abu Hureyra*. New York: Oxford University Press, 327–398.
- Hillman G, Davies MS. 1990.** Domestication rates in wild wheats and barley under primitive cultivation. *Biological Journal of the Linnean Society* **39**: 39–78.
- Hillman GC, Hedges R, Moore AMT, Colledge S, Pettitt P. 2001.** New evidence of Late Glacial cereal cultivation at Abu Hureyra on the Euphrates. *The Holocene* **11**: 383–393.
- Hopf M. 1973.** Frühe Kulturpflanzen aus Bulgarien. *Jahrb. Rom.-German. Zentralmus, Mainz* **20**: 1–47.
- Hoshikawa K. 1993.** Anthesis, fertilization and development of caryopsis. In: Matsuo T, Hoshikawa K, eds. *Science of the rice plant, Vol. 1. Morphology*. Tokyo: Food and Agriculture Policy Center, 339–376.

- Housely RA. 1981.** *The palaeoethnobotany of Servia: a Neolithic and Early Bronze Age site in western Macedonia, Greece.* MSc dissertation, Department of Archaeology, University of Southampton.
- Huang F, Zhang M. 2000.** Pollen and phytolith evidence for rice cultivation during the Neolithic at Longqiuzhuang, eastern Jainghuai, China. *Vegetation History and Archaeobotany* **9**: 161–168.
- Jesse F. 2003.** Early ceramics in the Sahara and the Nile Valley. In: Krzyzanski L, Kroeper K, Kobusiewicz M, eds. *Cultural markers in the later prehistory of northeastern Africa and recent Research.* Poznan: Poznan Archaeological Museum, 35–50.
- Jones MK. 1985.** Archaeobotany beyond subsistence reconstruction. In: Barker G, Gamble C eds. *Beyond domestication in Prehistoric Europe, investigations in subsistence archaeology and social complexity.* London: Academic Press, 107–128.
- Jones MK, Brown TA. 2000.** Agricultural origins: the evidence of modern and ancient DNA. *The Holocene* **10**: 775–782.
- Jones G, Valamoti S, Charles M. 2000.** Early Crop diversity: a ‘new’ glume wheat from northern Greece. *Vegetation History and Archaeobotany* **9**: 133–146.
- Jousse H. 2004.** A new contribution to the history of pastoralism in West Africa. *Journal of African Archaeology* **2**: 187–201.
- Kajale MD. 1977.** Ancient plant economy at Nevasa during Satavahana and Indo-Roman period. *Bulletin of the Deccan College Post-Graduate and Research Institute* **36**: 48–61.
- Kelly RL. 1995.** *The foraging spectrum: diversity in hunter-gatherer life-ways.* Washington, DC: Smithsonian Institution Press.
- Kislev ME. 1997.** Early agriculture and paleoecology of Netiv Hagdud. In: Bar-Yosef O, Gopher O eds. *An early Neolithic village in the Jordan Valley, Part I: The archaeology of Netiv Hagdud.* Cambridge, Mass.: Peabody Museum of Archaeology and Ethnology, Harvard University, 209–236.
- Kislev ME, Nadel D, Carmi I. 1992.** Epipalaeolithic (19,000 bp) cereal and fruit diet at Ohalu II, Sea of Galilee, Israel. *Review of Palaeobotany and Palynology* **73**: 161–166.
- Kislev ME, Weiss E, Hartmann A. 2004.** Impetus for sowing and the beginning of agriculture: ground collecting of wild cereals. *Proceedings of the National Academy of Sciences of the USA* **101**: 2692–2695.
- Kislev ME, Hartmann A, Bar-Yosef O. 2006.** Early domesticated fig in the Jordan Valley. *Science* **312**: 1372–1374.
- Klee M, Zach B, Neumann K. 2000.** Four thousand years of plant exploitation in the Chad Basin of northeast Nigeria I: The archaeobotany of Kursakata. *Vegetation History and Archaeobotany* **9**: 223–237.
- Klee M, Zach B, Stika H-P. 2004.** Four thousand years of plant exploitation in the Lake Chad Basin (Nigeria), part III: plant impressions in potsherds from the Final Stone Age Gajiganna Culture. *Vegetation History and Archaeobotany* **13**: 131–142.
- Köhler-Schneider M. 2003.** Contents of a storage pit from late Bronze Age Stillfried, Austria: another record of the ‘new’ glume wheat. *Vegetation History and Archaeobotany* **12**: 105–111.
- Krishnasamy V, Seshu DV. 1989.** Seed germination rate and associated characters in rice. *Crop Science* **29**: 904–908.
- Kreuz A, Boenke N. 2002.** The presence of two-grained einkorn at the time of the Bandkeramik culture. *Vegetation History and Archaeobotany* **11**: 233–240.
- Küster H. 2000.** Rye. In: Kiple KF, Ornelas KC eds. *Cambridge world history of food.* Cambridge: Cambridge University Press, 149–152.
- Ladzinsky G. 1987.** Pulse domestication before cultivation. *Economic Botany* **41**: 60–65.
- Ladzinsky G. 1993.** Lentil domestication: on the quality of evidence and arguments. *Economic Botany* **47**: 60–64.
- Londo JP, Chiang Y-C, Hung K-H, Chiang T-Y, Schaal BA. 2006.** Phylogeography of Asian wild rice, *Oryza rufipogon*, reveals multiple independent domestications of cultivated rice, *Oryza sativa*. *Proceedings of the National Academy of Sciences of the USA* **103**: 9578–9583.
- Lone FA, Khan M, Buth GM. 1993.** *Palaeoethnobotany – plants and ancient man in Kashmir.* Rotterdam: A. A. Balkema.
- Lu TLD. 1998.** Some botanical characteristics of green foxtail (*Setaria viridis*) and harvesting experiments on the grass. *Antiquity* **72**: 902–907.
- Lu TLD. 1999.** *The transition from foraging to farming and the origin of agriculture in China.* Oxford: BAR.
- MacDonald KC. 1996.** The Windè Koroji Complex: evidence for the peopling of the eastern Inland Niger Delta (2100–500 BC). *Prehistoire Anthropologie Méditerranéennes* **5**: 147–165.
- MacDonald KC, Vernet R, Fuller DQ, Woodhouse J. 2003.** New light on the Tichitt Tradition: a preliminary report on survey and excavation at Dhar Nema. In: Mitchell P, Haour A, Hobart J, eds. *Researching Africa’s past. New contributions from British archaeologists.* Oxford: Oxford University School of Archaeology, 73–80.
- Madella M, Fuller DQ. 2006.** Paleoecology and the Harappan Civilisation of South Asia: a reconsideration. *Quaternary Science Reviews* **25**: 1283–1301.
- Maranon T, Grubb PJ. 1993.** Physiological basis and ecological significance of the seed size and relative growth rate relationship in Mediterranean annuals. *Functional Ecology* **7**: 591–599.
- Marshall F, Hildebrand E. 2002.** Cattle before crops: the beginnings of food production in Africa. *Journal of World Prehistory* **16**: 99–143.
- McCorriston J. 2000.** Barley. In: Kiple KF, Ornelas KC eds. *Cambridge world history of food.* Cambridge: Cambridge University Press, 81–89.
- Miller LJ. 2003.** Secondary products and urbanism in South Asia: the evidence for traction at Harappa. In: Weber SA, Belcher WR eds. *Indus ethnobiology. New perspectives from the field.* Lanham: Lexington Books, 251–326.
- Mohamed HA, Clark JA, Ong CK. 1985.** The influence of temperature during seed development on the germination characteristics of millet seeds. *Plant, Cell and Environment* **8**: 361–362.
- Morrell PL, Toleno DM, Lundy KE, Clegg MT. 2005.** Low levels of linkage disequilibrium in wild barley (*Hordeum vulgare* ssp. *spontaneum*) despite high rates of self-fertilization. *Proceedings of the National Academy of Sciences of the USA* **102**: 2442–2447.
- Nesbitt M. 2006.** *Identification guide for near eastern grass seeds.* London: Institute of Archaeology, University College London.
- Neumann K. 2003.** The late emergence of agriculture in sub-Saharan Africa: archaeobotanical evidence and ecological considerations. In: Neumann K, Butler A, Kahlheber S, eds. *Food, fuel and fields. Progress in African archaeobotany.* Köln: Heinrich-Barth-Institute, 71–92.
- Oka H-I. 1988.** *Origins of cultivated rice.* Amsterdam: Elsevier.
- Oka H-I, Morishima H. 1967.** Variations in the breeding systems of a wild rice, *Oryza perennis*. *Evolution* **21**: 249–258.
- Oka H-I, Morishima H. 1971.** The dynamics of plant domestication: cultivation experiments with *Oryza perennis* and its hybrid with *O. sativa*. *Evolution* **25**: 356–364.
- Pearsall D. 2000.** *Paleoethnobotany. A handbook of procedures,* 2nd edn. New York: Academic Press.
- Pei A. 1998.** Notes on new advancements and revelations in the agricultural archaeology of early rice domestication in the Dongting Lake region. *Antiquity* **72**: 878–885.
- Pelling R. 2005.** Garamantean agriculture and its significance in a wider North Africa context: the evidence of plant remains from the Fazzan Project. *The Journal of North African Studies* **10**: 397–411.
- Plitman U, Kislev ME. 1989.** Reproductive changes induced by domestication. In: Stirton CH, Zarucchi JL, eds. *Advances in legume biology.* St. Louis: Missouri Botanical Garden, 487–503.
- Poncet V, Lamy F, Devos K, Gale M, Sarr A, Robert T. 2000.** Genetic control of domestication traits in pearl millet (*Pennisetum glaucum* L., Poaceae). *Theoretical and Applied Genetics* **100**: 147–159.
- Reimer PJ, Baillie MGL, Bard E, Bayliss A, Beck JW, Bertrand C, et al. 2004.** IntCal04 terrestrial radiocarbon age calibration, 0–26 Cal Kyr BP. *Radiocarbon* **46**: 1029–1058.
- Saraswat KS, Sharma NK, Saini DC. 1994.** Plant economy at Ancient Narhan (Ca. 1,300 B.C. – 300/400 A.D.). In: Singh P, ed. *Excavations at Narhan (1984–1989).* Varanasi: Banaras Hindu University, 255–346.
- Sato Y-I. 2002.** Origin of rice cultivation in the Yangtze River Basin. In: Yasuda Y, ed. *The origins of pottery and agriculture.* New Delhi: Lustre Press and Roli Books, 143–150.
- Sauer CO. 1958.** Jericho and composite sickles. *Antiquity* **32**: 187–189.
- Shinde V. 1987.** Farming in the Chalcolithic Deccan, India c. 2000–1000 B.C. *Tools and Tillage* **5**: 215–227.
- Smartt J. 1990.** *Grain legumes.* Cambridge: Cambridge University Press.
- Smith BD. 1989.** The origins of agriculture in eastern North America. *Science* **246**: 1566–1571.

- Steele WM, Bunting AH. 1982. Plant remains from the mortar 192.2.10 from Qasr Ibrim. In: Mills AJ, ed. *The cemeteries of Qasr Ibrim*. London: Egypt Exploration Society, 86–87.
- Stemler AB. 1990. A scanning electron microscopic analysis of plant impressions in pottery from sites of Kadero, El Zakiab, Um Direiwa and El Kadada. *Archéologie du Nil Moyen* 4: 87–106.
- Takahashi R, Hosoya LA. 2002. Nut exploitation in Jomon society. In: Mason SLR, Hather J, eds. *Hunter-gatherer archaeobotany. Perspectives from the northern temperate zone*. London: Institute of Archaeology, UCL, 146–155.
- Tomooka N, Vaughan DA, Moss H, Mixed N. 2003. *The Asian Vigna: genus Vigna subgenus Ceratotropis genetic resources*. New York: Kluwer.
- Tang L. 2003. The primitive rice remains from Chuodun Site. *Dongnan Wenhua [southeast Culture]* 2003 (special issue 1): 46–49.
- Tanno K-I, Willcox G. 2006a. How fast was wild wheat domesticated? *Science* 311: 1886.
- Tanno K-I, Willcox G. 2006b. The origins of cultivation of *Cicer arietinum* L. and *Vicia faba* L.: early finds from Tell el-Kerkh, north-west Syria, late 10th millennium B.P. *Vegetation History and Archaeobotany* 15: 197–204.
- Tao J, Chen M-T, Xu S. 2006. A Holocene environmental record from the southern Yangtze River delta, eastern China. *Palaeogeography, Palaeoclimatology, Palaeoecology* 230: 204–229.
- Tostain S. 1992. Enzyme diversity in pearl millet (*Pennisetum glaucum* L.) 3. Wild millet. *Theoretical and Applied Genetics* 83: 733–742.
- Unger-Hamilton R. 1989. The epi-Palaeolithic southern Levant and the origins of cultivation. *Current Anthropology* 30: 88–103.
- Van Zeist WA. 1999. Evidence for agricultural change in the Balikh basin, northern Syria. In: Gosden C, Hather J, eds. *The prehistory of food. Appetites for change*. London: Routledge, 350–373.
- Van Zeist W, Bakker-Heeres JH. 1985. Archaeobotanical studies in the Levant 1. Neolithic sites in the Damascus Basin: Aswad, Ghoraife, Ramad. *Palaeohistoria* 24: 165–256.
- Van Zeist W, Bakker-Heeres JH. 1986. Archaeobotanical studies in the Levant 3. Late Palaeolithic Mureybet. *Palaeohistoria* 26: 171–199.
- Van Zeist W, Bottema S. 1971. Plant husbandry at Neolithic Nea Nikomedeia, Greece. *Acta Botanica Neerlandica* 20: 524–538.
- Van Zeist W, Buitenhuis H. 1983. A palaeobotanical study of Neolithic Erbaba, Turkey. *Anatolica* 10: 47–89.
- Vaughan DA. 1994. *The wild relatives of rice: a genetic resources handbook*. Los Banos, Philippines: International Rice Research Institute.
- Wang C, Ding J. 2000. The plant opal analysis of Shaoqingshan Site, Kunshan City, Jiangsu Province. *Kaogu (Archaeology)* 2000: 87–93 [in Chinese].
- Wang L, Sarntheim M, Erlenkeuser H, Grootes PM, Grimalt JO, Peljero C, Linck G. 1999. Holocene variations in Asian monsoon moisture: a bidecadal sediment record from the South China Sea. *Geophysical Research Letters* 26: 2889–2892.
- Wasylikowa K, Dahlberg J. 1999. Sorghum in the economy of the Early Neolithic Nomadic Tribes at Nabta Playa, Southern Egypt. In: van der Veen M, ed. *The exploitation of plant resources in ancient Africa*. New York: Kluwer/Plenum, 11–32.
- Weiss E, Wetterstrom W, Nadel D, Bar-Yosef O. 2004. The broad spectrum revisited: evidence from plant remains. *Proceedings of the National Academy of Sciences of the USA* 101: 9551–9555.
- Weiss E, Kislev ME, Hartmann A. 2006. Autonomous cultivation before domestication. *Science* 312: 1608–1610.
- Westoby M, Leishman M, Lord J. 1996. Comparative ecology of seed size and dispersal. *Philosophical Transactions of the Royal Society B: Biological Sciences* 351: 1309–1318.
- Wilke PJ, Bettinger R, King TF, O'Connell JF. 1972. Harvest selection and domestication in seed plants. *Antiquity* 46: 203–209.
- Willcox G. 1992. Archaeobotanical significance of growing Near Eastern progenitors of domestic plants at Jalès (France). In: Anderson PC, ed. *Prehistoire de L'Agriculture*. Paris: CNRS, 159–177.
- Willcox G. 1999. Agrarian change and the beginnings of cultivation in the Near East: evidence from wild progenitors, experimental cultivation and archaeobotanical data. In: Gosden C, Hather J, eds. *The prehistory of food. Appetites for change*. London: Routledge, 478–500.
- Willcox G. 2002. Charred plant remains from a 10th millennium B.P. kitchen at Jerf el Ahmar (Syria). *Vegetation History and Archaeobotany* 11: 55–60.
- Willcox G. 2004. Measuring grain size and identifying Near Eastern cereal domestication: evidence from the Euphrates valley. *Journal of Archaeological Science* 31: 145–150.
- Willcox G. 2005. The distribution, natural habitats and availability of wild cereals in relation to their domestication in the Near East: multiple events, multiple centres. *Vegetation History and Archaeobotany* 14: 534–541.
- Willcox G. 2006. Magzalia 1979. <http://perso.orange.fr/g.willcox/magzalia-seeds.htm> (accessed 21 September, 2006).
- Yi S, Saito Y, Zhao Q, Wang P. 2003. Vegetation and climate changes in the Changjiang (Yangtze River) Delta, China, during the past 13,000 years inferred from pollen records. *Quaternary Science Reviews* 22: 1501–1519.
- Yu SY, Zhu C, Song J, Qu WZ. 2000. Role of climate in the rise and fall of Neolithic cultures on the Yangtze delta. *Boreas* 29: 157–165.
- Zach B, Klee M. 2003. Four thousand years of plant exploitation in the Chad Basin of NE Nigeria II: discussion on the morphology of carpopses of domesticated *Pennisetum* and complete catalogue of the fruits and seeds of Kursakata. *Vegetation History and Archaeobotany* 12: 187–204.
- Zhang W. 2002. The bi-peak tubercle of rice, the character of ancient rice and the origin of cultivated rice. In: Yasuda Y, ed. *The origins of pottery and agriculture*. New Delhi: Lustre Press and Roli Books, 205–216.
- Zhao Z. 2005. Palaeoethnobotany and its new achievements in China. *Kaogu [Archaeology]* 2005: 42–49 [in Chinese].
- Zhejiang Provincial Institute of Cultural Relics and Archaeology. 2003. *Hemudu: an excavation report of the Neolithic site*. Beijing: Cultural Relics Publishing House [in Chinese].
- Zhejiang Provincial Institute of Cultural Relics and Archaeology. 2004. *Kuahuqiao site report*. Beijing: Cultural Relics Publishing House [in Chinese].
- Zheng Y, You X, Xu J, Bian Q, Yu W. 1994. Rice phytolith analysis of Hemudu Site. *Journal of Zhejiang Agriculture University* 1994: 81–85 [in Chinese].
- Zheng Y, Dong Y, Matsui A, Udatu T, Fujiwara H. 2003. Molecular genetic basis of determining subspecies of ancient rice using shape of phytoliths. *Journal of Archaeological Science* 30: 1215–1221.
- Zheng Y, Matsui A, Fujiwara H. 2004a. Phytoliths of rice detected in the Neolithic sites in the Valley of the Taihu Lake in China. *Environmental Archaeology* 8: 177–183.
- Zheng Y, Jaing L-P, Zheng J-M. 2004b. Study on the remains of ancient rice from Kuahuqiao Aite in Zhejiang Province. *Chinese Journal of Rice Science* 18: 119–124 [in Chinese].
- Zohary D. 1969. The progenitors of wheat and barley in relation to domestication and agriculture dispersal in the Old World. In: Ucko PJ, Dimbleby GW, eds. *The domestication and exploitation of animals and plants*. London: Duckworth, 47–66.
- Zohary D. 1989. Pulse domestication and cereal domestication: how different are they? *Economic Botany* 43: 31–34.
- Zohary D, Hopf M. 2000. *Domestication of plants in the Old World*, 3rd edn. Oxford: Oxford University Press.
- Zohary D, Spiegel-Roy P. 1975. Beginnings of fruit growing in the Old World. *Science* 187: 319–327.
- Zou H, Gu J, Li M, Tang L, Ding J, Yao Q. 2000. Findings of paddies of Majiabang Culture at Caoxieshan, Jiangsu Province. In: Yan W, Yasuda Y, eds. *The origins of rice agriculture, pottery and cities*. Beijing: Cultural Relics Publishing House, 97–114 [in Chinese].