

1 **Title: Quantification of population sizes of large herbivores and their**
2 **long-term functional role in ecosystems using dung fungal spores**

3 Running title: Fossil dung spores and large herbivore numbers

4 Author list and addresses: Baker, Ambroise G. ^{1,2}, Cornelissen, Perry ³, Bhagwat, Shonil A. ^{4,5},

5 Vera, Fransciscus W. M. ⁶, Willis, Katherine J. ^{1,7,8}

6 ¹ Oxford Long-term Ecology Laboratory, Biodiversity Institute, Department of Zoology,

7 University of Oxford, United Kingdom

8 ² Department of Geography, University College London, London, United Kingdom.

9 ³ State Forestry Service, Driebergen, the Netherlands.

10 ⁴ Department of Geography, The Open University, United Kingdom

11 ⁵ School of Geography and the Environment, University of Oxford, United Kingdom

12 ⁶ Bureau Nature Development, University of Groningen, Wijk bij Duurstede, The Netherlands.

13 ⁷ Department of Biology, University of Bergen, Norway

14 ⁸ Royal Botanic Gardens, Kew, UK

15 Corresponding author:

16 Ambroise Baker, Department of Geography, University College London, Pearson Building,

17 Gower Street, London, WC1E 6BT +44 (0)20 7679 5439 ambroise.baker@ucl.ac.uk

18

19 Abstract:

- 20 1. The relationship between large herbivore numbers and landscape cover over time is
21 poorly understood. There are two schools of thought: one views large herbivores as
22 relatively passive elements upon the landscape and the other as ecosystem engineers
23 driving vegetation succession. The latter relationship has been used as an argument to
24 support reintroductions of large herbivores onto many landscapes in order to increase
25 vegetation heterogeneity and biodiversity through local-scale disturbance regimes.
26 Most of the research examining the relationship between large herbivores and their
27 impact on landscapes has used extant studies. An alternative approach is to estimate
28 the impact of variations in herbivore populations through time using fossil dung fungal
29 spores and pollen in sedimentary sequences. However, to date there has been little
30 quantification of fossil dung fungal spore records and their relationship to herbivore
31 numbers, leaving this method open to varied interpretations.
- 32 2. In this study we developed further the dung fungal spore method and determined the
33 relationship between spore abundance in sediments (number $\text{cm}^{-2} \text{ year}^{-1}$) and
34 herbivore biomass densities (kg ha^{-1}). To establish this relationship, we used the
35 following: i) the abundance of *Sporormiella* spp., *Sordaria* spp. and *Podospora* spp.
36 spores in modern sediments from ponds ii) weekly counts of contemporary wildlife
37 over a period of five years from the rewilded site, Oostvaardersplassen, in the
38 Netherlands.
- 39 3. Results from this study demonstrate that there is a highly significant relationship
40 between spore abundance and local biomass densities of herbivores that can be used
41 in the calibration of fossil records. Mammal biomass density (comprising Konik horses,
42 Heck cattle and red deer) predicts in a highly significant way the abundance of all dung
43 fungal spores amalgamated together. This relationship is apparent at a very local scale
44 (<10m), when the characteristics of the sampled ponds are taken into account (surface

45 area of pond, length of shoreline). In addition, we identify that dung fungal spores are
46 principally transported into ponds by surface run-off from the shores.

47 4. These results indicate that this method provides a robust quantitative measure of
48 herbivore population size over time.

49 Key Words: Ecosystem function, ecosystem engineers, grazing ecology, herbivory, land
50 management, large mammals, Oostvaardersplassen, palaeoecology, rewilding, Sporormiella

51

52 **Introduction**

53 Large herbivores and other vertebrates that live at high trophic levels play an important role in
54 shaping vegetation cover and community composition across most landscapes (Nuttall *et al.*
55 2011; Estes *et al.* 2011; Ripple & Beschta 2012; Tanentzap & Coomes 2012; Ritchie *et al.* 2012;
56 Peh & Lewis 2012; Cromsigt & te Beest 2014). However, what is still widely debated is when
57 and where top-down processes (by which large herbivore populations modify ecosystems)
58 override bottom-up processes (by which climate and soil productivity determine ecosystem
59 structures and composition, including herbivore population size). These processes and their
60 interaction with population dynamics are highly relevant to over 25% of land on earth that is
61 intentionally managed as grazing systems for food production (Asner *et al.* 2004; Steinfeld,
62 Gerber & Wassenaar 2006; Ellis *et al.* 2010) and to over 50% of land, covering semi-natural
63 and wild ecosystems (Ellis *et al.* 2010), where large grazers and browsers are present in
64 abundance. Therefore, quantifying the effects of wild and domesticated large herbivore
65 pressure on ecosystems is extremely important if we are to gain understanding of future land
66 cover changes and their impact on biodiversity and ecosystem services (Carpenter *et al.* 2009).
67 Increasingly conservation policy is looking to introduce large herbivores as ecosystem
68 engineers where they are absent (Seddon *et al.* 2014; Ceausu *et al.* 2015; Naundrup &
69 Svenning 2015). This policy is based on the premise that before the late Quaternary

70 extinctions of megafauna (Koch & Barnosky 2006; Stuart 2015) large herbivores were
71 important drivers of ecosystem disturbance, function and biodiversity (Donlan *et al.* 2005;
72 Vera 2000; Sandom *et al.* 2014; Corlett, 2013). However, this pre-human baseline scenario is
73 not without its critics, with an alternative view that large herbivores were 'passive' on early
74 landscapes and not ecosystem engineers (Bradshaw, Hannon & Lister, 2003; Birks, 2005).

75 One of the main obstacles in this debate has been a limited understanding of past population
76 dynamics of large herbivores (Bradshaw, Hannon & Lister 2003; de Bryun *et al.* 2011). A
77 number of previous studies have demonstrated the presence of large herbivores on past
78 landscapes and their impact on ecosystems using fossil spores from coprophilous ascomycetes
79 fungi (hereafter dung fungal spores) (Burney, Robinson & Burney 2003; Gill *et al.* 2009; Rule *et*
80 *al.* 2012; Baker, Bhagwat & Willis 2013; Froyd *et al.* 2014). These dung fungal spores are
81 unintentionally ingested by large herbivores while feeding on vegetation, and germinate after
82 digestion when deposited with dung. Mycelium growth and fructifications have species-
83 specific responses to different moisture levels, temperatures, microclimates, microhabitats
84 and types of dung and, when successful, the sticky spores are released explosively onto
85 surrounding vegetation, ready to be ingested (Dix & Webster 1995; Krug, Benny & Keller 2004;
86 Bell 2005). Without wind, dung fungal spores are typically ejected up to 30 cm away from the
87 fruiting body (e.g. Yafetto *et al.* 2008). With wind, the majority of spores are deposited within
88 meters (Jackson & Lyford 1999); however, occasionally spores can travel further away (e.g.
89 Gonianakis *et al.* 2005, Hernández Trejo *et al.* 2011). Because dung fungi are strictly reliant on
90 large herbivores' digestive tracts to complete their life cycle, the presence of these spores in
91 sediments when recovered during palaeoecological investigations are interpreted as a
92 compelling evidence for the presence of large herbivores (van Geel *et al.* 2003; Davis & Shafer
93 2006; Baker, Bhagwat & Willis 2013; Johnson *et al.* 2015).

94 Despite the increasing use of dung fungi to study past populations of large herbivores, it
95 remains as yet unclear whether and how the abundance of those spores in sediments can

196 indicate herbivore *densities* within a landscape (Raper & Bush 2009; Feranec *et al.* 2011; Wood
197 & Wilmshurst 2011; Parker & Williams 2011; Baker, Bhagwat & Willis 2013, Etienne *et al.*
198 2013). A number of factors can potentially obscure the relationship between spore abundance
199 and herbivore abundance. They include the differential spore production of dung fungal
200 species and their reliance on specific herbivore species, as well as other taphonomic processes
201 such as average distance travelled by spores and spatial patterns of deposition within waters.
202 These factors are under-researched and evidence can be contradictory. For instance, within
203 water bodies, higher dung fungal spore abundances are reported to be positively related to
204 inflow proximity (Etienne *et al.* 2013), shore proximity (Raper & Bush 2009) and shore distance
205 (Parker & Williams 2011).

206 The aim of this study was therefore to determine whether the spores can be used as a
207 *quantitative* proxy of herbivore density over time. We addressed this question by calibrating
208 dung fungal spore abundance in relation to contemporary herbivore presence, i.e. by
209 examining the density of dung spores in modern sediments in relation to known herbivore
210 biomass densities (biomass per surface area, kg ha⁻¹).

211 The objectives of the study were as follows:

- 212 (i) To determine the way dung fungal spores travel into sediments i.e. long distance
213 transportation by wind, short distance transportation by wind, transportation with
214 surface run-off, or a combination of these transportation mechanisms.
- 215 (ii) To understand the relationship between different spore types and different herbivore
216 species.
- 217 (iii) To ascertain whether changes in fossil spore abundances through time in a single
218 sequence can be used to infer changes in herbivore density on the surrounding
219 landscape.

220 **Methods**

121 *Study site and sample collection*

122 The Oostvaardersplassen nature reserve, The Netherlands, was established on polder land
123 reclaimed from Lake IJsselmeer in 1968. Re-wilding was initiated at this site from 1983 with
124 the introduction of free-ranging Heck cattle (*Bos taurus* Linnaeus) in 1983, Konik horses (*Equus*
125 *ferus caballus* Linnaeus) in 1984 and red deer (*Cervus elaphus* Linnaeus) in 1992. These
126 herbivores have access to the whole nature reserve but mainly use about 2000 ha of
127 grasslands (e.g. *Lolium perenne* L., *Poa trivialis* L., *Trifolium repens* L.), tall herbs (e.g. *Cirsium*
128 spp, *Urtica dioica* L.), reed (*Phragmites australis* (Cav.) Trin. ex Steud.), *Sambucus nigra* L.
129 scrub and *Salix* spp. shrubs (Figure 1) of high net primary productivity (Cornelissen & Vulink
130 2015). The grasslands are visited by large numbers of geese (thousands to tens of thousands of
131 greylag goose *Anser anser* Linnaeus; Barnacle goose *Branta leucopsis* Bechstein; white-fronted
132 goose *Anser albifrons* Scopoli). The site is managed with a policy of minimal intervention, i.e.
133 the population size of freely roaming large herbivores is not controlled by culling, no
134 supplementary feeding is given during winter and no management intervention is
135 implemented to maintain vegetation.

136 We sampled modern sediment from 16 ponds (mean surface area: 2573 m², see Table 2) to
137 obtain a good spatial spread across the reserve (see Figure 1) and a large amplitude of
138 herbivore abundances. These ponds were created between 1985 and 2000 for avian
139 biodiversity. The sediment samples were collected where water was the deepest within each
140 pond and were made up of top sediments representing contemporary deposition. We used a
141 simple tube sampler with sharpened edges. Sedimentary sequences were also collected using
142 a simplified Livingstone corer in order to obtain a sedimentary record of the lifespan of three
143 ponds since they were first created. To prevent further fungal growth, all samples were stored
144 in sealed plastic bags at 4⁰C until processed in laboratory.

145 *Response variable: spore abundance in sediments*

146 The spores were extracted from one 1 cm³ sub-samples per sediment sample collected from
147 the 16 ponds. We followed a standard extraction method in pollen analysis to isolate spores,
148 estimate their concentration using *Lycopodium* spore tablets (batch 938934, Lund University)
149 and carry out identifications at a 400x magnification (Willis & Bennett 2001). Spore
150 identification and spore association with obligate dung fungi were based on the literature
151 reviewed in Baker, Bhagwat & Willis (2013). The abundance of spores in sediments was
152 calculated as accumulation rates (spore cm⁻² year⁻¹) using the spore concentration (spore cm⁻³)
153 and the sedimentation rate (cm year⁻¹) (Maher 1981; Bennett 1994; Willis & Bennett 2001).
154 Using our three cores, we estimated sedimentation rates on the basis of the age of the ponds
155 and the depth of sediment deposited since creation as detailed in Appendix 1. We averaged
156 these sedimentation rates and applied the average to all samples from our 16 ponds. Applying
157 a constant sedimentation rate to all our samples means that the analysis undertaken returns
158 exactly the same results whether we use spore concentration or spore accumulation rates. We
159 opted for analysing and presenting realistic spore accumulation rates throughout this paper in
160 order to facilitate comparison with similar studies in the future.

161 *Explanatory variables: herbivore biomass densities and physical variables*

162 The distribution of herbivores (cattle, horses, deer and geese) was monitored by the reserve
163 wardens on a weekly basis at the Oostvaardersplassen (see Figure 1 B). The data analysed in
164 this paper span the period 2005-2009, for which we established the average number of
165 individuals for every species in each small, medium and large nested plot (see Figure 1 C).
166 Small plots were the basic unit for monitoring large herbivores. For ponds overlaying two small
167 plots, data from the relevant small plots were aggregated. Medium plots included the relevant
168 small plot(s) plus adjacent small plots freely accessible by large herbivores. Large plots
169 represented uninterrupted grasslands delimited by ditches or abrupt vegetation changes
170 known to be of relevance for large herbivore movement. Other animals including foxes and
171 large birds represent a negligible herbivore biomass in comparison with those monitored and

172 there are only incomplete associated data regarding their numbers. To account for the
173 difference of dung production in goose species, red deer, Konik horse and Heck cattle, we
174 used herbivore biomass as a proxy for the dung production. We transformed herbivore
175 numbers into herbivore biomass density (i.e. biomass per surface area, kg ha^{-1}) using an
176 average biomass of individuals per species. The biomass of herbivore species was compiled
177 from Dunning (2007) for geese, and from the long-term monitoring of large herbivore biomass
178 by the RWS Water Service and the State Forestry Office, The Netherlands, for cattle, horses
179 and deer (see Appendix 2).

180 Physical variables of each pond (i.e. pond surface area, pond shore length) and surrounding
181 habitat (i.e. total length of shores within small, medium and large nested plots, total surface
182 area of grassland within small, medium and large nested plots) were calculated on the basis of
183 0.5 x 0.5 m resolution georeferenced aerial photographs taken in 2010 (Ministry of
184 Infrastructure and the Environment and the Ministry of Economic Affairs, The Netherlands).

185 *Data analysis*

186 In order to determine the predominant mechanism of transport of dung fungal spores into the
187 ponds, we used spore abundance as the response variable and compared it to three spore
188 transportation mechanisms (surface run-off, wind-transportation from local shores and wind-
189 transportation from local grasslands) as three explanatory variables. The factors included in
190 each of the transportation mechanisms are detailed in Table 1. They all account for herbivore
191 biomass density around the ponds and for the surface area of the pond, in keeping with the
192 well-studied transportation of pollen grains from vegetation into water bodies (Sugita 1993;
193 Giesecke & Fontana 2008). They differ as follows. 'Surface run-off' accounts for the pond
194 perimeters, or shoreline lengths, to distinguish spores produced on the shore of the sampled
195 ponds. 'Wind-transportation from local shores' accounts for the total length of shorelines
196 around the pond, to distinguish spores produced from habitats with permanent moisture

197 supply, an important factor for dung fungal growth (Dix & Webster 1995; Krug, Benny & Keller
198 2004). 'Wind-transportation from local grasslands' accounts for the surface area of grassland
199 around the ponds to distinguish spores produced from the overall density of dung around the
200 ponds. An additional transportation mechanism, background spore rain (i.e. constant
201 deposition of spores across the reserve) was accounted for as the intercept of our models. We
202 used generalized linear models (GLM) in R (R Core Team, 2012) to examine this relationship.
203 Because over-dispersion of our count data was highlighted by our initial analyses using a link
204 function for Poisson distribution, we used negative binomial regression throughout the
205 analysis. This was chosen over the quasi-Poisson alternative, because more weight on
206 sampling points with higher spore counts was not deemed appropriate in our case (Hoef &
207 Boveng 2007; O'Hara & Kotze 2010). Models were fitted using all three explanatory variables
208 without interaction and stepwise-simplified using the function stepAIC of the MASS package
209 (Venables & Ripley 2002) in order to recover the minimal adequate model, aka optimum
210 model (Crawley 2007).

211 In order to understand the relationship between the different spore types and different
212 herbivore species, we analysed separately herbivore densities (kg ha^{-1}) of geese, Konik horses,
213 Heck cattle, red deer and the three mammalian large herbivores together (5 sets of
214 explanatory variables) against each of the three main dung fungal spore types (*Sporormiella*,
215 *Sordaria* and *Podospora*, see Baker, Bhagwat & Willis 2013) and their sum (therefore 4
216 response variables), resulting in 20 distinct optimum models. Each of these 20 models had
217 been initially selected, on the basis of AICs, out of the 27 optimum models representing all
218 combinations of scales the transportation mechanisms were available (three transportation
219 mechanisms measured each at three nested plot sizes).

220 **Results**

221 A total of 21 modern sediment samples was collected from 16 ponds of similar morphology
222 (mean surface area: 2573 m^2) across the reserve (Table 2). Throughout those samples, 17

223 fungal spore types were identified but only those from *Sporormiella* spp., *Sordaria* spp. and
224 *Podospora* spp. had regular occurrence and made up c. 70% of all 370 fungal spore identified.
225 Dung fungal spore abundance varied between 161 and 2049 (spore cm⁻² year⁻¹) (mean=945
226 sd=537 N=16) and herbivore biomass densities between 308 and 1863 (kg ha⁻¹) (mean=728
227 sd=448 N=16) around the ponds (small plot scale). The cores had an overall average
228 sedimentation rate of 1.14 (cm year⁻¹) and showed little variation within the
229 Oostvaardersplassen (see Appendix 1).

230 Our results demonstrate that there is a quantitative relationship between total dung fungal
231 spore abundance and total biomass density of large herbivores (Table 3). In particular, shore
232 run-off explained, in a highly significant way, total spore abundance. Plots of the significant
233 relationships between the spore types and transportation by surface run-offs are shown in
234 Figure 2. Local wind dispersal, whether from the grasslands or other nearby pond shorelines,
235 did not contribute to spore influx into the sediments.

236 The background spore deposition is highly significantly different from zero and positive,
237 implying a spatially constant atmospheric input of spores across the Oostvaardersplassen. The
238 maximum likelihood estimation of the background spore deposition was 318.7 (dung fungal
239 spore cm⁻² year⁻¹) (95% CI between 428.5 and 237.1) and was in the vicinity of this value for all
240 models presented in Tables 3. The absence of spatial autocorrelation for the spore abundance
241 in our sediment samples (Moran's I test, observed =0.05787573, expected = -0.06666667, sd =
242 0.1011752, p = 0.2183387), supports the very local origin of spore abundance in sediments.

243 Total mammal biomass density of large herbivores related better to total spore abundance
244 than any of the herbivore biomass densities taken individually. While biomass densities of
245 Heck cattle or Konik horses both showed a good fit with the models, biomass densities of red
246 deer were never significantly related to any of the spore types using the methodology

247 adopted. Total biomass densities of geese only showed significant relationship with *Sordaria*
248 abundance.
249 *Sordaria* and *Sporormiella* taken separately show very similar patterns. They are best
250 explained by total biomass density of large herbivores. However, the significance levels are
251 lower for *Sporormiella*. On the contrary, *Podospora* is better explained by biomass densities of
252 Konik horse alone. In this case though, the results should be interpreted cautiously because
253 two samples stood out as outliers on model checking plots, suggesting potential
254 heteroscedasticity and potential non-normal errors.

255 **Discussion**

256 In the introduction we highlighted the factors that can potentially obscure the relationship
257 between herbivore biomass densities and dung fungal spores in sediments. These factors
258 relate to spore production by different fungal species and whether they rely on specific
259 herbivores, as well as to the taphonomic processes that the spores experience between their
260 release in the air and their deposition into sediments. Our results demonstrate a highly
261 significant relationship between spore abundance and local biomass densities of large
262 herbivores when these biological and taphonomic factors are taken into account.

263 *Fungal and herbivore species*

264 Total spore abundance, *Sordaria* abundance and *Sporormiella* abundances were each best
265 explained by total biomass density of mammals (kg ha^{-1}). This demonstrates that dung fungal
266 spores as identified in this study, do not indicate any herbivore species in particular but
267 instead indicate Heck cattle, Konik horse and red deer collectively. This finding differs from
268 some other studies of dung fungal diversity (e.g. Richardson 1972, 2001); however, these
269 earlier studies identify dung fungi to the species level using whole living organisms, i.e. with a
270 greater taxonomic precision than is currently possible using spore morphology alone (Baker,
271 Bhagwat & Willis 2013). Therefore, it would appear that the current limitations in the

272 identification of dung fungal spores from sediments limits our ability to infer which specific
273 large herbivore species they are associated with. However, our data on dung fungal spore
274 abundance suggest that there is a direct link between spore production and the total biomass
275 density of large herbivores. *Podospora* abundance was associated with Konik horse biomass
276 density but the validity of the modelling method for this spore type should be interpreted with
277 some caution. This limitation is probably due to the low number of individual *Podospora*
278 spores recovered and does not concern *Sporormiella* and *Sordaria* types that were significantly
279 more abundant.

280 Similarly, although *Sordaria* spore explained goose abundance significantly, this herbivore is
281 less likely than mammals to be the main source of fungal spore according the AIC selection
282 method used. This result is congruent with bird dung being reported to be a substrate less
283 suitable for fungal growth (Richardson 2001; Doveri 2007). As a result, the important grazing
284 pressure from goose and other birds of similar size (Jano, Jefferies & Rockwell 1998; Jefferies &
285 Rockwell 2002) may not be well captured by the abundance of dung fungal spores. Application
286 of our method away from arctic wetlands or other areas favoured by geese would therefore
287 convey total grazing pressure with greater accuracy. However, there is clear evidence that the
288 past presence of larger flightless herbivorous birds in New Zealand can be tracked using the
289 spores of *Sporormiella* spp. (Wood *et al.* 2011). Therefore, more studies such as ours but in
290 different environments and featuring other species of herbivore (e.g. Froyd *et al.* 2014) will be
291 required to fully assess how our results can be applied to other situations.

292 Using our model selection methodology, it was not possible to statistically determine whether
293 it is preferable to aggregate all dung fungal spores into one indicator or to keep them
294 separate. This was because there is no widely accepted method to calculate absolute
295 goodness of fit, or R^2 , for GLM. Nevertheless, the confidence intervals plotted in Fig. 2
296 highlight that aggregated dung fungal spores would have a higher predictive power for

297 palaeoecological reconstructions than spore types taken individually. In fact, significant
298 increases in aggregated spore abundance appear to be a systematic reflection of an increase in
299 herbivore biomass densities. Subtle changes, and spore types taken individually, may be more
300 difficult to interpret. In addition, studies of dung fungal diversity (e.g. Richardson 1972, 2001)
301 highlight the preference of certain species for certain types of dung. Moreover, the large body
302 of evidence reviewed by Dix & Webster (1995) and Krug, Benny & Keller (2004) shows that
303 dung fungi have species-specific responses to different environmental conditions. The
304 consequent assumption is that the greater the diversity of spore types, the more likely it is to
305 capture all large herbivore activities. Dix & Webster (1995) also highlight the importance of
306 competition between species as a driver for the composition of the dung fungal community.
307 This suggests that dung fungal biomass, and thus ultimately spore production, is strongly
308 limited by factors such as space, nutrient and moisture availability. The ecology of dung fungi
309 therefore suggests that the sum of individual dung spores, irrespective of the type
310 encountered, provides the most appropriate measurement of herbivore biomass.

311 *Taphonomic factors*

312 Our analysis demonstrates that surface run-off from the shoreline and surrounding slopes (as
313 opposed to longer distances by wind) explains dung fungal spore abundances highly
314 significantly. This is the first time evidence is gained regarding the source area of dung fungal
315 spores from water bodies (Feranec *et al.* 2011; Baker, Bhagwat & Willis 2013). The main
316 implication of this finding is that a time-series of spore abundance tracks herbivore abundance
317 in the close proximity of the sampled water body (in our case less than 10 m away from pond
318 shore). At the same time, drinking water from water features such as those sampled in this
319 study directly determine the daily movements of wild and domesticated herbivores (e.g.
320 Putfarken *et al.* 2008; Shannon *et al.* 2009). Thus, water features are a strategic location to
321 sample and it can be postulated that the local herbivore abundance quantified with dung
322 fungal spore actually represents herbivore abundance in a broader landscape. In our samples,

323 there is in addition a significant influx of spores that is not related to the local distribution of
324 herbivores within the reserve and that we identify as background spore deposition. The
325 extremely high herbivore biomass density prevailing in the reserve (474.3 kg ha^{-1} , see
326 Appendix 2) in comparison to the surrounding land (mostly arable land, built-up areas and
327 open water, where large herbivores are overall in low density) suggest that much of this
328 background spore influx originates within the reserve. This indicates that the influx of wind-
329 dispersed spores in our case represents a signal from the overall abundance of large
330 herbivores within the reserve that is not specific to the exact location of sampling. This
331 contrasts with Gill *et al.*'s (2013) study in North America which demonstrated the importance
332 of short-distance wind dispersal (<100 meters) to explain the significant relationship between
333 bison local distribution and abundance of dung fungal spores. Their study was conducted in
334 terrestrial habitats away from water, so further research will be necessary to assess fully the
335 relative importance of run-off and wind transportation in different deposition environments.

336 We found that biomass density of large herbivores explains dung fungal spore abundance in a
337 highly significant way when accounted in conjunction with the morphological characteristics of
338 the sampled pond (i.e. pond surface area and pond shoreline length). Therefore, changes in
339 spore abundance through time can be used to indicate large herbivore population size
340 variation. However, this is with the caveat that the water body has stayed approximately the
341 same size during the same interval in time. If drastic hydrological changes are suspected, there
342 are several means to assess water level in palaeoecology, notably using macrofossils of aquatic
343 plants (Hannon & Gaillard 1997; Dieffenbacher-Krall & Halteman 2000).

344 **Conclusion**

345 There is much debate regarding the long-term impact of large herbivores on their
346 environment. As a consequence, there is great difficulty in predicting with certainty the impact
347 that wild and domestic large herbivores might have, particularly in conjunction with the
348 unpredictable effects of global change. Several factors can influence fluctuations in large

349 herbivore population dynamics worldwide: for instance, agricultural abandonment in marginal
350 areas, the growing need for food production and the adoption of novel conservation strategies
351 such as rewilding. Our aim was to develop a method for the measurement of long time-series
352 of large herbivore population sizes in relation to environmental factors affecting or impacted
353 by those populations because this is a critical step towards improving our understanding of
354 herbivore-dominated ecosystems. Based on an existing method in palaeoecology, we provide
355 here the foundations for the quantitative reconstruction of long time-series of herbivore
356 densities using fossil dung spores contained in sedimentary sequences.

357 Using modern surface sediments, we found that there is a significant relationship between
358 biomass density of large herbivores and dung fungal spore abundance in sediments. To
359 extrapolate this relationship into the past we ascertained that when the morphology of the
360 water body sampled remains the same, accurate quantitative reconstructions are possible.

361 **Acknowledgement**

362 We thank Hans Breeveld for access to the Oostvaardersplassen nature reserve. The herbivore
363 data were provided by the State Forestry Service of the Netherlands. This work was carried
364 out as part of a UK Natural Environment Research Council studentship to AGB. We thank John
365 Birks, David Etienne and an anonymous reviewer for their insightful comments and
366 recommendations.

367 **Data Accessibility**

368 The data used are archived in the appendices of Dr Ambroise Baker's D.Phil Thesis, University
369 of Oxford. See <http://ora.ox.ac.uk/>

370 **Literature cited**

371 Asner, G.P., Elmore, A.J., Olander, L.P., Martin, R.E. & Harris, A.T. (2004). Grazing Systems,
372 Ecosystem Responses, and Global Change. *Annual Review of Environment and Resources*,
373 **29**, 261–299.

- 374 Baker, A.G., Bhagwat, S.A. & Willis, K.J. (2013). Do dung fungal spores make a good proxy for
375 past distribution of large herbivores? *Quaternary Science Reviews*, **62**, 21–31.
- 376 Bell, A. (2005). *An Illustrated Guide to the Coprophilous Ascomycetes of Australia*. APS Press,
377 St-Paul.
- 378 Bennett, K.D. (1994). Confidence intervals for age estimates and deposition times in late-
379 Quaternary sediment sequences. *The Holocene*, **4**, 337–348.
- 380 Birks, H.J.B. (2005) Mind the gap: how open were European primeval forests? *Trends in*
381 *Ecology and Evolution*, **20**, 154–156.
- 382 Bradshaw, R.H.W., Hannon, G.E. & Lister, A.M. (2003). A long-term perspective on ungulate–
383 vegetation interactions. *Forest Ecology and Management*, **181**, 267–280.
- 384 Burney, D.A., Robinson, G.S. & Burney, L.P. (2003). *Sporormiella* and the late Holocene
385 extinctions in Madagascar. *Proceedings of the National Academy of Sciences of the*
386 *United States of America*, **100**, 10800–10805.
- 387 Carpenter, S.R., Mooney, H.A., Agard, J., Capistrano, D., Defries, R.S., Díaz, S., Dietz, T.,
388 Duraipappah, A.K., Oteng-Yeboah, A., Pereira, H.M., Perrings, C., Reid, W. V, Sarukhan, J.,
389 Scholes, R.J. & Whyte, A. (2009). Science for managing ecosystem services: Beyond the
390 Millennium Ecosystem Assessment. *Proceedings of the National Academy of Sciences of*
391 *the United States of America*, **106**, 1305–12.
- 392 Ceaușu, S., Hofmann, M., Navarro, L.M., Carver, S., Verburg, P.H. & Pereira, H.M. (2015).
393 Mapping opportunities and challenges for rewilding in Europe. *Conservation Biology*, **29**,
394 1017-1027.

- 395 Corlett, R.T. (2012). The shifted baseline: Prehistoric defaunation in the tropics and its
396 consequences for biodiversity conservation. *Biological Conservation*, **163**, 13–21.
- 397 Cornelissen, P. & Vulink, J.T. (2015). Density-dependent diet selection and body condition of
398 cattle and horses in heterogeneous landscapes. *Applied Animal Behaviour Science*, **163**,
399 28-38.
- 400 Crawley, M.J. (2007). *The R Book*. Wiley, Chichester, UK.
- 401 Cromsigt, J.P.G.M. & te Beest, M. (2014). Restoration of a megaherbivore: Landscape-level
402 impacts of white rhinoceros in Kruger National Park, South Africa. *Journal of Ecology*,
403 **102**, 566–575.
- 404 Davis, O.K. & Shafer, D.S. (2006). *Sporormiella* fungal spores, a palynological means of
405 detecting herbivore density. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **237**,
406 40–50.
- 407 De Bruyn, M., Hoelzel, A.R., Carvalho, G.R. & Hofreiter, M. (2011). Faunal histories from
408 Holocene ancient DNA. *Trends in ecology & evolution*, **26**, 405–13.
- 409 Dieffenbacher-Krall, A. & Halteman, W. (2000). The relationship of modern plant remains to
410 water depth in alkaline lakes in New England, USAe. *Journal of Paleolimnology*, **24**, 213–
411 229.
- 412 Dix, N.J. & Webster, J. (1995). *Fungal ecology*. Chapman & Hall, London, UK.
- 413 Donlan, J.C., Berger, J., Bock, C.E., Bock, J.H., Burney, D.A., Estes, J.A., Foreman, D., Martin,
414 P.S., Roemer, G.W., Smith, F.A., Soulé, M.E. & Greene, H.W. (2005). Re-wilding North
415 America. *Nature*, **436**, 913–914.

416 Doveri, F. (2007). *Fungi Fimicoli Italici: A Guide to the Recognition of Basidiomycetes and*
417 *Ascomycetes Living on Faecal Material*. Associazione Micologica Bresadola, Trento.

418 Dunning, J.B. (2007). *CRC Handbook of Avian Body Masses*, secondn. CRC Press, London.

419 Ellis, E.C., Klein Goldewijk, K., Siebert, S., Lightman, D. & Ramankutty, N. (2010). Anthropogenic
420 transformation of the biomes, 1700 to 2000. *Global Ecology and Biogeography*, **19**, 589–
421 606.

422 Estes, J.A., Terborgh, J., Brashares, J.S., Power, M.E., Berger, J., Bond, W.J., Carpenter, S.R.,
423 Essington, T.E., Holt, R.D., Jackson, J.B.C., Marquis, R.J., Oksanen, L., Oksanen, T., Paine,
424 R.T., Pickett, E.K., Ripple, W.J., Sandin, S.A., Scheffer, M., Schoener, T.W., Shurin, J.B.,
425 Sinclair, A.R.E., Soulé, M.E., Virtanen, R. & Wardle, D.A. (2011). Trophic downgrading of
426 planet Earth. *Science*, **333**, 301–6.

427 Etienne, D., Wilhelm, B., Sabatier, P., Reys, J.-L., Arnaud, F. (2013). Influence of sample
428 location and livestock numbers on *Sporormiella* concentrations and accumulation rates in
429 surface sediments of Lake Allos, French Alps. *Journal of Paleolimnology*, **49**, 117–127.

430 Feranec, R.S., Miller, N.G., Lothrop, J.C. & Graham, R.W. (2011). The *Sporormiella* proxy and
431 end-Pleistocene megafaunal extinction: A perspective. *Quaternary International*, **245**,
432 333–338.

433 Froyd, C.A., Coffey, E.E.D., van der Knaap, W.O., van Leeuwen, J.F.N., Tye, A. & Willis, K.J.
434 (2014). The ecological consequences of megafaunal loss: giant tortoises and wetland
435 biodiversity. *Ecology letters*, **17**, 144–54.

436 Giesecke, T. & Fontana, S.L. (2008). Revisiting pollen accumulation rates from Swedish lake
437 sediments. *The Holocene*, **18**, 293–305.

- 438 Gill, J.L., McLauchlan, K.K., Skibbe, A.M., Goring, S., Zirbel, C.R. & Williams, J.W. (2013). Linking
439 abundances of the dung fungus *Sporormiella* to the density of bison: implications for
440 assessing grazing by megaherbivores in palaeorecords. *Journal of Ecology*, **101**, 1125–
441 1136.
- 442 Gill, J.L., Williams, J.W., Jackson, S.T., Lininger, K.B. & Robinson, G.S. (2009). Pleistocene
443 megafaunal collapse, novel plant communities, and enhanced fire regimes in North
444 America. *Science*, **326**, 1100–1103.
- 445 Gonianakis, M., Neonakis, I., Darivianaki, E., Gonianakis, I., Bouros, D. & Kontou-Fili, K. (2005).
446 Airborne Ascomycotina on the island of Crete: Seasonal patterns based on an 8-year
447 volumetric survey. *Aerobiologia*, **21**, 69–74.
- 448 Hannon, G.E. & Gaillard, M.-J. (1997). The plant-macrofossil record of past lake-level changes.
449 *Journal of Paleolimnology*, **18**, 15–28.
- 450 Hernández Trejo, F., Muñoz Rodríguez, A.F., Tormo Molina, R. & Silva Palacios, I. (2011).
451 Airborne ascospores in Mérida (SW Spain) and the effect of rain and other
452 meteorological parameters on their concentration. *Aerobiologia*, **28**, 13–26.
- 453 Hoef, J. Ver & Boveng, P. (2007). Quasi-Poisson vs. negative binomial regression: how should
454 we model overdispersed count data? *Ecology*, **88**, 2766–2772.
- 455 Jackson, S.T. & Lyford, M.E. (1999). Pollen dispersal models in Quaternary plant ecology:
456 Assumptions, parameters, and prescriptions. *The Botanical Review*, **65**, 39–75.
- 457 Jano, A., Jefferies, R. & Rockwell, R. (1998). The detection of vegetational change by
458 multitemporal analysis of LANDSAT data: the effects of goose foraging. *Journal of*
459 *Ecology*, **86**, 93–99.

- 460 Jefferies, R. & Rockwell, R. (2002). Foraging geese, vegetation loss and soil degradation in an
461 Arctic salt marsh. *Applied Vegetation Science*, **5**, 7–16.
- 462 Johnson, C.N., Rule, S., Haberle, S.G., Turney, C.S.M., Kershaw, A.P. & Brook, B.W. (2015).
463 Using dung fungi to interpret decline and extinction of megaherbivores: problems and
464 solutions. *Quaternary Science Reviews*, **110**, 107–113.
- 465 Koch, P.L. & Barnosky, A.D. (2006). Late Quaternary Extinctions: State of the Debate. *Annual*
466 *Review of Ecology, Evolution, and Systematics*, **37**, 215–250.
- 467 Krug, J.C., Benny, G.L. & Keller, H.W. (2004). Coprophilous fungi. *Biodiversity of Fungi* (eds
468 G.M. Mueller, G.F. Bills & M.S. Foster), pp. 468–499. Elsevier, Amsterdam.
- 469 Maher, L.J. (1981). Statistics for microfossil concentration measurements employing samples
470 spiked with marker grains. *Review of Palaeobotany and Palynology*, **32**, 153–191.
- 471 Naundrup, P.J. & Svenning, J.-C. (2015). A Geographic Assessment of the Global Scope for
472 Rewilding with Wild-Living Horses (*Equus ferus*). *Plos One*, **10**, e0132359.
- 473 Nuttle, T., Yerger, E.H., Stoleson, S.H. & Ristau, T.E. (2011). Legacy of top-down herbivore
474 pressure ricochets back up multiple trophic levels in forest canopies over 30 years.
475 *Ecosphere*, **2**, art4.
- 476 O'Hara, R.B. & Kotze, D.J. (2010). Do not log-transform count data. *Methods in Ecology and*
477 *Evolution*, **1**, 118–122.
- 478 Parker, N.E. & Williams, J.W. (2011). Influences of climate, cattle density, and lake morphology
479 on *Sporormiella* abundances in modern lake sediments in the US Great Plains. *The*
480 *Holocene*, **22**, 475–483.

- 481 Peh, K.S.-H. & Lewis, S.L. (2012). Conservation implications of recent advances in biodiversity–
482 functioning research. *Biological Conservation*, **151**, 26–31.
- 483 Putfarken, D., Dengler, J., Lehmann, S. & Härdtle, W. (2008). Site use of grazing cattle and
484 sheep in a large-scale pasture landscape: A GPS/GIS assessment. *Applied Animal*
485 *Behaviour Science*, **111**, 54–67.
- 486 Raper, D. & Bush, M. (2009). A test of *Sporormiella* representation as a predictor of
487 megaherbivore presence and abundance. *Quaternary Research*, **71**, 490–496.
- 488 Richardson, M.J. (1972). Coprophilous ascomycetes on different dung types. *Transactions of*
489 *the British Mycological Society*, **58**, 37–48.
- 490 Richardson, M.J. (2001). Diversity and occurrence of coprophilous fungi. *Mycological Research*,
491 **105**, 387–402.
- 492 Ripple, W.J. & Beschta, R.L. (2012). Trophic cascades in Yellowstone: The first 15 years after
493 wolf reintroduction. *Biological Conservation*, **145**, 205–213.
- 494 Ritchie, E.G., Elmhagen, B., Glen, A.S., Letnic, M., Ludwig, G. & McDonald, R. a. (2012).
495 Ecosystem restoration with teeth: what role for predators? *Trends in ecology &*
496 *evolution*, **27**, 265–71.
- 497 Rule, S., Brook, B.W., Haberle, S.G., Turney, C.S.M., Kershaw, A.P. & Johnson, C.N. (2012). The
498 Aftermath of Megafaunal Extinction: Ecosystem Transformation in Pleistocene Australia.
499 *Science*, **335**, 1483–1486.
- 500 Sandom, C.J., Ejrnaes, R., Hansen, M.D.D. & Svenning, J.-C. (2014). High herbivore density
501 associated with vegetation diversity in interglacial ecosystems. *Proceedings of the*
502 *National Academy of Sciences*, **111**, 4162–4167.

- 503 Seddon, P.J., Griffiths, C.J., Soorae, P.S. & Armstrong, D.P. (2014). Reversing defaunation:
504 Restoring species in a changing world. *Science*, **345**, 406–412.
- 505 Shannon, G., Matthews, W.S., Page, B.R., Parker, G.E. & Smith, R.J. (2009). The affects of
506 artificial water availability on large herbivore ranging patterns in savanna habitats: a new
507 approach based on modelling elephant path distributions. *Diversity and Distributions*, **15**,
508 776–783.
- 509 Steinfeld, H., Gerber, P. & Wassenaar, T. (2006). *Livestock's long shadow*. FAO, Rome.
- 510 Stuart, A.J. (2015). Late Quaternary megafaunal extinctions on the continents: a short review.
511 *Geological Journal*, **50**, 338–363.
- 512 Sugita, S. (1993). A Model of Pollen Source Area for an Entire Lake Surface. *Quaternary*
513 *Research*, **39**, 239–244.
- 514 Tanentzap, A.J. & Coomes, D.A. (2012). Carbon storage in terrestrial ecosystems: do browsing
515 and grazing herbivores matter? *Biological reviews of the Cambridge Philosophical Society*,
516 **87**, 72–94.
- 517 Van Geel, B., Buurman, J., Brinkkemper, O., Schelvis, J., Aptroot, A., van Reenen, G.B.A. &
518 Hakbijl, T. (2003). Environmental reconstruction of a Roman Period settlement site in
519 Uitgeest (The Netherlands), with special reference to coprophilous fungi. *Journal of*
520 *Archaeological Science*, **30**, 873–883.
- 521 Venables, W.N. & Ripley, B.D. (2002). *Modern Applied Statistics with S*, Fourth Edin. Springer,
522 New York.
- 523 Vera, F.W.M. (2000). *Grazing ecology and forest history*. CABI, Wallingford.

524 Willis, K.J. & Bennett, K.D. (2001). Pollen. *Tracking environmental change using lake sediments*
525 *Volume 3 Terrestrial, Algal and Siliceous Indicators* (eds J.P. Smol, H.J.B. Birks & W.M.
526 Last), pp. 5–32. Kluwer Academic Publishers, Dordrecht.

527 Wood, J.R. & Wilmshurst, J.M. (2011). Wetland soil moisture complicates the use of
528 *Sporormiella* to trace past herbivore populations. *Journal of Quaternary Science*, **27**, 254–
529 259.

530 Wood, J.R., Wilmshurst, J.M., Worthy, T.H. & Cooper, A. (2011). *Sporormiella* as a proxy for
531 non-mammalian herbivores in island ecosystems. *Quaternary Science Reviews*, **30**, 915–
532 920.

533 Yafetto, L., Carroll, L., Cui, Y., Davis, D.J., Fischer, M.W.F., Henterly, A.C., Kessler, J.D., Kilroy,
534 H.A., Shidler, J.B., Stolze-Rybczynski, J.L., Sugawara, Z. & Money, N.P. (2008). The Fastest
535 Flights in Nature: High-Speed Spore Discharge Mechanisms among Fungi (A.S. Gladfelter,
536 Ed.). *PLoS ONE*, **3**, e3237.

537

538

539 Table 1: Transportation mechanism used for the analysis. 'distance A' can be either
540 small, medium or large nested plots for which we had herbivore densities, 'pond i' is
541 any of the 16 ponds sampled.

	Transport agent	Location of dung	Factors included
Run-off	Surface run-off, very short flight, erosion	Pond shoreline and slopes	$H_{A,i}$: Herbivore biomass density within distance A of pond i (kg ha^{-1}) P_i : Perimeter of pond i (m) S_i^{-1} : (Surface area of pond i) $^{-1}$ (m^{-2})
Wind (shore)	Turbulent air	Shorelines and slopes from nearby water bodies	$H_{A,i}$: Herbivore biomass density within distance A of pond i (kg ha^{-1}) $SL_{A,i}$: Shore length of other water bodies within distance A of pond i (m) S_i^{-1} : (Surface area of pond i) $^{-1}$ (m^{-2})
Wind (land)	Turbulent air	Nearby grasslands	$H_{A,i}$: Herbivore biomass density within distance A of pond i (kg ha^{-1}) $L_{A,i}$: Surface area of grassland within distance A of pond i (m^2) S_i^{-1} : (Surface area of pond i) $^{-1}$ (m^{-2})
542 Background	Turbulent air	Unspecified	(Model intercept)

543

544

545

546 Table 2: List of ponds, their characteristics and samples analysed. Note that five ponds
 547 were sampled twice and therefore have two sample years, two sample codes and two
 548 sample depths.

Pond	Surface (m square)	Shore length (m)	Total shore length within medium plot (m)	North	East	Year(s) sampled	Sample code(s)	Sample depth(s) (cm)
OO2	7081	414	1657	52.43050800	5.39363240	2009, 2010	Oost-2, Oost-213	20, 7
OO3	3092	487	4583	52.43812010	5.39735413	2009, 2010	Oost-3, Oost-222	20, 9
OO4	4403	525	6776	52.44220897	5.40135547	2009	Oost-4	20
OO5	4419	696	6776	52.44513021	5.39834122	2009	Oost-5	20
OO6	2687	322	2274	52.43912714	5.38580996	2009, 2010	Oost-6, Oost-3/1	20, 1.5
OO7	1521	204	4250	52.42868373	5.38800097	2009, 2010	Oost-7, Oost-217	20, 8
OO8	203	56	172	52.42444476	5.34973321	2009	Oost-8	20
OO9	1063	171	658	52.42969659	5.36165301	2009	Oost-9	20
OO10	5705	323	3209	52.43070174	5.31460139	2009	Oost-10	20
OO11	2011	169	5412	52.42800156	5.30843074	2009	Oost-11	20
OO12	1998	250	4250	52.42982508	5.39061816	2010	Oost-214	10
OO13	443	76	75	52.42389313	5.37887143	2010	Oost-216	10
OO14	2368	274	4338	52.42603232	5.38194373	2010	Oost-201	8
OO15	1206	178	2428	52.42438731	5.37715128	2010	Oost-202	7
OO16	1621	254	6776	52.44189417	5.40323788	2009, 2010	Oost-203, Oost-204	10, 5
549 OO17	1353	247	6209	52.43784175	5.39378056	2010	Oost-205	7

550

551


552

553 Table 3: Identification of transportation mechanisms. Each line summarises the
 554 optimum model with the lowest AIC out of a series of 27 optimum models (all
 555 combinations of 3 plot scales for 3 transportation mechanisms). Significance codes for
 556 p-values: '****' <0.001; '***' <0.01; '**' <0.05; ns otherwise. (-) indicates a negative
 557 relationship, and small, medium and large for the nested plots (see text and Figure 1
 558 C); x: mechanism excluded by model simplification.

Response variables	Biomass density (kg ha ⁻¹) for:	AIC	Significance levels				Optimum plot scales		
			run-offs (shore)	wind (shores)	wind (land)	Intercept	run-off (shore)	wind (shores)	wind (land)
Dung Fungal Spore Total	mammal	240.9	***	(-) ns		***	small	medium	x
Dung Fungal Spore Total	Heck cattle	242.8	***	(-) *		***	small	medium	x
Dung Fungal Spore Total	Konik horse	241.7	***	(-) **		***	small	large	x
Dung Fungal Spore Total	red deer	246.4		(-) *	ns	***	x	medium	medium
Dung Fungal Spore Total	goose	247.3		(-) ns	*	***	x	large	large
Sordaria -type	mammal	224.6	**			***	small	x	x
Sordaria -type	Heck cattle	226.4	**	(-) *		***	small	medium	x
Sordaria -type	Konik horse	226.5	**	(-) ns		***	small	large	x
Sordaria -type	red deer	228.9				***	x	x	x
Sordaria -type	goose	227.2	**	(-) **	*	***	medium	medium	large
Sporormiella -type	mammal	210.5	*			***	medium	x	x
Sporormiella -type	Heck cattle	211.7			ns	***	x	x	large
Sporormiella -type	Konik horse	212.1			ns	***	x	x	large
Sporormiella -type	red deer	211.8	ns			***	large	x	x
Sporormiella -type	goose	212.7				***	x	x	x
<i>Podospora</i> -type	mammal	199.0	ns			***	small	x	x
<i>Podospora</i> -type	Heck cattle	199.3				***	x	x	x
<i>Podospora</i> -type	Konik horse	197.5	**	(-) ns		***	small	large	x
<i>Podospora</i> -type	red deer	197.9		(-) *		***	x	medium	x
<i>Podospora</i> -type	goose	199.3				***	x	x	x

560

561



563


564 Figure 1: Maps of the Oostvaardersplassen, sample locations indicated with triangles.

565 A. Vegetation map highlighting the grasslands where the herbivores spend most of

566 their time. B. Monitoring method, the number of herbivores is recorded weekly for

567 each of the small plots following established routes, in red. C. The nested plots utilised

568 in this study (aggregation of the small plots shown in B).



569
 570 Figure 2: Optimum relationships between spore abundances and herbivore biomass
 571 densities. On the x axis is the run-off transportation mechanism, i.e. the product of
 572 local herbivore biomass density (only Konik horses for *Podospora*), pond shore length
 573 and pond surface area. Dots in red are the observed data, in black the best model
 574 prediction as in Table 3 (dashed, 0.95 confidence interval).