

## **Routine adherence monitoring after renal transplantation**

*Invited Commentary for Transplant International*

**Simon R Knight** <sup>1,2</sup>

1. Nuffield Department of Surgical Sciences, University of Oxford, UK
2. Centre for Evidence in Transplantation, University of Oxford, UK

*Correspondence to:*

Mr Simon R Knight  
Oxford Transplant Centre  
Churchill Hospital  
Old Road  
Oxford OX3 7LE  
United Kingdom

Tel : +44 1865 227131

E-mail: [simon.knight@nds.ox.ac.uk](mailto:simon.knight@nds.ox.ac.uk)

Successful renal transplantation requires lifelong immunosuppression, usually with a combination of drugs, to prevent rejection and immunologically-mediated damage to the graft. To achieve maximum effect, these immunosuppressive agents must be taken regularly as prescribed. This is particularly true of critical-dose drugs such as tacrolimus, with a narrow therapeutic window between efficacy and toxicity. There is good evidence that poor adherence to immunosuppression is deleterious to graft outcomes, with one systematic review suggesting a 7-fold increase in the risk of graft loss in poorly adherent patients (1).

Assessment of medication adherence is challenging. Non-adherence can take a number of forms, including missed doses, complete discontinuation, mistimed dosing or incorrect dosing. It may affect one or all drugs in the regimen and depend on a number of factors such as side effect profile, dose frequency, route of administration and lifestyle factors (peer pressure, work and social commitments). In everyday clinical practice adherence is rarely formally assessed, with clinicians relying on identifying patients from at-risk groups (younger age, male sex, lower education status (2)) or using surrogate markers for adherence (low or variable tacrolimus levels, early acute rejection episodes, poor clinic attendance).

Formal assessment of adherence is usually reserved for the clinical trial setting, with use of self-reporting questionnaires, smartphone monitoring, prescription refills or electronic pill-counting. Whilst these methods might give a better estimate of true medication compliance, they are often expensive, subject to manipulation and the measurement itself may have an effect on adherence (3).

In the current edition of *Transplant International*, Gustavsen and colleagues evaluate a number of possible methods for the assessment of adherence in a real-world renal transplant population for use in a national transplant registry. Patients were recruited to two cohorts – a prospective cohort randomised to more or less intensive adherence assessment, and a cross-sectional cohort to control for the potential effect of assessment itself on adherence. Adherence was assessed by a combination of the self-reported Basel Assessment of Adherence to Immunosuppressive medication Scale (BAASIS), tacrolimus trough inpatient variability (IPV) and an adherence score from the treating clinician. Interestingly, the adherence tools appeared to capture different patient populations, and whilst more intensive assessment captured more non-adherence events, it did not appear to affect adherence behaviour overall. Poor adherence, as measured by the tools used here, did not significantly increase the risk of biopsy-proven acute rejection, but was associated with an increased risk of de novo donor-specific antibody development.

The focus in this study was to identify tools that can be used in routine clinical practice for registry purposes. As such the validated BAASIS questionnaire is a simple to administer, fast self-reporting tool consisting of just 4 questions regarding missed doses and administration timing over the past 4 weeks. Return rates in the current study were good (82% at 1 year), although there may be an element of selection bias in those patients prepared to take part.

Measurement of tacrolimus IPV is also straightforward, requiring 3 or more trough tacrolimus measurements over a time-period that can be used to calculate a co-efficient of variation. High IPV has shown association with poor clinical outcomes (4,5), and is associated with other non-adherence attitudes and behaviours (6,7). However, it only captures adherence to tacrolimus, and adherence to other immunosuppressants may differ depending on dose frequency and side effect profile. Once

patients are more than 1-year post-transplant, the frequency of trough measurements decreases meaning that IPV is assessed over a longer period and may miss shorter-term fluctuations in levels. Despite this limitation, 88% patients in the current cohort had sufficient data for IPV calculation at 1 year.

The benefit of the addition of clinician adherence assessment is less clear. As already described, clinicians assess adherence based upon demographic risk, clinical events and drug levels. The latter has quite a large influence, and so there is likely to be considerable overlap with those non-adherent patients identified through high tacrolimus IPV. This is supported by the higher Cohen's kappa value reported for the association between clinician's score and IPV, although agreement was still only fair. At 1-year post-transplant, patient-clinician interaction is likely to be limited making assessment challenging. Indeed, rates of clinician-scored non-adherence at 1 year (7%) were lower than those assessed by the BAASIS questionnaire (32%) and tacrolimus IPV (13%).

One of the major drawbacks of the study from Gustavsen and colleagues is the lack of a gold-standard assessment for true non-adherence, making it difficult to identify the optimal tool or combination of tools for monitoring. Pill-counting was attempted in one arm of the study, but return rates were poor (43%) and further analysis was not possible. This highlights the difficulties of pill-counting in a real-world patient group, suggesting that it is not a suitable tool for adherence assessment outside the clinical trial setting.

Overall, the study from Gustavsen and colleagues suggests that routine adherence assessment for registry purposes is possible using any of these three tools, and that a combination of tools may be required for a complete picture. Ultimately, the usefulness of these assessments will depend on response rates, but incorporation of these measures in a national registry has the potential to provide a very useful dataset to assess factors influencing adherence in the long-term post-transplantation. These tools may also prove useful in adherence assessment in clinical trials where the primary purpose of the study is not to influence adherence and formal assessment (e.g. pill counting) is not feasible.

## References

1. Butler JA, Roderick P, Mullee M, Mason JC, Peveler RC. Frequency and impact of nonadherence to immunosuppressants after renal transplantation: a systematic review. *Transplantation* 2004; 77 (5): 769.
2. Prihodova L, Nagyova I, Rosenberger J et al. Adherence in patients in the first year after kidney transplantation and its impact on graft loss and mortality: a cross-sectional and prospective study. *Journal of Advanced Nursing* 2014; 70 (12): 2871.
3. Sutton S, Kinmonth A-L, Hardeman W et al. Does Electronic Monitoring Influence Adherence to Medication? Randomized Controlled Trial of Measurement Reactivity. *Annals of Behavioral Medicine* 2014; 48 (3): 293.
4. Shuker N, Shuker L, van Rosmalen J et al. A high inpatient variability in tacrolimus exposure is associated with poor long-term outcome of kidney transplantation. *Transplant International: Official Journal of the European Society for Organ Transplantation* 2016; 29 (11): 1158.

5. Sapir-Pichhadze R, Wang Y, Famure O, Li Y, Kim SJ. Time-dependent variability in tacrolimus trough blood levels is a risk factor for late kidney transplant failure. *Kidney International* 2014; 85 (6): 1404.
6. Tielen M, van Exel J, Laging M et al. Attitudes to Medication after Kidney Transplantation and Their Association with Medication Adherence and Graft Survival: A 2-Year Follow-Up Study, Attitudes to Medication after Kidney Transplantation and Their Association with Medication Adherence and Graft Survival: A 2-Year Follow-Up Study. *Journal of Transplantation, Journal of Transplantation* 2014; 2014, 2014: e675301.
7. Goodall DL, Willicombe M, McLean AG, Taube D. High Inpatient Variability of Tacrolimus Levels and Outpatient Clinic Nonattendance Are Associated With Inferior Outcomes in Renal Transplant Patients. *Transplantation Direct [Internet]* 2017; [cited 2018 Oct 24] 3 (8). Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5540630/>