Adsorption at the Air-water interface in Biosurfactant–Surfactant mixtures: Quantitative Analysis of Adsorption in a Five Component Mixture

Jessica R. Liley,^{†,||} Robert K. Thomas,^{*,†} Jeffrey Penfold,^{‡,⊥} Ian M. Tucker,[¶]
Jordan T. Petkov,^{¶,#} Paul S. Stevenson,[¶] Ibrahim M. Banat,[§] Roger Marchant,[§]
M Rudden,[§] and John R. P. Webster[‡]

[†]*Physical and Theoretical Chemistry Laboratory, South Parks Road, Oxford, OX1 3QZ,
United Kingdom*

[‡]*STFC, Rutherford-Appleton Laboratory, Chilton, Didcot, Oxfordshire, OX11 0QX, United
Kingdom*

[¶]*Unilever Research and Development Laboratory, Port Sunlight, Quarry Road East,
Bebington, Wirral CH63 3JW, United Kingdom*

[§]*School of Biomedical Sciences, University of Ulster, Coleraine, Northern Ireland*

^{||}*Current address: LGC, Queens Road, Teddington, Middlesex, TW11 0LY, United
Kingdom*

[⊥]*Physical and Theoretical Chemistry Laboratory, South Parks Road, Oxford, OX1 3QZ,
United Kingdom*

[#]*Current address: Lonza UK, GB-Blackley, Manchester, Lancs., M9 8ES, United Kingdom*

E-mail: robert.thomas@chem.ox.ac.uk

Abstract

The composition of the air–water adsorbed layer of a quinary mixture consisting of three conventional surfactants, octa–ethylene glycol monododecyl ether ($C_{12}E_8$), dodecane–6–p–sodium benzene sulfonate (LAS6), and diethylene glycol monododecyl ether sodium sulfate (SLE_2S) mixed with two biosurfactants, the rhamnolipids L–rhamnosyl–L–rhamnosyl– β –hydroxydecanoyl– β –hydroxydecanoyl, R2, and L–rhamnosyl– β –hydroxydecanoyl– β –hydroxydecanoyl, R1, has been measured over a range of compositions above the mixed critical micelle concentration (CMC). Additional measurements on some of the subsets of ternary and binary mixtures have also been measured by NR. The results have been analysed using the pseudophase approximation in conjunction with an excess free energy, G_E , that depends on quadratic and cubic terms in the composition. The compositions of the binary, ternary and quinary mixtures could all be fitted to two sets of interaction parameters between pairs of surfactants, one for micelles and one for adsorption. No ternary interactions or ternary corrections were required. Since the system contains two strongly anionic surfactants, this indicates that the pseudophase approximation can be extended in practice to ionic surfactants, contrary to the prevailing view. The values of the interaction parameters show that the quinary mixture, SLE_2S –LAS6– $C_{12}E_8$ –R1–R2, which is known to be a highly effective surfactant system, is characterized by a sequence of strong surface but weak micellar interactions. About half of the minima in G_E for the strong surface interactions occur well away from the regular solution value of 0.5.

Introduction

There is an increasing interest in the properties of biosurfactants as the development of bio-sustainable and biodegradable surfactant based products becomes more important.¹ A wide range of different biosurfactants is produced by many different organisms, of which one of the more commonly studied and most promising categories is the glycolipids, e.g. rhamnolipids

and sphorolipids. Glycolipids are disaccharides acetylated by long chain fatty acids and their production and basic characterisation have been studied by several authors.^{2,3} Apart from their biosustainability and biodegradability, these biosurfactants exhibit a high tolerance to pH, temperature and salinity, have anti-microbial and anti-bacterial properties, and can be synthesised from a variety of non-petrochemical sources. These attractive properties have already led to a diverse range of applications which include enhanced oil recovery,⁴ bioremediation of heavy metals,⁵ and some specialised applications in health care, cosmetics and food processing.⁶

The rhamnolipids are one of the most extensively studied biosurfactants,⁷ but their potential application has been hindered by the challenges of cheap large scale production and purification,⁸ as a result of which the most promising route to a more widespread incorporation in formulation is by blending them with different conventional surfactants. This requires a detailed understanding of their mixing behaviour in multi-component mixtures with a range of different conventional surfactants, and this is the focus of this paper. Rhamnolipids are produced by a variety of different strains of *Pseudomonas aeruginosa* and generally exist with a structure which has one or two rhamnose molecules linked to one or two molecules of β -hydroxydecanoic acid. The two most common forms, L-rhamnosyl-L-rhamnosyl- β -hydroxydecanoyl- β -hydroxydecanoyl, Rha₂C₁₀C₁₀ or R2, and L-rhamnosyl- β -hydroxydecanoyl- β -hydroxydecanoyl, RhaC₁₀C₁₀ or R1, are the ones studied here.

The surface adsorption⁹⁻¹⁴ and self-assembly^{13,15-18} of the rhamnolipids have been studied mainly by surface tension, although there are a small number of x-ray, light and neutron scattering studies. Chen et al.^{19,20} used neutron reflectivity (NR) to study the adsorption of the rhamnolipids R1 and R2, and their mixtures with each other and with dodecane-6-p-sodium benzene sulfonate (LAS6) at the air-water interface. R1 is more surface active than R2 and in mixed R1-R2 adsorption R1 preferentially partitions at the interface, R2 competing less favourably due to the steric or packing constraints associated with the bulkier R2 headgroup. The mixing behaviour is non-ideal and is not consistent with the straightforward

application of the regular solution approximation.²¹ Mixtures of R1 or R2 with the anionic surfactant LAS6 are weakly non-ideal, and only R1 competes strongly with LAS6 for the interface. In ternary R1–R2–LAS6 mixtures at pH 9 in buffer, the relative surface activities were found to be in the order LAS6 > R1 > R2. An unusual feature of the ternary mixtures is that there is a pronounced synergy in the total adsorption at a LAS6–rhamnolipid composition of 1:1, a synergy that is not observed in any of the associated binary mixtures.

Surfactant mixing has been extensively studied and thermodynamic treatments based on the pseudo phase approximation (PPA) are well established.^{21–25} The two main problems with the PPA are that it is generally thought to apply only to nonionic surfactants and that it is mainly used only with the regular solution approximation, which restricts the excess free energy of mixing, G_E , for a binary mixture to be Bx_1x_2 , where B is an interaction parameter and x_i the mole fraction. On the more positive side, the PPA provides an intermediate stage between the direct experimental data, from surface tension or neutron reflectometry (NR) measurements, and the interaction parameters. The latter are highly suitable for comparison with theoretical models. At present, theoretical models either use the PPA to link molecular level calculations with measurements on multi-component systems or they bypass it altogether, e.g.^{26,27} The ability of neutron reflectometry (NR) to determine directly the adsorbed amounts of each component in multi-component mixtures at the air-water interface both below and above the critical micelle concentration (CMC), allows a greater range of types of G_E to be fitted to experiment, e.g.,^{28–31} and this in turn allows some assessment of the range of validity of the PPA. Here we use NR to probe the surface mixing properties of a quinary (5-component) surfactant mixture, in which the rhamnolipids R1 and R2 are combined with the ternary mixture of conventional surfactants, octa-ethylene glycol monododecyl ether ($C_{12}E_8$), diethylene glycol monododecyl ether sodium sulfonate (SLE_2S) and LAS6. The measurements were all done above the CMC.

The ternary $C_{12}E_8$ –LAS6– SLE_2S surfactant mixture is a good model representation of surfactant based formulations typical of many home and personal care products. Recently

Liley et al.^{30,31} have used NR to make a detailed study of the surface mixing at the air-water interface of the C₁₂E₈–LAS6–SLE₂S mixture in the absence of electrolyte and in the presence of NaCl and CaCl₂. At surfactant concentrations in excess of the mixed CMC the surface mixing is highly non-ideal and the surface composition is still significantly different from the solution composition. The relative surface activities are in the order C₁₂E₈ > LAS6 > SLE₂S with the result that the surface is significantly depleted of the anionic SLE₂S, even at surfactant concentrations $\gg$ CMC. The non-ideality is consistent with the PPA when terms of higher order than the quadratic B terms (see above) are included in the expression for G_E . Addition of electrolyte drives the surface composition much closer to the solution composition for concentrations $\gg$ CMC, and there is a shift in the relative surface activities in the order LAS6 > SLE₂S > C₁₂E₈. The focus of this study is to understand how replacing part of the C₁₂E₈–LAS6–SLE₂S ternary mixture with rhamnolipids affects the adsorption and the balance in the relative surface mole fractions.

Experimental Details

The samples of h-C₁₂E₈, d-C₁₂E₈, h-LAS6, d-LAS6, h-SLE₂ and d-SLE₂, where d and h indicate perdeuterated (98%, unless stated otherwise) and protonated hydrophobic units respectively, were the same ones as used by Liley et al., who described in detail their synthesis and characterization.³⁰ The hydrogeneous rhamnolipids were obtained from Jeneil Biosurfactant Co. and separated into the pure R1 and R2 components as described by Chen et al.¹⁹ The deuterium labelled rhamnolipids, d-R1 and d-R2 were grown in a *Pseudomonas aeruginosa* culture fed with D₂O and glycerol-d₇. The initial extraction of the surface active components and subsequent purification have been described elsewhere.⁸ The pure d-R1 and d-R2 components were separated and characterised using the same procedure as used for the hydrogeneous surfactants and were approximately 90 % deuterium labelled. The purity of the surfactants was assessed by surface tension below and up to the CMC and by NR measurements at concentrations above the CMC. D₂O (99.9 %) was obtained from Fluorochem.

High purity water (Elga Ultrapure) was used throughout. All glassware for solution preparation and Teflon troughs for the NR experiments were cleaned in dilute Decon90 solution and rinsed in ultrapure water, ethanol, and acetone, and dried under a nitrogen gas flow.

The NR measurements were made at a surfactant concentration of 2 mM, well above the mixed CMC for all the mixtures studied. Measurements were made for the binary mixtures R1–C₁₂E₈, R2–C₁₂E₈, R1–SLE₂S, and R2–SLE₂S, and for the ternary mixtures R1–R2–C₁₂E₈ and R1–R2–SLE₂S. Data for the equivalent R1, R2 and LAS6 mixtures were taken from Chen et al.^{19,20} The measurements for the ternary mixtures were made at a fixed R1–R2 mole ratio of 1:1, and variable R (R1 + R2)–C for each of C = C₁₂E₈, LAS6 or SLE₂S mixtures. A series of measurements were made for the 5-component mixture of R1–R2–C₁₂E₈–LAS6–SLE₂S at a fixed mole ratio of overall R:overall C of 30 : 70, i.e. 30 mole % of the C₁₂E₈–LAS6–SLE₂S ternary mixture was replaced by rhamnolipid. The measurements were made for two rhamnolipid R1–R2 mole ratios of 1:1 and 1:2, shown in two triangular diagrams as Figure S1 in the Supporting Information. At the fixed R:C and R1:R2 mole ratios the measurements were made over a range of C₁₂E₈–LAS6–SLE₂S compositions. All the solutions in the present set of measurements were adjusted by the addition of small amounts of NaOH solution to give a final pH of approximately 6.5.

For experiments in null reflecting water (NRW) the neutron reflectivity R is given by

$$R = \frac{64\pi^2}{\kappa^4} \rho^2 \sin^2 \left(\frac{\kappa d}{2} \right) \quad (1)$$

where d and ρ are the thickness and scattering length density of the adsorbed layer, κ is the wave vector transfer in the direction perpendicular to the surface, z , which is given by $\kappa = 4\pi(\sin \theta)/\lambda$, where θ is the grazing angle of incidence and λ is the neutron wavelength. The scattering length density ρ equals $\sum_i b_i n_{iz}$ where n_i is the number density of species i and b_i its neutron scattering length. NRW is a 92 : 8 mole % H₂O:D₂O mixture with a scattering length density of zero, and hence a refractive index the same as air. For a

monolayer of deuterium labelled surfactant adsorbed at the interface the reflectivity then arises only from the adsorbed layer. For a single surfactant species at the interface the area per molecule A is then related to the product $d \times \rho$ and the $\sum_i b_{ij}$ value of the surfactant by

$$A = \frac{\sum_i b_{ij}}{d\rho} \quad (2)$$

where i are the component nuclei of surfactant j . The surface excess, Γ , is given by $\Gamma = 1/N_a A$. For a ternary mixture Eqn (2) becomes

$$d\rho = \frac{\sum_i b_{i1}}{A_1} + \frac{\sum_i b_{i2}}{A_2} + \frac{\sum_i b_{i3}}{A_3} \quad (3)$$

and this is similarly extended to a quinary mixture. For mixtures, a series of NR measurements with each of the components deuterium labelled in turn leads to a set of simultaneous equations, which can be solved to determine the adsorbed amounts of each component. The values of the sum of scattering lengths for the different surfactants studied here have been given in previous papers.^{19,30} For the binary, ternary and quinary mixtures measurements were made for the respective isotopic combinations of dd, dh, and hd, ddd, dhh, hdd, and hhd, and and ddddd, dhhhh, hddhh, hhdhh, hhhhd, and hhhhd in NRW, where d and h refer to the deuterium labelled and hydrogenous hydrophobic chains of the surfactants. In all cases the system is over determined and the sets of 3, 4 or 6 simultaneous equations based on Eqns (2) and (3) and their extension to five components were solved as described previously.³⁰

The NR measurements were made on the SURF and INTER reflectometers at ISIS³² and on FIGARO at the ILL.³³ The measurements were made at a fixed glancing angle of incidence θ using a range of wavelengths sorted by time of flight, to measure $R(\kappa)$. The reflected intensity was normalised to the direct beam and the absolute reflectivity values were calibrated by reference to the reflectivity from the surface of D₂O. The samples were contained in sealed Teflon troughs held at 298K and containing a sample volume of ≈ 25

mL.

Results

Analysis Procedure

The compositions of the adsorbed layers above the CMC were analysed using the PPA and an excess free energy, G_E , containing both quadratic and cubic terms, as described by Liley et al.³⁰ Each binary pair is characterized by four parameters, the quadratic and cubic terms in G_E , i.e. B^μ and C^μ for micellization, and B^σ and C^σ for surface adsorption. The primary aim of the experiments is to test whether parameters obtained for the binary mixtures can be used to describe quantitatively the adsorption behaviour in the ternary and quinary mixtures, i.e. we attempt to fit the ternary and quinary data without any additional parameters. However, for a two parameter G_E the sensitivity of the data to each of the four parameters varies with the range of surface compositions being probed, which may also vary significantly between binary, ternary and quinary mixtures. It is therefore more even-handed to follow the approach of Liley et al. which was to attempt to fit the whole set of data to a single set of parameters for micellization and adsorption.

The excess free energy of mixing, G_E , can be expressed as a series consisting of quadratic (regular solution) and cubic terms in the mole fraction x_i where i is numbered from 1 to 5³⁴

$$G_E = \sum_{j < k} x_j x_k B_{jk} + \sum_{j < k} x_j x_k (x_j - x_k) C_{jk} \quad (4)$$

where we note that the sign of C_{jk} must be opposite to that of C_{kj} because of the $(x_j - x_k)$ term. For a ternary mixture with just quadratic interactions the activity coefficient of component one is^{23,35}

$$RT \ln f_{1B} = x_2^2 B_{12} + x_2 x_3 (B_{12} - B_{23} + B_{31}) + x_3^2 B_{31} \quad (5)$$

and the other two coefficients are obtained by cycling the indices. If there are non-zero cubic terms there is an additional contribution for f_1 ^{36,37}

$$RT \ln f_{1C} = [2x_1x_2(1 - x_1 + x_2) - x_2^2] C_{12} + 2x_2x_3(x_3 - x_2)C_{23} - [2x_3x_1(1 - x_1 + x_3) - x_3^2] C_{31} \quad (6)$$

where f_{2C} and f_{3C} are again obtained by cycling the indices. The overall activity coefficient must be the product of the two terms, i.e. $f = f_{1B}f_{1C}$. The importance of introducing the cubic terms is that the optimum composition, i.e. minimum in G_E , of the mixed layer (or micelle) does not need to be equimolar as in the regular solution approximation. For a quinary mixture the activity coefficients can again be obtained from Eqn (4), as shown in the Supporting Information. As an example, the activity coefficient for component 1 is

$$\begin{aligned} RT \ln f_1 &= (1 - x_1)(x_2B_{12} + x_5B_{15} + x_3B_{13} + x_4B_{14}) \\ &\quad - x_2x_3B_{23} - x_2x_4B_{24} - x_2x_5B_{25} - x_3x_4B_{34} - x_3x_5B_{35} - x_4x_5B_{45} \\ &= [2x_1x_2(x_2 - x_1) + x_2(2x_1 - x_2)] C_{12} + [2x_1x_3(x_3 - x_1) + x_3(2x_1 - x_3)] C_{13} \\ &\quad + [2x_1x_4(x_4 - x_1) + x_4(2x_1 - x_4)] C_{14} + [2x_1x_5(x_5 - x_1) + x_5(2x_1 - x_5)] C_{15} \\ &\quad + 2x_2x_3(x_3 - x_2)C_{23} + 2x_2x_4(x_4 - x_2)C_{24} + 2x_2x_5(x_5 - x_2)C_{25} \\ &\quad + 2x_3x_4(x_4 - x_3)C_{34} + 2x_3x_5(x_5 - x_3)C_{35} + 2x_4x_5(x_5 - x_4)C_{45} \end{aligned} \quad (7)$$

and the full set of activity coefficients is given in the Supporting Information.

The use of the PPA assumption means that at equilibrium the chemical potential of each component is the same in micelle, solution, and surface layer. The micelle-solution equilibrium then leads to the set of fundamental equations for each component,²¹⁻²⁴

$$x_i = \frac{c_i^{mon}}{f_i c_i^\mu} \quad (8)$$

where x_i is the mole fraction of component i in the micelle (i again running from 1 to 5), c_i^{mon} the monomer concentration of i in the bulk solution, f_i its activity coefficient in the

micelle (the activity coefficient, or mean activity coefficient for an ionic surfactant in the bulk is taken to be 1), and c_i^μ its CMC. The assumption also implies that all the surfactants are undissociated, a feature that we will return to in the Discussion.

Above the CMC, where all the measurements were made, adsorption depends on the monomer activity (mole fraction) in the solution, as well as on factors determining the intrinsic strength of the adsorption. The monomer composition is obtained using an equation resulting from mass balance combined with Eqn (8)²²

$$x_i = \frac{\alpha_i c - f_i x_i c_i^\mu}{c - \sum_j f_j x_j c_j^\mu} \quad (9)$$

where α_i and c are the overall compositions and concentration of the surfactants, and x_i and f_i are again the mole fractions and activity coefficients in the micelle. The further substitution of $\sum x_i = 1$ allows one of the compositions to be eliminated resulting in a set of $n - 1$ quadratic equations where n is the number of components. For the ternary mixture the two equations are

$$\begin{aligned} A_{31}x_1^2 + A_{32}x_1x_2 + (c - A_{31})x_1 - \alpha_1c &= 0 \\ A_{32}x_2^2 + A_{31}x_1x_2 + (c - A_{32})x_2 - \alpha_2c &= 0 \end{aligned} \quad (10)$$

where

$$A_{ij} = f_i c_i^\mu - f_j c_j^\mu \quad (11)$$

All the relevant functions can be differentiated in Eqns (10), (5) and (6), which means that the equations can be solved iteratively to obtain the set of micelle compositions x_i^μ using the Newton–Raphson method.³⁰ The equivalent Eqns (9) and (10) for a quinary mixture lead to a set of equations, which can be written generally as

$$A_{ji}x_i^2 + \left[c - A_{ji} + \sum_{k \neq i, k \neq j} A_{jk} \right] x_i - \alpha_1 c = 0 \quad (12)$$

where $A_{ji} = f_j c_j^\mu - f_i c_i^\mu$. When for example x_5 is eliminated, the following four quadratic equations in the four independent compositions are obtained

$$\begin{aligned}
A_{51}x_1^2 + [A_{52}x_2 + A_{53}x_3 + A_{54}x_4 + c - A_{51}]x_1 - \alpha_1c &= 0 \\
A_{52}x_2^2 + [A_{53}x_3 + A_{54}x_4 + A_{51}x_1 + c - A_{52}]x_2 - \alpha_2c &= 0 \\
A_{53}x_3^2 + [A_{52}x_2 + A_{51}x_1 + A_{54}x_4 + c - A_{53}]x_3 - \alpha_3c &= 0 \\
A_{54}x_4^2 + [A_{51}x_1 + A_{52}x_2 + A_{53}x_3 + c - A_{54}]x_4 - \alpha_4c &= 0
\end{aligned} \tag{13}$$

Since any one of the other four components could also be selected for elimination, there are 20 possible equations of this form so there is a choice of possible ways of calculating the four unknown x_i . These equations have to be solved by iteration and, although the Newton–Raphson method was used for the ternary mixtures, it becomes too cumbersome to apply to the quinary. Holland and Rubingh used the the Nelder-Mead simplex technique for function minimization for their ternary mixtures.³⁷ However, we found that a relatively simple and short iteration could be used for the quinary system.

At concentrations well above the CMC the overall mole fraction becomes approximately equal to the fraction in the micelle, i.e. $x_i = \alpha_i$. Using this approximation as a starting point to obtain a set of x_i , the activity coefficients are calculated using Eqn (7) and the related equations for the f_i (see Supporting Information). This set of x_i and f_i are used to give a group of five quadratic equations from the set of Eqns (12), which are solved to give a new set of x_i and hence a new set of f_i . The iteration is damped by limiting the successive changes. Adjustment of the damping and a careful choice of which quadratic equations to select from Eqns (12), both of which are likely to depend on the system, gave us accurate convergence to the solution in under 20 steps. We were expecting to use the approximation $x_i = \alpha_i$ and then the simple substitution into Eqn (8) gives

$$c_i^{mon} \approx \alpha_i f_i c_i^\mu \tag{14}$$

The individual monomer mole fractions for a quinary mixture are then given directly, i.e. there is no iteration, as

$$x_i^{mon} \approx \frac{\alpha_i}{\alpha_i + \sum_{j \neq i} \alpha_j R_{ji}^f R_{ji}^\mu} \quad (15)$$

where R_{ji}^f and R_{ji}^μ are the ratios f_j/f_i and c_j^μ/c_i^μ respectively. However, with the accurate method we were able to show that for our quinary mixtures the total concentration would have needed to be 10 mM or greater for the approximate method to work, i.e. approximately 50× the highest mixed CMC. The use of the approximation is, however, likely to be system dependent.

The second step in the calculation is to determine the surface composition using the monomer compositions. Eqn (8) now becomes

$$x_i = \frac{c_i^{mon}}{f_i c_i^\pi} \quad (16)$$

where x_i and f_i now refer to the surface layer, the c_i^{mon} have been determined in step one, and c_i^π is a measure of the adsorption strength. We used the Holland approximation³⁷ for c_i^π , in which the assumption is made that a complete monolayer is adsorbed at the CMC, i.e. the CMC is the major determinant of the adsorption but it is modified by a surface pressure factor,

$$c_i^\pi = c_i^\mu \exp \left[\frac{(\pi^{mix} - \pi_i) A_i^{cmc}}{RT} \right] \quad (17)$$

where π_j and π^{mix} are respectively the surface pressures of the pure monomer and the mixture above the CMC, and A_i^{cmc} is the limiting molecular area of i at the CMC. The surface pressure correction factor can be experimentally determined and then an iterative solution of the set of Eqns (16) is used to determine the surface composition. Java programs were written in-house to carry out and check the two iterations.

Figure 1: Variation of adsorbed mole fraction with bulk mole fraction for the three binary mixtures (a) LAS6–R1, (b) LAS6–R2 and (c) R1–R2 at $\text{pH} = 9$ in buffer solution at a total concentration of 1 mM. The fitting parameters are given in Table 1, ideal surface mixing is shown as a thin red line and the monomer composition as a thin blue line.

Rhamnolipid–LAS6 Mixtures

Chen et al. measured the variation of composition of the surface with bulk composition for the binary mixtures LAS6–R1, LAS6–R2, R1–R2, and for the LAS6–R1–R2 ternary mixtures at three different R1:R2 ratios. The data have been published in the Supporting Information of the papers by Chen et al.^{19,20} The original experiments were done at a pH of 9 in a buffer solution of 23 mM borax and 8 mM HCl at a total concentration of 1 mM. At this pH the CMCs of the three surfactants are 0.34 mM for LAS6, 0.36 mM for R1, and 0.18 mM for R2, i.e. the measurements were all above the CMC. These values compare with 1.2, 0.18 and 0.11 mM respectively for LAS6, R1 and R2 at pH 7, also with buffer. For R1 and R2 the behaviour of the CMCs shows that the effect of pH strongly dominates the normal suppression of the CMC by electrolyte and the buffer seems to have little effect on these values at pH 7. The data were fitted using Eqns (10), (5) and (6) for the monomer concentrations in the ternary mixtures above the CMC, Eqns (16) and (17) for the surface, and the binary equivalents for the binary mixtures, all as described by Liley et al.^{30,31} The parameters needed in Eqn (17) were determined experimentally by Chen et al. and were as follows. For R1 and R2 A^{cmc} was found from NR to be respectively 60 and 88 Å² at pH 9 in buffer. For LAS6 NR gives values of A^{cmc} from 52–57 Å², which are not significantly affected by buffer, and we take an average value of 54 Å². The limiting surface tensions were 28.5, 31.2 and 37.4 mN m⁻¹ respectively for LAS6, R1 and R2. The final experimental input needed to apply Eqn (17) is the surface tension of the mixtures. Strictly, there is no fixed value because this varies with composition. However, it only varies over a narrow range and over this range it is not expected to have a large effect. We therefore take it to be constant and for further simplicity we assume it has the same value for all the mixtures, which we take to be 33 mN m⁻². We return to this assumption in the Discussion.

The data and the fits (thick black lines) for the three binary mixtures are shown in Figure 1 and for the three ternary mixtures in Figure 2. The fits use an identical set of B^μ , C^μ , B^σ , and C^σ parameters for each pair of surfactants, and the values are given in Table 1.

Figure 2: Variation of adsorbed mole fractions of each component with bulk mole fraction of LAS6 for three ternary mixtures (a) LAS6 - ($R1:R2 = 2:1$), (b) LAS6 - ($R1:R2 = 1:1$) and (c) LAS6 - ($R1:R2 = 1:2$). The conditions were $pH = 9$ in buffer solution at a total concentration of 1 mM. The fitting parameters are given in Table 1.

Two other sets of lines are included in all the binary diagrams, a thin red line, which is the composition that the adsorbed layer would have if there were ideal mixing at the surface, but not in the micelles, and a blue line, which is the monomer composition in the solution. We have shown elsewhere³¹ that the intersection of the ideal (red) line with the data marks the surface composition at which the minimum in G_E occurs. Thus, Figure 1 (a) shows that for LAS6–R1 the data are intersected by the red line at a composition of about 0.5 (mole fraction LAS6), i.e. C^σ must be close to zero in both cases and the solution is regular. This is not the case for LAS6–R2 nor for the stronger R1–R2 interaction where the intersections are both at about 2/3 (mole fraction of the first named component in each case).

Before further discussion of the fitting of the results for the binary mixtures, it is useful to assess the different contributions of micellization and adsorption to the red and blue lines in the figures. In the absence of any interactions, adsorption of a particular surfactant is driven by two factors, the mixed CMC (c^μ) which controls the monomer composition, and the c^π , which controls adsorption and which is given by Eqn (17). The monomer composition is lower for the component with the lower c^μ because it preferentially segregates into the micelles, and adsorption is stronger for a lower c^π because it adsorbs at a lower concentration. The values of c^μ and c^π For LAS6, R1 and R2 at pH 9 and in buffer are respectively 0.34, 0.36 and 0.18 mM (c^μ) and 0.19, 0.28 and 0.26 mM (c^π). For zero interaction in the micelles the values of c^μ and the total concentration c determine the monomer composition. In the special case where the c^μ are equal the blue line will be a straight diagonal between the points (0,0) and (1,1) for the surfactant that is being plotted. The effect of the preferential segregation into the micelles is that if this surfactant has the higher c^μ the line will become a curve above the diagonal, but below if it has the lower c^μ . However, the preferential segregation is strongest just above the mixed CMC and weakens as the concentration increases, i.e. any curvature in the plot caused by a difference in the c^μ will be exaggerated and distorted as the concentration decreases towards the mixed CMC. A non-zero value of B^μ for equal values of c^μ will cause the straight diagonal line to become sigmoidal (S shaped) if B^μ is

negative or reverse sigmoidal if it is positive, and non-zero values of C^μ will confer a wave like distortion on the diagonal, with correspondingly more complicated profiles if the C^μ are not equal. The change from the monomer composition (blue line) to the ideal adsorption (red line) is determined by the two c^π values. For the simple case of a monomer composition conforming to the diagonal straight line described above, the ideal adsorption curve will be above the line if the main surfactant has the lower c^π because it is more strongly adsorbed, and vice versa. Non-zero values of B^σ and C^σ have effects comparable to those of B^μ and C^μ on the monomer composition, but the signs are reversed. This means that in this simple case identical values of B^μ and B^σ cancel each other out. While this would be an unusual occurrence, there are occasions when this coupling may make it difficult to allocate the parameters between micellization and adsorption.

In practice, this is not as serious a problem as it might seem because (a) the differing pairs of c^π and c^μ usually mean that the distortions of the curves by the interaction parameters are displaced from each other, (b) independent measurements of the c^μ allow an independent determination of the red line, (c) surface measurements below the CMC can be used to determine the adsorption behaviour independently of micellization, and (d) the effects of micellization are altered by changing the overall concentration (the binary version of Eqn (12)). This last effect is large and can be effective for separating the micellization and adsorption parameters, although it has been little used.²⁹ It also comes into play in a more subtle way when mixtures higher than binary are examined, e.g. the ternary and quinary mixtures studied below. In general, the error is larger for the smaller overall interaction (B and C combined) and in most of the cases here the interaction in the micelles is weaker. This is the case for the R1–R2 mixture, where the cancellation effects are small because there is a mismatch in the two minima of G_E , that for the micelles being at 0.5 R1 whereas that for the surface is at 0.64 R1. The ± 0.4 and ± 0.3 errors for micelles and surface respectively in Table 1 are estimates of the errors arising from cancellation effects. For the binary mixtures in Figure 1 the mixed CMCs have also been determined and the fits of the parameters in

Table 1 are shown in Figure S2 of the Supporting Information.

We now apply some of this analysis to Figures 1 and 2. In Figure 1(a) the very similar CMCs of LAS6 and R1 and the weak interaction ($B^\mu = 0.3$) mean that the blue monomer line is almost exactly on the diagonal. However, as shown above, LAS6 has a significantly lower c^π than R1 and hence the red ideal adsorption line is above the blue monomer line. In addition, B^σ is negative and large enough to add a marked reverse sigmoidal shape to obtain the fitted line. C^σ is not exactly zero and this is enough to move the intersection slightly below a mole fraction of 0.5. In Figure 1(b) LAS6 has a significantly higher c^μ than R2 and the blue monomer curve is above the diagonal. The much lower value of c^π for LAS6 compared with R2 makes LAS6 much more surface active and the red ideal adsorption now shows that LAS6 dominates the surface. The reverse sigmoidal shape that has to be added to the ideal mixing to fit the data shows that there is an attractive interaction but this time the intersection of the red and black curves shows that G_E is displaced to 0.67. Finally, Figure 1(c) for R1–R2 is similar to LAS6–R2 except that B^μ is sufficiently large and negative to give some sigmoidal shape to the monomer curve. Nevertheless, R1 dominates the surface. These experiments were done at a concentration (1 mM) where the total concentration has a significant influence on the monomer composition.

It is less easy to disentangle the various effects in the ternary mixtures. First, it is important to note that the way the pairwise interactions are probed in the ternary mixtures is different to the binaries. Thus in the ternary mixture with equimolar R1 and R2, the limits of the range of bulk compositions for LAS6 and R1 are 1 and 0, as in the binaries, but at 0.5 LAS6 the R1 composition in the ternary is only 0.25. Not only is the functional form of the interaction then changed but the total concentration range relative to the mixed CMC is also changed. This is an example of how the factor (d) above can help to resolve ambiguities in cancellation effects between micellization and adsorption. LAS is slightly dominant at the surface in the ternary LAS6–R1–R2 mixtures (Figure 2) in that, at a solution mole fraction of 0.5 LAS6, the fractions of adsorbed LAS6 are 0.59, 0.60 and 0.57 (fitted curves) at the

Table 1: Parameters determining the excess free energy of mixing for the ten binary pairs of surfactants. Note that the sign of C_{ij} depends on the order of the surfactants and i refers to the first named surfactant in each pair. The first seven pairs of parameters are from the results in this paper. The remaining three, marked with asterisks, are from Liley et al.³⁰

system	B^μ/RT ± 0.4	C^μ/RT ± 0.4	B^σ/RT ± 0.3	C^σ/RT ± 0.3	G_{min} RT	x_1
LAS6-R1	-0.3	0	-	-	0.08	0.5
LAS6-R1	-	-	-1.2	0.2	-0.30	0.46
C ₁₂ E ₈ -R1	-0.5	-0.3	-	-	-0.13	0.62
C ₁₂ E ₈ -R1	-	-	-1.5	-0.3	-0.38	0.55
SLE ₂ S-R1	-1.0	-0.5	-	-	-0.26	0.61
SLE ₂ S-R1	-	-	-3.2	-0.6	-0.81	0.55
R1-R2	-0.5	0	-	-	-0.13	0.5
R1-R2	-	-	-1.7	-1.4	-0.48	0.65
LAS6-R2	0	0	-	-	-	-
LAS6-R2	-	-	-0.8	-0.8	-0.24	0.67
C ₁₂ E ₈ -R2	0.2	0	-	-	0.04	0.5
C ₁₂ E ₈ -R2	-	-	0	0	-	-
SLE ₂ S-R2	-0.7	0.7	-	-	-0.21	0.33
SLE ₂ S-R2	-	-	-0.8	0.8	-0.24	0.33
SLE ₂ S-LAS6*	0	0	-	-	-	-
SLE ₂ S-LAS6*	-	-	-2.2	-1.2	-0.59	0.61
LAS6-C ₁₂ E ₈ *	-2.6	1.8	-	-	-0.72	0.37
LAS6-C ₁₂ E ₈ *	-	-	-4.4	0.8	-1.11	0.46
SLE ₂ S-C ₁₂ E ₈ *	-1.8	1.8	-	-	-0.53	0.33
SLE ₂ S-C ₁₂ E ₈ *	-	-	-2.0	2.0	-0.59	0.33

R1:R2 ratios of 1:2, 1:1 and 2:1 respectively, compared with 0.54 in the LAS6–R1 binary, i.e. the LAS6 seems to have become more dominant. However, LAS6 is adsorbed much more strongly in its mixtures with R2 and the weighted average of the composition ratios LAS6:R1 and LAS6:R2 at this point ought to give higher values of 0.62, 0.65 and 0.68 respectively. Thus the adsorption of LAS6 appears to be weakened in the ternaries by a relative amount that increases as the bulk fraction of R1 increases. The origin of this is important. If there is a pair of surfactants in a ternary mixture that have a much stronger attractive interaction than any other pair (here R1 and R2), their compositions at the surface will be increased at the expense of the third component (here LAS6) and this is the basic effect operating in the LAS6–R1–R2 mixtures.^{30,31} Furthermore, the large negative value of C^σ enhances the interaction as the ratio R1:R2 increases, as observed. There is the slight complication in this case in that the moderate attractive interaction between R1 and R2 in the micelles has the opposite effect. However, it is the surface effect that dominates. Finally, the fit to the data in Figure 2 (a) is less good than the other two, and this is the result of a compromise between the fits to this figure, to the right hand side of the binary R1–R2 mixture (Figure 1(c)) and to the CMCs in Figure S2. The latter favours a smaller magnitude for B^μ than the other two.

Rhamnolipid Mixtures with C₁₂E₈ and SLE₂S

The compositions of the surface of the binary mixtures of R1 and R2 with C₁₂E₈ and SLE₂S at an overall concentration of 2 mM are given in Table 2. The monomer and the adsorbed layer compositions were fitted in the same way as described above for the LAS6–R1–R2 systems. The two B and two C parameters for each binary mixture are given in Table 1 and we have again been able to use a single set of values for these surfactant components in binary, ternary and quinary solutions. This means that, as already illustrated for LAS6–R1–R2, a better fit to a particular data set is sometimes sacrificed for a better overall fit. The R1–R2 pair of surfactants occur in the mixtures with LAS6 described in the previous section

Figure 3: Variation of adsorbed mole fraction with bulk mole fraction for the two binary mixtures (a) $C_{12}E_8$ -R1 and (b) $C_{12}E_8$ -R2, and (c) the ternary mixture $C_{12}E_8$ -R1-R2 with equimolar R1 and R2 at $pH = 6.5$ and a total concentration of 2 mM. The fitting parameters are those given in Table 1. The monomer composition is shown as a thin blue line in (a) and (b) and the ideal surface mixing curve as a thin red line in (a). It is obscured by the black line in (b).

and in the pres
previous section
However, their r
areas and limiti
these mixtures a
and $75 \text{ \AA}^2$, and l
all of which cha

Figure 4: Variation of adsorbed mole fraction with bulk mole fraction for the two binary mixtures (a) SLE₂S–R1 and (b) SLE₂S–R2, and (c) the ternary mixture SLE₂S–R1–R2 (equimolar R1 and R2) at pH = 6.5 and overall concentration 2 mM. The fitting parameters are given in Table 1. The ideal surface mixing curves and the monomer compositions for the binary mixtures are shown as thin lines in red and blue respectively.

The plots for the two binary mixtures C₁₂E₈–R1 and C₁₂E₈–R2 are shown in Figure 3 (a) and (b). They show both a fitted line (in black) using the parameters in Table 1 and the curves for ideal mixing and the monomer composition. The values of c^μ and c^π for R2

Table 2: Variation in surface composition and total adsorption at a total concentration of 2 mM for R1–C₁₂E₈, R1–SLE₂S, R2–C₁₂E₈ and R2–SLE₂S at a pH of 6.5.

system	mole fraction R1	surface mole fraction R1 ± 0.04	$10^6 \times \Gamma_{total}$ $\pm 0.1 \text{ mol m}^{-2}$
R1–C ₁₂ E ₈ (2 mM, pH 6.5)	0.10	0.26	3.16
	0.25	0.47	3.30
	0.50	0.62	3.35
R1–SLE ₂ S (2 mM, pH 6.5)	0.10	0.51	3.15
	0.25	0.63	3.04
	0.50	0.74	2.96
	0.75	0.85	2.83
R2–C ₁₂ E ₈ (2 mM, pH 6.5)	0.25	0.34	2.99
	0.50	0.54	2.83
	0.75	0.72	2.25
R2–SLE ₂ S (2 mM, pH 6.5)	0.25	0.61	2.79
	0.50	0.71	2.56
	0.75	0.84	2.20

and C₁₂E₈ are comparable and this, combined with mixing that is near ideal (micellization) and fully ideal (adsorption) results in a monomer curve just above the diagonal and an ideal adsorption curve identical with the fitted adsorption. The differences in c^μ and c^π are significantly larger for R1 ($c^\mu = 0.18$ and $c^\pi = 0.09$ mM) and C₁₂E₈ ($c^\mu = 0.09$ and $c^\pi = 0.14$ mM), so that the monomer line is well above the diagonal, but this is reversed to give the ideal adsorption line. The effect on the monomer line is reinforced by the weak attraction in the micelle. The moderately strong attractive interaction in the adsorbed layer then adds a reverse sigmoidal shape to give the final fitted curve. Thus the adsorption of C₁₂E₈–R1 shows significant synergy, in which the attractive surface interaction is dominated by the quadratic term, i.e. the minimum in G_E at a mole fraction of C₁₂E₈ of 0.54 is close to the value of 0.5 for the regular solution approximation.

As for the LAS6–R1–R2 mixtures the results for the ternary mixture probe the interaction differently from the binaries. However, the main effects here are the two strong surface interactions between C₁₂E₈ and R1 and between R1 and R2. The stronger of these is the R1–R2 interaction and it is markedly unsymmetrical and favours a 2:1 R1:R2 stoichiometry.

In the fitting it was found that the $C_{12}E_8$ –R1 interaction increased these two components at the expense of R2. However, the stronger R1–R2 interaction finally drives an overall increase of the adsorption of R1 and R2 and an overall small decrease of adsorbed $C_{12}E_8$. There is a pronounced difference in the R1:R2 ratio across the diagram when C^σ is non-zero. When the full value is used the R1:R2 ratio changes significantly to reach a fairly uniform 2:1 ratio over the whole range.

Table 3: Surface composition and total adsorbed amount for R1–R2– $C_{12}E_8$ and R1–R2– SLE_2S mixtures. The measurements were done at a total concentration of 2 mM at a pH of 6.5.

total mole fraction $\alpha_{R1} = \alpha_{R2}$			surface mole fraction ± 0.05			$10^6 \times \Gamma_{total}$ $\pm 0.1 \text{ mol m}^{-2}$
R1	R2	$C_{12}E_8$	R1	R2	$C_{12}E_8$	
0.05	0.05	0.90	0.14	0.08	0.76	3.2
0.125	0.125	0.75	0.31	0.05	0.64	3.2
0.25	0.25	0.50	0.45	0.20	0.35	2.9
0.375	0.375	0.25	0.58	0.30	0.12	2.8
0.45	0.45	0.10	0.60	0.34	0.06	2.5
R1	R2	SLE_2S	R1	R2	SLE_2S	
0.05	0.05	0.90	0.41	0.04	0.55	3.4
0.125	0.125	0.75	0.42	0.20	0.38	3.1
0.25	0.25	0.50	0.54	0.20	0.26	2.9
0.375	0.375	0.25	0.60	0.15	0.25	2.8
0.45	0.45	0.10	0.55	0.24	0.22	2.9

Using the same format as for $C_{12}E_8$ the fits and data for the binary SLE_2S –R1 and SLE_2S –R2 and ternary SLE_2S –R1–R2 are shown in Figure 4 using the fitting parameters in Table 1. The most striking features are that because SLE_2S has a much higher CMC than the other two surfactants there is a high concentration of SLE_2S monomer and the monomer curves are well above the diagonal. However, the differences in the c^π are even larger for both R1 and R2 and therefore the ideal adsorption curves reverse the effect of the high SLE_2S monomer concentration to give an ideal curve that is well below the diagonal. The large and negative B^σ imposes the strong characteristic deviation from the ideal adsorption curve. R1 so dominates the adsorption in its mixture with SLE_2S that the adsorbed SLE_2S does not

reach a surface composition of 0.5, where the minimum in G_E is, until its bulk mole fraction reaches 0.96. This is also a situation where the surface concentration range is significantly different from that probed for the micelles, which makes it fairly straightforward to fit the micellar and adsorption parameters reasonably independently.

The fits of the ternary SLE₂S–R1–R2 data are dominated by the two large interactions, between SLE₂S and R1, which is close to symmetrical, and between R1 and R2. The nett effect of these two interactions is that SLE₂S and R2 are more strongly adsorbed while R1 does not change very much. Again, it is noticeable that the effect of the large C^σ for R1–R2 drives these two surfactant into the composition ratio 2:1 at the surface. The fit of this ternary set of data is not as good as for the C₁₂E₈ set, especially at low bulk SLE₂S mole fractions. This cannot be attributed to a poor choice of parameters because the basic problem is that the observed values are higher than would be accounted for by any set of parameters, i.e. the problem probably lies in the experimental data and we return to this issue in the Discussion. A final point is that the R1–R2 interaction is an important contributor to the ternary fits in Figures 3 and 4 but the parameters are exactly the same as for the R1 and R2 data in Figure 1(c) and Figure 2, although the pH and the ionic strength are quite different. This suggests that their interaction is largely determined by packing rather than electrostatic forces.

Quinary Mixture

The set of compositions of each of the components at the air–water interface of the quinary mixture SLE₂S–LAS6–C₁₂E₈–R1–R2 and the total adsorption, all determined by NR, are given in Table 4. The use of a fixed bulk solution composition of both R1 and R2 means that the data are in a form that can be plotted on a conventional 2D plot, as shown in Figure 5, where the adsorbed mole fraction of each component is shown as a function of the total solution mole fraction of SLE₂S. In this figure the bulk mole fractions of R1 and R2 were fixed at 0.15 in (a) and at 0.1 and 0.2 respectively in (b). In both parts of the figure, the

Figure 5: Variation of adsorbed mole fraction of each component with bulk mole fraction of SLE₂S for the quinary mixture SLE₂S–LAS6–C₁₂E₈–R1–R2 containing fixed mole fractions of (a) 0.15 of each of R1 and R2 and (b) 0.1 R1 and 0.2 R2. The fits used the values given in Table 1 for all the B and C parameters. Some of the points have been laterally displaced for clarity.

Table 4: Variation in surface composition and total adsorption at an overall surfactant concentration of 2 mM for a quinary mixture of C₁₂E₈, LAS6, SLE₂S, R1 and R2. The solution compositions of R1 and R2 in the top half of the table were both fixed at a mole fraction of 0.15 and those in the bottom half at 0.1 and 0.2 respectively. The solutions were all equimolar in LAS6 and C₁₂E₈. Other measurements with LAS6 and C₁₂E₈ in varying ratios are given in Table S1 of the Supporting Information.

C ₁₂ E ₈	total mole fraction				surface mole fraction ± 0.05					$10^6 \times \Gamma_{total}$ $\pm 0.05 \text{ mol m}^{-2}$
	LAS6	SLE ₂ S	R1	R2	C ₁₂ E ₈	LAS6	SLE ₂ S	R1	R2	
0.02	0.02	0.66	0.15	0.15	0.04	0.04	0.29	0.48	0.16	3.07
0.035	0.035	0.63	0.15	0.15	0.02	0.06	0.31	0.43	0.18	3.21
0.09	0.09	0.52	0.15	0.15	0.05	0.12	0.22	0.46	0.15	3.00
0.14	0.14	0.42	0.15	0.15	0.08	0.19	0.19	0.39	0.15	3.18
0.233	0.233	0.233	0.15	0.15	0.17	0.20	0.06	0.41	0.14	3.23
0.26	0.26	0.18	0.15	0.15	0.17	0.22	0.08	0.38	0.14	3.19
0.04	0.04	0.62	0.10	0.20	0.03	0.07	0.32	0.38	0.19	3.18
0.09	0.09	0.52	0.10	0.20	0.11	0.15	0.2	0.35	0.17	3.17
0.23	0.23	0.23	0.10	0.20	0.17	0.26	0.11	0.30	0.16	3.06
0.26	0.26	0.18	0.10	0.20	0.14	0.29	0.12	0.28	0.17	3.05

maximum value of the bulk mole fraction of SLE₂S is therefore limited to 0.7. The total LAS6 + C₁₂E₈ is then also limited to a maximum of 0.7 and for the data plotted in Figure 5 LAS6 and C₁₂E₈ were also equimolar (there are some additional data given in Table S1 of the Supporting Information). The overall surfactant solution was again chosen to be 2 mM, which is well above the mixed CMC at any composition.

A quinary mixture is characterized by 10 sets of pairwise interactions, one set for the micelles and one set for adsorption, a B and a C parameter in each set and therefore 40 parameters in all. These have been determined by fitting adsorption profiles for each binary combination and for several ternary combinations for seven of the pairs in the present paper and the values are listed in Table 1. The remaining three sets, SLE₂S–LAS6, SLE₂S–C₁₂E₈, and LAS6–C₁₂E₈ have been determined by Liley et al.³⁰ using NR and their values are also included in Table 1. The iteration procedure for calculating the adsorption patterns of the different components from this set of 40 parameters has been described above and the outcome of the calculation is superimposed on the experimental data in Figure 5(a). We

emphasize that
adjusted to obtain
from the binary
found some tens
ternary data, and
this small element
reproduce the overall
in terms of the
ternary mixture.

Figure 6: The fit of three sets of interaction parameters to the quinary mixture SLE_2S – LAS6 – C_{12}E_8 – R1 – R2 containing a fixed mole fraction of 0.15 of each of R1 and R2 . (a) micellar interactions as in Table 1 and ideal adsorption, (b) micellar interactions plus the LAS6 – C_{12}E_8 , LAS6 – SLE_2S and SLE_2S – C_{12}E_8 surface interactions, (c) same as (b) but with the addition of the surface interactions for R1 – R2 , SLE_2S – R1 and SLE_2S – R2 .

This is an important result and we return to it in the Discussion section below. Here we

examine the extent to which the fitting is determined by the stronger interactions. Figure 6(a) gives the calculated lines obtained for ideal mixing at the surface. The lines are the equivalent of the thin red lines shown earlier for the binary mixtures and they have been calculated using just the micellar mixing parameters in Table 1 and the c^μ and c^π values. The latter are established directly by experiment, i.e. independently of any fitting of the adsorbed compositions, and, since the interaction parameters for micellization all happen to be generally small the calculated lines in the figure are largely determined by the c^μ and c^π . This fit to ideal adsorption evidently gives an approximate description of the surface composition, its main fault being that it slightly overestimates the fractions of R1 and R2 at the surface. However, what has been omitted from this calculation are the known surface interaction parameters for the trio of surfactants, LAS6, SLE₂S and C₁₂E₈. These have been determined by Liley et al. and are large, that between LAS6 and C₁₂E₈ being particularly large.³⁰ When these are included, as in Figure 6 (b) the fit is very poor. The fraction of R1 is now completely wrong as are those of LAS6 and C₁₂E₈. As explained earlier, when G_E for surface mixing for a pair of surfactants is large and negative they are preferentially adsorbed, displacing the other surfactants in the process. Compared with Figure 6 (a) the main effect is the strong interaction between LAS6 and C₁₂E₈, which enhances their adsorption strongly and drives R1 and R2 off the surface. SLE₂S is not significantly affected because of its own strong attractive interactions with LAS6 and C₁₂E₈. In Figure 6 (c) introduction of the strong R1-R2 interaction to the situation in (b) enhances both R1 and R2 at the expense of the other three surfactants and then the strong R1-SLE₂S interaction enhances R1 and SLE₂S at the expense of R2. The result is a large increase in R1 and SLE₂S at the expense of LAS6 and C₁₂E₈, while R2 is little changed, boosted also by its significant interaction with SLE₂S. Figure 6 (c) is reasonably close to the best fit in Figure 5 (a) except on the left hand side. The parameters still missing from (c) are the moderate interactions of LAS6 and C₁₂E₈ with R1 and R2 and these are the ones whose influence is mainly on the left hand side of the diagram. The final fit is therefore, not unexpectedly dominated by the strong

interactions in the system.

The reason for the resemblance of the ideal surface fit (Figure 6 (a)) to the best fit in Figure 5(a) is therefore a coincidental result of each of the five surfactants being part of a strongly attracting interacting pair. This can be viewed as a kind of cancellation of the attractive interactions. However, while this is how the interactions impact on the composition of the surface, it has a completely different effect on the overall adsorption and the surface tension. Eqn (17) shows that the two concentrations $c^\pi = c^\mu$ must become identical at the CMC for a single surfactant. Although this assumption does not strictly apply to the mixture, it shows that, if c^π is lowered relative to c^μ by large negative interaction parameters at the surface, then the adsorbed layer is stabilized relative to the micelle. The intersection of the two, which determines the CMC, is then lowered both in concentration and in surface tension. There is no similar gain when the surface is ideal. The synergy comes from attractive interactions not the intrinsic surface activity (c^π) of the pure components.

Discussion and Conclusions

We discuss three features of these results, (i) the implications that the electrostatic contributions to the interactions appear to be able to be treated as pairwise additive, (ii) causes of possible error in the analysis of the data, particularly the degree to which micellar and surface interactions can be separated, and (iii) the opportunities opened up by the experimental method of using adsorption above the CMC.

The PPA strictly only applies to nonionic surfactants. The original work of Rubingh and Holland sidestepped this issue by assuming that any contributions from electrostatics to the interactions may be handled as an implicit part of the overall interaction. This idea has not been adequately tested, partly because of the difficulty of studying mixtures with at least three components, which is where the issue arises, and partly because it has been assumed that the safest strategy is to screen out the electrostatics by adding electrolyte (see Nikishido for a full discussion of these issues²⁵). The difficulty of obtaining suitable information stems

partly from the unreliability often associated with the analysis of surface tension (ST) data to give compositional information, which causes authors to prefer to use the more direct ST information such as c^μ or c^π . However, as we have argued elsewhere,²⁹ the extraction of interaction parameters is more sensitive to composition than it is to values of c^μ or c^π (see also the discussion by Hoffmann and Posnecker³⁸). NR gives sufficiently accurate model independent results for the composition of binary mixtures that (a) it is relatively easy to apply a more realistic model for G_E than the regular solution approximation, and (b) it is not significantly more difficult to study mixtures with more than two components.

This and the two previous papers show that for the systems studied the pairwise addition of the binary interactions accounts for the mixing characteristics of a quinary mixture as well as for several of the associated ternary mixtures. The mixtures contained different strongly anionic surfactants, SDS, SLE₂S, LAS6. Furthermore, there is evidence that in one of these mixtures (SDS-C₁₂E₈) the anionic surfactant does indeed change its mean state of ionization on micelle formation,³⁹ and that addition of electrolyte significantly alters the interaction parameters in some of the binary and ternary mixtures that are constituents of this quinary system.³¹ The successful use of pairwise additivity suggests that electrostatic contributions to the interactions are either dominated by the Stern layer counterions or that both Stern layer counterions and diffuse layer counterions behave as though they are part of the pseudophase. The distribution of the counterions between these two states strongly affects the surface tension and the values of c^π and c^μ . For this set of systems any effect of a change of ionization on G_E evidently contributes at a level lower than the present experimental error. Because most previous studies have used the regular solution approximation and because several of the binary mixtures used here are far from behaving as regular mixtures, an important component of the present study is the higher accuracy available to the NR method.

Neutron reflection is a particularly easy measurement to make. A typical composition determination on a N-fold mixture at a single concentration takes N+1 measurements of 5

– 30 min each, the length of time being about 10 min for a binary mixture. Unlike surface tension (ST), the interpretation to obtain the surface coverage at a particular concentration is independent of measurements at other concentrations. Unlike ST it is also easy, direct and requires no assumptions. Measurements can be made equally well above or below the CMC. NR, like all techniques, is sensitive to surface active impurities but is much less vulnerable to them than ST. To a significant degree, impurity errors can be minimized by working above the CMC. The particular advantages of working above the CMC are that impurities are usually solubilized into the micelles, and that total concentration variation can be used indirectly to determine the composition of the micelles, a measurement that is well known to be otherwise very challenging.⁴⁰

Although the measurements of composition from NR above the CMC are straightforward, the analysis of these compositions in terms of the pseudophase assumption may be less so. Above the CMC adsorption depends on both the monomer composition, which is controlled by the micellar composition, and on the adsorption to the surface, i.e. the adsorption data has to be parameterized in terms of both micellar mixing and surface mixing. This is a disadvantage in that the interpretation becomes more complicated and there may be partial cancellation of the different effects of micellization and adsorption. However, since the monomer concentration varies strongly with the total concentration close to the CMC, there is the advantage that the otherwise largely inaccessible information about micellar composition can also in principle be extracted. Although the measurements described in this paper were mainly made at a single total concentration, the surface interaction has almost always dominated the overall interaction in the various binary, ternary and quinary mixtures studied. There is then little effect of cancellation errors on the surface interaction parameters, although this can significantly increase the relative error on the micellar interaction parameters. Finally, we have made the approximation that the surface tension at the mixed CMCs is constant at 33 mN m^{-1} . Eqn (17) at the mixed CMC shows that it must vary. However, the variation will be over only a small range, it will have a form that is

not easily parameterized, and tests of its variation on the fitting parameters with overall composition indicate that it tends to produce systematic effects that are inconsistent with any fit. The value of the surface tension must be partly described by Eqn (17), but it must also involve the total coverage and this is something that is not part of the PPA at present.

Acknowledgements

The provision of beam time at ISIS and the ILL, and the invaluable help and expertise of the Instrument Scientists, Max Skoda, and Arwell Hughes at ISIS, and Richard Campbell at the ILL, and the technical support is acknowledged.

Supporting Information

The supporting information includes (i) triangular concentration plots of the experimental compositions studied, (ii) a diagram of the fit of the CMCs of R1–R2, R1–LAS6 and R2–LAS6, (iii) a derivation of all the activity coefficients in a quinary mixture, and (iv) additional NR data on quinary mixtures, not analysed in the main text. The material is available free of charge via the internet at <http://pubs.acs.org>.

References

- (1) Muthusamy, K.; Gopalakrishnan, S.; Kockupappy, T.; Sivachidambaram, P. Biosurfactants: properties, commercial production and application. *Curr. Sci.* **2008**, *95*, 736–747.
- (2) Nitschke, M.; OCosta, S. G. V. A.; Contiero, J. Rhamnolipid surfactants: an update on the general aspects of these remarkable biomolecules. *Biotechnol. Prog.* **2005**, *21*, 1593–1600.

- (3) Ron, E. Z.; Rosenberg, E. Natural role of biosurfactants: mini review. *Environ. Microbiol.* **2001**, *3*, 229–236.
- (4) Banat, I. M.; Mukkar, R. S.; Cameotra, S. S. Potential commercial applications of microbial surfactants. *Appl. Microbiol. Biotechnol.* **2000**, *53*, 495–508.
- (5) Milligan, C. N. Environment applications of biosurfactants. *Environ. Pollut.* **2005**, *133*, 183–198.
- (6) Singh, P.; Cameotra, S. S. Potential applications of microbial surfactants in biomedical sciences. *Trends Biotechnol.* **2004**, *22*, 142–146.
- (7) Randhawa, K. K. S.; Rahman, P. K. S. M. Rhamnolipid biosurfactant—past, present and future scenario of the global market. *Frontiers Microbiol.* **2014**, *5*, 1–7.
- (8) Perfumo, A.; Rudden, M.; Smyth, T. J. P.; Marchant, R.; Stevenson, R. S.; Parry, N. J.; Banat, I. M. Rhamnolipids are conserved biosurfactants: implications for their biotechnological potential. *Appl. Microbiol. Biotechnol.* **2013**, *97*, 7297–7306.
- (9) Abolas, S.; Pinazo, A.; Infante, M. R.; Casals, M.; Garcia, F.; Manresa, A. Physico-chemical and antimicrobial properties of new rhamnolipids produced by *Pseudomonas aeruginosa* AT10 from soybean refinery wastes. *Langmuir* **2001**, *17*, 1367–1371.
- (10) Helvacı, S. A.; Peker, S.; Ozdemir, G. Effect of electrolytes on the surface behaviour of rhamnolipids R1 and R2. *Coll. Surf. B.* **2004**, *35*, 225–233.
- (11) Zhong, H.; Zeng, G. M.; Liu, J. X.; Xu, X. M.; Yuan, X. Z.; Fu, H. Y.; Huang, G. H.; Liu, Z.; Ding, Y. Adsorption of monorhamnolipid and dirhamnolipid from two *Pseudomonas aeruginosa* strains and the effect of cell hydrophobicity. *Appl. Microbiol. Biotechnol.* **2008**, *79*, 671–677.
- (12) Ozdemir, G.; Peker, S.; Helvacı, S. S. Effect of pH on the surface and interfacial behaviour of rhamnolipids R1 and R2. *Coll. Surf. A* **2004**, *234*, 135–143.

- (13) Guo, Y. P.; Hu, Y. Y.; Gu, R. R.; Lu, H. Characterisation and micellisation of rhamnolipidic fractions and crude extracts produced by *Pseudomonas aeruginosa* mutant MIG-N146. *J. Colloid Interface Sci.* **2009**, *331*, 356–363.
- (14) Wang, H.; Cross, C. S.; Mudalige, A.; Polt, R. L.; Pemberton, J. E. A PM-IRRAS investigation of monorhamnolipid orientation at the air-water interface. *Langmuir* **2013**, *29*, 4441–4450.
- (15) Sanchez, M.; Aranda, K. J.; Espuny, M. J.; Marques, A.; Terval, J. A.; Manresa, A.; Ortiz, A. Aggregation behaviour in a dirhamnolipid biosurfactant secreted by *Pseudomonas aeruginosa* in aqueous media. *J. Colloid Interface Sci.* **2007**, *307*, 246–253.
- (16) Dahrazma, B.; Mulligan, C. N.; Nieh, M. Effects of additives on the structure of rhamnolipid (biosurfactant): a small angle neutron scattering (SANS) study. *J. Colloid Interface Sci.* **2008**, *319*, 590–593.
- (17) Pornsunthorntawe, O.; Chavadej, S.; Rujiravanit, R. Solution properties and vesicle formation of rhamnolipid biosurfactants produced by *Pseudomonas aeruginosa* SP4. *Coll. Surf. B* **2009**, *72*, 6–15.
- (18) Song, D.; Li, Y.; Liang, S.; Wang, J. Micelle behaviours of sophorolipid / rhamnolipid binary mixed biosurfactant systems. *Coll. Surf. A* **2013**, *436*, 201–206.
- (19) Chen, M. L.; Penfold, J.; Thomas, R. K.; Smyth, T. J. P.; Perfumo, A.; Marchant, R.; Banat, I. M.; Stevenson, P.; Parry, A.; Tucker, I.; Grillo, I. Solution Self-Assembly and Adsorption at the Air-Water Interface of the Monorhamnose and Dirhamnose Rhamnolipids and Their Mixtures. *Langmuir* **2010**, *26*, 18281–18292.
- (20) Chen, M. L.; Penfold, J.; Thomas, R. K.; Smyth, T. J. P.; Perfumo, A.; Marchant, R.; Banat, I. M.; Stevenson, P.; Parry, A.; Tucker, I.; Grillo, I. Mixing Behavior of the Biosurfactant, Rhamnolipid, with a Conventional Anionic Surfactant, Sodium Dodecyl Benzene Sulfonate. *Langmuir* **2010**, *26*, 17958–17968.

- (21) Holland, P. M. Nonideal Mixed Micellar Solutions. *Adv. Colloid and Interface Sci.* **1986**, *26*, 111–129.
- (22) Rubingh, D. *Solution Chemistry of Surfactants*, Editor K. L. Mittal; Plenum Press: New York, 1979; pp 337–354.
- (23) Holland, P. M.; Rubingh, D. N. Mixed Surfactant Systems: An Overview. *ACS Symposium Series* **1992**, *501*, 2–30.
- (24) Holland, P. M. Modelling Mixed Surfactant Systems: Basic Introduction. *ACS Symposium Series* **1992**, *501*, 31–44.
- (25) Nikishido, N. In *Mixed Surfactant Systems*; Ogino, K., Abe, M., Eds.; Marcel Dekker Inc.: New York, 1993; pp 23–61.
- (26) Shiloach, A.; Blankschtein, D. Prediction of Critical Micelle Concentrations of Nonideal Ternary Surfactant Mixtures. *Langmuir* **1998**, *14*, 4105–4114.
- (27) Goldsipe, A.; Blankschtein, D. Molecular-Thermodynamic Theory of Micellization of Multicomponent Surfactant Mixtures: 1. Conventional (pH-Insensitive) Surfactants. *Langmuir* **2007**, *23*, 5942–5952.
- (28) Jackson, A. J.; Li, P. X.; Dong, C. C.; Thomas, R. K.; Penfold, J. Structure of Partially Fluorinated Surfactant Monolayers at the Air-Water Interface. *Langmuir* **2009**, *25*, 3957–3965.
- (29) Li, P. X.; Ma, K.; Thomas, R. K.; Penfold, J. Analysis of the Asymmetric Synergy in the Adsorption of Zwitterionic-Ionic Surfactant Mixtures at the Air-Water Interface below and above the Critical Micelle Concentration. *J. Phys. Chem.* **2016**, *120*, 3677–3691.
- (30) Liley, J. R.; Thomas, R. K.; Penfold, J.; Tucker, I. M.; Petkov, J. T.; Stevenson, P.; Webster, J. R. P. Surface Adsorption in Ternary Surfactant Mixtures above the Critical

- Micelle Concentration: Effects of Asymmetry on the Composition Dependence of the Excess Free Energy. *J. Phys. Chem. B* **2017**, *121*, 2825–2838.
- (31) Liley, J.; Thomas, R. K.; Penfold, J.; Tucker, I. M.; Petkov, J. T.; Stevenson, P.; Webster, J. R. P. The Impact of Electrolyte on Adsorption at the Air–Water Interface for Ternary Surfactant Mixtures above the Critical Micelle Concentration. *Langmuir* **2017**, *0*, 00–00.
- (32) Webster, J. R. P.; Langridge, S.; Dalglish, R. M.; Charlton, T. R. Reflectometry techniques on the Second Target Station at ISIS: Methods and science. *Eur. Physical J. Plus* **2011**, *126*, 112.
- (33) Campbell, R. A.; Wacklin, H. P.; Sutton, I.; Cubitt, R.; Fragneto, G. FIGARO: the new horizontal neutron reflectometer at the ILL. *Eur. Physical. J. Plus* **2011**, *126*, 107.
- (34) Redlich, O.; Kister, A. T. Algebraic Representation of Thermodynamic Properties and the Classification of Solutions. *Ind. Eng. Chem.* **1948**, *40*, 345–348.
- (35) Porter, A. W. On the Vapour Pressures of Mixtures. *Trans. Faraday Soc.* **1920**, *16*, 336–345.
- (36) Barker, J. A. Determination of Activity Coefficients from Total Pressure Measurements. *Australian J. Chem.* **2005**, *6*, 207–210.
- (37) Holland, P. M.; Rubingh, D. N. Nonideal Multicomponent Mixed Micelle Model. *J. Phys. Chem.* **1983**, *87*, 1984–1990.
- (38) Hoffmann, H.; Possnecker, G. The Mixing Behavior of Surfactants. *Langmuir* **1994**, *10*, 381–389.
- (39) Penfold, J.; Tucker, I.; Thomas, R. K.; Staples, E.; Schuermann, R. Structure of Mixed Anionic/Nonionic Surfactant Micelles: Experimental Observations Relating to the Role

of H
Phy
 (40) Iyer
 lizat
 tant

Figure 7: Table of contents graphic.