

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37

Title: Genomic and archaeological evidence suggest a dual origin of domestic dogs

Authors: Laurent A. F. Frantz^{1†*}, Victoria E. Mullin^{2†}, Maud Pionnier-Capitan^{3,4}, Ophélie Lebrasseur¹, Morgane Ollivier³, Angela Perri⁵, Anna Linderholm^{1,6}, Valeria Mattiangeli², Matthew D. Teasdale², Evangelos A. Dimopoulos^{1,7}, Anne Tresset⁴, Marilyne Duffraisse³, Finbar McCormick⁸, László Bartosiewicz⁹, Erika Gál¹⁰, Éva A. Nyerges¹⁰, Mikhail V. Sablin¹¹, Stéphanie Bréhard⁴, Marjan Mashkour⁴, Adrian Bălăşescu¹², Benjamin Gillet³, Sandrine Hughes³, Olivier Chassaing³, Christophe Hitte¹³, Jean-Denis Vigne⁴, Keith Dobney¹⁴, Catherine Hänni³, Daniel G. Bradley^{2*} and Greger Larson^{1*}

Affiliations:

¹ The Palaeogenomics & Bio-Archaeology Research Network, Research Laboratory for Archaeology and History of Art, The University of Oxford, Oxford, UK.

² Smurfit Institute of Genetics, Trinity College Dublin, Dublin, Dublin 2, Ireland.

³ CNRS/ENS de Lyon, IGFL, UMR 5242 and French National Platform of Paleogenetics, PALGENE, Ecole Normale Supérieure de Lyon, 46 allée d'Italie, 69364 Lyon Cedex 07, France / Université Grenoble Alpes, Laboratoire d'Ecologie Alpine (LECA), F-38000, Grenoble, France.

⁴ CNRS/MNHN/SUs – UMR 7209, Archéozoologie et Archéobotanique, Sociétés, Pratiques et Environnements, Département Ecologie et Gestion de la Biodiversité, 55 rue Buffon, 75005 Paris, France.

⁵ Department of Human Evolution, Max Planck Institute for Evolutionary Anthropology, 04103 Leipzig, Germany.

⁶ Department of Anthropology, Texas A&M University, College Station, TX 77843-4352, USA.

⁷ School of Biology, Aristotle University of Thessaloniki, Thessaloniki, Greece.

⁸ School of Geography, Archaeology and Palaeoecology, Queen's University Belfast, University Road, Belfast, Northern Ireland, UK.

⁹ Osteoarchaeological Research Laboratory, University of Stockholm, Stockholm, Sweden.

¹⁰ Archaeological Institute, Research Centre for the Humanities, Hungarian Academy of Sciences, Budapest, Hungary.

¹¹ Zoological Institute RAS, Universitetskaya nab. 1, 199034 Saint-Petersburg, Russia

¹² The National Museum of Romanian History, 12 Calea Victoriei, 030026 Bucharest, Romania.

¹³ Institut de Génétique et Développement de Rennes, CNRS-UMR6290, Université de Rennes 1, Rennes, France.

38 ¹⁴ Department of Archaeology, School of Geosciences, University of Aberdeen, St.
39 Mary's, Elphinstone Road, AB24 3UF, UK.

40 * Corresponding authors: Laurent A. F. Frantz – laurent.frantz@arch.ox.ac.uk; Greger
41 Larson – greger.larson@arch.ox.ac.uk; Dan G. Bradley - dbradley@tcd.ie

42 † Contributed equally

43 **Abstract:** The geographic and temporal origins of dogs remain controversial. Here, we generated
44 genetic sequences from 59 ancient dogs and a complete (28x) genome of a late Neolithic dog
45 (~4,800 calBP) from Ireland. Our analyses revealed a deep split separating modern East Asian
46 and Western Eurasian dogs. Surprisingly, the date of this divergence (~14,000-6,400 years ago)
47 occurs commensurate or several millennia after the first appearance of dogs in Europe and East
48 Asia. Additional analyses of ancient and modern mitochondrial DNA revealed a sharp
49 discontinuity in haplotype frequencies in Europe. Combined, these results suggest that dogs may
50 have been domesticated independently in Eastern and Western Eurasia from distinct wolf
51 populations. East Eurasia dogs were then possibly transported alongside people where they
52 partially replaced European Palaeolithic dogs.

53 **One Sentence Summary:** Genomics and archaeology reveal both a possible dual origin of
54 domestic dogs and a subsequent translocation of East Asian dogs into Europe.

55 **Main Text:** Dogs were the first domestic animal and the only animal domesticated prior to the
56 advent of settled agriculture (1). Despite their importance in human history, no consensus has
57 emerged with regard to their geographic and temporal origins, or whether dogs were
58 domesticated just once or independently on more than one occasion. Though several claims have
59 been made for an initial appearance of dogs in the early Upper Palaeolithic (~30,000 years ago;
60 *e.g.* 2), the first remains confidently assigned to dogs appear in Europe ~15,000 years ago and in
61 Far East Asia over 12,500 years ago (1, 3). While archaeologists remain open to the idea that
62 there was more than one geographic origin for dogs (*e.g.* (4, 5), most genetic studies have
63 concluded that dogs were likely domesticated just once (6) – disagreeing on whether this
64 occurred in Europe (7), Central Asia (8), or East Asia (9).

65 Recent palaeogenetic studies have had a tremendous impact on our understanding of
66 early human evolution (*e.g.* (10, 11)). Here we apply a similar approach to reconstruct the
67 evolutionary history of dogs. We generated 59 ancient mtDNA sequences from European dogs
68 (from 14,000 to 3,000 years ago) as well as a high coverage nuclear genome (~28x) of an ancient
69 dog ~4,800 calBP (12) from the Neolithic passage grave complex of Newgrange (*Sí an Bhrú*) in
70 Ireland. We combined our ancient sample with 80 modern publically available full genome
71 sequences and 605 modern dogs (including village dogs and 48 breeds) genotyped on the 170k
72 HD SNP array (12).

73 We first assessed characteristics of the Newgrange dog by typing SNPs associated with
74 specific phenotypic traits and by inferring its level of inbreeding, compared to other breed and
75 village dogs (12). Our results suggest that the degree of artificial selection and controlled
76 breeding during the Neolithic was similar to that observed in modern free-living dogs. In
77 addition, the Newgrange dog did not possess variants associated with modern breed-defining
78 traits including hair length or coat color. And though this dog was likely able to digest starch less
79 efficiently than modern dogs, it was more efficient than wolves (12).

80 A phylogenetic analysis, based on 170k SNPs revealed a deep split separating the modern
81 Sarloos breed from other dogs (Fig. 1a). This breed - created in the 1930s in the Netherlands -
82 involved breeding German Shepherds with captive wolves (13), thus explaining the breed's
83 topological placement. Interestingly, the second deepest split (evident on the basis of both the
84 170K SNP panel – Fig 1a - and genome-wide SNPs - Fig. S4) separates modern East Asian and
85 Western Eurasian (Europe and the Middle East) dogs. Moreover, the Newgrange dog clusters
86 tightly with Western Eurasian dogs. We used Principal Component Analysis (PCA), D-statistics
87 and *TreeMix* (12) to further test this pattern. Each of these analyses unequivocally placed the
88 Newgrange dog with modern European dogs (Figs. S5, S6, S7). These findings demonstrate that
89 the node separating the East Asian and Western Eurasian clades is older than the Newgrange
90 individual; directly radiocarbon dated to ~4,800 years ago.

91 Other nodes leading to multiple dog populations and breeds (including the basal breeds
92 (1) such as Greenland Sledge dogs or Siberian Husky; Fig. 1a) are poorly supported, suggesting
93 that these breeds likely possess mixed ancestry from both Western Eurasian and East Asian dog
94 lineages. To further assess the robustness of the deep split and those nodes associated with the
95 potentially admixed lineages, we defined Western Eurasian and East Asian “core” groups (Fig.
96 1a) supported by the strength of the node leading to each cluster (12). We then used D-statistics
97 to assess the affinity of each population to either Western Eurasian or East Asian core groups
98 (12). The results of this analysis again revealed a clear East-West geographic pattern across
99 Eurasia associated with the deep phylogenetic split (Fig. 1b). Breeds such as the Eurasier,
100 Greenland Sledge dogs and Siberian Huskies (all basal breeds from Northern regions(1)),
101 however, possess strong signatures of admixture with the East Asian core samples (Fig. S11), as
102 do populations sampled in East Asia that clustered alongside Western Eurasian dogs (e.g. Papua
103 New Guinean village dog; Fig. 1a).

104 We used the Multiple Sequentially Markovian Coalescent (*MSMC*)(12, 14) to reconstruct
105 the population history of East Asian and Western Eurasia dogs. An analysis of individual high
106 coverage genomes demonstrated a long, shared population history between the Newgrange dog
107 and modern dogs from both Western Eurasia and East Asia (Fig. S15). A reconstruction using
108 two genomes per group improved the resolution for recent time periods (Fig. 2a) and revealed a
109 bottleneck in the Western Eurasian population, following its divergence from the East Asian
110 core. A similar bottleneck observed in non-African human populations has been interpreted as a
111 signature of a migration out of Africa (15). We therefore speculate that the analogous bottleneck
112 observed in our dataset could be the result of a divergence and subsequent migration from east to
113 west; supporting suggestions drawn from recent analyses of modern dog genomes (8, 9, 16).

114 We then used *MSMC* to compute divergence times as a mean to assess the time frame of
115 the shared population history among dogs, and between dogs and wolves. To obtain reliable time
116 estimates, we used the radiocarbon age of the Newgrange dog to calibrate the mutation rate for
117 dogs (12)(Fig. S16). This resulted in a mutation rate estimate of between 0.3×10^{-8} and 0.45×10^{-8}
118 per generation - similar to that obtained with an ancient grey wolf genome (17). Using this
119 mutation rate, we calculated the divergence time between the two modern Russian wolves (18)
120 used in this study and the modern dogs to be 60,000-20,000 years ago (Fig. S17; Fig. 2b).
121 Importantly, this date should not be interpreted as a time frame for domestication, since the
122 wolves we examined may not have been closely related to the population that gave rise to dogs
123 (6).

124 These analyses also suggested that the divergence between the East Asian and Western
125 Eurasian core groups (~14,000-6,400 years ago) occurred commensurate, or several millennia
126 after the earliest known appearance of domestic dogs in both Europe (>15,000 years) and East
127 Asia (>12,500 years) (1) (Figs. S17, 2b). In addition, admixture signatures from wolves into
128 Western Eurasian dogs most likely pushed this estimated time of divergence deeper into the past
129 (12) meaning that the expected time of divergence between East and Western cores is likely
130 younger than our estimate. These results imply that indigenous populations of dogs were already
131 present in Europe and East Asia during the Palaeolithic (prior to this genomic divergence).
132 Under this hypothesis, this early indigenous dog population in Europe was replaced (at least
133 partially) by the arrival of East Eurasian dogs.

134 To investigate this potential replacement, we sequenced and analyzed 59 hyper-variable
135 mtDNA fragments from ancient dogs spread across Europe and combined those with 167
136 modern sequences (12). Each sequence was then assigned to one of four major well-supported
137 haplogroups (A-D) (19). While the majority of ancient European dogs belonged to either
138 haplogroup C or D (63% and 20%, respectively), most modern European dogs possess sequences
139 within haplogroups A and B (64 and 22% respectively) (Fig. 2c, d, e). Using simulations, we
140 showed that this finding cannot be explained by drift alone (12). Instead, this pattern arose from
141 clear turnover in the mitochondrial ancestry of European dogs, most likely as a result of an
142 arrival of East Asian dogs. This migration led to a partial replacement of ancient dog lineages in
143 Europe that were present by at least 15,000 years ago (1).

144 Though the mtDNA turnover is obvious, the nuclear signature reveals an apparent long-
145 term continuity. Assessments of ancestry in humans have demonstrated that major (nuclear)
146 turnovers can be difficult to detect without samples from the admixing population (11). A
147 genome-wide PCA analysis revealed that PC2 clearly discriminates the Newgrange dog from
148 other modern dogs (Fig. S8), suggesting that this individual possessed ancestry from an
149 unsampled population.

150 Our MSMC analysis reveals that the population split between the Newgrange dog and the
151 East Asian core (as measured by cross coalescence rate [CCR]) is older (on average) than the
152 split between modern Western Eurasian and East Asian lineages (Fig. 2b). Simulations suggest
153 that this pattern could be explained by a partial replacement model in which the Newgrange dog
154 retained a degree of ancestry from an outgroup population (Fig. S20a,b), that was different from
155 modern wolves (12). Alternatively, this pattern could also be explained by secondary gene flow
156 from Asian dogs into modern European dogs (Fig. S20c). Nevertheless, simulations show that
157 secondary gene flow has a smaller effect on CCR than the partial replacement model (Fig.
158 S20b,d). Moreover, secondary gene flow cannot explain the placement of the Newgrange dog on
159 our genome-wide PCA (Fig. S8). Overall, these observations are consistent with a scenario in
160 which the Newgrange dog retained a degree of ancestry from an ancient canid population that
161 falls outside of the variation of modern dogs, but that is also different from modern wolves. This
162 pattern also suggests that the replacement of European indigenous Palaeolithic dogs may not
163 have been complete.

164 To assess the consilience between our results and the archaeological record, we compiled
165 evidence for the earliest dog remains across Eurasia (Fig. 3a). We found that while dogs are
166 present at sites as old as 12,500 years in Eastern Eurasia (China, Kamchatka and East Siberia)
167 and 15,000 years in Western Eurasia (Europe and Near East) dog remains older than 8,000 years
168 have yet to be recovered in Central Eurasia (Fig. 3a; Table S7). Combined with our DNA

169 analyses, this observation suggests that two distinct populations of dogs were present in Eastern
170 and Western Eurasia during the Palaeolithic.

171 The establishment of these populations is consistent with two scenarios: a single origin of
172 Eurasian dogs followed by early transportation, founder effects, isolation and drift, or two
173 independent domestication processes on either side of Eurasia. In the first scenario, the
174 archaeological record should reveal a temporal cline of the first appearance of dogs across
175 Eurasia stemming from a single source. Given the current lack of dog remains prior to 8,000
176 years ago in Central Eurasia, a scenario involving a single origin followed by an early
177 transportation seems less likely.

178 Given our combined results, we suggest the following hypothesis: two genetically
179 differentiated and potentially extinct wolf populations in Eastern (8, 9) and Western Eurasia (7)
180 may have been independently domesticated prior to the advent of settled agriculture (Fig. 3a).
181 The eastern dog population then dispersed westward alongside humans, between 6,400 and
182 14,000 years ago, into Western Europe (10, 11, 20) whereupon they partially replaced an
183 indigenous Palaeolithic dog population. Our hypothesis reconciles previous studies that have
184 suggested domestic dogs originated in East Asia (9, 19) and Europe (7). For numerous reasons,
185 the null hypothesis should be that individual animal species were domesticated just once (21).
186 The combined genetic and archaeological results presented here, however, suggest that dogs, like
187 pigs (22), may have been domesticated twice. Nevertheless, given the complexity of the
188 evolutionary history of dogs and uncertainties related to mutation rates, generation times and the
189 incomplete nature of the archaeological record, our scenario remains hypothetical. Genome
190 sequences derived from ancient Eurasian dogs and wolves will provide the necessary means to
191 assess whether dog domestication occurred more than once.

192 **References and Notes:**

- 193 1. G. Larson *et al.*, *Proc. Natl. Acad. Sci.* **109**, 8878–83 (2012).
- 194 2. M. Germonpré, M. Lázničková-Galetová, M. V. Sablin, *J. Archaeol. Sci.* **39**, 184–202
195 (2012).
- 196 3. M. Pionnier-Capitan *et al.*, *J. Archaeol. Sci.* **38**, 2123–2140 (2011).
- 197 4. T. Dayan, *J. Archaeol. Sci.* **21**, 633–640 (1994).
- 198 5. M. Ollivier *et al.*, *PLoS One.* **8**, e75110 (2013).
- 199 6. A. H. Freedman *et al.*, *PLoS Genet.* **10**, e1004016 (2014).
- 200 7. O. Thalmann *et al.*, *Science.* **342**, 871–4 (2013).
- 201 8. L. M. Shannon *et al.*, *Proc. Natl. Acad. Sci. U. S. A.* **112**, 13639–13644 (2015).
- 202 9. G.-D. Wang *et al.*, *Cell Res.* **26**, 21–33 (2015).
- 203 10. M. E. Allentoft *et al.*, *Nature.* **522**, 167–172 (2015).
- 204 11. W. Haak *et al.*, *Nature.* **522**, 207–211 (2015).
- 205 12. Supplementary Material.
- 206 13. D. Morris, *Dogs: The Ultimate Dictionary of Over 1,000 Dog Breeds* (Trafalgar Square,
207 2008).

- 208 14. S. Schiffels, R. Durbin, *Nat. Genet.* **46**, 919–25 (2014).
- 209 15. H. Li, R. Durbin, *Nature.* **475**, 493–6 (2011).
- 210 16. M. Pilot *et al.*, *Proc. Biol. Sci.* **282**, 20152189– (2015).
- 211 17. P. Skoglund, E. Ersmark, E. Palkopoulou, L. Dalén, *Curr. Biol.* **25**, 1515–1519 (2015).
- 212 18. G. Wang *et al.*, *Nat. Commun.* **4**, 1860 (2013).
- 213 19. P. Savolainen, Y. Zhang, J. Luo, J. Lundeberg, T. Leitner, *Science.* **298**, 1610–3 (2002).
- 214 20. L. M. Cassidy *et al.*, *Proc. Natl. Acad. Sci.* **113**, 201518445 (2015).
- 215 21. G. Larson, J. Burger, *Trends Genet.* **29**, 197–205 (2013).
- 216 22. L. A. F. Frantz *et al.*, *Nat. Genet.* **47**, 1141–1148 (2015).
- 217 23. C. Gamba *et al.*, *Nat. Commun.* **5**, 5257 (2014).
- 218 24. L. H. van Wijngaarden-Bakker, *Proc. R. Irish Acad. Sect. C Archaeol. Celt. Stud. Hist.*
219 *Linguist. Lit.* **86C**, 17–111 (1986).
- 220 25. M. J. O’Kelly, R. M. Cleary, D. Lehan, *Newgrange, Co. Meath, Ireland: The Late*
221 *Neolithic/Beaker Period Settlement* (B.A.R., 1983).
- 222 26. F. McCormick, in *The Holocene History of the European Vertebrate Fauna: Modern*
223 *Aspects of Research*, N. Benecke, Ed. (1998).
- 224 27. P. Reimer, *Radiocarbon.* **55**, 1869–1887 (2013).
- 225 28. S. Pääbo *et al.*, *Annu. Rev. Genet.* **38**, 645–679 (2004).
- 226 29. L. Orlando *et al.*, *Genome Res.* **21**, 1705–19 (2011).
- 227 30. J. T. Vilstrup *et al.*, *PLoS One.* **8**, e55950 (2013).
- 228 31. M. Meyer, M. Kircher, *Cold Spring Harb. Protoc.* **2010**, pdb.prot5448 (2010).
- 229 32. M. Martin, *EMBnet.journal.* **17**, 10 (2011).
- 230 33. H. Li, R. Durbin, *Bioinformatics.* **25**, 1754–60 (2009).
- 231 34. K. Lindblad-Toh *et al.*, *Nature.* **438**, 803–19 (2005).
- 232 35. M. Schubert *et al.*, *Nat. Protoc.* **9**, 1056–82 (2014).
- 233 36. H. Li *et al.*, *Bioinformatics.* **25**, 2078-9 (2009).
- 234 37. A. McKenna *et al.*, *Genome Res.* **20**, 1297–1303 (2010).
- 235 38. K. Okonechnikov, A. Conesa, F. García-Alcalde, *Bioinformatics.* **32**, 292-4 (2015).
- 236 39. H. Jónsson, A. Ginolhac, M. Schubert, P. L. F. Johnson, L. Orlando, *Bioinformatics.* **29**,
237 1682–4 (2013).
- 238 40. B. Bai *et al.*, *Nucleic Acids Res.* **43**, 777-83 (2014).
- 239 41. A. Auton *et al.*, *Science.* **336**, 193-8 (2012).
- 240 42. A. R. Quinlan, I. M. Hall, *Bioinformatics.* **26**, 841–2 (2010).
- 241 43. E. Han, J. S. Sinsheimer, J. Novembre, *Mol. Biol. Evol.* **31**, 723–35 (2014).

- 242 44. A. Vaysse *et al.*, *PLoS Genet.* **7**, e1002316 (2011).
- 243 45. M. L. Speir *et al.*, *Nucleic Acids Res.*, **44** 717–25 (2016).
- 244 46. S. Purcell *et al.*, *Am. J. Hum. Genet.* **81**, 559–75 (2007).
- 245 47. E. Paradis, J. Claude, K. Strimmer, *Bioinformatics.* **20**, 289–290 (2004).
- 246 48. N. Patterson, A. L. Price, D. Reich, *PLoS Genet.* **2**, e190 (2006).
- 247 49. J. K. Pickrell, J. K. Pritchard, *PLoS Genet.* **8**, e1002967 (2012).
- 248 50. E. Y. Durand, N. Patterson, D. Reich, M. Slatkin, *Mol. Biol. Evol.* **28**, 2239–2252 (2011).
- 249 51. N. Patterson *et al.*, *Genetics.* **192**, 1065–93 (2012).
- 250 52. S. R. Browning, B. L. Browning, *Am. J. Hum. Genet.* **81**, 1084–97 (2007).
- 251 53. Z. Fan *et al.*, *Genome Res.* **26**, 163–73 (2016).
- 252 54. H. Angleby, P. Savolainen, *Forensic Sci. Int.* **154**, 99–110 (2005).
- 253 55. A. Ardalan *et al.*, *Ecol. Evol.* **1**, 373–85 (2011).
- 254 56. A. R. Boyko *et al.*, *Proc. Natl. Acad. Sci. U. S. A.* **106**, 13903–8 (2009).
- 255 57. S. K. Brown *et al.*, *PLoS One.* **6**, e28496 (2011).
- 256 58. S. Castroviejo-Fisher, P. Skoglund, R. Valadez, C. Vilà, J. A. Leonard, *BMC Evol. Biol.*
257 **11**, 73 (2011).
- 258 59. V. Muñoz-Fuentes, C. T. Darimont, P. C. Paquet, J. A. Leonard, *Conserv. Genet.* **11**, 547–
259 556 (2009).
- 260 60. N. Okumura, N. Ishiguro, M. Nakano, A. Matsui, M. Sahara, *Anim. Genet.* **27**, 397–405
261 (2009).
- 262 61. M. C. R. Oskarsson *et al.*, *Proc. Biol. Sci.* **279**, 967–74 (2012).
- 263 62. J.-F. Pang *et al.*, *Mol. Biol. Evol.* **26**, 2849–64 (2009).
- 264 63. P. Savolainen, T. Leitner, A. N. Wilton, E. Matisoo-Smith, J. Lundeberg, *Proc. Natl.*
265 *Acad. Sci. U. S. A.* **101**, 12387–90 (2004).
- 266 64. K. Tsuda, Y. Kikkawa, H. Yonekawa, Y. Tanabe, *Genes Genet. Syst.* **72**, 229–38 (1997).
- 267 65. J. A. Leonard *et al.*, *Science.* **298**, 1613–6 (2002).
- 268 66. M. F. Deguilloux, J. Moquel, M. H. Pemonge, G. Colombeau, *J. Archaeol. Sci.* **36**, 513–
269 519 (2009).
- 270 67. F. Verginelli *et al.*, *Mol. Biol. Evol.* **22**, 2541–51 (2005).
- 271 68. M. Pionnier-Capitan, thesis, ENS Lyon (2010).
- 272 69. O. Lebrasseur, thesis, Durham University (2014).
- 273 70. N. Rohland, M. Hofreiter, *Nat. Protoc.* **2**, 1756–62 (2007).
- 274 71. A. Cooper, *Science (80-.)*. **289**, 1139b–1139 (2000).
- 275 72. M. T. P. Gilbert, H.-J. Bandelt, M. Hofreiter, I. Barnes, *Trends Ecol. Evol.* **20**, 541–4

- 276 (2005).
- 277 73. K. S. Kim, S. E. Lee, H. W. Jeong, J. H. Ha, *Mol. Phylogenet. Evol.* **10**, 210–20 (1998).
- 278 74. M.-S. Peng *et al.*, *Mol. Ecol. Resour.* **15**, 1238–42 (2015).
- 279 75. R. C. Edgar, *Nucleic Acids Res.* **32**, 1792–7 (2004).
- 280 76. J. Leigh, PopART (Population Analysis with Reticulate Trees) (2015).
- 281 77. R. R. Hudson, *Bioinformatics.* **18**, 337–338 (2002).
- 282 78. G. K. Chen, P. Marjoram, J. D. Wall, *Genome Res.* **19**, 136–42 (2009).
- 283 79. D. H. Alexander, J. Novembre, K. Lange, *Genome Res.* **19**, 1655–64 (2009).
- 284 80. M. Boudadi-Maligne, J.-B. Mallye, M. Langlais, C. Barshay-Szmidt, *PALEO*, 39–54
285 (2012).
- 286 81. R. J. Losey *et al.*, *PLoS One.* **8**, e63740 (2013).
- 287 82. N. Benecke, *J. Archaeol. Sci.* **14**, 31–49 (1987).
- 288 83. M. Germonpré *et al.*, *J. Archaeol. Sci.* **36**, 473–490 (2009).
- 289 84. R. Flad, *J. East Asian Archaeol.* **3**, 23–51 (2001).
- 290 85. L. Barton *et al.*, *Proc. Natl. Acad. Sci. U. S. A.* **106**, 5523–8 (2009).
- 291 86. A. Lasota-Moskalewska, K. Szymczak, M. Khudzhazarov, *Archaeol. Balt.* **11** (2009).
- 292 87. A. Razzokov. *Sarazm. Dushanbe: Institute of History, Archaeology, and Ethnography.*
293 *Dushanbe, Tajikistan: Academy of Sciences of Tajikistan A. Donish History, Archaeology*
294 *and Ethnographic Institute.* (2008).
- 295 88. H. Bocherens, M. Mashkour, D. Billiou, *Environ. Archaeol.* **33**, 253–64 (2013).
- 296 89. A. K. Kasparov, *Paléorient.* **22**, 161–167 (1996).
- 297 90. M. Frachetti, N. Benecke, *Antiquity.* **83**, 1023–1037 (2015).
- 298 91. C. Chang, N. Benecke, F. Grigoriev, A. Rosen, P. Tourtellotte, *Antiquity.* **77**, 298–312
299 (2003)
- 300 92. V. Bakhshaliev, in *The Archaeology of Nakhichevan. Ten Years of New Discoveries* (Ege
301 Yayinlari, Istanbul, 1997).
- 302 93. R. Berthon *et al.*, *Environ. Archaeol.* **18**, 191–200 (2013).
- 303 94. S. J. Pawankar and P. K. Thomas, *Bulletin of the Deccan College Research Institute* **56**,
304 363–370 (1996).
- 305 95. B. & R. Allchin, F. R. Allchin, in *The Birth of Indian Civilization: India and Pakistan*
306 *Before 500 B.C.* (1968).
- 307 96. R. V. Joshi, *Archaeol. Surv. India, New Delhi, India*, 12–16 (1961).
- 308 97. P. K. Thomas, in *The Walking larder: patterns of domestication, pastoralism, and*
309 *predation* (Routledge, 2014).
- 310 98. B. P. Sahu, *From hunters to breeders: faunal background of early India* (Anamika

- 311 Prakashan, 1988).
- 312 99. N. Bholá, G. V. S. Rao, *Animal remains from Lothal excavations* (Zoological Survey of
313 India, 1962).
- 314 100. R. B. . Sewell, B. S. Guha, in *Mohenjo-Daro and the Indus* (1931).
- 315 101. G. Stein, in *Early Animal Domestication and Its Cultural Context*, P. J. Crabtree, D. V.
316 Campana, K. Ryan, Eds. (UPenn Museum of Archaeology, 1989).
- 317 102. Telegin D.Ya., *redno-Stogivska kultura epokhi midi*. (1973).
- 318 103. K. N. Wilkinson *et al.*, *J. F. Archaeol.* **37**, 20-33 (2013).
- 319 104. R. D. Barnett, W. Watson, *Iraq.* **14**, 132 (1952).
- 320 105. M. D. Frachetti, N. Benecke, A. N. Mar'yashev, P. N. Doumani, *World Archaeol.* **42**,
321 622–646 (2010).
- 322 106. P. M. Dolukhanov, in *Hunters in transition: Mesolithic societies of temperate Eurasia and*
323 *their transition to farming* (1986).
- 324 107. G. Matyushin, *The mesolithic and neolithic in the southern Urals and central Asia*.
325 (1986).
- 326 108. N. D. Ovodov *et al.*, *PLoS One.* **6**, e22821 (2011).
- 327 109. M. Germonpré, M. Lázničková-Galetová, R. J. Losey, J. Rääkkönen, M. V. Sablin, *Quat.*
328 *Int.* **359-360**, 261–279 (2015).
- 329 110. M.-A. Garcia, *Bull. la Société préhistorique française.* **102**, 103–108 (2005).
- 330 111. M. V. Sablin, G. A. Khlopachev, *Curr. Anthropol.* **43**, 795–799 (2002).
- 331 112. A. Perri, *J. Archaeol. Sci.* **68**, 1–4 (2016).
- 332 113. S. J. Crockford, Y. V. Kuzmin, *J. Archaeol. Sci.* **39**, 2797–2801 (2012).
- 333 114. M. Boudadi-Maligne, G. Escarguel, *J. Archaeol. Sci.* **45**, 80–89 (2014).
- 334 115. A. G. Drake, M. Coquerelle, G. Colombeau, *Sci. Rep.* **5**, 8299 (2015).
- 335 116. D. P. Howrigan, M. A. Simonson, M. C. Keller, *BMC Genomics.* **12**, 460 (2011).
- 336 117. E. Axelsson *et al.*, *Nature.* **495**, 360–4 (2013).
- 337 118. E. Cadieu *et al.*, *Science.* **326**, 150–3 (2009).
- 338 119. H. G. Parker *et al.*, *Science.* **325**, 995–8 (2009).

339

340 **Acknowledgments:** Raw reads of the Newgrange dog have been deposited at the European
341 Nucleotide Archive (ENA) with project number: PRJEB13070. Mitochondrial sequences as well
342 as genotype files (in plink format) were deposited on Dryad
343 (<http://datadryad.org/review?doi=doi:10.5061/dryad.8gp06>). We would like to thank G. Wang, J.
344 Schraiber, L. Orlando, L. Dalén, R.E. Green, P. Savolainen, E. Loftus for their valuable input.
345 We are also grateful to A. Osztás and I. Zalai-Gaál R.-M. Arbogast, A. Beeching, A. Boroneant,
346 O. Lecomte, S. Madeleine, C. & D. Mordant, M. Patou-Mathis, P. Pétrequin, L. Salanova, J.
347 Schibler, A. Tsuneki, F. Valla for providing archaeological material. L.A.F.F., O.L., A.L. and

348 G.L. were supported by a European Research Council grant (ERC-2013-StG-337574-UNDEAD)
349 and Natural Environmental Research Council grants (NE/K005243/1 and NE/K003259/1).
350 L.A.F.F. was supported by a Junior Research Fellowship (Wolfson College, University of
351 Oxford). V.E.M, V.M , M.D.T. and the sequencing of the Newgrange dog genome were funded
352 by ERC Investigator grant 295729-CodeX awarded to D.G.B. We would also like to
353 acknowledge the National Museum of Ireland for providing the petrous bone of the Newgrange
354 dog and the Science Foundation Ireland Award 12/ERC/B2227 and Trinseq. AB was supported
355 by a Romanian National Authority for Scientific Research (PN-II-ID-PCE-2011-3-1015). The
356 work at ENS Lyon and at MNHN Paris was also supported by PURINA-NESTLE. The authors
357 declare no conflict of interest.

358

359 **Fig. 1: Deep split between East Asian and Western Eurasian dogs.** *a.* A neighbour-joining
360 tree (with bootstrap values) based on Identity by State (*I*₂) of 605 dogs. Red and yellow clades
361 represent the East Asian and Western Asian core groups respectively (*I*₂). *b.* A map showing the
362 location and relative proportion of ancestry (mean *D*-values) of dogs (Fig. S10). Positive values
363 (red) indicate that the population shares more derived alleles with the East Asian core while
364 negative values (yellow) indicate a closer association with the Western Eurasian core.

365 **Fig. 2: Effective population size, divergence times and mtDNA.** *a.* Effective population size
366 through time of East and Western Eurasian dogs and wolves with MSMC. *b.* Cross-coalescence
367 rate (CCR) per year for each population pair in Fig. 2a. The CCR represents the ratio of within
368 and between population coalescence rates (CR). The ratio measures the age and pace of
369 divergence between two populations. Values close to 1 indicate that both within and between CR
370 are equal meaning the two populations have not yet diverged. Values close to 0 indicate that the
371 populations have completely diverged. *c.* Bar plot representing the proportion of mtDNA
372 haplogroups at different time periods. *d.* Locations of archaeological sites with haplogroup
373 proportions. *e.* Location of modern samples with haplogroup proportions.

374 **Fig. 3: Archaeological evidence for the first appearance of dogs across Eurasia and a model**
375 **of dog domestication.** *a.* Map representing the geographic origin and age of the oldest
376 archaeological dog remains in Eurasia (*I*₂). *b.* A suggested model of dog domestication under
377 the dual origin hypothesis. An initial wolf population split into East and West Eurasian wolves
378 that were then domesticated independently before going extinct (as indicated by the † symbol).
379 The Western Eurasian dog population (European) was then partially replaced by a human-
380 mediated translocation of Asian dogs at least 6,400 years ago, a process that took place gradually
381 after the arrival of the eastern dog population.

382

383 **Supplementary Materials:**

384 Materials and Methods

385 Figs. S1-S29

386 Tables S1-S7

387 References (23-110)