

Published in final edited form as:

Diabetes. 2018 June ; 67(6): 1200–1205. doi:10.2337/db17-1268.

Circulating Fetuin-A and Risk of Type 2 Diabetes: A Mendelian Randomization Analysis

Janine Kröger^{1,2}, Karina Meidtner^{1,2}, Norbert Stefan^{2,3,4}, Marcela Guevara^{5,6,7}, Nicola D Kerrison⁸, Eva Ardanaz^{5,6,7}, Dagfinn Aune^{9,10}, Heiner Boeing¹¹, Miren Dorronsoro^{12,13,7}, Courtney Dow^{14,15,16}, Guy Fagherazzi^{14,15,16}, Paul W Franks^{17,18}, Heinz Freisling¹⁹, Marc J Gunter¹⁹, José María Huerta^{20,7}, Rudolf Kaaks²¹, Timothy J Key²², Kay Tee Khaw²³, Vittorio Krogh²⁴, Tilman Kühn²⁵, Francesca Romana Mancini^{14,26,16}, Amalia Mattiello²⁷, Peter M Nilsson¹⁷, Anja Olsen²⁸, Kim Overvad^{29,30}, Domenico Palli³¹, J. Ramón Quirós³², Olov Rolandsson¹⁸, Carlotta Sacerdote^{33,34}, Núria Sala³⁵, Elena Salamanca-Fernández^{36,37,7}, Ivonne Sluijs³⁸, Annemieke MW Spijkerman³⁹, Anne Tjønneland²⁸, Konstantinos K Tsilidis^{40,41}, Rosario Tumino^{42,43}, Yvonne T van der Schouw³⁸, Nita G Forouhi⁸, Stephen J Sharp⁸, Claudia Langenberg⁸, Elio Riboli⁴⁴, Matthias B Schulze^{1,2}, and Nicholas J Wareham⁸

¹Department of Molecular Epidemiology, German Institute of Human Nutrition Potsdam-Rehbruecke, Germany ²German Center for Diabetes Research (DZD), München-Neuherberg, Germany ³Department of Internal Medicine IV, University Hospital Tübingen, Tübingen, Germany ⁴Institute of Diabetes Research and Metabolic Diseases (IDM), Helmholtz Centre Munich at the University of Tübingen, Tübingen, Germany ⁵Navarre Public Health Institute (ISPN), Pamplona, Spain ⁶IdiSNA, Navarra Institute for Health Research, Pamplona, Spain ⁷CIBER de Epidemiología y Salud Pública (CIBERESP), Spain ⁸MRC Epidemiology Unit, University of Cambridge, Cambridge, United Kingdom ⁹Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London, London, United Kingdom ¹⁰Björknes University College, Oslo, Norway ¹¹German Institute of Human Nutrition Potsdam-Rehbruecke, Germany ¹²Public Health Division of Gipuzkoa, San Sebastian, Spain ¹³Instituto BIO-Donostia, Basque Government, San Sebastian, Spain ¹⁴INSERM U1018, Center for Research in Epidemiology and Population Health, Villejuif, France ¹⁵Paris South-Paris Saclay University, Villejuif, France ¹⁶Gustave Roussy Institute, Villejuif, France ¹⁷Lund University, Malmö, Sweden ¹⁸Umeå University, Umeå, Sweden ¹⁹Section of Nutrition and Metabolism, International Agency for Research on Cancer (IARC-WHO), Lyon, France ²⁰Department of Epidemiology, Murcia Regional Health Council, IMIB-Arrixaca, Murcia,

Corresponding author: Matthias B. Schulze, DrPH, German Institute of Human Nutrition Potsdam-Rehbruecke, Department of Molecular Epidemiology, Arthur-Scheunert-Allee 114-116, 14558 Nuthetal, Germany, Ph: +49 (0) 33200 88 2434, Fax:+49 (0)33200 88 2437, mschulze@dife.de.

Duality of Interest. No conflicts of interest relevant to this article were reported.

Author Contributions. J.K. and M.B.S. conceptualized the project. J.K. had access to all data for this study, analyzed the data, drafted the manuscript, and takes responsibility for the manuscript contents. K.M. provided support with the statistical analysis. N.S. was responsible for the measurement of fetuin-A. All authors qualify for authorship according to American Diabetes Association criteria. They all contributed to conception and design, interpretation of data, revising the article critically for important intellectual content, and final approval of the version to be published. J.K. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Prior Presentation. Parts of this study were presented in German at the 12th Annual Meeting of the German Society for Epidemiology, Lübeck, Germany, 5-8 September 2017.

Spain ²¹Division of Cancer Epidemiology, German Cancer Research Center (DKFZ), Heidelberg, Germany ²²University of Oxford, United Kingdom ²³University of Cambridge, Cambridge, UK ²⁴Fondazione IRCCS Istituto Nazionale dei Tumori, Milano Italy ²⁵German Cancer Research Centre (DKFZ), Heidelberg, Germany ²⁶University Paris-Saclay, University Paris-Sud, Villejuif, France ²⁷Dipartimento di Medicina Clinica e Chirurgia, Federico II University, Naples, Italy ²⁸Danish Cancer Society Research Center, Copenhagen, Denmark ²⁹Department of Public Health, Section for Epidemiology, Aarhus University, Aarhus, Denmark ³⁰Aalborg University Hospital, Aalborg, Denmark ³¹Cancer Research and Prevention Institute (ISPO), Florence, Italy ³²Public Health Directorate, Asturias, Spain ³³Unit of Cancer Epidemiology, Citta' della Salute e della Scienza Hospital-University of Turin and Center for Cancer Prevention (CPO), Torino, Italy ³⁴Human Genetics Foundation (HuGeF), Torino, Italy ³⁵Unit of Nutrition and Cancer, Cancer Epidemiology Research Program, and Translational Research Laboratory, Catalan Institute of Oncology (IDIBELL), Barcelona, Spain ³⁶Andalusian School of Public Health, Granada, Spain ³⁷Instituto de Investigación Biosanitaria de Granada (Granada.ibs), Granada (Spain) ³⁸Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht University, Utrecht, the Netherlands ³⁹National Institute for Public Health and the Environment (RIVM), Bilthoven, The Netherlands ⁴⁰Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London, London, UK ⁴¹Department of Hygiene and Epidemiology, University of Ioannina School of Medicine, Ioannina, Greece ⁴²ASP Ragusa, Italy ⁴³AIRE - ONLUS, Ragusa, Italy ⁴⁴School of Public Health, Imperial College London, UK

Abstract

Fetuin-A, a hepatic-origin protein, is strongly positively associated with risk of type 2 diabetes in human observational studies, but it is unknown whether this association is causal. We aimed to study the potential causal relation of circulating fetuin-A to risk of type 2 diabetes in a Mendelian Randomization study with SNPs located in the fetuin-A-encoding *AHSG* gene. We used data from eight European countries of the prospective EPIC-InterAct case-cohort study including 10,020 incident cases. Plasma fetuin-A concentration was measured in a subset of 965 subcohort participants and 654 cases. A genetic score of the *AHSG* SNPs was strongly associated with fetuin-A (28% explained variation). Using the genetic score as instrumental variable of fetuin-A, we observed no significant association of a 50 µg/ml higher fetuin-A concentration with diabetes risk (HR 1.02 [95%-CI 0.97, 1.07]). Combining our results with those from the Diabetes Genetics Replication And Meta-analysis (DIAGRAM) consortium (12,171 cases) also did not suggest a clear significant relation of fetuin-A with diabetes risk. In conclusion, although there is mechanistical evidence for an effect of fetuin-A on insulin sensitivity and secretion, this study doesn't support a strong, relevant relationship between circulating fetuin-A and diabetes risk in the general population.

Fetuin-A is a protein produced primarily by the liver, particularly in nonalcoholic fatty liver disease (1; 2). Fetuin-A is an inhibitor of the insulin receptor at the tyrosine kinase level (3; 4) Fetuin-A knockout mice were protected against insulin resistance (5; 6) and strong direct associations between fetuin-A levels and diabetes risk have been observed in prospective observational studies (7–13).

Fetuin-A is encoded by the *AHSG* gene. Earlier studies have shown strong associations of *AHSG* SNPs with fetuin-A levels (14; 15). The Mendelian Randomization approach, which involves the use of genetic information as an instrumental variable of the exposure, can be applied to estimate the potential causal effect of fetuin-A on diabetes risk (16). So far, only one small study has applied the Mendelian randomization approach to investigate fetuin-A in relation to risk of type 2 diabetes (17). This study suggested no causal involvement of fetuin-A (17), however, it was limited by low power to detect associations.

We therefore used data of the multicenter European Prospective Investigation into Cancer and Nutrition (EPIC)-InterAct study, with more than 10,000 incident cases of type 2 diabetes, to estimate the unconfounded effect of fetuin-A on diabetes risk with a Mendelian randomization approach. To maximize power, we additionally incorporated data from the Diabetes Genetics Replication And Meta-analysis (DIAGRAM) consortium involving further 12,171 type 2 diabetes cases.

Research Design and Methods

EPIC-InterAct study

The EPIC-InterAct study is a case-cohort study nested within the prospective EPIC study (18). In brief, EPIC includes 519,978 men and women, mostly aged 35-70 years, who were recruited between 1991 and 2000 at 23 centers in ten European countries participating in EPIC (19). Each EPIC center obtained individual written informed consent and local ethics approval.

For the EPIC-InterAct study, incident cases of type 2 diabetes occurring in the EPIC cohort were ascertained and verified. All EPIC countries except Norway and Greece contributed to EPIC-InterAct ($n=455,680$). Individuals without stored blood ($n=109,625$) or without information on reported diabetes status ($n=5,821$) were excluded, leaving 340,234 participants eligible for inclusion in EPIC-InterAct (18).

Case-cohort construction and case ascertainment

A center-stratified, random subcohort of 16,835 individuals was selected. After exclusion of 548 individuals with prevalent diabetes and 133 with uncertain diabetes status, the subcohort included 16,154 individuals for analysis (18).

Details about the ascertainment of cases can be found in the supplemental material. Altogether, 12,403 verified incident cases were identified (18). In total, the EPIC-InterAct study involves 27,779 participants (16,154 subcohort members, 12,403 incident cases including 778 cases within the subcohort) (18).

Study Population for the Present Analysis

Figure 1 provides an overview of the study populations. As fetuin-A had been measured in Potsdam only, analyses involving fetuin-A were based on data of 1,593 participants from Potsdam (965 subcohort members, 654 incident cases including 26 cases within the subcohort). Instrumental variable analysis was performed in the whole of EPIC-InterAct, excluding participants whose genetic data was not usable ($n=5,287$ due to insufficient

amount of DNA, low call rate, X chromosome heterozygosity concordance with self-reported sex, outliers for heterozygosity, or lack of concordance with previous genotyping results) as well as EPIC-Potsdam participants with missing ($n=16$) or implausible ($n=97$) fetuin-A data, resulting in a dataset of 22,379 men and women (12,975 subcohort members, 10,020 incident cases). Analyses of the association of SNPs with potential confounders or mediators were performed after further exclusion of individuals ($n=3,946$) with missing values for these variables (dataset of 18,433 participants with 10,850 subcohort members and 8,108 cases).

Measurement of Fetuin-A and Other Biomarkers

Blood samples were taken at baseline. For the Potsdam participants, plasma levels of fetuin-A were measured in Tübingen, Germany, with the automatic ADVIA 1650 analyzer (Siemens Medical Solutions, Erlangen, Germany). An immunoturbidimetric method was used with specific polyclonal goat antihuman fetuin-A antibodies to human fetuin-A (BioVendor Laboratory Medicine, Modreci, Czech Republic). This method was evaluated in a side-by-side comparison with an ELISA (intra-assay CV: 3.5%, inter-assay: CV 5.4%, BioVendor) showing a correlation of $r=0.93$.

All other laboratory measures were done for all EPIC-InterAct participants by the Stichting Huisartsen Laboratorium Groep (Etten-Leur, the Netherlands, see Supplemental Material).

Genotyping, Identification of Tagging SNPs, and Construction of Genetic Scores

Details about genotyping methods can be found in the Supplemental material.

We selected five tagging SNPs of the *AHSG* gene (rs4917, rs2070635, rs2070633, rs2248690, rs4831) using HapMap 22/phaseII CEPH population data applying stringent criteria (minor allele frequency $>5\%$, pairwise $r^2 \geq 0.80$) as similarly done earlier (14). A genetic score was created by summing the number of fetuin-A-raising alleles of the single SNPs. We also constructed a weighted genetic score to account for the different strength of association between SNPs and fetuin-A (details of instrumental variables described in the Supplemental Material).

Assessment of Other Covariables

Standardized questionnaires were used at baseline to collect information on sociodemographic characteristics and lifestyle including age, education level, smoking status, occupational and leisure-time physical activity, and history of previous illness. Height, weight, and waist circumference of participants were obtained by trained staff during the baseline examination using standardized protocols (20). However, for participants from France and some participants from Oxford (UK), self-reported anthropometric data were collected.

Statistical Analysis

The associations of SNPs with fetuin-A were determined by linear regression in the Potsdam subcohort with SNPs modeled per fetuin-A-increasing allele (additive model). First, the individual SNPs were analyzed in separate models. Then, all SNPs were included in the

same model. SNPs showing a significant association with fetuin-A in this model were used to generate the genetic scores, with beta coefficients used as weights.

The associations of the genetic score with potential confounders or mediators were evaluated by linear regression for continuous and logistic regression for dichotomous confounders/mediators.

To study the association of genetic variables with diabetes, we performed Cox regression with Prentice weighting (21). Age was used as underlying time scale. Models were stratified by integers of age (years), study center, and genotyping source and further adjusted for sex. Heterogeneity between countries in the association of the genetic variables with diabetes risk was investigated by computing country-specific hazard ratios and pooling these with random-effects meta-analyses.

We calculated instrumental variable estimates to estimate the unconfounded effect of an increase of 50 µg/ml in fetuin-A on diabetes risk. We assumed very similar associations of *AHSG* SNPs and fetuin-A concentration between the Potsdam center and the other EPIC-InterAct centers and applied the two-stage least squares (2-SLS) procedure (22) for calculating instrumental variable estimates.

To increase power, we further included data from the DIAGRAM consortium (see Supplemental Material). Summary statistics were meta-analyzed with the OR and 95% CI of the respective SNP derived from EPIC-InterAct (after excluding the Norfolk center because this is included in DIAGRAM) using a fixed-effect model.

Statistical analyses were performed with SAS (version 9.4, Enterprise Guide 6.1; SAS Institute, Cary, NC, USA).

Furthermore, we performed a power calculation using the online tool mRnd (<http://glimmer.rstudio.com/kn3in/mRnd/>)(23), as described in the Supplemental Material.

Results

Table 1 shows baseline characteristics for the subcohort members of the EPIC-InterAct study and of the Potsdam center.

Observational analysis of fetuin-A and diabetes risk

In EPIC-Potsdam, circulating fetuin-A was positively associated with diabetes, with a HR of 1.18 (95% CI: 1.05, 1.33) per 50 µg/ml in the multivariable-adjusted model (Supplemental Table 4). Additional adjustment for BMI and waist circumference did not alter the observed association.

Association of *AHSG* SNPs and genetic scores with fetuin-A and diabetes risk

In the Potsdam subcohort, four of the five *AHSG* SNPs were significantly associated with fetuin-A in univariate analyses (Supplemental Table 1). Including all five SNPs simultaneously in one model revealed significant associations for three SNPs (Table 2), which were subsequently used to create the genetic scores. Both the weighted (weighted by

beta coefficients from linear regression) and the unweighted genetic score showed a strong association with fetuin-A explaining 28% and 27% of variation in fetuin-A, respectively. Both genetic scores were not associated with diabetes risk (Supplemental Figures 1, 2).

We detected no meaningful association of the weighted genetic score with potential confounders or mediators in the EPIC-InterAct subcohort (Supplemental Tables 2, 3).

Instrumental variable analysis of fetuin-A and diabetes

Using instrumental variable analysis to estimate the unconfounded association for a 50µg/ml higher fetuin-A level did not indicate an effect on diabetes risk in EPIC-InterAct. The HR for the weighted genetic score was 1.02 (95% CI 0.97, 1.07). Also for single SNPs, the HRs were generally very low in magnitude (between 1.01 and 1.04) and nonsignificant (Table 3). When further combining EPIC-InterAct with DIAGRAM, the ORs for the single SNPs remained the same, but ORs for two SNPs became borderline significant (OR for rs4917=1.02 [95% CI 1.00, 1.06], OR for rs2248690=1.04 [95% CI 1.00, 1.08]).

We also analyzed potential differences in the association of the weighted genetic score with diabetes risk in EPIC-InterAct. Stratification by age, sex, waist circumference, or HbA_{1c} did not reveal meaningful differences between subgroups (Supplemental Table 4).

Discussion

In this large European case-cohort study, instrumental variables reflecting circulating fetuin-A were not significantly associated with diabetes risk. Further meta-analyzing results with those from the DIAGRAM consortium to bolster statistical power also did not indicate a strong association of genetically-elevated fetuin-A with diabetes.

Several prospective observational studies have detected a significant and strong positive association between plasma fetuin-A levels and the risk of type 2 diabetes (7–13). We could not confirm this strong association in a Mendelian Randomization approach. Although we found borderline significant estimates for two SNPs (OR for rs4917=1.02 [95% CI 1.00, 1.06], OR for rs2248690=1.04 [95% CI 1.00, 1.08]), our results do not suggest a strong causal association because an allele increase corresponds to a high difference in fetuin A (50µg/ml, SD for fetuin A=61µg/ml). According to our results of the observational analysis, we would have expected an OR of 1.18. Due to the use of different assays to determine concentrations of fetuin-A across studies, we could not calculate expected ORs based on the other observational studies on fetuin-A and diabetes risk. the slightly stronger associations of AHSG SNPs with fetuin-a (17) than in EPIC-Potsdam and results of quantile comparisons (7–9; 11; 12) suggest even higher expected ORs based on these studies.

Our study was not designed to unravel specific factors that explain the results of earlier observational studies, but it suggests that fetuin-A concentration more strongly reflects an adverse metabolic profile than being an important causal risk factor in the pathogenesis of diabetes.

Our results are in agreement with those from an earlier small study on *AHSG* SNPs and diabetes risk involving older adults (17). Fetuin-A was positively associated with fasting

glucose levels, but *AHSG* SNPs did not support an association between genetically predicted fetuin-A concentrations and fasting glucose or diabetes risk. A major limitation of this study is the small study size resulting in a low power for the instrumental variable analysis (3,435 participants including 259 incident cases).

In experiments of mice and human adipocytes, it has been observed that free fatty acids (FFAs) require the presence of fetuin-A to induce an inflammatory signaling pathway resulting in insulin resistance and subclinical inflammation (24). Fetuin-A might indeed contribute to the development of type 2 diabetes in already predisposed individuals with high concentrations of FFAs. Supporting this assumption, an interaction of fetuin-A and FFAs in determining insulin sensitivity was observed in a study of 347 participants at increased risk for type 2 diabetes and CVD (25). In our study, the unavailability of FFA blood concentrations precluded us from investigating a potential effect modification by FFAs.

A major strength of our study relates to the large sample size for analyzing *AHSG* SNPs and genetic scores in relation to diabetes risk. *AHSG* SNPs and genetic scores showed strong associations with circulating fetuin-A resulting in powerful instrumental variables. All SNPs are located in the fetuin-A-encoding *AHSG* gene, making pleiotropic roles of the SNPs rather unlikely. As limitation, we have information on circulating fetuin-A only in the Potsdam center of EPIC-InterAct.

In conclusion, although there is mechanistical evidence for an effect of fetuin-A on insulin sensitivity and secretion, this study does not support a strong, relevant relationship between circulating fetuin A and diabetes risk in the general population.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank all EPIC participants and staff for their contribution to the study. We thank N. Kerrison (MRC Epidemiology Unit, Cambridge, UK) for managing the data for the EPIC-InterAct project.

Funding. Funding for the EPIC-InterAct project was provided by the European Union Sixth Framework Programme (grant number LSHM_CT_2006_037197). In addition, InterAct investigators acknowledge funding from the following agencies: MBS & KM: Supported by a Grant from the German Federal Ministry of Education and Research (BMBF) to the German Center for Diabetes Research (DZD) and the State of Brandenburg; MG: Regional Government of Navarre; EA: Health Research Fund (FIS) of the Spanish Ministry of Health and Navarre Regional Government; MG, EA, ESF: Instituto de Salud Carlos III (PIE14/00045); MD: We thank the participants of the Spanish EPIC cohort for their contribution to the study as well as to the team of trained nurses who participated in the recruitment; PWF: Swedish Research Council, Novo Nordisk, Swedish Diabetes Association, Swedish Heart-Lung Foundation; JMH: Health Research Fund of the Spanish Ministry of Health; Murcia Regional Government (Nº 6236); RK: German Cancer Aid, German Ministry of Research (BMBF); TJK: Cancer Research UK C8221/A19170 and Medical Research Council MR/M012190/1; KTK: Medical Research Council UK, Cancer Research UK; TK: German Cancer Aid, German Cancer Research Center (DKFZ), German Federal Ministry of Education and Research (BMBF); PMN: Swedish Research Council; KO: Danish Cancer Society; JRQ: Regional Government of Asturias; OR: The Västerbotten County Council; NS: Spanish Ministry of Health network RTICCC (ISCIII RD12/0036/0018) cofunded by FEDER funds /European Regional Development Fund (ERDF), "a way to build Europe"; and Generalitat de Catalunya, AGAUR 2014SGR726; IS: Dutch Ministry of Public Health, Welfare and Sports (VWS), Netherlands Cancer Registry (NKR), LK Research Funds, Dutch Prevention Funds, Dutch ZON (Zorg Onderzoek Nederland), World Cancer Research Fund (WCRF), Statistics Netherlands; Verification of diabetes cases in EPIC-NL was additionally funded by NL Agency grant IGE05012 and an Incentive Grant from

the Board of the UMC Utrecht.; AMWS: EPIC Bilthoven and Utrecht acknowledge the Dutch Ministry of Public Health, Welfare and Sports (VWS), Netherlands Cancer Registry (NKR), LK Research Funds, Dutch Prevention Funds, Dutch ZON (Zorg Onderzoek Nederland), World Cancer Research Fund (WCRF), Statistics Netherlands (The Netherlands); AT: Danish Cancer Society; RT: EPIC Ragusa acknowledges for funding SICILIAN REGIONAL GOVERNMENT and AIRE-ONLUS RAGUSA. EPIC Ragusa acknowledges for their participation blood donors of AVIS RAGUSA (local blood donors association); YTvdS: EPIC Bilthoven and Utrecht acknowledge the Dutch Ministry of Public Health, Welfare and Sports (VWS), Netherlands Cancer Registry (NKR), LK Research Funds, Dutch Prevention Funds, Dutch ZON (Zorg Onderzoek Nederland), World Cancer Research Fund (WCRF), Statistics Netherlands. In EPIC Utrecht verification of the Dutch diabetes cases was additionally funded by NL Agency (grant IGE05012) and an Incentive Grant from the Board of the UMC Utrecht.; ER: Imperial College Biomedical Research Centre; MBS: Supported by a Grant from the German Federal Ministry of Education and Research (BMBF) to the German Center for Diabetes Research (DZD) and the State of Brandenburg

References

1. Stefan N, Hennige AM, Staiger H, Machann J, Schick F, Krober SM, Machicao F, Fritsche A, Haring HU. Alpha2-Heremans-Schmid glycoprotein/fetuin-A is associated with insulin resistance and fat accumulation in the liver in humans. *Diabetes Care*. 2006; 29:853–857. [PubMed: 16567827]
2. Denecke B, Graber S, Schafer C, Heiss A, Woltje M, Jahnen-Dechent W. Tissue distribution and activity testing suggest a similar but not identical function of fetuin-B and fetuin-A. *Biochem J*. 2003; 376:135–145. [PubMed: 12943536]
3. Auberger P, Falquerho L, Contreres JO, Pages G, Le Cam G, Rossi B, Le Cam A. Characterization of a natural inhibitor of the insulin receptor tyrosine kinase: cDNA cloning, purification, and anti-mitogenic activity. *Cell*. 1989; 58:631–640. [PubMed: 2766355]
4. Rauth G, Poschke O, Fink E, Eulitz M, Tippmer S, Kellerer M, Haring HU, Nawratil P, Haasemann M, Jahnen-Dechent W, et al. The nucleotide and partial amino acid sequences of rat fetuin. Identity with the natural tyrosine kinase inhibitor of the rat insulin receptor. *European journal of biochemistry / FEBS*. 1992; 204:523–529.
5. Mathews ST, Singh GP, Ranalletta M, Cintron VJ, Qiang X, Goustin AS, Jen KL, Charron MJ, Jahnen-Dechent W, Grunberger G. Improved insulin sensitivity and resistance to weight gain in mice null for the Ahsq gene. *Diabetes*. 2002; 51:2450–2458. [PubMed: 12145157]
6. Mathews ST, Rakhade S, Zhou X, Parker GC, Coscina DV, Grunberger G. Fetuin-null mice are protected against obesity and insulin resistance associated with aging. *Biochem Biophys Res Commun*. 2006; 350:437–443. [PubMed: 17011519]
7. Ix JH, Biggs ML, Mukamal KJ, Kizer JR, Zieman SJ, Siscovick DS, Mozaffarian D, Jensen MK, Nelson L, Ruderman N, Djousse L. Association of fetuin-a with incident diabetes mellitus in community-living older adults: the cardiovascular health study. *Circulation*. 2012; 125:2316–2322. [PubMed: 22511752]
8. Ix JH, Wassel CL, Kanaya AM, Vittinghoff E, Johnson KC, Koster A, Cauley JA, Harris TB, Cummings SR, Shlipak MG. Fetuin-A and incident diabetes mellitus in older persons. *JAMA*. 2008; 300:182–188. [PubMed: 18612115]
9. Laughlin GA, Barrett-Connor E, Cummins KM, Daniels LB, Wassel CL, Ix JH. Sex-specific association of fetuin-A with type 2 diabetes in older community-dwelling adults: the Rancho Bernardo study. *Diabetes care*. 2013; 36:1994–2000. [PubMed: 23315604]
10. Stefan N, Fritsche A, Weikert C, Boeing H, Joost HG, Haring HU, Schulze MB. Plasma fetuin-A levels and the risk of type 2 diabetes. *Diabetes*. 2008; 57:2762–2767. [PubMed: 18633113]
11. Sun Q, Cornelis MC, Manson JE, Hu FB. Plasma Levels of Fetuin-A and Hepatic Enzymes and Risk of Type 2 Diabetes in Women in the U.S. *Diabetes*. 2013; 62:49–55. [PubMed: 22923470]
12. Aroner SA, Mukamal KJ, St-Jules DE, Budoff MJ, Katz R, Criqui MH, Allison MA, de Boer IH, Siscovick DS, Ix JH, Jensen MK. Fetuin-A and Risk of Diabetes Independent of Liver Fat Content: The Multi-Ethnic Study of Atherosclerosis. *Am J Epidemiol*. 2017; 185:54–64. [PubMed: 27856445]
13. Roshanzamir F, Miraghajani M, Rouhani MH, Mansourian M, Ghiasvand R, Safavi SM. The association between circulating fetuin-A levels and type 2 diabetes mellitus risk: systematic review

- and meta-analysis of observational studies. *J Endocrinol Invest.* 2018; 41:33–47. [PubMed: 28643299]
14. Fisher E, Stefan N, Saar K, Drogan D, Schulze MB, Fritsche A, Joost HG, Haring HU, Hubner N, Boeing H, Weikert C. Association of AHSR gene polymorphisms with fetuin-A plasma levels and cardiovascular diseases in the EPIC-Potsdam study. *Circ Cardiovasc Genet.* 2009; 2:607–13. [PubMed: 20031641]
 15. Jensen MK, Jensen RA, Mukamal KJ, Guo X, Yao J, Sun Q, Cornelis M, Liu Y, Chen MH, Kizer JR, Djousse L, et al. Detection of genetic loci associated with plasma fetuin-A: A meta-analysis of genome-wide association studies from the CHARGE Consortium. *Hum Mol Genet.* 2017; 26:2156–2163. [PubMed: 28379451]
 16. Davey Smith G, Ebrahim S. 'Mendelian randomization': can genetic epidemiology contribute to understanding environmental determinants of disease? *Int J Epidemiol.* 2003; 32:1–22. [PubMed: 12689998]
 17. Jensen MK, Bartz TM, Djousse L, Kizer JR, Zieman SJ, Rimm EB, Siscovick DS, Psaty BM, Ix JH, Mukamal KJ. Genetically elevated fetuin-A levels, fasting glucose levels, and risk of type 2 diabetes: the cardiovascular health study. *Diabetes Care.* 2013; 36:3121–3127. [PubMed: 23801724]
 18. Langenberg C, Sharp S, Forouhi NG, Franks PW, Schulze MB, Kerrison N, Ekelund U, Barroso I, Panico S, Tormo MJ, Spranger J, et al. Design and cohort description of the InterAct Project: an examination of the interaction of genetic and lifestyle factors on the incidence of type 2 diabetes in the EPIC Study. *Diabetologia.* 2011; 54:2272–2282. [PubMed: 21717116]
 19. Riboli E, Hunt KJ, Slimani N, Ferrari P, Norat T, Fahey M, Charrondiere UR, Hemon B, Casagrande C, Vignat J, Overvad K, et al. European Prospective Investigation into Cancer and Nutrition (EPIC): study populations and data collection. *Public Health Nutr.* 2002; 5:1113–1124. [PubMed: 12639222]
 20. Haftenberger M, Schuit AJ, Tormo MJ, Boeing H, Wareham N, Bueno-de-Mesquita HB, Kumle M, Hjartaker A, Chirlaque MD, Ardanaz E, Andren C, et al. Physical activity of subjects aged 50–64 years involved in the European Prospective Investigation into Cancer and Nutrition (EPIC). *Public Health Nutr.* 2002; 5:1163–1176. [PubMed: 12639225]
 21. Barlow WE, Ichikawa L, Rosner D, Izumi S. Analysis of case-cohort designs. *J Clin Epidemiol.* 1999; 52:1165–1172. [PubMed: 10580779]
 22. Palmer TM, Sterne JA, Harbord RM, Lawlor DA, Sheehan NA, Meng S, Granell R, Smith GD, Didelez V. Instrumental variable estimation of causal risk ratios and causal odds ratios in Mendelian randomization analyses. *Am J Epidemiol.* 2011; 173:1392–1403. [PubMed: 21555716]
 23. Brion MJ, Shakhbazov K, Visscher PM. Calculating statistical power in Mendelian randomization studies. *Int J Epidemiol.* 2013; 42:1497–1501. [PubMed: 24159078]
 24. Pal D, Dasgupta S, Kundu R, Maitra S, Das G, Mukhopadhyay S, Ray S, Majumdar SS, Bhattacharya S. Fetuin-A acts as an endogenous ligand of TLR4 to promote lipid-induced insulin resistance. *Nat Med.* 2012; 18:1279–1285. [PubMed: 22842477]
 25. Stefan N, Haring HU. Circulating fetuin-A and free fatty acids interact to predict insulin resistance in humans. *Nat Med.* 2013; 19:394–395. [PubMed: 23558619]




Figure 1.
Construction of the EPIC-InterAct case-cohort study and the study populations used for the present analysis

Table 1
Baseline characteristics of subcohort participants of the EPIC-InterAct study ($n=10,850$)
and the Potsdam center ($n=965$)

	EPIC-InterAct	Potsdam center
<i>General characteristics</i>		
Age, years	53 (9)	50 (9)
Male	37	39
BMI (kg/m^2)	26.1 (4.2)	25.7 (4.1)
Current smoking	27	19
Physically active	21	16
Longer education	21	38
Alcohol intake (g/d)	7.4 (1.2, 20.1)	8.2 (3.1, 19.5)
<i>Biomarkers</i>		
Fetuin-A, $\mu\text{g/ml}$	-	267 (61)
GGT, U/l	20 (14, 32)	21 (15, 35)
ALT, U/l	18 (14, 25)	19 (14, 27)
Albumin, g/l	46 (3)	46 (3)
Creatinine, $\mu\text{mol/l}$	68 (60, 78)	69 (61, 80)
CRP, mg/l	1.1 (0.6, 2.4)	1.1 (0.6, 2.3)
HDL cholesterol, mmol/l	1.5 (0.4)	1.5 (0.4)
Total cholesterol, mmol/l	6.0 (1.1)	5.9 (1.1)
Triglycerides, mmol/l	1.1 (0.8, 1.7)	1.1 (0.8, 1.7)
HbA _{1c} , % (mmol/mol)	5.4 [5.2, 5.7] (36 [33, 39])	5.4 [5.1, 5.5] (35 [32, 37])

Values are mean (SD), median (interquartile range), or %

Table 2
Association of individual *AHSG* SNPs and genetic scores with fetuin-A concentration in the Potsdam part of the EPIC-InterAct study subcohort – Simultaneous adjustment of SNPs

	Allele	β (SE) for fetuin concentration	<i>p</i>	F statistic
Individual SNPs				
rs4917	T→C	16.9 (8.4)	0.0446	<0.0001
rs2070635	G→A	-0.6 (5.1)	0.9013	
rs2070633	T→C	23.7 (8.2)	0.0040	
rs2248690	T→A	12.3 (4.3)	0.0040	
rs4831	G→C	1.1 (8.6)	0.9008	
Genetic scores*				
Weighted score		52.4 (2.8)	<0.0001	<0.0001
Unweighted score		53.0 (2.8)	<0.0001	<0.0001

n=965

β for individual SNPs obtained from one linear regression model including all five SNPs simultaneously, β for genetic scores obtained from single univariate linear regression models, SNPs and genetic scores modelled per fetuin-A-increasing allele and fetuin-A expressed in $\mu\text{g/ml}$

* Genetic scores were created as sum of fetuin-A-increasing alleles of rs4917, rs2070633 and rs2248690, divided by three. Betas of the three SNPs from linear regression have been used as weights for the weighted genetic score.

Table 3
Instrumental variable estimates for the weighted genetic score and single *AHSG* SNPs reflecting the association of a 50µg/ml increase in fetuin-A concentration with diabetes risk

	Weighted genetic score	rs4917	rs2070633	rs2248690
EPIC-InterAct [*]	1.02 (0.97, 1.07)	1.02 (0.98, 1.07)	1.01 (0.95, 1.08)	1.04 (0.96, 1.10)
EPIC-InterAct and DIAGRAM combined [†]		1.02 (1.00, 1.06)	1.01 (0.98, 1.05)	1.04 (1.00, 1.08)

Data are HRs (for EPIC-InterAct) or ORs (for EPIC-InterAct and DIAGRAM combined) and 95% CIs per 50µg/ml increase in fetuin-A concentration. Instrumental variable estimates were obtained from two-stage least squares procedure. Estimates for EPIC-InterAct and DIAGRAM have been combined with a fixed-effects meta-analysis.

^{*} *n*=22,379 including 12,975 incident cases of type 2 diabetes

[†] DIAGRAM: *n*=69,033 including 12,171 incident cases of type 2 diabetes