

ARTICLE


Recent progress in the total synthesis of pyrrole-containing natural products (2011-2020)

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Nidhi Singh^a, Snigdha Singh^a, Sahil Kohli^a, Aarushi Singh^a, Hannah Asiki^c, Garima Rathee^a, Ramesh Chandra^{a,b*} and Edward A. Anderson^{c*}

Natural products have long served as a rich resource for drug discovery in the treatment of illnesses and chronic diseases. Among many families exhibiting fascinating structural complexity and impressive bioactivity, pyrrole-containing natural products frequently present a significant challenge to practitioners of organic synthesis. As a result, synthetic chemists have paid intense attention to the construction of such frameworks, not least as the amounts of natural product isolated from their natural source is often far less than that needed for biological testing. This review discusses total syntheses of pyrrole-containing natural products over the last ten years, highlighting recent advances in the chemistry of pyrroles both in the context of their innate reactivity, and their preparation in complex settings.



1. Introduction
2. Total syntheses of natural products utilizing premade pyrrole motifs
 - 2.1. Premade pyrrole motifs in syntheses of natural products containing a simple pyrrolic moiety
 - 2.1.1. Brietfussins A and B
 - 2.1.2. Tambjamine K
 - 2.1.3. Zyzzyanones A-D
 - 2.1.4. Marinopyrroles A and B
 - 2.2. Premade pyrrole motifs in asymmetric syntheses of natural products containing a simple pyrrolic moiety
 - 2.2.1. Heronapyrroles A, C and D
 - 2.2.2. Roseophilin
 - 2.2.3. Nitropyrrolin A
 - 2.2.4. Mukanadin F
 - 2.2.5. Axinellamines A and B, and massadine
 - 2.3. Premade pyrrole motifs in syntheses of natural products containing a fused pyrrolic moiety
 - 2.3.1. Sanguinolentaquinone and mycenaflavin B
 - 2.3.2. Thiaplakortone A
 - 2.3.3. Aplidipiosamine A
 - 2.3.4. Lukianols A and B
 - 2.3.5. Dictyodendrins A and B
 - 2.3.6. Chlorizidine A dimethyl ether
 - 2.3.7. 2-Debromohymenol
 - 2.4. Premade pyrrole motif in asymmetric syntheses of natural products containing a fused pyrrolic moiety
 - 2.4.1. Cylindradines A and B
 - 2.4.2. Rhazinal and kopsiunnanes C1-3
 - 2.4.3. Acortatarins A and B
 - 2.4.4. Mycenarubin A
 - 2.4.5. Curvulamine
 - 2.4.6. Exiguamines A and B
 - 2.4.7. Pollenopyrroside A and capparisine B
3. Total Syntheses of natural products entailing en route construction of the pyrrole motif

^a Department of Chemistry, University of Delhi, Delhi 110007, India

^b Dr. B. R. Ambedkar Center for Biomedical Research, University of Delhi, Delhi 110007, India

^c Chemistry Research Laboratory, University of Oxford, 12 Mansfield Road, Oxford, OX1 3TA, U.K.

*Corresponding author: Prof. Edward A. Anderson

Chemistry Research Laboratory,
University of Oxford, 12 Mansfield Road,
Oxford, OX1 3TA, U.K.

Email: edward.anderson@chem.ox.ac.uk

*Corresponding author: Prof. Ramesh Chandra,

Drug Discovery & Development Laboratory,
Department of Chemistry, University of Delhi and
Dr.B.R.Ambedkar Center for Biomedical Research,
University of Delhi, Delhi- 110007, India
Email: acbrdu@hotmail.com, rameshchandragroup@gmail.com

- 3.1. En route generation of a pyrrole in syntheses of natural products containing a simple pyrrole motif
 - 3.1.1. Magnolamide and lobechine
 - 3.1.2. Pyrrolostatin
- 3.2. En route generation of a pyrrole in asymmetric syntheses of natural products containing a simple pyrrole motif
 - 3.2.1. Hemerocallisamine
- 3.3. En route generation of a pyrrole in syntheses of natural products containing a fused pyrrole motif
 - 3.3.1. Makaluvamine O, batzelline, damirone C and makaluvone
 - 3.3.2. Makaluvamine A and D, damirone B, batzelline C, makaluvone and isobatzelline C
 - 3.3.3. Spiroindimicins B and C
- 3.4. En route generation of a pyrrole in asymmetric syntheses of natural products containing a fused pyrrole motif
 - 3.4.1. Rhazinicine
 - 3.4.2. Palau'amine
 - 3.4.3. Pollenopyrroside A
 - 3.4.4. Xylapyrrosides A and B
 - 3.4.5. Cycloprodigiosin
 - 3.4.6. Marineosin A
 - 3.4.7. Cyclooroidin
 - 3.4.8. Streptorubin B
4. Conclusions
5. Conflicts of Interest
6. Acknowledgements
7. Notes and References

1. Introduction

The diverse structures and biological properties of natural products render them prime candidates for the discovery of novel lead compounds against human diseases. Among nitrogen-containing aromatic heterocycles, which are of high importance in medicinal chemistry research, the parent 5-membered heterocycle pyrrole was first extracted through the distillation of bone oil,¹ and first identified in 1834 by F.F. Runge as a constituent of coal tar.² It is a volatile, colourless liquid that darkens upon exposure to air, and has lower basicity than other nitrogen-containing compounds such as amines and pyridines due to the incorporation of its lone pair into its aromatic system. Pyrroles are well-known as biologically active scaffolds that possess a wide range of activities, and are found in an equally large number of natural products. Some of the most common natural molecules containing the pyrrole nucleus include bile pigments such as bilirubin, porphyrins of heme, porphyrinogens, chlorophyll and Vitamin B12.

Marketed drugs incorporating a pyrrole ring system exhibit diverse biological activities such as anticancer, antibacterial, anti-inflammatory, antimalarial and antipsychotic properties, among others. Prime examples include the multi-billion dollar drugs atorvastatin, zomipirac, and tolmetin.⁵ Owing to this importance of the pyrrole nucleus in medicinal chemistry, the

research community has extensively explored biological applications of naturally occurring pyrroles. Examples include the lamellarins,³ isolated from marine invertebrates, which have been found to possess anti-HIV and antitumor activities, and halitulin,⁴ a marine sponge alkaloid isolated from *Haliclona tulearenis* which showed wide range of activity against several tumor cell lines. Hundreds of marine natural products belong to the pyrrole-imidazole alkaloid family, the parent member of which is oroidin.⁵ The marinopyrroles⁶ are another important class of marine alkaloids which have gained significance due to their potent activity against methicillin-resistant bacteria. The tripyrrolic prodiginine alkaloids have been isolated from various bacteria (e.g. *Hahella chejuensis* KCTC 2396 and *Pseudoalteromonas denitrificans*) and found to display antifungal, antibacterial, antiprotozoal and antimalarial bioactivities.^{7,8} In other cases, the amounts of natural products obtained through isolation is not enough to carry out extensive biological studies, and here total synthesis can provide a solution to the supply problem, facilitating biological research.

The well known tendency of pyrrole to polymerize, especially under acidic conditions, means that its synthetic chemistry carries an intrinsic challenge – one that in many ways serves to heighten the interest of organic chemists in this fascinating class of molecule. The development of strategies for the assembly of pyrroles has been reviewed a number of times over recent years.^{9–14} The last review on the synthesis of pyrrole containing natural products was reported in 2010.¹⁵ This review offers a comprehensive coverage of endeavours towards total syntheses of natural products containing pyrroles over the last ten years. It is divided into two main sections: firstly, strategies employing premade pyrrolic moieties, and secondly those that feature en route construction of the pyrrole motif. Within each section, syntheses involving simple (monocyclic) pyrrolic moieties are presented first, followed by syntheses of natural products containing fused (polycyclic) pyrroles. This distinction is further divided into non-asymmetric and asymmetric syntheses, while within the category of fused pyrroles, we have included only those target compounds where the pyrrole heterocycle is fused with another ring in such a way that the fusion does not lead to the formation of another common category of heterocycle (e.g. indoles). Due to the space limitations of this treatise, we have focussed mainly on key transformations and strategies within the syntheses, and have also omitted synthetic work that generates only a pyrrole core rather than a complete natural product.

2. Total syntheses of natural products utilizing premade pyrrole motifs


This section discusses synthetic routes that employ a 'premade' pyrrole motif in the synthetic sequence. Initially we discuss total syntheses of natural products containing simple pyrrolic moieties, followed by natural products incorporating a fused pyrrole. As noted above, within each category the syntheses of achiral/racemic natural products are presented initially, followed by asymmetric syntheses.

2.1 Premade pyrrole motifs in syntheses of natural products containing a simple pyrrolic moiety


2.1.1 Brietfussins A and B

The halogenated natural products brietfussin A **1** and B **2** were isolated from Arctic hydrozoan *Thuiaria breitfussi*.¹⁶ In 2015, Bayer *et al.*¹⁷ described their first total syntheses (Scheme 1). The synthetic approach commenced with the reaction of readily available phenol derivative **3** with iodomethane and caesium carbonate in DMF. Subsequent treatment with DMF:DMA and pyrrolidine, and then Zn powder, afforded **4** in moderate yield. Iodination at the C3 position of indole **4** and TIPS protection furnished **5**, which on Suzuki coupling with boronic ester **6** gave **7** in good yield. **7** was treated with aqueous HCl to selectively remove the TIPS protection at C2; metalation of this position and iodination resulted in the intermediate diiodooxazole **8**.¹⁷ This underwent a further Suzuki coupling with *N*-Boc-2-pyrroleboronic acid **9** to give **10** in moderate yield; Boc removal followed by desilylation afforded brietfussin A **1**. Alternatively, bromination at the C2 position of the pyrrole in compound **10** afforded product **11**, which provided brietfussin B **2** in two further steps.

Sperry *et al.*¹⁸ developed an iridium-catalyzed indole triborylation / alkoxylation process to install the C4 methoxy substituent in a formal synthesis of brietfussin B **2** (Scheme 2). Hydrolysis of the resulting protected tryptamine **12**, followed by amide coupling with 2-(chloroacetyl)pyrrole **13**, provided amide intermediate **14**, which intercepts with a previous synthesis of brietfussin B **2** by Khan and Chen.¹⁹ In Chen's work, this was converted to **15** in two steps using DDQ and IBX, which underwent Robinson-Gabriel cyclisation ion treatment with POCl₃ to obtain **16**. This compound could be converted to the natural product via bromination. Sperry *et al.*¹⁸ synthesized the intermediate **14** in fewer steps compared to the synthesis of Chen and coworkers.¹⁹



Scheme 1: Synthesis of brietfussin A **1** and B **2** (Bayer *et al.*, 2015)¹⁷




Scheme 2: Synthesis of brietfussin B **2** (Sperry *et al.*, 2018; Khan and Chen, 2015)^{18,19}

2.1.2 Tambjamine K


Tambjamines A-J are members of a 2,2-bipyrrole family of alkaloids which differ in their aliphatic termini.²⁰⁻²⁴ The tambjamines show a broad range of bioactivities including anti-cancer, antimicrobial, and immunosuppressive properties. Tambjamine K **17** (Scheme 3), isolated from the Azorean nudibranch *Tambja Ceutae* and the subject of this section,²⁵ showed antiproliferative activity and cytotoxicity against various cancer and noncancer cell lines. Lindsley *et al.*²⁶ disclosed the first total synthesis of tambjamine K, starting from bromoenamine **18**. This underwent Suzuki coupling with Boc-1*H*-pyrrol-2-yl boronic acid **9** to deliver the Boc-protected product **19**. Acid-catalyzed condensation between **19** and *iso*-pentylamine provided tambjamine K **17** in good yield.

2.1.3 Zyzzyanones A-D

Zyzzyanones A-D (**20-23**, Scheme 4) are tetracyclic bis-pyrroloquinone alkaloids isolated from *Zyzzya fuliginosa*.²⁷⁻²⁹ the first total syntheses of which were reported by Velu *et al.*²⁹ in 2013 (Scheme 4). The zyzzyanone framework was prepared from pyrroloquinone **24**, which first underwent Mn(OAc)₃-mediated oxidative coupling with acetal **25** at 80 °C to give bis-pyrroloquinone **26**. This was further treated with iodomethane under basic conditions to afford the methylated products **27** and **28**, in 34% and 46% yield respectively. Compound **27** was treated with Pd black and ammonium formate to provide **29** in moderate yield, which upon Boc removal using TFA afforded zyzzyanone A **20** as its TFA salt. Alternatively, removal of the *N*-benzyl, *N*-tosyl, *N*-Boc and *O*-benzyl protecting groups from **28** afforded zyzzyanone B **21**. Selective Boc deprotection of **27** and **28**, followed by *N*-formylation and debenzylation afforded zyzzyanones C and D (**22, 23**) respectively.



Scheme 3: Synthesis of tamjamine K **17** (Lindsley *et al.*, 2010)²⁶

Scheme 6: Synthesis of (±)-marinopyrrole B 42 (Clive et al., 2013)³⁶

2.1.4 Marinopyrroles A and B

The antimicrobial marinopyrroles A and B were first isolated by Fenical *et al.*^{30,31} in 2008 from the marine *Streptomyces* strain CNQ-418. Their promising biological activity against MRSA strains and colon cancer cell lines make them attractive targets for drug development.^{32–34} Gulder *et al.*³⁵ reported a synthesis of marinopyrrole A **33** in 2016 (Scheme 5), which began with preparation of bipyrrrole **36** via condensation of aminopyrrole **34** with δ -ketoacetal **35**. The ester groups were converted to bis-Weinreb amide **37**, which was reacted with *ortho*-lithiated anisole **38** to give diketone **39** in 86% yield. Perchlorination of



the pyrrole rings was accomplished using NCS with a catalytic amount of iodobenzamide **40** (10 mol%), which provided *O,O*-dimethylmarinopyrrole **41** in 84% yield. Cleavage of the *O*-methyl groups under Lewis acidic conditions afforded **33**.


Clive *et al.*³⁶ described the first synthesis of marinopyrrole B (**42**, Scheme 6). Their route commenced with the commercially available trichloromethylketone **13**, which via a sequence of chlorination and bromination steps afforded bromodichloropyrrole **43**. Reaction of **43** with allylic acetate **44** gave pyrrole **45** in 95% yield, ozonolysis of which, followed by alkylation with allyl bromide, provided *O*-allyl enol ether **46**. An elegant Claisen rearrangement / ozonolysis sequence gave dicarbonyl **47**, which underwent Paal-Knorr pyrrole cyclization giving **48** and oxidation state adjustment to give the dialdehyde **49**. Double addition of *o*-methoxyphenylmagnesium bromide to this compound afforded diol **50**, which was then converted to marinopyrrole B **42** in a further five steps.

2.2 Premade pyrrole motifs in asymmetric syntheses of natural products containing a simple pyrrolic moiety

2.2.1 Heronapyrroles A, C and D

The heronapyrroles are a family of antibiotic natural products that were first isolated in 2010 from *Streptomyces* sp.³⁷


Scheme 4: Synthesis of zyzzyanone (A-D) 20-23 (Velu et al., 2013)²⁹Scheme 5: Synthesis of marinopyrrole A **33** (Gulder *et al.*, 2016)³⁵



Scheme 7: Synthesis of (+)-heronapyrrole A **51** (Morimoto *et al.*, 2015)³⁸

Morimoto *et al.*³⁸ described a total synthesis of (+)-heronapyrrole A **51** in 2015 (Scheme 7). Their strategy commenced with the preparation of epoxy bromide **54** from known (15S)-diol **52**³⁸ through a five step sequence, including a chemoselective bromoetherification to form the tetrahydrofuran ring as a protecting group for the central trisubstituted alkene. The bromopyrrole **55** was alkylated with **54** following lithiation in THF/HMPA, and upon desilylation afforded pyrrole **56**. After Boc protection of the pyrrole nitrogen atom, treatment with acetone / $\text{BF}_3 \cdot \text{OEt}_2$ effected epoxide ring opening / acetalization to give acetonide **57**. Boc deprotection, and bromoether elimination using sodium metal generated the C11-C12 trisubstituted double bond in **58**. Temporary silylation of the resulting hydroxyl group enabled pyrrole nitration; finally removal of the protecting groups yielded (+)-heronapyrrole A **51**.

Subsequent to this work, Brimble *et al.*³⁹ reported total syntheses of (+)-heronapyrrole C **60** and (–)-heronapyrrole D **59** (Scheme 8). Their strategy focused on sp^2 - sp^3 Stille coupling to attach the pyrrole motif to the tetrahydrofuran-containing side chain, and involved an early stage nitration. Iodomethyl nitropyrrole **63** was first efficiently obtained via a formylation / reduction / iodination sequence from 2-nitropyrrole **61**. The stannane partner for the planned coupling was prepared from triene **64**, which underwent a chemoselective asymmetric dihydroxylation of the terminal trisubstituted alkene **65**, followed by Shi epoxidation of the C7–C8 olefin of **65** and acid-catalyzed cyclization to tetrahydrofuran **66**. This was converted to stannane **67** using a Hunsdiecker decarboxylation to transform the carboxyl group into an alkenyl iodide, and subsequent lithiation / stannylation. The key cross-coupling of




Scheme 8: Synthesis of (+)-heronapyrrole C **60** and (–)-heronapyrrole D **59** (Brimble *et al.*, 2016)³⁹

alkenyl stannane **67** with iodomethylpyrrole **63** afforded compound **68**, deprotection of which gave (–)-heronapyrrole D **59**. A further asymmetric epoxidation / CSA-catalyzed epoxide ring-opening / cyclization afforded diastereomerically pure (+)-heronapyrrole C **60**.

2.2.2 Roseophilin

Ansa-bridged prodiginines are lipochromophores which are produced by marine and terrestrial bacteria.^{40–42} Roseophilin (**69**, Scheme 9) is a relatively new addition to this family; its structure contains two C–C σ bonds connecting its hydrocarbon core to its bis-heterocyclic tail. Harran *et al.*⁴³ reported an asymmetric synthesis of (+)-roseophilin **69** in 2013 (Scheme 9). This began with linked pyrrole-furan **70**, which was obtained by lithiation of the corresponding 3-methoxyfuran at the 2-position, followed by treatment with ZnBr_2 and Pd-catalyzed carboxylation. Activation of the carboxylic acid with benzotriazole **71** provided an intermediate benzotriazole amide, which underwent TiCl_4 -mediated acylation with 2-(8-nonenyl)pyrrole **72** to give the *bis*-heteroarylketone **73** in good yield. This pyrrole was protected by treatment with diethylchlorophosphite to provide the phosphoramidate **74**, presumably via aerobic oxidation of the intermediate *N*-phosphinyl derivative. Cross metathesis of **74** with ketone **75** using Grubbs II catalyst (5 mol%), followed by *in situ* reduction of the resultant enone through Pd-catalyzed hydrosilylation, yielded diketone **76**. The macrocycle **77** was then obtained in good yield (66%) by aldol condensation on treatment of **76** with KHMDS/18-crown-6. At this point, prochiral pyrrolaphane **77** was subjected to hydrogenation using $\text{Rh}(\text{cod})_2\text{OTf}$ (5 mol%) / Josiphos ligand **78** (5 mol%) as catalyst, to produce **79** in 67% ee




Scheme 9: Asymmetric synthesis of (+)-roseophilin **69** (Harran *et al.*, 2013)⁴³

and excellent diastereoselectivity. This was treated with 10 mol% of a rhenium tricarbonyl Lewis acid, which triggered a nucleophilic cyclization by the pyrrole onto the proximal ketone to produce an unstable 2-azafulvene intermediate. Protonation of this product with dry HCl / *t*-BuOH (25 mol%) afforded (+)-roseophilin hydrochloride salt **69**.

2.2.3 Nitropyrrolin A

Fenical and coworkers isolated nitropyrrolins A–E from a marine sediment collected from La Jolla, California in 2010.⁴⁴ Produced by the MAR4 strain CNQ-509,⁴⁴ nitropyrrolins A, B and D showed cytotoxicity towards human colon carcinoma cells. Morimoto *et al.*⁴⁵ reported the first total synthesis of these nitropyrrolins in 2016 using a strategy similar to that of their heronapyrrole syntheses described previously (Scheme 7). Brimble *et al.*⁴⁶ later described the total synthesis of (+)-nitropyrrolin A **80**, which is shown in Scheme 10. This involved a Sonagashira coupling between iodopyrrole **81** and terminal alkyne **82** to install the pyrrole C3 side chain; the key hydroxyketone motif was then constructed by a high-yielding silver-catalyzed carboxylative cyclization of **83** to give enol carbonate **84**. Cleavage of this cyclic carbonate was achieved by formation of a sulfonyl carbamate on treatment with TsNH₂, then pyridine-promoted methanolysis to give **85**. Diastereoselective reduction of **85** gave



Scheme 10: Synthesis of (+)-nitropyrrolin A **80** (Brimble *et al.*, 2017)⁴⁶


a 6:1 mixture of alcohols **86a** and **86b**, with methanolysis of the *N*-Boz (*N*-Benzyloxymethyl) group delivering (+)-nitropyrrolin A **80**.

2.2.4 Mukanadin F


Numerous linear C9-substituted bromopyrrole alkaloids have been isolated from sponges of the *Agelasida* and *Axinellida* genera. Barker *et al.*⁴⁷ reported an enantioselective synthesis of (*S*)-mukanadin F **87** (Scheme 11), which began with alcohol (*R*)-**88**.⁴⁷ This was subjected to Swern oxidation, followed by Horner-Wadsworth-Emmons reaction with phosphonate **89** to afford a diastereomeric mixture of enamides (*S*)-**90** (*E/Z* 1:2). The PMB and Boc groups in **90** were deprotected under acidic conditions to give an amine salt, which underwent coupling with trichloromethyl ketopyrrole **91** to afford the natural product **87**.

2.2.5 Axinellamines A and B and massadine

Dimeric pyrrole-imidazole alkaloids represent a topologically rich class of bromopyrrole alkaloids.⁴⁸ In 1999, Quinn *et al.*⁴⁹ reported the first members of this family, namely axinellamines A and B (**92** and **93**). A few years later, Fusetani *et al.*⁵⁰ isolated the tetracyclic pyrrole-imidazole alkaloid massadine **98**, which features a different heterocycle connectivity to the axinellamines. The densely functionalized cores display a high nitrogen content, and eight contiguous stereocenters, as well as sensitive functional groups such as halogens and hemiaminals.



Scheme 11: Enantioselective synthesis of (S)-mukanadin F **87** (Barker *et al.*, 2018)⁴⁷




Scheme 12: Enantioselective synthesis of (-)-axinellamine A **92** and (-)-axinellamine B **93** (Baran *et al.*, 2011)⁵¹

Baran *et al.*⁵¹ reported enantioselective total syntheses of axinellamines A and B, and massadine, in 2011 (Schemes 12 and 13 respectively). Among many challenges, the synthesis of the axinellamines presents two significant problems: formation of the the polycyclic core, and oxidative installation of the C20 hemiaminal. The former challenge was solved by an oxidative cyclization of the aminoimidazole **94**, the cyclopentane ring of which was originally derived from a Pauson-Khand synthesis. Introduction of the hemiaminal was achieved after a thorough screen of oxidants (e.g. halogenating agents, hypervalent iodine, potassium persulfate, potassium ferrate), with the most satisfactory results obtained by employing silver(II) picolinate **96** in aqueous trifluoroacetic acid, which delivered **97a** and **97b** in modest yield. The reduction of the azides in these compounds required conditions that could accommodate the highly polar and sensitive nature of the substrate, a challenge that was overcome by using propanedithiol and triethylamine. This

provided the corresponding diamines in sufficient purity to undergo double acylation with (trichloroacetyl)pyrrole **91**, to give (-)-axinellamine A **92** and (-)-axinellamine B **93**.

The main challenge in the approach to massadine **99** (Scheme 13) is control over the C20 oxidation state of the massadine skeleton. Use of the silver(II) picolinate salt **96** again provided the solution, enabling oxidation of **100** to **101**. Interestingly, the optimal conditions for the synthesis of the 2-aminoimidazole ring employing brine as the solvent, which reduced the extent of Cl/OH exchange at C17, and provided **102** in good yield. Oxidation of **102** to a transient epoxide with DMDO, followed by acid-promoted cyclization, gave the desired



Scheme 13: Enantioselective synthesis of (-)-massadine **99** (Baran *et al.*, 2011)⁵¹




epimer **103a** in low yield. Azide hydrogenolysis, followed by acylation of the resulting amines with pyrrole **91**, provided the natural product **98** and **99** in a 2:1 ratio.

2.3. Premade pyrrole motifs in syntheses of natural products containing a fused pyrrolic moiety

2.3.1 Sanguiolentaquinone and mycenaflavin B

Sanguiolentaquinone **104** (Scheme 14), isolated from *Mycena sanguinolenta*⁵², and mycenaflavins A and B **105** (Scheme 15), isolated from *Mycena haematopus*,⁵³ feature interesting pyrrole-*ortho*-quinone cores. Spiteller *et al.*⁵⁴ reported the first total synthesis of sanguiolentaquinone and mycenaflavin B in 2018. The approach to **104** (Scheme 14) started with bis-benzyloxyindole **106**, which was converted to quinone **109** in 6 steps. This underwent Michael addition with 3,4-dimethoxybenzyl (DMB) protected 3-aminopropanol **110**, followed by aerobic oxidation to give protected sanguiolentaquinone. The Boc group was removed under standard acidic conditions to provide the natural product **104**.

The synthesis of **105** used the same indole starting material **106**, which in this case underwent C3-alkylation with Eschenmoser's salt, and then Somei-Kametani substitution of the dimethylamino group with imine **111** to give **112**. Treatment with aqueous HCl liberated the amine group, and the side chain was introduced by reductive amination with aldehyde **113** to afford amine **114**. Hydrogenolysis of the benzyl groups, and then aerobic oxidation gave the corresponding quinone **115**. Saponification of the ethyl ester group in **115**, followed by

Scheme 14: Synthesis of sanguinolentaquinone **104** (Spiteller *et al.*, 2018)⁵⁴Scheme 15: Synthesis of mycenaflavin B **105** (Spiteller *et al.*, 2018)⁵⁴Scheme 16: Synthesis of thiaplakortone A **116** (Quinn *et al.*, 2013)⁵⁶

decarboxylation under acidic conditions and THP deprotection, afforded mycenaflavin B **105**.

2.3.2 Thiaplakortone A

Thiaplakortone A **116** (Scheme 16) was isolated from the marine sponge *Plakortis lita*.⁵⁵ Quinn *et al.*⁵⁶ described a total synthesis of this potent antimalarial natural product in 2013. From indole **117**, hydrogenolysis of the benzyl protecting group was followed by oxidation of the unstable intermediate phenol with Fremy's salt. The resulting quinone then underwent a double conjugate addition / oxidation sequence on reaction with 2-aminoethanesulfonic acid to give dihydrothiazine dioxide **118**. The dihydrothiazine ring was then oxidized, with Boc deprotection affording the HCl salt of thiaplakortone A **116**.


2.3.3 Aplidiopsamine A

The alkaloid aplidiopsamine A **119** (Scheme 17), obtained from the Australian ascidian, *Aplidiopsis confluata*,⁵⁷ exhibited significant growth inhibition of drug-resistant strains of *Plasmodium falciparum*. Takasu *et al.*⁵⁷ described a synthetic strategy to aplidiopsamine A beginning with ynamide **120**, treatment of which with triflic acid at -78 °C triggered cyclization

and NBS-mediated bromination to give **121**. Reaction of bromide **121** with di-Boc-protected adenine **122**⁵⁸ in the presence of caesium carbonate gave **123**, which on acidic deprotection afforded aplidiopsamine A **119** in high yield.

2.3.4 Lukianols A and B

Scheuer *et al.*⁵⁹ isolated lukianols A **124** (Scheme 18) and B **125** (Scheme 19) from a tunicate collected from a lagoon of the Palmyra atoll. Lukianol B **125** exhibited high h-ALR2 inhibitory activity. Iwao *et al.*⁶⁰ developed approaches to lukianol A **124** and lukianol B **125**; the key steps in the total syntheses are pyrrole lithiation, and Pd-catalyzed cross-coupling to install the phenol side chains. Thus, lithiation of **126**⁶¹ at -78 °C (Scheme

Scheme 17: Synthesis of aplidiopsamine A **119** (Takasu *et al.*, 2014)⁵⁷


18) followed by trapping with benzylchloroformate gave 2-benzylloxycarbonylpyrrole **127**. This was subjected to Suzuki coupling with *p*-isopropoxybenzylboronic acid, followed by removal of the sulfonyl protecting group to afford **128**. Alkylation of **128** with *p*-methoxyphenylacetyl bromide afforded **129**, and hydrogenolysis of the benzyl ester resulted in **130**. Finally, heating in Ac₂O followed by deprotection of the aryl isopropyl and methyl ethers with BBr₃ afforded lukianol A **124**.

The synthesis of lukianol B **125** requires differentiation of the phenol side chains and commenced with a Pd-catalyzed coupling of **126** with *p*-isopropoxyphenylboronic acid. The mono coupling product **131** was treated with LDA / benzyl chloroformate to give **132**, which underwent a second cross-coupling with *p*-methoxyphenylboronic acid to produce **133**. The resultant bis-arylated compound **133** was transformed into pyrrolooxazinone **134** in the same manner as described for the synthesis of lukianol A **124**. **134** was treated with BCl₃ to selectively deprotect the isopropyl ether, followed by double iodination of the resultant phenol with *N*-iodosaccharin (NISac) to give diiodide **136**. Finally, reaction of **136** with Zn-Cu couple, followed by demethylation with BBr₃, afforded lukianol B **125**.

aldehyde **161**. Reaction of this aldehyde with bromoketone **162**⁶⁷ under Reformatsky conditions led to β -hydroxy ketone **163** in moderate yield. The endgame consisted of a Mitsunobu cyclization to **164**, followed by reduction of the ketone, and oxidation of the upper pyrrolopyrrolidine ring using KMnO_4 to afford methyl-protected chlorozidine A **155**.

2.3.7 2-Debromohymenin

2-Debromohymenin **165** (Scheme 23) was isolated from *Stylissa carteri* (syn. *Axinella carteri*).⁶⁸ Lovely *et al.*⁶⁹ reported a racemic synthesis of **165** in 2020 using an Au-catalyzed alkyne hydroarylation. Hydroxylamine derivative **166** (Scheme 23A) was first treated with TFA to remove the Boc group, and the resulting *N*-methoxyamine was acylated with pyrrolecarbonyl chloride **167** to afford pyrrole amide **168**. Treatment of **168** with AuCl_3 in dioxane led effected cyclization to pyrroloazepinone **169** along with minor regioisomer **170**, and a small amount of trans acylated derivative **171**. The desired product **169** was then hydrogenated with Pd/C and H_2 ; subsequent lithiation with LDA and reaction with TsN_3 resulted in the synthesis of azide **172**.



Scheme 21: Synthesis of dictyodendrin B **146** (Ohno *et al.*, 2017)⁶⁵


Low yielding dibromination was achieved upon treatment of **172** with NBS (to give **173**), treatment of which with HCl

followed by Lindlar reduction, and N–O reduction with Sml_2 in THF, afforded 2-debromohymenin **165** in good yield. An alternative route from **172** to **165** involved reaction with NBS followed by HCl, and then Lindlar reduction to afford **174**; reduction of the N–O bond using $\text{Mo}(\text{CO})_6$ in acetonitrile gave 2-debromohymenin **165**.


2.4. Premade pyrrole motifs in asymmetric syntheses of natural products containing a fused pyrrolic moiety

2.4.1 Cylindradines A and B

Pyrrole-imidazole alkaloids (PIAs) are members of the wider oroidin-derived marine natural products,^{70–74} and exhibit diverse biological activities including immunosuppressive, adrenoceptor agonistic and anticancer activities. Uno *et al.*⁷⁵ isolated cylindradine A **175** (Scheme 24) and B **176** (Scheme 25) from *Axinella cylindratus*. Unlike other PIAs which consist of a 2-carbamoylpyrrole unit, cylindradines A and B contain an unusual 3-carbamoylpyrrole.¹⁴ As with other members of the family, cylindradine A **175** also features a characteristic *N,N'*-aminal in the cyclic guanidine moiety; the enantioselective




Scheme 22: Synthesis of chlorozidine A **155** (Mhaske *et al.*, 2017)⁶⁷

Scheme 23: Synthesis of (±)-2-debromohymenin 165 (Lovely *et al.*, 2020)⁶⁹

construction of this aminal is the key challenge⁷⁶⁻⁷⁷ in the construction of the cylindradine skeleton.



The first total synthesis of (+)-cylindradine A **175** (Scheme

Scheme 24: Enantioselective synthesis of (+)-cylindradine A **175** (Nagasawa *et al.*, 2014)⁷⁸

24) was disclosed by Nagasawa *et al.*⁷⁸ Their synthesis is based on an intramolecular Friedel-Crafts cyclization followed by oxidative cyclization. The synthetic approach to cylindradine A commenced with amide bond formation between pyrrolidine **177** and acid **178**⁷⁹ using EDC.HCl and DMAP, which gave **179**. This was converted into pyrrole-aldehyde **180** in four steps. Friedel-Crafts cyclization was achieved in good yield on

treatment of **180** with CSA to give **182** as the major product isomer. Stereoselective azidation of **182** with DPPA and DBU, followed by reduction under Staudinger conditions gave amine **183**. Guanidine **185** was then synthesised by reaction with 2,2,2-trichloroethoxysulfonyl (Tces) protected imine **184**,⁸⁰ which on oxidation with PIDA in the presence of MgO gave tetracyclic guanidine **186**. This product was transformed into cylindradine A **175** in a further four steps with a 39% yield.

Cylindradine B **176** (Scheme 25) features an *anti*-1,2-diol on the pyrrolidine D ring. Nagasawa *et al.*⁸¹ described the first total synthesis of (+)-cylindradine B **176**, based on an improved version of their synthesis of cylindradine A.⁷⁸ The strategy involves a diastereoselective Pictet-Spengler reaction for the

Scheme 25: Synthesis of (+)-cylindradine B **176** (Nagasawa *et al.*, 2017)⁸¹Scheme 26: Synthesis of (+)-rhazinal **197** and (+)-kopsiyunnanine C3 **196** (Gu *et al.*, 2016)⁸³


synthesis of amine **190** from pyrrole **188**, employing phosphoric acid catalyst **189**. Amine **190** was treated with H₂ and 10% Pd/C to deprotect the Cbz group, followed by reaction of the resultant pyrrolidine with **191** to give **192** in 75% yield. After desilylation, the *S*-methylisothiourea was converted into the

target guanidine motif by reaction with NH_3/MeOH and mercury(II) chloride, followed by oxidative cyclization with PhIO. A four step sequence including dibromination of the pyrrole moiety afforded (+)-cylindradine B **176**.


2.4.2 Rhazinal and kopsiyunnanines C1-3

Three rhazinilam derived alkaloids, kopsiyunnanines C1-C3, were isolated from *Yunnan Kopsia Arbores* in 2009.⁸² Kopsiyunnanines C1 and C2 (**194** and **195**, Scheme 26) contain ethoxymethyl or methoxymethyl sidechains, while C3 (**196**, Scheme 27) features the corresponding free alcohol. Gu *et al.*⁸³ presented the first asymmetric total synthesis of kopsiyunnanines C1-C3 (Schemes 26 and 27), along with (+)-rhazinal (**197**, Scheme 26). The latter two natural products were formed from an asymmetric Heck reaction of pyrrole iodide **201** (Scheme 26) under the influence of ligand **203**, and then Pd-catalyzed C3 arylation with **202**, to give fused ring pyrrole **204**. Both the nitro group and alkene of **204** were reduced with Pd/C and H_2 ; the resulting intermediate was treated with TFA, followed by addition of the Mukaiyama reagent **200** to achieve cyclization leading to the formation of (+)-rhazinal **197**. The aldehyde group in **197** was reduced with $\text{NaBH}_4/\text{EtOH}$ to afford (+)-kopsiyunnanine C3 **196**.

The same group pursued a number of strategies to



Scheme 27: Synthesis of (+)-kopsiyunnanine C1 **194** and (+)-kopsiyunnanine C2 **195** (Gu *et al.*, 2016)⁸³




Scheme 28: Stereoselective synthesis of (+)-acortatarin A **205** (Tan *et al.*, 2012)⁸⁵

synthesize (+)-kopsiyunnanines C1 and 2 from *t*-butyl ester **198** (Scheme 27); eventually it was found that the *t*-butyl group could be hydrolyzed by TFA, followed by reduction of the aldehyde with NaBH_4 and acidification with HCl to give ethers **199a** and **199b**. These were treated with the Mukaiyama reagent **200** to afford (+)-kopsiyunnanines C1 **194** and C2 **195** in good yield.


2.4.3 Acortatarins A and B

The acortatarins are spiroketal pyrrole alkaloids isolated from the roots of *Acorus tatarinowii*.⁸⁴ Tan *et al.*⁸⁵ reported an efficient synthesis of (+)-acortatarin A **205** (Scheme 28) and (–)-acortatarin B **206** (Scheme 29) through stereoselective glycal spirocyclizations. The synthesis commenced with D-thymidine **207**, which underwent TIPS protection of hydroxyl groups, followed by hydrolysis of the *N,O*-acetal with ammonium sulfate to give the corresponding TIPS protected D-ribal. This intermediate underwent C1 formylation and reduction to give TIPS-protected C1-hydroxymethyl-D-ribal which was then converted to D-ribal iodide **208** upon treatment with iodine and PPh_3 . **208** was alkylated with pyrrole dicarboxaldehyde **209** under biphasic conditions to give **210**, which in turn was reduced to alcohol **211**. Oxidative spirocyclisation promoted by Hg(II) acetate afforded the β -mercurial spiroketals, which were reduced by NaBH_4 to give spiroketals **212** and **213**. Desilylation of the mixture of **212** and **213** afforded C1-epi-acortatarin A **214** and (+)-acortatarin A **205**. In subsequent work, the same group applied an epoxidation-spirocyclisation approach for the synthesis of acortatarin B **206**, in which pyrrologlycal **210** underwent epoxidation using DMDO; treatment with Bu_4NBH_4 afforded the desired β -spiroketal **215** in good yield and diastereoselectivity at the C3 position. Desilylation provided (–)-acortatarin B **206** and its C1-epimer.

In 2017, Pale *et al.*⁸⁶ developed a different synthetic route to (+)-



Scheme 29: Stereoselective synthesis of (–)-acortatarin B **206** (Tan *et al.*, 2012)⁸⁵



Scheme 30: Zeolite catalyzed synthesis of (+)-acortatarin A **205** (Pale *et al.*, 2017)⁸⁶

acortatarin A **205** using a variety of zeolite catalysts (Scheme 30), which provide an interesting alternative to solution-based catalysts in organic synthesis. Bromoalkyne **220** was first synthesized from 2-deoxy-D-ribose **218** in six steps, two of which (glycoside formation and hydrolysis) were acid-catalyzed using the protic

zeolite H-ZSM5. The bromoalkyne itself was formed by Ramirez olefination of the intermediate **219**, followed by protection of the resulting hydroxyl group, and E2 elimination from the dibromoalkene. Its coupling partner **221** was easily obtained from available ethyl pyrrole-2-carboxylate.⁸⁷ Ullmann coupling between **220** and **221** was achieved using a CuI-zeolite catalyst to give the aryl ynamine **222**. Two-step spiroketalisation was achieved using three further zeolite-based catalysts to give **224**; a further three steps yielded (+)-acortatarin A **205**. Another asymmetric synthesis of acortatarin A has been reported by Brimble *et al.*⁸⁸ in 2014.


2.4.4 Mycenarubin A

The first total synthesis of (+)-mycenarubin A **226** (Scheme 31), isolated from *Mycena rosea*⁸⁹, was reported by Spiteller *et al.* in 2018 (Scheme 31).⁵⁴ Their strategy commenced with the synthesis of bromomethyl indole **228** from known compound **227** in three steps. **228** was combined with *N*-

(diphenylmethylene)glycine ethyl ester **229** in the presence of Corey's cinchona catalyst **230** (1.4 mol%) and CsOH.H₂O to afford product **231** in high yield and enantiomeric excess. Treatment of **231** with HCl, followed by reductive amination of the resultant aldehyde with amine **232** afforded the amine **233**. Hydrogenolysis of the benzyl groups, followed by oxidation to the pyrroloquinoline **234** using MnO₂ allowed access to (+)-mycenarubin A **226** in two further steps.


2.4.5 Curvulamine I

Tan *et al.*⁹⁰ isolated the bispyrrole alkaloid curvulamine **236** (Scheme 32) from *Curvularia* sp. IFB Z10 in 2014. Marimone *et al.*⁹¹ reported the first enantioselective synthesis of (–)-curvulamine and related alkaloids, which was completed in just ten steps. The synthesis commenced with the preparation of racemic cyanohydrin **237** and pyrrole **238**. The former was



Scheme 32: Enantioselective synthesis of (–)-curvulamine **236** (Marimone *et al.*, 2020)⁹¹

assembled by S_N2 reaction of the sodium salt of 2-methyl pyrrole **239** with methyl 2-bromopropionate **240**, followed by methyl ester reduction to yield an intermediate aldehyde, which was then transformed to cyanohydrin (±)-**237**. Aldol condensation of Boc-protected pyrrole **241** and (*E*)-4-methoxybut-3-en-2-one **242** afforded a dienone, which under microwave irradiation with 1,8-diazabicyclo[5.4.0]undec-7-ene cyclized to **238** in 60% yield. With these partners in hand, cyanohydrin **237** was treated with NaHMDS / LiCl followed by addition to heterocycle **238** to form a key congested C–C bond. The intermediate enolate was quenched with NIS to generate




Scheme 31: Synthesis of (+)-mycenarubin A **226** (Spiteller *et al.*, 2018)⁵⁴

iodide **243**. Irradiation of iodide **243** in *t*-BuOH triggered radical cyclization onto the pyrrole, affording product **244**. This was treated with excess lithiated ethyl vinyl ether **245** in THF to afford hemiketal **246**. Epimerization of this lactol was required to transform the methyl-bearing stereocentre into the desired configuration. As such, **246** was heated with NaOMe in methanol to form targeted epimer **247** in high yield. **247** was deprotonated with KHMDS, then reacted with ClCSOPh to afford the mixture of thiocarbamate epimers **248** and **249**. Radical deoxygenation of **248**, followed by hydrolysis of the enol ether afforded racemic methylketone **250**. Treatment of **250** with (*R*)-2-methyl-CBS-oxazaborolidine **251** and BH₃.DMS afforded the natural product (–)-**236** and the epimer of its enantiomer, (+)-**252**.

2.4.6. Exiguamines A and B

Exiguamine A **253** and B **254** (Scheme 33) were isolated from *Neopetrosia exigua*. Trauner *et al.*⁹² reported the total syntheses of exiguamine A **253** and exiguamine B **254**. Their strategy began with fused pyrrole cyclohexanone **255**, which was doubly brominated adjacent to the ketone, aromatized via elimination of HBr, and finally protected as its benzyl ether **256**. Desulfonation, formylation, and Henry reaction with nitromethane gave **257**, treatment of which with borane in THF resulted in the reduction of the nitrovinyl moiety. After Boc protection, **258** underwent Negishi coupling with aryl zinc **259** to give biaryl **260**, which was then debenzylated to afford 7-hydroxyindole **261**. Oxidation of **261** proceeded smoothly to yield indole *p*-quinone **262**. This intermediate was treated with an excess of *N,N*-dimethylhydantoin **263**, followed by BBr₃, to afford **264**. Reaction of this catechol with silver(II) oxide effected double cyclization to give exiguamine A **253** in moderate yield. In an attempt to improve the yield of **253**, **264**

was treated with a 20-fold excess of silver(II) oxide, which afforded exiguamine B **254** via further benzylic oxidation.



Scheme 33: Syntheses of exiguamines A **253** and B **254** (Trauner *et al.*, 2012)⁹²


2.4.7. Pollenopyrroside A and capparisine B

The pyrrole spiroketal alkaloids (PSAs) capparins A and B were isolated from powdered fruits of *Capparis spinosa*,⁹³ and pollenpyrroside A and B were collected from *Brassica campestris* pollen. The PSAs feature a pyrrole ring fused to a spiroketal. Pollenpyrroside A **265** and capparisine B **266** are closely related, being epimeric at the spiroketal carbon atom. Zhao *et al.*⁹⁴ also reported total syntheses of (+)-pollenpyrroside A **265** and (+)-capparisine B **266** (Scheme 34). The synthesis began with the known preparation of pyrrole **268**⁹⁵ from D-fructose **267**. **268** was transformed into bis-hydroxymethyl pyrrole **269** under microwave irradiation with formaldehyde. **269** was oxidized using MnO₂, followed by a series of protecting group manipulations to provide **272**. Reduction of one of the formyl groups in **272** with sodium borohydride, followed by PTSA-promoted spiroketalization, afforded compounds **273** and **274**. These underwent deoxygenation and two step debenzylation to deliver **265** and **266** respectively.

3. Total Syntheses of natural products entailing en route construction of the pyrrole motif

This section of the review discusses synthetic strategies wherein the pyrrole motif is constructed as a key part of the synthesis. As before, the syntheses of natural products containing a simple pyrrolic moiety are discussed first, followed by syntheses of

natural products incorporating a fused pyrrolic moiety. Within each category, the syntheses of achiral and racemic natural




Scheme 34: Syntheses of (+)-pollenopyrroside A **265** and (+)-capparisine B **266** (Zhao *et al.*, 2015)⁹⁴

products are presented initially, followed by asymmetric syntheses of natural products.


3.1. En route generation of a pyrrole in syntheses of natural products containing a simple pyrrole motif

3.1.1 Magnolamide and lobecheine

Magnolamide **275** and lobecheine **276** (Scheme 35) were isolated from *Magnolia coco* and *Lobelia chinensis* respectively. Brimble *et al.*⁹⁶ disclosed the first synthetic approach to lobecheine **276**, and a route to magnolamide **275**, which employed Maillard condensations to form the pyrrole rings of the natural products. The route to magnolamide **275** commenced with a condensation reaction between amine **277** and dihydropyranone **278**, prepared from furfuryl alcohol in four steps. The resulting amine salt **279** underwent coupling with succinimidyl ester **280** (which was prepared from ferulic acid) to afford magnolamide **275**. The synthesis of lobecheine **276** followed an equivalent path, using aminoester **281** as the initial coupling partner. In this case, the TBS-protected product **282** was desilylated using TFA, followed by saponification to afford lobecheine **276**.



Scheme 35: Synthesis of magnolamide **275** and lobecheine **276** (Brimble *et al.*, 2014)⁹⁶



Scheme 36: Synthesis of pyrrolostatin **283** (Knight *et al.*, 2016)⁹⁷

3.1.2 Pyrrolostatin

Pyrrolostatin **283** (Scheme 36) is another member of the relatively rare mono-pyrrole natural product class. Knight *et al.*⁹⁷ developed a novel approach to pyrrolostatin which improved on previous methods. Their route features construction of the pyrrole ring in the final step of the synthesis by a Ag(I)-catalyzed 5-endo-dig-cyclisation. The strategy commenced with double deprotonation of dithiane-carbamate **284** using *n*-BuLi / DMPU, followed by alkylation with bromide **285**. Ketone **286** was then revealed using *N*-chlorosuccinimide / silver nitrate. Nucleophilic addition of *O*-TBS-propargyl alcohol **287** to **286** produced propargylic alcohol **288** in good yield. **288** was found to be inert to silver nitrate on silica gel, potentially due to steric crowding. However, after deprotection (TBAF), a quantitative cyclization reaction was observed to afford **289**. This was converted into pyrrolostatin **283** in three further steps with 80% yield.

3.2. En route generation of a pyrrole in asymmetric syntheses of natural products containing a simple pyrrole motif

3.2.1 Hemerocallisamine


Hemerocallisamine **290** (Scheme 37) contains a 2-formylpyrrole and 4-hydroxyglutamine side chain. Brimble and coworkers⁹⁸ applied the Maillard condensation used in earlier work as described above for the synthesis of this natural product. The synthesis commenced with construction of amine **294**⁹⁹ from (4*R*)-hydroxyproline **291** which was converted to pyroglutamate **293** by various functional group protections followed by oxidation with RuO₂ (20 mol%) / NaIO₄. **293** was in turn converted to amine **294** by aminolysis with *p*-

methoxybenzylamine (PMBNH₂), followed by Boc-deprotection of the α -amino group using TFA. Maillard condensation between **294** and **278** yielded 2-formylpyrrole **295**, which was treated with PTSA followed by DDQ to give (–)-hemerocallisamine **290**.

3.3 En route generation of a pyrrole in syntheses of natural products containing a fused pyrrole motif

3.3.1 Makaluvamine O, batzelline, damirone C and makaluvone

The pyrroloquinone alkaloids makaluvamine O, isobatzellines, and damirones, which are isolated from various marine organisms, are of considerable biological interest. Spiteller *et al.*¹⁰⁰ reported total syntheses of damirone C **297**, batzelline C




Scheme 37: Synthesis of (–)-hemerocallisamine **290** (Brimble *et al.*, 2017)⁹⁸


298, batzelline D **299**, makaluvone **300**, and makaluvamine O **301** (Scheme 38). The protected indole **304** was synthesised from vanillin **302** in seven steps,¹⁰¹ the last of which involved a reductive cyclisation of the dinitro compound **303** using Fe powder / silica gel in AcOH / H₂O to form the pyrrole ring. Vilsmeier formylation of indole **304** afforded aldehyde **305** which, with or without methylation (**306**), underwent Henry reaction followed by reduction of the nitroalkene to afford tryptamines **307** and **308**. Hydrogenolysis of the benzyl groups using Pd/BaSO₄ followed by aerobic oxidation to the *ortho*-quinone triggered an intramolecular aza-Michael addition. Reoxidation with oxygen afforded damirone C **297** and its *N*-methyl derivative **309**. Selective halogenation was carried out by adding one equivalent of NCS or NBS to compounds **309** and **297** to obtain the four natural products **298–301**.

3.3.2 Makaluvamine A and D, damirone B, batzelline C, makaluvone and isobatzelline C

Tokuyama *et al.*¹⁰² also described the total syntheses of a series of these pyrroloquinoline natural products, namely makaluvamines A **310**, makaluvamine D **311**, damirone B **312**,



Scheme 38: Syntheses of damirone C **297**, batzelline C **298**, batzelline D **299**, makaluvone **300** and makaluvamine O **301** (Spiteller *et al.*, 2017)¹⁰⁰



Scheme 39: Syntheses of makaluvamine A **310**, makaluvamine D **311** and damirone B **312** (Tokuyama *et al.*, 2012)¹⁰²

isobatzelline C **313**, makaluvone **314** and batzelline C **315** (Scheme 39 and 40). The syntheses of **310–312** commenced with bromoaniline **316**, which after double allylation underwent zirconium-mediated cyclization / iodination to indoline **318**, with concurrent metalation / iodination of the arene. Chain extension of the iodomethyl group to dihydrotryptamine **319** was followed by construction of the tricyclic pyrrolo[4,3,2-de]quinoline skeleton via cyclization of the carbamate nitrogen atom onto a benzyne intermediate. This underwent aromatization and deprotection of the Boc and ethoxycarbonyl groups to give tricyclic product **320**. To access makaluvamine A **310**, indole **320** was subjected to selective methylation on the indole nitrogen, followed by treatment with salcomine, to afford an intermediate pyrroloiminoquinone. Treatment of this product with methanolic NH₄Cl gave makaluvamine A **310**.


Makaluvamine D **311** was synthesized through oxidation of

320 using Fremy's salt, then substitution of the methyl ether in product **321** by tyramine to give makaluvamine D. Alternatively, treatment of **321** with MeI and KI achieved selective methylation on the imine nitrogen atom; subsequent cleavage of the methyl ether afforded damirone B **312**.

Compound **319** could also undergo cyclization / halogenation (Scheme 40) using LiTMP at -78°C , followed by chlorination/bromination using $\text{Cl}(\text{CCl}_2)_2\text{Cl}$ or $\text{Br}(\text{CCl}_2)_2\text{Br}$. The halogenated compounds **322** and **323** were then transformed to the unprotected dihydropyrroloquinoline **324** and **325** in three steps. As before, oxidation with Fremy's salt, followed by pyrrole *N*-methylation afforded **326** and **327**. Reaction of iminoquinone **326** with $\text{NH}_4\text{Cl}/\text{EtOH}$ gave isobatzelline C **313**, while cleavage of the methyl ether with BBr_3 followed by spontaneous isomerization, led to makaluvone **314** and batzelline C **315**.


3.3.3 Spiroindimicins B and C

The natural products spiroindimicins A–D were isolated from marine actinomycete *Streptomyces* sp. SCSIO03032.¹⁰³ Spiroindimicins B–D exhibit moderate cytotoxicity against




Scheme 40: Syntheses of isobatzelline C **313**, makaluvone **314** and batzelline C **315** (Tokuyama *et al.*, 2012)


various cancer cell lines, while spiroindimicin A does not show any activity. Sperry *et al.*¹⁰⁴ reported the first total syntheses of ($\pm$)-spiroindimicins B **328** and C **329** in 2016 (Scheme 41). Beginning with iodoaniline **330**, alkylation with bromide **331** gave **332**, which underwent spirocyclization via a Heck reaction, followed by alkene hydrogenation to give intermediate **333**. A Fischer indole synthesis between **333** and 4-chlorophenylhydrazine **334** afforded **335** in good yield. After conversion to ketone **336**, treatment with excess TiCl_4 , Et_3N , and 4-methylbenzenethiol afforded dithioacetal **337**. This underwent double oxidation / sulfoxide elimination to deliver a



In 1993, Scheuer *et al.*¹⁰⁷ isolated palau'amine from the sponge *Stylotella agminate*. Baran *et al.*⁵¹ reported the first enantioselective total synthesis of palau'amine **348** in 2011 (Scheme 43). Compound **102** (see Scheme 13) was brominated by treatment with bromine and trifluoroacetic anhydride, affording **349** in moderate yield. The pyrrole precursor **350** was then attached to **349** under acidic conditions, affording **351** in 44% yield. The azide moieties in **351** were hydrogenated over palladium acetate, giving diamine **352**, which underwent cyclization mediated by EDC.HCl. Exposure of the intermediate 9-membered lactam **353** to hot TFA afforded (–)-palau'amine **348**.



Scheme 42: Synthesis of (–)-rhazinicine **342** (Tokuyama *et al.*, 2013)¹⁰⁶



Scheme 43: Enantioselective synthesis of (–)-palau'amine **348** (Baran *et al.*, 2011)⁵¹


3.4.3 Pollenopyrroside A

Brimble *et al.*¹⁰⁸ reported another application of the Maillard condensation described above in a convergent synthesis of (+)-pollenopyrroside A **265** (Scheme 44) in 2016. In this case, it involved a reaction between primary amine **356** and dihydropyranone **278**, the former being prepared in a number of steps from deoxy-D-ribose **354**. In the event, it was found that condensation of **356** with **278** was sensitive to temperature and pH. Reaction selectivity was improved by initial transformation of dihydropyranone **278** into its acetate derivative, followed by reaction with PPTS, which afforded 2-formylpyrrole **357** in good yield. Ley-Griffith oxidation of the secondary alcohol, TBS removal, and treatment with catalytic PPTS promoted cyclization to spiroketals **358** as a 1.5:1 mixture


of diastereomers. Acetonide removal using PPTS in methanol afforded a 7:1 ratio of (+) pollenopyrroside A **265** and (–) 9-epi-pollenopyrroside A **359**.

3.4.4 Xylapyrrosides A and B


In 2015 Hu *et al.*¹⁰⁹ isolated the pyrrole alkaloids xylapyrrosides A (**360**, Scheme 45) and B (**361**, Scheme 46), along with two previously known PSAs (pollenopyrrosides A and acortatarin A) from *Xylaria nigripes*. They also reported total syntheses of (–)-xylapyrrosides A **360**, (–)-xylapyrrosides B **361**, and (+)-



Scheme 44: Synthesis of (+)-pollenopyrroside A **265** (Brimble *et al.*, 2016)¹⁰⁸



Scheme 45: Stereoselective synthesis of (–)-xylapyrroside A **360** (Hu *et al.*, 2015)¹⁰⁹




Scheme 46: Stereoselective syntheses of (–)-xylapyrroside B **361** and (+)-acortatarin A **205** (Hu *et al.*, 2015)¹⁰⁹

acortatarin A **205**, all of which showed moderate antioxidant activity. These syntheses began with treatment of commercially available (*R*)-2,2-dimethyl-1,3-dioxolane-4-carbaldehyde **362** with allylmagnesium bromide, followed by protecting group adjustment to give alkene **363**. Epoxidation of the terminal olefin in **363**, followed by epoxide ring opening with cerium chloride heptahydrate and sodium iodide, then oxidation of the hydroxyl group delivered ketone **364**. This α -iodo ketone **364** was used to alkylate pyrrole **365**, giving **366** in 87% yield. This compound underwent intramolecular spiroketalization under acidic conditions to afford a pair of C-9 epimers, which enabled the synthesis of (–)-xylapyrroside A **360** in 70% yield on treatment with TiCl_4 .

(–)-Xylapyrroside B **361** and (+)-acortatarin A **205** were accessed using a very similar route (Scheme 46), which differs only in the arrangement of benzyl and TBS protecting groups. Specifically, ketone **368**¹⁰⁹ was prepared using equivalent chemistry and was converted to **369** via pyrrole alkylation. However, in this case the 5,6-PSA natural products resulted via desilylation, then debenzoylation.

3.4.5 Cycloprodigiosin

The natural product cycloprodigiosin **370** (Scheme 47) is produced by various bacteria including *Pseudoalteromonas denitrificans*, *Pseudoalteromonas* (*Alteromonas*) *rubra* and *Vibrio gazogenes*,¹¹⁰ with its structure first identified in 1983.^{111,112} This natural product has emerged as a potent anticancer and immunosuppressant agent.¹¹³ Over the past decades, only Fukuyama¹¹⁴ had reported a synthesis of



Scheme 47: Enantioselective synthesis of (*R*)-cycloprodigiosin **370** (Sarpong *et al.*, 2013)¹¹⁵


cycloprodigiosin **370** (in 1984) until the first enantioselective synthesis of (*R*)-cycloprodigiosin **370** was described by Sarpong *et al.*¹¹⁵ Their synthetic strategy commenced with enantioenriched allenyl alkyne **373**, which was prepared in six steps from known alkyne **372**.¹¹⁶ **373** was converted into a 1:1 mixture of imines **374** and the desired pyrrole **375**, via treatment with copper (II) thiophene carboxylate and tosyl azide followed by $\text{Rh}_2(\text{oct})_4$ in chloroform. The tosyl group in **375** was removed using lithium aluminium hydride to afford **376**. Under Lindsley conditions, condensation of **376** with **377** afforded cycloprodigiosin **370** in good yield.

3.4.6 Marineosin A

Marineosins A **378** and B **394** were isolated from a *Streptomyces*-related actinomycete in 2008.¹¹⁷ Marineosins A and B consist of two pyrrole moieties, a macrocyclic ring, and a spiro tetrahydropyran-dihydropyrrole iminal. Xu *et al.*¹¹⁸ reported the first total synthesis of marineosin A **378**; their synthetic approach was completed from the readily available (*S*)-6-methyl-5,6-dihydro-2-pyrone, but with low overall yield (1.2%). Harran *et al.*¹¹⁹ recently reported an asymmetric synthesis of marineosin A **378** and also reassigned its stereochemistry (Scheme 48). The route employed a photochemical rearrangement of pyridine *N*-oxide **382** (prepared in two steps by the Ni-catalyzed cross-coupling of 2,6-dichloropyridine with a bis-Grignard) to construct cyclophane pyrrole **383**. After enolate alkylation with epoxide **386**, the resultant putative oxocarbenium ion intermediate underwent addition of bipyrrrole **388**¹²⁰ to build the C-9 linked ansa-bridge **389**. Cleavage of the sulfonamide group (Mg / MeOH) gave **390**, which under acidic conditions rearranged to **391**. This unstable intermediate was converted to spiroiminales **392** and **393** in very low yield on oxidation with MnO_2 . Hydrogenation using Pd/BaSO_4 afforded (+)-marineosin A **378**.

3.4.7 Cyclooroidin

Hoveyda *et al.*¹²¹ reported an enantioselective synthesis of (–)-cyclooroidin **395** (Scheme 49) in 2014. Treatment of **396** with NaHCO_3 in dichloromethane generated the phosphinoylimine, direct subjection of which to enantioselective allenylation with propargyl boronic ester **397** (catalyzed by a copper complex with ligand **398**) gave homoallenylamide **399** in 62% yield. The phosphinoyl group present in **399** was selectively removed on

[illegible]

Scheme 50: Enantioselective synthesis of (*R*)-streptorubin B **404** (Thomson *et al.*, 2011)¹²⁶

of vinyl anion **408** and exposure of the resulting alcohol to KHMDS and 18-crown-6 produced the 10-membered ring **409** via a Cope rearrangement ring expansion. This intermediate then underwent alkene reduction along with simultaneous cleavage of the benzyl ethers; oxidation of the intermediate diol and paal-Knorr pyrrole synthesis affording pyrrole core **410** in 67% yield. **410** underwent an acid-mediated condensation reaction with aldehyde **377**,¹²⁷ followed by deprotection of the Boc group to obtain a 10:1 mixture of two compounds. The major product did not match the spectroscopic data of the natural product, but after 10 days the mixture had entirely transformed to **404**, as revealed by the reexamination of the NMR sample.

4 Conclusions

The discovery of natural products with useful biological activities has an important role in the future of human health. Pyrrole containing natural products have established medicinal importance, and hence provide the chemist with an array of fascinating targets. In past years, significant advances have been made in the development of effective methods for the

syntheses of pyrrole containing natural products, and various new natural products incorporating pyrrole motifs have been isolated and successfully prepared using such tactics. In this review, we have described synthetic methodologies for the construction of achiral and chiral pyrrole-containing natural products. Some of the more innovative strategies employed include C2-symmetric bithiourea catalysis, arene–ynamide cyclization, intramolecular Friedel–Crafts type cyclizations, oxidative cyclizations, gold catalysed annulation reactions, benzyne-mediated cyclizations and so on, which enable synthesis of the target molecule in fewer steps. For further details on strategy, **Table S1** (see the Supporting Information) summarises the various key transformations which have been used in this review for the total syntheses of pyrrole containing natural products. As ever, challenges nonetheless remain which we hope will inspire the synthetic community to seek ever more efficient approaches to these useful molecules.

5 Conflicts of Interest

The authors declare no conflict of interest.

6 Acknowledgements

RC and AS are thankful to DST-SERB (EEQ/2016/000489) for providing financial assistance for conducting research. NS would like to acknowledge CSIR-SRF for providing the fellowship. AS is thankful to DST-SERB (EEQ/2016/000489) for providing Senior Research Fellowship. SS is thankful to ICMR (45/66/2018-PHA/BMS/OL) for providing a Senior Research Fellowship. HA thanks the Wellcome Trust for support (218514/Z/19/Z). EA thanks the EPSRC for support (EP/S013172/1).

7 Notes and References

1. A.L. Harreus, Pyrrole, *Ullmann's Encyclopedia of Industrial Chemistry*, 2000.
2. F.F. Runge, Ueber einige produkte der steinkohlendestillation, *Annalen der Physik*, 1834, **107**, 65–78.
3. M. Iwao, T. Fukuda and F. Ishibashi, Synthesis and biological activity of lamellarin alkaloids: an overview, *Heterocycles*, 2011, **83**, 491–529.
4. Y. Kashman, G. Koren-Goldshlager, M.G. Gravalos and M. Schleyer, Halitulin, a new cytotoxic alkaloid from the marine sponge *Haliclona tulearensis*, *Tetrahedron Lett.*, 1999, **40**, 997–1000.
5. V. Estévez, M. Villacampa and J.C. Menéndez, Recent advances in the synthesis of pyrroles by multicomponent reactions, *Chem. Soc. Rev.*, 2014, **43**, 4633–4657.
6. R.A. Rane, N.U. Sahu, C.P. Shah and N.K. Shah, Design, synthesis and antistaphylococcal activity of marine pyrrole alkaloid derivatives, *J. Enzyme. Inhib. Med. Chem.*, 2014, **29**, 401–407.
7. N.R. Williamson, P.C. Fineran, F.J. Leeper and G.P. Salmond, The biosynthesis and regulation of bacterial prodiginines, *Nat. Rev. Microbiol.*, 2006, **4**, 887–899.
8. N.R. Williamson, H.T. Simonsen, R.A. Ahmed, G. Goldet, H. Slater, L. Woodley, F.J. Leeper and G.P. Salmond, Biosynthesis of the red antibiotic, prodigiosin, in *Serratia*: identification of a novel 2-methyl-3-n-amylyl-pyrrole (MAP) assembly pathway, definition of the terminal condensing enzyme, and implications for undecylprodigiosin biosynthesis in *Streptomyces*, *Mol. Microbiol.*, 2005, **56**, 971–989.
9. S. D Joshi, U. A More, V. H Kulkarni and T. M Aminabhavi, Pyrrole: chemical synthesis, microwave assisted synthesis, reactions and applications: a review, *Curr. Org. Chem.*, 2013, **17**, 2279–2304.
10. V. Estevez, M. Villacampa and J.C. Menendez, Multicomponent reactions for the synthesis of pyrroles, *Chem. Soc. Rev.*, 2010, **39**, 4402–4421.
11. M. S Mohamed and S. S Fathallah, Pyrroles and fused pyrroles: synthesis and therapeutic activities, *Mini. Rev. Org. Chem.*, 2014, **11**, 477–507.
12. I. Bauer and H.J. Knölker, Synthesis of pyrrole and carbazole alkaloids, In *Alkaloid Synthesis*, 2011, 203–253. Springer, Berlin, Heidelberg.
13. N. Yoshikai and Y. Wei, Synthesis of Pyrroles, Indoles, and Carbazoles through Transition-Metal-Catalyzed C H Functionalization, *Asian J. Org. Chem.*, 2013, **2**, 466–478.
14. A. Al-Mourabit, M.A. Zancanella, S. Tilvi and D. Romo, Biosynthesis, asymmetric synthesis, and pharmacology, including cellular targets, of the pyrrole-2-aminoimidazole marine alkaloids, *Nat. Prod. Rep.*, 2011, **28**, 1229–1260.
15. I.S. Young, P.D. Thornton and A. Thompson, Synthesis of natural products containing the pyrrolic ring, *Nat. Prod. Rep.*, 2010, **27**, 1801–1839.
16. K.Ø. Hanssen, B. Schuler, A.J. Williams, T.B. Demissie, E. Hansen, J.H. Andersen, J. Svenson, K. Blinov, M. Repisky, F. Mohn and G. Meyer, A combined atomic force microscopy and computational approach for the structural elucidation of breitfussin A and B: Highly modified halogenated dipeptides from *Thuiaria breitfussi*, *Angew. Chem. Int. Ed.*, 2012, **51**, 12238–12241.
17. S.K. Pandey, Y. Guttormsen, B.E. Haug, C. Hedberg and A. Bayer, A concise total synthesis of breitfussin A and B, *Org. Lett.*, 2015, **17**, 122–125.
18. A.A. Nabi, J. Liyu, A.C. Lindsay and J. Sperry, C4–H alkoxylation of 6-bromoindole and its application to the synthesis of breitfussin B, *Tetrahedron*, 2018, **74**, 1199–1202.
19. A.H. Khan and J.S. Chen, Synthesis of breitfussin B by late-stage bromination, *Org. Lett.*, 2015, **17**, 3718–3721.
20. N. Lindquist and W. Fenical, New tambjamine class alkaloids from the marine ascidian *Atapozoa* sp. and its nudibranch predators. Origin of the tambjamines in *Atapozoa*, *Experientia*, 1991, **47**, 504–506.

21. B. Carté and D.J. Faulkner, Role of secondary metabolites in feeding associations between a predatory nudibranch, two grazing nudibranchs, and a bryozoan, *J. Chem. Ecol.*, 1986, **12**, 795-804.
22. D.M. Pinkerton, M.G. Banwell and A.C. Willis, Total syntheses of tambjamines C, E, F, G, H, I and J, BE-18591, and a related alkaloid from the marine bacterium *Pseudoalteromonas tunicata*, *Org. Lett.*, 2007, **9**, 5127-5130.
23. C. Boonlarppradab, C.A. Kauffman, P.R. Jensen and W. Fenical, Marineosins A and B, cytotoxic spiroaminals from a marine-derived actinomycete, *Org. Lett.*, 2008, **10**, 5505-5508.
24. L.N. Aldrich, E.S. Dawson and C.W. Lindsley, Evaluation of the Biosynthetic Proposal for the Synthesis of Marineosins A and B, *Org. Lett.*, 2010, **12**, 1048-1051.
25. M. Carbone, C. Irace, F. Costagliola, F. Castelluccio, G. Villani, G. Calado, V. Padula, G. Cimino, J.L. Cervera, R. Santamaria and M. Gavagnin, A new cytotoxic tambjamine alkaloid from the Azorean nudibranch *Tambja ceutae*, *Bioorg. Med. Chem. Lett.*, 2010, **20**, 2668-2670.
26. L.N. Aldrich, S.L. Stoops, B.C. Crews, L.J. Marnett and C.W. Lindsley, Total synthesis and biological evaluation of tambjamine K and a library of unnatural analogs, *Bioorg. Med. Chem. Lett.*, 2010, **20**, 5207-5211.
27. N.K. Utkina, A.E. Makarchenko and V.A. Denisenko, Zyzzyanones B–D, Dipyrroloquinones from the Marine Sponge *Zyzya fuliginosa*, *J. Nat. Prod.*, 2005, **68**, 1424-1427.
28. N.K. Utkina, A.E. Makarchenko, V.A. Denisenko and P.S. Dmitrenok, Zyzzyanone A, a novel pyrrolo [3, 2-f] indole alkaloid from the Australian marine sponge *Zyzya fuliginosa*, *Tetrahedron Lett.*, 2004, **45**, 7491-7494.
29. D.H. Nadkarni, S. Murugesan and S.E. Velu, Total synthesis of zyzzyanones A–D, *Tetrahedron*, 2013, **69**, 4105-4113.
30. C.C. Hughes, C.A. Kauffman, P.R. Jensen and W. Fenical, Structures, reactivities, and antibiotic properties of the marinopyrroles A–F, *J. Org. Chem.*, 2010, **75**, 3240-3250.
31. C.C. Hughes, A. Prieto-Davo, P.R. Jensen and W. Fenical, The marinopyrroles, antibiotics of an unprecedented structure class from a marine *Streptomyces* sp, *Org. Lett.*, 2008, **10**, 629-631.
32. C. Cheng, Y. Liu, H. Song, L. Pan, J. Li, Y. Qin and R. Li, Marinopyrrole derivatives as potential antibiotic agents against methicillin-resistant *Staphylococcus aureus* (II), *Mar. Drugs*, 2013, **11**, 2927-2948.
33. K. Doi, R. Li, S.S. Sung, H. Wu, Y. Liu, W. Manieri, G. Krishnegowda, A. Awwad, A. Dewey, X. Liu and S. Amin, Discovery of marinopyrrole A (maritoclax) as a selective Mcl-1 antagonist that overcomes ABT-737 resistance by binding to and targeting Mcl-1 for proteasomal degradation, *J. Biol. Chem.*, 2012, **287**, 10224-10235.
34. C.C. Hughes, Y.L. Yang, W.T. Liu, P.C. Dorrestein, J.J.L. Clair and W. Fenical, Marinopyrrole A target elucidation by acyl dye transfer, *J. Am. Chem. Soc.*, 2009, **131**, 12094-12096.
35. M. Stodulski, S.V. Kohlhepp, G. Raabe and T. Gulder, Exploration of the Bis (thio) urea-Catalyzed Atropselective Synthesis of Marinopyrrole A, *Eur. J. Org. Chem.*, 2016, 2170-2176.
36. P. Cheng, D.L. Clive, S. Fernandopulle and Z. Chen, Racemic marinopyrrole B by total synthesis, *Chem. Commun.*, 2013, **49**, 558-560.
37. R. Raju, A.M. Piggott, L.X. Barrientos Diaz, Z. Khalil and R.J. Capon, Heronapyrroles A– C: Farnesylated 2-Nitropyrroles from an Australian Marine-Derived *Streptomyces* sp., *Org. Lett.*, 2010, **12**, 5158-5161.
38. T. Matsuo, S. Hashimoto, K. Nishikawa, T. Kodama, S. Kikuchi, Y. Tachi and Y. Morimoto, Total synthesis and complete stereochemical assignment of heronapyrroles A and B, *Tetrahedron Lett.*, 2015, **56**, 5345-5348.
39. X.B. Ding, D.P. Furkert and M.A. Brimble, 2-Nitropyrrole cross-coupling enables a second generation synthesis of the heronapyrrole antibiotic natural product family, *Chem. Commun.*, 2016, **52**, 12638-12641.
40. A.E. Stanley, L.J. Walton, M.K. Zerikly, C. Corre and G.L. Challis, Elucidation of the *Streptomyces coelicolor* pathway to 4-methoxy-2, 2'-bipyrrole-5-carboxaldehyde, an intermediate in prodiginine biosynthesis, *Chem. Commun.*, 2006, 3981-3983.
41. S. Mo, P.K. Sydor, C. Corre, M.M. Alhamadsheh, A.E. Stanley, S.W. Haynes, L. Song, K.A. Reynolds and G.L. Challis, Elucidation of the *Streptomyces coelicolor* pathway to 2-undecylpyrrole, a key intermediate in undecylprodiginine and streptorubin B biosynthesis, *Chem. Biol.*, 2008, **15**, 137-148.
42. N.R. Williamson, P.C. Fineran, F.J. Leeper and G.P. Salmond, The biosynthesis and regulation of bacterial prodiginines, *Nat. Rev. Microbiol.*, 2006, **4**, 887-899.
43. J.H. Frederich and P.G. Harran, Modular access to complex prodiginines: total synthesis of (+)-roseophilin via its 2-azafulvene prototropisomer, *J. Am. Chem. Soc.*, 2013, **135**, 3788-3791.
44. H.C. Kwon, A.P.D. Espindola, J.S. Park, A. Prieto-Davó, M. Rose, P.R. Jensen and W. Fenical, Nitropyrrolins A– E, cytotoxic farnesyl- α -nitropyrroles from a marine-derived bacterium within the actinomycete family *Streptomycetaceae*, *J. Nat. Prod.*, 2010, **73**, 2047-2052.
45. H. Mitani, T. Matsuo, T. Kodama, K. Nishikawa, Y. Tachi and Y. Morimoto, Total synthesis of nitropyrrolins A, B, and D, *Tetrahedron*, 2016, **72**, 7179-7184.
46. X.B. Ding, D.P. Furkert and M.A. Brimble, General Synthesis of the Nitropyrrolin Family of Natural Products via Regioselective CO₂-Mediated Alkyne Hydration, *Org. Lett.*, 2017, **19**, 5418-5421.
47. M. van Rensburg, B.R. Copp and D. Barker, Synthesis and Absolute Stereochemical Reassignment of Mukanadin F: A

- Study of Isomerization of Bromopyrrole Alkaloids with Implications on Marine Natural Product Isolation, *Eur. J. Org. Chem.*, 2018, 3065-3074.
48. R.P. Walker, D.J. Faulkner, D. Van Engen and J. Clardy, Scepterin, an antimicrobial agent from the sponge *Agelas scepterum*, *J. Am. Chem.*, 1981, **103**, 6772-6773.
 49. S. Urban, P. de Almeida Leone, A.R. Carroll, G.A. Fechner, J. Smith, J.N. Hooper and R.J. Quinn, Axinellamines A–D, Novel Imidazo–Azolo–Imidazole Alkaloids from the Australian Marine Sponge *Axinella* sp., *J. Org. Chem.*, 1999, **64**, 731-735.
 50. S. Nishimura, S. Matsunaga, M. Shibazaki, K. Suzuki, K. Furihata, R.W. van Soest and N. Fusetani, Massadine, a Novel Geranylgeranyltransferase Type I Inhibitor from the Marine Sponge *Stylissa* aff. *massa*, *Org. Lett.*, 2003, **5**, 2255-2257.
 51. I.B. Seiple, S. Su, I.S. Young, A. Nakamura, J. Yamaguchi, L. Jørgensen, R.A. Rodriguez, D.P. O'Malley, T. Gaich, M. Köck and P.S. Baran, Enantioselective total syntheses of (–)-palau'amine, (–)-axinellamines, and (–)-massadines, *J. Am. Chem. Soc.*, 2011, **133**, 14710-14726.
 52. S. Peters and P. Spiteller, Sanguinones A and B, blue pyrroloquinoline alkaloids from the fruiting bodies of the mushroom *Mycena sanguinolenta*, *J. Nat. Prod.*, 2007, **70**, 1274-1277.
 53. J.S. Lohmann, S. Wagner, M. von Nussbaum, A. Pulte, W. Steglich and P. Spiteller, Mycenaflavin A, B, C, and D: pyrroloquinoline alkaloids from the fruiting bodies of the mushroom *Mycena haematopus*, *Chem. Eur. J.*, 2018, **24**, 8609-8614.
 54. J. Backenköhler, B. Reck, M. Plaumann and P. Spiteller, Total Synthesis of Mycenarubin A, Sanguinolentaquinone and Mycenaflavin B and their Cytotoxic Activities, *Eur. J. Org. Chem.*, 2018, **22**, 2806-2816.
 55. R.A. Davis, S. Duffy, S. Fletcher, V.M. Avery and R.J. Quinn, Thiaplakortones A–D: Antimalarial thiazine alkaloids from the Australian marine sponge *plakortis lita*, *J. Org. Chem.*, 2013, **78**, 9608-9613.
 56. R.H. Pouwer, S.M. Deydier, P.V. Le, B.D. Schwartz, N.C. Franken, R.A. Davis, M.J. Coster, S.A. Charman, M.D. Edstein, T.S. Skinner-Adams and K.T. Andrews, Total synthesis of thiaplakortone A: Derivatives as metabolically stable leads for the treatment of malaria, *ACS Med. Chem. Lett.*, 2014, **5**, 178-182.
 57. Y. Yamaoka, T. Yoshida, M. Shinozaki, K.I. Yamada and K. Takasu, Development of a Brønsted acid-promoted Arene–Ynamide Cyclization toward the total syntheses of Marinoquinolines A and C and aplidiopsamine A, *J. Org. Chem.*, 2015, **80**, 957-964.
 58. S. Dey and P. Garner, Synthesis of tert-butoxycarbonyl (Boc)-protected purines, *J. Org. Chem.*, 2000, **65**, 7697-7699.
 59. W.Y. Yoshida, K.K. Lee, A.R. Carroll and P.J. Scheuer, A complex pyrrolo-oxazinone and its iodo derivative isolated from a tunicate, *Helv. Chim. Acta*, 1992, **75**, 1721-1725.
 60. K. Takamura, H. Matsuo, A. Tanaka, J. Tanaka, T. Fukuda, F. Ishibashi and M. Iwao, Total synthesis of the marine natural products lukianols A and B, *Tetrahedron*, 2013, **69**, 2782-2788.
 61. T. Fukuda, E.I. Sudo, K. Shimokawa and M. Iwao, Palladium-catalyzed cross-coupling of N-benzenesulfonyl-3, 4-dibromopyrrole and its application to the total syntheses of lamellarins O, P, Q, and R, *Tetrahedron*, 2008, **64**, 328-338.
 62. A.D. Yamaguchi, K.M. Chepiga, J. Yamaguchi, K. Itami and H.M. Davies, Concise syntheses of dictyodendrins A and F by a sequential C–H functionalization strategy, *J. Am. Chem. Soc.*, 2015, **137**, 644-647.
 63. K. Okano, H. Fujiwara, T. Noji, T. Fukuyama and H. Tokuyama, Total synthesis of dictyodendrin A and B, *Angew. Chem. Int. Ed.*, 2010, **49**, 5925-5929.
 64. H. Tokuyama, K. Okano, H. Fujiwara, T. Noji and T. Fukuyama, Total synthesis of dictyodendrins AE, *Chem. Asian J.*, 2011, **6**, 560-572.
 65. J. Matsuoka, Y. Matsuda, Y. Kawada, S. Oishi and H. Ohno, Total Synthesis of Dictyodendrins by the Gold-Catalyzed Cascade Cyclization of Conjugated Dienes with Pyrroles, *Angew. Chem. Int. Ed.*, 2017, **56**, 7444-7448.
 66. X. Alvarez-Mico, P.R. Jensen, W. Fenical and C.C. Hughes, Chlorizidine, a Cytotoxic 5 H-Pyrrolo [2, 1-a] isoindol-5-one-Containing Alkaloid from a Marine *Streptomyces* sp., *Org. Lett.*, 2013, **15**, 988-991.
 67. J.P. Mahajan and S.B. Mhaske, Synthesis of Methyl-Protected (±)-Chlorizidine A, *Org. Lett.*, 2017, **19**, 2774-2776.
 68. C. Eder, P. Proksch, V. Wray, K. Steube, G. Bringmann, R.W. van Soest, Sudarsono, X, E. Ferdinandus, L.A. Pattisina, S. Wiryowidagdo and W. Moka, New alkaloids from the indopacific sponge *Stylissa carteri*, *J. Nat. Prod.*, 1999, **62**, 184-187.
 69. R.P. Singh, M.R. Bhandari, F.M. Torres, T. Doundoulakis, D. Gout and C.J. Lovely, Total Synthesis of (±)-2-Debromohymenine via Gold-Catalyzed Intramolecular Alkyne Hydroarylation, *Org. Lett.*, 2020, **22**, 3412-3417.
 70. A. Al Mourabit and P. Potier, Sponge's molecular diversity through the ambivalent reactivity of 2-aminoimidazole: a universal chemical pathway to the oroidin-based pyrrole-imidazole alkaloids and their palau'amine congeners, *Eur. J. Org. Chem.*, 2001, 237-243.
 71. H. Hoffmann and T. Lindel, Synthesis of the pyrrole-imidazole alkaloids, *Synthesis*, 2003, 1753-1783.
 72. B.H. Northrop, D.P. O'Malley, A.L. Zografos, P.S. Baran and K.N. Houk, Mechanism of the vinylcyclobutane rearrangement of scepterin to ageliferin and nagelamide E, *Angew. Chem. Int. Ed.*, 2006, **45**, 4126-4130.

73. M. Köck, A. Grube, I.B. Seiple and P.S. Baran, The pursuit of palau'amine, *Angew. Chem. Int. Ed.*, 2007, **46**, 6586-6594.
74. D.E. Jacquot and T. Lindel, Challenge palau'amine: Current standings, *Curr. Org. Chem.*, 2005, **9**, 1551-1565.
75. M. Kuramoto, N. Miyake, Y. Ishimaru, N. Ono and H. Uno, Cylindradines A and B: novel bromopyrrole alkaloids from the marine sponge *Axinella cylindratus*, *Org. Lett.*, 2008, **10**, 5465-5468.
76. (a) K.G. Poullennec and D. Romo, Enantioselective total synthesis of (+)-dibromophakellstatin, *J. Am. Chem. Soc.*, 2003, **125**, 6344-6345. (b) S. Wang and D. Romo, Enantioselective Synthesis of (+)-Monobromophakellin and (+)-Phakellin: A Concise Phakellin Annulation Strategy Applicable to Palau'amine, *Angew. Chem. Int. Ed.*, 2008, **47**, 1284-1286.
77. (a) T. Imaoka, O. Iwamoto, K.I. Noguchi and K. Nagasawa, Total Synthesis of (+)-Dibromophakellin, *Angew. Chem. Int. Ed.*, 2009, **48**, 3799-3801. (b) T. Imaoka, T. Akimoto, O. Iwamoto and K. Nagasawa, Total synthesis of (+)-dibromophakellin and (+)-dibromophakellstatin, *Chem. Asian J.*, 2010, **5**, 1810-1816.
78. M. Iwata, K. Kanoh, T. Imaoka and K. Nagasawa, Total synthesis of (+)-cylindradine A, *Chem. Commun.*, 2014, **50**, 6991-6994.
79. B. P. J. de Lacy Costello, P. Evans, N. Guernion, N. M. Ratcliffe, P. S. Sivanand and G. C. Teare, The synthesis of a number of 3-alkyl and 3-carboxy substituted pyrroles; their chemical polymerisation onto poly (vinylidene fluoride) membranes, and their use as gas sensitive resistors, *Synth. Met.*, 2000, **114**, 181-188.
80. M. Kim, J. V. Mulcahy, C. G. Espino and J. Du Bois, Expanding the substrate scope for C–H amination reactions: oxidative cyclization of urea and guanidine derivatives, *Org. Lett.*, 2006, **8**, 1073-1076. Tces: 2,2,2-trichloroethoxysulfonyl.
81. M. Iwata, Y. Kamijoh, E. Yamamoto, M. Yamanaka and K. Nagasawa, Total Synthesis of Pyrrole–Imidazole Alkaloid (+)-Cylindradine B, *Org. Lett.*, 2017, **19**, 420-423.
82. Y. Wu, M. Suehiro, M. Kitajima, T. Matsuzaki, S. Hashimoto, M. Nagaoka, R. Zhang and H. Takayama, Rhazinilam and quebrachamine derivatives from *Yunnan Kopsia arborea*, *J. Nat. Prod.*, 2009, **72**, 204-209.
83. K. Zhao, S. Xu, C. Pan, X. Sui and Z. Gu, Catalytically asymmetric Pd/norbornene catalysis: enantioselective synthesis of (+)-rhazinal, (+)-rhazinilam, and (+)-kopsiunnanine C1–3, *Org. Lett.*, 2016, **18**, 3782-3785.
84. X.G. Tong, L.L. Zhou, Y.H. Wang, C. Xia, Y. Wang, M. Liang, F.F. Hou and Y.X. Cheng, Acortatarins A and B, two novel antioxidative spiroalkaloids with a naturally unusual morpholine motif from *Acorus tatarinowii*, *Org. Lett.*, 2010, **12**, 1844-1847.
85. J.M. Wurst, A. L. Verano and D.S. Tan, Stereoselective synthesis of acortatarins A and B, *Org. Lett.*, 2012, **14**, 4442-4445.
86. E. Wimmer, S. Borghèse, A. Blanc, V. Bénétteau and P. Pale, Zeolite-Based Organic Synthesis (ZeoBOS) of Acortatarin A: First Total Synthesis Based on Native and Metal-Doped Zeolite-Catalyzed Steps, *Chem. Eur. J.*, 2017, **23**, 1484-1489.
87. M. Hellwig and T. Henle, Baking, ageing, diabetes: a short history of the Maillard reaction, *Angew. Chem. Int. Ed.*, 2014, **53**, 10316-10329.
88. H.M. Geng, L.A. Stubbing, J. Li-yang Chen, D.P. Furkert and M.A. Brimble, Synthesis of the revised structure of acortatarin A, *Eur. J. Org. Chem.*, 2014, 6227-6241.
89. S. Peters and P. Spiteller, Mycenarubins A and B, red pyrroloquinoline alkaloids from the mushroom *Mycena rosea*, *Eur. J. Org. Chem.*, 2007, 1571-1576.
90. W. B. Han, Y. H. Lu, A. H. Zhang, G. F. Zhang, Y. N. Mei, N. Jiang, X. Lei, Y. C. Song, S. W. Ng and R. X. Tan, Curvulamine, a New Antibacterial Alkaloid Incorporating Two Undescribed Units from a *Curvularia* Species, *Org. Lett.*, 2014, **16**, 5366.
91. K.T. Haelsig, J. Xuan and T.J. Maimone, Total Synthesis of (–)-Curvulamine, *J. Am. Chem. Soc.*, 2020, **142**, 1206-1210.
92. V. Sofiyev, J.P. Lumb, M. Volgraf and D. Trauner, Total synthesis of exiguamines A and B inspired by catecholamine chemistry, *Chem. Eur. J.*, 2012, **18**, 4999-5005.
93. T. Yang, C.H. Wang, G.X. Chou, T. Wu, X.M. Cheng and Z.T. Wang, New alkaloids from *Capparis spinosa*: Structure and X-ray crystallographic analysis, *Food Chem.*, 2010, **123**, 705-710.
94. Z. Cao, Y. Li, S. Wang, X. Guo, L. Wang and W. Zhao, Total synthesis of two pyrrole spiroketal alkaloids: pollenopyrroside A and capparisine B, *Synlett*, 2015, **26**, 921-926.
95. S. Huo, Y. Li, C. Liang, J. Liu and W. Zhao, Synthesis of novel N-carbohydrate-derived heterocyclic compounds by nucleophilic substitution reaction of carbohydrate imidazole-1-sulfonates, *J. Carbohydr. Chem.*, 2011, **30**, 75-84.
96. T.Y. Yuen, S.E. Eaton, T.M. Woods, D.P. Furkert, K.W. Choi and M.A. Brimble, A Maillard Approach to 2-Formylpyrroles: Synthesis of Magnolamide, Lobechine and Funebral, *Eur. J. Org. Chem.*, 2014, 1431-1437.
97. D.G. Dunford and D.W. Knight, A brief total synthesis of Pyrrolostatin, *Tetrahedron Lett.*, 2016, **57**, 2746-2748.
98. J.M. Wood, D.P. Furkert and M.A. Brimble, Total synthesis and stereochemical revision of the 2-formylpyrrole alkaloid hemerocallisamine I, *J. Nat. Prod.*, 2017, **80**, 1926-1929.
99. X. Zhang, A.C. Schmitt and W. Jiang, A convenient and high yield method to prepare 4-hydroxypyroglutamic acids, *Tetrahedron Lett.*, 2001, **42**, 5335-5338.

100. J. Backenköhler, S. Spindler and P. Spiteller, Total Synthesis of Damirone C, Makaluvamine O, Makaluvone, Batzelline C and Batzelline D, *ChemistrySelect*, 2017, **2**, 2589-2592.
101. F.G.H. Lee, D.E. Dickson, J. Suzuki, A. Zirnis and A.A. Martian, Synthesis of 5, 7-and 6, 7-disubstituted tryptamines and analogs, *J. Heterocycl. Chem.*, 1973, **10**, 649-654.
102. T. Oshiyama, T. Satoh, K. Okano and H. Tokuyama, Total synthesis of makaluvamine A/D, damirone B, batzelline C, makaluvone, and isobatzelline C featuring one-pot benzyne-mediated cyclization–functionalization, *Tetrahedron*, 2012, **68**, 9376-9383.
103. W. Zhang, L. Ma, S. Li, Z. Liu, Y. Chen, H. Zhang, G. Zhang, Q. Zhang, X. Tian, C. Yuan and S. Zhang, Indimicins A–E, bisindole alkaloids from the deep-sea-derived *Streptomyces* sp. SCSIO 03032, *J. Nat. Prod.*, 2014, **77**, 1887-1892.
104. L.M. Blair and J. Sperry, Total syntheses of (±)-spiroindimicins B and C enabled by a late-stage Schöllkopf–Magnus–Barton–Zard (SMBZ) reaction, *Chem. Commun.*, 2016, **52**, 800-802.
105. I. Gerasimenko, Y. Sheludko and J. Stöckigt, 3-oxo-rhazinilam: a new indole alkaloid from *Rauvolfia serpentina* × *Rhazya stricta* hybrid plant cell cultures, *J. Nat. Prod.*, 2001, **64**, 114-116.
106. K. Sugimoto, K. Toyoshima, S. Nonaka, K. Kotaki, H. Ueda and H. Tokuyama, Protecting-Group-Free Total Synthesis of (–)-Rhazinilam and (–)-Rhazinicine using a Gold-Catalyzed Cascade Cyclization, *Angew. Chem. Int. Ed.*, 2013, **52**, 7168-7171.
107. R.B. Kinnel, H.P. Gehrken and P.J. Scheuer, Palau'amine: a cytotoxic and immunosuppressive hexacyclic bisguanidine antibiotic from the sponge *Stylotella agminata*, *J. Am. Chem. Soc.*, 1993, **115**, 3376-3377.
108. J.M. Wood, D.P. Furkert and M.A. Brimble, Synthesis of the 2-formylpyrrole spiroketal pollenopyrroside A and structural elucidation of xylapyrroside A, shensongine A and capparisine B, *Org. Biomol. Chem.*, 2016, **14**, 7659-7664.
109. M. Li, J. Xiong, Y. Huang, L.J. Wang, Y. Tang, G.X. Yang, X.H. Liu, B.G. Wei, H. Fan, Y. Zhao, W.Z. Zhai and Z.F. Hu, Xylapyrrosides A and B, two rare sugar-morpholine spiroketal pyrrole-derived alkaloids from *Xylaria nigripes*: Isolation, complete structure elucidation, and total syntheses, *Tetrahedron*, 2015, **71**, 5285-5295.
110. S. Lee, Y.S. Kim, S. Park, J. Kim, S.J. Kang, M.H. Lee, S. Ryu, J.M. Choi, T.K. Oh and J.H. Yoon, Exceptional production of both prodigiosin and cycloprodigiosin as major metabolic constituents by a novel marine bacterium, *Zooshikella rubidus* S1-1, *Appl. Environ. Microbiol.*, 2011, **77**, 4967-4973.
111. N.N. Gerber, Cycloprodigiosin from *Beneckea gazogenes*, *Tetrahedron Lett.*, 1983, **24**, 2797-2798.
112. H. Lattasch and R.H. Thomson, A revised structure for cycloprodigiosin, *Tetrahedron Lett.*, 1983, **24**, 2701-2704.
113. R. Pandey, R. Chander and K.B. Sainis, Prodigiosins as anti cancer agents: living upto their name, *Curr. Pharm. Des.*, 2009, **15**, 732-741.
114. H.H. Wasserman and J.M. Fukuyama, The synthesis of (±)-cycloprodigiosin, *Tetrahedron Lett.*, 1984, **25**, 1387-1388.
115. E.E. Schultz and R. Sarpong, Application of in situ-generated Rh-bound trimethylenemethane variants to the synthesis of 3, 4-fused pyrroles, *J. Am. Chem. Soc.*, 2013, **135**, 4696-4699.
116. T. Magauer, H.J. Martin and J. Mulzer, Total Synthesis of the Antibiotic Kendomycin by Macrocyclization using Photo-Fries Rearrangement and Ring-Closing Metathesis, *Angew. Chem. Int. Ed.*, 2009, **48**, 6032-6036.
117. C. Boonlarpradab, C.A. Kauffman, P.R. Jensen and W. Fenical, Marineosins A and B, cytotoxic spiroaminals from a marine-derived actinomycete, *Org. Lett.*, 2008, **10**, 5505-5508.
118. B. Xu, G. Li, J. Li and Y. Shi, Total synthesis of the proposed structure of marineosin A, *Org. Lett.*, 2016, **18**, 2028-2031.
119. Z. Feng, T.K. Allred, E.E. Hurlow and P.G. Harran, Anomalous chromophore disruption enables an eight-step synthesis and stereochemical reassignment of (+)-marineosin A, *J. Am. Chem. Soc.*, 2019, **141**, 2274-2278.
120. J. H. Frederich, J. K. Matsui, R. O. Chang and P. G. Harran, Substituted 2,2'-bipyrroles and pyrrolylfurans via intermediate isoxazolylopyrroles, *Tetrahedron Lett.*, 2013, **54**, 2645-2647.
121. N.W. Mszar, F. Haeffner and A.H. Hoveyda, NHC–Cu-Catalyzed Addition of Propargylboron Reagents to Phosphinoylimines. Enantioselective Synthesis of Trimethylsilyl-Substituted Homoallenylamides and Application to the Synthesis of S-(–)-Cyclooroidin, *J. Am. Chem. Soc.*, 2014, **136**, 3362-3365.
122. C.C. Hughes, C.A. Kauffman, P.R. Jensen, and W. Fenical, Structures, reactivities, and antibiotic properties of the marinopyrroles A–F, *J. Org. Chem.*, 2010, **75**, 3240-3250.
123. J. Patel, N. Pelloux-Léon, F. Minassian and Y. Vallée, Total synthesis of (S)-(–)-cyclooroidin, *Tetrahedron Lett.*, 2006, **47**, 5561-5563.
124. S. Mukherjee, R. Sivappa, M. Yousufuddin and C.J. Lovely, Asymmetric total synthesis of ent-cyclooroidin, *Org. Lett.*, 2010, **12**, 4940-4943.
125. H. Laatsch, M. Kellner and H. Weyland, Butyl-meta-cycloheptyl prodiginines—a revision of the structure of the former ortho-isomer, *J. Antibiot.*, 1991, **44**, 187-191.
126. D.X. Hu, M.D. Clift, K.E. Lazarski and R.J. Thomson, Enantioselective total synthesis and confirmation of the absolute and relative stereochemistry of streptorubin B, *J. Am. Chem. Soc.*, 2011, **133**, 1799-1804.

127. L.N. Aldrich, E.S. Dawson and C.W. Lindsley, Evaluation of the Biosynthetic Proposal for the Synthesis of Marineosins A and B, *Org. Lett.*, 2010, **12**, 1048-1051.