

Saponin adsorption at the air-water interface – neutron reflectivity and surface tension study

J. Penfold^{1,2,*}, R. K. Thomas¹, I. Tucker³, J. T. Petkov^{3†}, S. D. Stoyanov⁴, N. Denkov⁵, K. Golemanov^{4,‡}, S. Tcholakova⁵, J. R. P. Webster²

1. Physical and Theoretical Chemistry Laboratory, Oxford University, South Parks Road, Oxford, UK,
2. ISIS Facility, STFC, Rutherford Appleton Laboratory, Chilton, Didcot, OXON, UK
3. Unilever Research and Development, Port Sunlight Laboratory, Quarry Road East, Bebington, Wirral, UK
4. Unilever Research and Development, Vlaardingen, The Netherlands
5. Department of Chemical and Pharmaceutical Engineering, Faculty of Chemistry and Pharmacy, Sofia University, 1 J. Bourchier Avenue, 1164 Sofia, Bulgaria

† Current address: Lonza UK, GB-Blackley, Manchester, Lancs, M9 8ES, UK

‡ Current Address: Department of Chemical and Pharmaceutical Engineering, Faculty of Chemistry and Pharmacy, Sofia University, 1 J. Bourchier Avenue, 1164 Sofia, Bulgaria

*Corresponding author:

Prof. Jeff Penfold

Phone: +44 1235 445681

E-mail: jeff.penfold@stfc.ac.uk

Keywords: Saponin, Escin, Tea saponin, Quillaja saponin, neutron scattering, surfactant adsorption layer.

ABSTRACT

Saponins are a large group of glycosides present in many plant species. They exhibit high surface activity, which arises from a hydrophobic scaffold of triterpenoid or steroid groups and attached hydrophilic saccharide chains. The diversity of molecular structures, present in various plants, gives rise to a rich variety of physicochemical properties and biological activity, and results in a wide range of applications in foods, cosmetics, medicine and several other industrial sectors. Saponin surface activity is a key property in such applications and here the adsorption of three triterpenoid saponins, Escin, Tea saponins and Quillaja saponin, is studied at the air-water interface by neutron reflectivity and surface tension. All these saponins form adsorption layers with very high surface visco-elasticity. The structure of the adsorbed layers has been determined from the neutron reflectivity data, and is related to the molecular structure of the saponins. The results indicate that the structure of the saturated adsorption layers, is governed by densely packed hydrophilic saccharide groups. The tight molecular packing and the strong hydrogen bonds between the neighbouring saccharide groups are the main reasons for the unusual rheological properties of the saponin adsorption layers.

INTRODUCTION.

Saponins are a large group of glycosides present in more than 500 plant species.¹⁻⁴ Their intrinsic surface activity distinguishes them from other glycosides. This surface activity arises from a hydrophobic scaffold comprising a triterpenoid, steroid or steroid-alkaloid group and a hydrophilic part consisting of saccharide residues, attached to the hydrophobic scaffold via glycoside bonds. The wide range of different molecular structures of the saponins, found in various plant species, gives rise to a rich variety of physicochemical properties and biological activity.

The intrinsic surface activity of the saponins is the basis of their traditional use as emulsifiers and foaming agents in foods^{5,6} and their biological properties are recognised in natural medication.^{7,8} More recently the surface and biological activities have resulted in the identification and development of a variety of new applications in medicine, foods and cosmetics. Saponins are currently used as foam and emulsion stabilisers in beer and soft drinks,^{2,6} and as solubilising agents for food additives.⁹ They are used as stabilizers in cosmetic emulsions, shampoos and conditioners, and in skin anti-ageing products¹⁰. Saponins possess anti-inflammatory, anti-fungal, anti-bacterial, anti-cancer and cholesterol lowering functions¹¹. These properties make the saponins potentially important for an even wider range of new applications and technologies. Hence some aspects of the adsorption behaviour of saponins,¹²⁻¹⁶ saponin/surfactant¹⁷ and saponin/protein¹⁸⁻²¹ mixtures have been studied recently, and their self-assembly in solution has been characterised.^{22,23}

The saponins exhibit some very unusual surface properties, such as an extremely high surface modulus and shear visco-elasticity. These properties have an important impact upon the mechanisms associated with foam and emulsion stabilisation by these molecules. Hence several detailed studies have been focussed recently on the surface rheological properties of saponin solutions.¹²⁻¹⁵ Stanimirova et al.¹² and Golemanov et al.¹³ showed that the *Quillaja* saponin adsorption layers, subjected to small deformations, exhibit a very high surface dilatational elasticity, $\sim 280 \text{ mNm}^{-1}$, lower shear elasticity, $\sim 26 \text{ mNm}^{-1}$, and a negligible dilatational viscosity. The high surface elasticity is indicative of densely packed solid-like adsorbed layers, with strong inter-molecular interactions, arising from multiple hydrogen bonds between neighbouring sugar groups. Golemanov et al.¹³ showed that the surface layers of the *Quillaja* triterpenoid saponins were highly elastic, whereas the steroid based *Yucca*

saponin layers were purely viscous. In a subsequent paper Golemanov et al.¹⁴ and Paguerva et al.¹⁵ studied the surface rheology of a wider range of triterpenoid and steroid saponins, and showed that all steroid saponins exhibited no shear elasticity and had negligible surface viscosity. In contrast, most of the triterpenoid saponins showed complex visco-elastic surface behavior with extremely high surface elastic modulus (up to 1100 mN/m) and surface viscosity (130 N.s/m). The saponin extracts, showing the highest elastic moduli, were those of Escin, Tea saponins and Berry saponins, all containing predominantly monodesmosidic triterpenoid saponins (with a single oligosaccharide chain attached to the triterpenoid group). Hence, to explain the reasons for the unusually high visco-elasticity of the adsorption layers observed in some saponins, it is important to characterise these adsorbed layers in more detail. Both the total adsorbed amount and the structure of the adsorbed layer will provide important information about the interactions between adsorbed saponin molecules.

Hence three different saponins, Escin, Tea saponins and *Quillaja* saponin, were studied and are reported in this paper. The molecular structures of these saponins are shown in Figure 1. These are all triterpenoid saponins which exhibit rather high surface visco-elasticity. The main difference between these saponins is in the number and length of the oligosaccharide hydrophilic chains. Escin and Tea saponins have only one oligosaccharide chain attached to the triterpenoid group (monodesmosidic saponins) with Tea saponin having one more saccharide group, whereas the *Quillaja* saponins have two oligosaccharide chains attached to the triterpenoid group (bidesmosidic saponin).

It is now well established that neutron reflectivity can be used to determine adsorption at interfaces,²⁴ and has been demonstrated for surfactants,²⁴ polymers,²⁵ proteins,²⁶ and their associated mixtures at the air-water interface. Although isotopic substitution, using H/D exchange, is often used to optimise the selectivity and sensitivity of neutron reflectivity to the adsorption, it has been shown that for proteins and some other large bio-molecules there is often sufficient contrast without the need to use deuterium labelling.^{26,27} This is particularly important here in the context of studying the adsorption of saponin at the air-water interface, where the composition and size of the saponin molecules ensures a sufficient contrast to observe their adsorption at the air-water interface using neutron reflectivity without the need for deuterium labelling. However, deuterium labelling or partial labelling of the solvent can help to provide additional structural in-sights, as demonstrated in this study.

Measurements were made at the air-water interface, at concentrations from below to above the critical micelle concentration (cmc) of the saponins in order to establish the adsorption isotherms. Neutron reflectivity data were obtained and surface tension measurements made in order to quantify the amount of adsorbed saponin. For Escin and Tea saponin above the cmc, neutron reflectivity measurements were made with different solvent contrasts, in D₂O and H₂O/ D₂O mixture, in order to determine in more detail the structure of the adsorbed layer. For Escin the effect of pH on the adsorption was also studied, as it contains a carboxylic group in one of its saccharide groups. The neutron reflectivity and surface tension measurements were made on the same systems and under the same solution conditions.




Figure 1. Molecular structure of the saponins studied, (A)Escin, (B)Tea Saponin, and (C) Quillaja saponin. The Escin and Tea saponin have only one oligosaccharide chain, attached to the triterpenoid scaffold, but differ in the number and the type of sugar residues. The Quillaja

saponin is a larger molecule with more sugar residues, combined in two separate oligosaccharide chains.

EXPERIMENTAL SECTION.

Materials used.

The Escin, also known as Aescin, was obtained from Sigma (cat. num. E1378, CAS Number 6805-41-0, molecular formula $C_{54}H_{84}O_{23}$). The Tea saponin extract was obtained from Zhejiang Yuhong Import & Export Co., Ltd, and used as supplied. According to the product data sheets the concentrations of saponins in these extracts were 95% for Escin and 96.2% for the Tea saponin. The *Quillaja* saponin extract (Supersap) was obtained from Desert King, Chile. It is a white powder produced for pharmaceutical applications, with very high purity, and contains 91 wt % saponins. The rest is ~ 8 wt % moisture, and traces of electrolytes and other organic ingredients. As such only Escin is a monomolecular substance. The Tea saponin and Supersap extracts studied contain a variety of molecular species, which share the same triterpenoid scaffold but differ to some extent in the number and type of the attached saccharide groups. In Figure 1 we show and in the following consideration we use the predominant component in the respective extracts as representative for the whole extract.

All solutions contained 10 mM NaCl to maintain a fixed ionic strength. The pH of the working solutions was adjusted by adding small aliquots of 0.1M NaOH or 0.1M HCl solutions, prepared from NaOH with purity ≥ 95 % and 32 % HCl acid (both products of Sigma-Aldrich).

Experimental Methods.

The neutron reflectivity measurements were made on the INTER reflectometer at the ISIS neutron source.²⁸ The neutron beam was incident at the air-water interface with a grazing angle of incidence, $\theta = 2.3^\circ$, to cover a wave vector transfer, $Q = 4\pi\sin\theta/\lambda$, in the direction normal to the surface in the range between 0.03 and 0.5 \AA^{-1} , using neutron wavelengths from 0.5 to 15 \AA . The reflectivity, $R(Q)$, was calibrated with respect to the direct beam intensity and reflection from a D_2O surface. The measurements were made in sealed Teflon troughs with sample volumes ~ 25 mL and maintained at a temperature of 25 $^\circ\text{C}$. Each reflectivity profile took ~ 60 min, and repeated measurements over a 3-4 hour period showed

no further changes with time. Hence it is assumed that the data presented represent equilibrium adsorption values.

In the kinematic or Born approximation the reflectivity $R(Q)$ is related to the square of the Fourier transform of the scattering length density profile, $\rho(z)$,²⁴:

$$R(Q) = \frac{16\pi^2}{Q^2} \left| \int \rho(z) e^{-iQz} dz \right|^2 \quad (1)$$

where $\rho(z) = \sum_i n_i(z) b_i$, $n_i(z)$ is the number density of the i -th component and b_i - its scattering length. The scattering length, molecular weight and molecular volumes of the three saponins studied are listed in Table 1. However in the context of this study the standard optical matrix method is used to analyse the data (24).

Table 1. Saponin parameters used for analysis of neutron reflectivity data.

Saponin type	$\sum b$ (x10 ⁻⁴ nm)	Molecular weight	ρ (x10 ⁻⁸ nm ⁻²)
Escin	1.78	1101	1.6
Tea	1.97	1259	1.5
Quillaja saponin	2.00	1650	1.2

The adsorption measurements were made in null reflecting water, nrw, (92 mol% H₂O / 8 mol% D₂O) with a scattering length density of zero, the same as air. The data are consistent with a single monolayer at the interface. Analysing the data as a single layer of uniform composition yields a layer thickness, d , and a scattering length density, ρ . The area / molecule is then given by,

$$A = \sum b_i / \rho d \quad (2)$$

and the adsorbed amount, Γ [mol.m⁻²], is then

$$\Gamma = 1/N_a A \quad (3)$$

where N_a is the Avogadro number. Additional neutron reflectivity measurements were made for the Escin and Tea saponins at a concentration higher than the cmc in D_2O and a D_2O/H_2O mixture, index matched to $\rho=4 \times 10^{-8} \text{ nm}^{-2}$, in order to obtain a more detailed description of the adsorbed layer structure. From this sequence of measurements the data were simultaneously analysed with the simplest model consistent with the data, in this case a two-layer model. In both the adsorption measurements and the more detailed structural measurements the key refinable model parameters are the thickness and scattering length density of the layers, and a flat sample / instrumental background (~ 5 to 8×10^{-6}). In both the single and two layer characterisations, the simplest model consistent with all the data is used; and additional parameters, such as roughness which can be quite arbitrary and difficult to assign and physical significance to, are not included.

The Wilhelmy plate method ^{29, 30} was used to determine the equilibrium surface tension of the saponin solutions, σ_e . These measurements were made on a K100 tensiometer (Kruss GmbH, Hamburg, Germany) using a platinum plate. Before each measurement, this plate was cleaned by heating on a flame, followed by abundant rinsing with deionized water. The measurements were all made at a constant temperature of 20 °C.

RESULTS and DISCUSSION.

Neutron reflectivity data.

The adsorption of the Escin saponin was measured at the air-water interface in nrw, in the concentration range of 2×10^{-6} to $1.5 \times 10^{-3} \text{ M/l}$ by neutron reflectivity. The data are modelled as a single uniform layer with a thickness of $\approx 2.6 \text{ nm}$. The variation in the adsorbed amount with surfactant concentration, at natural pH, pH 4 and pH 8, is shown in Figure 2. The saturated adsorbed amounts at the different pH values are evaluated as 2.2, 2.4 and $2.5 \pm 0.3 \text{ } \mu\text{mol m}^{-2}$ at pH 8, natural pH and pH 4 respectively; corresponding to area/molecule of 0.75, 0.69 and $0.61 \pm 0.04 \text{ nm}^2$. However, within experimental error, the adsorption at and above the cmc is independent of pH, and has a mean value $\approx 2.4 \pm 0.3 \text{ } \mu\text{mol.m}^{-2}$ (area/molecule $\approx 0.69 \pm 0.04 \text{ nm}^2$). A clear effect of changing the solution pH is

evident at lower saponin concentrations and as the pH decreases from 8 to 4 the onset of adsorption saturation occurs at about a 5-fold lower concentration.

The latter results indicate that Escin does not behave entirely as a simple nonionic surfactant. This is due to the presence of a carboxylic group in one of its sugar groups, see Figure 1A. The carboxylic group is expected to be protonated at pH = 4, ionized at pH = 8, and probably it is partially ionized at the natural pH, which varies between ≈ 5.2 and ≈ 4.7 at a saponin concentration between 1.82 $\mu\text{mol/L}$ and 7.26 $\mu\text{mol/L}$. Nevertheless, as discussed above, the saturation saponin adsorption is, within experimental error, not affected by pH. The results shows that the saturation adsorption is controlled by the molecular dimensions which determine the area per molecule at close packing in the layer, while the presence of the ionic charge has a major effect in diluted adsorption layers only. The triterpenoid scaffold in the saponin molecules is rather large (it contains 30 carbon atoms) and, therefore, the surface activity of these molecules remains very high, even in the presence of ionic charge.




Figure 2. Adsorbed amount versus surfactant concentration at the air-water interface for Escin saponin at natural pH (black triangles), pH = 8 (red circles), and pH = 4 (blue squares).

Similar neutron reflectivity measurements were made in nrw, at natural pH only, for the Tea saponin and for *Quillaja* saponin (Supersap) in the concentration range between 10^{-3}

and 0.5 mM. The respective saponin adsorption isotherms are shown in Figure 3, and compared with that measured for Escin at natural pH. The mean adsorption for the Tea saponin above the cmc is $\approx 2.1 \mu\text{mol.m}^{-2}$ (area/molecule $\approx 0.79 \pm 0.04 \text{ nm}^2$) and for Supersap it is $\approx 1.9 \mu\text{mol.m}^{-2}$ (area/molecule $\approx 0.87 \pm 0.0 \text{ nm}^2$). As expected, the area per molecule at saturation increases with the molecular mass of the saponin molecules. The surface saturation occurs at comparable bulk molecular concentrations of Escin and Tea saponins, because their hydrophobic fragments are very similar, see the comparison in Figures 1A and 1B. The surface saturation occurs at a much lower concentration for Supersap, due to the additional hydrophobic chain present in bidesmosidic Quillaja saponin molecules, see Figure 1C, which leads to higher surface activity. However, given the higher level of impurity in this saponin, further more detailed structural measurements were not pursued here.






Figure 3. Adsorbed amount versus surfactant concentration at the air-water interface for Tea saponin (black circles), Escin saponin (red squares) and Quillaja saponin (blue triangles) at natural pH.

To determine the total adsorbed amount from the neutron reflectivity data, the product $d \cdot \rho$ is sufficient, see eq. 2. For the Escin and Tea saponins, the data at saturation adsorption were further analysed by systematically varying d to obtain a more accurate estimate of the thickness of the adsorbed layer in the frame of the homogeneous monolayer model. The variation in χ^2 (“goodness of fit”) with d provides a more accurate evaluation of the

monolayer thickness and its associated error. This is shown in Figure S1 for Escin and Tea saponin in the Supporting Information. By minimizing the variation in χ^2 values for the mean adsorbed layer thickness of 2.8 ± 0.2 nm for Escin and 3.0 ± 0.2 nm for the Tea saponin were derived. A more detailed characterization of the structure of the adsorbed layer for the Escin and Tea saponins was made by making a series of neutron reflection measurements with different solvent contrasts, in nrw, D₂O, and a D₂O/H₂O mixture with scattering length density $\approx 4 \times 10^{-8}$ nm⁻² (cm4.0). This latter scattering length density was estimated to be intermediate between that for D₂O and the hydrated headgroup region, and should provide the greatest sensitivity to the structural details in combination with the measurements in nrw and D₂O. The experimental data, obtained with 0.3 mM Escin solutions ($>$ cmc), at natural pH, are shown in Figure 4a along with the associated model fits. The data for the different contrasts were analysed simultaneously using the same structural model, in which only the scattering length densities of the layers were independent variables. The single layer model, used for the data in nrw to estimate the adsorbed amount, was not sufficient when the solvent contrast was altered and the different regions of the surface were highlighted differently. The simplest model consistent with these data was a two-layer model. The corresponding scattering length density distributions are shown in Figure 4b, and the associated parameters are summarised in Table 2.



(a)



(b)

Figure 4. (a) Neutron reflectivity profile for Escin saponin at 0.3 mM in nrw, D₂O and cm4.0 solvents. The solid lines are model fits using the profiles in b; (b) Scattering length density profiles from model fits, (—) nrw, (...) cm4.0, and (--) D₂O, for key model parameters summarised in table 2.

The results in Table 2 show an outer layer, adjacent to the air phase, with thickness ~ 0.8 nm, volume fraction of saponin ~ 0.3 , and no solvent. This outer layer corresponds to the triterpenoid hydrophobic region of the saponin molecules. Taking into account that the total length of this hydrophobic region is ≈ 1.4 nm, we interpret this result as an indication that a fraction of this scaffold is embedded in the aqueous subphase and/or that the triterpenoid scaffold is strongly tilted with respect to the air-water interface. Both these explanations are consistent with a recent simulation of Escin adsorption by molecular dynamics³¹. The inner layer, adjacent to the aqueous phase, is ≈ 1.4 nm thick. It contains both saponin and water, with volume fractions of ≈ 0.8 and 0.2 , respectively. The high volume fraction of the saponin fragments in this inner layer (80 vol. %) reveals that it is composed mostly of tightly packed, hydrated sugar groups.

Saponin type	Concentration (mM)	Solvent contrast	$d_1 [\pm 0.1 \text{ nm}]$	$\rho_1 (\pm 0.04 \times 10^{-8} \text{ nm}^{-2})$	$d_2 [\pm 0.1 \text{ nm}]$	$\rho_2 (\pm 0.04 \times 10^{-8} \text{ nm}^{-2})$
Escin	0.3	nrw	0.8	0.58	1.4	1.3

Escin	0.3	D ₂ O	0.8	0.58	1.4	3.0
Escin	0.3	Cm4.0	0.8	0.58	1.4	2.2
Tea	0.13	nrw	1.4	0.7	1.3	1.0
Tea	0.13	D ₂ O	1.4	0.7	1.3	2.8
Tea	0.13	Cm4.0	1.4	0.7	1.3	2.8

Table 2. Key model parameters from analysis of neutron reflectivity in figures 4a and S2a

A similar sequence of neutron reflection measurements was made for the Tea saponin at natural pH and a saponin concentration of 0.13 mM (above cmc). The neutron reflection data and the corresponding model fits are shown in Figure S2a in the Supporting Information. The corresponding scattering length density profiles are shown in Figure S2b, and the key model parameters are listed in Table 2. The two layer model is also the simplest one consistent with the data, and is different to that observed for Escin. The outer layer, adjacent to the air phase is ≈ 1.4 nm thick, has a saponin volume fraction ≈ 0.4 , and contains no solvent. The inner layer, adjacent to the aqueous phase, is ≈ 1.3 nm thick, has a volume fraction of saponin ≈ 0.65 , and a corresponding water volume fraction ≈ 0.35 .




Figure 5. Schematic representation of the structure of the saponin adsorption layer: (A) Escin; (B) Tea Saponin; consistent with the measured scattering length density profiles.

Figure 5 presents schematically the structure of adsorption layers of Escin and Tea saponin. These results indicate qualitatively similar structure and packing of the Escin and Tea saponin molecules in their adsorption layers, however, with some noticeable differences. The thickness of the solvated hydrophilic region is similar for both saponins, and the slight difference in the saponin volume fraction in that inner layer reflects the higher adsorption of the Escin compared to Tea saponin. For both saponins the volume fraction in the inner layer is relatively high. This is consistent with the strong inter-molecular interactions which arise from multiple hydrogen bonding between neighbouring sugar groups, and which are assumed to be responsible for the high surface elasticity, observed with these adsorption layers.¹²⁻¹⁴ The most important difference between the two saponins is that the hydrophobic part of the Tea saponin adsorption layer is much thicker and seems to be composed of triterpenoid scaffold which is directed perpendicularly to the surface and is entirely exposed to the air phase. Whether this difference is due to the extra hydroxyl group present in the triterpenoid part of the Escin (which is absent in Tea saponin) or to the extra sugar residue in the Tea saponin, remains unclear for the moment. For both saponins, the hydrophobic outer layer is not densely packed and contains a significant fraction of empty space between the hydrophobic scaffolds. This important result demonstrates that the total saponin adsorption is predominantly governed by the close-packing of the hydrated sugar groups which are densely populating the inner layer, adjacent to the water sub-phase.

Surface tension data.

In addition to the neutron reflectivity measurements, complementary surface tension measurements were made. The surface tension of the saponin solutions were measured by the Wilhelmy plate method, as described earlier. For these solutions the surface tension decreases with time, and the kinetics of surface tension, $\sigma(t)$, and approach to equilibrium are relatively slow and can take many minutes¹². Here an approach which describes $\sigma(t)$ by a bi-exponential equation with two relaxation times (t_1, t_2)¹² was used,

$$\sigma(t) - \sigma_e = \Delta\sigma_1 e^{-t/t_1} + \Delta\sigma_2 e^{-t/t_2} \quad (4)$$

where $\sigma(t)$ and σ_e are the current and the equilibrium values of the surface tension, respectively. The time dependence of the surface tension data was described by eq. 4. From the best fits the values of the equilibrium surface tension, σ_e , were determined, and used to plot the variation in surface tension with surfactant concentration, as shown in Figure S3 in the Supporting Information. The Gibbs adsorption equation^{32,33} was used to obtain an estimate of the adsorption at saturation,

$$\frac{d\sigma_e}{d\ln C_s} = -k_B T \Gamma \quad (5)$$

where k_B is Boltzmann constant, T is absolute temperature, C_s is surfactant concentration in the bulk solution, and Γ is surfactant adsorption. The experimental points in the concentration range corresponding to $\sigma_e < 65$ mN/m and cmc were interpolated by eq. 5 to determine the saponin adsorption, as summarised in Table 3.

Table 3. Critical micelle concentration (cmc), area / molecule and saponin adsorption at cmc, for air-water interface, as determined from surface tension.

Saponin type	pH	cmc (mM)	Area/molecule (nm ²)	Adsorbed amount, Γ (μmol.m ⁻²)
Escin in light water	4	0.06	0.63 ± 0.05	2.6 ± 0.2
Escin in light water	natural pH	0.11	0.57 ± 0.06	2.9 ± 0.2
Escin in D ₂ O/H ₂ O (nrw)	natural pH	0.11	0.54 ± 0.06	3.1 ± 0.2
Escin in light water	8	0.072	0.47 ± 0.05	3.2 ± 0.3
Tea saponin in light water	natural pH	0.30	0.86 ± 0.08	2.3 ± 0.2
Tea saponin in D ₂ O/H ₂ O (nrw)	natural pH	0.30	0.66 ± 0.06	2.5 ± 0.2
Supersap in light water*	natural pH	0.13	1.1 ± 0.05	1.5 ± 0.05

*-data taken from Stanimirova et al. [12].

A comparison of the adsorbed amounts for the Escin and Tea saponins obtained from neutron reflectivity and surface tension data show some differences. The mean values from neutron reflectivity for the Escin and Tea saponins are 2.4 and 2.1 $\mu\text{mol m}^{-2}$, compared to values from surface tension of 3.0 and 2.4. In both the neutron reflectivity and surface tension data for the Escin saponin there is an increase in the adsorption as the pH increases from 4 to 8. From the neutron reflectivity data it increases from 2.2 to 2.7 $\mu\text{mol m}^{-2}$, and from the equivalent surface tension data the change is from 2.6 to 3.2 $\mu\text{mol m}^{-2}$. However, in both cases the changes are within experimental error. Even at pH 4, where Escin is expected to be non-ionic, the adsorbed amounts from neutron reflectivity and surface tension are still systematically different. For the Supersap saponin the adsorbed amount from surface tension is 1.5 $\mu\text{mol m}^{-2}$ compared to value of 1.9 $\mu\text{mol m}^{-2}$ from neutron reflectivity. It has been shown for non-ionic surfactants³⁴, where generally the adsorption saturates before the cmc, that the adsorption determined from surface tension can be in good agreement with the directly determined surface excesses from neutron reflectivity. In broadest terms the results from surface tension and neutron reflectivity for Escin at different pH, for Tea saponin and for Supersap saponin are in agreement when the errors in the measurements are taken into account. Indeed the general trends; that is, the adsorption is highest for Escin, intermediate for Tea saponin and lowest for Supersap saponin, are consistent for both the surface tension and neutron reflectivity data.

For ionic or weakly ionic surfactants, Xu et al³⁵ have shown that compared to the neutron reflectivity data, the application of equation 5 to surface tension data generally underestimates the saturation adsorption. This is as a result of the variation in the activity in the region of the cmc due to the onset of micellisation. Xu et al³⁵ showed that this can be accounted for by applying the Mass Action model of micellisation to the surface tension data in that region, where the variation largely depends upon the degree of ionisation of the micelles. From the form of the adsorption isotherms measured by neutron reflectivity (see figures 2 and 3) the saponins are most likely only weakly ionic at any of the solution conditions. However, taking into account such factors would not necessarily account for the differences encountered. For example, surface tension overestimates the adsorption compared to neutron reflectivity for the Escin and Tea saponins, which is opposite to the trends

normally observed ³⁵. Whilst for the Supersap saponin surface tension underestimates the adsorption compared to neutron reflectivity. For the saponins there is another factor, in that it is assumed that the saponins are a single non-ionic species, whereas the Tea and Supersap saponin extracts contain a range of molecular species with different sugar residues and structures. In both the surface tension and neutron reflectivity data analysis the saponins are assumed to be a single molecular species, as shown in figure 1. It is possible that the heterogeneity of the materials has a greater impact on the interpretation of the surface tension data than the neutron reflectivity data.

The surface tension measurements for the Escin and Tea saponins were also made in two different solvent isotopic compositions, in H₂O and nrw (see table 3 and figure S3 in the Supporting Information). Generally the evidence in the literature is that in surfactant adsorption there is no measurable isotope effect when H is replaced by D, as demonstrated on a wide range of surfactants ²⁴, and more recently on some different biosurfactants³⁶. Although nrw contains only 8 mole % D₂O the occurrence of strong hydrogen bonding between the saponin sugar groups raises the potential for a deuterium isotope effect, as discussed by Wade³⁶. The results summarised in table 3 show that the differences encountered in H₂O and nrw are within experimental error and that any isotope effect is not significant.

SUMMARY and CONCLUSIONS

The structural measurements with neutron reflectivity show some interesting similarities and differences between the adsorption layers of Escin and Tea saponins at air-water interface, as illustrated in figures 4, S2 and 5. The surface structure is dominated by a relatively dense and hydrated headgroup region. The different headgroup structure, with an additional sugar group in the Tea saponin, compared to Escin, does not result in significant differences in the hydrophilic region. The thicknesses are similar, and the density difference is in part accounted for by the difference in the saturation adsorption for these two saponins. The hydrophobic triterpenoid region is not densely packed and contains no solvent. For Escin, this region is rather thin, when compared to the total length of the triterpenoid scaffold, which shows that this hydrophobic region is partially embedded in the aqueous phase and / or tilted with respect to the surface normal. For Tea saponin this layer is thicker which indicates

fully stretched triterpenoid groups, oriented almost perpendicularly to the solution surface. These observations, combined with the fact that both these saponins form highly visco-elastic adsorption layers, support the assumption that strong inter-molecular hydrogen bonds occur between the neighbouring sugar groups in the adsorption layers, and that these bonds give rise to the observed high surface elasticity.

The molar adsorption decreases with the molecular mass of the saponins, and is highest for Escin, intermediate for Tea saponin, and lowest for Quillaja saponin. The total adsorption determined from surface tension is in a broad agreement with the results from the neutron reflectivity. However, differences up to 20-30 % in the measured adsorption were observed for the results, obtained by the two methods; and such differences are often encountered, as discussed by Li et al ³⁴ and Xu et al ³⁵. Here these differences have a further contributing factor as saponin extracts studied are typically multicomponent mixtures, containing ionisable carboxylic groups. The simple approach for interpretation of surface tension data, based on eq. 5, may not be appropriate for these systems. Nevertheless, the surface tension data give reasonable estimate of the saponin adsorption and the main trends for the three saponins studied are captured.

The neutron reflection results show that relatively subtle differences in the structure of the adsorbed layers are visible for saponin molecules, without the need for deuterium labelling of the saponin. These structural properties can be potentially correlated with macroscopic surface properties, such as the surface rheology. This approach could be applied to a wider range of similar systems, provided that relatively high purity samples are available.

REFERENCES

- (1) Hostettmann, K.; Marston, A. *Saponins*; Cambridge University Press: New York, 1995.
- (2) Guglu-Ustundag, O.; Mazza, G. Saponins: properties, application, and processing. *Crit. Rev. Food Sci. Nutr.* **2007**, *47*, 231-258.
- (3) Vincken, J. P.; Heng, L.; De Groot, A.; Gruppen, H. Saponins, classification and occurrence in the plant kingdom. *Phytochemistry* **2007**, *68*, 275-297.
- (4) Sparg, S. G.; Light, M. E.; van Staden, J. Biological activities and distribution of plant saponins. *J. Ethnopharmacology* **2004**, *94*, 219-243.
- (5) Oakenfull, D. Saponins in food – a review. *Food Chem.* **1981**, *7*, 19-40.
- (6) Cheeke, P. R. Actual and potential applications of *Yucca schidigera* and *Quillaja saponaria* saponins in human and animal nutrition. *Proc. Am. Soc. Anim. Sci.* **1999**, *E9*, 1-10.
- (7) Liu, J.; Henkel, T. Traditional Chinese medicine: are polyphenols and saponins the key ingredients triggering biological activities. *Curr. Med. Chem.*, **2002**, *9*, 1483-1485.
- (8) Fukuda, N.; Tanaka, H.; Shoyama, Y. Isolation of the pharmacologically active saponin Ginsenoside Rb1 from Ginseng by immunoaffinity column chromatography. *J. Nat. Prod.* **2000**, *62*, 283-285.
- (9) Jenkins, K. J.; Atwal, A. S. Effects of dietary Saponins on fecal bile acids and neutral steroids, and availability of vitamins A and E in the chick. *J. Nutr. Biochem.* **1994**, *5*, 134-137.
- (10) Brown, R. The natural way in cosmetics and skin care. *Chem. Mark. Rep.* **1998**, *254*, FR8.
- (11) Sirtori, C. R. Aescin: pharmacology, pharmacokinetics and therapeutic profile. *Pharmacol. Res.* **2001**, *44*, 183-193.
- (12) Stanimirova, R.; Marinova, K.; Tcholakova, S.; Denkov, N. D.; Stoyanov, S.; Pelan, E. Surface rheology of saponin adsorption layers. *Langmuir* **2011**, *27*, 12486-12498.
- (13) Golemanov, K.; Tcholakova, S.; Denkov, N.; Pelan, E.; Stoyanov, S. D. Surface shear rheology of saponin adsorption layers. *Langmuir* **2012**, *28*, 12071-12084.
- (14) Golemanov, K.; Tcholakova, S.; Denkov, N. D.; Pelan, E.; Stoyanov, S. D. Remarkably high surface visco-elasticity of adsorption layers of triterpenoid saponins. *Soft Matter* **2013**, *9*, 5738-5752.

- (15) Pagureva, N.; Tcholakova, S.; Golemanov, K.; Denkov, N.; Pelan, E.; Stoyanov, S. D. Surface properties of adsorption layers formed from triterpenoid and steroid saponins, *Coll. Surf. A*, **2016**, *491*, 18-28
- (16) Wojciechowski, K.; Orczyk, M.; Marcinkowski, K.; Kobiela, T.; Trapp M.; Gutberlet, T.; Geue T. Effect of hydration of sugar groups on adsorption of Quillaja bark saponin at air/water and Si/water interfaces. *Colloids Surf. B*. **2014**, *117*, 60-67.
- (17) Jian, H.; Liao, X.; Zhu, L.; Zhang, W.; Jiang, J. Synergism and foaming properties in binary mixtures of a biosurfactant derived from *Camellia oleifera* Abel and synthetic surfactants. *J. Coll. Int. Sci.* **2011**, *359*, 487-492.
- (18) Wojciechowski, K.; Piotrowski, M.; Popielarz, W.; Sosnowski, T. R. Short and mid-term adsorption of Quillaja Bark saponin and its mixtures with lysozyme. *Food Hydrocolloids* **2011**, *25*, 687-693.
- (19) Piotrowski, M.; Lewandowska, J.; Wojciechowski, K. Biosurfactant-protein mixtures: Quillaja Bark saponins at the water/air and water/oil interfaces in the presence of β -lactaglobulin. *J. Phys. Chem. B* **2012**, *116*, 4843-4850.
- (20) Kezwon, A.; Wojciechowski, K. Interaction of Quillaja bark saponins with food-relevant proteins. *Adv. Coll. Int. Sci.* **2014**, *209*, 185-195.
- (21) Wojciechowski, K.; Orczyk, M.; Gutberlet, T.; Geue, T. Complexation of phospholipids and cholesterol by triterpenic saponins in bulk and in monolayers, *Biochimica et Biophysica Acta*, **2016**, *1858*, 363-373
- (22) Mitra, S.; Dungan, S. R. Micellar properties of Quillaja saponin. 1. Effects of temperature, salt, and pH on solution properties. *J. Agric. Food Chem.* **1997**, *45*, 1587-1595.
- (23) Oakenfull, D. G. Aggregation of saponins and bile acids in aqueous solution. *Aust. J. Chem.* **1986**, *39*, 1671-1683.
- (24) Lu, J. R.; Thomas, R.K.; Penfold, J. Surfactant layers at the air/water interface: structure and composition. *Adv. Coll. Int. Sci.* **2000**, *84*, 143-304.
- (25) Russell, T. P. X-ray and neutron reflectivity investigation of polymers. *Mat. Sci. Rep.* **1990**, *5*, 171-271.
- (26) Lu, J. R.; Zhao, X. B.; Yassen, M. Protein adsorption studied by neutron reflectivity. *Curr. Opin. Coll. Int. Sci.* **2007**, *12*, 9-16.
- (27) Zhang, X. L.; Penfold, J. Thomas, R. K.; Tucker, I. M.; Petkov, J. T.; Bent, J.; Cox, A.; Campbell, R. A. Adsorption behaviour of hydrophobin and hydrophobin / surfactant mixtures at the air-water interface. *Langmuir* **2011**, *27*, 11316-11323.
- (28) INTER reflectometer at the ISIS facility, <http://www.isis.stfc.ac.uk/instruments/>
- (29) Dukhin, S. S.; Kretschmar, G.; Miller, R. *Dynamics of Adsorption at Liquid Interfaces*; Elsevier: Amsterdam, 1995.
- (30) Adamson, A. W.; Gast, A. P. *Physical Chemistry of Surfaces*, 6th ed.; Wiley: New York, 1997.

- (31) Tsibranska, S.; Ivanova, A.; Tcholakova, S.; Denkov, N., Self-assembly of ESCIN molecules at the air-water interface as studied by molecular dynamics, *Langmuir*, **2017**, 33, 8330-8341
- (32) Rosen, M. J. *Surfactants and Interfacial Phenomena*; Wiley: New York, 1989.
- (33) Gurkov, T. D.; Dimitrova, D. T.; Marinova, K. G.; Bilke-Crause, C.; Gerber, C.; Ivanov, I. B. Ionic surfactants on fluid interfaces: determination of the adsorption; role of the salt and the type of the hydrophobic phase. *Colloids Surf. A* **2005**, 261, 29.
- (34) Li, P. X.; Li, Z. X.; Shen, H. H.; Thomas, R. K.; Penfold, J.; Lu, J. R. The use of the Gibbs equation to determine surface excess of non-ionic surfactants and polymers at the air-water interface, comparison with values determined directly using neutron reflection, *Langmuir*, **2013**, 29, 9324-9334
- (35) Xu, H.; Ma, K.; Li, P. X.; Thomas, R. K.; Penfold, J.; Lu, J. R. Limitations in the application of the Gibbs equation to anionic surfactants at the air-water interface: SDS and SLES above and at the cmc, *Langmuir*, **2013**, 29, 9335-9351
- (36) Penfold, J.; Thomas, R. K.; Shen, H. H. Adsorption of naturally occurring biosurfactants, glycolipids and lipopeptides studied by neutron reflection, *Soft Matter*, **2012**, 8, 578-591
- (37) Wade, D. Deuterium isotope effects on noncovalent interactions between molecules. *Chem. Biol. Interact.* **1999**, 117, 191-217.

ACKNOWLEDGEMENTS

The provision of beam time on the INTER reflectometer at ISIS is acknowledged. The invaluable scientific and technical assistance of the Instrument Scientists, and support staff is gratefully recognized. This study falls under the umbrellas of COST MP 1106 action “Smart and Green Interfaces” and of the FP7 project “Beyond Everest”. The authors are grateful to Ms. Nevena Borisova for the measurement of the surface tension isotherms.

SUPPORTING INFORMATION.

Some additional experimental data are available in the Supporting Information. This material is available free of charge via the internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author: Jeff Penfold, jeff.penfold@stfc.ac.uk

Author Contributions

All the authors have given their approval of the final version of the manuscript

Funding Sources

Funded through neutron beam time at the ISIS Facility, UK (STFC).

TABLE OF CONTENTS GRAPHICS

Saponin adsorption at the air-water interface

J Penfold^{1,2}, RK Thomas¹, I Tucker³, JT Petkov³, SD Stoyanov⁴, N Denkov⁵, K Golemanov^{4,5}, S Tcholakova⁵, JRP Webster²

