

The faint radio source population at 15.7 GHz – IV. The dominance of core emission in faint radio galaxies

I. H. Whittam¹,^{1,2}★ D. A. Green,³ M. J. Jarvis^{1,2} and J. M. Riley³

¹Department of Physics and Astronomy, University of the Western Cape, Robert Sobukwe Road, Bellville 7535, South Africa

²Department of Astrophysics, University of Oxford, Denys Wilkinson Building, Keble Road, Oxford OX1 3RH, UK

³Astrophysics Group, Cavendish Laboratory, 19 J. J. Thomson Avenue, Cambridge CB3 0HE, UK

Accepted 2020 January 29. Received 2020 January 29; in original form 2019 November 12

ABSTRACT

We present 15-GHz Karl G. Jansky Very Large Array observations of a complete sample of radio galaxies selected at 15.7 GHz from the Tenth Cambridge (10C) survey. 67 out of the 95 sources (71 per cent) are unresolved in the new observations and lower frequency radio observations, placing an upper limit on their angular size of ~ 2 arcsec. Thus, compact radio galaxies, or radio galaxies with very faint jets, are the dominant population in the 10C survey. This provides support for the suggestion in our previous work that low-luminosity ($L < 10^{25} \text{ W Hz}^{-1}$) radio galaxies are core dominated, although higher resolution observations are required to confirm this directly. The 10C sample of compact, high-frequency selected radio galaxies is a mixture of high-excitation and low-excitation radio galaxies and displays a range of radio spectral shapes, demonstrating that they are a mixed population of objects.

Key words: catalogues – surveys – galaxies: active – radio continuum: galaxies.

1 INTRODUCTION

Most deep studies of the extragalactic radio sky have focused on frequencies around 1 GHz, due to the increased telescope time required to conduct a survey at higher frequencies, leaving the faint, high-frequency sky as a relatively unexplored parameter space. One major exception to this is the Tenth Cambridge (10C) survey (AMI Consortium: Davies et al. 2011; AMI Consortium: Franzen et al. 2011; Whittam et al. 2016a) at 15.7 GHz, which is complete to 0.5 mJy in 10 different fields, and extended to 0.1 mJy in 2 of these fields, making it the deepest high-frequency radio survey to date.

In the first three papers in this series (Whittam et al. 2013, 2015, 2016b, hereafter Papers I, II, and III, respectively), we studied the properties of a complete sample of faint (~ 1 mJy) sources selected at 15.7 GHz from the 10C survey in the Lockman Hole. We have used a combination of observations at other radio frequencies and data from across the electromagnetic spectrum to extensively study the properties of these sources and found that they are essentially all (>94 per cent) radio-loud active galactic nuclei (AGNs) with a median redshift $z \sim 0.9$ (Paper II). While the brighter sources in this sample have steep radio spectra, as expected, the fainter radio galaxies have flatter spectra than previously assumed; the median spectral index¹ between 15.7 GHz and 610 MHz changes from 0.75 for sources with $S_{15 \text{ GHz}} > 1.5$ mJy to 0.08 for $S_{15 \text{ GHz}} <$

0.8 mJy (Paper I). This suggests that these fainter radio galaxies are dominated by emission from the core, which generally has a flatter radio spectrum (due to the superposition of several synchrotron self-absorbed components from the base of the jet) than the optically thin synchrotron emission from any extended lobes. Some of the sources display extended morphologies typical of Fanaroff and Riley type I or II (FRI or FRII) sources (Fanaroff & Riley 1974), based on 610-MHz Giant Meterwave Radio Telescope (GMRT) observations with a resolution of 5 arcsec (Paper III). The majority (68 out of 96, 71 per cent), however, appear to be compact radio galaxies (Paper III), which are referred to as ‘FR0’ sources. [The FR0 classification was introduced by Ghisellini et al. (2011) to describe the class of weak, compact radio sources described by Baldi & Capetti (2009), but also studied by Slee et al. (1994) and possibly even earlier.] The sources are a mixture of high-excitation and low-excitation radio galaxies (HERGs and LERGs; see e.g. Best & Heckman 2012), and the HERGs tend to have flatter spectra and be more core dominated than the LERGs (Paper III).

Most models of the high-frequency extragalactic radio sky (e.g. de Zotti et al. 2005; Wilman et al. 2008; Tucci et al. 2011) are based on extrapolations from lower frequencies, and are a poor fit to the observed 15-GHz source counts below $\lesssim 10$ mJy. One such model is the SKA Simulated Skies (S³; Wilman et al. 2008, 2010), which is a set of simulations of the radio and sub-mm universe. One component is a semi-analytic simulation of a $20 \times 20 \text{ deg}^2$ patch of the extragalactic sky out to a redshift of 20, containing 320 million radio components. While this simulation reproduces the observed source distributions at lower frequencies, it underpredicts

★ E-mail: imogen.whittam@physics.ox.ac.uk

¹The convention $S \propto \nu^{-\alpha}$, for flux density S , frequency ν , and spectral index α , is used throughout this work.

the number of sources observed in the Ninth Cambridge (9C; Waldrum et al. 2003, 2010) and 10C surveys with $S_{18\text{ GHz}} < 10\text{ mJy}$ by a factor of 2, and fails to reproduce the observed spectral index distribution. In Whittam et al. (2017), we showed that by applying a simple modification to the simulation and assuming that the cores of the fainter FRI sources in the simulation are more dominant than previously thought, the observed high-frequency source counts can be reproduced. The observations are best matched by assuming that the fraction of the total 1.4-GHz flux density that originates from the core varies with 1.4-GHz luminosity; sources with 1.4-GHz luminosities $< 10^{25}\text{ W Hz}^{-1}$ require a core fraction $S_{\text{core}}/S_{\text{tot}} \sim 0.3$, while the more luminous sources require a much smaller core fraction of 5×10^{-4} . The more recent Tiered Radio Extragalactic Continuum Simulation (T-RECS; Bonaldi et al. 2019) models steep-spectrum sources, flat-spectrum radio quasars, and BL Lacs separately and is a significantly better fit to the observed high-frequency source counts. The primary reason for the better fit to the data is the inclusion of a significant population of low-powered radio-loud AGNs with flat spectra, which are present in the observed 10C sample but not included in S^3 . This is discussed further in Section 6.

In this paper, we describe new, higher resolution, observations of a subsample of 10C sources made with the Karl G. Jansky Very Large Array (VLA). The observations were made at 15 GHz in C configuration, so are matched in frequency to the original 10C observations but have a significantly higher resolution of ~ 2 arcsec compared to 30 arcsec for 10C. They allow us to constrain the fraction of the total flux density originating from the cores of the radio galaxies, and therefore test our proposed modifications to S^3 .

These new observations also allow us to further investigate the properties of this unique sample of high-frequency selected faint, compact radio galaxies. Despite the fact that they dominate lower powered radio AGN samples (e.g. Sadler et al. 2014; Baldi, Capetti & Massaro 2018), the nature of compact radio galaxies (FR0 sources) is not well understood. While some compact radio galaxies are young sources that may go on to become more extended FRI or FRII galaxies, the number density of compact sources compared to that of extended sources implies that most must be older sources that have failed to produce large extended structures (Sadler et al. 2014; Baldi et al. 2018). Baldi, Capetti & Giovannini (2015) found no difference in the properties of the host galaxies (e.g. optical morphology, colour, stellar mass) of FRI sources compared to the more compact FR0 sources, although there is more recent evidence that FR0 sources are hosted by less massive galaxies than FRIs (Miraghaei & Best 2017; Baldi et al. 2018). The reason for the lack of extended emission in some radio galaxies is therefore not clear; possibilities include intermittent activity of the central engine (Reynolds & Begelman 1997; Conway 2002; Czerny et al. 2009), jets that are easily interrupted due to a lower bulk speed, or interactions with surrounding gas (e.g. Baldi, Capetti & Giovannini 2015, 2019; Sadler 2016; Torresi et al. 2018).

In Section 2, we outline the sample selection and the radio and multiwavelength data that are already available. Section 3 describes the new VLA observations and data reduction, and in Section 4 we update the multiwavelength counterparts associated with the radio sources based on these observations. The new observations are compared to the 10C observations in Section 5. In Section 6, we examine our measurements of radio core dominance in relation to the modification to S^3 we suggested in Whittam et al. (2017) and the T-RECS simulation, and discuss what these observations tell us about the properties of faint, compact radio galaxies. We then present our conclusions in Section 7.

Throughout this paper, we assume $H_0 = 70\text{ km s}^{-1}\text{ Mpc}^{-1}$, $\Omega_M = 0.3$, and $\Omega_\Lambda = 0.7$ and use J2000 coordinates.

2 SAMPLE USED

This work is based on a sample selected from the 10C survey at 15.7 GHz. The 10C survey was observed with the Arcminute Microkelvin Imager (AMI; Zwart et al. 2008) Large Array and covers 27 deg^2 complete to 1 mJy in 10 different fields, and a further 12 deg^2 complete to 0.5 mJy contained within these fields. Here, we use a complete sample of 96 sources selected from a region of the 10C survey in the Lockman Hole with particularly good low-frequency data available. These 96 sources form a subsample of the full 10C Lockman Hole sample discussed in Paper I; full details of the sample selection are given in Paper II.

In Paper I, the sample was matched to several lower frequency (and higher resolution) radio catalogues available in the field. These are a deep 610-MHz GMRT image (Garn et al. 2008, 2010), a 1.4-GHz Westerbork Synthesis Radio Telescope (WSRT) image (Guglielmino et al. 2012; Prandoni et al. 2018), two deep VLA images at 1.4 GHz that only cover part of the field (Biggs & Ivison 2006; Owen & Morrison 2008), the National Radio Astronomy Observatory (NRAO) VLA Sky Survey (NVSS; Condon et al. 1998), and the Faint Images of the Radio Sky at Twenty Centimetres (FIRST; White et al. 1997). All but one of the 96 sources in this sample have a match in at least one of the lower frequency catalogues, allowing radio spectral indices and sizes to be found for these sources.

There is a wealth of multiwavelength data available in the Lockman Hole, most of which is contained in the ‘SERVS Data Fusion’ catalogue (Vaccari 2015), a multiwavelength catalogue selected in the mid-infrared. In the Lockman Hole, this catalogue contains data from the *Spitzer* Wide-Area Infrared Extragalactic Survey (SWIRE; see Lonsdale et al. 2003), the *Spitzer* Extragalactic Representative Volume Survey (SERVS; see Mauduit et al. 2012), and the United Kingdom Infrared Telescope Infrared Deep Sky Survey (see Lawrence et al. 2007) with deep optical photometric data taken by González-Solares et al. (2011) at the Isaac Newton Telescope and the Kitt Peak National Observatory. In order to be included in the catalogue, a source must be detected in either the *Spitzer* IRAC1 or IRAC2 bands. In Paper II, the radio sources in the sample were matched to the SERVS Data Fusion catalogue; these matches are reviewed in Section 4 with the addition of the new, higher resolution, radio data presented here.

There are also two X-ray surveys that each cover part of this field: one with *Chandra* (Wilkes et al. 2009) that covers 0.7 deg^2 and one with *XMM-Newton* (Brunner et al. 2008) that covers $\sim 0.2\text{ deg}^2$. The 10C sources are matched to these two catalogues in Paper III.

The 96 sources in this sample have 15.7-GHz flux densities in the range $0.5 < S_{10C}/\text{mJy} < 45$ and a redshift distribution peaking at $z \sim 1$ (Paper II). This sample probes a wide luminosity range $10^{21} < L_{1.4\text{ GHz}}/\text{W Hz}^{-1} < 10^{28}$.

3 OBSERVATIONS AND DATA PROCESSING

3.1 The VLA observations

The new data² were taken with the Karl G. Jansky VLA in C configuration in the Ku band. Pointed observations were made of

²Project code: 17A-183.

the 96 sources in the sample described in Section 2. The 75 sources with $S_{10C} > 0.75$ mJy (S_{10C} is the 15-GHz peak flux density from the 10C catalogue) were observed for 50 s on source. The fainter sources with $S_{10C} < 0.75$ mJy (21 sources) were observed for a time scaled according to the 10C peak flux density of each source, to allow the fainter sources to be observed for longer. The time for each source was calculated to achieve an rms noise level of $S_{10C}/25$. For example, the faintest source in the sample has $S_{10C} = 0.35$ mJy and in order to achieve the required rms noise level of $14 \mu\text{Jy}$ per beam it was observed for 187 s.

The observations were split into two scheduling blocks (SBs) containing 42 and 54 sources, and these were observed on 2017 May 31 and 2017 June 11, respectively. 3C 286 was observed once at the start of each SB for flux density and bandpass calibration. J1035+5628 was observed approximately every 15 min for phase calibration. The Ku band receiver was used with a central frequency of 15 GHz and 48×128 MHz spectral windows in full polarization.

The data were initially processed using the NRAO VLA pipeline,³ which consists of a series of CASA⁴ commands. The pipeline performs some initial flagging; this includes the removal of the initial few integration points of a scan where the antennas may not all be on source, the removal of data from shadowed antennas, and automatic radio frequency interference flagging. The pipeline then performs delay and bandpass corrections.

After the execution of the NRAO pipeline, the observations were averaged using 8 s integration times. The sources were imaged using the CASA task CLEAN. For the five sources with significant extended emission, clean boxes were placed by hand; the remaining sources were imaged automatically using 500 iterations, with cycle factor = 5. Sources with a signal-to-noise ratio >20 were self-calibrated. Two rounds of self-calibration were performed, both phase-only, with a first solution interval of 60 s and a second of 30 s. In some cases, errors were reported by the CASA task GAINCAL due to insufficient signal-to-noise ratio; for these sources, the non-self-calibrated images were used. The signal-to-noise ratio in each image was compared after each stage of self-calibration; if it went down at any point, then the non-self-calibrated images were used. The final resolution is $\approx 1.6 \times 2.4 \text{ arcsec}^2$ for sources observed in the first SB and $\approx 1.6 \times 4.0 \text{ arcsec}^2$ for those observed in the second SB. The resolutions were different for the two SBs as they were observed at different elevations. The nominal largest scale structure for the VLA in C array at 15 GHz is 97 arcsec, but as these are snapshot observations the largest angular scale they are sensitive to is around 50 arcsec.

3.2 Source properties

The flux density of each source was estimated by fitting one or more Gaussians to each source using the CASA task IMFIT. 22 sources were resolved by these observations. 17 of these are significantly extended, so were poorly fit by a Gaussian; the integrated flux densities of these sources were found interactively using the CASA task IMVIEW. In cases where the core and extended emission could be distinguished from each other, the flux density of the core was also found separately. The ratio of integrated to peak flux densities is shown in Fig. 1 as a function of signal-to-noise ratio; it is clear that many of the sources in this sample are unresolved on arcsec




Figure 1. Ratio of the integrated to peak flux densities as a function of signal-to-noise ratio (σ_{local} = local rms noise). The red circles are sources with more than one component.

scales. The 17 sources resolved into more than one component are indicated by red circles in Fig. 1.

For single-component sources, angular sizes were estimated using the Gaussian fit from IMFIT. For those that could be resolved, the deconvolved full width at half-maximum (FWHM) major axis was used as a measure of the angular size. If, however, the source could not be deconvolved by IMFIT, the FWHM major axis of the restoring beam was taken as an upper limit on the angular size of the source. This is a conservative upper limit, as a source must be intrinsically smaller than this to appear unresolved. This was the case for 73 sources (26 in SB 1 and 47 in SB 2). As the synthesized beam size and shape are different for the two SBs ($1.6 \times 2.4 \text{ arcsec}^2$ compared to $1.6 \times 4.0 \text{ arcsec}^2$), there are different upper limits on the angular sizes of the unresolved sources observed in the different SBs. Note that for the second SB the unresolved sources are probably smaller than the 4 arcsec upper limit suggesting that the synthesized beam is highly elliptical and the sources are not resolved in either direction.

For the 17 sources resolved into more than one component, the angular size was measured from the contour map by taking the maximum straight line extent of the source within the 3σ contours. The angular sizes of the sample of 95 sources measured from the 15-GHz VLA observations are shown in Fig. 2 and listed in Table 1.

Six of the sources that are unresolved in the new VLA observations have extended ($>10 \text{ arcsec}$) emission visible in the lower frequency observations (at 1.4 GHz and/or 610 MHz, described in Section 2) and are therefore clearly not genuinely compact. Therefore, 28 sources (22 resolved at 15 GHz and a further 6 resolved at lower frequencies) in this sample are extended and 67 are compact on arcsec scales. Since the extended emission typically has a steep spectrum, the lower frequency observations provide a better measure of any extended emission (where present) than the 15-GHz observations. Therefore, when considering the overall linear sizes of the sources, the measurements from the lower frequency surveys (defined in section 6.4 of Paper II) are used for the extended sources. For the sources that are compact in both the lower frequency and 15-GHz observations (67 sources), the 15-GHz measurement is used as an upper limit on the size of the source as this provides a smaller limit. However, note that it is possible that these sources may have some extended emission that is resolved out of the new VLA observations; this is discussed further in Section 5.

³<https://science.nrao.edu/facilities/vla/data-processing/pipeline>

⁴<http://casa.nrao.edu/>




Figure 2. Distribution of angular sizes measured from the 15-GHz VLA images for all the sources in the sample. Upper limits are shown in white. Note that six sources that are unresolved in these observations have extended emission visible in 1.4-GHz and/or 610-MHz observations and are therefore not genuinely compact.

In [Paper III](#), the sources in this sample were classified according to their morphology and radio spectra using 610-MHz GMRT data with a resolution of 5×6 arcsec². Each source was classified as FR0, FR0/I, FRI, FRI/II, or FRII based on its morphology; the compact FR0 sources were further classified based on their radio spectra and linear sizes as FR0c [candidate compact steep spectrum (CSS) sources], FR0g [candidate GHz-peaked spectrum (GPS) sources], or FR0u (uncertain) (detailed descriptions of the classification criteria are given in [Paper III](#)). These classifications were reviewed in light of the new data by examining each source by eye. Nine sources that were previously classified as FR0u are resolved by the new observations and therefore reclassified. A further three sources with uncertain classifications (FR0/I or FRI/II) are also reclassified. Note that none of the sources previously identified as candidate CSS or GPS sources are resolved in the new observations, providing support for these classifications. For the six sources that appear compact in

the new observations where only the radio core is detected but that have extended emission visible at 610 MHz, the original 610-MHz FRI or FRII classifications are retained.

When examining all available images, we realized that two nearby sources (10C J105144+573313 and 10C J105148+573245, see [Fig. 3](#)) that had previously been classified as two FR0 sources are more likely the core and hotspot of one source. We have therefore combined these sources into one entry in the catalogue, labelled as 10C J105148+573245, as this source is the core. The FR classification has also changed for one other source, 10C J105009+570724, which was previously thought to be one FRI source but is now classified as two separate FR0 sources. This is discussed further in Section 4. These two cases illustrate the difficulties in correctly classifying and identifying host galaxies for radio sources.

The properties of the sources in this sample derived from the new VLA observations are summarized in [Table 1](#), which is available in the online supplementary material in machine-readable format. Images of the 22 sources that are resolved in the observations are shown in [Fig. 3](#).

3.3 False detection

One source in the sample, 10C J104927+583830, was not detected in the new VLA observations. This source had a peak flux density of 0.45 ± 0.08 mJy per beam in the 10C catalogue but is undetected in the new image that has an rms noise of 0.018 mJy, giving a 5σ upper limit on the peak flux density of the source of 0.09 mJy per beam. This suggests that if the source is real, its peak flux density must have decreased by more than a factor of 5 over a ~ 9 -yr period.

However, we now believe that this source is a false detection. It is the only source in the sample that was not detected in any of the other lower frequency radio catalogues that cover this field, and it also does not have a counterpart in the SERVS Data Fusion multiwavelength catalogue ([Paper II](#)). When extracting sources for the 10C catalogue, exclusion zones were defined around bright (> 15 mJy) sources due to the increased noise close to these sources caused by amplitude, phase, and deconvolution errors (see AMI

Table 1. Properties of the sources in this sample based on the new VLA observations. A machine-readable version of this table is available online as supplementary material.

10C ID (1)	RA (2)	Dec. (3)	S_{peak} (mJy) (4)	ΔS_{peak} (mJy) (5)	S_{int} (mJy) (6)	ΔS_{int} (mJy) (7)	N_{comp} (8)	rms (μ Jy per beam) (9)	S_{core} (mJy) (10)	θ (arcsec) (11)	FR class (12)
10C J104320+585621	10:43:20.515	+58:56:20.780	1.043	0.022	1.052	0.046	1	34		<4.6	FRII
10C J104328+590312	10:43:28.435	+59:03:14.080	1.193	0.020	1.262	0.044	1	37		<4.0	FR0u
10C J104344+591503	10:43:44.677	+59:15:02.550	2.686	0.020	2.724	0.042	1	27		<4.7	FR0u
10C J104428+591540	10:44:28.382	+59:15:41.490	0.512	0.020	0.571	0.045	1	35		<5.1	FR0g
10C J104441+591949	10:44:41.132	+59:19:46.100	0.643	0.019	0.639	0.038	1	26		<4.5	FR0g
10C J104451+591929	10:44:50.388	+59:19:26.350	1.116	0.023	1.533	0.055	1	29		<4.0	FR0u
10C J104528+591328	10:45:28.206	+59:13:26.190	1.768	0.022	1.800	0.045	1	27		<4.5	FR0c
10C J104539+585730	10:45:39.678	+58:57:29.550	0.417	0.046	0.731	0.203	1	32		<5.1	FRII
10C J104551+590838	10:45:50.787	+59:08:41.420	0.695	0.016	0.757	0.036	1	23		<4.0	FR0u
10C J104624+590447	10:46:24.841	+59:04:45.760	0.896	0.025	1.214	0.059	1	23		<4.0	FR0u

Notes: (1) Source name from the 10C catalogue. (2, 3) Right ascension and declination of peak flux in VLA image (J2000). (4) Peak flux density in VLA image. (5) Uncertainty in (4). (6) Integrated flux density of source in VLA image. (7) Uncertainty in (6). (8) Number of components source is resolved into. (9) Local rms noise in VLA image, measured in a 100×100 pixel box away from the source. (10) Flux density of core component; left blank if core is not distinguishable from the extended emission. (11) Angular size of source measured at 15 GHz (see Section 3.2); upper limits are marked with '<'. (12) Morphological classification.




Figure 3. The new 15-GHz VLA observations of the 22 sources resolved in the VLA observations.

Consortium: Davies et al. 2011 for details). This source is close to a bright source and fell just outside one such exclusion zone; the exclusion zone in question had a radius of 3.35 arcmin and this source is located 3.42 arcmin from centre of the zone. On examining

the original 10C map, the noise level appears to still be elevated in the region around this source. Thus, all the evidence indicates that this source is a false detection, and it is therefore removed from the sample discussed in the remainder of this paper.




Figure 3 – continued

4 MULTIWAVELENGTH COUNTERPARTS

This sample was matched to the SERVS Data Fusion multiwavelength catalogue in [Paper II](#), where we found that 80 out of the 95 sources had a multiwavelength counterpart. Here, we review these matches with the additional information provided by the new, higher resolution 15-GHz VLA data. In order to check the original matching process, we overlaid contours of the new VLA images on to optical images of the area around each source, along with the positions of entries in the SERVS Data Fusion catalogue.

Eight sources were classified as ‘confused’ when matching in [Paper II](#), as there were several possible optical counterparts within the 3σ (where σ = local rms noise) radio contours and we were unable to decide upon the correct match. With the higher resolution VLA data, we are now able to identify the multiwavelength counterpart for all eight of these sources. Example images of two of these sources are shown in Appendix A. One previously confused source, 10C J105009+570724, is resolved into two separate components in the new data, each with its own separate optical counterpart (see Fig. A2). We therefore split this source into two separate entries in the updated catalogue. This source was previously classified as an FR II source, and is now classified as two separate FR 0 sources. Both of these sources have integrated flux densities at 15 GHz greater than 0.5 mJy, the limit originally used to select the sample.

For source 10C J105327+574546, with the additional information provided by the higher resolution radio data, we now believe the original match is incorrect. This is discussed in more detail in Appendix A. Additionally, two sources that were originally classified as not having a match now have a counterpart in the

multiwavelength catalogue. For the remaining 84 sources, the new data confirm the original classification.

In summary, there are now 95 sources in the sample. Of the original 96 sources, one has been removed (as we believe it is a false detection), and one has been split into two sources, adding one source. A further two have been combined into one source as we now believe they are components of the same radio galaxy. 89 out of 95 sources now have a multiwavelength counterpart. There are 11 original sources with a new match: 10 that previously did not have a match (8 ‘confused’ and 2 ‘no match’) and 1 where the previous match was incorrect. One of these original source is actually 2 separate sources, both of which have a match, meaning there are a total of 12 sources with a new match.

4.1 Updated redshifts

Spectroscopic redshifts are available for four of the new optical counterparts. For the remaining eight new matches, photometric redshift values are used. These are assigned in the same way as described in [Paper II](#). Four sources have photometric redshifts in the Rowan-Robinson et al. (2013) catalogue, so these values are used, and for the remaining sources photometric redshifts are estimated using the publicly available LE PHARE code (Arnouts et al. 1999; Ilbert et al. 2006; Arnouts & Ilbert 2011); see section 4 of [Paper II](#) for full details of this process.

Additionally, spectroscopic redshifts are now available for five more objects that only had photometric redshift estimates when [Paper II](#) was published. We therefore now have spectroscopic redshifts for 33 sources.

4.2 Updated HERG/LERG classification

The 12 sources with new multiwavelength counterparts are classified as HERGs or LERGs in the same way as described in Paper III. Three of these sources have optical spectra available, so information from these spectra is used to classify them. For the remaining nine sources, a combination of mid-infrared colour–colour diagrams, optical compactness, and X-ray information is used to classify the sources in the same way as outlined in section 3 of Paper III. This resulted in one new HERG and nine new LERGs; the remaining two sources with a new match did not have enough information available to be classified.

Eight sources that were classified as HERGs or LERGs based on their mid-infrared colours/optical compactness/X-ray information in Paper III now have optical spectra available, providing an additional check on these classifications. For seven of these eight sources, the classifications from the optical spectra confirm the classifications given in Paper III (three HERGs and four LERGs). One source (10C J105142+573447), however, was classified as an HERG based on its X-ray emission in Paper III, but its optical spectrum clearly resembles that of a low-excitation galaxy. Mingo et al. (2014) found that 4 of the 46 sources in their sample are classified as LERGs based on their optical emission lines but display HERG-like behaviour in the X-ray. As classifications based on optical spectra take precedence in our classification scheme, we have changed the classification of this source from HERG to LERG in the updated catalogue.

The numbers of sources with the different classifications are therefore as follows: 32 HERGs, 43 LERGs, and 20 unclassified sources (95 total). A table summarizing the properties of the multiwavelength counterparts and redshift values of all the sources in this sample is available online as supplementary material. A description of the table is given in Appendix B.

5 COMPARISON WITH THE 10C SURVEY

The observations contributing to the 10C survey were taken between 2008 August and 2010 June and have a synthesized beam size of 30 arcsec, while the VLA observations were taken in 2017 May and June and have a synthesized beam size of ~ 2 arcsec. Both sets of observations are at 15 GHz, so any differences in the flux densities between the two sets of observations could be due to (i) uncertainties in the measurements, (ii) the different resolutions of the observations, or (iii) intrinsic source variability.

The flux densities of the sources in the 10C survey and the new VLA observations are compared in Fig. 4. Note that for the purposes of comparing the two surveys sources 10C J105009+570724A and 10C J105009+570724B are combined as they cannot be separated in the 10C data. Although the values from the two sets of observations are broadly consistent for the majority of sources, for nearly one-third of the sources the values differ by more than 3σ , suggesting that another factor besides the uncertainties in the measurements is contributing towards these differences.

Differences in the measured flux densities between the two sets of observations could be due to the different beam sizes and array configurations of the AMI telescope and the VLA in C configuration. It is possible that diffuse emission from the sources may have a surface brightness too low to be detected in the VLA observations, or be resolved out, resulting in a measured VLA flux density that is lower than the measured 10C flux density. Low surface brightness emission is clearly more likely to be missed at low signal-to-noise ratio, so this may well explain the large positive




Figure 4. Comparison of flux densities in the original 10C catalogue and the new VLA observations. Integrated flux densities are used for the VLA observations, and in the 10C observations peak flux density is used for unresolved sources and integrated flux density for resolved sources, in the same way as in the 10C survey (see AMI Consortium: Davies et al. 2011). Sources where the two values differ by more than 3σ are marked by red circles.

flux ratios seen for several sources at low signal-to-noise ratio. This suggests that although these sources appear unresolved in both the 10C and VLA observations, they may have some extended emission on scales between 2 and 30 arcsec. 14 unresolved sources have a 10C flux density that is larger than the VLA flux density by more than 3σ , so may have some emission on these scales. There is evidence for this from lower frequency observations of the 10C sample; 2 of the 14 unresolved sources with 10C flux densities that are larger by more than 3σ are extended in either the GMRT (610 MHz, resolution 6 arcsec; Garn et al. 2008, 2010) or WSRT (1.4 GHz, resolution 11 arcsec; Prandoni et al. 2018) observations (see Paper I for further details of these observations). A further four sources that are compact in both the 10C and VLA observations have extended emission detected at lower frequencies; the flux densities of these sources from the two sets of observations agree within 3σ .

Alternatively, other nearby sources in the AMI beam may be contributing to the measured flux, causing it to be larger in the 10C observations than in the new observations. However, if a source were bright enough to make a significant contribution to the measured flux of a nearby source, we would expect it to be detected in the new VLA observations. In one case, 10C J105009+570724, the 10C detection clearly has contributions from two separate sources as already discussed, both of which are detected in the VLA observation. None of the other sources with higher flux densities in the 10C observations than the VLA observations show evidence of nearby sources in the VLA image.

There was an interval of ~ 9 yr between the two sets of observations, so the flux density differences seen could be due to variability. While there have been no studies of the variability of 15-GHz sources in our flux density range, Bolton et al. (2006) studied the variability of 51 9C sources with $S_{15\text{ GHz}} > 25$ mJy over a 3-yr period and found that 29 per cent of sources showed some variability. Sadler et al. (2014) found that one-third of the FR0 sources in their sample of nearby 20-GHz selected sources varied by more than 10 per cent over a time period of between 10 and 15 yr. We therefore expect some variability for the sources in our sample, so this could be the cause of some of the flux density differences seen. Flat-spectrum sources are expected to be more




Figure 5. Ratio of flux densities in the 10C and VLA images as a function of spectral index. For the 10C flux densities, peak flux densities are used for sources that appear point-like in the 10C observations, and integrated flux densities are used otherwise, in the same way as in AMI Consortium: Franzen et al. (2011). For the VLA data, integrated flux densities are used for all sources. The dashed line indicates where the ratio of the flux densities from the two surveys is equal to 1, and the dotted line is at $\alpha = 0.5$, the standard value used to classify sources as having either steep ($\alpha > 0.5$) or flat ($\alpha < 0.5$) spectra. Source 10C J105009 + 570724, which has been split into two, has been recombined for the purpose of this comparison as the two sources are not resolved separately in the 10C observations.

variable than those with steep spectra, so if variability was the main reason for the difference in the flux densities between the two sets of observations, we would expect these differences to be larger for sources with flat spectra. Fig. 5 shows the ratio of flux densities as a function of spectral index; the number of steep- and flat-spectrum sources having flux densities that differ by more than 3σ is very similar (15 flat-spectrum sources, defined as having $\alpha < 0.5$, and 16 steep-spectrum sources, with $\alpha > 0.5$). However, four out of the five sources with flux density values that vary by more than 10σ have flat spectra, suggesting that intrinsic source variability could be the cause of these large differences.

In summary, the differences in the measured flux densities between the two sets of observations displayed by some sources could be due to intrinsic source variability or extended emission on scales between 2 and 30 arcsec not detected in the VLA observations (or a combination of these two factors). Further observations sensitive to emission on scales between 2 and 30 arcsec are required to distinguish between these two scenarios.

6 THE DOMINANCE OF CORE EMISSION IN FAINT RADIO GALAXIES

While there have been several studies of the core emission in powerful radio galaxies (e.g. Broderick & Fender 2011), the cores of fainter radio galaxies ($S \lesssim 1$ mJy) have not been well studied. In Whittam et al. (2017), we showed that if the cores of less-powerful ($L_{1.4 \text{ GHz}} < 10^{25} \text{ W Hz}^{-1}$) FRI galaxies are more dominant than assumed in the S^3 (Wilman et al. 2008, 2010), the simulated source counts are a much better fit to observations at higher radio frequencies ($\gtrsim 15$ GHz).

As described fully in Whittam et al. (2017) and summarized here, we found that the observed source counts at >15 GHz can be better reproduced by making the following simple modification to the simulated catalogue. A fraction x of the total 1.4-GHz flux density of each FRI source in the simulation is assumed to be in




Figure 6. Core fraction at 15 GHz as a function of 1.4-GHz luminosity for extended sources with an identifiable core. The dashed line is the core fraction at 15 GHz used in our proposed modification to the S^3 (Whittam et al. 2017) and the dotted line is the mean core fraction in the original simulation. These values are extrapolated to 15 GHz by assuming the spectral indices of the core emission and extended emission are 0.0 and 0.75, respectively.

the core, and assigned a spectral index between 1.4 and 18 GHz of $\alpha = 0$. The remaining flux density is assumed to be extended emission and therefore given a spectral index of 0.75. This is then used to produce a revised flux density for each FRI source at 4.8 and 18 GHz. We found that the best-fitting values for x were 0.31 for FRI sources with $L_{1.4 \text{ GHz}} < 10^{25} \text{ W Hz}^{-1}$ and 2.5×10^{-3} for sources with $L_{1.4 \text{ GHz}} > 10^{25} \text{ W Hz}^{-1}$. This proposed modification is clearly a simplification, and for example does not include effects such as spectral ageing; however, it demonstrates that the high-frequency source counts can be better reproduced by assuming that the FRI sources in the simulation with $L_{1.4 \text{ GHz}} < 10^{25} \text{ W Hz}^{-1}$ are more core dominated than previously assumed in the model used to generate the simulation. The observations described in this paper allow us to measure the core dominance in this sample of faint, high-frequency selected radio galaxies and therefore test this proposed modification.

Out of 95 sources in the sample, 17 sources are resolved into more than one component and 12 of the 17 have an identifiable core. For these sources, we can directly measure the core fraction, x , defined as $x = S_{\text{core}}/S_{\text{tot}}$. These core fractions are compared to those from the modified S^3 presented in Whittam et al. (2017) in Figs 6 and 7. The modified S^3 core fractions are average values for the whole population; clearly we expect the values for individual sources to display considerable scatter. Most of the lower luminosity 10C sources (with $L < 10^{25} \text{ W Hz}^{-1}$) are not resolved in the VLA observations, so cannot be included in this plot. These low-luminosity sources generally have flatter spectra (as demonstrated in Paper I) and therefore are likely to have higher core fractions. The low-luminosity sources that can be plotted clearly have larger core fractions than assumed in the original simulation, providing support for our proposed modification to the simulation. Note, however, that the more luminous sources also seem to have larger core fractions than assumed in either the original simulation or the modified version.

The newly released T-RECS simulation (Bonaldi et al. 2019) is a much better fit to the observed high-frequency source counts than S^3 (see fig. 6 of Bonaldi et al. 2019). While Bonaldi et al. do not provide the information required to calculate core fractions, we are able to compare the spectral indices of the sources in the T-RECS




Figure 7. Core fraction at 15 GHz as a function of spectral index between 1.4 and 15 GHz for sources with an identifiable core. The core fractions at 15 GHz used in our proposed modification to the S^3 (Whittam et al. 2017) are plotted as squares (red in the online version), and the mean value in the original simulation is shown as a diamond (orange in the online version). These values are extrapolated to 15 GHz by assuming the spectral indices of the core emission and extended emission are 0.0 and 0.75, respectively.




Figure 8. Median spectral index in a series of 15-GHz flux density bins. The median values from the full Lockman Hole 10C sample are shown, along with the original and modified S^3 simulation and the T-RECS simulation.

simulated catalogue to those in S^3 and in our 10C sample, as shown in Fig. 8. Below 1 mJy, the T-RECS spectral indices are close to those measured from the 10C sample, which is consistent with the fact that T-RECS is able to reproduce the observed high-frequency source counts, while S^3 is not. For the bright sources, however, the T-RECS median spectral indices appear to be lower (i.e. the spectra are flatter) than those in the 10C sample.

The spectral shapes of the sources in our sample are illustrated by the radio colour–colour plot shown in Fig. 9, where the sources that are resolved in the new VLA observations are shown separately. The majority of the resolved sources have steep spectra (as expected), while the unresolved sources display a range of different spectral shapes. In particular, a number of compact sources are found in the bottom left and right quadrants, where high-frequency peakers and GPS sources, respectively, are expected to lie. This suggests that these sources could be young radio galaxies.




Figure 9. Radio colour–colour diagram. Sources that are resolved and unresolved in the new 15-GHz VLA observations are shown separately. The six sources that are unresolved in the new observations but are extended at 1.4 GHz/610 MHz are shown as cyan crosses. Sources with an upper limit on their 610-MHz flux density are marked with black triangles (these are all unresolved in the 15-GHz and lower frequency observations).

The 1.4-GHz luminosities of the radio galaxies in this sample are shown in Fig. 10 as a function of linear size, in which the unresolved compact sources are shown as triangles. The resolved sources all have 1.4-GHz luminosities greater than 10^{24} W Hz $^{-1}$, while the unresolved sources span the full range of luminosities $10^{21} < L_{1.4 \text{ GHz}}/\text{W Hz}^{-1} < 10^{29}$. Evolutionary tracks from An & Baan (2012) for FRI and FRII sources are shown in Fig. 10; the majority of the sources lie below the FRI track, although since most of these sources have upper limits on their linear sizes they could in fact move to the left and lie in the region of the diagram indicative of young sources. Note that many of the FRII sources in our sample have relatively low luminosities, lying well below the evolutionary track for FRII sources.

Similar numbers of the compact sources are classified as LERGs and HERGs (23 and 27 out of 67 sources, respectively), while a far greater proportion of the extended sources are classified as LERGs than HERGs (19 are LERGs, while only 5 are HERGs).

While the extended sources mostly have $L_{1.4 \text{ GHz}} > 10^{24}$ W Hz $^{-1}$, steep spectra, and are LERGs, the compact 10C sources display a range of spectral shapes and radio powers and are a mixture of HERGs and LERGs. This suggests that they are a composite population containing a mixture of different source types. 31 per cent of the FR0 sources in this sample are candidate GPS or CSS sources, implying that they may be young radio galaxies that are yet to develop powerful extended emission. Alternatively, the compact sources may be intermittent AGNs that are currently in an active phase, but this active phase is not sustained for long enough to develop extended radio jets. Another possibility is that the intrinsic properties of the jets (e.g. lower bulk speeds) may mean they are more easily disrupted, preventing them from forming extended structures. Further study is required to distinguish between these scenarios.




Figure 10. Radio power–linear size diagram. The lines are example evolutionary tracks for FRI (dashed line) and FRII (dotted line) sources from An & Baan (2012). For the compact sources, sizes are measured from the VLA observations and for the extended sources they are taken from the lower frequency observations. Triangles indicate upper limits on linear sizes. In the top panel, HERGs, LERGs, and unclassified sources are shown separately. In the bottom panel, sources are coloured according to their FR morphological classification; these classifications are based on 1.4-GHz/610-MHz observations and are described in Paper III.

It has been suggested (e.g. Sadler 2016) that some FR0 sources do have extended radio jets, but the emission is below the surface brightness limit of most large-scale radio surveys, meaning that these sources appear compact. Shabala et al. (2017) modelled the emission from three AGNs and showed that the surface brightness of their lobes would be too low to be detected by most surveys with the VLA and the Low Frequency Array (LOFAR). This idea is supported by the fact that, as discussed in Section 5, there is evidence that some of the unresolved sources in our sample may have emission on scales between 2 and 30 arcsec, the surface brightness of which is too low to be detected in our VLA observations. The MeerKAT radio telescope in South Africa would be the ideal instrument with which to investigate this as the large number of short baselines in the array provide excellent sensitivity to diffuse, low surface brightness emission. Future work will involve performing such a study using MeerKAT data in the COSMOS field.

The results in this paper show that compact radio galaxies, or those with only very weak extended emission, are the dominant population in the 10C survey (making up 67/95 sources in this sample), rather than FRI sources dominated by extended emission as predicted by S^3 . Sources with compact morphologies (FR0s) have been found to be the dominant population in lower powered

($L_{1.4 \text{ GHz}}/\text{WHz}^{-1} < 10^{25}$) radio-loud AGN samples in the nearby universe ($z < 0.1$). For example, both ‘FROCAT’ selected from FIRST and NVSS at $z < 0.05$ (Baldi et al. 2018) and the Sadler et al. (2014) study of AT20G-6dFGS sources selected at 20 GHz with $z < 0.1$ are dominated by compact radio galaxies. As the median redshift for our sample is 0.9, our results show that this is also the case at higher redshifts. Recent high-resolution VLA observations of 18 nearby FR0 sources by Baldi et al. (2019) found that 14 were still unresolved on scales of ~ 0.3 arcsec, constraining their linear sizes to < 1 kpc. Higher resolution observations are required to determine whether our higher redshift sources are compact on these scales.

The compact radio galaxies in our sample are a mixed population; the Sadler et al. (2014) study produced similar results, finding that the compact radio galaxies in their sample display a range of spectral indices and are a mixture of different types of objects. Additionally, recent Very Long Baseline Interferometry observations of 14 nearby FR0 sources by Cheng & An (2018) found that their sample is a heterogeneous population consisting of a mixture of CSS and GPS sources, displaying a range of spectral indices and other source properties.

We have previously compared the properties of 10C sources to those in the AT20G-6dFGS sample (Paper III), and suggested that the compact radio galaxies in our sample may be higher redshift analogues of the AT20G-6dFGS sources. Both our sample and the AT20G-6dFGS sample contain a mixture of HERGs and LERGs, while the sources in the Baldi et al. (2018) sample are almost exclusively hosted by LERGs. Our sample therefore allows us to investigate the properties of compact HERGs, which have not been as well studied as the compact LERGs prevalent at lower redshifts. In the local universe, most HERGs are FRII sources, so these compact HERGs at higher redshifts could be young sources that will later develop into FRII sources. Some support for this comes from the fact that in the current sample 34 per cent (11/32) of the HERGs are candidate CSS or GPS sources (FR0c or FR0g), providing evidence that these may be young sources, compared to 14 per cent (6/42) of the LERGs.

The properties of the sources in our sample therefore suggest that these compact, high-frequency selected radio galaxies are a mixed population containing young radio galaxies, older sources that have failed to produce large jets, and, potentially, sources with extended emission with a surface brightness that is too low to be detected in most observations.

7 CONCLUSIONS

We have presented new 15-GHz VLA observations with arcsec resolution of a complete sample of 95 sources selected from the 10C survey at 15.7 GHz. With these higher resolution observations, we were able to identify multiwavelength counterparts for 8 additional sources, meaning that we now have multiwavelength counterparts for 89 out of the 95 sources in the sample. The main results of this work are as follows.

(i) 73 out of 95 (77 per cent) of this complete sample of 10C sources appear unresolved in the VLA observations. However, 6 of these unresolved sources show some extended emission at lower frequencies and are therefore not genuinely compact, meaning 67 out of the 95 sources (71 per cent) are compact on arcsec scales.

(ii) The 10C survey is therefore dominated by compact radio galaxies, or radio galaxies with only very weak extended emission. Sadler et al. (2014) found that compact radio galaxies are the dominant population in high-frequency selected samples in the local universe, so this result shows that this is also the case at $z \sim 1$.

(iii) This provides support for our proposed modification to the S^3 (Whittam et al. 2017), where we showed that if the flat-spectrum cores of faint ($L < 10^{25} \text{ W Hz}^{-1}$) radio galaxies are assumed to be more dominant than previously thought, the high-frequency (18 GHz) source counts and spectral index distributions can be reproduced. However, higher resolution imaging is required to separate the core emission from any small-scale jet emission and fully constrain the fraction of the emission originating from the core.

(iv) The compact radio galaxies in the 10C survey are a mixture of HERGs (40 per cent of compact sources) and LERGs (34 per cent) and display a wide range of spectral indices ($-0.6 < \alpha_{0.61}^{15.7} < 1.2$) and radio powers ($10^{21} < L_{1.4 \text{ GHz}}/\text{W Hz}^{-1} < 10^{29}$), indicating that they are a mixed population. These sources may be the higher redshift ($z \sim 1$) analogues of the compact FR0 sources found in the local universe ($z \sim 0.1$) studied by Sadler et al. (2014), Cheng & An (2018), and Baldi et al. (2018).

(v) In contrast, the extended 10C sources generally have steep spectra, $L_{1.4 \text{ GHz}} > 10^{24} \text{ W Hz}^{-1}$, and are LERGs.

Most studies of compact radio galaxies have been at low redshift and are dominated by LERGs. This sample, on the other hand, contains a significant number of compact HERGs and further studies of these objects have the potential to provide insights into the origin and evolution of HERGs.

ACKNOWLEDGEMENTS

The authors thank the anonymous referee for their helpful comments. IHW acknowledges the financial assistance of the South African Radio Astronomy Observatory (SARAO) towards this research (<http://www.ska.ac.za>). IHW thanks the South African Astronomical Observatory, where some of this work was carried out. The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc. This research has made use of ASTROPY, a community-developed core PYTHON package for astronomy (Astropy Collaboration 2013); APLPY, an open-source plotting package for PYTHON (Robitaille & Bressert 2012); and NASA's Astrophysics Data System.

REFERENCES

AMI Consortium: Davies et al., 2011, *MNRAS*, 415, 2708
 AMI Consortium: Franzen et al., 2011, *MNRAS*, 415, 2699
 An T., Baan W. A., 2012, *ApJ*, 760, 77
 Arnouts S., Ilbert O., 2011, *Astrophysics Source Code Library*, record ascl:1108.009
 Arnouts S., Cristiani S., Moscardini L., Matarrese S., Lucchin F., Fontana A., Giallongo E., 1999, *MNRAS*, 310, 540
 Astropy Collaboration, 2013, *A&A*, 558, A33
 Baldi R. D., Capetti A., 2009, *A&A*, 508, 603
 Baldi R. D., Capetti A., Giovannini G., 2015, *A&A*, 576, A38
 Baldi R. D., Capetti A., Massaro F., 2018, *A&A*, 609, A1
 Baldi R. D., Capetti A., Giovannini G., 2019, *MNRAS*, 482, 2294
 Best P. N., Heckman T. M., 2012, *MNRAS*, 421, 1569
 Biggs A. D., Ivison R. J., 2006, *MNRAS*, 371, 963
 Bolton R. C., Chandler C. J., Cotter G., Pearson T. J., Pooley G. G., Readhead A. C. S., Riley J. M., Waldram E. M., 2006, *MNRAS*, 370, 1556
 Bonaldi A., Bonato M., Galluzzi V., Harrison I., Massardi M., Kay S., De Zotti G., Brown M. L., 2019, *MNRAS*, 482, 2
 Broderick J. W., Fender R. P., 2011, *MNRAS*, 417, 184
 Brunner H., Cappelluti N., Hasinger G., Barcons X., Fabian A. C., Mainieri V., Szokoly G., 2008, *A&A*, 479, 283
 Cheng X.-P., An T., 2018, *ApJ*, 863, 155

Condon J. J., Cotton W. D., Greisen E. W., Yin Q. F., Perley R. A., Taylor G. B., Broderick J. J., 1998, *AJ*, 115, 1693
 Conway J. E., 2002, *New Astron. Rev.*, 46, 263
 Czerny B., Siemiginowska A., Janiuk A., Nikiel-Wroczyński B., Stawarz Ł., 2009, *ApJ*, 698, 840
 de Zotti G., Ricci R., Mesa D., Silva L., Mazzotta P., Toffolatti L., González-Nuevo J., 2005, *A&A*, 431, 893
 Fanaroff B. L., Riley J. M., 1974, *MNRAS*, 167, 31P
 Fotopoulou S. et al., 2012, *ApJS*, 198, 1
 Garn T., Green D. A., Riley J. M., Alexander P., 2008, *MNRAS*, 387, 1037
 Garn T. S., Green D. A., Riley J. M., Alexander P., 2010, *Bull. Astron. Soc. India*, 38, 103
 Ghisellini G., 2011, in Aharonian F. A., Hofmann W., Rieger, F. M., eds, *AIP Conf. Proc. Vol. 180, 25th Texas Symposium on Relativistic Astrophysics*. Am. Inst. Phys., New York, p. 1381
 González-Solares E. A. et al., 2011, *MNRAS*, 416, 927
 Guglielmino G., Prandoni I., Morganti R., Heald G., 2012, *Proc. Sci., Resolving the Sky – Radio Interferometry: Past, Present and Future*. SISSA, Trieste, PoS#22
 Ilbert O. et al., 2006, *A&A*, 457, 841
 Lacy M. et al., 2004, *ApJS*, 154, 166
 Lawrence A. et al., 2007, *MNRAS*, 379, 1599
 Lonsdale C. J. et al., 2003, *PASP*, 115, 897
 Mahony E. K. et al., 2011, *MNRAS*, 417, 2651
 Mauduit J.-C. et al., 2012, *PASP*, 124, 714
 Mingo B., Hardcastle M. J., Croston J. H., Dicken D., Evans D. A., Morganti R., Tadhunter C., 2014, *MNRAS*, 440, 269
 Miraghaei H., Best P. N., 2017, *MNRAS*, 466, 4346
 Owen F. N., Morrison G. E., 2008, *AJ*, 136, 1889
 Prandoni I., Guglielmino G., Morganti R., Vaccari M., Maini A., Röttgering H. J. A., Jarvis M. J., Garrett M. A., 2018, *MNRAS*, 481, 4548
 Reynolds C. S., Begelman M. C., 1997, *ApJ*, 487, L135
 Robitaille T., Bressert E., 2012, *Astrophysics Source Code Library*, record ascl:1208.017
 Rowan-Robinson M., Gonzalez-Solares E., Vaccari M., Marchetti L., 2013, *MNRAS*, 428, 1958
 Sadler E. M., 2016, *Astron. Nachr.*, 337, 105
 Sadler E. M., Ekers R. D., Mahony E. K., Mauch T., Murphy T., 2014, *MNRAS*, 438, 796
 Shabala S. S., Deller A., Kaviraj S., Middelberg E., Turner R. J., Ting Y. S., Allison J. R., Davis T. A., 2017, *MNRAS*, 464, 4706
 Slee O. B., Sadler E. M., Reynolds J. E., Ekers R. D., 1994, *MNRAS*, 269, 928
 Torresi E., Grandi P., Capetti A., Baldi R. D., Giovannini G., 2018, *MNRAS*, 476, 5535
 Tucci M., Toffolatti L., de Zotti G., Martínez-González E., 2011, *A&A*, 533, A57
 Vaccari M., 2015, *Proc. Sci., The Many Facets of Extragalactic Radio Surveys: Towards New Scientific Challenges*. SISSA, Trieste, PoS#27
 Waldram E. M., Pooley G. G., Grainge K. J. B., Jones M. E., Saunders R. D. E., Scott P. F., Taylor A. C., 2003, *MNRAS*, 342, 915
 Waldram E. M., Pooley G. G., Davies M. L., Grainge K. J. B., Scott P. F., 2010, *MNRAS*, 404, 1005
 White R. L., Becker R. H., Helfand D. J., Gregg M. D., 1997, *ApJ*, 475, 479
 Whittam I. H. et al., 2013, *MNRAS*, 429, 2080(Paper I)
 Whittam I. H., Riley J. M., Green D. A., Jarvis M. J., Vaccari M., 2015, *MNRAS*, 453, 4244(Paper II)
 Whittam I. H., Riley J. M., Green D. A., Davies M. L., Franzen T. M. O., Rumsey C., Schammel M. P., Waldram E. M., 2016a, *MNRAS*, 457, 1496
 Whittam I. H., Riley J. M., Green D. A., Jarvis M. J., 2016b, *MNRAS*, 462, 2122(Paper III)
 Whittam I. H., Jarvis M. J., Green D. A., Heywood I., Riley J. M., 2017, *MNRAS*, 471, 908
 Wilkes B. J. et al., 2009, *ApJS*, 185, 433
 Wilman R. J. et al., 2008, *MNRAS*, 388, 1335
 Wilman R. J., Jarvis M. J., Mauch T., Rawlings S., Hickey S., 2010, *MNRAS*, 405, 447
 Zwart J. T. L. et al., 2008, *MNRAS*, 391, 1545

SUPPORTING INFORMATION

Supplementary data are available at [MNRAS](https://www.mnras.org/online) online.

Table 1. Properties of the sources in this sample based on the new VLA observations.

Table B1. The columns of the table summarizing the properties of the host galaxies for the sources in this sample.

Please note: Oxford University Press is not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.

APPENDIX A: DETAILS OF UPDATED OPTICAL COUNTERPARTS

In Section 4, we reviewed, with the addition of the higher resolution VLA data, the multiwavelength counterparts originally assigned to




Figure A1. Two sources previously classified as ‘confused’. The grey scale is the *r*-band optical Sloan Digital Sky Survey image, the blue contours are 1.4-GHz FIRST data, and the green contours are the new 15-GHz VLA data. In both cases, the contours start at 2σ , where σ is the local rms noise in the relevant radio image. The red circles indicate the positions of sources in the multiwavelength Data Fusion catalogue. Note that this catalogue is selected in the mid-infrared, which is why objects are not always visible in the optical grey-scale image shown here. The images are centred on the position of the source in the original 10C catalogue. The images are 0.3×0.3 arcmin².




Figure A2. Source 10C J105009+570724, which was previously classified as confused. We now think this source is two separate sources, which match to the objects labelled 5081 and 5091. Contours and labels are the same as in Fig. A1.




Figure A3. Source 10C J105327+574546. The background grey scale is the new 15-GHz VLA data and the contours are 610-MHz GMRT data from Garn et al. (2008). The red circles indicate the positions of sources in the multiwavelength Data Fusion catalogue (Vaccari 2015). The original match was the object marked with a blue square. With the addition of the new VLA data, we now believe that the match is actually the object marked by the cyan triangle, as this appears to be the core of the source, and the more diffuse emission to the left (which is brighter at 1.4 GHz, indicating that it has a steep spectrum) is a lobe.

these sources in Paper II. Some notes on individual sources are presented in this appendix.

There are eight sources that were classified as ‘confused’ in Paper II; we are now able to identify an optical counterpart for all eight sources. Two examples are shown in Fig. A1.

Source 10C J105009+570724, which was previously classified as confused, is shown in Fig. A2. This source is resolved into two separate components in the VLA image, both of which have a match in the Data Fusion catalogue. We therefore believe that this source is actually two separate objects, and have split it into two entries in the final catalogue (10C J105009+570724A and 10C J105009+570724B).

Source 10C J105327+574546 is shown in Fig. A3. We now believe that the optical counterpart originally assigned to this source is incorrect. Based on the new information from the VLA observations, the correct optical counterpart is actually the object marked with the orange star, as this appears to be coincident with the core of the source. The more diffuse emission to the left is brighter at 1.4 GHz, indicating that it has a steep spectrum, which is consistent with this being one lobe of the radio source.

APPENDIX B: HOST GALAXY CATALOGUE

A table summarizing the multiwavelength data, redshift estimates, and HERG/LERG classifications for the sources in the sample discussed in this paper is available online. This is an update of the information provided in table A1 in Paper II and tables 2 and 5 in Paper III in light of the revised multiwavelength counterparts discussed in Section 4 and Appendix A. Table B1 provides a summary of the columns in the catalogue; the full catalogue is available online.

Table B1. The columns of the table summarizing the properties of the host galaxies for the sources in this sample, which is available online as supplementary material. This is an update of the information provided in table A1 in Paper II and tables 2 and 5 in Paper III in light of the revised multiwavelength counterparts discussed in Section 4 and Appendix A.

Column heading	Description (if applicable)
10C ID	ID from the 10C catalogue
<i>g</i>	<i>g</i> -band magnitude
<i>i</i>	<i>i</i> -band magnitude
<i>r</i>	<i>r</i> -band magnitude
<i>z</i>	<i>z</i> -band magnitude
<i>J</i>	<i>J</i> -band magnitude
<i>K</i>	<i>K</i> -band magnitude
SERVS1 (μ Jy)	
SERVS2 (μ Jy)	
SWIRE1 (μ Jy)	
SWIRE2 (μ Jy)	
SWIRE3 (μ Jy)	
SWIRE4 (μ Jy)	
Best <i>z</i>	Final redshift value (see section 4.3 in Paper II). Null if no redshift value is available.
<i>z</i> flag	Origin of final redshift value: 1 = spectroscopic; 2 = Rowan-Robinson et al. (2013); 3 = Fotopoulou et al. (2012); 4 = LE PHARE. Null if no redshift value is available.
Optical compact	E = source is classified as extended; P = source is classified as point-like (see section 3.1 of Paper III). Null if information is not available.
Lacy et al. AGN area	Y = source is located inside the Lacy et al. (2004) area on the mid-infrared colour–colour diagram; N = source is located outside this region (including sources with limits that must lie outside this region); see section 3.2 in Paper III. Null if information is not available.
X-ray detection	Y = source is detected by an X-ray survey; N = source lies inside X-ray survey area but is not detected (see section 3.3 in Paper III). Null if source is outside the X-ray survey area.
Spectral class	H = source is classified as an HERG based on its optical spectrum; L = source is classified as an LERG based on its optical spectrum (see section 3.5 in Paper III). Null if no spectrum is available.
HERG class	Overall HERG or LERG classification. Null if there is insufficient information available to classify the source.

This paper has been typeset from a \LaTeX file prepared by the author.