



Cite this article: Crellen T, Haswell M, Sithithaworn P, Sayasone S, Odermatt P, Lamberton PHL, Spencer SEF, Déirdre Hollingsworth T. 2023 Diagnosis of helminths depends on worm fecundity and the distribution of parasites within hosts. *Proc. R. Soc. B* **290**: 20222204. <https://doi.org/10.1098/rspb.2022.2204>

Received: 3 November 2022

Accepted: 16 December 2022

Subject Category:

Ecology

Subject Areas:

health and disease and epidemiology, ecology

Keywords:

neglected tropical diseases, foodborne trematodiasis, epidemiology, macroparasite, Bayesian inference

Author for correspondence:

Thomas Crellen

e-mail: thomas.crellen@glasgow.ac.uk

Diagnosis of helminths depends on worm fecundity and the distribution of parasites within hosts

Thomas Crellen^{1,2,3}, Melissa Haswell^{4,5}, Paiboon Sithithaworn⁶, Somphou Sayasone⁷, Peter Odermatt^{8,9}, Poppy H. L. Lamberton^{1,2}, Simon E. F. Spencer¹⁰ and T. Déirdre Hollingsworth³

¹School of Biodiversity One Health and Veterinary Medicine, Graham Kerr Building, University of Glasgow, 82 Hillhead Street, Glasgow G12 8QQ, UK

²Wellcome Centre for Integrative Parasitology, Sir Graeme Davies Building, University of Glasgow, 120 University Place, Glasgow G12 8TA, UK

³Big Data Institute, Li Ka Shing Centre for Health Information and Discovery, University of Oxford, Oxford OX3 7LF, UK

⁴Office of the Deputy Vice Chancellor, Indigenous Strategy and Services and School of Geosciences, John Woolley Building, University of Sydney, Sydney, New South Wales 2050, Australia

⁵School of Public Health and Social Work, Kelvin Grove Campus, Queensland University of Technology, Brisbane City, Queensland 4000, Australia

⁶Department of Parasitology, Khon Kaen University, 123 Thanon Mittraphap, Khon Kaen 40002, Thailand

⁷Lao Tropical and Public Health Institute, Samsenthai Road, Sisattanak district, Vientiane, Lao PDR

⁸Department of Public Health and Epidemiology, Swiss Tropical and Public Health Institute, Kreuzstrasse 2, Allschwil 4123, Switzerland

⁹University of Basel, Petersplatz 1, Basel 4001, Switzerland

¹⁰Department of Statistics, University of Warwick, Coventry CV4 7AL, UK

ID TC, 0000-0003-2934-1063; MH, 0000-0002-4067-7951; PS, 0000-0002-9480-1778; SS, 0000-0001-8219-9567; PO, 0000-0002-0296-2508; PHLL, 0000-0003-1048-6318; SEFS, 0000-0002-8375-5542; TDH, 0000-0001-5962-4238

Helminth transmission and morbidity are dependent on the number of mature parasites within a host; however, observing adult worms is impossible for many natural infections. An outstanding challenge is therefore relating routine diagnostics, such as faecal egg counts, to the underlying worm burden. This relationship is complicated by density-dependent fecundity (egg output per worm reduces due to crowding at high burdens) and the skewed distribution of parasites (majority of helminths aggregated in a small fraction of hosts). We address these questions for the carcinogenic liver fluke *Opisthorchis viverrini*, which infects approximately 10 million people across Southeast Asia, by analysing five epidemiological surveys ($n = 641$) where adult flukes were recovered. Using a mechanistic model, we show that parasite fecundity varies between populations, with surveys from Thailand and Laos demonstrating distinct patterns of egg output and density-dependence. As the probability of observing faecal eggs increases with the number of mature parasites within a host, we quantify diagnostic sensitivity as a function of the worm burden and find that greater than 50% of cases are misdiagnosed as false negative in communities close to elimination. Finally, we demonstrate that the relationship between observed prevalence from routine diagnostics and true prevalence is non-linear and strongly influenced by parasite aggregation.

1. Introduction

Parasitic helminths continue to cause diseases of major medical and veterinary significance. The frequency distribution of mature parasites is the key quantity of interest in the epidemiology of helminths as higher worm burdens are linked

with increased host infectiousness, more severe morbidity, and are a marker of past exposure [1–3]. The transmission dynamics of helminths, also classified as macroparasites, can be described through changes to the mean worm burden in the population over time. This is in contrast to many protozoal, bacterial and viral infections (collectively known as microparasites), where the trajectory of epidemics can be characterized by the number of hosts in certain disease states; for instance susceptible, infected or recovered [2,4]. In populations of humans [2] and wildlife [5], helminths are typically aggregated in a small fraction of hosts. This makes control using mass deworming more challenging compared to randomly distributed parasites [6], however if heavily infected hosts can be identified it presents an opportunity to drive down transmission rates. Worm burden is also a critical driver of chronic morbidity. While the mechanisms for helminth-induced pathology are species-specific, in many cases, chronic damage is attributable to the intensity and duration of past infection with adult worms [7].

Despite the importance of characterizing the distribution of parasites within a population, it is challenging to directly observe mature worms from natural hosts for most species of helminths, with the roundworm *Ascaris lumbricoides* as a notable exception [8–10]. Diagnosis typically relies on indirect coprological or serological measures of the worm burden, such as faecal egg counts, worm antigen concentrations or molecular detection of parasite DNA in blood or stool. Historical autopsies provide critical insights into the relationship between worm burdens and indirect diagnostics for several helminth species [11–13]; however, these studies may not be reflective of current parasite fecundity in endemic settings due to host population heterogeneity. For instance, the fecundity of *A. lumbricoides* differs markedly between hosts in Bangladesh and Nigeria [14]. The relationship may also alter over time as the nutritional status of populations changes and control programmes reduce the intensity of parasite infections through repeated mass treatment [15,16]. In the absence of data on worm burdens, statistical frameworks have been proposed to infer the number of adult worms at the population level from indirect diagnostics, usually counts of parasite eggs in stools [17–19]. However, accurate inference of the worm burden is dependent on substantial prior knowledge of parasite ecology including (i) the relationship between worm burden and egg output in the population of interest; (ii) a probability distribution which captures the variability in egg output from a given number of worms, and (iii) the sensitivity and specificity of diagnostics.

The liver fluke *Opisthorchis viverrini* infects around 10 million people across Southeast Asia, predominantly in Laos, Cambodia and the North and Northeast regions of Thailand [20]. The parasite has a complex life cycle which includes both *Bithynia* snails and freshwater fish as intermediate hosts. Humans become infected by consuming raw or undercooked fish encysted with metacercariae. After ingestion, adult worms migrate to the bile duct where they cause chronic damage to host tissue through a combination of mechanical damage, inflammation and the secretion of proteins; most notably the cell growth factor granulin [21,22]. The pathology progresses to cholangiocarcinoma (bile duct cancer) in around 1–5% of infected people, with higher worm burdens increasing the risk of developing cancer [23,24]. Humans do not develop immunity to

O. viverrini and repeated reinfection is common [25]. Nevertheless, regular anthelmintic treatment to kill adult worms is likely to slow the progression of liver pathology and reduce the risk of developing cholangiocarcinoma [26]. The resulting sequela is considered a neglected tropical disease (NTD), under the grouping of foodborne trematodiasis, and is targeted for enhanced control globally by 2030 [27]. Endemic countries have initiated control programmes, with Thailand the first country to start an opisthorchiasis control programme in 1950, aiming to halt parasite transmission and reduce cases of the resulting cholangiocarcinoma [28]. Central to achieving these objectives is a quantitative understanding of parasite transmission to inform control strategies [29,30].

The overdispersed pattern of helminth parasites in a population is best captured with a negative binomial distribution, also known as a gamma–Poisson mixture, parameterized by the mean worm burden M and a dispersion factor k that scales inversely with the variance [2]. While previous studies have reported an aggregated distribution of *O. viverrini* worms within human hosts [31–33], the negative binomial distribution has never been fitted to data on *O. viverrini* and therefore there are no likelihood-based estimates of the aggregation parameter k [30]. The relationship between *O. viverrini* adult worm burdens and faecal egg counts has previously been reported from either (i) autopsies in which the liver is dissected, or (ii) surveys where worms are expelled by administration of a saline purgative following anthelmintic treatment with praziquantel (table 1 and figure 1a). Typically, a linear regression between log-transformed worm burden (x) and total egg counts per host (y) is used; $\log_{10}(y+1) = \beta \cdot \log_{10}(x+1)$, or $y = (x+1)^{\beta} - 1$ when untransformed. Previous values of β are 1.67 [34] and 2.0 [31] which indicate inverse density-dependence. As this is biologically implausible and contrary to the reported findings in these studies, the log-log linear model is likely inappropriate. Alternative parametric functions have been proposed to model the relationship between total worm burden and egg counts, including power law [40] and algebraic decay functions [41].

The sensitivity of parasitological diagnostics is known to covary with the number of mature worms within hosts (or pairs of mated adult worms in the case of dioecious helminths) as the probability of observing at least one egg in stool or urine increases with higher worm burdens [42]. Diagnostic sensitivity is rarely considered as a function of worm burden, however, this provides considerable insights into the relationship between the true and observed prevalence and how the reliability of diagnostics evolves as control programmes reduce infection intensity [43,44].

Here, we infer the *O. viverrini* worm burden at an individual and community level (figure 1b) using a mechanistic statistical model. We quantify the relationship between worm burden and egg output in separate populations (figure 1c); provide the first fitting of the negative binomial distribution to *O. viverrini* worms at the population level using partial pooling across surveys to estimate the aggregation parameter k (figure 2a); estimate the sensitivity of faecal egg diagnostics as a function of the worm burden (figure 2c,d); and show the relationship between the true and observed prevalence (figure 2e,f). Our study provides a foundation for accurately characterizing the infection intensity of helminths in endemic communities




Figure 1. Observed data on *Opisthorchis viverrini*, worm burden inference and parasite fecundity. (a) Relationship between observed *O. viverrini* adult worms (w) and eggs per gram of stool (y) in five surveys (table 1) shown on the log scale. (b) Relationship between inferred worm burdens (x) and faecal egg counts. Eggs per gram for TH1 and TH2 have been adjusted by the model. (c) As (b) with points coloured by country of survey. The dashed lines are a power law function with parameter values taken from mean posterior estimates. The shaded area indicates the 95% credible interval of the mean. (d) As (c) with the shaded area representing the 90% prediction interval based on a zero-truncated negative binomial error distribution. Validation data are shown as square points (table 1).

Table 1. Cross-sectional surveys investigating the relationship between adult *Opisthorchis viverrini* worm burden and faecal egg counts in humans included in the analysis. Surveys TH1–4 and LAO1 had individual-level data used for model fitting. Surveys TH5 and LAO2–3 had aggregated data used for model validation. The table shows the method of adult worm collection, either from liver examination during autopsy or expulsion with a saline purgative after anthelmintic treatment with praziquantel; the study location and date; the number of human participants (sample size); the arithmetic mean number of *O. viverrini* adult worms recovered; and the diagnostic method used for parasite egg identification and counting. TH relates to surveys in Thailand and LAO to Laos.

survey and method	location and year(s)	sample size	worms recovered			faecal egg diagnostic
			mean	range	s.d. ^a	
TH1 autopsy [34]	Khon Kaen 1982–1999	139	160	0–2954	390	FECT ^b
TH2 expulsion [35]	Khon Kaen 1987	33	85	0–565	154	Stoll's dilution
TH3 expulsion [31]	Kalasin 1989	231	39	0–832	105	FECT ^b
TH4 expulsion [36]	Khon Kaen 1991	141	49	0–874	102	FECT ^b
LAO1 expulsion [37]	Savannakhet 2005	97	183	0–2178	286	FECT ^b
TH5 autopsy ^c [38]	Khon Kaen 1966	9	2588	20–11856	NA ^d	Stoll's dilution
LAO2 expulsion ^c [39]	Savannakhet 2008	125	11	0–111	29	Kato–Katz
LAO3 expulsion ^c [39]	Savannakhet 2011	82	3	0–66	9	Kato–Katz

^aStandard deviation.

^bFormalin–ether concentration technique.

^cSurvey used for validation only.

^dData not available.




Figure 2. Distribution of adult *O. viverrini* worms and diagnostic sensitivity. (a) Negative binomial dispersion parameter k by survey (table 1) where points show the posterior mean and lines the 95% credible interval, 90% prediction interval, and validation datasets shown as squares. (b) Relationship between estimates of k and the estimated mean worm burden (M) by survey. (c) The probability of detecting ≥ 1 faecal egg by parasitological diagnostic (formalin–ether concentration technique; FECT) as a function of an individual's worm burden (x), the shaded area gives the 95% credible interval. (d) Sensitivity of parasitological diagnostics (FECT) at the population level for different frequency distributions of adult worms, given by the aggregation parameter k . (e) Relationship between the observed prevalence by parasitological diagnostic (FECT) and the true prevalence for different values of the aggregation parameter k (see legend in (d)). Long dashed line gives equality between axes. Short dashed lines indicate the true prevalence when the observed prevalence is 20%. (f) As (e) with a varying specificity of parasitological diagnostic, where worm aggregation $k = 0.36$ for all curves.

and estimating epidemiological parameters to facilitate evidence-based control.

2. Material and methods

(a) Epidemiological framework

We developed and applied a mechanistic statistical model to individual-level data on adult worm burdens and egg counts which incorporates key aspects of parasite ecology. Our observed data are the counts of recovered *O. viverrini* adult

worms from each individual i in survey j , denoted w_{ij} , and the eggs per gram of stool, denoted y_{ij} . We start from the assumption that the true number of *O. viverrini* adult worms per individual, x_{ij} , follow the negative binomial distribution within community j , which is parameterized by a mean worm burden M_j and dispersion factor k_j [2,5]

$$\Pr(X_{ij} = x | k_j, M_j) = \binom{x + k_j - 1}{x} \left(\frac{k_j}{M_j + k_j} \right)^{k_j} \left(\frac{M_j}{M_j + k_j} \right)^x. \quad (2.1)$$

The value of k_j from each study is themselves normally distributed with an overall mean of μ_k and a standard deviation of σ_k , meaning that estimates of k_j are informed by partial pooling. In worm expulsion studies, participants are treated with the anthelmintic praziquantel followed by a saline purgative, either magnesium sulfate solution or sodium sulfate solution, several hours later. This is known to result in imperfect recovery as worms can be degraded and unrecoverable following anthelmintic treatment; remain blocked in the biliary tracked; or stool sample collections may be missed [45,46]. Therefore, we allow the true worm burden to be greater or equal to the observed number of worms for individuals in expulsion studies ($x_{ij} \geq w_{ij}$) and the probability of observing w_{ij} worms given a true count of x_{ij} is a binomial sampling process with a probability of worm recovery r that is constant across surveys. During autopsy, adult *O. viverrini* were carefully removed from cross sections of liver and worm recovery is likely close to 100% [34]. Therefore, we consider that the true worm count for each individual is equal to the recovered worms in autopsy studies ($x_{ij} = w_{ij}$, $r = 1$).

The expected *O. viverrini* eggs per gram of stool for each person, π_{ij} , is given by the generalized function $\Lambda(x, \theta)$, where θ is the set of parameters specific to that function. Here, we show results from the power law function as the impact of density-dependent regulation on helminth fecundity is directly quantified by the γ parameter; $\pi_{ij} = (\lambda_{c(j)} x_{ij})^{\gamma_{c(j)}}$, where $c(j)$ denotes the country of survey j . As *O. viverrini* is hermaphroditic [47], we assume that a single worm is capable of producing eggs. Observed egg counts for an individual, y_{ij} , are a realization of an error distribution with mean π_{ij} . Our model uses a negative binomial hurdle distribution with dispersion parameter $h_{c(j)}$ and a separate function for false-negative egg counts, which gives the probability of observing zero eggs as a function of the worm burden within an individual

$$\Pr(Y_{ij} = y | X_{ij} = x, \pi_{ij}, h_{c(j)}, b) = \begin{cases} 1 & y = 0, x = 0 \\ 0 & y \geq 1, x = 0 \\ 1 - \frac{x}{x+b} & y = 0, x \geq 1 \\ \frac{x}{x+b} \binom{y+h-1}{y} \frac{(h/(\pi+h))^h (\pi/(\pi+h))^y}{1 - (h/(\pi+h))^h} & y \geq 1, x \geq 1. \end{cases} \quad (2.2)$$

Bringing these probabilities together, we obtain an expression for the likelihood

$$\begin{aligned} \Pr(Y_{ij} = y, W_{ij} = w | \pi_{ij}, h_{c(j)}, b, r, M_j, k_j) \\ = \sum_{x=w}^{\infty} \Pr(Y_{ij} = y | X_{ij} = x, \pi_{ij}, h_{c(j)}, b) \cdot \\ \Pr(W_{ij} = w | x, r) \cdot \Pr(X_{ij} = x | M_j, k_j). \end{aligned} \quad (2.3)$$

The diagnostic sensitivity S varies by survey j , and is dependent on individual sensitivity, or the probability of observing at least one faecal egg as a function of an individual's worm burden $se(x) = x/(x+b)$, and the frequency distribution of worms in the community

$$S(M, k) = \sum_{x=1}^{\infty} se(x) \binom{x+k-1}{x} \frac{(k/(M+k))^k (M/(M+k))^x}{p(M, k)}, \quad (2.4)$$

where $p(M, k)$ indicates the true prevalence and is given by

$$p(M, k) = 1 - \left(\frac{k}{M+k} \right)^k. \quad (2.5)$$

The observed prevalence p' for survey j is therefore

$$p' = p(M, k) \cdot S(M, k) + (1 - p(M, k)) \cdot (1 - sp), \quad (2.6)$$

where sp gives the diagnostic specificity.

(b) Data sources

We accessed individual-level data from one autopsy ($n = 139$) and four worm expulsion surveys ($n = 502$), which are summarized in table 1. We contacted the study authors to access the original data. Where these were unavailable, we extracted data from publication figures using webplotdigitizer (v. 4.5) and obtained additional information, e.g. the number of (0, 0) points, from text and tables. We confirmed that extracted data showed near identical values to the original reports. Two studies used different methodologies to transform observed egg counts into the reported eggs per gram of stool (EPG). In survey TH1 egg counts were multiplied by an additional factor between 1 and 4 depending on stool consistency [34]. As this was not performed in other surveys, we introduced an integer factor for each individual in TH1, ξ_i , which can take on values $\in \{1, 2, 3, 4\}$ and the expected egg count is modified by this factor; $\pi_i = \Lambda(x, \theta) \xi_i$. Another survey (TH2) used an older technique, Stoll's dilution, to calculate EPG. We, therefore, modified the expected egg counts from TH2 with a real factor; $\pi = \Lambda(x, \theta) \rho$, to ensure that our estimates of diagnostic parameters are consistent for FECT. We obtained aggregated data from a further three surveys where *O. viverrini* worms were recovered, which were used for model validation (TH5, LAO2 and LAO3; see table 1). For the expulsion surveys (LAO2 and LAO3), we corrected the mean number of worms recovered by dividing the observed values by the probability of worm recovery (r) estimated in the model. Values of k for LAO2 and LAO3 (figure 2a,b) were approximated using the negative binomial relationship between prevalence and corrected mean worm burden (equation (2.5)). Egg counts in TH5 were reported as perworm per day (3160), which was converted to eggs per gram of stool by multiplying by the mean number of worms, then dividing by the expected daily mass of stool (250 g in developing countries [48]) and the correction factor for Stoll's dilution (ρ).

(c) Model fitting

We fitted statistical models to data using Hamiltonian Monte Carlo implemented in Stan v. 2.30.1 [49,50]. Integer parameters were marginalized out of the likelihood function (equation 2.3) and estimated by computing the normalized likelihood. As the model fitting was performed in a Bayesian framework, parameters were assigned weakly or moderately informative prior distributions based on the available literature. Each model was run with four parallel chains with a burn-in of 2750 iterations per chain and a total of 1000 samples from the posterior distribution. Model fitting diagnostics used to determine successful chain convergence were the Gelman–Rubin diagnostic $\hat{r} \leq 1.01$ and effective sample size greater than 500 [51]. We report the posterior mean and 95% credible interval (CrI) for estimated

parameters. Data were processed and figures produced in R v. 4.2.1 [52].

3. Results

(a) Individual worm burdens

Expulsion of liver flukes following anthelmintic treatment is known to result in imperfect recovery [31,45] and we quantified the proportion of adult flukes recovered as less than half ($r=0.44$ [95% CrI 0.42–0.45]). The true *O. viverrini* worm burden for each individual (x_{ij}) was estimated as greater than the observed count (w_{ij}) for 421/502 people in worm expulsion studies, with a median increase of 10 worms in the inferred burden and a mean increase of 92 worms (figure 1b). Across all positive cases for *O. viverrini*, the inferred adult worm burden ranges from 1 to 4677 with a median of 38 and a mean of 192.

(b) Worm fecundity and variance in faecal egg counts

The expected parasite fecundity, measured as eggs per gram of stool, was quantified using a power law function (see Methods), which directly estimates the number of eggs produced by a single fluke (λ), and the extent of density-dependence (γ), where values $\gamma < 1$ indicate a decline in *per capita* egg output at higher worm burdens. From surveys in Thailand, one adult *O. viverrini* was estimated to produce a mean of 72 eggs per gram of stool (95% CrI 66–78), which is substantially higher than the 20 eggs per gram of stool (95% CrI 13–30) estimated from a single fluke in Laos. Our results suggest that density-dependence regulates egg output at higher worm burdens in Laos ($\gamma=0.82$ [95% CrI 0.74–0.90]) and to a lesser extent in Thailand, where the posterior mean for γ was close to unity ($\gamma=0.98$ [95% CrI 0.96–0.99]), see figure 1c. The functional relationship between worm burden (x) and expected egg output (π) is therefore $\pi=72x^{-0.98}$ in Thailand and $\pi=20x^{-0.82}$ in Laos. Expected eggs per gram of stool for survey TH2, which used an alternative diagnostic method (table 1), were corrected by a factor estimated in the model ($\rho=6.2$). There was substantial variation around the expected values for egg counts, which is consistent with reports from other helminths [53]. We quantified this variance using a zero-truncated negative binomial distribution, also known as a hurdle model (equation (2.2)), characterized by a mean and an aggregation parameter h , which scales inversely with variance. Estimated values of h for Thailand ($h=0.40$ [95% CrI 0.32–0.48]) indicate substantially higher variation in egg output compared with Laos ($h=0.63$ [95% CrI 0.47–0.79]), which is partly due to heterogeneity across the multiple surveys in Thailand. We calculated 90% prediction intervals for *O. viverrini* fecundity (figure 1d), which were validated using additional survey data where faecal egg counts and worm burdens were aggregated (TH5, LAO2 and LAO3; see table 1) and which were not used in our model fitting. We adjusted the reported mean worms expelled in surveys LAO2 and LAO3 by dividing by the recovery parameter $r=0.44$, giving mean worm burdens of 25 and 6, respectively. The validation survey data are shown in figure 1d, and the relationships between worm burdens and faecal egg counts are within our 90% prediction intervals.

(c) Parasite distribution and aggregation

Adult worm burdens of *O. viverrini* have been described as overdispersed within endemic populations. This motivated our fitting of the negative binomial distribution (equation (2.1)), with values of the aggregation parameter k below one indicating a significant departure from randomly distributed worms (Poisson variation). Our posterior mean estimates of the dispersion parameter k_j range from 0.24 to 0.56 by survey (figure 2a). Values of k_j across surveys were assumed to be normally distributed with a global mean (μ_k) of 0.36 (95% CrI 0.12–0.55). Given our estimate of the standard deviation between surveys ($\sigma_k=0.18$), our 90% prediction interval for k is 0.07–0.66. These values are indicative of high levels of parasite aggregation and are consistent with findings that most helminth species have k values between 0.1 and 1 [2,5,8]. Estimates for the mean *O. viverrini* burden by survey (M_j) were all estimated to be higher than the observed mean flukes expelled (shown in table 1) and increased to 176 (TH1), 115 (TH2), 81 (TH3), 91 (TH4) and 245 (LAO1), respectively. The mean *O. viverrini* burden for the autopsy study (TH1) was estimated to be slightly higher than the recovered mean count of worms due to fitting the negative binomial distribution. Examining the relationship between values of k and mean worm burden by survey, we observe that parasite dispersion increases at lower worm burdens (figure 2b), as has also been reported for other helminths [54,55], presenting challenges for helminth elimination as parasites are aggregated in fewer hosts as control programmes progress.

(d) Diagnostic sensitivity

We quantify the probability of observing parasite eggs for an individual as a function of the worm burden using a strictly increasing function; $se(x)=x/(b+x)$. Given our mean estimate for the parameter $b=1.7$ (95% CrI 1.6–1.9), the probability of observing at least one parasite egg when an individual is infected with one *O. viverrini* fluke is 0.37 (95% CrI 0.34–0.39), five flukes is 0.74 (95% CrI 0.72–0.76), 20 flukes is 0.92 (95% CrI 0.92–0.93), and reaches 0.99 at burdens of 115 flukes and higher, see figure 2c.

To calculate the sensitivity of faecal egg diagnostics at the population level, the probability of observing worms per individual is considered along with the distribution of adult worms (equation 2.4). Diagnostic sensitivity covaries with the adult worm burden, however, the relationship is nonlinear and depends on adult worm aggregation described by the negative binomial parameter k (figure 2d). Given the average worm dispersion observed in this study ($k=0.36$), diagnostic sensitivity at a high mean worm burden, $M=100$, is 0.88, reducing M to 20, 5, 1 or 0.1 results in population diagnostic sensitivities of 0.78, 0.66, 0.50 and 0.37, respectively. Therefore, when close to the threshold for elimination the proportion of false-negative cases is greater than or equal to 50%, which presents challenges both for case detection and the cost per detected case in low-intensity settings [56].

(e) Prevalence

The relationship between the true prevalence of infection and the observed prevalence from faecal eggs is nonlinear and depends on the population diagnostic sensitivity; worm burden and aggregation; and assumptions on diagnostic

Table 2. Summary of epidemiological parameters for *Opisthorchis viverrini* estimated in this study. TH relates to parameters estimated from surveys in Thailand and LAO relates to surveys in Laos (table 1).

parameter	symbol	posterior mean	95% credible interval
worm aggregation	μ_k	0.36	0.12–0.55
proportion of worms recovered by expulsion	r	0.44	0.42–0.45
egg output from 1 worm	λ (TH)	72	66–78
	λ (LAO)	20	13–30
density-dependence	γ (TH)	0.98	0.96–0.99
	γ (LAO)	0.82	0.74–0.90
egg dispersion	h (TH)	0.40	0.32–0.48
	h (LAO)	0.63	0.47–0.79

specificity (equation 2.6). At an observed prevalence of 20%, which is the threshold for annual or biannual community deworming of *O. viverrini* [57], the true prevalence can vary from 29 to 42% depending on worm aggregation in the range $k=0.1$ – 0.7 (figure 2e). Relaxing the assumption of perfect specificity for parasitological diagnosis has implications for interpreting observed parasite prevalence, with false-positive results dominating at low worm burdens (figure 2f).

4. Discussion

Control of helminths in humans is increasingly influenced by results from quantitative transmission models, which inform recommendations for national control programmes [29,58,59]. Predictions made by these models, such as the frequency and coverage of mass treatment required to achieve elimination, are strongly influenced by model structure and the assumed parameter values. This study brings together multiple datasets of individual-level parasite worm burdens and egg counts in conjunction with an epidemiological framework which quantifies key aspects of parasite ecology for the liver fluke *Opisthorchis viverrini*. Our estimates of the aggregation of adult worms, worm fecundity and diagnostic sensitivity as a function of worm burden (table 2) are essential for interpreting routinely collected parasitological survey data, understanding parasite transmission dynamics, and estimating the worm burden from indirect diagnostics [17,19]. Our inference framework is broadly applicable across helminth species and can be used to estimate epidemiological parameters for other parasites of medical and veterinary significance.

Macroparasite population dynamics are sensitive to small differences in the aggregation of adult worms, given by the negative binomial parameter k [2,60]. Here, we have quantified k and its variation between surveys using a hierarchical model which pools information between five surveys, making the posterior estimates robust to outlier values. An analysis of *O. viverrini* transmission dynamics in Southern Laos assumed that the negative binomial dispersion of adult worms given by the parameter k was 0.1 [61]. Given our prediction interval for k , we estimate there is a 93% probability that the value of k is higher than 0.1, and our best estimate is 0.36. However, given the strong observed relationship between mean worm burden and k [54,55], a value of 0.1

is plausible in lower-intensity settings (figure 2b). Parasite aggregation is caused by heterogeneity in host exposure and/or susceptibility. For *O. viverrini* aggregated worm burdens are ultimately driven by variance in individual feeding rates and the density of metacercariae in local fish [30]. There is a cultural component to transmission as consumption of raw fish is perceived to be linked with male virility and is a marker for local identity and traditional practices in the Northeast ‘Isan’ region of Thailand [62,63]. The distribution of *O. viverrini* metacercariae in cyprinid fish intermediate hosts is also overdispersed [33], which further contributes to variance in the adult worm distribution within definitive human hosts; however, this has not yet been quantified by fitting the negative binomial distribution.

The relationship between mature worms and indirect diagnostics, such as faecal egg counts, is central to estimating the worm burden and describing transmission potential. Our models support different patterns of helminth fecundity in surveys from Thailand over the period 1966–1992 versus more recent surveys from Laos conducted between 2005 and 2011 (figure 1). These findings bolster the claim that helminth fecundity varies between host populations, as previously reported for *A. lumbricoides* [14]. The mechanisms for these differences are poorly understood and may comprise multiple host and parasite factors, however ecological studies of helminths show an important role for host nutrition [64].

Imperfect diagnostic observations bias estimates of epidemiological parameters [44]. Here, we explicitly quantify these inaccuracies by estimating the proportion of worms recovered by worm expulsion and the probability of detecting faecal eggs as a function of individual worm burden. Our sensitivity estimates at the population level (figure 2d) are most relevant for the formalin–ether concentration technique (FECT) used in the majority of surveys (table 1) and we accounted for the alternative diagnostic used in survey TH2 (Methods). Our results at lower mean worm burdens are consistent with estimates of FECT sensitivity validated with PCR and worm expulsion as the gold standard [65]. Understanding how diagnostic sensitivity changes in response to a declining worm burden has important implications for control programmes as the number of false negatives increases when a programme is close to elimination and so larger sample sizes are required to estimate endemicity. There is a nonlinear relationship between the observed and true prevalence, which is dependent on the population worm burden

and the degree of aggregation (figure 2e and equation (2.6)). At an observed prevalence of 20% for *O. viverrini*, below which the World Health Organization advises mass drug administration to reduce from biannual to annual treatment [57], we estimate that the true prevalence is actually twofold higher at 40% where $k > 0.5$. These findings demonstrate that prevalence of infection is an unreliable indicator on which to base thresholds for helminth control programmes.

There were limitations to our analysis. Recruitment in expulsion surveys was not always random and if individuals with low or zero egg counts are underrepresented in the sample, this may influence our estimation of sensitivity, as there are fewer observations from individuals with low worm burdens, and inflate values of k , as the community worm burden appears more homogeneous; although by using partial pooling we likely reduced the bias in estimating k . An additional challenge is the presence of minute intestinal flukes; helminths with the same transmission route as *O. viverrini* which frequently occur as co-infections [39,65]. These flukes produce visually similar eggs to *O. viverrini* leading to imperfect specificity by parasitological diagnostics. Our ability to account for misdiagnosis in model fitting was limited as most surveys in Thailand did not clarify whether intestinal flukes were present. We explore the impact of imperfect diagnostic specificity on the observed prevalence under simplistic scenarios shown in figure 2f; in reality, we expect specificity for *O. viverrini* to covary with the endemicity of minute intestinal flukes. The increased use of molecular diagnostics enables the species of parasite egg to be identified [66], and recent improvements to *O. viverrini* antigen tests also provides a method to discriminate cases [67,68].

Future studies on macroparasite epidemiology would benefit from a renewed focus on the worm burden to glean insights into transmission dynamics, parasite-induced morbidity, the impact of interventions, and the efficacy of anthelmintic treatment. For species where it is difficult to observe worm burdens directly, population genetics also provides a means to assess the relatedness between offspring and thus estimate the number of mature adult worms [69,70].

Research on the epidemiology of parasitic helminths experienced a period of substantial development between roughly 1978–2000, instigated by the collaboration between Roy Anderson and Robert May who drew together ideas from ecology, parasitology and mathematics into a highly influential series of deterministic models [1–3]. This framework for infectious diseases was further developed by a generation of researchers, many of whom were graduate students or postdocs of Anderson & May [71]. In recent

years, the epidemiology of microparasites has advanced significantly, driven by both theoretical work [4] and improved data (in particular, whole-genome sequencing makes it possible to infer transmission networks for certain pathogens [72,73]); however, research on macroparasites has received less attention [74,75]. Many scientific questions raised in foundational studies, including the extent to which adult worm overdispersion is driven by host or parasite genetic factors, have arguably remained unanswered and the focus on worm burdens has been neglected by many in the field with insights from disease ecology replaced by less appropriate techniques from medical statistics [43,76]. The literature is replete with criticisms of ‘classical’ frequentist statistics and null-hypothesis testing [77–79]. While reiterating these arguments goes beyond the scope of this article, we emphasize that estimating biologically meaningful parameters is a powerful framework for understanding parasite ecology.

In this study, we have quantified the distribution of adult worms in host populations and worm fecundity, and in doing so provide a inference framework to convert readily available diagnostic data into worm burdens to further our understanding of the transmission, evolution and control of helminth parasites.

Ethics. As an analysis of previously published and anonymous human data, this study met the criteria for exemption from ethical review at the Universities of Oxford and Glasgow.

Data accessibility. Data used for model fitting (individual worm and egg counts) are available from Dryad <https://doi.org/10.5061/dryad.q83bk3jn6> [80]. Code to reproduce the analysis is available at the GitHub repository <https://github.com/tc13/worm-inference>.

Authors' contributions. T.C.: conceptualization, data curation, formal analysis, funding acquisition, investigation, methodology, project administration, software, validation, visualization, writing—original draft, writing—review and editing; M.H.: conceptualization, resources; P.S.: conceptualization, resources; S.S.: resources, validation; P.O.: resources, validation; P.H.L.L.: supervision, writing—review and editing; S.E.F.S.: conceptualization, methodology, supervision, writing—review and editing; T.D.H.: conceptualization, methodology, supervision, writing—original draft, writing—review and editing.

All authors gave final approval for publication and agreed to be held accountable for the work performed therein.

Conflict of interest declaration. The authors declare no competing interests.

Funding. This research was funded in whole, or in part, by the Wellcome Trust (grant no. 215919/B/19/Z). For the purpose of open access, the author has applied a CC BY public copyright licence to any Author Accepted Manuscript version arising from this submission.

Acknowledgements. We thank Daniel T. Haydon, Joaquin M. Prada and Anastasia Hernandez Koutoucheva for helpful input and discussions.

References

- May RM, Anderson RM. 1979 Population biology of infectious diseases: part II. *Nature* **280**, 455–461. (doi:10.1038/280455a0)
- Anderson RM, May RM 1991 *Infectious diseases of humans: dynamics and control*. Oxford, UK: Oxford University Press.
- Wilson K, Bjørnstad ON, Dobson AP, Merler S, Poglayen G, Randolph SE, Read AF, Skorpington A. 2002 Heterogeneities in macroparasite infections: patterns and processes. In *The ecology of wildlife diseases* (eds PJ Hudson, A Rizzoli, BT Grenfell, H Heesterbeek, AP Dobson), ch. 2, pp. 6–44. Oxford, UK: Oxford University Press.
- Diekmann O, Heesterbeek H, Britton T 2013 *Mathematical tools for understanding infectious disease dynamics*. Princeton, NJ: Princeton University Press.
- Shaw DJ, Grenfell BT, Dobson AP. 1998 Patterns of macroparasite aggregation in wildlife host populations. *Parasitology* **117**, 597–610. (doi:10.1017/S0031182098003448)
- Irvine MA, Reimer LJ, Njenga SM, Gunawardena S, Kelly-Hope L, Bockarie M, Hollingsworth TD. 2015 Modelling strategies to break transmission of lymphatic filariasis—aggregation, adherence and vector competence greatly alter elimination. *Parasit. Vectors* **8**, 1–19. (doi:10.1186/s13071-015-1152-3)
- Wiegand RE et al. 2021 Associations between infection intensity categories and morbidity prevalence in school-age children are much stronger for *Schistosoma haematobium* than for *S. mansoni*.

- PLoS Negl. Trop. Dis.* **15**, e0009444. (doi:10.1371/journal.pntd.0009444)
8. Guyatt HL, Bundy DAP, Medley GF, Grenfell BT. 1990 The relationship between the frequency distribution of *Ascaris lumbricoides* and the prevalence and intensity of infection in human communities. *Parasitology* **101**, 139–143. (doi:10.1017/S003118200079841)
 9. Hall A, Selim Anwar K, Tomkins A, Rahman L. 1999 The distribution of *Ascaris lumbricoides* in human hosts: a study of 1765 people in Bangladesh. *Trans. R. Soc. Trop. Med. Hyg.* **93**, 503–510. (doi:10.1016/S0035-9203(99)90351-6)
 10. Easton AV *et al.* 2016 Multi-parallel qPCR provides increased sensitivity and diagnostic breadth for gastrointestinal parasites of humans: field-based inferences on the impact of mass deworming. *Parasit. Vectors* **9**, 1–12. (doi:10.1186/s13071-016-1314-y)
 11. Hou PC, Pang LSC. 1964 *Clonorchis sinensis* infestation in man in Hong Kong. *J. Pathol. Bacteriol.* **87**, 245–250. (doi:10.1002/path.1700870204)
 12. Cheever AW. 1968 A quantitative post-mortem study of schistosomiasis mansoni in man. *Am. J. Trop. Med. Hyg.* **17**, 38–64. (doi:10.4269/ajtmh.1968.17.38)
 13. Kamel IA, Cheever AW, Elwi AM, Mosimann JE, Danner R. 1977 *Schistosoma mansoni* and *S. haematobium* infections in Egypt. I. Evaluation of techniques for recovery of worms and eggs at necropsy. *Am. J. Trop. Med. Hyg.* **26**, 696–701. (doi:10.4269/ajtmh.1977.26.696)
 14. Hall A, Holland C. 2000 Geographical variation in *Ascaris lumbricoides* fecundity and its implications for helminth control. *Parasitol. Today* **16**, 540–544. (doi:10.1016/S0169-4758(00)01779-8)
 15. Bundy DAP, Golden MHN. 1987 The impact of host nutrition on gastrointestinal helminth populations. *Parasitology* **95**, 623–635. (doi:10.1017/S003118200058042)
 16. Elkins DB, Haswell-Elkins M, Anderson RM. 1988 The importance of host age and sex to patterns of reinfection with *Ascaris lumbricoides* following mass anthelmintic treatment in a South Indian fishing community. *Parasitology* **96**, 171–184. (doi:10.1017/S0031182000081749)
 17. De Vlas SJ, Gryseels B, Van Oortmarssen GJ, Polderman AM, Habbema JDF. 1992 A model for variations in single and repeated egg counts in *Schistosoma mansoni* infections. *Parasitology* **104**, 451–460. (doi:10.1017/S003118200006371X)
 18. Irvine MA, Njenga SM, Gunawardena S, Njeri Wamae C, Cano J, Brooker SJ, Deirdre Hollingsworth T. 2016 Understanding the relationship between prevalence of microfilariae and antigenaemia using a model of lymphatic filariasis infection. *Trans. R. Soc. Trop. Med. Hyg.* **110**, 118–124. (doi:10.1093/trstmh/trv096)
 19. Borlase A, Rudge JW, Léger E, Diouf ND, Fall CB, Diop SD, Catalano S, Sène M, Webster JP. 2021 Spillover, hybridization, and persistence in schistosome transmission dynamics at the human–animal interface. *Proc. Natl Acad. Sci. USA* **118**, e2110711118. (doi:10.1073/pnas.2110711118)
 20. Suwannatrai A, Saichua P, Haswell M. 2018 Epidemiology of *Opisthorchis viverrini* infection. *Adv. Parasitol.* **101**, 41–67. (doi:10.1016/bs.apar.2018.05.002)
 21. Sripa B, Brindley PJ, Mulvenna J, Laha T, Smout MJ, Mairiang E, Bethony JM, Loukas A. 2012 The tumorigenic liver fluke *Opisthorchis viverrini*—multiple pathways to cancer. *Trends Parasitol.* **28**, 395–407. (doi:10.1016/j.pt.2012.07.006)
 22. Arunsan P *et al.* 2019 Programmed knockout mutation of liver fluke granulin attenuates virulence of infection-induced hepatobiliary morbidity. *Elife* **8**, e41463. (doi:10.7554/eLife.41463)
 23. Haswell-Elkins MR, Mairiang E, Mairiang P, Chaikakum J, Chamadol N, Loapaiboon V, Sithithaworn P, Elkins DB. 1994 Cross-sectional study of *Opisthorchis viverrini* infection and cholangiocarcinoma in communities within a high-risk area in northeast Thailand. *Int. J. Cancer* **59**, 505–509. (doi:10.1002/ijc.2910590412)
 24. Sriamporn S, Pisani P, Pipitgool V, Suwanrungruang K, Kamsa-Ard S, Parkin DM. 2004 Prevalence of *Opisthorchis viverrini* infection and incidence of cholangiocarcinoma in Khon Kaen, Northeast Thailand. *Trop. Med. Int. Health* **9**, 588–594. (doi:10.1111/j.1365-3156.2004.01234.x)
 25. Upatham ES, Vivanant V, Brockelman WY, Kurathong S, Lee P, Kraengraeng R. 1988 Rate of reinfection by *Opisthorchis viverrini* in an endemic northeast Thai community after chemotherapy. *Int. J. Parasitol.* **18**, 643–649. (doi:10.1016/0020-7519(88)90099-9)
 26. Mairiang E, Haswell-Elkins MR, Mairiang P, Sithithaworn P, Elkins DB. 1993 Reversal of biliary tract abnormalities associated with *Opisthorchis viverrini* infection following praziquantel treatment. *Trans. R. Soc. Trop. Med. Hyg.* **87**, 194–197. (doi:10.1016/0035-9203(93)90489-D)
 27. World Health Organization. 2021 Ending the neglect to attain the sustainable development goals: a sustainability framework for action against Neglected Tropical Diseases 2021–2030. Technical report, World Health Organization.
 28. Wattanawong O *et al.* 2021 Current status of helminthiasis in Thailand: a cross-sectional, nationwide survey, 2019. *Acta Trop.* **223**, 106082. (doi:10.1016/j.actatropica.2021.106082)
 29. Déirdre Hollingsworth T. 2018 Counting down the 2020 goals for 9 neglected tropical diseases: what have we learned from quantitative analysis and transmission modeling? *Clin. Infect. Dis.* **66**, S237–S244. (doi:10.1093/cid/ciy284)
 30. Crellen T, Sithithaworn P, Pitaksakulrat O, Khuntikeo N, Medley GF, Déirdre Hollingsworth T. 2021 Towards evidence-based control of *Opisthorchis viverrini*. *Trends Parasitol.* **37**, 370–380. (doi:10.1016/j.pt.2020.12.007)
 31. Elkins DB, Sithithaworn P, Haswell-Elkins M, Kaewkes S, Awacharagan P, Wongratanchewin S. 1991 *Opisthorchis viverrini*: relationships between egg counts, worms recovered and antibody levels within an endemic community in northeast Thailand. *Parasitology* **102**, 283–288. (doi:10.1017/S0031182000062600)
 32. Sithithaworn P *et al.* 1991 Quantitative post-mortem study of *Opisthorchis viverrini* in man in North-East Thailand. *Trans. R. Soc. Trop. Med. Hyg.* **85**, 765–768. (doi:10.1016/0035-9203(91)90449-9)
 33. Sithithaworn P, Yongvanit P, Tesana S, Pairajkul C. 2007 Liver flukes. In *Food-borne parasitic zoonoses* (eds DK Murrell, B Fried), pp. 3–52. Berlin, Germany: Springer.
 34. Sithithaworn P, Tesana S, Pipitgool V, Kaewkes S, Pairajkul C, Sripa B, Paupairoj A, Thaiklar K. 1991 Relationship between faecal egg count and worm burden of *Opisthorchis viverrini* in human autopsy cases. *Parasitology* **102**, 277–281. (doi:10.1017/S0031182000062594)
 35. Ramsay RJ, Sithithaworn P, Prociw P, Moorhouse DE, Methaphat C. 1989 Density-dependent fecundity of *Opisthorchis viverrini* in humans, based on faecal recovery of flukes. *Trans. R. Soc. Trop. Med. Hyg.* **83**, 241–242. (doi:10.1016/0035-9203(89)90662-7)
 36. Haswell-Elkins MR *et al.* 1994 Liver fluke infection and cholangiocarcinoma: model of endogenous nitric oxide and extragastric nitrosation in human carcinogenesis. *Mutat. Res. Fundam. Mol. Mech. Mutagen.* **305**, 241–252. (doi:10.1016/0027-5107(94)90244-5)
 37. Sayasone S, Vonghajack Y, Vanmany M, Rasphone O, Tesana S, Utzinger J, Akkhavong K, Odermatt P. 2009 Diversity of human intestinal helminthiasis in Lao PDR. *Trans. R. Soc. Trop. Med. Hyg.* **103**, 247–254. (doi:10.1016/j.trstmh.2008.10.011)
 38. Wykoff DE, Ariyaprakai K. 1966 *Opisthorchis viverrini* in Thailand—egg production in man and laboratory animals. *J. Parasitol.* **52**, 631. (doi:10.2307/3276418)
 39. Sato M *et al.* 2015 Patterns of trematode infections of *Opisthorchis viverrini* (Opisthorchiidae) and *Haplorchis taichui* (Heterophyidae) in human populations from two villages in Savannakhet Province, Lao PDR. *J. Helminthol.* **89**, 439–445. (doi:10.1017/S0022149X14000261)
 40. Walker M, Hall A, Anderson RM, Basáñez M-G. 2009 Density-dependent effects on the weight of female *Ascaris lumbricoides* infections of humans and its impact on patterns of egg production. *Parasit. Vectors* **2**, 1–18. (doi:10.1186/1756-3305-2-1)
 41. Fowler AC, Déirdre Hollingsworth T. 2016 The dynamics of *Ascaris lumbricoides* infections. *Bull. Math. Biol.* **78**, 815–833. (doi:10.1007/s11538-016-0164-2)
 42. Bärenbold O, Raso G, Coulibaly JT, N'Goran EK, Utzinger J, Vounatsou P. 2017 Estimating sensitivity of the Kato–Katz technique for the diagnosis of *Schistosoma mansoni* and hookworm in relation to infection intensity. *PLoS Negl. Trop. Dis.* **11**, e0005953. (doi:10.1371/journal.pntd.0005953)
 43. Walker M, Hall A, Basáñez M-G. 2013 *Ascaris lumbricoides*: new epidemiological insights and mathematical approaches. In *Ascaris: the neglected parasite*, pp. 155–201. London, UK: Academic Press.

44. Medley GF, Turner HC, Baggeley RF, Holland C, Hollingsworth TD. 2016 The role of more sensitive helminth diagnostics in mass drug administration campaigns: elimination and health impacts. *Adv. Parasitol.* **94**, 343–392. (doi:10.1016/bs.apar.2016.08.005)
45. Shen C, Kim J-H, Lee J-K, Bae YM, Choi M-H, Oh J-K, Lim MK, Shin H-R, Hong S-T. 2007 Collection of *Clonorchis sinensis* adult worms from infected humans after praziquantel treatment. *Korean J. Parasitol.* **45**, 149. (doi:10.3347/kjp.2007.45.2.149)
46. Kim J-H, Choi M-H, Bae YM, Oh J-K, Lim MK, Hong S-T. 2011 Correlation between discharged worms and fecal egg counts in human clonorchiasis. *PLoS Negl. Trop. Dis.* **5**, e1339. (doi:10.1371/journal.pntd.0001339)
47. Upatham ES, Viyanant V. 2003 *Opisthorchis viverrini* and opisthorchiasis: a historical review and future perspective. *Acta Trop.* **88**, 171–176. (doi:10.1016/j.actatropica.2003.01.001)
48. Rose C, Parker A, Jefferson B, Cartmell E. 2015 The characterization of feces and urine: a review of the literature to inform advanced treatment technology. *Crit. Rev. Environ. Sci. Technol.* **45**, 1827–1879. (doi:10.1080/10643389.2014.1000761)
49. Carpenter B *et al.* 2017 Stan: a probabilistic programming language. *J. Stat. Softw.* **76**, 1–32. (doi:10.18637/jss.v076.i01)
50. Stan Development Team. 2021 CmdStanR: the R interface to CmdStan. R package version 0.4.0. See <http://mc-stan.org/>.
51. McElreath R. 2020 *Statistical rethinking: a Bayesian course with examples in R and Stan*. London, UK: Chapman and Hall/CRC.
52. R Core Team. 2022 *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. See www.R-project.org/.
53. Hall A. 1981 Quantitative variability of nematode egg counts in faeces: a study among rural Kenyans. *Trans. R. Soc. Trop. Med. Hyg.* **75**, 682–687. (doi:10.1016/0035-9203(81)90148-6)
54. Basáñez M-G, Rodríguez-Pérez MA, Reyes-Villanueva F, Collins RC, Henry Rodríguez M. 1998 Determination of sample sizes for the estimation of *Onchocerca volvulus* (Filarioidea: Onchocercidae) infection rates in biting populations of *Simulium ochraceum* s.l. (Diptera: Simuliidae) and its application to ivermectin control programs. *J. Med. Entomol.* **35**, 745–757. (doi:10.1093/jmedent/35.5.745)
55. Kura K, Truscott JE, Collyer BS, Phillips A, Garba A, Anderson RM. 2022 The observed relationship between the degree of parasite aggregation and the prevalence of infection within human host populations for soil-transmitted helminth and schistosome infections. *Trans. R. Soc. Trop. Med. Hyg.* **116**, 1226–1229. (doi:10.1093/trstmh/tra033)
56. Turner HC, Bettis AA, Dunn JC, Whitton JM, Déirdre Hollingsworth T, Fleming FM, Anderson RM. 2017 Economic considerations for moving beyond the Kato–Katzen technique for diagnosing intestinal parasites as we move towards elimination. *Trends Parasitol.* **33**, 435–443. (doi:10.1016/j.pt.2017.01.007)
57. World Health Organization. 2017 Expert consultation to accelerate control of foodborne trematode infections, taeniasis and cysticercosis, Seoul, Republic of Korea, 17–19 May 2017: meeting report. World Health Organization.
58. Anderson R, Truscott J, Déirdre Hollingsworth T. 2014 The coverage and frequency of mass drug administration required to eliminate persistent transmission of soil-transmitted helminths. *Phil. Trans. R. Soc. B* **369**, 20130435. (doi:10.1098/rstb.2013.0435)
59. Minter A, Pellis L, Medley GF, Déirdre Hollingsworth T. 2021 What can modeling tell us about sustainable end points for neglected tropical diseases? *Clin. Infect. Dis.* **72**(S129–S133). (doi:10.1093/cid/ciab188)
60. Churcher TS, Ferguson NM, Basanez M-G. 2005 Density dependence and overdispersion in the transmission of helminth parasites. *Parasitology* **131**, 121–132. (doi:10.1017/S0031182005007341)
61. Bürlü C, Harbrecht H, Odermatt P, Sayasone S, Chitnis N. 2018 Mathematical analysis of the transmission dynamics of the liver fluke, *Opisthorchis viverrini*. *J. Theor. Biol.* **439**, 181–194. (doi:10.1016/j.jtbi.2017.11.020)
62. Asavarut P, Norsworthy PJ, Cook J, Taylor-Robinson SD, Harrison RV. 2016 Diet and disease: transgressing boundaries between science and society—understanding neglected diseases through the lens of cultural studies and anthropology. *Med. Humanit.* **42**, 181–183. (doi:10.1136/medhum-2016-010893)
63. Suwannatitorn P, Webster J, Riley S, Mungthin M, Donnelly CA. 2019 Uncooked fish consumption among those at risk of *Opisthorchis viverrini* infection in central Thailand. *PLoS ONE* **14**, e0211540. (doi:10.1371/journal.pone.0211540)
64. Sweeny AR, Clerc M, Pontifex PA, Venkatesan S, Babayan SA, Pedersen AB. 2021 Supplemented nutrition decreases helminth burden and increases drug efficacy in a natural host–helminth system. *Proc. R. Soc. B* **288**, 20202722. (doi:10.1098/rspb.2020.2722)
65. Lovis L, Mak TK, Phongluxa K, Soukhathammavong P, Sayasone S, Akkhavong K, Odermatt P, Keiser J, Felger I. 2009 PCR diagnosis of *Opisthorchis viverrini* and *Haplorchis taichui* infections in a Lao community in an area of endemicity and comparison of diagnostic methods for parasitological field surveys. *J. Clin. Microbiol.* **47**, 1517–1523. (doi:10.1128/JCM.02011-08)
66. Buathong S, Phaiphilai K, Ruang-Areerate T, Sittichot N, Thita T, Mungthin M, Suwannatitorn P. 2020 Genetic differentiation of *Opisthorchis*-like eggs in Northern Thailand using stool specimens under national strategic plan to control liver fluke infection and cholangiocarcinoma. *Am. J. Trop. Med. Hyg.* **103**, 1118–1124. (doi:10.4269/ajtmh.20-0231)
67. Worasith C *et al.* 2019 Comparing the performance of urine and copro-antigen detection in evaluating *Opisthorchis viverrini* infection in communities with different transmission levels in Northeast Thailand. *PLoS Negl. Trop. Dis.* **13**, e0007186. (doi:10.1371/journal.pntd.0007186)
68. Worasith C *et al.* 2022 Effects of day-to-day variation of *Opisthorchis viverrini* antigen in urine on the accuracy of diagnosing opisthorchiasis in Northeast Thailand. *PLoS ONE* **17**, e0271553. (doi:10.1371/journal.pone.0271553)
69. Inês Neves M, Webster JP, Walker M. 2019 Estimating helminth burdens using sibship reconstruction. *Parasit. Vectors* **12**, 1–12. (doi:10.1186/s13071-019-3687-1)
70. Berger DJ *et al.* 2021 Whole-genome sequencing of *Schistosoma mansoni* reveals extensive diversity with limited selection despite mass drug administration. *Nat. Commun.* **12**, 1–14. (doi:10.1038/s41467-021-24958-0)
71. Heesterbeek JAP, Roberts MG. 2015 How mathematical epidemiology became a field of biology: a commentary on Anderson and May (1981) 'The population dynamics of microparasites and their invertebrate hosts'. *Phil. Trans. R. Soc. B* **370**, 20140307. (doi:10.1098/rstb.2014.0307)
72. Worby CJ, Lipsitch M, Hanage WP. 2017 Shared genomic variants: identification of transmission routes using pathogen deep-sequence data. *Am. J. Epidemiol.* **186**, 1209–1216. (doi:10.1093/aje/kwx182)
73. Grubaugh ND, Ladner JT, Lemey P, Pybus OG, Rambaut A, Holmes EC, Andersen KG. 2019 Tracking virus outbreaks in the twenty-first century. *Nat. Microbiol.* **4**, 10–19. (doi:10.1038/s41564-018-0296-2)
74. Basanez M-G, McCarthy JS, French MD, Yang G-J, Walker M, Gambhir M, Prichard RK, Churcher TS. 2012 A research agenda for helminth diseases of humans: modelling for control and elimination. *PLoS Negl. Trop. Dis.* **6**, e1548. (doi:10.1371/journal.pntd.0001548)
75. Déirdre Hollingsworth T *et al.* 2015 Quantitative analyses and modelling to support achievement of the 2020 goals for nine neglected tropical diseases. *Parasit. Vectors* **8**, 1–28.
76. O'Hara RB, Kotze DJ. 2010 Do not log-transform count data. *Methods Ecol. Evol.* **1**, 118–122. (doi:10.1111/j.2041-210X.2010.00021.x)
77. Hilborn R, Mangel M. 1997 *The ecological detective: confronting models with data*. Princeton, NJ: Princeton University Press.
78. Anderson DR, Burnham KP, Thompson WL. 2000 Null hypothesis testing: problems, prevalence, and an alternative. *J. Wildl. Manage.* **64**, 912–923. (doi:10.2307/3803199)
79. Bolker BM. 2008 *Ecological models and data in R*. Princeton, NJ: Princeton University Press.
80. Crellen T, Haswell M, Sithithaworn P, Sayasone S, Odermatt P, Lamberton PHL, Spencer SEF, Déirdre Hollingsworth T. 2023 Data from: Diagnosis of helminths depends on worm fecundity and the distribution of parasites within hosts. Dryad Digital Repository. (doi:10.5061/dryad.q83bk3jn6)