



Open Access

INVITED COMMENTARY

Teasing apart the multiple roles of Shp2 (*Ptpn11*) in spermatogenesis

Geoffrey J Maher*, Anne Goriely

Asian Journal of Andrology (2020) 22, 122; doi: 10.4103/aja.aja_79_19; published online: 30 July 2019

Male germ cells are the only adult cells to undergo the intricate process of meiosis, which is preceded by the complex molecular transition from mitotic spermatogonia to spermatocytes. Recent single-cell transcriptomic studies have shown that this transition involves differential expression of thousands of genes.^{1,2} Pinpointing the specific roles played by key regulators of this process is crucial to further our understanding of male fertility and, in the long-term, to support the development of infertility therapy and *in vitro* spermatogenesis.

The tyrosine phosphatase Shp2 (encoded by the gene *Ptpn11*) has previously been shown to have multiple roles during mouse spermatogenesis: loss of Shp2 expression in Sertoli cells disrupts the integrity of the blood–testis barrier and also results in a reduction of spermatogonia number and increased expression of differentiation-promoting factors.³ Knockout of Shp2 in the embryonic precursor germ cells (gonocytes) demonstrated that this protein is essential for the production of undifferentiated spermatogonia.⁴ However, although both Shp2 inhibition or global knockout suggested a role in adult germ cells,⁵ its specific function in this cell population has yet to be established.

In *Asian Journal of Andrology*, Li *et al.*⁶ address this aspect by investigating the role of Shp2 in the postnatal germ cells using a *Stra8-Cre* driver strain. They show that this germ cell-specific knockout causes increased differentiation of spermatogonia and reduced numbers of meiotic cells. Abnormal expression of numerous genes/proteins associated with meiotic recombination and synapsis was also identified, consistent with a role for Shp2 in the transition from spermatogonia to spermatocytes.

In humans, a role for SHP2 in spermatogenesis has previously been inferred from genetic studies which revealed that pathogenic gain-of-function mutations in *PTPN11* are associated with conferring a “selfish” selective advantage to spermatogonia. This results in the formation of mutant clones that spread within seminiferous tubules with age.⁷ If passed on to progeny, these mutations cause the developmental disorder Noonan syndrome. Given this, it would be interesting to assess the effects of *Ptpn11* gain-of-function mutations

in murine adult spermatogonia – which may provide further clues on the role of SHP2 in spermatogenesis and human disease.

Although Li *et al.*⁶ propose that activation of SHP2 could be a potential target for the treatment of male infertility, given that *PTPN11* is a known oncogene, implementing such an approach could carry some risks. On the other hand, data from Li *et al.*⁶ and from previous studies⁵ suggest that permanent infertility could be a side effect of therapies being developed to inhibit SHP2.⁸

Delineating the role of *Ptpn11*/Shp2 in adult spermatogenesis provides a valuable insight into the multifaceted functions of this key disease gene.

COMPETING INTERESTS

Both authors declare no competing interests.

REFERENCES

- 1 Guo J, Grow EJ, Mlcochova H, Maher GJ, Lindskog C, *et al.* The adult human testis transcriptional cell atlas. *Cell Res* 2018; 28: 1141–57.
- 2 Hermann BP, Cheng K, Singh A, Roa-De La Cruz L, Mutoji KN, *et al.* The mammalian spermatogenesis single-cell transcriptome, from spermatogonial stem cells to spermatids. *Cell Rep* 2018; 25: 1650–67.
- 3 Hu X, Tang Z, Li Y, Liu W, Zhang S, *et al.* Deletion of the tyrosine phosphatase Shp2 in Sertoli cells causes infertility in mice. *Sci Rep* 2015; 5: 12982.
- 4 Puri P, Phillips BT, Suzuki H, Orwig KE, Rajkovic A, *et al.* The transition from stem cell to progenitor spermatogonia and male fertility requires the SHP2 protein tyrosine phosphatase. *Stem Cells* 2014; 32: 741–53.
- 5 Puri P, Walker WH. The tyrosine phosphatase SHP2 regulates Sertoli cell junction complexes. *Biol Reprod* 2013; 88: 59.
- 6 Li Y, Liu WS, Yi J, Kong SB, Ding JC, *et al.* The role of tyrosine phosphatase Shp2 in spermatogonial differentiation and spermatocyte meiosis. *Asian J Androl* 2019; Doi: 10.4103/aja.aja_49_19. [Epub ahead of print].
- 7 Maher GJ, Ralph HK, Ding Z, Koelling N, Mlcochova H, *et al.* Selfish mutations dysregulating RAS-MAPK signaling are pervasive in aged human testes. *Genome Res* 2018; 28: 1779–90.
- 8 Frankson R, Yu ZH, Bai Y, Li Q, Zhang RY, *et al.* Therapeutic targeting of oncogenic tyrosine phosphatases. *Cancer Res* 2017; 77: 5701–5.

Clinical Genetics Group, MRC Weatherall Institute of Molecular Medicine, University of Oxford, Oxford OX3 9DS, UK.

*Current Address: ICTEM Building, Hammersmith Campus, Imperial College London, Du Cane Road, London, W12 0NN, UK.

Correspondence: Dr. GJ Maher (geoffrey.maher@imperial.ac.uk)

Received: 20 May 2019; Accepted: 28 May 2019

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

©The Author(s) (2019)