

1 *Tansley Review*

2 ***Senecio* as a model system for integrating studies of genotype, phenotype and fitness**

3 Greg M. Walter<sup>1\*</sup> <https://orcid.org/0000-0002-0883-3440> @natureantics

4 Richard J. Abbott<sup>2</sup> <https://orcid.org/0000-0002-4146-5969>

5 Adrian C. Brennan<sup>3</sup> <https://orcid.org/0000-0002-8171-769X>

6 Jon R. Bridle<sup>1</sup> <https://orcid.org/0000-0002-5999-0307> @jonrbridle

7 Mark Chapman<sup>4</sup> <https://orcid.org/0000-0002-7151-723X> @CUCsoton

8 James Clark<sup>5</sup> <https://orcid.org/0000-0003-2896-1631> @boynamedjames

9 Dmitry Filatov<sup>5</sup> <https://orcid.org/0000-0001-8077-5452>

10 Bruno Nevado<sup>5</sup> <https://orcid.org/0000-0002-9765-2907>

11 Daniel Ortiz-Barrientos<sup>6</sup> <https://orcid.org/0000-0002-7493-416X> @dortizba

12 Simon J. Hiscock<sup>5\*</sup>

13 <sup>1</sup>University of Bristol, School of Biological Sciences, Bristol BS8 1TQ, UK

14 <sup>2</sup>University of St Andrews, School of Biology, St. Andrews, Fife KY16 9TH, UK

15 <sup>3</sup>University of Durham, School of Biological and Biomedical Sciences, Durham, DH1 3LE, UK

16 <sup>4</sup>University of Southampton, School of Biological Sciences, Southampton, SO17 1BJ, UK

17 <sup>5</sup>University of Oxford, Department of Plant Sciences, Oxford, OX1 3RB, UK

18 <sup>6</sup>University of Queensland, School of Biological Sciences, St. Lucia QLD 4072, Australia

19 \* Corresponding Authors:

20 Greg M Walter  
21 [g.walter@bristol.ac.uk](mailto:g.walter@bristol.ac.uk)  
Ph: +44 (0)7305 866305

22 Simon J. Hiscock  
23 [simon.hiscock@obg.ox.ac.uk](mailto:simon.hiscock@obg.ox.ac.uk)  
24 Ph: +44 (0)1865 610302

25

<b>Word count</b> Total word count = 7,998 Summary = 199 No. figures = 4 (all colour) No. tables = 1 No. boxes = 1 No. supporting information = 1	<b>Content list</b> Summary I. Model systems for experimental ecology and evolution II. <i>Senecio</i> as an experimental model system III. Mating system diversity and evolution in <i>Senecio</i> IV. Mechanisms and consequences of hybrid speciation V. Adaptation and speciation in <i>Senecio</i> VI. Genomics of adaptation and speciation VII. Integrating studies of genotype, phenotype and fitness VIII. Conspectus Acknowledgements References
---	---

28 **Summary**

29 Two major developments have made it possible to use examples of ecological radiations as model systems to  
30 understand evolution and ecology. First, the integration of quantitative genetics with ecological experiments  
31 allows detailed connections to be made between genotype, phenotype and fitness in the field. Second,  
32 dramatic advances in molecular genetics have created new possibilities for integrating field and laboratory  
33 experiments with detailed genetic sequencing. Combining these approaches allows evolutionary biologists to  
34 better study the interplay between genotype, phenotype and fitness to explore a wide range of evolutionary  
35 processes. Here, we present the genus *Senecio* (Asteraceae) as an excellent system to integrate these  
36 developments, and to address fundamental questions in ecology and evolution. *Senecio* is one of the largest  
37 and most phenotypically diverse genera of flowering plants, containing species ranging from woody  
38 perennials to herbaceous annuals. These *Senecio* species exhibit many growth habits, life histories and  
39 morphologies, and occupy a multitude of environments. Common within the genus are species that have  
40 hybridised naturally, undergone polyploidisation, and colonised diverse environments, often through rapid  
41 phenotypic divergence and adaptive radiation. These diverse experimental attributes make *Senecio* an  
42 attractive model system in which to address a broad range of questions in evolution and ecology.

43

44 **Keywords:** adaptive radiation, Asteraceae, hybrid speciation, quantitative genetics, genomics, model system,  
45 *Senecio*, self-incompatibility

46

## 1. Model systems for experimental ecology and evolution

Recent technological advances in genomic and computational technologies have allowed genetic analyses of many non-model organisms, providing important insights into genome evolution, the structure of developmental pathways and the control of gene expression. However, to answer fundamental questions in ecology and evolutionary biology we need to develop model systems that integrate these novel genetic resources with experimentally flexible systems that are also ecologically diverse. Using ecologically divergent systems is critical for understanding how genotypes relate to phenotypes across different environments, and to test the processes driving adaptation, speciation and biogeographical patterns. Nevertheless, many model systems used in evolutionary biology are only studied in laboratory settings because despite possessing key experimental attributes (e.g. short life-cycles, self-fertility), they show limited variation in life history, mating system, or habitat. The utility of such systems for detailed lab or greenhouse experiments therefore limits their value in field experiments testing hypotheses relating to evolution and ecology, such as the interaction between phenotypic plasticity and natural selection, and how and when populations specialise to particular environments. This limitation has become increasingly relevant given the urgent need to predict how, where, and how quickly populations and ecological communities will respond to ongoing environmental change. Research needs to focus on model systems with populations and species that inhabit different environments and ecosystems, but that remain amenable to large field experiments and quantitative genetic crossing designs. Specifically, we need model systems that can connect genotype, phenotype and fitness under a range of environmental conditions. In this review, we argue that *Senecio* is one of very few model systems that can fill this role.

## 2. *Senecio* as an experimental model system

*Senecio* (Linnaeus, 1753) is a large genus of flowering plants within the tribe Senecioneae of the daisy family, Asteraceae. Containing in excess of 1,200 species distributed across the globe, *Senecio* radiated rapidly during the Miocene c.10.7 million years ago to become one of the largest genera of flowering plants (Kandziora *et al.*, 2017). Species of *Senecio*, commonly known as ragworts and groundsels, are generally herbaceous, but also occur as vines, shrubs, succulents, epiphytes and small trees (**Fig. 1**). The genus is aptly named after the tendency for species to produce senescent flower heads resembling tufts of white hair (Latin ‘senex’ = ‘old man’) formed by the pappus, which facilitate airborne seed dispersal.

Many species of *Senecio* have useful attributes of a model system, including: small to medium-sized genomes (mean c.5 Gbp; range 1.13-51.25 Gbp; Leitch *et al.*, 2019), a rapid sexual cycle (3-5 months from seed-to-seed), ease of crossing because of widespread self-incompatibility. Species are also often perennial or short-lived perennials so can be regenerated from cuttings to create genetically identical clones, are

79 genetically transformable (Kim *et al.*, 2008), and are easily grown in glasshouses and field plots. The fact  
80 that many species show these attributes, combined with their recent radiation and occupation of a diverse  
81 array of habitats, make *Senecio* species more amenable than many other systems for quantitative genetic  
82 designs. Such designs allow researchers to combine genetic and phenotypic studies in both the field and  
83 under controlled conditions. The genus *Mimulus* shares some of these experimental characteristics (e.g., short  
84 generation times, easy to grow, clonal propagation) and has a widespread distribution across different  
85 environments (Wu *et al.*, 2008), but *Senecio* has the added benefit of a worldwide distribution, and a greater  
86 diversity of species that include a range of mating systems and a multitude of growth forms (**Fig. 1**). Such  
87 diversity in morphology and habitat makes *Senecio* attractive to a similar diversity of scientists, from  
88 ecologists to phylogeneticists, molecular geneticists and quantitative geneticists. We therefore propose  
89 *Senecio* as an attractive model system for answering fundamental questions in ecology and evolution, joining  
90 the ranks of *Antirrhinum*, *Arabidopsis* (including *Boechera* and other relatives), *Helianthus*, *Mimulus*, *Silene*  
91 and *Solanum*.

92 *What questions can we answer using Senecio?*

93 Future work will benefit from the imminent completion of high coverage and highly contiguous annotated  
94 genomes for *S. squalidus* and *S. lautus* (B. Nevado *et al.* unpublished; Wilkinson *et al.*, 2019). These will be  
95 used as references for further genomic analyses of other *Senecio* species, including whole genome  
96 resequencing, DNA methylation and RNA-seq approaches. These new resources will make *Senecio* a  
97 powerful system to study the genomic architecture of selection, adaptation and divergence across  
98 environments. Combining such genetic association techniques with quantitative genetic crossing designs is a  
99 powerful way to explore how the relationship between genotype, phenotype and fitness changes across  
100 environments. Of special importance is the ease and speed at which species of *Senecio* can be propagated  
101 from cuttings. This ability to generate clones allows researchers to control for genotype and transplant large  
102 numbers of clones across multiple environments, and sets *Senecio* apart from other model systems. Such  
103 approaches can address an array of questions in ecology, evolution and evolutionary genomics (see: **Box 1**).

## 104 **2.1 Origins and diversity of *Senecio***

105 The presence of repeated radiations in the Americas, Australasia, and across Northern Africa into Europe  
106 highlights the ability of *Senecio* to colonise and rapidly adapt to multiple environments (Comes & Abbott,  
107 2001; Roda *et al.*, 2013; Liew *et al.*, 2018). This broad geographical distribution is reflected in species'  
108 occupation of contrasting environments, from harsh coastlines to rainforests, deserts, mountains and alpine  
109 areas. Associated with this diversity of habitats is an array of different phenotypes (**Fig. 1**) typified by the  
110 Australian *Senecio lautus* species complex, which provides a fascinating example where recent radiation into

more than 10 contrasting environments across much of the continent has resulted in the evolution of ecotypes with a broad diversity of plant architecture and leaf form (**Fig. 2**). Among the numerous recorded radiations within *Senecio* there is evidence of local adaptation (Walter *et al.*, 2016), parallel evolution (Roda *et al.*, 2013; Comes *et al.*, 2017), polyploidy (Alexander, 1979; Chapman & Abbott, 2010), introgressive hybridization (Kim *et al.*, 2008; Osborne *et al.*, 2016), hybrid speciation (James & Abbott, 2005; Hegarty *et al.*, 2012) and apparently stable hybrid zones (Brennan *et al.*, 2009; Abbott *et al.*, 2018).

## 2.2 Historical use of *Senecio* for research

With its wealth of morphological, ecological and reproductive diversity, *Senecio* has attracted studies addressing many different questions in ecology and evolution. Experimental research using *Senecio* began in the late 19<sup>th</sup> Century (**Table 1**) when Anna Bateson, sister of the geneticist, William Bateson, showed that progeny generated by crossing *S. vulgaris* plants were taller and produced more seed per capitulum than those generated by self-fertilization, suggesting an heterotic effect (Bateson, 1888). Because *S. vulgaris* has a short generation time and is self-compatible but easy to cross, Trow recognised its value for genetic research and used it to establish the genetic basis of several discontinuous and semi-continuous traits within the species in the early 20th Century (Trow, 1912, 1916). Approximately 40 years then passed, before Harland emphasised the value of *Senecio* in both genetic and evolutionary research by using interspecific crosses to show that *Senecio cambrensis* was the recent allohexaploid of *S. vulgaris* and *S. squalidus* (Harland, 1954a, b). Through this early work, Bateson, Trow and Harland laid the foundation for more extensive works that followed (**Table 1**).

## 3. Mating system diversity and evolution in *Senecio*

### 3.1 Outcrossing and self-incompatibility

As a member of the Asteraceae, *Senecio* inflorescences are capitula (‘flower heads’), comprising multiple uni-ovulate flowers (florets) tightly gathered across a disc-shaped receptacle. Florets are frequently differentiated into two types dependent on their position within the capitulum (‘disc’). Centrally located disc florets are usually radially symmetrical while outermost ray florets are bilaterally symmetrical with petals fused into a single large ‘ray’ (Abbott & Schmitt, 1985; Gillies *et al.*, 2002; Kim *et al.*, 2008; Garcês *et al.*, 2016), which facilitates outcrossing by attracting generalist insect pollinators (Abbott & Irwin, 1988; Andersson & Widén, 1993).

The genetic control of flower development has been studied in *S. vulgaris* (ssp. *vulgaris*), which is mostly ray-less throughout its range but has a rayed variant in the UK (*S. vulgaris* var. *hibernicus*) that originated via introgressive hybridization with *S. squalidus* (Abbott *et al.*, 1992; Kim *et al.*, 2008). The emergence of a

rayed form increased the attractiveness of *S. vulgaris* to insect pollinators (Abbott & Irwin, 1988) and increased maternal outcrossing in the newly formed radiate variant (Marshall & Abbott, 1982, 1984). This provides a rare example of a shift towards outcrossing in a selfing species (Abbott *et al.*, 1998), in contrast to the more typical trend for selfing to evolve from outcrossing (Barrett, 2013). Floret development in this rayed variant is controlled by localised expression of, and interactions between, at least three *cycloidea* (*CYC*)-like genes (Kim *et al.*, 2008; Garcês *et al.*, 2016). There is evidence for selection on a duplicated cluster of these genes, collectively known as *Ray*, in the form of gene transfer between species and clinal patterns of variation across ecotones, reflecting the adaptive importance of ray florets as a modulator of outcrossing (Kim *et al.*, 2008; Chapman & Abbott, 2010).

Outcrossing in many *Senecio* species is maintained by the presence of self-incompatibility (SI), a molecular mechanism of self-pollen recognition and rejection (Hiscock, 2000a, b) that is common among species within the Asteraceae (Ferrer & Good-Avila, 2007). SI in *Senecio* is controlled sporophytically with a single genetic locus, *S*, determining both the pollen and stigma recognition phenotypes (Hiscock, 2000a, b; Brennan *et al.*, 2002; 2006; 2011). Interestingly, the expression of sporophytic SI (SSI) in *Senecio* appears to be affected by additional unlinked modifier loci (Hiscock, 2000a, b; Brennan *et al.*, 2011), which raises intriguing questions about its underlying molecular mechanism. To date however, unravelling the molecular basis of SSI in *Senecio* has proved challenging (Allen *et al.*, 2011). Those studies that have been undertaken provide clear evidence that this molecular mechanism is distinct from the well-characterised SSI system of the Brassicaceae (Tabah *et al.*, 2004; McInnis *et al.*, 2006; Allen *et al.*, 2011), indicating the evolution of a different mechanism of SSI in the Asteraceae and further supporting phylogenetic analyses that have shown that SI has evolved multiple times during the radiation of flowering plants (Allen & Hiscock, 2008). Encouraging progress towards characterising the *S*-locus in members of the Asteraceae is being made through mapping genomic regions linked to SI in sunflower (Gandhi *et al.*, 2005; Gonthier *et al.*, 2013), which are also proving useful in identifying syntenic regions in the *Senecio* genome (B. Nevado *et al.*, unpublished). Nevertheless, it remains a possibility that the Asteraceae harbours more than one SI system, especially as elements of an underlying gametophytic SI system have been described in certain crosses between SSI species of *Parthenium* (Lewis, 1994) and also in *Senecio* (Hiscock, 2000b; Tabah, 2004). This important area of mating system evolution will benefit from further research as more genomic tools emerge for *Senecio*.

### 3.2 Implications of self-incompatibility for genomic structure and regulation

#### *Suppressed recombination around the S locus*

Since both pollen and pistil *S* determinants are required for functional SI, all *S* loci characterised to date have

a haplotypic structure containing at least two tightly linked polyallelic genes responsible for male and female SI function, respectively (Iwano & Takayama, 2012). Recombination between haplotypes is necessarily minimised through extensive structural differences between haplotypes and/or high sequence divergence (Boyes *et al.*, 1997; Shiba *et al.*, 2003; Goubet *et al.*, 2012). This extensive sequence divergence is created by balancing selection that maintains many functionally distinct *S* haplotypes to maximise population mate availability (Schierup & Vekemans, 2008; Roux *et al.*, 2013). The availability of sequenced genomes for *Senecio* species will allow testing of predictions about the genomic structure of the *S* locus in this novel SSI system thereby broadening the taxonomic breadth of studies of the evolution of SI and *S* loci.

### *Dominance in SI systems*

Sporophytic expression of male determinant *S* alleles in pollen creates dominance interactions between *S* alleles that optimise mate availability in populations while effectively restricting self-pollination, even when allelic diversity is relatively low (Vekemans *et al.*, 1998; Brennan *et al.*, 2003; Billiard *et al.*, 2007). The presence of *S* allele dominance interactions (both for pollen and stigma *S* determinants) is thought to have contributed to the colonising success of self-incompatible *S. squalidus* in the UK and *S. inaequidens* in continental Europe despite the presence of limited functional *S* haplotype diversity (Brennan *et al.*, 2002; Brennan *et al.*, 2003, 2006; Lafuma & Maurice, 2007). Moreover, evidence for differences in *S* allele dominance interactions across different UK *S. squalidus* populations suggests that *S* allele dominance is labile and can evolve (Brennan *et al.*, 2006). *Senecio* can be used to determine whether independently evolved systems of SSI have also independently evolved this flexible mechanism of *S* allele dominance regulation. Future research on SSI in *Senecio* could explore the relevance of recent discoveries in the Brassicaceae, such as the control of pollen *S* allele dominance by *S* allele-specific sRNAs (Durand *et al.*, 2014) and the accumulation of inbreeding depression associated with the *S* locus (Billiard *et al.*, 2007; Llaurens *et al.*, 2009), generating an important test of whether such mechanisms apply beyond a single SSI system.

### **3.3 Implications of self-incompatibility for hybridization and polyploidy**

As a mechanism promoting outcrossing, SI also increases opportunities for interspecific hybridization (Brennan *et al.*, 2013; Vallejo-Marin & Hiscock, 2016). Hybridization is recognised as an important evolutionary force that rapidly generates new genetic combinations for selection to act upon, which enhances the spread of adaptive traits across related groups (Abbott *et al.*, 2013; Harrison & Larson, 2014). Strong balancing selection promoting *S* haplotype diversity increases the likelihood of gene flow between taxa at the *S* locus (Castric *et al.*, 2008), as reflected by extensive *S* haplotype sharing between hybridizing *Senecio* species (Brennan *et al.*, 2013).



206 *Senecio* provides many examples of allopolyploid hybrid species, created when interspecific hybridization is  
 207 accompanied by a doubling of a hybrid's chromosome number (Abbott & Lowe, 2004; Kadereit *et al.*, 2006;  
 208 Hegarty *et al.*, 2012; Pelser *et al.*, 2012). Different ploidy levels between species need not present an  
 209 insurmountable barrier to gene flow following hybridization, as shown by examples of gene transfer of the  
 210 *Ray* locus from diploid *S. squalidus* to tetraploid radiate *S. vulgaris* (**Fig. 3**; Kim *et al.*, 2008; Chapman &  
 211 Abbott, 2010). Sometimes ploidy differences can contribute to genetic swamping from the dominant pollen  
 212 donor as sterile progeny with unbalanced chromosome numbers are produced. This process is thought to be  
 213 contributing to the invasive success of diploid *S. madagascariensis* in encroaching upon the habitat occupied  
 214 by native tetraploid *S. pinnatifolius* in Australia (Prentis *et al.*, 2007).

215 Hybridization between *S. squalidus* (SI) and self-compatible (SC) *S. vulgaris* within the last 100 years has  
 216 resulted in the evolution of the allohexaploid hybrid species *S. cambrensis* (SC) on at least two occasions in  
 217 north Wales and Edinburgh (**Fig. 3**; Abbott & Lowe, 2004; Brennan & Hiscock, 2010; Hegarty *et al.*, 2012).  
 218 The process of genome duplication itself might have led to the disruption of the SI system inherited from *S.*  
 219 *squalidus*, resulting in the SC condition in *S. cambrensis*. SC might therefore be selected for in new  
 220 polyploids because they are initially rare with limited *S* haplotype diversity soon after origin, which could  
 221 cause problems with mate availability under obligate outcrossing. However, many polyploid self-  
 222 incompatible *Senecio* species are known (Andersson, 2001; Lafuma & Maurice, 2007) and in resynthesized  
 223 *S. cambrensis* allohexaploids it was found that SI reappeared among some offspring in the F2 generation  
 224 (Brennan & Hiscock, 2010) suggesting complex control of SSI in neopolyploids. These examples  
 225 demonstrate that *Senecio* provides a fascinating study system to better understand how breeding system,  
 226 hybridization, and polyploidy interact in evolution to generate and maintain genetic, taxonomic and  
 227 ecological diversity.

### 228 **3.4 Implications of self-incompatibility for invasiveness and colonisation**

229 According to Baker's Law, self-compatibility is favoured in invasive populations because of enhanced  
 230 reproductive assurance during colonization (Baker, 1967; Pannell & Barrett, 1998). However, many  
 231 exceptions exist, and it is clear that the interactions between breeding systems and colonisation or  
 232 invasiveness can be complex. In the UK, investigations of invasive *S. squalidus* populations have revealed  
 233 individual variation in the strength of SI within and between populations; a phenomenon often referred to as  
 234 pseudo-self-compatibility (PSC; Brennan *et al.*, 2011). Despite SI being predominant in all populations of *S.*  
 235 *squalidus*, a low frequency of PSC (~6-8%) sometimes higher (40% SC, 13% PSC in an Exeter sample; A.  
 236 Brennan & S. Hiscock, unpublished) was detected in all UK populations sampled, and may be contributing to  
 237 mating and colonizing success (Brennan *et al.*, 2005). PSC might be of considerable adaptive importance in

conferring reproductive assurance through breeding system ‘flexibility’, particularly during periods of population disturbance or establishment when the relative advantages of selfing outweigh those of outcrossing (Levin, 1996; Hiscock, 2000a, b; Stephenson *et al.*, 2000).

PSC and SC in *S. squalidus* are heritable and appear regulated by multiple genetic loci that interact with the *S* locus to modify its expression (Hiscock, 2000b; Tabah, 2004; Brennan *et al.*, 2011) similar to the control of unilateral interspecific incompatibility, which regulates the compatibility of crosses between closely related SI and SC species (Hancock *et al.*, 2003). Environmental factors, such as salt concentration, also affect the expression of SI and PSC (Hiscock, 2000a, b), potentially providing a mechanism for the environment to have direct effects on recombination rates in natural populations (Stevison *et al.*, 2017). Furthermore, it has been demonstrated experimentally that recombination can ‘resurrect’ SI to varying levels of penetrance in SC lines of synthetic *Senecio* allopolyploids generated by crossing an SC (*S. vulgaris*: 4n) with an SI species (*S. squalidus*: 2n) (Brennan & Hiscock, 2010). Therefore, rather than a qualitative trait (as it has traditionally been viewed), SI *sensu lato* could be viewed as a quantitative trait at the end of a potentially flexible continuum from SI through PSC to SC. These studies highlight the potential of *Senecio* as an informative model system for investigating evolutionary interactions between breeding systems and invasiveness.

#### 4. Mechanisms and consequences of hybrid speciation

Historical and current interspecific hybridization is common in *Senecio* (e.g., Comes & Abbott, 2001; Pelser *et al.*, 2012; Osborne *et al.*, 2016), occurring both within and across ploidy levels, and between selfing and outcrossing species (e.g., Kadereit *et al.*, 2006; Chapman & Abbott, 2010). Indeed, *Senecio* has been highlighted for its exceptional recent rates of hybrid speciation, particularly as opportunities for new interspecific interactions arise due to human-mediated change (Thomas, 2015). *Senecio* is notable for containing examples of both recent homoploid hybrid speciation (involving no change in chromosome number) and allopolyploid speciation (inheritance of more than two sets of chromosomes), serving as excellent models for investigating hybrid speciation (**Fig. 3**).

The homoploid hybrid species *S. squalidus* (Oxford ragwort) originated from material introduced to the Oxford Botanic Garden (UK) at the end of the 17th Century (Harris, 2002) derived from hybridization between two species native to Mount Etna, Sicily, *S. aethnensis* and *S. chrysanthemifolius* (**Fig. 3a-b**; James & Abbott, 2005). Following almost a century of cultivation in the Oxford Botanic Garden, the new homoploid hybrid species rapidly spread across Britain and into Ireland over the next 200 years (Abbott *et al.*, 2009). *Senecio squalidus* is genetically and phenotypically divergent from its parents and their hybrids, which are restricted to Sicily (Brennan *et al.*, 2012), with evidence of extreme divergence in gene expression (Hegarty *et al.*, 2008, 2009). In addition to its ecogeographical isolation, recent genetic mapping studies

indicate that *S. squalidus* has inherited a combination of genetic incompatibilities from its parent species contributing further to its reproductive isolation (Brennan *et al.*, 2019). Together this genetic, phenotypic and transcriptomic divergence has likely contributed to its local adaptation to the UK (Allan & Pannell, 2009; Ross, 2010), which is the target of current transcriptomic (RNA-seq) research aimed at identifying candidate genes for local adaptation (Nevado and Hiscock, pers. comm.).

During its spread in Britain, *S. squalidus* has repeatedly hybridised with the native tetraploid, *S. vulgaris*, resulting in the origin of the introgressant *S. vulgaris* var. *hibernicus*, a new tetraploid species, *S. eboracensis* (Irwin & Abbott, 1992; Lowe & Abbott, 2003), and multiple origins of the neoallohexaploid, *S. cambrensis* (Figs. 3-4; Abbott & Lowe, 2004; Hegarty *et al.*, 2012). These hybridisation events were associated with marked changes in gene expression (Hegarty *et al.*, 2006; 2012; 2013) that may have had adaptive consequences by facilitating the establishment of new hybrid taxa through ecological differentiation (see also section 6.3). Multiple independent hybridisation events, as observed in *Senecio*, provide a natural experiment with which to study the consequences of hybridisation and polyploidy in populations. Species of *Senecio* therefore provide abundant systems for future detailed genomic analyses of mechanisms favouring the origin and establishment of hybrid taxa and introgression in the wild (e.g., Alexander-Webber *et al.*, 2016).

## 5. Adaptation and speciation in *Senecio*

### 5.1 The evolution of reproductive isolation during the formation of ecotypes

Ecotypes originate when adaptation to contrasting habitats leads to a build-up of extrinsic reproductive isolation between populations. Understanding how and when ecotypes become independent species remains a fundamental issue in evolutionary biology (Abbott & Comes, 2007; Lowry, 2012). Of particular importance is how ecological divergence becomes associated with intrinsic barriers to gene flow, created by genetic divergence that produce genetic incompatibilities between ecotypes.

In the *S. lautus* complex, which consists of multiple taxonomic ecotypes and species (see Roda *et al.*, 2013), a strong correlation is observed between plant morphology and habitat (Fig. 2; Radford *et al.*, 2004; Walter *et al.*, 2018a). Coastal populations grow tall on sand dunes (Dune ecotype), but prostrate on rocky headlands (Headland ecotype) and cliffs along the Australian coastline (Fig. 2; Roda *et al.*, 2013). Dune and Headland populations often occur in adjacent habitats and are self-incompatible and interfertile. Transplant experiments have revealed that neither Dune nor Headland ecotypes, nor their hybrids, can colonise each other's habitat (Melo *et al.*, 2014; Richards & Ortiz-Barrientos, 2016; Richards *et al.*, 2016; Walter *et al.*, 2016). The probability of surviving in the alternative environment is a function of the proportion of alleles of local origin that a hybrid individual carries (Richards *et al.*, 2016). Together, these observations suggest that limited

dispersal and natural selection against migrants and hybrids keep populations distinct despite their parapatric distribution, and that selection acting in opposing directions has played a fundamental role in the formation of ecotypes in this system, likely leading to the formation of new plant species.

Barriers to gene flow that act after pollination, such as F1 hybrid inviability or sterility, are often masked by heterosis (Lowry *et al.*, 2008). This can bias perception as to how intrinsic and extrinsic reproductive barriers evolve in a system that has diverged recently. For instance, in transplant experiments of *S. lautus*, F1 hybrids show strong hybrid vigour (Walter *et al.*, 2016), suggesting that heterosis masks the deleterious effects of gene flow between populations locally adapted to contrasting habitats. However, a more detailed reciprocal transplant including all combinations of F1 hybrids amongst four ecotypes (**Fig. 2**, Woodland ecotype - dry sclerophyll woodland, and Tableland ecotype - subtropical rainforest edges), showed that field fitness depended on the chromosomal origins in a hybrid (Walter *et al.*, 2016). For each of the four transplant environments, F1 hybrids with a chromosome native to the transplant environment (e.g., Dune in the dune environment) performed better than F1 hybrids where both chromosomes were from foreign ecotypes (e.g., F1<sub>Headland×Tableland</sub> in the dune environment). This suggests that alleles associated with heterosis were also involved in adaptation. F1 hybrids also show vigour in glasshouses when compared to parents, while F2 hybrids perform poorly. The presence of F2 hybrid breakdown suggests that intrinsic postzygotic isolation is created by alleles that reduce fitness when recombined into other genetic backgrounds (Walter *et al.*, 2016).

Although reproductive compatibility seems to be high among ecotypes of *S. lautus*, geographical patterns of reproductive isolation suggest that different populations of the same ecotype are becoming reproductively isolated (Melo *et al.*, 2019). Such patterns provide the opportunity to study the evolution of reproductive isolation along a speciation continuum to identify generalisations for speciation in plants (Melo *et al.*, 2014). Recent work has identified molecular links between adaptation and speciation in *S. lautus* (Melo *et al.*, 2019), and found evidence that adaptive divergence also creates genetic incompatibilities between populations inhabiting contrasting habitats (Walter *et al.*, 2019). Overall, current evidence suggests that natural selection plays a fundamental role in creating barriers to gene flow between *S. lautus* populations and that extrinsic barriers seem to have evolved early during divergence, while intrinsic barriers were initially weak but have accumulated. This work highlights the potential for using *Senecio* to better understand how adaptation and speciation occur across environments and geographical ranges.

## 5.2 Ecological speciation on Mount Etna

Genomic analyses have made it clear that in the presence of divergent selection, evolutionary divergence can occur despite populations continuing to exchange alleles (whether in sympatry or parapatry), with intrinsic reproductive isolation arising as a consequence later (Rundle & Nosil, 2005). This ‘divergence with gene

flow' scenario is of interest to evolutionary biologists because the relative roles of major mechanisms of evolution (ecology, selection, genetic drift and gene flow) can be studied together to understand how phenotypes, genotypes, and alleles of diverging populations interact during speciation.

On Mount Etna, Sicily, two sister *Senecio* species inhabit ecologically distinct, high and low elevation environments. *Senecio aethnensis* (**Fig. 4a**) is found above ca. 2,000m above sea level (a.s.l.) on old lava flows, and is covered by snow each winter, while *Senecio chrysanthemifolius* (**Fig.4b**) occurs in disturbed habitats and along roadsides below ca. 1,000m a.s.l., and endures intense summer heat. At intermediate elevations the species form a hybrid zone (James & Abbott, 2005; Brennan *et al.*, 2009). The parental species show divergence in leaf and capitulum shape and size, with plants in the hybrid zone exhibiting a range of intermediate phenotypes, with clines for different traits varying significantly in width, suggesting selection against at least some intermediate traits (Brennan *et al.*, 2009). The two species are almost completely interfertile with hybrids generally fit and fertile, even in later generations. It is therefore likely that selection is responsible for maintaining the distinctness of the two parental species (Chapman *et al.*, 2005; but see Brennan *et al.* [2014] and Chapman *et al.* [2016] for evidence of incompatibility due to hybrid breakdown).

Using transcriptome sequences from one individual from each of *S. aethnensis* and *S. chrysanthemifolius* (and *S. vernalis* as an outgroup), Osborne *et al.* (2013) found evidence that speciation occurred with gene flow (i.e. in parapatry versus allopatry). Supported by Chapman *et al.* (2013), both studies found similar estimates in mean divergence time between the two species of ca. 108,000 years ago (bootstrap range: 53,000 - 187,000), which coincides with the uplift of Mount Etna that likely produced a novel high altitude habitat in which *S. aethnensis* could have originated (Chapman *et al.*, 2013; Osborne *et al.*, 2013). The well characterised demographic history of the Mt Etna hybrid zone provides an intriguing system and an ideal basis for future work to study the genetic architecture of traits underlying adaptation, speciation and hybrid breakdown, especially in combination with fully annotated genomes for these species.

### 5.3 Introgression and clinal analysis in the formation of a hybrid zone during adaptive divergence

The hybrid zone between populations of *S. aethnensis* and *S. chrysanthemifolius* on Mount Etna has been studied by quantifying how traits (associated with reproduction, growth form and leaf shape) and neutral microsatellite markers change across a hybrid zone (Brennan *et al.*, 2009). This analysis provides evidence for strong selection on leaf traits (relative to neutral markers), which have narrower cline widths (strong selection) than genetic markers (1.49 km compared to 3.24 km), and for selection against hybrids associated with the elevational gradient (1,515-1,795 m), which determines the position of the cline centre. There were also significant increases of covariances and variances of genotypes and ecological variation at the cline centre, suggesting a high frequency of relatively recent hybrids at the centre. A significant displacement of

the cline centre of floral traits compared to genetic markers of 0.41 km suggests that alleles associated with variation in some traits may be moving to higher elevations. Reciprocal transplants throughout the hybrid zone on Mt. Etna (or other *Senecio* hybrid zones), at different spatial scales, and in combination with high-throughput sequencing, can provide an important way to explore the genetic and genomic architecture of traits affecting fitness in *Senecio* in contrasting environments.

## 6. Genomics of adaptation and speciation

### 6.1 Genomics of adaptation and parallel evolution

The genetic basis of adaptation in plants remains largely unexplored outside model and crop systems. Notable exceptions are wild sunflowers (*Helianthus* spp.), rockcress (*Boechera stricta*) and monkeyflowers (*Mimulus* spp.), where studies have revealed the genetic architecture and molecular basis of adaptive traits, including flowering time (Blackman *et al.*, 2011), flowering phenology (Anderson *et al.*, 2013), physiological adaptations to contrasting environments (Rieseberg *et al.*, 2003) and drought resistance (Lowry & Willis, 2010; VanWallendael *et al.*, 2019). In *S. lautus*, where multiple pairs of dune and headland populations have evolved repeatedly and independently along the Australian coast, genomic approaches are providing important insights into the genetic changes associated with rapid adaptive divergence.

Using single nucleotide polymorphisms (SNPs) distributed across the genome, Roda *et al.* (2013) showed that different pairs of parapatric dune and headland populations had unique patterns of genetic differentiation, suggesting independent adaptive differentiation. Although patterns of population pair differentiation were unique at the SNP level, they were similar at the level of predicted function, notably for development and growth via auxin. This suggests that paths towards adaptation and patterns of repeated speciation (between ecotypes for multiple population pairs) are often different but lead to similar end results in terms of phenotypes. Current research using *S. lautus* is exploring the molecular basis of auxin and gravity responses (Wilkinson *et al.*, 2019). Within *Senecio*, the parallel evolution of traits across populations occurs in many other species (e.g., Ni accumulation in *S. coronatus* [Meier *et al.* 2018] and ecotype formation in Mediterranean *Senecio* [Comes *et al.* 2017]). The existence of multiple adaptive radiations in *Senecio*, each involving adaptation to similar or contrasting environmental stresses highlights the potential of the genus to understand the ecological genetics of adaptation and speciation.

### 6.2 Population genomics of ecological speciation

The interaction between recombination, gene flow and selection during speciation are still poorly understood but actively researched both theoretically (e.g., Schilling *et al.*, 2018) and empirically (e.g., Chapman *et al.*, 2016; Safran *et al.*, 2016; Doellman *et al.*, 2018; Stankowski *et al.*, 2019). *Senecio* contains multiple systems

where these evolutionary dynamics can be studied. For example, despite differing profoundly in phenotypes, life history, and ecology, *S. aethnensis* and *S. chrysanthemifolius* on Mount Etna show surprisingly little genomic divergence (Chapman *et al.*, 2013, 2016; Filatov *et al.*, 2016). Osborne *et al.* (2016) demonstrated extensive contemporary gene flow between the two species, supported by a demographic analysis of the species (Chapman *et al.*, 2013), with more gene flow from *S. aethnensis* to *S. chrysanthemifolius* than in the opposite direction. Genomic comparisons between *S. aethnensis* and *S. chrysanthemifolius* revealed the clustering of highly differentiated genetic markers (identified by outlier scan; Chapman *et al.*, 2013) in a handful of small regions of the genome (Chapman *et al.*, 2016). However, their clustering appeared to be associated with low recombination regions and not islands of divergence. QTL mapping demonstrated that outlier genetic markers tended to underlie morphological QTL, suggesting a link between genomic divergence, morphological divergence and adaptation to altitude (Chapman *et al.*, 2016). In *S. lautus*, Roda *et al.* (2017) found evidence for clustering of adaptive alleles associated with variation in both architectural and reproductive traits. This suggests that pre- and post-pollination reproductive barriers can evolve concomitantly and facilitate speciation.

### 6.3 Differential gene expression associated with hybrid speciation and adaptation.

The extent and role of changed patterns of gene expression in local adaptation or the formation of new species via hybridisation is not well understood (Vallejo-Marin & Hiscock, 2016). However, there is considerable interest in the idea that the regions of the genome that show environmental (or trans-generational) mediation of gene expression may be disproportionately involved in local adaptation (West-Eberhard, 2005). Gene expression can also be altered dramatically during the rapid changes in gene number associated with speciation by hybridisation. To investigate how gene expression changes during homoploid and allopolyploid hybrid speciation, Hegarty *et al.* (2005) performed microarray experiments and sequenced genes that showed over or under expression in *Senecio* hybrids, relative to one or both parents. Gene expression was explored in natural and resynthesised allopolyploid hybrids between *Senecio vulgaris* and *S. squalidus* (**Fig. 3**), their primary triploid hybrid *S. x baxteri* ( $3n=3x=30$ ) and their allohexaploid hybrid *S. cambrensis* ( $2n=6x=60$ ) (Hegarty *et al.*, 2005; 2006). Early generation hybrids exhibited massive changes in gene expression affecting hundreds of genes, a phenomenon described as ‘transcriptome shock’ (Hegarty *et al.*, 2005; 2006). Interestingly, changes in levels of gene expression were greater in triploid *S. x baxteri* than in the allohexaploid *S. cambrensis* suggesting that genome duplication, leading to the formation of the allohexaploid, has an ‘ameliorating’ effect on hybridisation-induced ‘transcriptome shock’ (Hegarty *et al.*, 2006).

Transcriptome shock was also observed in the homoploid hybrid *S. squalidus* and in artificial ‘*S. squalidus*’ hybrids (propagated for five generations) produced by crossing its parental taxa (Hegarty *et al.*, 2009). The effect of transcriptome shock was more extreme in the artificial homoploid hybrids. In a subset of genes, transgressive expression (up or down regulation with respect to both parents) was maintained in subsequent generations, while other affected genes tended to move towards a pattern of expression consistent with wild *S. squalidus*. These transgressive genes were involved in photosynthesis, catalytic activity, lipid binding, protein biosynthesis and protein folding (Hegarty *et al.*, 2009). The finding of transgressive expression was consistent with similar microarray analysis in hybrid sunflowers (Lai *et al.*, 2006), which identified transgressive levels of gene expression in wild hybrids adapted to environmental conditions not tolerated by their parent species. Both studies suggest that new patterns of transgressive gene expression arising from hybridization could provide genetic novelty for selection to facilitate adaptive divergence.

Comparisons of gene expression and transcriptome sequences between *S. aethnensis* and *S. chrysanthemifolius* identified genes potentially involved in adaptation to high and low altitudes, and the relative roles of expression and sequence divergence in the genomic basis of adaptive divergence (Chapman *et al.*, 2013). However, in sequence comparisons, alleles that showed high interspecific divergence (ca. 200) were just a small fraction of the loci that were analysed (ca. 18,000) suggesting that species boundaries due to morphological differences and ecological adaptations might be maintained by just a small portion of the genome (Chapman *et al.*, 2013), a pattern consistent with recently diverged species as part of the ‘genic view’ of speciation (Wu, 2001). Future work should exploit the ability to clone and transplant individuals across environments. In this way, field experiments and the power of recent advances in NGS transcriptome analysis can be combined to explore the adaptive potential of changes in gene expression among *Senecio* taxa (e.g., Alexander-Webber *et al.*, 2016), alongside those associated with the generation of hybrids, as well as the loci showing plastic responses to environmental variation within species.

## **7. Integrating studies of genotype, phenotype and fitness**

### **7.1 The challenge of integrating genotype, phenotype and fitness information in the field.**

Quantifying genetic variation in fitness, and its relationship with quantitative traits, allows the responses of groups of traits to natural selection to be predicted. This information is central to understanding how populations and ecological communities will respond to the reductions in fitness predicted by ongoing global change, as well to explore how selection varies in strength and form among populations, species and in different ecological contexts (Shaw & Etterson, 2012). Knowledge on genetic and ecological correlations among traits will also help us to understand what drives populations to specialise, forming new species with



158 narrow ecological tolerances, rather than creating ecologically widespread species that show local adaptation  
159 across their range.

160 Quantitative genetic analyses of traits and fitness across ecological context demand large sample sizes and  
161 large experiments that integrate genetic, phenotypic and fitness information, ideally across a range of natural  
162 environments. To estimate genetic variation in fitness, selection should be measured on individuals of known  
163 relatedness (pedigrees or crossing designs) by quantifying phenotypic traits and fitness under field  
164 conditions, and estimating the additive genetic covariance between traits and fitness (Morrissey *et al.*, 2012;  
165 Pujol *et al.*, 2018; Shaw, 2019). The combination of appealing experimental attributes (short generation  
166 times, recent genetic developments and the ease to which they can be propagated) and the range of ecologies  
167 in which the fitness of phenotypes can be assessed, mean that such integrative experiments are more tractable  
168 (and powerful) in *Senecio* than in many other model systems, especially when incorporated with five  
169 powerful experimental approaches outlined below.

170 *1) Full-sibling families:* Controlled crosses can be carried out easily in most *Senecio* species. In most cases  
171 they can also produce sufficient seed/genetic resources to conduct the large-scale field experiments with  
172 sufficient power to quantify evolutionary parameters (e.g., Walter *et al.*, 2019). Crossing designs containing  
173 both full and half-sibling families (i.e., individuals sharing only a father) allow partitioning of the phenotypic  
174 information to estimate the additive genetic covariation between traits, or between traits and fitness (Lynch &  
175 Walsh, 1998). The benefit of using families within a breeding design is two-fold. Firstly, transplanting  
176 multiple individuals (siblings) from the same family in different environments allows them to be tested in  
177 different environments to identify how fitness varies across environments. Secondly, with sufficient seed, the  
178 same families can be propagated in separate glasshouse and field experiments. In this scenario, we can  
179 connect the mean of each family across separate experiments, making it possible to estimate the mean  
180 phenotype in the glasshouse, and the mean fitness in the field. Connecting phenotype with fitness using  
181 family means is logistically simpler than measuring phenotypes in the field where selection may have  
182 removed variation, and plasticity may mask genotypic effects. In addition, because phenotypes are measured  
183 under controlled conditions, the number of phenotypes that can be measured can be maximised, and each  
184 trait measured more precisely by minimising environmental variation. Traits that are difficult to measure in  
185 the field, such as physiological traits, can then be included more easily in largescale experiments. Fieldwork  
186 can then focus on quantifying different and more comprehensive measures of fitness to characterise how  
187 these performance measures change with ecological, geographic or temporal variation.

188 *2) Clones:* Ideally, the phenotype and fitness of a given individual within a pedigreed population (or crossing  
189 design) should be assessed across different environments, controlling for genetic variation. However, this is

only possible where genotypes can be replicated (i.e. cloned) and transplanted into multiple environments, an approach that assesses the reaction of the same genotype to environmental variation. Such an approach can be implemented in very few other model systems (e.g., *Mimulus*). However, many *Senecio* species produce roots readily from cuttings, allowing the use of clones of known genotypes in field and molecular experiments, at sufficiently high replication to allow useful estimates of genetic variation in fitness to be made under field conditions, as well as exploration of the genomic bases of such variation.

3) *Artificial hybridization*: Detecting ongoing natural selection, and genetic variation in fitness, is difficult if populations are already close to their optimal phenotype and stabilizing selection is strong, or genetic variation low (Walsh & Lynch, 2018). The benefit of using *Senecio* as a model system to overcome such difficulties is two-fold. Firstly, many *Senecio* species show evidence of recent rapid adaptive radiation, providing the opportunity to characterise the strength and direction of selection during the early stages of adaptive diversification using transplant experiments (Walter *et al.*, 2016; Richards *et al.*, 2019). Secondly, *Senecio* species often display high levels of intraspecific variation, often with a well-defined structure among populations (i.e., ecotypes). In many cases, species also produce viable hybrids (unlike highly divergent taxa), allowing their use in multi-generation artificial hybridisation experiments (e.g., Richards *et al.*, 2019; Walter *et al.*, 2019; Wilkinson *et al.*, 2019). These crosses can be used to recreate genetic and phenotypic variation that is currently absent in natural populations and may reflect earlier stages of divergence among sister species. If transplanted into the field, the fitness of these alleles and genotypes can be used to explore the ecological consequences of such variation, as has also been conducted with great success to study hybrid speciation in *Helianthus* sunflowers (Lexer *et al.*, 2003a; Lexer *et al.*, 2003b).

4) *Molecular methods for relating genotype to phenotype*: The transcriptome is a critical intermediary between genetic and phenotypic variation. Appropriately designed experiments can answer fundamental questions about the genomic architecture underlying phenotypic responses to the environment. *Senecio* can be used in experiments that propagate large numbers of clones of a single genotype, which are then transplanted across environmental conditions to produce powerful analyses of differential gene expression. Changes in levels of gene expression across the genome can then quantify the effects of the environment, tissue-type, sex, or developmental stage. RNA-seq has been used to study differential expression between divergent species on Mt Etna (Hegarty *et al.*, 2008; Chapman *et al.*, 2013), between polyploids and their diploid progenitors (Alexander-Webber *et al.*, 2016), and also intraspecific variation of specific traits, such as nickel hyper accumulation in *S. coronatus* (Meier *et al.*, 2018). Differential expression analyses tend to yield lists of tens to hundreds of genes that differ in the number of transcripts produced between conditions. However, functional annotation of genes provides a means of deriving meaningful results from such lists. For example, Meier and colleagues found significant enrichment for transporter proteins, and qRT-PCR

confirmed elevated expression of two known nickel transporters (IRT1 and IREG2), which were associated with intra-specific variation in nickel accumulation.

5) *Genetic transformation and gene editing*: More explicit tests of function are possible in *Senecio* using *Agrobacterium*-mediated transformation (Kim *et al.*, 2008; Garcês *et al.*, 2016). By combining gene expression and functional analyses with phylogenetic analyses, Chen *et al.* (2018) showed that ray florets in *S. vulgaris* are regulated by three CYC2-like paralogs. The phylogenetic history of this gene family across *Senecio*, and Asteraceae more broadly, can show how gene duplication and introgression can give rise to greater floral complexity. Finally, the development of the CRISPR/Cas9 gene editing technology for non-model organisms (Shan *et al.*, 2018) will allow fast targeted mutagenesis in *Senecio* to test the function of loci identified from comparative genomic studies and field experiments. Such technologies will provide an excellent opportunity to relate gene function to evolutionary ecology and the gain and loss of key traits across such a diverse plant genus.

## 7.2 Exploring the evolution of environmental sensitivity

The ability of a single genotype to produce different phenotypes in response to environmental variation (phenotypic plasticity) allows complex organisms to survive in a variable world. Across environments where this plasticity is adaptive, it reduces the exposure of alleles to selection (Ghalambor *et al.*, 2007), acting against the formation of clines (local adaptation) at many loci. Such responses of genotypes to environmental variation in time and space determines the distribution of populations, their range limits and how they will respond to future environmental disturbances (Chevin *et al.*, 2010). The relatively recent radiation of *Senecio* (10.7My), their occupation of contrasting environments, and their ability to hybridise makes *Senecio* a powerful system to study how environmental sensitivity evolves in different environmental contexts.

Local adaptation in many plant systems (Hereford, 2009), including numerous *Senecio* species (e.g., Walter *et al.*, 2016), confirms that plasticity within a given species can only be adaptive across the limited range of environments that typically define their ecological niche (DeWitt *et al.*, 1998). Studies have also identified heritable variation in plasticity for several important traits (Nussey *et al.*, 2005), suggesting that genetic variation for plasticity persists that may permit future evolution of different forms of environmental sensitivity. However, such studies are rare in plants, and require the development of model systems that can integrate the approaches described above (especially clonal propagation).

In other systems, reciprocal transplants of clones, seeds, or seedlings within and outside existing habitats has been used to explore genetic variation in phenological traits at range margins (e.g., Sheth & Angert, 2016). Ongoing *Senecio* research on Mount Etna is complementing these studies by testing how adaptation to

contrasting environments affects their capacity for phenotypic plasticity when exposed to environmental variation (G. Walter *et al.*, unpublished). Therefore, *Senecio* can be used to understand how local adaptation (i.e. specialisation) may reduce the forms and magnitude of plasticity possible within a species, with concomitant consequences for fitness and resilience to future environmental change.

### 7.3 Understanding trade-offs in life history and among environments

Understanding life history trade-offs is crucial for understanding how organisms acquire and utilise resources throughout their lifetime, and why they form species with narrow niches over longer timescales (Futuyma & Moreno, 1988; Agrawal *et al.*, 2010). Field transplants of ecotypes of *S. lautus* have shown developmental mismatches with the environment that results in mortality during seedling establishment. Such mortality reflects fitness trade-offs between contrasting environments and life stages, suggesting specialisation for environment-specific resource allocation strategies (Walter *et al.*, 2018b). How variation in resource allocation across environments is mediated by biotic challenges (e.g., immunity or resistance to fungi) remains a key issue in understanding ecological range margins.

### 7.4 Genetic constraints to adaptation

Trait correlations bias the distribution of genetic variation towards certain multivariate phenotypes, at the expense of other phenotypes. If genetic variance is orientated away from the direction of natural selection, there will be little genetic variance for fitness, and the rate of adaptation will be slowed (Lande, 1979; Walsh & Blows, 2009). For much of the 20<sup>th</sup> century, genetic correlations were thought to be stable, making it difficult to see how rapid evolution in novel directions could occur (Arnold *et al.*, 2008). However, recent comparative evidence has shown that genetic correlations can evolve rapidly (Doroszuk *et al.*, 2008; Eroukhmanoff & Svensson, 2011), and experiments with *Senecio* suggest that changes in genetic correlations could overcome constraints to adaptation, allowing rapid ecological diversification (Walter *et al.*, 2018a). By integrating the approaches outlined above, experiments can capitalise on the tractability of *Senecio* to identify how the environment affects genetic variation in traits and fitness, identify the conditions that alter genetic correlations, and assess how and when the evolution of trait correlations leads to rapid evolutionary responses to novel environments.

## 8. Conspectus

*Senecio* is a powerful, but as yet underutilised model system that has a unique potential to answer many fundamental questions in ecology and evolutionary biology. Most recently, four studies using Australian species of *Senecio* in field and glasshouse experiments combined artificial hybridization with molecular and quantitative genetics to identify mechanisms underlying adaptive divergence leading to speciation (Roda *et*

al., 2017; Melo *et al.*, 2019; Walter *et al.*, 2019; Wilkinson *et al.*, 2019). In Europe, the development of genetic and genomic resources for *S. squalidus* and its close relatives has facilitated comparative genomics of species adapted to contrasting environments, and the identification of candidate genes for local adaptation (Hegarty *et al.*, 2008; Chapman *et al.*, 2013; B. Nevado *et al.* unpublished). Meanwhile, detailed genomic and ecological analyses of closely related but ecologically divergent species in Sicily have revealed how selection acts across different genomic and ecological backgrounds (Brennan *et al.*, 2009), the role of gene flow and geological processes in promoting adaptive divergence (Chapman *et al.*, 2013; Osborne *et al.*, 2016), and the way that gene expression variation may be involved in these processes. Alongside these studies, focused ecological analyses can use *Senecio* to better understand the biotic and abiotic factors that limit species distributions across taxa and biogeography. While ongoing next-generation sequencing can further develop *Senecio* as a system for studying genome evolution within a phylogenetic context. Quantitative genetic research can develop *Senecio* by taking clones of individuals from breeding designs, and transplanting them across environments to identify genetic variation that can promote adaptive evolutionary responses to environmental change. Combining transcriptomic analyses with such experiments can then identify the genetic basis of plasticity. Such experiments using the *Senecio* model will ultimately yield an integrative ‘triangle’ linking genotype, phenotype and fitness.

## 301 **Acknowledgements**

302 We would like to thank Lynda Delph, Jeannine Cavender-Bares and three anonymous reviewers for valuable  
303 comments on earlier versions of this paper. We apologise that we cannot name all contributors, but we are  
304 especially thankful to those scientists who have contributed to advancing *Senecio* as a system for research in  
305 Australasia: L. Ambrose, B. Britain, D. Bernal, M. James, H. Liu, M. Melo, H. North, P. Pelser, P. Prentis, T.  
306 Richards, F. Roda and M. Wilkinson, and in Europe: P. Ashton, M. Coleman, H-P. Comes, S. Cozzolino, P.  
307 Crisp, A. Cristaudo, D. Forbes, P. Gibbs, A. Gillies, S. Harris, M. Hegarty, R. Ingram, J. Irwin, J. James, J.  
308 Kadereit, M. Kim, G. Liu, A. Lowe, D. Tabah, S. McInnis, O. Osborne, G. Barker, A. Allen and D. Marshall.  
309 We also acknowledge the national funding bodies that have supported this work, including NERC, The  
310 Genetics Society, BBSRC and the Leverhulme Trust in the UK, NSF in the USA, and ARC in Australia.

## 312 **References**

- 313 **Abbott R, Albach D, Ansell S, Arntzen JW, Baird SJE, Bierne N, Boughman JW, Brelsford A, Buerkle**  
314 **CA, Buggs R, et al. 2013.** Hybridization and speciation. *Journal of Evolutionary Biology* **26**: 229-  
315 246.
- 316 **Abbott RJ, Ashton PA, Forbes DG. 1992.** Introgressive origin of the radiate groundsel, *Senecio vulgaris* L.  
317 var. *hibernicus* Syme - *Aat-3* evidence. *Heredity* **68**: 425-435.

- 18 **Abbott RJ, Brennan AC, James JK, Forbes DG, Hegarty MJ, Hiscock SJ. 2009.** Recent hybrid origin  
19 and invasion of the British Isles by a self-incompatible species, Oxford ragwort (*Senecio squalidus*  
20 L., Asteraceae). *Biological Invasions* **11**: 1145-1158.
- 21 **Abbott RJ, Bretagnolle FC, Thebaud C. 1998.** Evolution of a polymorphism for outcrossing rate in  
22 *Senecio vulgaris*: Influence of germination behavior. *Evolution* **52**: 1593-1601.
- 23 **Abbott RJ, Comes HP. 2007.** Blowin' in the wind - the transition from ecotype to species. *New Phytologist*  
24 **175**: 197-200.
- 25 **Abbott RJ, Comes HP, Goodwin ZA, Brennan AC. 2018.** Hybridisation and detection of a hybrid zone  
26 between mesic and desert ragworts (*Senecio*) across an aridity gradient in the eastern Mediterranean.  
27 *Plant Ecology & Diversity* **11**: 267-281.
- 28 **Abbott RJ, Irwin JA. 1988.** Pollinator movements and the polymorphism for outcrossing rate at the ray  
29 floret locus in groundsel, *Senecio vulgaris* L. *Heredity* **60**: 295-298.
- 30 **Abbott RJ, Lowe AJ. 2004.** Origins, establishment and evolution of new polyploid species: *Senecio*  
31 *cambrensis* and *S. eboracensis* in the British Isles. *Biological Journal of the Linnean Society* **82**: 467-  
32 474.
- 33 **Abbott RJ, Rieseberg LH 2012.** Hybrid speciation. *eLS*. Chichester: John Wiley & Sons, Ltd.
- 34 **Abbott RJ, Schmitt J. 1985.** Effect of environment on percentage female ray florets per capitulum and  
35 outcrossing potential in a self-compatible composite (*Senecio vulgaris* L. var. *hibernicus* Syme). *New*  
36 *Phytologist* **101**: 219-229.
- 37 **Agrawal AA, Conner JK, Rasmann S 2010.** Tradeoffs and adaptive negative correlations in evolutionary  
38 ecology. In: Bell M, Eanes W, Futuyma D, Levinton J eds. *Evolution After Darwin: The First 150*  
39 *Years*. Sunderland, MA: Sinauer Associates, pp 243-268.
- 40 **Alexander JC. 1979.** The Mediterranean species of *Senecio* sections *Senecio* and *Delphinifolius*. *Notes*  
41 *Royal Botanic Garden Edinburgh* **37**: 387-428.
- 42 **Alexander-Webber D, Abbott RJ, Chapman MA. 2016.** Morphological convergence between an  
43 allopolyploid and one of its parental species correlates with biased gene expression and DNA loss.  
44 *Journal of Heredity* **107**: 445-454.
- 45 **Allan E, Pannell JR. 2009.** Rapid divergence in physiological and life-history traits between northern and  
46 southern populations of the British introduced neo-species, *Senecio squalidus*. *Oikos* **118**: 1053-1061.
- 47 **Allen AM, Hiscock S 2008.** The evolution and phylogeny of self-incompatibility systems in flowering  
48 plants. In: Franklin-Tong VE ed. *Self-incompatibility in Angiosperms*. New York, Berlin: Springer,  
49 pp 73-101.
- 50 **Allen AM, Thorogood CJ, Hegarty MJ, Lexer C, Hiscock SJ. 2011.** Pollen-pistil interactions and self-  
51 incompatibility in the Asteraceae: new insights from studies of *Senecio squalidus* (Oxford ragwort).  
52 *Annals of Botany* **108**: 687-698.
- 53 **Anderson JT, Lee CR, Rushworth CA, Colautti RI, Mitchell-Olds T. 2013.** Genetic trade-offs and  
54 conditional neutrality contribute to local adaptation. *Molecular Ecology* **22**: 699-708.

- 355 **Andersson S. 2001.** Fitness consequences of floral variation in *Senecio jacobaea* (Asteraceae): evidence  
 356 from a segregating hybrid population and a resource manipulation experiment. *Biological Journal of*  
 357 *the Linnean Society* **74**: 17-24.
- 358 **Andersson S, Widén B. 1993.** Pollinator-mediated selection on floral traits in a synthetic population of  
 359 *Senecio integrifolius* (Asteraceae). *Oikos* **66**: 72-79.
- 360 **Arnold SJ, Burger R, Hohenlohe PA, Ajie BC, Jones AG. 2008.** Understanding the evolution and stability  
 361 of the G-Matrix. *Evolution* **62**: 2451-2461.
- 362 **Baker HG. 1967.** Support for Baker's law - as a rule. *Evolution* **21**: 853-856.
- 363 **Barrett SCH. 2013.** The evolution of plant reproductive systems: how often are transitions irreversible?  
 364 *Proceedings of the Royal Society B-Biological Sciences* **280**: 20130913.
- 365 **Bateson A. 1888.** The effect of cross-fertilization on inconspicuous flowers. *Annals of Botany* **1**: 255-261.
- 366 **Billiard S, Castric V, Vekemans X. 2007.** A general model to explore complex dominance patterns in plant  
 367 sporophytic self-incompatibility systems. *Genetics* **175**: 1351-1369.
- 368 **Blackman BK, Michaels SD, Rieseberg LH. 2011.** Connecting the sun to flowering in sunflower  
 369 adaptation. *Molecular Ecology* **20**: 3503-3512.
- 370 **Boyes DC, Nasrallah ME, Vrebalov J, Nasrallah JB. 1997.** The self-incompatibility (S) haplotypes of  
 371 *Brassica* contain highly divergent and rearranged sequences of ancient origin. *Plant Cell* **9**: 237-247.
- 372 **Brennan A, Harris SA, Tabah DA, Hiscock SJ. 2002.** The population genetics of sporophytic self-  
 373 incompatibility in *Senecio squalidus* L. (Asteraceae) I: S allele diversity in a natural population.  
 374 *Heredity* **89**: 430-438.
- 375 **Brennan AC, Barker D, Hiscock SJ, Abbott RJ. 2012.** Molecular genetic and quantitative trait divergence  
 376 associated with recent homoploid hybrid speciation: a study of *Senecio squalidus* (Asteraceae).  
 377 *Heredity* **108**: 87-95.
- 378 **Brennan AC, Bridle JR, Wang AL, Hiscock SJ, Abbott RJ. 2009.** Adaptation and selection in the *Senecio*  
 379 (Asteraceae) hybrid zone on Mount Etna, Sicily. *New Phytologist* **183**: 702-717.
- 380 **Brennan AC, Harris SA, Hiscock SJ. 2003.** The population genetics of sporophytic self-incompatibility in  
 381 *Senecio squalidus* L. (Asteraceae): avoidance of mating constraints imposed by low S-allele number.  
 382 *Philosophical Transactions of the Royal Society of London Series B-Biological Sciences* **358**: 1047-  
 383 1050.
- 384 **Brennan AC, Harris SA, Hiscock SJ. 2005.** Modes and rates of selfing and associated inbreeding  
 385 depression in the self-incompatible plant *Senecio squalidus* (Asteraceae): a successful colonizing  
 386 species in the British Isles. *New Phytologist* **168**: 475-486.
- 387 **Brennan AC, Harris SA, Hiscock SJ. 2006.** The population genetics of sporophytic self-incompatibility in  
 388 *Senecio squalidus* L. (Asteraceae): The number, frequency, and dominance interactions of S alleles  
 389 across its British range. *Evolution* **60**: 213-224.

'90 **Brennan AC, Harris SA, Hiscock SJ. 2013.** The population genetics of sporophytic self-incompatibility in  
 '91 three hybridizing *Senecio* (Asteraceae) species with contrasting population histories. *Evolution* **67**:  
 '92 1347-1367.

'93 **Brennan AC, Hiscock SJ. 2010.** Expression and inheritance of sporophytic self-incompatibility in synthetic  
 '94 allohexaploid *Senecio cambrensis* (Asteraceae). *New Phytologist* **186**: 251-261.

'95 **Brennan AC, Hiscock SJ, Abbott RJ. 2019.** Completing the hybridization triangle: The inheritance of  
 '96 genetic incompatibilities during homoploid hybrid speciation in ragworts (*Senecio*). *AoB Plants* **11**:  
 '97 ply078.

'98 **Brennan AC, Tabah DA, Harris SA, Hiscock SJ. 2011.** Sporophytic self-incompatibility in *Senecio*  
 '99 *squalidus* (Asteraceae): S allele dominance interactions and modifiers of cross-compatibility and  
 '00 selfing rates. *Heredity* **106**: 113-123.

'01 **Castric V, Bechsgaard J, Schierup MH, Vekemans X. 2008.** Repeated adaptive introgression at a gene  
 '02 under multiallelic balancing selection. *PLoS Genetics* **4**: e1000168.

'03 **Chapman MA, Abbott RJ. 2010.** Introgression of fitness genes across a ploidy barrier. *New Phytologist*  
 '04 **186**: 63-71.

'05 **Chapman MA, Forbes DG, Abbott RJ. 2005.** Pollen competition among two species of *Senecio*  
 '06 (Asteraceae) that form a hybrid zone on Mt. Etna, Sicily. *American Journal of Botany* **92**: 730-735.

'07 **Chapman MA, Hiscock SJ, Filatov DA. 2013.** Genomic divergence during speciation driven by adaptation  
 '08 to altitude. *Molecular Biology and Evolution* **30**: 2553-2567.

'09 **Chapman MA, Hiscock SJ, Filatov DA. 2016.** The genomic bases of morphological divergence and  
 '10 reproductive isolation driven by ecological speciation in *Senecio* (Asteraceae). *Journal of*  
 '11 *Evolutionary Biology* **29**: 98-113.

'12 **Chen J, Shen CZ, Guo YP, Rao GY. 2018.** Patterning the Asteraceae capitulum: Duplications and  
 '13 differential expression of the flower symmetry *CYC2*-like genes. *Frontiers in Plant Science* **9**: 551.

'14 **Chevin LM, Lande R, Mace GM. 2010.** Adaptation, plasticity, and extinction in a changing environment:  
 '15 towards a predictive theory. *PLoS Biology* **8**: e1000357.

'16 **Comes HP, Abbott RJ. 2001.** Molecular phylogeography, reticulation, and lineage sorting in Mediterranean  
 '17 *Senecio* sect. *Senecio* (Asteraceae). *Evolution* **55**: 1943-1962.

'18 **Comes HP, Coleman M, Abbott RJ. 2017.** Recurrent origin of peripheral, coastal (sub)species in  
 '19 Mediterranean *Senecio* (Asteraceae). *Plant Ecology & Diversity* **10**: 253-271.

'20 **DeWitt TJ, Sih A, Wilson DS. 1998.** Costs and limits of phenotypic plasticity. *Trends in Ecology &*  
 '21 *Evolution* **13**: 77-81.

'22 **Doellman MM, Ragland GJ, Hood GR, Meyers PJ, Egan SP, Powell THQ, Lazorchak P, Glover MM,**  
 '23 **Tait C, Schuler H, et al. 2018.** Genomic differentiation during speciation-with-gene-flow:  
 '24 Comparing geographic and host-related variation in divergent life history adaptation in *Rhagoletis*  
 '25 *pomonella*. *Genes* **9**: 262.



- '26 **Doroszuk A, Wojewodziec MW, Gort G, Kammenga JE. 2008.** Rapid divergence of genetic variance-  
'27 covariance matrix within a natural population. *The American Naturalist* **171**: 291-304.
- '28 **Durand E, Méheust R, Soucaze M, Goubet PM, Gallina S, Poux C, Fobis-Loisy I, Guillon E, Gaude T,**  
'29 **Sarazin A, et al. 2014.** Dominance hierarchy arising from the evolution of a complex small RNA  
'30 regulatory network. *Science* **346**: 1200-1205.
- '31 **Eroukhmanoff F, Svensson EI. 2011.** Evolution and stability of the G-matrix during the colonization of a  
'32 novel environment. *Journal of Evolutionary Biology* **24**: 1363-1373.
- '33 **Ferrer MM, Good-Avila SV. 2007.** Macrophylogenetic analyses of the gain and loss of self-incompatibility  
'34 in the Asteraceae. *New Phytologist* **173**: 401-414.
- '35 **Filatov DA, Osborne OG, Papadopoulos AS. 2016.** Demographic history of speciation in a *Senecio*  
'36 altitudinal hybrid zone on Mt. Etna. *Molecular Ecology* **25**: 2467-2481.
- '37 **Futuyma DJ, Moreno G. 1988.** The evolution of ecological specialization. *Annual Review of Ecology,*  
'38 *Evolution, and Systematics* **19**: 207-233.
- '39 **Gandhi SD, Heesacker AF, Freeman CA, Argyris J, Bradford K, Knapp SJ. 2005.** The self-  
'40 incompatibility locus (S) and quantitative trait loci for self-pollination and seed dormancy in  
'41 sunflower. *Theoretical and Applied Genetics* **111**: 619-629.
- '42 **Garcès HMP, Spencer VMR, Kim M. 2016.** Control of floret symmetry by *RAY3*, *SvDIV1B*, and *SvRAD* in  
'43 the capitulum of *Senecio vulgaris*. *Plant Physiology* **171**: 2055-2068.
- '44 **Ghalambor CK, McKay JK, Carroll SP, Reznick DN. 2007.** Adaptive versus non-adaptive phenotypic  
'45 plasticity and the potential for contemporary adaptation in new environments. *Functional Ecology* **21**:  
'46 394-407.
- '47 **Gillies ACM, Cubas P, Coen E, Abbott RJ 2002.** Making rays in the Asteraceae: genetics and evolution of  
'48 radiate versus discoid flower heads. In: Cronk QCB, Bateman RM, Hawkins JA eds. *Developmental*  
'49 *Genetics and Plant Evolution*. London: Taylor & Francis, pp 233-246.
- '50 **Gonthier L, Blassiau C, Morchen M, Cadalen T, Poiret M, Hendriks T, Quillet MC. 2013.** High-density  
'51 genetic maps for loci involved in nuclear male sterility (*NMS1*) and sporophytic self-incompatibility  
'52 (S-locus) in chicory (*Cichorium intybus* L., Asteraceae). *Theoretical and Applied Genetics* **126**: 2103-  
'53 2121.
- '54 **Goubet PM, Bergés H, Bellec A, Prat E, Helmstetter N, Mangenot S, Gallina S, Holl AC, Fobis-Loisy I,**  
'55 **Vekemans X, et al. 2012.** Contrasted patterns of molecular evolution in dominant and recessive self-  
'56 incompatibility haplotypes in *Arabidopsis*. *PLoS Genetics* **8**: e1002495.
- '57 **Hancock CN, Kondo K, Beecher B, McClure B. 2003.** The S-locus and unilateral incompatibility.  
'58 *Philosophical Transactions of the Royal Society of London Series B-Biological Sciences* **358**: 1133-  
'59 1140.
- '60 **Harland SC. 1954a.** The genus *Senecio* as a subject for cytogenetical investigation. *Proceedings of the*  
'61 *Botanical Society of the British Isles* **1**: 1:256.
- '62 **Harland SC. 1954b.** Two *Senecio* hybrids. *Proceedings of the Botanical Society of the British Isles* **1**: 260-  
'63 261.

- '64 **Harris SA. 2002.** Introduction of Oxford ragwort, *Senecio squalidus* L. (Asteraceae), to the United  
'65 Kingdom. *Watsonia* **24**: 31-43.
- '66 **Harrison RG, Larson EL. 2014.** Hybridization, introgression, and the nature of species boundaries. *Journal*  
'67 *of Heredity* **105**: 795-809.
- '68 **Hegarty M, Coate J, Sherman-Broyles S, Abbott R, Hiscock S, Doyle J. 2013.** Lessons from natural and  
'69 artificial polyploids in higher plants. *Cytogenetic and Genome Research* **140**: 204-225.
- '70 **Hegarty MJ, Abbott RJ, Hiscock SJ 2012.** Allopolyploid speciation in action: The origins and evolution of  
'71 *Senecio cambrensis*. In: Soltis P, Soltis D eds. *Polyploidy and Genome Evolution*. Berlin, Heidelberg:  
'72 Springer, pp 245-270.
- '73 **Hegarty MJ, Barker GL, Brennan AC, Edwards KJ, Abbott RJ, Hiscock SJ. 2008.** Changes to gene  
'74 expression associated with hybrid speciation in plants: further insights from transcriptomic studies in  
'75 *Senecio*. *Philosophical Transactions of the Royal Society B-Biological Sciences* **363**: 3055-3069.
- '76 **Hegarty MJ, Barker GL, Brennan AC, Edwards KJ, Abbott RJ, Hiscock SJ. 2009.** Extreme changes to  
'77 gene expression associated with homoploid hybrid speciation. *Molecular Ecology* **18**: 877-889.
- '78 **Hegarty MJ, Barker GL, Wilson ID, Abbott RJ, Edwards KJ, Hiscock SJ. 2006.** Transcriptome shock  
'79 after interspecific hybridization in *Senecio* is ameliorated by genome duplication. *Current Biology* **16**:  
'80 1652-1659.
- '81 **Hegarty MJ, Jones JM, Wilson ID, Barker GL, Coghill JA, Sanchez-Baracaldo P, Liu GQ, Buggs**  
'82 **RJA, Abbott RJ, Edwards KJ, et al. 2005.** Development of anonymous cDNA microarrays to study  
'83 changes to the *Senecio* floral transcriptome during hybrid speciation. *Molecular Ecology* **14**: 2493-  
'84 2510.
- '85 **Hereford J. 2009.** A quantitative survey of local adaptation and fitness trade-offs. *The American Naturalist*  
'86 **173**: 579-588.
- '87 **Hiscock SJ. 2000a.** Genetic control of self-incompatibility in *Senecio squalidus* L. (Asteraceae): a successful  
'88 colonizing species. *Heredity* **85**: 10-19.
- '89 **Hiscock SJ. 2000b.** Self-incompatibility in *Senecio squalidus* L (Asteraceae). *Annals of Botany* **85**: 181-190.
- '90 **Irwin JA, Abbott RJ. 1992.** Morphometric and isozyme evidence for the hybrid origin of a new tetraploid  
'91 radiate groundsel in York, England. *Heredity* **69**: 431-439.
- '92 **Iwano M, Takayama S. 2012.** Self/non-self discrimination in angiosperm self-incompatibility. *Current*  
'93 *Opinion in Plant Biology* **15**: 78-83.
- '94 **James JK, Abbott RJ. 2005.** Recent, allopatric, homoploid hybrid speciation: The origin of *Senecio*  
'95 *squalidus* (Asteraceae) in the British Isles from a hybrid zone on Mount Etna, Sicily. *Evolution* **59**:  
'96 2533-2547.
- '97 **Kadereit JW, Uribe-Convers S, Westberg E, Comes HP. 2006.** Reciprocal hybridization at different times  
'98 between *Senecio flavus* and *Senecio glaucus* gave rise to two polyploid species in north Africa and  
'99 south-west Asia. *New Phytologist* **169**: 431-441.

- 300 **Kandziora M, Kadereit JW, Gehrke B. 2017.** Dual colonization of the Palaearctic from different regions in  
 301 the Afrotropics by *Senecio*. *Journal of Biogeography* **44**: 147-157.
- 302 **Kim M, Cui ML, Cubas P, Gillies A, Lee K, Chapman MA, Abbott RJ, Coen E. 2008.** Regulatory genes  
 303 control a key morphological and ecological trait transferred between species. *Science* **322**: 1116-  
 304 1119.
- 305 **Lafuma L, Maurice S. 2007.** Increase in mate availability without loss of self-incompatibility in the  
 306 invasive species *Senecio inaequidens* (Asteraceae). *Oikos* **116**: 201-208.
- 307 **Lai Z, Gross BL, Zou Y, Andrews J, Rieseberg LH. 2006.** Microarray analysis reveals differential gene  
 308 expression in hybrid sunflower species. *Molecular Ecology* **15**: 1213-1227.
- 309 **Lande R. 1979.** Quantitative genetic analysis of multivariate evolution, applied to brain:body size allometry.  
 310 *Evolution* **33**: 402-416.
- 311 **Leitch IJ, Johnston E, Pellicer J, Hidalgo O, Bennett MD. 2019.** Angiosperm DNA C-values database.  
 312 Kew gardens. <https://cvalues.science.kew.org/> [accessed 25 November 2019].
- 313 **Levin DA. 1996.** The evolutionary significance of pseudo-self-fertility. *American Naturalist* **148**: 321-332.
- 314 **Lewis D 1994.** Gametophytic-sporophytic incompatibility. In: Williams EG, Clarke AE, Knox RB eds.  
 315 *Genetic control of self-incompatibility and reproductive development in flowering plants*. Dordrecht:  
 316 Springer, pp 88-101.
- 317 **Lexer C, Randell RA, Rieseberg LH. 2003a.** Experimental hybridization as a tool for studying selection in  
 318 the wild. *Ecology* **84**: 1688-1699.
- 319 **Lexer C, Welch ME, Raymond O, Rieseberg LH. 2003b.** The origin of ecological divergence in  
 320 *Helianthus paradoxus* (Asteraceae): Selection on transgressive characters in a novel hybrid habitat.  
 321 *Evolution* **57**: 1989-2000.
- 322 **Liew CS, Memory AE, Ortiz-Barrientos D, de Lange PJ, Pelser PB. 2018.** The delimitation and  
 323 evolutionary history of the Australasian Lautusoid group of *Senecio* (Asteraceae: Senecioneae).  
 324 *Taxon* **67**: 130-148.
- 325 **Llaurens V, Billiard S, Castric V, Vekemans X. 2009.** Evolution of dominance in sporophytic self-  
 326 incompatibility systems: I. Genetic load and coevolution of levels of dominance in pollen and pistil.  
 327 *Evolution* **63**: 2427-2437.
- 328 **Lowe AJ, Abbott RJ. 2003.** A new British species, *Senecio eboracensis* (Asteraceae), another hybrid  
 329 derivative of *S. vulgaris* L. and *S. squalidus* L. *Watsonia* **24**: 375-388.
- 330 **Lowry DB. 2012.** Ecotypes and the controversy over stages in the formation of new species. *Biological*  
 331 *Journal of the Linnean Society* **106**: 241-257.
- 332 **Lowry DB, Modliszewski JL, Wright KM, Wu CA, Willis JH. 2008.** The strength and genetic basis of  
 333 reproductive isolating barriers in flowering plants. *Philosophical Transactions of the Royal Society of*  
 334 *London Series B-Biological Sciences* **363**: 3009-3021.
- 335 **Lowry DB, Willis JH. 2010.** A widespread chromosomal inversion polymorphism contributes to a major  
 336 life-history transition, local adaptation, and reproductive isolation. *PLoS Biology* **8**: 1-14.

- 337 **Lynch M, Walsh B. 1998.** *Genetics and analysis of quantitative traits*. Sunderland: Sinauer Associates, Inc.
- 338 **Marshall DF, Abbott RJ. 1982.** Polymorphism for outcrossing frequency at the ray floret locus in *Senecio*  
339 *vulgaris* L. I. Evidence. *Heredity* **48**: 227-235.
- 340 **Marshall DF, Abbott RJ. 1984.** Polymorphism for outcrossing frequency at the ray floret locus in *Senecio*  
341 *vulgaris* L. III. Causes. *Heredity* **53**: 145-149.
- 342 **McInnis SM, Emery DC, Porter R, Desikan R, Hancock JT, Hiscock SJ. 2006.** The role of stigma  
343 peroxidases in flowering plants: insights from further characterization of a stigma-specific peroxidase  
344 (SSP) from *Senecio squalidus* (Asteraceae). *Journal of Experimental Botany* **57**: 1835-1846.
- 345 **Meier SK, Adams N, Wolf M, Balkwill K, Muasya AM, Gehring CA, Bishop JM, Ingle RA. 2018.**  
346 Comparative RNA-seq analysis of nickel hyperaccumulating and non-accumulating populations of  
347 *Senecio coronatus* (Asteraceae). *The Plant Journal* **95**: 1023-1038.
- 348 **Melo MC, Grealy A, Brittain B, Walter GM, Ortiz-Barrientos D. 2014.** Strong extrinsic reproductive  
349 isolation between parapatric populations of an Australian groundsel. *New Phytologist* **203**: 323-334.
- 350 **Melo MC, James ME, Roda F, Bernal-Franco D, Wilkinson MJ, Liu H, Walter GM, Ortiz-Barrientos**  
351 **D. 2019.** Evidence for mutation-order speciation in Australian wildflower. *bioRxiv*. DOI:  
352 <https://doi.org/10.1101/692673>.
- 353 **Morrissey MB, Parker DJ, Korsten P, Pemberton JM, Kruuk LE, Wilson AJ. 2012.** The prediction of  
354 adaptive evolution: empirical application of the secondary theorem of selection and comparison to the  
355 breeder's equation. *Evolution* **66**: 2399-2410.
- 356 **Nussey DH, Postma E, Gienapp P, Visser ME. 2005.** Selection on heritable phenotypic plasticity in a wild  
357 bird population. *Science* **310**: 304-306.
- 358 **Osborne OG, Batstone TE, Hiscock SJ, Filatov DA. 2013.** Rapid speciation with gene flow following the  
359 formation of Mt. Etna. *Genome Biology and Evolution* **5**: 1704-1715.
- 360 **Osborne OG, Chapman MA, Nevado B, Filatov DA. 2016.** Maintenance of species boundaries despite  
361 ongoing gene flow in ragworts. *Genome Biology & Evolution* **8**: 1038-1047.
- 362 **Pannell JR, Barrett SCH. 1998.** Baker's law revisited: Reproductive assurance in a metapopulation.  
363 *Evolution* **52**: 657-668.
- 364 **Pelser PB, Abbott RJ, Comes HP, Milton JJ, Moller M, Looseley ME, Cron GV, Barcelona JF,**  
365 **Kennedy AH, Watson LE, et al. 2012.** The genetic ghost of an invasion past: Colonization and  
366 extinction revealed by historical hybridization in *Senecio*. *Molecular Ecology* **21**: 369-387.
- 367 **Prentis PJ, White EM, Radford IJ, Lowe AJ, Clarke AR. 2007.** Can hybridization cause local extinction:  
368 A case for demographic swamping of the Australian native *Senecio pinnatifolius* by the invasive  
369 *Senecio madagascariensis*? *New Phytologist* **176**: 902-912.
- 370 **Pujol B, Blanchet S, Charmantier A, Danchin E, Facon B, Marrot P, Roux F, Scotti I, Teplitsky C,**  
371 **Thomson CE, et al. 2018.** The missing response to selection in the wild. *Trends in Ecology &*  
372 *Evolution* **33**: 337-346.

- Radford IJ, Cousens RD, Michael PW. 2004. Morphological and genetic variation in the *Senecio pinnatifolius* complex: are variants worthy of taxonomic recognition? *Australian Systematic Botany* 17: 29-48.
- Richards TJ, Ortiz-Barrientos D. 2016. Immigrant inviability produces a strong barrier to gene flow between parapatric ecotypes of *Senecio laetus*. *Evolution* 70: 1239-1248.
- Richards TJ, Ortiz-Barrientos D, McGuigan K. 2019. Natural selection drives leaf divergence in experimental populations of *Senecio laetus* under natural conditions. *Ecology and Evolution* 9: 6959–6967.
- Richards TJ, Walter GM, McGuigan K, Ortiz-Barrientos D. 2016. Divergent natural selection drives the evolution of reproductive isolation in an Australian wildflower. *Evolution* 70: 1993-2003.
- Rieseberg LH, Raymond O, Rosenthal DM, Lai Z, Livingstone K, Nakazato T, Durphy JL, Schwarzbach AE, Donovan LA, Lexer C. 2003. Major ecological transitions in wild sunflowers facilitated by hybridization. *Science* 301: 1211-1216.
- Roda F, Ambrose L, Walter GM, Liu HLL, Schaul A, Lowe A, Pelsner PB, Prentis P, Rieseberg LH, Ortiz-Barrientos D. 2013. Genomic evidence for the parallel evolution of coastal forms in the *Senecio laetus* complex. *Molecular Ecology* 22: 2941-2952.
- Roda F, Walter GM, Nipper R, Ortiz-Barrientos D. 2017. Genomic clustering of adaptive loci during parallel evolution of an Australian wildflower. *Molecular Ecology* 26: 3687-3699.
- Ross RI. 2010. *Local adaptation and adaptive divergence in a hybrid species complex in Senecio*. DPhil, University of Oxford, UK.
- Roux C, Pauwels M, Ruggiero MV, Charlesworth D, Castric V, Vekemans X. 2013. Recent and ancient signature of balancing selection around the S-locus in *Arabidopsis halleri* and *A. lyrata*. *Molecular Biology and Evolution* 30: 435-447.
- Rundle HD, Nosil P. 2005. Ecological speciation. *Ecology Letters* 8: 336-352.
- Safran RJ, Scordato ES, Wilkins MR, Hubbard JK, Jenkins BR, Albrecht T, Flaxman SM, Karaardic H, Vortman Y, Lotem A, et al. 2016. Genome-wide differentiation in closely related populations: the roles of selection and geographic isolation. *Molecular Ecology* 25: 3865-3883.
- Schierup MH, Vekemans X. 2008. Genomic consequences of selection on self-incompatibility genes. *Current Opinion in Plant Biology* 11: 116-122.
- Schilling MP, Mullen SP, Kronforst M, Safran RJ, Nosil P, Feder JL, Gompert Z, Flaxman SM. 2018. Transitions from single- to multi-Locus processes during speciation with gene flow. *Genes* 9: 274.
- Shan SC, Mavrodiev EV, Li RQ, Zhang ZZ, Hauser BA, Soltis PS, Soltis DE, Yang B. 2018. Application of CRISPR/Cas9 to *Tragopogon* (Asteraceae), an evolutionary model for the study of polyploidy. *Molecular Ecology Resources* 18: 1427-1443.
- Shaw RG. 2019. From the past to the future: Considering the value and limits of evolutionary prediction. *The American Naturalist* 193: 1-10.

- Shaw RG, Etterson JR. 2012. Rapid climate change and the rate of adaptation: insight from experimental quantitative genetics. *New Phytologist* **195**: 752-765.
- Sheth SN, Angert AL. 2016. Artificial selection reveals high genetic variation in phenology at the trailing edge of a species range. *The American Naturalist* **187**: 182-193.
- Shiba H, Kenmochi M, Sugihara M, Iwano M, Kawasaki S, Suzuki G, Watanabe M, Isogai A, Takayama S. 2003. Genomic organization of the S-locus region of *Brassica*. *Bioscience Biotechnology and Biochemistry* **67**: 622-626.
- Stankowski S, Chase MA, Fuiten AM, Rodrigues MF, Ralph PL, Streisfeld MA. 2019. Widespread selection and gene flow shape the genomic landscape during a radiation of monkeyflowers. *Plos Biology* **17**: e3000391.
- Stephenson AG, Good SV, Vogler DW. 2000. Interrelationships among inbreeding depression, plasticity in the self-incompatibility system, and the breeding system of *Campanula rapunculoides* L- (Campanulaceae). *Annals of Botany* **85**: 211-219.
- Stevison LS, Sefick S, Rushton C, Graze RM. 2017. Recombination rate plasticity: revealing mechanisms by design. *Philosophical Transactions of the Royal Society B-Biological Sciences* **372**: 20160459.
- Tabah DA. 2004. *Studies of self-incompatibility in Senecio squalidus* L. (Asteraceae). . PhD Dissertation, University of Bristol, UK.
- Tabah DA, McInnis SM, Hiscock SJ. 2004. Members of the S-receptor kinase multigene family in *Senecio squalidus* L. (Asteraceae), a species with sporophytic self-incompatibility. *Sexual Plant Reproduction* **17**: 131-140.
- Thomas CD. 2015. Rapid acceleration of plant speciation during the Anthropocene. *Trends in Ecology & Evolution* **30**: 448-455.
- Trow AH. 1912. On the inheritance of certain characters in the common groundsel - *Senecio vulgaris*, Linn - And its segregates. *Journal of Genetics* **2**: 239-276.
- Trow AH. 1916. On the number of nodes and their distribution along the main axis in *Senecio vulgaris* and its segregates. *Journal of Genetics* **6**: 1-63.
- Vallejo-Marin M, Hiscock SJ. 2016. Hybridization and hybrid speciation under global change. *New Phytologist* **211**: 1170-1187.
- VanWallendael A, Soltani A, Emery NC, Peixoto MM, Olsen J, Lowry DB. 2019. A molecular view of plant local adaptation: Incorporating stress-response networks. *Annual Review of Plant Biology* **70**: 559-583.
- Vekemans X, Schierup MH, Christiansen FB. 1998. Mate availability and fecundity selection in multi-allelic self-incompatibility systems in plants. *Evolution* **52**: 19-29.
- Walsh B, Blows MW. 2009. Abundant genetic variation plus strong selection = multivariate genetic constraints: A geometric view of adaptation. *Annual Review of Ecology, Evolution, and Systematics* **40**: 41-59.
- Walsh B, Lynch M. 2018. *Evolution and selection of quantitative traits*. Oxford: Oxford University Press.

- 146 **Walter GM, Aguirre JD, Blows MW, Ortiz-Barrientos D. 2018a.** Evolution of genetic variance during  
 147 adaptive radiation. *The American Naturalist* **191**: E108–E128.
- 148 **Walter GM, Aguirre JD, Wilkinson MJ, Richards TJ, Blows MW, Ortiz-Barrientos D. 2019.** Local  
 149 adaptation and hybrid failure share a common genetic basis. *bioRxiv*. DOI:  
 150 <https://doi.org/10.1101/520809>.
- 151 **Walter GM, Wilkinson MJ, Aguirre JD, Blows MW, Ortiz-Barrientos D. 2018b.** Environmentally  
 152 induced development costs underlie fitness tradeoffs. *Ecology* **99**: 1391–1401.
- 153 **Walter GM, Wilkinson MJ, James ME, Richards TJ, Aguirre JD, Ortiz-Barrientos D. 2016.**  
 154 Diversification across a heterogeneous landscape. *Evolution* **70**: 1979–1992.
- 155 **West-Eberhard MJ. 2005.** Developmental plasticity and the origin of species differences. *Proceedings of*  
 156 *the National Academy of Sciences of the United States of America* **102 Suppl 1**: 6543–6549.
- 157 **Wilkinson MJ, Roda F, Walter GM, James ME, Nipper R, Walsh J, Allen SL, Bernal DM, North HL,**  
 158 **Beveridge CA, et al. 2019.** Divergence in hormone signalling links local adaptation and hybrid  
 159 failure. *bioRxiv*. DOI: <https://doi.org/10.1101/845354>.
- 160 **Wu CA, Lowry DB, Cooley AM, Wright KM, Lee YW, Willis JH. 2008.** *Mimulus* is an emerging model  
 161 system for the integration of ecological and genomic studies. *Heredity* **100**: 220–230.
- 162 **Wu CI. 2001.** The genic view of the process of speciation. *Journal of Evolutionary Biology* **14**: 851–865.
- 163
- 164 **Supporting information**
- 165 **Note S1:** Photo credits for Fig. 1 and Fig. 4.
- 166 **Note S2:** Details of the methods to produce Fig. 2.
- 167 **Note S3:** References cited in Table 1.

168     **Tables, Boxes and figures**

169



370 **Table 1.** Use of *Senecio* in genetic, evolutionary, systematic and ecological research. Within each section of the table, studies are  
371 listed in approximate historical sequence. Full references of citations are listed in Note S3. Research on taxa formerly within  
372 *Senecio* but now placed within the genus *Jacobaea* is not cited in the table.

Topic	Taxa	References
<b>Genetics, Cytology, Development</b>		
Heterosis	<i>S. vulgaris</i>	Bateson (1888)
Genetics of traits	<i>S. vulgaris</i>	Trow (1912, 1916ab); Skinner (1952); Basford (1961ab); Comes (1998); Brennan <i>et al.</i> (2016)
	<i>S. lautus</i>	Roda <i>et al.</i> (2017)
	<i>S. squalidus</i>	Ingram & Taylor (1982); Hiscock (2000a); Hiscock <i>et al.</i> (2003); Tabah <i>et al.</i> (2004); McInnis <i>et al.</i> (2005)
Genome size variation	Diverse <i>Senecio</i>	Ornduff <i>et al.</i> (1963); Lawrence (1980, 1985); Lafuma & Maurice (2007); Lopez <i>et al.</i> (2013)
Maternal effects	<i>S. vulgaris</i>	Aarssen & Burton (1990)
Heteroplasmy	<i>S. vulgaris</i>	Frey (1998); Frey <i>et al.</i> (2005)
<b>Evolution</b>		
Hybridization & introgression	<i>S. vulgaris</i> x <i>S. squalidus</i>	Harland (1954a); Ingram (1977, 1978); Ingram <i>et al.</i> (1980); Marshall & Abbott (1980); Abbott <i>et al.</i> (1992); Kim <i>et al.</i> (2008); Lowe & Abbott (2015)
	<i>S. vulgaris</i> x <i>S. viscosus</i>	Gibbs (1971)
	<i>S. viscosus</i> x <i>S. squalidus</i>	Crisp & Jones (1978)
	<i>S. vulgaris</i> x <i>S. vernalis</i>	Comes & Kadereit (1990); Comes (1994a)
	<i>S. pinnatifolius</i> x <i>S. madagascariensis</i>	Prentis <i>et al.</i> (2007)
Hybrid speciation – Allopolyploidy	<i>S. cambrensis</i>	Harland (1954b); Rosser (1955); Weir & Ingram (1980); Ingram & Noltie (1984, 1986, 1987, 1995); Ashton & Abbott (1992); Lowe & Abbott (1996); Abbott & Forbes (2002); Hegarty <i>et al.</i> (2006, 2008, 2011, 2012)
	<i>S. mohavensis</i>	Coleman <i>et al.</i> (2001); Kadereit <i>et al.</i> (2006); Alexander-Webber <i>et al.</i> (2016)
	<i>S. hoggariensis</i>	Kadereit <i>et al.</i> (2006)
Hybrid speciation – Homoploidy	<i>S. eboracensis</i>	Irwin & Abbott (1992); Lowe & Abbott (2000, 2003, 2004)
	<i>S. squalidus</i>	James & Abbott (2005); Brennan <i>et al.</i> (2012, 2019); Hegarty <i>et al.</i> (2009)
Hybrid zone structure & maintenance	<i>S. aethnensis</i> x <i>S. chysanthemifolius</i>	James & Abbott (2005); Chapman <i>et al.</i> (2005); Brennan <i>et al.</i> (2009); Ross <i>et al.</i> (2012)
	<i>S. vernalis</i> x <i>S. glaucus</i>	Abbott <i>et al.</i> (2018)
	<i>S. hercynicus</i> x <i>S. ovatus</i>	Raudnitscka <i>et al.</i> (2007); Bog <i>et al.</i> (2017a)
Floral polymorphism	<i>S. vulgaris</i>	Richards (1975); Oxford & Andrews (1977); Marshall & Abbott (1982, 1984ab, 1987); Abbott (1985, 1986); Ross & Abbott (1987); Abbott & Irwin (1988); Oxford <i>et al.</i> (1996); Abbott <i>et al.</i> (1998); Kim <i>et al.</i> (2008); Garcés <i>et al.</i> (2016)
Breeding/Mating system variation	Diverse <i>Senecio</i>	Gibbs <i>et al.</i> (1975)
	<i>S. vulgaris</i>	Hull (1974); Campbell & Abbott (1976); Marshall & Abbott (1982, 1984ab, 1987); Ross & Abbott (1987); Warren (1988); Damgaard & Abbott (1995); Irwin <i>et al.</i> (2016); Love <i>et al.</i> (2016)
	<i>S. vernalis</i>	Comes (1994b)
	<i>S. squalidus</i>	Abbott & Forbes (1993); Hiscock (2000ab); Hiscock <i>et al.</i> (2003); Brennan <i>et al.</i> (2002, 2003ab; 2005, 2006); Tabah <i>et al.</i> (2004); McInnis <i>et al.</i> (2006); Allen <i>et al.</i> (2010, 2011)
	<i>S. inaequidens</i>	Lafuma & Maurice (2007)
	<i>S. cambrensis</i>	Brennan & Hiscock (2010)
	<i>S. vulgaris</i>	Crawford (1966); Abbott (1976ab); Briggs (1976, 1978); Ren & Abbott (1991); Theaker & Briggs (1993); Müller-Schärer & Fischer M (2001)
Ecotypic divergence & speciation	<i>S. sylvaticus</i>	Kumler (1969)
	<i>S. viscosus</i>	Akeroyd <i>et al.</i> (1978)
	<i>S. lautus</i>	Radford & Cousens (2004); Melo <i>et al.</i> (2014); Richards <i>et al.</i> (2016, 2019);

		Richards & Ortiz-Barrientos (2016); Walter <i>et al.</i> (2016, 2018ab, 2019)
	<i>S. squalidus</i>	Allan & Pannell (2009)
	<i>S. aethnensis</i> – <i>S. chrysanthemifolius</i>	Chapman <i>et al.</i> (2013); Muir <i>et al.</i> (2013); Osborne <i>et al.</i> (2013); Brennan <i>et al.</i> (2014, 2016)
Parallel divergence & speciation	<i>S. lautus</i> complex	Roda <i>et al.</i> (2013ab, 2017)
	Mediterranean <i>Senecio</i>	Comes <i>et al.</i> (2017)
Herbicide resistance	<i>S. vulgaris</i>	Ryan (1970); Holliday & Putwain (1977); Scott & Putwain (1981); Holt (1988)
<b>Systematics</b>		
Biosystematics	European and African <i>Senecio</i>	Crisp (1972); Alexander (1979); Kadereit (1984ab); Gilmer & Kadereit 1989; Emig & Kadereit (1993); Comes (1995a); Hodálová & Marhold (1996)
Molecular systematics	British <i>Senecio</i>	Harris & Ingram (1992ab)
Phylogeography & speciation	European & African <i>Senecio</i>	Kadereit <i>et al.</i> (1995); Comes (1995); Comes <i>et al.</i> (1997); Comes & Abbott (1998, 1999, 2000, 2001); Purps & Kadereit (1998); Coleman & Abbott (2003); Chapman & Abbott (2005)
Phylogeny	Representatives of genus	Pelser <i>et al.</i> (2002, 2007, 2010, 2012); Coleman <i>et al.</i> (2003); Calvo <i>et al.</i> (2013); Osborne <i>et al.</i> (2016); Kandziora <i>et al.</i> (2016ab)
	South American <i>Senecio</i>	Montesinos-Tubée (2014); Dušková <i>et al.</i> (2017)
	Australasian <i>Senecio</i>	Liew <i>et al.</i> (2018)
<b>Ecology</b>		
Plant invasions	<i>S. squalidus</i>	Kent (1956, 1960); Harris (2002); Abbott <i>et al.</i> (2009); Allan & Pannell (2009)
	<i>S. vernalis</i>	Kadereit (1983)
	<i>S. inaequidens</i>	Ernst (1998); Lafuma <i>et al.</i> (2003); Heger & Böhmer (2005); Bossdorf <i>et al.</i> (2008); Monty & Mahy (2009)
	<i>S. madagascariensis</i>	Scott <i>et al.</i> (1998); Kinoshita <i>et al.</i> (1999); López-García & Maillet J. (2005); Le roux <i>et al.</i> (2006, 2010); Prentis <i>et al.</i> (2007); Dormontt <i>et al.</i> (2014)
	<i>S. pterophorus</i>	Scott & Delfosse (1992); Pino <i>et al.</i> (2000); Caño <i>et al.</i> (2008)
	<i>S. vulgaris</i>	Zhu <i>et al.</i> (2017)
Physiological variation	<i>Senecio</i> species adapted to dry and wet habitats	Crawford (1966)
Life history	<i>S. sylvaticus</i> , <i>S. viscosus</i>	Palmblad (1968)
	<i>S. vulgaris</i>	Harper & Ogden (1970); Popay & Roberts (1970ab); Kadereit (1984c); Abbott (1986); Fenner (1986ab); Ren & Abbott (1991); Leiss & Müller-Schärer (2001)
	<i>S. vernalis</i>	Comes (1995)
	<i>S. lautus</i>	Walter <i>et al.</i> (2018b)
Chemical defence by pyrrolizidine alkaloids	<i>S. vulgaris</i>	Aplin <i>et al.</i> (1968); Hartmann <i>et al.</i> (1989); Frischknecht <i>et al.</i> (2001); Cheng <i>et al.</i> (2017)
	<i>S. vulgaris</i> , <i>S. vernalis</i>	Hartmann & Zimmer (1986)
	<i>S. pterophorus</i>	Castells <i>et al.</i> (2014)
	<i>S. nemorensis</i> complex	Bog <i>et al.</i> (2017b)
	Diverse <i>Senecio</i>	Langel <i>et al.</i> (2011)
Disease susceptibility to:		
- <i>Erysiphe fischeri</i>	<i>S. vulgaris</i>	Ben-Kalio & Clarke (1979); Harry & Clarke (1986)
- <i>Puccinia lagenophorae</i>	<i>S. vulgaris</i>	Paul & Ayres (1984)
- <i>Puccinia expansa</i>	Diverse <i>Senecio</i>	Alber <i>et al.</i> (1986)
Plant-pollinator interactions	Diverse <i>Senecio</i>	Schmitt (1980, 1983); Abbott & Irwin (1988)
Interspecific competition	<i>S. vulgaris</i>	Aarssen (1989); Bergelson & Perry (1989); Bergelson (1990ab); Bergelson <i>et al.</i> (1993)
Long distance dispersal	<i>S. flavus</i> , <i>S. mohavensis</i>	Liston <i>et al.</i> (1989); Liston & Kadereit (1995); Coleman <i>et al.</i> (2003)
	<i>S. inaequidens</i>	Pelser <i>et al.</i> (2012)

### Box 1 – Future research using *Senecio* as a model system

Below we detail research directions we consider likely to be particularly productive using *Senecio* as a model system.

**1. Responses to climate change.** Due to its widespread distribution and occupation of numerous habitats, *Senecio* allows the response of populations and communities to climate change to be tested across a range of biotic and abiotic interactions. Such tests could include: **(a)** Responses in ecologically similar but phylogenetically different *Senecio* (e.g., across Mediterranean or Alpine habitats in Europe, Africa, and Australasia). **(b)** Comparisons of ecologically different but phylogenetically similar species: e.g. Alpine species can be compared to closely related taxa at lower elevations, as on Mt Etna. **(c)** Field transplants across elevational gradients to simulate future climate scenarios. These transplants can also assess how transitions in plant, mycorrhizal, microbial and insect communities across elevational gradients interact with changes in the fitness and phenotypes of different *Senecio* species.

**2. Factors determining patterns of adaptive radiation.** Adaptive radiation appears rapid and common in *Senecio*. A major challenge is to identify the factors that have driven adaptive diversification, leading to its broad distribution across environments and geography. Potential approaches could be: **(a)** Bringing phylogenetic and morphological data together to test whether innovation in key traits created species radiations in the Asteraceae. This could include traits associated with long-distance dispersal (i.e., the pappus), variation in leaf morphology, or the production of toxic alkaloids that deter herbivores. **(b)** Studying the genetics and evolution of barriers to gene flow in *Senecio* using examples of ecotypic divergence, recent hybrid speciation and stable hybrid zones. **(c)** Using the many self-incompatible and self-compatible species of *Senecio* to study the evolution of diverse mating systems, especially within a broad phylogenetic context. **(d)** Using the repeated examples of morphological convergence and divergence and trait gains and losses in *Senecio* to test for genomic convergence in the evolution of complex traits within a single plant genus; for example, the evolution of growth habit in ecotypes of *S. lautus* could be expanded by studying how dramatic variation in growth habit has evolved in varied, but closely related, *Senecio* species.

**3. Ecological and evolutionary importance of phenotypic plasticity.** Many species of *Senecio* show considerable variation in phenotypic plasticity in response to environmental heterogeneity (e.g. in *S. vulgaris*) and can be used in experiments to determine the relative importance of phenotypic plasticity versus adaptation in responding to such heterogeneity. The utility of *Senecio* in glasshouse and field experiments, especially when combined with quantitative genetic breeding designs, makes it possible to combine large experimental studies with emerging genetic approaches: **(a)** Using genetic transformation based on existing *Agrobacterium*-mediated methods or the development of CRISPR/Cas9 gene editing, it will be possible to conduct functional analyses of genes associated with specific traits, as well as characterise the function of loci identified through field and -omics studies. **(b)** Transplanting a breeding design across environmental variation allows us to study changes in relative fitness, which can then be related to transcriptomic data to identify how differential gene expression contributes to fitness variation. Quantitative PCR could then be used to study the specific regions of the genome underlying differential expression, and the role of small RNAs (siRNA and miRNA) in determining patterns of differential expression to identify the regulatory basis of local adaptation and phenotypic plasticity.

**4. Developmental genetics of trait adaptations.** The diverse array of traits and habitats of *Senecio* species provide a rich resource for understanding the developmental and genetic pathways of many traits, for example, leaf shape variation. Studying the ecological variation underlying already identified pathways could complement work on *Arabidopsis* and relatives.

**5. Life history evolution.** With a diverse array of life history strategies in closely related species, *Senecio* can be used to study the ecological genetics of developmental plasticity, and the evolution of life history and resource allocation trade-offs, particularly where some species have overlapping generations, and others do not. For example, *Senecio* species with diverse growth habits could be used to study transgenerational effects on adaptive phenotypes, and how the nature of such transgenerational effects is shaped by environmental predictability and variability.

**6. The ecology and evolution of invasive species.** Notable examples of invasive species within *Senecio* are prime models for detailed analysis, especially where they are invading habitat already dominated by other *Senecio*: **(a)** In Australia, *S. madagascariensis* is colonising ecosystems occupied by a closely related native *Senecio* species. **(b)** In the UK, the recently-formed hybrid species *S. squalidus* has colonised much of the British Isles in the last 200 years, providing an important model system to study how invasive species form, spread and adapt rapidly to their new geographic range. **(c)** South African *Senecio inaequidens* has spread rapidly across Europe, the Americas and Australia, and shows differences in chromosome number across its range, allowing detailed investigation of how ploidy level correlates with invasiveness.

**7. Ecological and evolutionary importance of hybridisation.** *Senecio* contains many species that have been formed by recent hybridisation, with associated effects on chromosome number, genome size, and gene regulation. These include multiple examples of homoploid hybrid and allopolyploid species, and provide excellent opportunities to test how introgression changes the architecture of genetic variation across geography, and promotes hybrid speciation.

**8. How biotic and abiotic factors shape the distribution of widespread versus more specialised species.** In *S. lautus*, populations of ecotypes often exhibit very narrow distributions despite a wide distribution of their habitat, allowing studies to investigate what determines the distribution of populations, and their fundamental versus realised niches. For example, to test how abiotic stress tolerances and biotic interactions together determine range limits, and whether this changes across habitats.

## Figure legends

**Fig. 1:** With a worldwide distribution, species of *Senecio* show dramatic diversity in morphology and the habitats they occupy. Images are protected by copyright; this information is provided in parentheses and Note S1. 1. *S. wootonii* (CC BY-NC-SA 3.0 W. Anderson), 2. *S. pseudoarnica* (CC BY-NC 2.0 K. McFarland), 3. *S. soldanella* (CC BY-NC 3.0 M.E. Harte), 4. *S. flaccidus* (CC BY C.K. Kelly), 5. *S. clausenii* (CC BY N. Taylor & D. Zappi), 6. *S. petasitis* (© J. Moura), 7. *S. moqueguensis* (CC BY D.B. Montesinos-Tubée), 8. *S. candidans* (CC BY S.J. Tonge), 9. *S. barbertonicus* (CC BY-SA O. Leillinger), 10. *S. haworthii* (CC BY D. Avery), 11. *S. rowleyanus* (© Surreal Succulents, UK), 12. *S. nemorensis* (CC BY-NC C.T. Lin), 13. *S. halleri* (© I. Rey), 14. *S. leucanthemifolius* (CC BY-SA Petr Brož), 15. *S. graciliflorus* (CC BY Prashant Awale), 16. *S. cannabifolius* (CC BY Wikimedia), 17. *S. scandens* (CC BY-NC S. Duan), 18. *S. gregorii* (© P. Campbell).

**Fig. 2:** Australian *S. lautus* complex contains 12 ecotypes that have diverged rapidly from each other. (a) Ecotypes vary dramatically in leaf morphology and growth form, reflecting the diversity of habitats they occupy. (b-c) Ecotypes are distributed widely across Australia. Clades (represented by uppercase letters) group by geography, rather than ecology. (d) Morphological divergence of the ecotypes. Points and confidence intervals represent the mean and one standard error for each population measured. Colours represent ecotypes from panel c. See Note S2 for details of the data analysis.

**Fig. 3:** Hybrid origins of new *Senecio* taxa in the British Isles. Dates of first records are in red. For *S. vulgaris*, the non-radiate (nr) and radiate (r) forms are presented. Figure reproduced from Abbott and Rieseberg (2012) with permission.

**Fig. 4:** European *Senecio* species that have been studied extensively. *Senecio aethnensis* (a) and *S. chrysanthemifolius* (b) occur on Mount Etna (Sicily, Italy) at different altitudes. Hybrids between these two species were cultivated at the Oxford Botanic Garden (UK), from which they escaped in the late 18<sup>th</sup> century and spread across the UK, becoming a new species, *S. squalidus* (c). Hybridisation between *S. squalidus* and *S. vulgaris* (d) created the neoallohexaploid, *S. cambrensis* (e) and the tetraploid *S. eboracensis* (Fig. 3). Image of *S. vulgaris* © Botanical Society of Britain & Ireland, *S. cambrensis* © Frau-Doktor.