

Bipartite Sachdev-Ye models with Read-Saleur symmetriesJ. Classen-Howes¹, P. Fendley^{1,2}, A. Pandey³, and S. A. Parameswaran¹¹*Rudolf Peierls Centre for Theoretical Physics, Parks Road, Oxford OX1 3PU, United Kingdom*²*All Souls College, University of Oxford, Oxford OX1 4AL, United Kingdom*³*Department of Physics, Stanford University, Stanford, California 93405, USA*

(Received 27 June 2024; revised 30 August 2024; accepted 30 August 2024; published 19 September 2024)

We introduce an $SU(M)$ -symmetric disordered bipartite spin model with unusual characteristics. Although superficially similar to the Sachdev-Ye (SY) model, it has several markedly different properties for $M \geq 3$. In particular, it has a large nontrivial nullspace whose dimension grows exponentially with system size. The states in this nullspace are frustration-free and are ground states when the interactions are ferromagnetic. The exponential growth of the nullspace leads to Hilbert-space fragmentation and a violation of the eigenstate thermalization hypothesis. We demonstrate that the commutant algebra responsible for this fragmentation is a nontrivial subalgebra of the Read-Saleur commutant algebra of certain nearest-neighbor models such as the spin-1 biquadratic spin chain. We also discuss the low-energy behavior of correlations for the disordered version of this model in the limit of a large number of spins and large M , using techniques similar to those applied to the SY model. We conclude by generalizing the Shiraishi-Mori embedding formalism to nonlocal models, and apply it to turn some of our nullspace states into quantum many-body scars.

DOI: [10.1103/PhysRevB.110.125140](https://doi.org/10.1103/PhysRevB.110.125140)**I. INTRODUCTION**

Systems that violate the eigenstate thermalization hypothesis (ETH) have been intensely studied in recent years [1–5]. ETH is a set of constraints on the eigenstates of a quantum time-evolution operator such that for (almost) any initial state, observables evolve to a steady state at late times and exhibit only those correlations dictated by thermal equilibrium at the initial energy density. Such systems in effect “act as their own heat bath.” Generic isolated many-body systems are believed to satisfy ETH, giving an intuitive justification for applying various standard techniques of equilibrium statistical mechanics to time evolution. ETH can be violated by the presence of an extensive set of conservation laws, as occurs in integrable models [6]. Similar phenomena are believed to emerge in many-body-localized systems as a consequence of strong disorder [7]. In both cases, ETH is expected to be strongly violated, meaning that at most a set of measure zero of the eigenstates of such a quantum model obeys ETH.

A distinct symmetry-related mechanism for breaking ETH is Hilbert-space fragmentation (HSF) [2,8–11]. HSF occurs when the dimension of the commutant algebra of a Hamiltonian grows exponentially with system size, “fragmenting” the Hilbert space into exponentially many dynamically disconnected subspaces. The commutant algebra is defined for a family of Hamiltonians of the form $H = \sum_j J_j h_j$

as the set of all operators that commute with *each* of the local few-body operators h_j , and hence with H for any choice of the J_j . Strong HSF occurs when the size of each subspace is a vanishing fraction of the total size, leading to a strong violation of ETH. Weak HSF (and consequently a weak ETH violation) occurs when almost all eigenstates belong to a single subspace, i.e., the collective measure of all other subspaces tends to zero.

In this paper, we introduce an $SU(M)$ -symmetric bipartite spin model which exhibits HSF even in the presence of random and long-range couplings. Half of the $2N$ spins are in the fundamental $\mathbf{M}$ representation of $SU(M)$, and half in the conjugate $\bar{\mathbf{M}}$. The Hamiltonian consists of a sum over projectors onto the spin-singlet state for each $\mathbf{M}, \bar{\mathbf{M}}$ pair. This Hamiltonian generalizes the nearest-neighbor spin chains of Refs. [12,13], the most famous example being the “biquadratic” spin-1 chain.

We show by explicit construction that our Hamiltonian has a nullspace whose dimension grows exponentially with increasing system size. States in this nullspace are frustration-free, i.e., are annihilated by each of the projectors. When all the coefficients of the projectors are positive (the “ferromagnetic” case), they are ground states. Some of these states are product states, and the remainder can be found by exploiting the fact that the nullspace of the clean model (uniform couplings) is the same. For spin $M \geq 3$, we show that the resulting nullspace dimension grows exponentially with increasing system size. We identify precisely the $SU(M)$ representations in the nullspace and find their dimensions increase only polynomially with N . The exponential growth of the nullspace size therefore results in HSF.

This large nullspace arises as a consequence of a nontrivial commutant algebra. This algebra is a subalgebra of the

Read-Saleur (RS) algebra [14], whose elements commute with all the generators of certain presentations of the Temperley-Lieb algebra, including those used to build the nearest-neighbor analogs of our model [2]. Remarkably, here we find that some of the HSF and the commutant algebra present in the nearest-neighbor chains survives their generalization to our model as long as we preserve bipartiteness.

Our model can be viewed as a bipartite generalization of the Sachdev-Ye (SY) model [15,16], which has random all-to-all couplings and $SU(M)$ symmetry. The SY model, along with the closely related Sachdev-Ye-Kitaev (SYK) model of quartically coupled Majorana fermions, have received much attention for their links to quantum chaos. Many aspects of their thermalizing dynamics may be accessed using field-theoretic techniques. We also deploy such techniques in the limit of large M and N , and we find that the resulting equations are similar to those obtained for the bipartite SYK model [17]. They thus indicate similar non-Fermi-liquid behavior and emergent conformal symmetry at low energies for the appropriate disorder-averaged observables. However, our model exhibits the additional feature of HSF not apparent in the disorder-averaged field theory and not believed to be present in the SY model.

We show how to modify our Hamiltonian using a long-range variant of Shiraishi-Mori spectral embedding [18]. The product states are then spread throughout the spectrum and constitute quantum many-body scars [3]. The existence of these states results in a weak violation of ETH, and can be arranged so as to lead to periodic revivals, a hallmark of scars in experimental setups. Our model thus provides a rare example of a nonlocal model that violates ETH via fragmentation or scarring.

Our paper is organized as follows. In Sec. II, we define our Hamiltonian and demonstrate its key properties, namely the exponential growth of its nullspace with system size and the resulting Hilbert-space fragmentation. In Sec. III, we solve the uniform-coupling version of our model, and use this solution to derive an exact formula for the dimension of the nullspace of the full model. We show that the fragmented sectors can be identified solely by the $SU(M)$ symmetry: states in certain $SU(M)$ representations are always part of the nullspace, while others never belong to the nullspace. We find the commutant algebra in Sec. IV and show how it is a remnant of the Read-Saleur algebra. In Sec. V we exploit the bipartite Sachdev-Ye structure to discuss the structure of the disorder-averaged correlations of this model at low energies. We generalize the Shiraishi-Mori embedding formalism to nonlocal Hamiltonians in Sec. VI and show that the resulting scar states violate ETH. We apply this formalism to yield a nonlocal model with an exponentially growing number of scar states, leading to weak fragmentation. In Sec. VII, we provide concluding remarks.

II. BASIC PROPERTIES

In this section, we introduce our Hamiltonian H_M and derive some of its key properties. We explain how it can be thought of as a bipartite version of the well-known Sachdev-Ye model, an all-to-all $SU(M)$ -symmetric model. Despite the resemblance, we show that it has some significantly

different properties. Namely, we demonstrate that the dimension of the nullspace of H_M grows exponentially with increasing system size. The states in this nullspace are frustration-free and correspond to ground states for ferromagnetic couplings. A consequence is that our model is Hilbert-space fragmented, with each map between two nullspace eigenstates constituting an element of the commutant algebra.

A. Hamiltonian

We study a spin- S $SU(M)$ -symmetric bipartite system, with $M = 2S + 1$. The system is composed of $2N$ spins, with N spins transforming in the fundamental representation $\mathbf{M}$ of $SU(M)$ and N transforming in the conjugate representation $\bar{\mathbf{M}}$. We will usually label the spins of the former by upper indices i and the latter by lower indices j with $1 \leq i, j \leq N$. The corresponding states for each spin are labeled, respectively, as $|a^i\rangle$ and $|a_j\rangle$ with $1 \leq a \leq M$. We work in a basis where the symmetry generators are built from operators acting on a given spin as

$$T_b^{a,i} = |a^i\rangle\langle b^i|, \quad S_{b,j}^a = -|b_j\rangle\langle a_j|, \quad (1)$$

while acting trivially on the others. The operators

$$J_b^a = \sum_{i=1}^N T_b^{a,i} + \sum_{j=1}^N S_{b,j}^a \quad (2)$$

generate a $U(M)$ algebra, but the $U(1)$ generator $\sum_a J_a^a$ vanishes on any state in our Hilbert space. Thus, effectively the symmetry is $SU(M)$.

We study the bipartite model with all two-spin interactions between each pair (i, j) with i in the $\mathbf{M}$ half and j in the $\bar{\mathbf{M}}$ half. The spins in each half do not interact amongst themselves. The $SU(M)$ -invariant two-spin interaction can be written in terms of the (unnormalized) projector onto a singlet, which is given in this basis by

$$P_j^i = - \sum_{a,b=1}^M T_b^{a,i} S_{a,j}^b = \sum_{a,b=1}^M |a^i a_j\rangle\langle b^i b_j|. \quad (3)$$

The eigenstate of P_j^i with nonvanishing eigenvalue M is

$$|\psi_j^{0,i}\rangle = \sum_{a=1}^M |a^i a_j\rangle. \quad (4)$$

The most general such Hermitian Hamiltonian is composed of a sum over singlet projectors for each pair with arbitrary real coefficients r_j^i :

$$H_M = \sum_{i,j=1}^N r_j^i P_j^i. \quad (5)$$

For $M = 2$, the projector is the usual spin- $\frac{1}{2}$ Heisenberg interaction. When $M = 3$, a familiar way to rewrite the Hamiltonian is in terms of the “biquadratic” spin-1 interaction. Namely, we let $S^i = ((S^x)^i, (S^y)^i, (S^z)^i)$ denote the usual spin-1 $SU(2)$ generators acting on spin i , and likewise for S_j acting on spin j . The projector then can be unitarily

transformed into

$$U_3 P_i^j U_3^\dagger = (S^i \cdot S_j)^2 - \mathbb{1},$$

$$\text{where } U_3 = \prod_{j=1}^N \exp(i\pi S_j^z) \exp(i\pi S_j^x). \quad (6)$$

As has long been known [12], the symmetry of the biquadratic spin chain is enhanced to $SU(3)$ with generators given by Eq. (2).

The nearest-neighbor uniform chain analog of H_M is found by setting $r_j^i = 0$ unless $j = i, i+1$, yielding a chain with $2N$ sites and nearest-neighbor interactions:

$$H_{nn} = \sum_{i=1}^N (r_i^i P_i^i + r_{i+1}^i P_{i+1}^i). \quad (7)$$

This Hamiltonian has long been studied (see, e.g., Refs. [12,13]), because its generators obey the Temperley-Lieb algebra. With uniform couplings, the model is integrable, and many properties can be computed utilising the XXZ chain whose generators satisfy the same algebra.

Remarkably, H_{nn} has a large commutant algebra whose generators were derived by Read and Saleur (RS) [14]. Each element of this algebra commutes with each projector from Eq. (3) *individually*, and so commutes with H_{nn} for any couplings. For $M = 2$, where H_{nn} is the Heisenberg chain, the commutant algebra is simply $SU(2)$. However, for $M \geq 3$, the dimension of the RS algebra grows exponentially with system size, leading to exponentially large degeneracies in the spectrum. A heuristic way of understanding why is to note that the spectrum in essence follows from properties of the Temperley-Lieb algebra, while the dimensions of the Hilbert spaces of the XXZ chain and H_{nn} grow as 2^{2N} and M^{2N} , respectively. Thus, for $M \geq 3$, it is natural to expect large degeneracies in the spectrum of the latter. We discuss the ensuing RS algebra in more detail in Sec. IV.

B. Hilbert-space fragmentation

One of the central results of our paper is that H_M with completely arbitrary couplings r_j^i has for $M \geq 3$ a nullspace whose dimension grows exponentially with N , and as a consequence the model exhibits Hilbert-space fragmentation. We show these facts here by a rather elementary analysis. We derive an exact formula for the dimension of this nullspace in Sec. III, and show how the $SU(M)$ symmetry allows us to precisely characterize the corresponding states. In Sec. IV, we connect this result to the survival of a nontrivial subalgebra of the RS algebra.

Since H_M from Eq. (5) is a sum of projectors, its nullspace $\mathcal{N}$ for arbitrary couplings consists of those states annihilated by each projector P_j^i individually. Such states are said to be frustration-free. An exponentially large number of states in $\mathcal{N}$ are product states, and so can be found easily. We write a product state $|\psi_{\mathcal{N}}\rangle$ as

$$|\psi_{\mathcal{N}}\rangle = \otimes |t^1 t^2 \dots t^N s_1 s_2 \dots s_N\rangle, \quad (8)$$

where each $t^i, s_j \in \{1, \dots, M\}$ denotes the state of the spin i, j in the $\mathbf{M}$ and $\overline{\mathbf{M}}$ irreps, respectively. As apparent from

Eq. (3), if we choose t^i and s_j such that

$$t^i \leq z < s_j, \quad \text{for any } z \in \{1, \dots, M-1\}, \quad (9)$$

then $|\psi_{\mathcal{N}}\rangle$ will be annihilated by each projector in Eq. (5). The total number of possible values for all the levels t^i and s_j such that the above constraint is satisfied is $[z(M-z)]^N$ for each value of z .

Obviously, for $M \geq 3$, the number of product states in the nullspace obeying Eq. (9) grows exponentially with N . Since all these product states belong to nontrivial $SU(M)$ multiplets, the number of them provides a lower bound to the dimension $D_{\mathcal{N}}$ of the nullspace:

$$D_{\mathcal{N}} \geq \sum_{z=1}^{M-1} z^N (M-z)^N. \quad (10)$$

For N large, this bound is sharply peaked around $z \approx M/2$, so we expect the exponential growth to be

$$D_{\mathcal{N}} \propto \begin{cases} \left(\frac{M}{2}\right)^{2N}, & M \text{ even,} \\ \left(\frac{M^2-1}{4}\right)^N, & M \text{ odd.} \end{cases} \quad (11)$$

For ferromagnetic couplings ($r_j^i > 0$ for all i and j), each state in the nullspace is a ground state, yielding an exponentially growing ground-state degeneracy.

For any couplings, the large nullspace results in Hilbert-space fragmentation. In its analysis we follow the approach of Refs. [2,19,20]. The bond algebra $\mathcal{A}$ of H_M is generated by the projectors P_j^i and the identity operator, with elements given by linear combinations of arbitrary products of generators. The commutant algebra $\mathcal{C}$ is the algebra of operators $\widehat{O}$ that commute with each generator of $\mathcal{A}$ individually:

$$\mathcal{C} = \{\widehat{O} : [\widehat{O}, P_j^i] = 0, \text{ for } 1 \leq i, j \leq N\}. \quad (12)$$

Under $\mathcal{C}$, the Hilbert space splits into dynamically disconnected Krylov subspaces. The total number K of Krylov subspaces is bounded by the dimension of $\mathcal{C}$ [2] as

$$\sqrt{\dim(\mathcal{C})} \leq K \leq \dim(\mathcal{C}). \quad (13)$$

Hilbert-space fragmentation occurs in local models when $\log(\dim(\mathcal{C}))$, and hence $\log(K)$, grows as a volume law with system size. The fragmentation is nontrivial when $\mathcal{A}$ is non-Abelian, as in our case. (If $\mathcal{A}$ were abelian, the model would be trivially solvable.) As H_M is effectively zero-dimensional, we define fragmentation to occur when $\log(\dim(\mathcal{C}))$ grows linearly with N . We also require that this exponential growth does not arise solely from the degeneracies arising from the $SU(M)$ symmetry, i.e., the number of $SU(M)$ multiplets in the nullspace must grow exponentially.

Since the states in the nullspace $\mathcal{N}$ are annihilated by each P_j^i individually, maps between them that leave all other states invariant are elements of $\mathcal{C}$. Indeed, for any two states $|\psi_A\rangle, |\psi_B\rangle \in \mathcal{N}$, we have that

$$[P_j^i, |\psi_A\rangle\langle\psi_B|] = 0, \quad \text{for } 1 \leq i, j \leq N. \quad (14)$$

Hence, $\dim(\mathcal{C}) \geq (\dim(\mathcal{N}))^2$, and so the size of the commutant algebra grows exponentially with N . This leads to

Hilbert-space fragmentation, with each state in $\mathcal{N}$ constituting its own one-dimensional Krylov subspace.

In Sec. III D we derive an exact formula for the dimension of the nullspace, and find precisely which $SU(M)$ representations are present in it. This analysis allows us to show that the degeneracies are exponentially larger than those arising solely from the $SU(M)$ symmetry. Thus, our model exhibits Hilbert-space fragmentation.

C. Comparing H_M to the Sachdev-Ye Model

As an $SU(M)$ -symmetric spin model with disordered long-range interactions, H_M is reminiscent of the well-known Sachdev-Ye (SY) model [15,16]. The SY model is a disordered all-to-all variation on the $SU(M)$ -symmetric Heisenberg model [15]. Its Hamiltonian is given by

$$H_{SY} = \frac{1}{\sqrt{MN}} \sum_{i < j} \sum_{a,b=1}^M g_{ij} \mathcal{S}_{b,i}^a \mathcal{S}_{a,j}^b, \quad (15)$$

where $\mathcal{S}_{b,i}^a$ are generators of $SU(M)$ acting nontrivially at site i , and the g_{ij} are independent and identically distributed Gaussian random variables with vanishing means and variances that do not scale with N or M .

The crucial distinction between the SY model and ours is that all sites in the former transform in the same $SU(M)$ irreducible representation. When that representation is the fundamental $\mathbf{M}$, each operator $\mathcal{S}_{b,i}^a \mathcal{S}_{a,j}^b$ is equal (up to a constant shift) to the permutation operator $\mathbb{P}_{ij}$ swapping the spins on sites i and j .

As a consequence, H_{SY} and H_M have rather different physical properties. The permutation operators $\mathbb{P}_{ij}$ in SY split the M^2 states into $M(M+1)/2$ and $M(M-1)/2$ -dimensional subspaces. The singlet-projector operators P_j^i comprising H_M have rank 1, annihilating all but one of the basis states in the M^2 -dimensional Hilbert space formed by two spins. It is therefore not surprising that the H_M has a nontrivial nullspace, unlike SY. We would also expect the ranks of the projectors P_j^i to influence the rest of the spectrum, with H_M having a higher density of states close to energy zero than this SY model. We have checked this behavior numerically and indeed confirmed it at small N .

The SY model is therefore expected to obey ETH, as tested in the fermionic analog, the Sachdev-Ye-Kitaev (SYK) model [21,22]. A reason is that when all sites are in the fundamental representation, the SY model has the same bond algebra as the $SU(M)$ Heisenberg model. Indeed any $\mathbb{P}_{ij}$ is the product of nearest-neighbor permutation operators, e.g., for $j > i$, $\mathbb{P}_{ij} = \mathbb{P}_{i,i+1} \mathbb{P}_{i+1,i+2} \cdots \mathbb{P}_{j-1,j}$ [23]. The same bond algebra implies the same commutant algebra as well, and including random all-to-all couplings will not increase the symmetry. The $SU(M)$ Heisenberg model for $M > 2$ is not integrable and has no known symmetry algebra beyond $SU(M)$. It therefore obeys ETH, and the SY model thus should as well.

III. EXACT RESULTS

We here obtain the nullspace $\mathcal{N}$ of H_M . As $\mathcal{N}$ is defined as the set of states annihilated by all the projectors, it is independent of the couplings. We thus can find it by solving the

“clean” case, where all couplings are the same. We derive an exact formula for its exponentially large dimension, and show precisely which $SU(M)$ representations appear in $\mathcal{N}$. We also consider the impact of $SU(M)$ -symmetric perturbations.

A. Energies of the clean model

The clean version of H_M is given by setting all couplings r_j^i in Eq. (5) to one:

$$H_{\text{clean}} = \sum_{i,j=1}^N P_j^i = - \sum_{i,j=1}^N \sum_{a,b=1}^M T_b^{a,i} \mathcal{S}_{a,j}^b. \quad (16)$$

This Hamiltonian is invariant under permutations of the N spins in the $\mathbf{M}$ representation amongst themselves, and likewise for those in the $\bar{\mathbf{M}}$. This symmetry enhancement allows the spectrum of H_{clean} to be computed from a purely group-theoretical analysis.

The method for solving H_{clean} is to write it as a sum of Casimir invariants. Any representation of a Lie algebra possesses a quadratic Casimir operator that commutes with all the generators of the algebra in that representation. The full system we are studying is composed of N spins in the $\mathbf{M}$ representation of $SU(M)$ and N in the $\bar{\mathbf{M}}$, i.e., the representation is $\mathbf{M}^{\otimes N} \otimes \bar{\mathbf{M}}^{\otimes N}$ with generators given in Eqs. (1) and (2). The corresponding Casimir operator is

$$C_{\mathbf{M}^{\otimes N} \otimes \bar{\mathbf{M}}^{\otimes N}} = \sum_{a,b=1}^M J_b^a J_a^b. \quad (17)$$

We also need the $SU(M)$ Casimir operators for just the $\mathbf{M}$ spins and for just the $\bar{\mathbf{M}}$ spins, which are

$$\begin{aligned} C_{\mathbf{M}^{\otimes N}} &= \sum_{a,b=1}^M \mathcal{T}_b^a \mathcal{T}_a^b = -\frac{N^2}{M} + \sum_{i=1}^N \sum_{a,b=1}^M T_b^{a,i} T_a^{b,i}, \\ C_{\bar{\mathbf{M}}^{\otimes N}} &= \sum_{a,b=1}^M \mathcal{S}_b^a \mathcal{S}_a^b = -\frac{N^2}{M} + \sum_{a,b=1}^M \sum_{j=1}^N \mathcal{S}_{b,j}^a \mathcal{S}_{a,j}^b, \end{aligned} \quad (18)$$

where the $SU(M)$ generators acting on the $\mathbf{M}^{\otimes N}$ and $\bar{\mathbf{M}}^{\otimes N}$ subspaces are, respectively,

$$\mathcal{T}_b^a = \sum_{i=1}^N \left(T_b^{a,i} - \frac{1}{M} \delta_b^a \right), \quad \mathcal{S}_b^a = \sum_{j=1}^N \left(\mathcal{S}_{b,j}^a + \frac{1}{M} \delta_b^a \right),$$

giving traceless operators as needed.

The clean Hamiltonian therefore can be written in terms of these three Casimir operators as

$$H_{\text{clean}} = \frac{N^2}{M} + \frac{1}{2} (C_{\mathbf{M}^{\otimes N}} + C_{\bar{\mathbf{M}}^{\otimes N}} - C_{\mathbf{M}^{\otimes N} \otimes \bar{\mathbf{M}}^{\otimes N}}). \quad (19)$$

Other models consisting of a sum of $SU(M)$ Casimirs have been analyzed [24–27], but all spins in these models are in the $\mathbf{M}$ representation.

The key property of the quadratic Casimir operator is that for an irreducible representation it is proportional to the identity. Irreducible representations of $SU(M)$ are characterized by an M -dimensional vector $\rho = [\rho_1, \rho_2, \dots, \rho_M]$ composed of integers satisfying

$$\rho_1 \geq \rho_2 \geq \dots \geq \rho_M \geq 0. \quad (20)$$

Each such vector can be written as a partition $P_{S,k}$ of $S = \sum_a \rho_a$ into k positive integers, where $k \leq M$. For any partition with $k < M$, we set $\lambda_b = 0$ for $b > k$. The vector corresponding to the fundamental $\mathbf{M}$ representation is $[1, 0, 0, \dots, 0]$, while for the conjugate $\overline{\mathbf{M}}$ it is $[1, 1, \dots, 1, 0]$. $SU(M)$ representations related by shifting all $\rho_a \rightarrow \rho_a - 1$ (until $\rho_M = 0$) are equivalent.

The quadratic Casimir operator obeys

$$C_\rho = c_\rho \mathbb{1} \quad \text{for } \rho \text{ irreducible.} \quad (21)$$

For any representation of a semisimple Lie algebra, the number c_ρ can be determined by standard techniques. For $SU(M)$, it is [28]

$$c_\rho = \sum_a \left(\rho_a^2 + (M+1-2a)\rho_a \right) - \frac{1}{M} \left(\sum_a \rho_a \right)^2, \quad (22)$$

where here and everywhere unlabeled sums over a run from 1 to M . The Casimir c_ρ is indeed invariant under the shift $\rho_a \rightarrow \rho_a - 1$.

Group theory alone yields all the eigenvalues of H_{clean} . Finding them requires decomposing the Hilbert space $\mathbf{M}^{\otimes N} \otimes \overline{\mathbf{M}}^{\otimes N}$ into irreducible representations of $SU(M)$. Since Eq. (19) involves also the Casimir operators for the two types of spins alone, we must also keep track of the representations of $SU(M)$ in the corresponding subspaces. The simplest way to understand tensor products of $SU(M)$ representations is via Young diagrams. The Young diagram for an irreducible representation ρ is a collection of boxes glued together such that the a th row has ρ_a boxes in it. To take the tensor product of two representations, one glues the boxes from the corresponding Young diagrams together subject to certain rules (see, e.g., Ref. [29] or any text on Lie algebras).

We first need to decompose the tensor products $\mathbf{M}^{\otimes N}$ and $\overline{\mathbf{M}}^{\otimes N}$ into a direct sum over irreducible representations λ and $\overline{\gamma}$, respectively, and then decompose the tensor product of each λ and $\overline{\gamma}$ pair. The irreducible representations λ in the decomposition of $\mathbf{M}^{\otimes N}$ obey $\sum_a \lambda_a = N$. They therefore can be written as integer partitions of N into at most M parts:

$$\mathbf{M}^{\otimes N} = \oplus m(\lambda) \lambda, \quad \lambda \in P_{N,k} \text{ for } k \leq M, \quad (23)$$

where $m(\lambda)$ is the ‘‘multiplicity,’’ the number of times λ appears in this direct sum. We give its value below. The decomposition of $\overline{\mathbf{M}}^{\otimes N}$ can be characterized similarly by taking advantage of conjugate representations. The conjugate $\overline{\gamma}$ is defined via $\overline{\gamma}_a = \gamma_1 - \gamma_{M+1-a}$. The conjugate of a tensor product is the tensor product of the conjugates. We then utilize Eq. (23) to get

$$\overline{\mathbf{M}}^{\otimes N} = \oplus m(\gamma) \overline{\gamma}, \quad \gamma \in P_{N,l} \text{ for } l \leq M. \quad (24)$$

Here $\sum_a \overline{\gamma}_a = M\gamma_1 - N$.

To decompose the full Hilbert space into irreducible representations, we then take the tensor product of each pair to give $\lambda \otimes \overline{\gamma} = \oplus \rho$. The Hilbert space thus can be decomposed into sectors labeled by $(\lambda, \overline{\gamma}, \rho)$. The clean Hamiltonian is diagonal in such a basis, with any state $|\psi_{\lambda, \overline{\gamma}, \rho}\rangle$ in a given sector obeying

$$H_{\text{clean}} |\psi_{\lambda, \overline{\gamma}, \rho}\rangle = \left(\frac{N^2}{M} + \frac{1}{2}(c_\lambda + c_{\overline{\gamma}} - c_\rho) \right) |\psi_{\lambda, \overline{\gamma}, \rho}\rangle. \quad (25)$$

This spectrum of the clean model is highly degenerate, and below we compute its ground-state degeneracy.

B. Ground states of the clean model

To find the dimension of the nullspace of H_M , we first construct the states annihilated by H_{clean} . Since eigenvalues of H_{clean} and Casimirs of $SU(M)$ are always nonnegative, from Eq. (25) we have

$$c_\rho \leq 2\frac{N^2}{M} + c_\lambda + c_{\overline{\gamma}}, \quad (26)$$

with ground states given by those representations $\rho \in \lambda \otimes \overline{\gamma}$ satisfying the equality.

Thus, for a given pair $\lambda, \overline{\gamma}$, candidates for ground states must maximize c_ρ . To find this maximum, we exploit a useful inequality for $SU(M)$ representations: for any $\rho \in \lambda \otimes \overline{\gamma}$, then [30]

$$\sum_{a=1}^b (\lambda_a + \overline{\gamma}_a) \geq \sum_{a=1}^b \rho_a \quad (27)$$

for any b . It follows from the explicit expression (22) that this c_ρ is maximized when ρ obeys the equality in Eq. (27), i.e., $\rho = \lambda + \overline{\gamma}$. Here the Young diagram for ρ is given by gluing those for λ and $\overline{\gamma}$ together horizontally.

The zero-energy ground states of H_{clean} are therefore those λ and $\overline{\gamma}$ for which

$$c_{\lambda+\overline{\gamma}} = 2\frac{N^2}{M} + c_\lambda + c_{\overline{\gamma}}. \quad (28)$$

By construction $\sum_a \lambda_a = N$ and $\sum_a \overline{\gamma}_a = M\gamma_1 - N$, so that Eq. (22) yields

$$\begin{aligned} c_{\lambda+\overline{\gamma}} - c_\lambda - c_{\overline{\gamma}} &= 2 \sum_a \lambda_a \overline{\gamma}_a - 2N\gamma_1 + 2\frac{N^2}{M} \\ &= -2 \sum_a \lambda_a \gamma_{M+1-a} + 2\frac{N^2}{M}. \end{aligned} \quad (29)$$

The condition (28) then reduces to

$$\sum_a \lambda_a \gamma_{M+1-a} = 0. \quad (30)$$

In Eqs. (23) and (24) we defined λ and γ as partitions of N . Since both λ_a and γ_{M+1-a} are nonnegative, one of these two must vanish for all a for Eq. (30) to be satisfied.

The condition (30) therefore is equivalent to the remarkably simple constraint $k+l \leq M$. We thus find that the ground states of H_{clean} and the nullspace of H_M are all those $|\psi_{\lambda, \overline{\gamma}, \lambda+\overline{\gamma}}\rangle$ with

$$\boxed{\lambda \in P_{N,k}, \quad \gamma \in P_{N,l}, \quad k+l \leq M.} \quad (31)$$

This constraint is both necessary and sufficient for obtaining a zero-energy ground state of H_{clean} .

We thus have proved that the Hilbert space of H_M is fragmented: All states satisfying Eq. (31) are part of the null space. Since there are many such $SU(M)$ representations, this degeneracy goes well beyond the consequences of this global symmetry. In Sec. III D, we prove an even stronger statement: any state in a representation ρ that obeys $\rho = \lambda + \overline{\gamma}$ is in the nullspace, as long as λ and γ satisfy the conditions of Eq. (31).

C. Exact dimension of the nullspace

We here present an exact formula for the dimension of the nullspace $\mathcal{N}$ of H_M as a function of M and N . As discussed in Sec. IIB, all states in the nullspace are annihilated by each projector P_j^i . Because H_{clean} is a sum over these projection operators with positive coefficients, its eigenvalues are nonnegative, and all eigenstates with zero eigenvalue are annihilated by all P_j^i . Hence, the null states for any r_j^i are given by the zero-energy ground states of the clean model. We therefore may make use of the solution of the latter to derive the nullspace dimension $D_{\mathcal{N}}$. It therefore follows from purely group-theoretical calculations.

The nullspace is spanned by all the states in Eq. (31). The multiplicities $m(\lambda)$ and $m(\gamma)$ in Eqs. (23) and (24) give rise to one kind of degeneracy: any of the copies of λ and γ can be used to make $\rho = \lambda + \gamma$. The second kind of degeneracy comes from the fact that any of the states in the ρ representation give rise to a ground state. The number of these is $d(\lambda + \gamma)$, the dimension of this representation. The resulting number of ground states, and hence the dimension of the null space $\mathcal{N}$, is then

$$D_{\mathcal{N}} = \sum_{\substack{k, \ell=1 \\ k+\ell \leq M}}^{M-1} \sum_{\lambda \in P_{N,k}} \sum_{\gamma \in P_{N,\ell}} m(\lambda) m(\gamma) d(\lambda + \gamma), \quad (32)$$

which indicates that this degeneracy is significantly greater than can be explained via any obvious symmetry of the model.

The multiplicities and dimensions of these representations follow from standard group-theory methods. The N spins in the $\mathbf{M}$ representation are exchanged amongst each other by elements of the permutation group S_N , which in the clean model is a symmetry. The multiplicities $m(\lambda)$ in Eq. (23) are given by the representation of the corresponding irreducible representation of this S_N . Both this representation and the $SU(M)$ representation λ are labeled by the same Young diagram, a fact known as Schur-Weyl duality. The dimension of an S_N representation is given by

$$m(\lambda) = \frac{N!}{h(\lambda)}, \quad h(\lambda) \equiv \prod_{a,b} h_{a,b}, \quad (33)$$

where $h_{a,b}$ denotes the ‘‘hook length’’ of the box at row a , column b of the Young diagram for λ , and the product in $h(\lambda)$ is over all boxes. A hook comprises the box (a, b) , all the boxes to the right of it in the same row a , and all the boxes below it in the same column b . The hook length $h_{a,b}$ is the number of such boxes. The dimension $d(\rho)$ of a $SU(M)$ representation also involves the product of hook lengths. It is

$$d(\rho) = \frac{1}{h(\rho)} \prod_{a=1}^M \frac{(\rho_a + M - a)!}{(M - a)!}. \quad (34)$$

For $M = 2$, the formula (32) exhibits no great surprises. Here both k and ℓ must be 1, and the only partition $P_{N,1}$ is $[N, 0]$, which has $m([N, 0]) = 1$. The corresponding $SU(2)$ representation has dimension $d([2N, 0]) = 2N + 1$, so we find $D_{\mathcal{N}} = 2N + 1$, the dimension of the usual ferromagnetic multiplet.

As we noted in Eq. (11), for $M \geq 3$ the nullspace dimension grows exponentially with N . For $M = 3$ the precise

asymptotic is

$$D_{\mathcal{N}}|_{M=3} \approx \frac{9}{4} 2^N N^2 \text{ for large } N. \quad (35)$$

To prove Eq. (35), we first note that using the explicit expressions (33) and (34) for the multiplicities and the dimension in the exact formula (32) for $M = 3$ yields

$$D_{\mathcal{N}}|_{M=3} = (N+1)^3 + \sum_{n=1}^{\lfloor N/2 \rfloor} \mathcal{P}_n \binom{N}{n},$$

$$\mathcal{P}_n = (1+N-2n)^2(2+2N-n) \frac{1+N+n}{1+N-n}. \quad (36)$$

We define α via $n = N/2 - \alpha N$, so that the binomial is sharply peaked at α small. We take N large and α small, and use Stirling’s formula to approximate it as

$$\binom{N}{\frac{N}{2} - \alpha N} = 2^N \sqrt{\frac{2}{N\pi}} e^{-2\alpha^2 N} (1 + \mathcal{O}(\alpha^2, \alpha^4 N, N^{-1})).$$

This expression is exponentially small unless α is order $1/\sqrt{N}$ or smaller, justifying neglecting the other terms. We then can approximate

$$\mathcal{P}_n = 18\alpha^2 N^3 + \mathcal{O}(\alpha N^2, \alpha^3 N^3) \quad (37)$$

and replace the sum over n in Eq. (36) with an integral over $Nd\alpha$, giving

$$D_{\mathcal{N}}|_{M=3} \approx 2^N \cdot 18N^3 \sqrt{\frac{2N}{\pi}} \int_0^{\frac{1}{2}} d\alpha \alpha^2 e^{-2N\alpha^2}.$$

Since the integrand is sharply peaked around $\alpha = 0$, we can extend the upper limit of integration to ∞ and then do the Gaussian integral, yielding Eq. (35). The corrections are suppressed by $1/N$: the terms in Eq. (36) suppressed by only $1/\sqrt{N}$ cancel after doing the integral.

Another limit of Eq. (32) worth considering is the large M limit for fixed N . Since the rank of each projector P_j^i remains fixed at 1, and the total number of projectors is fixed at N^2 , we expect the nullspace to dominate in the large M limit. This is indeed the case: we show in App. VII that the limiting form is the dimension of the total Hilbert space:

$$\lim_{M \rightarrow \infty} \frac{D_{\mathcal{N}}}{M^{2N}} = 1. \quad (38)$$

We note that this effect does not happen in the SY model with spins in the fundamental representation: the permutation operators $\mathbb{P}^{ij}$ split the Hilbert space of two spins into symmetric and antisymmetric subspaces, which have comparable dimensions in the large M limit.

D. $SU(M)$ and the nullspace

We have demonstrated degeneracies in the spectrum of H_M going well beyond $SU(M)$, as for $M > 2$ many different representations obey the nullspace condition (31). The number of each such representation is the product of multiplicities $m(\lambda)m(\gamma)$, and it grows exponentially with N . For example, the binomial in the $SU(3)$ formula (36) arises from the multiplicities, not the degeneracies coming from a particular $SU(M)$ representation.

However, we see no evidence for degeneracies and fragmentation outside the nullspace. Thus, despite its being exponentially large, the nullspace for H_M still comprises a vanishing fraction of the full Hilbert space for M finite. Thus, fragmentation occurs only in a set of measure zero of the full Hilbert space, and almost all states obey ETH.

The $SU(M)$ representations appearing in the nullspace defined by Eq. (31) are rather special. The reason is that all the projectors P_j^i must annihilate states in the nullspace, and when forming $\rho \in \lambda \otimes \bar{\gamma}$, no singlets can appear.

This property leads to a rather interesting feature of H_M . We show here that determining whether a given eigenstate is in the nullspace requires only knowing how it transforms under $SU(M)$. Namely, any eigenstate of H_M in the $SU(M)$ representation $\rho = \lambda + \bar{\gamma}$ satisfying the conditions of Eq. (31) must be in the nullspace. Any eigenstates of H_M in any other representations are not in the nullspace.

Precisely, any eigenstate of H_M can be labeled nonuniquely by the triple λ, γ, ρ where $\lambda, \gamma \in P_{N,M}$ and $\rho \in \lambda \otimes \bar{\gamma}$, just as shown in Eq. (25) for H_{clean} . In general the eigenvalues depend on more data than just these representations, but eigenstates in the nullspace of H_M satisfy the more restrictive properties given in Eq. (31). We prove that the only way to have $\rho = \rho$ for any such ρ is for the corresponding $\lambda = \lambda$ and $\gamma = \gamma$ as well.

Assume λ and γ satisfy Eq. (31), with $\rho = \lambda + \bar{\gamma}$. Say ρ also occurs in the decomposition of some other $\lambda \otimes \bar{\gamma}$. In the decomposition, the boxes in the Young diagram for λ will be rearranged according to the tensor product rules into rows of length $n_1, \dots, n_M$, with $\sum_{a=1}^M n_a = N$, as they are attached to the rows in the diagram of $\bar{\gamma}$ to construct ρ . Row by row in the diagram of ρ , this yields the set of equations

$$\gamma_1 - \gamma_{M-a+1} + \lambda_a + m = \gamma_1 - \gamma_{M-a+1} + n_a, \quad (39)$$

where the integer m accounts for the fact $SU(M)$ irreps are equivalent up to a constant shift in the lengths of all rows. Summing the left and right hand sides of Eq. (39) yields $\gamma_1 + m = \gamma_1$, giving

$$-\gamma_{M-a+1} + \lambda_a = -\gamma_{M-a+1} + n_a \quad (40)$$

for each $a = 1 \dots M$. Let k be the number of nonzero elements in λ , and ℓ the number of nonzero elements in γ , with $k + \ell \leq M$ by Eq. (31). Summing the first k equations in Eq. (40) then yields

$$N = \sum_{a=1}^k n_a - \sum_{a=1}^k \gamma_{M-a+1}. \quad (41)$$

Because the first sum here is at largest N and the second nonnegative, we must have $\sum_{a=1}^k n_a = N$ and so $n_a = 0$ for $a \geq k + 1$. Using this result in Eq. (40) yields $m = 0$, $\gamma = \gamma$, and $n_a = \lambda_a$ for all values of a .

The final step comes from noting that the tensor product rules do not allow boxes in a Young diagram to move upward from their original row. Hence, the Young diagram of λ must have k rows or fewer, and the boxes in these rows can only be moved downward in the construction of the Young diagram of ρ to obtain $n_a = \lambda_a$ for $a \leq k$. However, since $\bar{\gamma}_a = \bar{\gamma}_a = \bar{\gamma}_1$ for $a \leq k$, any such downward movement would not be consistent with the tensor product rules: boxes originally in

the same row cannot end up in the same column after taking the tensor product. (The reason is that tensor indices are symmetrized in the rows of Young diagram, antisymmetrized in the columns.) Thus, the only consistent possibility for $\rho = \lambda + \bar{\gamma} \in \lambda \otimes \bar{\gamma}$ satisfying Eq. (31) is to have both $\gamma = \gamma$ and $\lambda = \lambda$, as claimed.

E. $SU(M)$ -symmetric perturbations

It is possible to add certain $SU(M)$ -symmetric perturbations to H_M while maintaining Hilbert-space fragmentation. The simplest such perturbation is the addition of permutation operators $\mathbb{P}_{ij}$ linking some spins which share a same sublattice. For example, setting N to be even, we consider the perturbed Hamiltonian

$$H'_M = H_M + \sum_{i=1}^{N/2} c^i \mathbb{P}^{2i-1, 2i} + \sum_{j=1}^{N/2} d_j \mathbb{P}_{2j-1, 2j}, \quad (42)$$

where the c^i and d_j are arbitrary distinct real coefficients, and upper and lower indices denote different spin sublattices as before. The permutation operators introduced in H'_M split the original nullspace of H_M into degenerate subspaces, the eigenvalues of which depend on how the interacting spins on each sublattice are divided into symmetrised and anti-symmetrised pairs. It is apparent that projectors onto the resulting eigenstates will form part of the commutant algebra. It is also simple to show that many of these subspaces will grow exponentially in dimension with N . For example, the dimension of the subspace where each interacting pair of spins on a same sublattice is symmetrised can be lower bounded by

$$\sum_{\substack{k, \ell=1 \\ k+\ell \leq M}}^{M-1} \sum_{\lambda \in P_{N/2, k}} \sum_{\gamma \in P_{N/2, \ell}} m(\lambda) m(\gamma) d(2\lambda + 2\bar{\gamma}),$$

where 2λ refers to the irrep obtained by gluing two copies of λ together horizontally. Hence the perturbed model H'_M will also be Hilbert-space-fragmented.

It is also possible to maintain fragmentation while adding in $SU(M)$ -symmetric interactions involving multiple spins on a same sublattice. In this scenario, a sufficient condition for fragmentation is that it must be possible to divide at least one of the two sublattices of the perturbed Hamiltonian into a linear-in- N number of further sublattices, such that none of the newly formed sublattices are directly linked by a term in the Hamiltonian. In this case, one can again show the presence of fragmentation by considering the subspace in which all spins in each of the newly formed sublattices are symmetrised amongst themselves. One can then demonstrate that the number of ways of constructing suitable irreps satisfying these symmetrisation constraints and with the same eigenvalue still grows exponentially. On the other hand, we have numerically verified at small N that adding all-to-all permutation interactions on both of the sublattices removes any exponentially growing degeneracies, and hence also removes Hilbert space fragmentation.

IV. READ-SALEUR SYMMETRY ALGEBRA AND ITS REMNANT

A remarkable feature of the nearest-neighbor chain (7) is the presence of a large commutant algebra discovered by Read and Saleur (RS) [14]. The resulting degeneracies occur throughout almost all the spectrum. Their number grows exponentially with system size, and so results in Hilbert-space fragmentation [2]. We have shown that the exponentially large dimension of the nullspace and the ensuing fragmentation survive the introduction of the many additional couplings in H_M . As described in Sec. II B, this behavior automatically results in a commutant algebra acting within this nullspace. In this section we demonstrate that this algebra is a subalgebra of the RS algebra.

A. Read-Saleur algebra

We here describe the Read-Saleur commutant algebra of the nearest-neighbor chain (7). To make the structure more apparent, we renumber the sites as $p = 1 \dots 2N$, with odd p labeling spins in the $\mathbf{M}$ representation and even p labeling those in the $\bar{\mathbf{M}}$. We also relabel the generators of H_{nn} as $e_{2i-1} \equiv P_i^i$ and $e_{2i} \equiv P_{i+1}^i$ so

$$H_{\text{nn}} = \sum_{p=1}^{2N} r_p e_p, \quad e_p = \sum_{a,b=1}^M |a_p a_{p+1}\rangle \langle b_p b_{p+1}|. \quad (43)$$

with r_p the nonvanishing random couplings. Each projector e_p onto a nearest-neighbor $\text{SU}(M)$ singlet has only one nonvanishing eigenvalue out of the M^2 possibilities. Since the nearest-neighbor chain has only $2N$ such projectors, one expects large degeneracies for large enough M .

Comparison with the XXZ chain suggests that exponentially large degeneracies occur for any $M \geq 3$. The generators e_p satisfy the Temperley-Lieb algebra [12]:

$$(e_p)^2 = M e_p, \quad e_p e_{p \pm 1} e_p = e_p, \quad (44)$$

and $e_p e_{p'} = e_{p'} e_p$ with $|p - p'| > 1$. Many interesting lattice Hamiltonians are written in terms of the e_p ; one prominent example is the XXZ chain. In the uniform-coupling case any such chain is integrable, and so the relation between the two spectra can be determined exactly (see Ref. [13] and references therein). Since the Hilbert spaces for XXZ and H_{nn} are 2^{2N} - and M^{2N} -dimensional, respectively, there should be exponentially large degeneracies in the spectrum of H_{nn} when $M > 3$.

Read and Saleur [14] showed how the degeneracies are independent of the couplings r_p in H_{nn} . They did so by constructing the irreducible representations of the Temperley-Lieb algebra in this chain. The resulting degeneracies are completely independent of the couplings, following solely from understanding how the Temperley-Lieb algebra works here. Not surprisingly, the nullspace has the largest degeneracy, but large degeneracies are found for all but a small set of states.

A key tool of Read and Saleur's results is the construction of the commutant algebra $\mathcal{C}^{\text{RS}}$ of the e_p . It is generated by $\mathbf{J}^{(s)}$ that satisfy

$$e_p \mathbf{J}^{(s)} = \mathbf{J}^{(s)} e_p = 0 \quad (45)$$

for all p . RS find a linear basis set for the generators in terms of the same building blocks (1) used to construct the $\text{SU}(M)$ generators, namely

$$J_{b,2i-1}^a = T_b^{a,i}, \quad J_{b,2j}^a = S_{b,j}^a. \quad (46)$$

Generators of $\mathcal{C}^{\text{RS}}$ are of the form

$$\begin{aligned} \mathbf{J}^{(s)} &= \sum_{a_1, b_1, \dots} \tilde{g}_{a_1 a_2 \dots a_s}^{b_1 b_2 \dots b_s} \mathbf{J}_{b_1 b_2 \dots b_s}^{a_1 a_2 \dots a_s}, \\ \mathbf{J}_{b_1 b_2 \dots b_s}^{a_1 a_2 \dots a_s} &\equiv \sum_{1 \leq p_1 < p_2 < \dots < p_s \leq 2N} J_{b_1, p_1}^{a_1} J_{b_2, p_2}^{a_2} \dots J_{b_s, p_s}^{a_s} \end{aligned} \quad (47)$$

for $0 \leq s \leq 2N$. Setting $s = 1$ yields the $\text{SU}(M)$ generators. For Eq. (45) to be satisfied, the necessary and sufficient conditions on the coefficients are

$$\sum_{\hat{a}=1}^M \tilde{g}_{\dots \hat{a}, b_{p+1} \dots}^{a_p, \hat{a}, \dots} = \sum_{\hat{a}=1}^M \tilde{g}_{\dots \hat{a}, a_{p+1} \dots}^{b_p, \hat{a}, \dots} = 0 \quad (48)$$

for all p . More explicit expressions for the $\mathbf{J}^{(s)}$ can be found in Ref. [14].

As apparent from Eq. (47), the RS algebra includes the universal enveloping algebra $\mathcal{U}(\text{SU}(M))$ [i.e., products of the $\text{SU}(M)$ generators in this representation]. It is easy to check that the $\mathbf{J}^{(s)}$ map between different representations of $\text{SU}(M)$, and so go beyond $\mathcal{U}(\text{SU}(M))$. RS derive a variety of interesting properties of $\mathcal{C}^{\text{RS}}$, for example showing how gluing together multiple systems allows one to define a tensor-product structure. More generally, they show the algebra is Morita equivalent to $\mathcal{U}_q(\mathfrak{sl}(2))$, the commutant algebra of some other Temperley-Lieb systems such as the XXZ chain. However, many interesting questions remain; even the minimal set of generators of $\mathcal{C}^{\text{RS}}$ and their commutation relations is not known.

B. Commutant algebra

We have shown explicitly in Sec. III C that degeneracies in the nullspace survive the inclusion of all N^2 couplings in H_M , and that one can define elements of the commutant algebra mapping in between them. We show here that these elements form a subalgebra of the RS algebra, in particular of the algebra generated by the $\mathbf{J}^{(2N)}$.

In the nearest-neighbor case, the $\mathbf{J}^{(2N)}$ act nontrivially on the nullspace of H_{nn} and annihilate all other states. The reason is that the $\mathbf{J}^{(2N)}$ annihilate any state with at least one nearest-neighbor $\text{SU}(M)$ singlet, and the eigenstates of H_{nn} with nonzero eigenvalues are solely composed of such states [14]. Furthermore, for any two states in the nullspace of H_{nn} , there is an operator amongst the $\mathbf{J}^{(2N)}$ that maps between them.

We generalize this result here. Since all nullstates of H_M are also nullstates of H_{nn} , maps between the former are given by operators $\mathbf{J}^{(2N)}$ that satisfy certain additional constraints. For notational ease, we relabel the indices as

$$\mathbf{J}^{(2N)} = \sum g_{a^1 \dots a^N c_1 \dots c_N}^{b^1 \dots b^N d_1 \dots d_N} T_{b_1}^{a_1, 1} \dots T_{b_N}^{a_N, N} S_{d_1, 1}^{c_1} \dots S_{d_N, N}^{c_N}, \quad (49)$$

where this sum and all those in this section are over the appropriate a^i, b^i, c_j, d_j from 1 to M . To be in the commutant

algebra of H_M , these operators obey

$$\mathbf{J}^{(2N)} P_j^i = P_j^i \mathbf{J}^{(2N)} = 0, \text{ for } 1 \leq i, j \leq N. \quad (50)$$

These constraints require that the coefficients satisfy

$$\sum_{\{b\}, \{c\}} \delta_{bj}^{c_i} g_{\dots c_i \dots}^{b^j \dots} = \sum_{\{a\}, \{d\}} \delta_{dj}^{a_i} g_{\dots a_i \dots}^{d^j \dots} = 0 \quad (51)$$

for all $1 \leq i, j \leq N$.

A general null state of H_M is given by

$$|\psi\rangle = \sum_{\{a\}, \{d\}} \psi_{a^1 \dots a^N}^{d_1 \dots d_N} |a^1 \dots a^N d_1 \dots d_N\rangle, \quad (52)$$

where the a^i and d_j transform in the $\mathbf{M}$ and $\overline{\mathbf{M}}$ representations, respectively, and the coefficients satisfy

$$\sum_{\{a\}, \{d\}} \delta_{dj}^{a_i} \psi_{a^1 \dots a^N}^{d_1 \dots d_N} = 0, \text{ for } 1 \leq i, j \leq N. \quad (53)$$

The resemblance to the constraints on $\mathbf{J}$ is not coincidental. We can find a solution to Eq. (51) by taking any two solutions ψ, χ of Eq. (53) and defining

$$g_{a^1 \dots a^N}^{b^1 \dots b^N} d_1 \dots d_N = \psi_{a^1 \dots a^N}^{d_1 \dots d_N} (\chi_{b^1 \dots b^N}^{c_1 \dots c_N})^*. \quad (54)$$

We thus have proved that $|\psi\rangle\langle\chi|$ can be written as an operator $\mathbf{J}^{(2N)}$ from Eq. (47).

Regarding the other energy levels of H_M , our numerics found no sign of degeneracies other than those arising from the global $SU(M)$ symmetry. This suggests that the only operators $\mathbf{J}^{(s)}$ which act nontrivially on eigenstates outside the nullspace are elements of $\mathfrak{U}(SU(M))$. This suggestion is in harmony with Read and Saleur's derivation of the symmetry generators for the nearest-neighbor chain [14]. Their construction utilizes a “dimer” basis for the Hilbert space, where each state is written as a product of spin-singlet (dimer) states. Operators in the Temperley-Lieb algebra act nontrivially on the dimers, whereas those in the RS algebra leave the dimers untouched. The nonlocal projectors P_j^i in H_M , however, mix the dimer states in a much more complicated way, and so there is no reason to expect degeneracies beyond $SU(M)$. Indeed, from this point of view it is rather surprising that even the nontrivial nullspace survives.

V. LOW-ENERGY FIELD-THEORY DESCRIPTION

In Sec. IIC we discussed how H_M resembles a bipartite version of the SY model but with Hilbert-space fragmentation. Here we sharpen the comparison by analyzing a field-theory limit using manipulations familiar from studies of the SY/SYK models [15,31]. We will take the limit of a large number of spins and of large M , and subsequently the limit of frequencies small compared to microscopic scales when we explicitly discuss constraints on conformal *ansatze* for the low-energy correlations. As we explain next, we also need to generalize the spins to larger representations of $SU(M)$.

A. Spins in antisymmetric representations

A natural generalization of H_M is to take spins in representations other than the fundamental ones. We utilize the antisymmetric representation $\mathbf{A}_m$ of $SU(M)$, which has a

Young diagram with one column and m rows. We consider N spins in the $\mathbf{A}_m$ representation and $\tilde{N}$ in the $\overline{\mathbf{A}}_m = \mathbf{A}_{M-m}$ representation. The Hamiltonian is the obvious generalization of Eq. (5), namely,

$$H_{M,m} = \frac{1}{\sqrt{MN}} \sum_{i=1}^N \sum_{j=1}^{\tilde{N}} g_j^i P_j^i, \quad (55)$$

where the projectors are as in Eq. (3) with the sum running instead to $d_m = \binom{M}{m}$, the number of states per site. We have included random couplings g_j^i and allow for $N \neq \tilde{N}$.

The generalized Hamiltonian $H_{M,m}$ remains a sum of positive semidefinite operators, so the nullspace still consists of those states annihilated by each two-site term individually. The calculation of the nullspace dimension thus proceeds as in Sec. III. The clean model Hamiltonian is written in terms of Casimirs as

$$H_{M,m}^{\text{clean}} = \frac{m^2(N^2 + \tilde{N}^2)}{2M} + \frac{1}{2} (C_{\mathbf{A}_m^{\otimes N}} + C_{\overline{\mathbf{A}}_m^{\otimes \tilde{N}}} - C_{\mathbf{A}_m^{\otimes N} \otimes \overline{\mathbf{A}}_m^{\otimes \tilde{N}}})$$

just like Eq. (25). The tensor products can be decomposed into irreducible representations $\lambda \in \mathbf{A}_m^{\otimes N}$, $\bar{\gamma} \in \overline{\mathbf{A}}_m^{\otimes \tilde{N}}$ and $\rho \in \gamma \otimes \lambda$. One then finds as in Eq. (31) that the ground states are states where $\rho = \lambda + \bar{\gamma}$ and $k + l \leq M$, with Young diagrams for λ and γ having k and l rows, respectively.

For $m \neq M/2$, similar arguments to those for $m = 1$ show that the dimension of the nullspace grows exponentially with N . For $m = M/2$, the model is equivalent to the SY model with spins in the representation $\mathbf{A}_{M/2}$. We thus expect no Hilbert-space fragmentation. Indeed, all but one of the possible λ have more than $M/2$ rows, and likewise for γ . Hence there is only one $SU(M)$ representation in the nullspace, that with a rectangular Young diagram with $m = M/2$ rows and $N + \tilde{N}$ columns. It has multiplicity one in the tensor products, so fragmentation does not occur.

B. Field-theory description at large M, m

Here we utilize field-theory techniques [15,31] to show how some of the physics closely resembles that of the SY and SYK models. We study the Hamiltonian $H_{M,m}$ from Eq. (55) with spins in antisymmetric representations. Here we set $\tilde{N}/N = b$, and $m/M = c$ with b and c order 1.

We represent the spins in the $\mathbf{A}_m$ and $\overline{\mathbf{A}}_m$ representations using fermionic operators $f^i, \tilde{f}_j$:

$$\begin{aligned} \mathcal{S}_b^{a,i} &= f^{a,i\dagger} f_b^i; & \sum_{a=1}^M f^{a,i\dagger} f_a^i &= m, \\ \tilde{\mathcal{S}}_{b,j}^a &= \tilde{f}_j^{a\dagger} \tilde{f}_{b,j}; & \sum_{a=1}^M \tilde{f}_j^{a\dagger} \tilde{f}_{a,j} &= M - m, \end{aligned} \quad (56)$$

where the fermions obey standard anticommutation relations: $\{f^{a,i}, f^{a,i\dagger}\} = \{\tilde{f}_{b,j}, \tilde{f}_{b,j}^\dagger\} = 1$, and all other anticommutators vanish. The Hamiltonian (55) written in terms of spins contains interactions of the SY form between the two groups of spins

$$H = \frac{1}{\sqrt{MN}} \sum_{i=1}^N \sum_{j=1}^{\tilde{N}} \sum_{a,b=1}^M g_j^i \mathcal{S}_b^{a,i} \tilde{\mathcal{S}}_{a,j}^b. \quad (57)$$

The coefficients g_j^i are independent and identically distributed Gaussian random variables with $\overline{g_j^i} = 0$ and $\overline{(g_j^i)^2} = g^2$.

We introduce the onsite (and diagonal in $SU(M)$ index) Green's functions at $T = 0$:

$$G(\tau) = -T_\tau \langle f^{a,i}(\tau) f^{a,i\dagger}(0) \rangle, \quad \tilde{G}(\tau) = -T_\tau \langle \tilde{f}_{a,j}(\tau) \tilde{f}_{a,j}^\dagger(0) \rangle. \quad (58)$$

That we calculate disorder-averaged quantities is implicit throughout this section; whether the quantities of interest are well-represented by their disorder averages is a separate question, and one which we shall not confront. One can calculate these Green's functions by writing the Hamiltonian of Eq. (57) in terms of fermions using Eq. (56), with chemical potentials $\mu, \tilde{\mu}$ to fix the respective fillings. We then observe that the diagrams which do not vanish in the large N, M limit are of the melon type, just like in the SYK model. The Schwinger-Dyson equations are

$$G(i\omega) = \frac{1}{i\omega + \mu - \Sigma(i\omega)}, \quad \tilde{G}(i\omega) = \frac{1}{i\omega + \tilde{\mu} - \tilde{\Sigma}(i\omega)}, \quad (59)$$

with

$$\Sigma(\tau) = \text{---} \text{---} \text{---} \text{---} = -U^2 G(\tau) \tilde{G}(\tau) \tilde{G}(-\tau), \quad (60)$$

where $U^2 = bg^2/2$ and

$$\tilde{\Sigma}(\tau) = \text{---} \text{---} \text{---} \text{---} = -\tilde{U}^2 \tilde{G}(\tau) G(\tau) G(-\tau), \quad (61)$$

where $\tilde{U}^2 = g^2/2$. In the above diagrams, solid and dashed lines denote the propagators G and $\tilde{G}$. The disorder average which couples the two vertices is implicit. Finally, the filling constraints of Eqs. (56) translate to

$$G(\tau = 0^-) = c, \quad \tilde{G}(\tau = 0^-) = 1 - c. \quad (62)$$

All other propagators are smaller by factors of N, M relative to these, and so will be ignored. Equations with a similar structure have appeared in the solution of a bipartite Majorana SYK model in Ref. [17] and an all-to-all $t - J$ model in Ref. [32].

Equations (59) through (62) in general need to be solved numerically. However, in looking for solutions in the $\omega \rightarrow 0$ (consequently $|\tau| \rightarrow \infty$) limit some further progress can be made. One starts by guessing SYK-like Green's functions in the upper half frequency plane ($\text{Im } z > 0$):

$$G(z) = C \frac{e^{-i(\pi\Delta+\theta)}}{z^{1-2\Delta}}, \quad \tilde{G}(z) = \tilde{C} \frac{e^{-i(\pi\tilde{\Delta}+\tilde{\theta})}}{z^{1-2\tilde{\Delta}}}. \quad (63)$$

These allow us to find the corresponding spectral functions, which yield, via Laplace transforms, $G(\tau)$ and $\tilde{G}(\tau)$:

$$G(\tau) = \begin{cases} -\frac{C\Gamma(2\Delta)}{\pi} \frac{\sin(\pi\Delta+\theta)}{|\tau|^{2\Delta}}, & \tau > 0, \\ \frac{C\Gamma(2\tilde{\Delta})}{\pi} \frac{\sin(\pi\tilde{\Delta}-\theta)}{|\tau|^{2\tilde{\Delta}}}, & \tau < 0, \end{cases} \quad (64)$$

and an identical expression with tildes for $\tilde{G}(\tau)$. Defining

$$\alpha_\pm = \frac{C\Gamma(2\Delta)}{\pi} \sin(\pi\Delta \pm \theta) \quad (65)$$

(and similar for $\tilde{\alpha}_\pm$) Eqs. (60) and (61) become, respectively,

$$\Sigma(\tau) = \begin{cases} -U^2 \alpha_+ \tilde{\alpha}_+ \tilde{\alpha}_- |\tau|^{-2\Delta-4\tilde{\Delta}}, & \tau > 0, \\ U^2 \alpha_- \tilde{\alpha}_+ \tilde{\alpha}_- |\tau|^{-2\Delta-4\tilde{\Delta}}, & \tau < 0, \end{cases} \quad (66)$$

and its tilded counterpart.

Again using the inverse Laplace transforms as an intermediate object, we can find $\Sigma(i\omega)$ and $\tilde{\Sigma}(i\omega)$. Then, assuming that the small-frequency behavior of the denominators in Eq. (59) is dominated by the self energies, equating the powers of ω in the same equations gives $1 - 2\Delta = 2\tilde{\Delta} + 4\tilde{\Delta} - 1$, i.e.,

$$\Delta + \tilde{\Delta} = \frac{1}{2}, \quad (67)$$

and equating the coefficients then gives

$$-\frac{1}{C} = \frac{\Gamma(2\Delta)\Gamma^2(2\tilde{\Delta})}{2\pi^3\Gamma(2\Delta+4\tilde{\Delta})} C \tilde{C}^2 U^2 (\cos 2\pi\tilde{\Delta} - \cos 2\pi\tilde{\theta}) \quad (68)$$

and the tilded version. Dividing these two equations, one arrives at

$$\frac{\tilde{U}^2}{U^2} = \frac{1}{b} = \frac{2\Delta}{1-2\Delta} \frac{\cos 2\tilde{\theta} + \cos 2\pi\Delta}{\cos 2\theta - \cos 2\pi\Delta}. \quad (69)$$

Setting $\theta = \tilde{\theta} = 0$, which enforces particle-hole symmetry, one recovers the equation for Δ in the bipartite SYK (i.e., quartic-in-Majoranas) model studied in Ref. [17]. Finally, we have equations that fix the fillings in terms of the parameters that enter the Green's functions [31,33]:

$$c = \frac{1}{2} - \frac{\theta}{\pi} - \left(\frac{1}{2} - \Delta \right) \frac{\sin 2\theta}{\sin 2\pi\Delta}, \quad (70)$$

$$1 - c = \frac{1}{2} - \frac{\tilde{\theta}}{\pi} - \Delta \frac{\sin 2\tilde{\theta}}{\sin 2\pi\Delta}. \quad (71)$$

Note that these ‘‘Luttinger relations’’ cannot be found by substituting the Green's functions of Eq. (64) (which are valid only at large $|\tau|$) into Eq. (62) (which concerns the $\tau \rightarrow 0$ behavior of the true Green's functions). Equations (69), (70), and (71) are three equations in the variables $\Delta, \theta, \tilde{\theta}$, and may be solved numerically.

We should stress that the conformal *ansatz*, while demonstrably a valid solution to the Schwinger-Dyson equations as $\omega \rightarrow 0$, is not guaranteed to be associated with any solution that is valid at higher energies as well. (See, e.g., Ref. [34] for a treatment of this issue in the SYK context.) In this sense the approach described here does not represent a complete solution for the Green's function even in the $N, M \rightarrow \infty$ limit.

We expect that the states in the nullspace, the focus of the rest of this paper, are invisible to the solutions of the Schwinger-Dyson equations discussed here, and would also be invisible to their finite-temperature version—even if analytic continuation to real time can be performed. This is because these states are exponentially (in N) rare in the spectrum, and so taking the $N \rightarrow \infty$ limit (which one has to do for any analytical control) will erase their contribution. While $1/M$ corrections may be possible to treat, and are likely to indicate a spin-glass state [35], the $N \rightarrow \infty$ limit must be taken first in such calculations; we are not aware of a way to

calculate, say, $1/N$ corrections at large but finite M . Furthermore, all the quantities one can calculate in the field-theoretic approach are disorder-averaged. It is not clear *a priori*, even at finite N , if disorder-averaged correlation functions should be expected to inherit any nontrivial features from the nonergodicity of the Hamiltonian.

VI. SCARS FROM THE SHIRAISHI-MORI CONSTRUCTION

We have established that each eigenstate in the nullspace of H_M does not mix with the others under time evolution. As such, it is tempting to label these eigenstates as quantum many-body scars [20]. The numerous product eigenstates identified in Sec. II B have area-law entanglement entropy scaling, a signal of a scar. However, typically a scar is a rare ETH-violating exception in an otherwise thermal spectrum. Here the situation is rather different, as states in certain $SU(M)$ representations are in the nullspace, while all others are not.

We show in this section how to perturb H_M to yield more conventional scars. We utilize the Shiraishi-Mori (SM) embedding formalism [18] to break the $SU(M)$ symmetry, converting some of the original nullspace eigenstates into quantum many-body scars while also maintaining Hilbert space fragmentation. To do so, we must address subtleties involved in applying this formalism to nonlocal models.

A quantum many-body scar for a local Hamiltonian violates ETH. Since global symmetries also constrain time evolution in the obvious fashion, we must split the Hilbert space into sectors $\mathcal{H}_n$, labeled by the corresponding quantum numbers. Defining $\mathcal{H}_{\text{thermal},n}$ to be largest Krylov subspace of $\mathcal{H}_n$, two conditions are sufficient for an eigenstate $|\psi_{\text{scar}}\rangle$ to be a quantum many-body scar [20], namely

$$|\psi_{\text{scar}}\rangle\langle\psi_{\text{scar}}| \in \mathcal{C}, \quad \frac{\dim(\mathcal{H}_{\text{thermal},n})}{\dim(\mathcal{H}_n)} \xrightarrow{N \rightarrow \infty} 1, \quad (72)$$

where $\mathcal{C}$ is a nontrivial commutant algebra. The first condition ensures that the scar state violates ETH, since it implies that the scar state is an eigenstate of a family of Hamiltonians, whereas ETH-obeying eigenstates generically have a unique parent Hamiltonian [20,36,37]. The second condition ensures that the scar state is a rare exception in a spectrum dominated by thermal eigenstates. We note that $|\psi_{\text{scar}}\rangle$ need not have area-law entanglement entropy [20,38]. Additionally, if the number of scars in a model grows exponentially with N , then by Eq. (72) the dimension of the commutant algebra grows exponentially as well, leading to Hilbert-space fragmentation. The fragmentation in this case is weak, because the second condition of Eq. (72) says that the spectrum is dominated by one large thermal Krylov subspace.

Following Refs. [18,20], we perturb H_M as

$$H_{\text{SM}} = \sum_{i,j=1}^N P_j^i h_{[j]}^{[i]} P_j^i + H_{\text{pert}}, \quad (73)$$

where the $h_{[j]}^{[i]}$ are finite-support operators that are sufficiently general to break the $SU(M)$ symmetry, and H_{pert} is chosen

to split the degeneracies between the nullspace eigenstates. Each operator $h_{[j]}^{[i]}$ need not have support on the spins i and j , so these indices merely serve as labels. The original Hamiltonian H_M from Eq. (5) corresponds to setting $h_{[j]}^{[i]} = r_j^i \mathbb{1}$ and $H_{\text{pert}} = 0$.

Because the $SU(M)$ symmetry of H_M is broken, the perturbed Hamiltonian H_{SM} acquires a dominant thermal subspace, and hence satisfies the second condition of Eq. (72). Furthermore, when $H_{\text{pert}} = 0$, the nullspaces of H_{SM} and H_M are the same and so states in it satisfy the first condition of Eq. (72) as well. However, satisfying these conditions does not automatically guarantee that states in the nullspace of a nonlocal model are quantum many-body scars. This is because a thermal eigenstate is generally understood to uniquely specify a parent Hamiltonian only for spatially local systems [36,37]. In the absence of locality, we thus cannot conclude that the first condition is thus sufficient to imply that an eigenstate violates ETH.

However, the product eigenstates in the nullspace identified in Sec. II B have area-law scaling and so violate ETH. Moreover, we can choose H_{pert} such that the product eigenstates of H_M are split from the remainder of the nullspace. Namely, we perturb by the diagonal onsite $U(M)$ generators defined in Eq. (1) via

$$H_{\text{pert}} = \sum_{\alpha=1}^M \left(\sum_{i=1}^N c^{\alpha,i} T_{\alpha}^{\alpha,i} + \sum_{j=1}^N d_j^{\alpha} S_{\alpha,j}^{\alpha} \right), \quad (74)$$

where the $c^{\alpha,i}$ and d_j^{α} are arbitrary distinct real coefficients. The product states from $\mathcal{N}$ remain eigenstates of H_{SM} , and now are situated in quantum-number sectors of H_{SM} that have a dominant thermal Krylov subspace. They thus satisfy the conditions in Eq. (72), and furthermore violate ETH due to their area-law entanglement entropy scaling. The product states hence constitute quantum many-body scars of H_{SM} . In particular, when the coefficients $c^{\alpha,i}$ and d_j^{α} are rational, any initial state composed of a sum of these product states would exhibit revivals when time evolved with H_{SM} , a hallmark of scars [3]. Additionally, since the number of such product states grows exponentially with N , the perturbed model H_{SM} has an exponentially growing number of quantum many-body scars and is Hilbert space fragmented.

VII. CONCLUDING REMARKS

The nonlocal model introduced in this paper possesses a number of remarkable properties. In spite of its all-to-all interactions, the dimension of its nullspace grows exponentially with increasing system size. Furthermore, maps between eigenstates in this nullspace are elements of the commutant algebra. The exponentially growing degeneracy thus leads to Hilbert-space fragmentation in some of the $SU(M)$ quantum number sectors of the spectrum, violating ETH in a weak sense. Much like the SY model, our model can in appropriate limits be attacked by field-theoretic methods, at least for disorder-averaged low-energy observables. However these techniques cannot access the special states in the nullspace. The Shiraishi-Mori embedding formalism can be utilised to

perturb many of the eigenstates in the nullspace into quantum many-body scars.

Nonlocal models with Hilbert-space fragmentation are rarer in the literature than those with quantum many-body scars. Nonlocal models have a large bond algebra: for example, the number of generators of the bond algebra of H_M grows quadratically with N , as opposed to the linear growth associated with a 1D spin chain. This strongly constrains the dimension of the commutant algebra, making fragmentation less likely.

Previous works have considered how scars and fragmentation can be stable against the addition of longer-range interactions to local Hamiltonians [39–44]. However, the model studied here is apparently a rare example of one where the nonlocal terms are the leading contribution to the problem, and cannot be ascribed a notion of spatial distance: it is intrinsically “zero dimensional.” It may be of interest to explore other problems of that nature which share some features of our model, for instance the model of Ref. [45] which is also SYK-like and has extensive degeneracies in its eigenspectrum. We leave this question for the future.

ACKNOWLEDGMENTS

We thank Sounak Biswas, Maine Christos, and Pablo Sala for helpful discussions. This work was supported by the European Research Council under the European Union Horizon 2020 Research and Innovation Programme via Grant Agreement No. 804213-TMCS (S.A.P.) and by the Fonds de recherche du Québec via Doctoral Level Grant No. 302311 (J.C.-H.). We acknowledge support from EPSRC via Grant No. EP/S020527/1.

APPENDIX: LIMITING FORM FOR M LARGE, N FIXED

In this Appendix, we show that for $M \geq 2N$, the Hilbert space is dominated by the nullspace, as displayed in Eq. (38):

$$\lim_{M \rightarrow \infty} \frac{D_{\mathcal{N}}}{M^{2N}} = 1.$$

The proof follows from two facts: (1) for all states in the nullspace as given in Eq. (31), the dimension of the corresponding $SU(M)$ multiplet grows as M^{2N} for fixed N , while (2) all other irreducible representations have dimension growing as a lower power of M .

We first note that $k + \ell \leq 2N < M$ when $M > 2N$. Since the restriction in the sum in Eq. (32) is always satisfied, the nullspace dimension simplifies to

$$D_{\mathcal{N}} = \sum_{k, \ell=1}^N \sum_{\lambda \in P_{N,k}} \sum_{\gamma \in P_{N,\ell}} m(\lambda) m(\gamma) d(\lambda + \bar{\gamma}). \quad (\text{A1})$$

The only dependence on M here is through the $SU(M)$ dimension $d(\rho)$. The formula (34) for this dimension then can be factored into two pieces by splitting this product into one piece involving the first k rows of the Young diagram for ρ , and another involving the other rows. The latter part of this diagram can be thought of as a separate

Young diagram with entries $\rho^{(M-k)} = [\rho_{k+1}, \rho_{k+2}, \dots, \rho_M] = [\bar{\gamma}_{k+1}, \bar{\gamma}_{k+2}, \dots, \bar{\gamma}_M]$. The corresponding contributions of this piece to $d(\rho)$ then amount to the dimension $d^{(M-k)}$ of the representation $\rho^{(M-k)}$ in the group $SU(M-k)$, yielding

$$d(\rho) = d^{(M-k)}(\rho^{(M-k)}) \frac{\prod_{a=1}^k (\rho_a + M - a)!}{\prod_{a=1}^k ((M-a)! \prod_{b=1}^{\rho_a} h_{a,b})}. \quad (\text{A2})$$

The resulting large- M behavior is straightforward to determine. We have

$$\prod_{a=1}^k \frac{(\rho_a + M - a)!}{(M-a)!} = \prod_{a=1}^k \frac{(\lambda_a + \gamma_1 + M - a)!}{(M-a)!} \xrightarrow{M \text{ large}} M^{N+k\gamma_1}, \quad (\text{A3})$$

where we exploit the facts that $\sum_{a=1}^k \lambda_a = N$ and $\bar{\gamma}_a = \gamma_1$ for $a \leq M-l$. Because of the latter fact along with $\gamma \in P_{N,\ell}$, the first γ_1 columns in the Young diagram of $\bar{\gamma}$ and hence ρ must be at least of length $M-l$. The remaining columns arise only from λ , and so the corresponding hook lengths are independent of M , yielding

$$\prod_{a=1}^k \prod_{b=1}^{\rho_a} h_{a,b} \xrightarrow{M \text{ large}} M^{k\gamma_1} h(\lambda). \quad (\text{A4})$$

Finally, we use the fact the dimensions of an $SU(M-k)$ representation and its conjugate are the same to derive

$$d^{(M-k)}(\rho^{(M-k)}) = d^{(M-k)}(\overline{\rho^{(M-k)}}) = d^{(M-k)}(\gamma).$$

Here the hook lengths are independent of M , yielding

$$d^{(M-k)}(\rho^{(M-k)}) = \frac{1}{h(\gamma)} \prod_{a=1}^{\ell} \frac{(\gamma_a + (M-k) - a)!}{(M-k-a)!} \xrightarrow{M \text{ large}} M^N \frac{1}{h(\gamma)}. \quad (\text{A5})$$

Putting the above results together, we see that

$$d(\lambda + \bar{\gamma}) \xrightarrow{M \text{ large}} \frac{1}{h(\lambda)h(\gamma)} M^{2N}. \quad (\text{A6})$$

We proved that states in any $SU(M)$ representation other than $\rho = \lambda + \bar{\gamma}$ are not in the nullspace. For $M \geq 2N$, these states must be in representations $\rho \in \lambda \otimes \bar{\gamma}$ in the tensor product, as there are no restrictions on $\lambda \in P_{N,k}$ and $\gamma \in P_{N,\ell}$. Since $\bar{\gamma}_a = \gamma_1$ for $a \leq M-l$ and $k < M-l$, the inequality (27) means that at least one box in λ must move downward when forming ρ . Moreover, it must move to a row a with $a > M - \ell + 1$, as boxes originally in the same row cannot end up in the same column after taking the tensor product. We thus have $\sum_{i=1}^k \rho_a < N + k\gamma_1$, as opposed to the equality used in the proof of Eq. (A3). The number of boxes in $\rho^{(M-k)}$ must therefore increase, so that $d^{(M-k)}(\rho^{(M-k)})$ must decrease relative to Eq. (A5). Meanwhile, M in Eq. (A4) is unchanged. Thus, $d(\rho)$ for $\rho \neq \lambda + \bar{\gamma}$ increases at large M at most with a power M^{2N-2} .

We have shown that the only $SU(M)$ representations with dimension growing as M^{2N} that appear are those in the

nullspace. Since the total dimension of the Hilbert space is M^{2N} , the relation (38) must therefore hold. An amusing by-product of this derivation is the identity

$$\sum_{k=0}^N \sum_{\lambda \in P_{N,k}} \frac{1}{h(\lambda)^2} = \frac{1}{N!}. \quad (\text{A7})$$

The proof of Eq. (A7) comes from noting that combining Eqs. (A6) and (38) yields

$$\sum_{k,l=0}^N \sum_{\lambda \in P_{N,k}} \sum_{\gamma \in P_{N,l}} \frac{m(\lambda)m(\gamma)}{h(\lambda)h(\gamma)} = 1$$

and then using the first part of Eq. (33).

-
- [1] S. Moudgalya, B. A. Bernevig, and N. Regnault, Quantum many-body scars and Hilbert space fragmentation: A review of exact results, *Rep. Prog. Phys.* **85**, 086501 (2022).
- [2] S. Moudgalya and O. I. Motrunich, Hilbert space fragmentation and commutant algebras, *Phys. Rev. X* **12**, 011050 (2022).
- [3] M. Serbyn, D. A. Abanin, and Z. Papić, Quantum many-body scars and weak breaking of ergodicity, *Nat. Phys.* **17**, 675 (2021).
- [4] Z. Papić, Weak ergodicity breaking through the lens of quantum entanglement, *arXiv:2108.03460*.
- [5] A. Chandran, T. Iadecola, V. Khemani, and R. Moessner, Quantum many-body scars: A quasiparticle perspective, *Annu. Rev. Condens. Matter Phys.* **14**, 443 (2023).
- [6] J. M. Deutsch, Eigenstate thermalization hypothesis, *Rep. Prog. Phys.* **81**, 082001 (2018).
- [7] R. Nandkishore and D. A. Huse, Many-body localization and thermalization in quantum statistical mechanics, *Annu. Rev. Condens. Matter Phys.* **6**, 15 (2015).
- [8] P. Sala, T. Rakovszky, R. Verresen, M. Knap, and F. Pollmann, Ergodicity breaking arising from Hilbert space fragmentation in dipole-conserving Hamiltonians, *Phys. Rev. X* **10**, 011047 (2020).
- [9] T. Rakovszky, P. Sala, R. Verresen, M. Knap, and F. Pollmann, Statistical localization: From strong fragmentation to strong edge modes, *Phys. Rev. B* **101**, 125126 (2020).
- [10] S. Moudgalya and O. I. Motrunich, Numerical methods for detecting symmetries and commutant algebras, *Phys. Rev. B* **107**, 224312 (2023).
- [11] A. Khudorozhkov, A. Tiwari, C. Chamon, and T. Neupert, Hilbert space fragmentation in a 2D quantum spin system with subsystem symmetries, *SciPost Phys.* **13**, 098 (2022).
- [12] M. T. Batchelor and M. N. Barber, Spin- s quantum chains and Temperley-Lieb algebras, *J. Phys. A: Math. Gen.* **23**, L15 (1990).
- [13] B. Aufgebauer and A. Klümper, Quantum spin chains of Temperley-Lieb type: Periodic boundary conditions, spectral multiplicities and finite temperature, *J. Stat. Mech.: Theory Exp.* (2010) P05018.
- [14] N. Read and H. Saleur, Enlarged symmetry algebras of spin chains, loop models, and S-matrices, *Nucl. Phys. B* **777**, 263 (2007).
- [15] S. Sachdev and J. Ye, Gapless spin-fluid ground state in a random quantum Heisenberg magnet, *Phys. Rev. Lett.* **70**, 3339 (1993).
- [16] A. Georges, O. Parcollet, and S. Sachdev, Quantum fluctuations of a nearly critical Heisenberg spin glass, *Phys. Rev. B* **63**, 134406 (2001).
- [17] M. Fremling, M. Haque, and L. Fritz, Bipartite Sachdev-Ye-Kitaev model: Conformal limit and level statistics, *Phys. Rev. D* **105**, 066017 (2022).
- [18] N. Shiraishi and T. Mori, Systematic construction of counterexamples to the eigenstate thermalization hypothesis, *Phys. Rev. Lett.* **119**, 030601 (2017).
- [19] S. Moudgalya and O. I. Motrunich, From symmetries to commutant algebras in standard Hamiltonians, *Ann. Phys.* **455**, 169384 (2023).
- [20] S. Moudgalya and O. I. Motrunich, Exhaustive characterization of quantum many-body scars using commutant algebras *arXiv:2209.03377*.
- [21] J. Sonner and M. Vielma, Eigenstate thermalization in the Sachdev-Ye-Kitaev model, *J. High Energy Phys.* **11** (2017) 149.
- [22] M. Haque and P. A. McClarty, Eigenstate thermalization scaling in Majorana clusters: From chaotic to integrable Sachdev-Ye-Kitaev models, *Phys. Rev. B* **100**, 115122 (2019).
- [23] B. E. Sagan, *The Symmetric Group: Representations, Combinatorial Algorithms, and Symmetric Functions*, 2nd ed., Graduate Texts in Mathematics (Springer, New York, NY, 2001).
- [24] M. E. Beverland, G. Alagic, M. J. Martin, A. P. Koller, A. M. Rey, and A. V. Gorshkov, Realizing exactly solvable $SU(N)$ magnets with thermal atoms, *Phys. Rev. A* **93**, 051601(R) (2016).
- [25] M. A. Perlin, D. Barberena, M. Mamaev, B. Sundar, R. J. Lewis-Swan, and A. M. Rey, Engineering infinite-range $SU(N)$ interactions with spin-orbit-coupled fermions in an optical lattice, *Phys. Rev. A* **105**, 023326 (2022).
- [26] D. Jakab, G. Szirmai, and Z. Zimborás, The bilinear-biquadratic model on the complete graph, *J. Phys. A: Math. Theor.* **51**, 105201 (2018).
- [27] D. Jakab and Z. Zimborás, Quantum phases of collective $SU(3)$ spin systems with bipartite symmetry, *Phys. Rev. B* **103**, 214448 (2021).
- [28] A. M. Perelomov and V. S. Popov, Casimir operators for semisimple Lie groups, *Mathematics of the USSR-Izvestiya* **2**, 1313 (1968).
- [29] M. Hamermesh, *Group Theory and Its Application to Physical Problems*, Addison Wesley Series in Physics (Dover Publications, Oxford, UK, 1989).
- [30] H. Schlosser, A closed formula for the decomposition of the Kronecker product of irreducible representations of $SU(N)$, *Mathematische Nachrichten* **134**, 237 (1987).
- [31] D. Chowdhury, A. Georges, O. Parcollet, and S. Sachdev, Sachdev-Ye-Kitaev models and beyond: Window into non-Fermi liquids, *Rev. Mod. Phys.* **94**, 035004 (2022).
- [32] M. Christos, D. G. Joshi, S. Sachdev, and M. Tikhonovskaya, Critical metallic phase in the overdoped random t - J model, *Proc. Natl. Acad. Sci. USA* **119**, e2206921119 (2022).
- [33] Y. Gu, A. Kitaev, S. Sachdev, and G. Tarnopolsky, Notes on the complex Sachdev-Ye-Kitaev model, *J. High Energy Phys.* **02** (2020) 157.

- [34] T. Azeyanagi, F. Ferrari, and F. I. Schaposnik Massolo, Phase diagram of planar matrix quantum mechanics, tensor, and Sachdev-Ye-Kitaev models, *Phys. Rev. Lett.* **120**, 061602 (2018).
- [35] M. Christos, F. M. Haehl, and S. Sachdev, Spin liquid to spin glass crossover in the random quantum Heisenberg magnet, *Phys. Rev. B* **105**, 085120 (2022).
- [36] J. R. Garrison and T. Grover, Does a single eigenstate encode the full Hamiltonian? *Phys. Rev. X* **8**, 021026 (2018).
- [37] X.-L. Qi and D. Ranard, Determining a local Hamiltonian from a single eigenstate, *Quantum* **3**, 159 (2019).
- [38] C. M. Langlett, Z.-C. Yang, J. Wildeboer, A. V. Gorshkov, T. Iadecola, and S. Xu, Rainbow scars: From area to volume law, *Phys. Rev. B* **105**, L060301 (2022).
- [39] A. Lerose, T. Parolini, R. Fazio, D. A. Abanin, and S. Pappalardi, Theory of robust quantum many-body scars in long-range interacting systems, [arXiv:2309.12504](https://arxiv.org/abs/2309.12504).
- [40] D. T. Stephen, O. Hart, and R. M. Nandkishore, Ergodicity breaking provably robust to arbitrary perturbations, *Phys. Rev. Lett.* **132**, 040401 (2024).
- [41] G. Francica and L. Dell'Anna, Hilbert space fragmentation in a long-range system, *Phys. Rev. B* **108**, 045127 (2023).
- [42] G. Zhang and Z. Song, Quantum scars in spin-isotropic Heisenberg clusters, *New J. Phys.* **25**, 053025 (2023).
- [43] K. Pakrouski, P. N. Pallegar, F. K. Popov, and I. R. Klebanov, Many-body scars as a group invariant sector of Hilbert space, *Phys. Rev. Lett.* **125**, 230602 (2020).
- [44] K. Pakrouski, P. N. Pallegar, F. K. Popov, and I. R. Klebanov, Group theoretic approach to many-body scar states in fermionic lattice models, *Phys. Rev. Res.* **3**, 043156 (2021).
- [45] E. Iyoda, H. Katsura, and T. Sagawa, Effective dimension, level statistics, and integrability of Sachdev-Ye-Kitaev-like models, *Phys. Rev. D* **98**, 086020 (2018).