

Automorphisms of Automorphism Groups of Free Groups

Martin R. Bridson¹

Mathematical Institute, 24-29 St Giles', Oxford OX1 3LB, United Kingdom
E-mail: bridson@maths.ox.ac.uk

and

Karen Vogtmann²

Mathematics Department, 555 Malott Hall, Cornell University, Ithaca, New York 14850
E-mail: vogtmann@math.cornell.edu

Communicated by Peter M. Neumann

Received November 19, 1999

If $n \geq 3$ and F_n is free of rank n , then $\text{Out}(\text{Aut}(F_n)) = \text{Out}(\text{Out}(F_n)) = \{1\}$.

© 2000 Academic Press

1. INTRODUCTION

There is a well-developed analogy between lattices in semisimple Lie groups on the one hand and mapping class groups and automorphism groups of free groups on the other. A recently developed part of this analogy involves generalizations of the theorem of Tits [9] which states that all automorphisms of a spherical building stem from the underlying algebraic group. Specifically, Ivanov [5] proved that the natural map from the mapping class group to the group of automorphisms of the curve complex is an isomorphism, and the present authors proved that the natural map from $\text{Out}(F_n)$ to the group of automorphisms of Outer space is an isomorphism. Tits's theorem leads to a proof that the outer automor-

¹Research supported by an EPSRC Advanced Fellowship.

²Research supported in part by NSF Grant DMS-9307313.

phism group of any irreducible lattice in higher rank is finite. In the same way, Ivanov's theorem implies that the outer automorphism group of the mapping class group is finite. In the case of $\text{Out}(F_n)$ we have been unable to identify a direct implication of the same type. However, in this paper we shall show that some of the ideas that we used in [1] to analyze the symmetries of Outer space can also be used to prove:

THEOREM. *For $n \geq 3$, every automorphism of $\text{Out}(F_n)$ is inner.*

The same techniques show that all automorphisms of $\text{Aut}(F_n)$ are inner. In [4] Dyer and Formanek give a very different proof that $\text{Out}(\text{Aut}(F_n))$ is trivial. Their proof relies on the following classical result of Burnside: if the center of G is trivial, then $\text{Out}(\text{Aut}(G)) = \{1\}$ if and only if $\text{Inn}(G) \subset \text{Aut}(G)$ is a characteristic subgroup.

Outline of the Proof

We shall assume throughout this paper that $n \geq 3$. We shall prove the theorem by considering the action of $\text{Out}(F_n)$ on Outer space. (The corresponding result for $\text{Aut}(F_n)$ is proved by an entirely similar analysis of the action of $\text{Aut}(F_n)$ on Auter space.) The relevant facts and vocabulary concerning the geometry of Outer space can be found in [3].

We first consider roses in Outer space. The stabilizers of roses are characterized among the finite subgroups of $\text{Out}(F_n)$ by their abstract isomorphism type, and hence there is a natural action of $\text{Aut}(\text{Out}(F_n))$ on the set of such stabilizers. Since $\text{Out}(F_n)$ acts transitively on the set of roses, $\text{Inn}(\text{Out}(F_n))$ acts transitively on the set of stabilizers of roses. Thus, given an automorphism ϕ of $\text{Out}(F_n)$, we may adjust ϕ within its outer automorphism class to ensure that ϕ leaves the stabilizer W_n of the standard rose invariant. After further adjustment, we may assume that ϕ induces one of the small number of possible outer automorphisms of W_n (Lemma 2.1). An analysis of the action of ϕ on the stabilizers of the graphs adjacent to the standard rose enables us to conclude that in fact the restriction of ϕ to W_n is the identity (Lemma 3.3). We now observe that $\text{Out}(F_n)$ is generated by W_n together with one other automorphism r , which arises as a symmetry of a certain graph in the link of the standard rose. By analyzing the stabilizer of this graph we shall see that there are only a small number of possibilities for $\phi(r)$. A further analysis shows that each of the putative images $\phi(r) \neq r$ leads to inconsistent relations in $\text{Out}(F_n)$. Thus we conclude that $\phi(r) = r$ and hence $\phi = 1$ in $\text{Out}(\text{Out}(F_n))$.

2. AUTOMORPHISMS OF THE STABILIZER OF A ROSE

Fix an ordered basis $\{x_1, \dots, x_n\}$ of F_n . For $i = 1, \dots, n$, let e_i be the automorphism of F_n that sends x_i to x_i^{-1} and fixes the other basis elements. For $i = 1, \dots, n-1$, let τ_i be the automorphism that interchanges x_i and x_{i+1} while leaving the other basis elements fixed. Let $W_n \cong (\mathbb{Z}/2)^n \rtimes S_n$ be the subgroup of $\text{Aut}(F_n)$ generated by the e_i and the τ_i . The group $\text{Aut}(F_n)$ (and therefore $\text{Out}(F_n)$) is generated by W_n together with the automorphism u which sends $x_1 \mapsto x_1 x_2$ and fixes x_i for $i > 1$ (see [8], p. 163).

There is a faithful representation $W_n \rightarrow GL(n, \mathbb{Z})$ whose image consists of all matrices of the form PD , where P is a permutation matrix and D is a diagonal matrix. The center of W_n has order 2; it is generated by $z = \prod_{i=1}^n e_i$, which is represented by the matrix $-I$.

Define $\alpha: W_n \rightarrow W_n$ by $\alpha(PD) = (\det P)PD$, and $\beta: W_n \rightarrow W_n$ by $\beta(PD) = (\det D)PD$. It is easy to check that α is an automorphism for all n , and β is an automorphism if and only if n is even. Both α and β have order 2, as does their product $\alpha\beta$. None of α , β , or $\alpha\beta$ is inner, since an inner automorphism would preserve traces, while $\text{tr}(\tau_1) \neq \text{tr}(\alpha(\tau_1))$, $\text{tr}(e_1) \neq \text{tr}(\beta(e_1))$, and $\text{tr}(\tau_1) \neq \text{tr}(\alpha\beta(\tau_1))$.

We are grateful to Peter Neumann for his comments on the following lemma.

LEMMA 2.1. *If n is odd, $\text{Out}(W_n) \cong \mathbb{Z}/2$ is generated by α . If n is even, $\text{Out}(W_n) \cong \mathbb{Z}/2 \times \mathbb{Z}/2$ is generated by α and β .*

Proof. Let $N = (\mathbb{Z}/2)^n$ be the normal subgroup generated by the e_i . We claim that N is a characteristic subgroup of W_n . To see this, note that if $A = \phi(N) \neq N$ for some automorphism ϕ of W_n , then A would have non-trivial image in the quotient $S_n = W_n/N$. Since the image would be a normal 2-group this is impossible unless $n = 4$ and the image is the Klein 4-group V . But the action of V on N has z as its only fixed non-zero vector, contradicting the fact that the intersection of the abelian group A with N has order 4.

Since N is characteristic, any automorphism of W_n induces an automorphism of $S_n = W_n/N$. If $n \neq 6$, this automorphism must be inner, since $\text{Out}(S_n)$ is trivial; we continue under the assumption that $n \neq 6$. Composing a given automorphism ϕ of W_n with conjugation by an element of $S_n = 1 \rtimes S_n$, we may assume that for all $\sigma \in S_n$ we have $\phi(\sigma) = g_\sigma \sigma$ for some $g_\sigma \in N$.

Let $\text{Fix}(\sigma)$ denote the fixed point set of the action of $\sigma \in S_n$ on N . For all $g \in N$ and $\sigma \in S_n$ we have $\phi(\sigma^{-1}g\sigma) = \sigma^{-1}\phi(g)\sigma$, so if $g \in \text{Fix}(\sigma)$ then $\phi(g) \in \text{Fix}(\sigma)$. Let σ be the $n-1$ cycle $(23 \dots n)$. Since $\text{Fix}(\sigma) = \{0, e_1, e_1 z, z\}$ and $\phi(e_1) \neq z$, we either have $\phi(e_1) = e_1$ or $\phi(e_1) = e_1 z$.

Given $i \neq 1$ we have $\phi(e_i) = \phi(e_1^{(1i)}) = \phi(e_1)^{(1i)}$ (writing the action of S_n as exponentiation) and hence we either have $\phi(e_i) = e_i$ for $i = 1, \dots, n$ or else $\phi(e_i) = e_i z$ for $i = 1, \dots, n$. This latter possibility is fine if n is even but does not define a homomorphism if n is odd because in that case $\phi(z) = \sum_i z \phi(e_i) = z \sum_i \phi(e_i) = z^2 = 0$.

After possibly composing with the outer automorphism β if n is even, we may now assume that $\phi|_N$ is the identity, as well as that ϕ induces the identity on the quotient $W_n/N = S_n$; thus we are reduced to finding all splittings $\phi: S_n \rightarrow W_n$. Such splittings are in one-to-one correspondence with the elements of $H^1(S_n, N)$. Here S_n acts on $N \cong (\mathbb{Z}/2)^n$ by permuting the factors, thus N is induced from the trivial action of S_{n-1} on $\mathbb{Z}/2$, that is $N = \text{Ind}_{S_{n-1}}^{S_n}(\mathbb{Z}/2)$. By Shapiro's lemma we have $H^1(S_n, N) \cong H^1(S_{n-1}, \mathbb{Z}/2) \cong \mathbb{Z}/2$. Thus the only splittings $S_n \rightarrow W_n$ are the one fixing the generators τ_i and the one sending each τ_i to $z\tau_i$.

It only remains to see that the non-trivial outer automorphism of S_6 does not extend to an automorphism of W_6 . But this is obvious from the fact that this outer automorphism interchanges the set of 3-cycles and the set of products of disjoint 3-cycles, and these different types of elements have stabilizers of different rank in the characteristic $N \subset W_6$. ■

3. GEOMETRIC REALIZATIONS OF FINITE SUBGROUPS

In this section we shall make liberal use of the vocabulary associated with the action of $\text{Out}(F_n)$ on Outer space. This action is related to the action of $\text{Aut}(\text{Out}(F_n))$ on the finite subgroups of $\text{Out}(F_n)$ by the following facts.

REALIZATION THEOREM (See [2, 11]). Every finite subgroup of $\text{Out}(F_n)$ fixes a vertex (marked graph) $[\Gamma, f]$ in the spine of Outer space, and there is a natural isomorphism from the stabilizer of $[\Gamma, f]$ to the group of graph-isomorphisms of Γ .

Unravelling the definitions, this amounts to the following: for every finite $G < \text{Out}(F_n)$ one can find a connected graph Γ (with no vertices of valence less than 3 and no separating edges), an isomorphism $f_*: F_n \rightarrow \pi_1(\Gamma)$ and a group $\tilde{G} = \{\tilde{g} \mid g \in G\}$ of automorphisms of Γ such that $g = f_*^{-1} \circ \tilde{g}_* \circ f_*$ for all $g \in G$. We say that $[\Gamma, f]$ realizes G . (One can realize finite subgroups $G < \text{Aut}(F_n)$ by graph automorphisms that fix a basepoint.)

Stabilizers of Roses are Characteristic

For $n \geq 3$, the subgroups of $\text{Out}(F_n)$ isomorphic to W_n are the maximal subgroups that can be realized by a rose; in other words, they are the stabilizers of the roses in Outer space (see, e.g., [7] or [10]).

LEMMA 3.1. *Let ϕ be an automorphism of $\text{Out}(F_n)$. The outer automorphism class of ϕ contains an automorphism ϕ' such that $\phi'(W_n) = W_n$ and $\phi'|_{W_n} \in \{\text{id}_{W_n}, \alpha, \beta, \alpha\beta\}$.*

Proof. We identify W_n with the stabilizer in $\text{Out}(F_n)$ of the rose ρ_0 associated with our fixed basis x_1, \dots, x_n of F_n ("the standard rose"). It follows from the second of the facts stated above that $\phi(W_n)$ is the stabilizer of some rose ρ_1 in Outer space. The action of $\text{Out}(F_n)$ is transitive on roses, so $\rho_1 = \psi(\rho_0)$ for some $\psi \in \text{Out}(F_n)$. Thus $\phi'(W_n) = W_n$, where ϕ' is the product of ϕ and $\text{ad}(\psi) \in \text{Inn}(\text{Out}(F_n))$. Lemma 2.1 finishes the proof. ■

We wish to eliminate the possibilities $\phi'|_{W_n} \in \{\alpha, \beta, \alpha\beta\}$. We shall do so by considering the action of ϕ' on the subgroup $H \subset W_n$ generated by $\{e_i, \tau_i \mid i = 3, \dots, n\}$ and τ_1 . The structure of H is $H = W_{n-2} \times \mathbb{Z}/2$, where the factor $\mathbb{Z}/2$ is generated by τ_1 and W_{n-2} is generated by the remaining τ_i and e_i .

We shall also need the subgroup $G = W_{n-2} \times S_3$, where W_{n-2} is as above and S_3 is the symmetric group of order 6 generated by τ_1 and the automorphism

$$r = ue_2: \quad x_1 \mapsto x_1x_2^{-1} \quad \text{and} \quad x_2 \mapsto x_2^{-1}.$$

G is the stabilizer of the marked graph γ shown in Fig. 1, and $G \cap W_n = H$.




FIGURE 1.

LEMMA 3.2. *The fixed point set for the action of H on the spine of Outer space consists of the subcomplex spanned by the vertices represented by the marked graphs shown in Fig. 1, and γ is the unique point fixed by G .*

Proof. According to [6] we can find all graphs realizing H by identifying one such graph and then successively collapsing and inserting H -invariant forests. (The process of inserting equivariant forests is called *equivalent blowing-up*.) One can perform an equivariant blowup of a graph Γ that realizes H if and only if the graph contains a set of oriented edges I with the following properties:

- (i) all edges in I terminate at the same vertex v ;
- (ii) I contains at least two and at most all but two of the edges terminating at v ;
- (iii) for $h \in H$, either $hI \cap I = I$ or $hI \cap I = \emptyset$;
- (iv) if v is a cut vertex, there are two edges e, f terminating at v which are in the same component of $\gamma - I$ such that $e \in I$ and $f \notin I$.

Since ρ_0 realizes H , we begin there. There are no invariant forests to collapse. There are four edge sets I which meet the criteria (i)–(iv) above, namely $I_1 = \{x_1, x_2\}$, $I_2 = \{\bar{x}_1, \bar{x}_2\}$, $I_3 = \{\bar{x}_1, x_2\}$, and $I_4 = \{x_1, \bar{x}_2\}$. Blowing up I_1 results in the graph γ , blowing up I_2 results in $z\gamma$, and blowing up either I_3 or I_4 results in $e_2\delta$. The graph δ is obtained from γ by blowing up I_2 (or from $z\gamma$ by blowing up I_1). No other graphs are obtainable by blowing up. Since collapsing invariant forests in these graphs produces no new marked graphs, this is the complete list of marked graphs fixed by H .

Any graph that realizes G also realizes $H \subset G$. The automorphism groups of δ and $e_2\delta$ do not contain subgroups isomorphic to G , and G is not contained in W_n , the stabilizer of ρ_0 , nor in $z^{-1}Gz$, the stabilizer of $z\gamma$. Thus G fixes only γ . ■

LEMMA 3.3. *Every class $[\phi] \in \text{Out}(\text{Out}(F_n))$ has a representative $\phi \in \text{Aut}(\text{Out}(F_n))$ such that $\phi|_{W_n} = \text{id}_{W_n}$.*

Proof. In the light of Lemma 3.1, we may assume that ϕ leaves W_n invariant, and we will be done if we can show that the “bad” possibilities $\phi|_{W_n} \in \{\alpha, \beta, \alpha\beta\}$ do not arise. If $n \geq 4$, by arguing as in Lemma 3.2 we see that for each bad possibility the only graphs fixed by $\phi|_{W_n}(H)$ are ρ_0 , δ , and $e_2\delta$. Thus if there were an automorphism ϕ with bad $\phi|_{W_n}$ then one of $\{\rho_0, \delta, e_2\delta\}$ would have to realize $\phi(G) \supset \phi(H)$.

But $\phi(H) = \phi(G \cap W_n) = \phi(G) \cap W_n$ is a proper subgroup of $\phi(G)$, and W_n is the stabilizer of ρ_0 , so ρ_0 does not realize $\phi(G)$. And the automorphism groups of δ and $e_2\delta$ do not contain subgroups isomorphic to G .

It remains to consider the case $n = 3$, supposing that $\phi|_{W_n} = \alpha$ (β does not give an automorphism of W_n since n is odd). In this case $\text{Fix}(\phi(H))$ is the subcomplex spanned by ρ_0 , δ , $e_2\delta$, $e_2\gamma$, and $ze_2\gamma$, so it is possible that $\phi(G)$ is the stabilizer of either $e_2\gamma$ or $ze_2\gamma$. After conjugating by z if necessary, we may assume $\phi(G) = e_2Ge_2$, the stabilizer of $e_2\gamma$.

The group G is generated by e_3 , τ_1 , and r . The only order 3 elements of G are $r\tau_1$ and τ_1r , so that $\phi: G \rightarrow e_2Ge_2$ must take $r\tau_1$ to $e_2\tau_1re_2$ or $e_2r\tau_1e_2$. Since we know that $\phi(\tau_1) = \alpha(\tau_1) = z\tau_1$ and $\phi(e_2) = e_2$, this implies that $\phi(r) = e_2e_3re_2$ or $e_2e_3\tau_1r\tau_1e_2$. In each of these cases we compute that, though $u = re_2$ (right multiplication of x_1 by x_2) commutes with $v = \tau_2zuz\tau_2$ (left multiplication of x_1 by x_3), the values forced for $\phi(u)$ and $\phi(v)$ do not commute, even up to inner automorphism. Thus there is no outer automorphism ϕ which restricts to α on W_3 . ■

4. REMAINDER OF THE PROOF

In the previous section we showed that every class $[\phi] \in \text{Out}(\text{Out}(F_n))$ contains a representative ϕ such that $\phi|_{W_n}$ is the identity. We also saw that any graph realizing $\phi(G) \supset \phi(H) = H$ must be in $\text{Fix}(H)$ and not equal to δ , $e_2\delta$, or ρ_0 . The only possibilities for such a graph are γ , in which case $\phi(G) = \text{stab}(\gamma) = G$, and $z\gamma$, in which case $\phi(G) = \text{stab}(z\gamma) = zGz^{-1}$. After composing by conjugation with z if necessary, we may assume $\phi(G) = G$.

Recall that $G = W_{n-2} \times S_3$ that $H = W_{n-2} \times \langle \tau_1 \rangle$, and that G is generated by H and the element r defined in Section 3. Recall also that $\text{Out}(F_n)$ is generated by W_n and r , so if we can show that $\phi(r) = r$ then we shall be done.

Since $\phi|_H$ is the identity, ϕ restricts to an automorphism of the centralizer of W_{n-2} in G . This centralizer is $\mathbb{Z}/2 \times S_3$, where $\mathbb{Z}/2 = \langle z' \rangle$ is the center of W_{n-2} (explicitly, $z' = e_3 \dots e_n$) and S_3 is generated by r and τ_1 . The only order 3 elements of this centralizer are $r\tau_1$ and τ_1r , so that $\phi(r\tau_1) = r\tau_1$ or $\phi(r\tau_1) = \tau_1r$. Since ϕ fixes τ_1 , this implies that $\phi(r) = r$ or $\phi(r) = \tau_1r\tau_1$.

Suppose that $\phi(r) = \tau_1r\tau_1$. We compute that $u = re_2$ commutes with e_1ue_1 but that $\phi(u) = \tau_1r\tau_1e_2$ does not commute with $\phi(e_1ue_1) = e_1\tau_1r\tau_1e_2e_1$. Furthermore, the commutator of $\phi(u)$ and $\phi(e_1ue_1)$ is not inner, contradicting the fact that ϕ is a homomorphism. ■

The proof that $\text{Out}(\text{Aut}(F_n))$ is trivial is essentially identical to the above proof; the only difference is that we consider each graph used as a graph with basepoint—the basepoint is at the (unique) vertex of maximal valence.

REFERENCES

1. M. R. Bridson and K. Vogtmann, The symmetries of Outer space, *Duke Math. J.*, to appear.
2. M. Culler, Finite groups of automorphisms of free groups, in "Contributions to Group Theory," Contemporary Mathematics, Vol. 33, pp. 197–207, Am. Math. Soc., Providence, 1984.
3. M. Culler and K. Vogtmann, Moduli of graphs and automorphisms of free groups, *Invent. Math.* **84** (1986), 91–119.
4. J. Dyer and E. Formanek, The automorphism group of a free group in complete, *J. London Math. Soc.* **11** (1975), 181–190.
5. N. V. Ivanov, Automorphisms of complexes of curves and of Teichmüller spaces, *Internat. Math. Res. Notices* No. 14 (1997), 651–666.
6. S. Krstić, Actions of finite groups on graphs and related automorphisms of free groups, *J. Algebra* **124** (1989), 119–138.
7. R. Kulkarni, Lattices on trees, automorphisms of graphs, free groups and surfaces, preprint, C.U.N.Y., 1988.
8. W. Magnus, A. Karrass, and D. Solitar, "Combinatorial Group Theory," Interscience, New York, 1966.
9. J. Tits, "Buildings of Spherical Type and Finite BN-Pairs," Lecture Notes in Mathematics, Vol. 386, Springer-Verlag, Berlin/Heidelberg/New York, 1974.
10. S. C. Wang and B. Zimmermann, The maximum order of finite groups of outer automorphisms of free groups, *Math. Z.* **216**, No. 1 (1994), 83–87.
11. B. Zimmermann, Über Homeomorphismen n -dimensionaler Henkelkörper und endliche Erweiterungen von Schottky-Gruppen [On homeomorphisms of n -dimensional handlebodies and finite extensions of Schottky groups], *Comment. Math. Helv.* **56**, No. 3 (1981), 474–486.