Arthroplasties for hip fracture in adults

Protocol information

Review type: Intervention

Review number: 284

Authors

Ashwini Sreekanta¹, Martyn J Parker², Hannah Wood³, Julie M Glanville³, MingHui Yang⁴, Jonathan Cook¹, Xavier L Griffin¹

¹Nuffield Department of Orthopaedics, Rheumatology and Musculoskeletal Sciences (NDORMS), University of Oxford, Oxford, UK

²Department of Orthopaedics, Peterborough and Stamford Hospitals NHS Foundation Trust, Peterborough, UK

³York Health Economics Consortium, York, UK

⁴Department of Orthopedics and Traumatology, Beijing Jishuitan Hospital, Beijing, China

Citation example: Sreekanta A, Parker MJ, Wood H, Glanville JM, Yang M, Cook J, Griffin XL. Arthroplasties for hip fracture in adults. Cochrane Database of Systematic Reviews , Issue . Art. No.: . DOI: .

Contact person

Ashwini Sreekanta

Systematic Reviewer

Nuffield Department of Orthopaedics, Rheumatology and Musculoskeletal Sciences (NDORMS)

University of Oxford

Kadoorie Centre

John Radcliffe Hospital, Headley Way

Oxford

OX3 9DU

UK

E-mail: ashwini.sreekanta@ndorms.ox.ac.uk

E-mail 2: ashcanbereached@gmail.com

Dates

Assessed as Up-to-date: Not provided

Date of Search: Not provided

Next Stage Expected: 31 March 2020

Protocol First Published: Not specified

Review First Published: Not specified

Last Citation Issue: Not specified

What's new

Date	Event	Description
------	-------	-------------

History

Date	Event	Description
------	-------	-------------

Abstract

Background

Objectives

Search methods

Selection criteria

Data collection and analysis

Main results

Authors' conclusions

Plain language summary

[Summary title]

[Summary text]

Background

Description of the condition

Epidemiology

A hip fracture, or proximal femoral fracture, is a break in the upper region of the femur (thigh bone) between the subcapital region (the area just under the femoral head) and 5 cm below the lesser trochanter (a bony projection of the upper femur). The incidence of hip fractures increases with age, and are most common in the older adult population ([Court-Brown 2017](#); [Kanis 2001](#)). Hip fractures in younger adults are usually associated with poor bone health ([Karantana 2011](#); [Rogmark 2018](#)). A small proportion of fractures occurring in younger people are as a result of high-energy trauma, such as road traffic collisions and sports injuries. Most hip fractures are fragility fractures associated with osteoporosis, such fractures resulting from mechanical forces that would not ordinarily result in fracture. The World Health Organization (WHO) has defined fragility fracture as those sustained from injuries equivalent to a fall from a standing height or less ([Kanis 2001](#)). In the UK, the mean age of a person with hip fracture is 83 years and approximately two-thirds occur in women ([NHFD 2017](#)).

Hip fractures are a major healthcare problem at the individual and population level, and present a huge challenge and burden to patients, healthcare systems, and society. The increased proportion of older adults in the world population means that the absolute number of hip fractures is rising rapidly worldwide. For example, in 2016 there were 65,645 new presentations of hip fracture to 177 trauma units in England, Wales, and Northern Ireland ([NHFD 2017](#)). Based on mid-2016 population estimates for these regions, this equates to an incidence rate of 108 cases per 100,000 population ([ONS 2018](#)). By 2050, the annual worldwide incidence is estimated to be six million hip fractures ([Cooper 2011](#); [Johnell 2004](#)). Incident hip fracture rates are higher in industrialised countries compared to low- or middle-income countries. The highest hip fracture rates are seen across northern Europe and the USA, and the lowest in Latin America and Africa ([Dhanwal 2011](#)). There is also a north-south gradient seen in European studies, and similarly more fractures are seen in the north of the USA than in the south ([Dhanwal 2011](#)). The factors responsible for this variation are thought to be population demographics (with more elderly populations in countries with higher incidence rates) and the influence of ethnicity, latitude, and environmental factors such as socioeconomic deprivation ([Bardsley 2013](#); [Cooper 2011](#); [Dhanwal 2011](#); [Kanis 2012](#)).

Burden of disease

Hip fractures are also associated with a high risk of death. For example, in England, Wales, and Northern Ireland, the 30-day mortality rate in 2016 remained high at 6.7%, despite a decline from 8.5% in 2011 and 7.1% in 2015 ([NHFD 2017](#)). Mortality at one year following a hip fracture is approximately 30%; however, fewer than half of deaths are attributable to the fracture itself, reflecting the frailty of the patients and associated high prevalence of co-morbidities and complications ([Parker 1991](#); [SIGN 2009](#)). Morbidity associated with hip fractures is similar to stroke in terms of impact, with a substantial loss of healthy life-years in older people ([Griffin 2015](#)). As such, hip fractures commonly result in reduced mobility and greater dependency, with many people failing to return to their pre-injury residence. In addition, the public health impact of hip fractures is significant: data from large prospective cohorts show the burden of disease due to hip fracture is 27 disability-adjusted life years (DALYs) per 1000 individuals, which equates to an average loss of 2.7% of the healthy life expectancy in this population at risk of fragility hip fracture ([Papadimitriou 2017](#)).

The direct economic burden of hip fractures is also substantial. Hip fractures are among the most expensive conditions seen in hospitals, with an aggregated cost of nearly USD 4.9 billion for 316,000 inpatient episodes in the USA in 2011 ([Torio 2011](#)). In England, Wales, and Northern Ireland, hip fracture patients occupy 1.5 million hospital bed days each year, and cost the National Health Service (NHS) and social care GBP 1 billion ([NHFD 2017](#)). Combined health and social care costs incurred during the first year following a hip fracture have been estimated at USD 43,669, which is greater than the cost for non-communicable diseases such as acute coronary syndrome (USD 32,345) and ischaemic stroke (USD 34,772) ([Williamson 2017](#)). In established market economies, hip fractures represent 1.4% of the total healthcare burden ([Johnell 2004](#)).

Types of hip fracture

Hip fractures either involve the region of the femur that is enveloped by the ligamentous hip joint capsule (intracapsular), or that is outside the capsule (extracapsular).

Intracapsular fractures include subcapital (immediately below the femoral head), transcervical (across the mid-femoral neck), or basicervical (across the base of the femoral neck). These injuries are also commonly termed fractures of the 'neck of femur' ([Lloyd-Jones 2015](#)). Intracapsular fractures can be further subdivided by fracture morphology using several different classification systems, such as the Garden ([Garden 1961](#)) or Pauwels classifications ([Pauwels 1935](#)). The reliability of these various classifications is poor ([Parker 1993](#); [Parker 1998](#)). A more appropriate grouping distinguishes only those fractures that are displaced, where the anatomy of the bone has been disrupted at the fracture site, and those that are undisplaced ([Blundell 1998](#); [Parker 1999](#)). This system broadly corresponds with prognosis; the more displaced, the more likely

the blood supply to the femoral head is compromised, which can lead to complications such as avascular necrosis and collapse of the femoral head. Furthermore, displaced fractures are less stable, so that treatments involving fixation have a higher risk of failure compared with displaced fractures. Approximately 60% of hip fractures are intracapsular, of these approximately 70% to 90% are displaced ([Keating 2010](#); [NHFD 2017](#)).

Extracapsular fractures traverse the femur within the area of bone bounded by the intertrochanteric line proximally up to a distance of 5 cm from the distal part of the lesser trochanter. Several classification methods have been proposed to define different types of extracapsular fractures ([AO Foundation 2018](#); [Evans 1949](#); [Jensen 1980](#)). They are generally subdivided depending on their relationship to the greater and lesser trochanters, the two bony projections present at the upper end of the femur, and the complexity of the fracture configuration. It is increasingly clear that each of these classifications is limited in its generalisability since inter- and intra-observer agreement is poor. [Table 1](#) provides a description of the most recent classification of trochanteric fractures ([AO Foundation 2018](#)). For this Cochrane Review, we plan to use a pragmatic simplification of these classifications as follows.

- Trochanteric fractures: those that lie mostly between the intertrochanteric line and a transverse line at the level of the lesser trochanter. These can be further divided into simple two-part stable fractures and comminuted or reverse obliquity unstable fractures.
- Subtrochanteric fractures: those that mostly lie in the region bordered by the lesser trochanter and 5 cm distal to the lesser trochanter.

Approximately 40% of hip fractures are extracapsular, of which 90% are trochanteric and 10% are subtrochanteric ([NHFD 2017](#)).

Description of the intervention

Internationally, many guidelines exist concerning hip fracture management (e.g. [AAOS 2014](#); [NICE 2011](#); [SIGN 2009](#)). Each recommends that early surgical management, generally within 24 to 48 hours, is the mainstay of care for most hip fractures. The overall goal of surgery in the older population is to facilitate early rehabilitation, enabling early mobilisation and the return to premorbid function while minimising the complication risk. This approach has been associated with reductions in mortality in many worldwide registries ([Neufeld 2016](#); [Sayers 2017](#)). A proposed grouping of arthroplasty interventions is given in [Table 2](#).

Arthroplasty

Arthroplasty entails replacing part or all of the hip joint with an endoprosthesis: an implant constructed of non-biological materials such as metal, ceramic, or polyethylene. Arthroplasties can be grouped into two main categories: hemiarthroplasty (HA) where only the femoral head and neck are replaced, and total hip replacement (THR) where both the femoral head and the acetabulum or socket are replaced.

Hemiarthroplasty

Hemiarthroplasty involves replacing the femoral head with a prosthesis whilst retaining the natural acetabulum and acetabular cartilage. The type of HA can be broadly divided into two groups: unipolar and bipolar. In unipolar HAs, the femoral head is a solid block of metal. Bipolar femoral heads include a single articulation that allows movement to occur, not only between the acetabulum and the prosthesis, but also at this joint within the prosthesis itself.

The best known of the early HA designs are the Moore prosthesis (1952) and the FR Thompson Hip Prosthesis (1954). These are both monoblock implants and were designed before the development of polymethylmethacrylate bone cement. They were therefore originally inserted as a 'press fit'. The Moore prosthesis has a femoral stem, which is fenestrated and also has a square stem with a shoulder to enable stabilisation within the femur; this resists rotation within the femoral canal. It is generally used without cement and, in the long term, bone in-growth into the fenestrations can occur. The Thompson prosthesis has a smaller stem without fenestrations and now often used in conjunction with cement. Numerous other designs of unipolar HAs exist based on stems that have been used for THRs.

In bipolar prostheses, there is an articulation within the femoral head component itself. In this type of prosthesis there is a spherical inner metal head with a size between 22 to 36 mm in diameter. This fits into a polyethylene shell, which in turn is enclosed by a metal cap. The objective of the second joint is to reduce acetabular wear by promoting movement at the intraprosthetic articulation rather than with the native acetabulum. There are a number of different types of prostheses with different stem designs. Examples of bipolar prostheses are the Charnley-Hastings, Bateman, Giliberty, and the Monk prostheses, but many other types with different stem designs exist.

Total hip replacement

Total hip replacement involves the replacement of the acetabulum in addition to the femoral head. The first successful THR was developed by John Charnley, using metal alloy femoral heads articulating with polyethylene acetabular components. Subsequently the articulating materials have diversified and designs using metal alloys, ceramics, and various polyethylenes in various combinations have all been used.

Component fixation

Irrespective of the nature of the articulating surfaces, the components must be fixed to the bone to ensure longevity of the arthroplasty. The two approaches used to achieve this fixation are cemented and uncemented designs.

Cemented systems

Polymethylmethacrylate bone cement may be inserted at the time of surgery. It sets hard and acts a grout between the prosthesis and the implant the time of surgery. Potential advantages of cement are a reduced risk of intraoperative fracture, later peri-prosthetic fracture and not relying on integration of the prosthesis with osteoporotic bone. Major side effects of cement are cardiac arrhythmias and cardiorespiratory collapse, which occasionally occur following its insertion. These complications may be fatal. The cause of this is either embolism from marrow contents forced into the circulation ([Christie 1994](#)), or a direct toxic effect of the cement.

Uncemented systems

Uncemented systems rely on osseous integration forming a direct mechanical linkage between the bone and the implant. A prosthesis may be coated with a substance, such as hydroxyapatite, which promotes bone growth into the prosthesis. Alternatively, the surface of the prosthesis may be macroscopically and microscopically roughened so that bone grows onto the surface of the implant.

The complications of arthroplasty are those that are general to surgical management of hip fracture, e.g. pneumonia, venous thromboembolism, infection, acute coronary syndrome, and cerebrovascular accident; and those that are specific to arthroplasty including dislocation of the prosthesis, loosening of the components, acetabular wear, and periprosthetic fracture.

Why it is important to do this review

This review will replace the Cochrane Review, [Parker 2010](#), on the same topic. We will use up-to-date review methods and optimise current relevance in terms of patient population, implants used, and outcomes for policymaking bodies such as the National Institute for Health and Care Excellence (NICE) in the UK, as well as international audiences. Since [Parker 2010](#), clinical uncertainty remains as to the optimum implant for this patient population; moreover, further studies have been reported since the last literature search in September 2009.

Appraisal and synthesis of contemporary evidence may enable more robust conclusions to be made to better inform practice. Furthermore, for displaced intracapsular fractures, the recommended treatment is either HA or total hip arthroplasty (THA) ([Parker 2010](#); [Hopley 2010](#); [NICE 2011](#)). However, there is a lack of evidence regarding whether this patient group experiences better outcomes with THA and HA. Recent research has also found interhospital variation and systematic inequalities observed in the provision of THA ([Perry 2016](#)). Further evidence is necessary to verify which patients gain the most from THA. For treatment of undisplaced intracapsular fractures, there is also a gap in the evidence that resulted in the recently updated NICE guideline being unable to make an evidence-based recommendation on the best surgical management strategy ([NICE 2011](#)). Other reviews are in preparation that will address other types of interventions; we will focus on arthroplasty in this review.

Objectives

To determine the relative effects (benefits and harms) of different designs, articulations, and fixation techniques of arthroplasties for treating hip fractures in adults.

This will include assessment of the relative effects (benefits and harms) of the following.

- THR versus HA
- Cemented versus uncemented arthroplasties
- Alternative articulation designs in THA
- Different designs of HA, e.g. bipolar versus unipolar

Methods

Criteria for considering studies for this review

Types of studies

We will include randomised controlled trials (RCTs) and quasi-RCTs that assess surgical interventions for the management of patients with hip fracture. Quasi-RCTs are defined as trials in which the methods of allocating people to a trial are not properly random, but were intended to produce similar groups when used to allocate participants ([Cochrane 2018](#)). We will include trials published as conference abstracts provided the trial authors report sufficient data relating to the methods and outcomes of interest. Unpublished data will also be considered for inclusion.

Types of participants

Population

In order to report the generality of evidence available for these patients, we plan to take a wide and pragmatic

approach to defining the eligibility criteria. We will report the details of the population in detail in the 'Characteristics of included studies' table.

As a benchmark, thus representative of the general hip fracture population, we would expect trial populations to have a mean age of between 80 to 85 years, and include 70% women, 30% with chronic cognitive impairment, and 50% with an American Society of Anesthesiologists (ASA) score greater than 2 ([NHFD 2017](#); [NICE 2011](#)).

Inclusion criteria

- All adults undergoing surgery for a fragility (low energy trauma) hip fracture, whether intracapsular or extracapsular

Exclusion criteria

Studies focusing exclusively on the treatment of the following.

- Patients younger than 16 years
- Patients with fractures caused by specific pathologies other than osteoporosis
- Patients with high energy fractures

Studies with mixed populations (fragility and other mechanisms, ages, fracture subtypes, or pathologies) will also be eligible for inclusion. Where data are reported separately, we will extract those subgroup data. Where a study has a mixed population, but subgroups are not reported, the proportion of patients who have standard fragility fracture is likely to hugely outnumber the numbers with high energy or local pathological fractures; therefore the results will be generalisable to fragility fracture population. We will consider sensitivity analyses, where possible, to test this assumption (see [Sensitivity analysis](#)).

We will not pool studies in which the fracture type is mixed (intracapsular and extracapsular).

Healthcare setting

The expected healthcare setting will be hospitals where operative acute care is undertaken.

Types of interventions

All hip prostheses: unipolar HA, bipolar HA, or THR (small and large head), applied with or without cement.

Comparisons

- Prostheses inserted with cement versus without cement (stratified by THR versus HA; HA group subgrouped by modern versus old uncemented stems)
- Bipolar HA versus unipolar HA
- HAs with each other (subgrouped by modern stem design ('ODEP 3A rating') and old stem design (e.g. Austin Moore or Thompson) (new comparison)
- THR versus HA (cemented or uncemented, subgrouped by old versus new as per above)
- Single versus multiple (dual/triple) articulations of THR
- Large-head THR (36 mm diameter or larger) versus other arthroplasty (stratified by THR versus HA)

Direction of comparison: in each comparison, the intervention is named first and the control second.

We plan to create a more detailed table of interventions, grouping them into those that can be reasonably grouped and indicating which are in worldwide use. We will perform this with clinical authors and with the International Fragility Fracture Network (www.fragilityfracturenetwork.org/). A preliminary exercise resulted in the proposed implant groupings given in [Table 2](#).

Types of outcome measures

Critical outcomes

We will extract information on the following 7 critical outcomes where reported. These are listed, which are in alphabetical order below, where reported. Depending on the length of follow-up reported, the endpoints for each of these 7 each critical outcomes will be categorised into early (before or up to 4 months) and late (after 4 months). Haywood 2014 Griffin 2015 Haywood 2014 We have selected 4 months as the definition of early, since most of early recovery has been achieved at this timepoint (Griffin 2015), and in accordance with the core outcome set for hip fracture (Haywood 2014) which prioritises early outcome over late recovery. Although priority will be given to outcome at 4 months, we will consider data availability in our selection of data for presentation.

- Activities of daily living (e.g. Barthel Index (BI), Functional Independence Measure (FIM))
- Delirium using recognised assessment scores, such as Mini mental test score or 4AT
- Functional status (region specific) (e.g. hip rating questionnaire, Harris Hip Score, Oxford Hip Score)
- Health-related Quality-of-Life (HRQoL) (e.g. SF36, EQ-5D)
- Mobility (e.g. indoor/outdoor walking status, Cumulated Ambulation Score, Elderly Mobility Scale Score, Timed up and go, Short Physical Performance Battery, self-reported walking scores (e.g. Mobility Assessment Tool – short form))

- Mortality
- Unplanned return to theatre: secondary procedure required for a complication resulting directly or indirectly from the index operation/primary procedure

~~Haywood 2014 Griffin 2015~~

Other important outcomes

In addition we will also report the following important outcomes which we will also categorise into early (up to and including 4 months) and late (after 4 months):

~~We will collect for these outcomes data up to 12 months, if reported:~~

- Pain (verbal rating or visual analogue scale) ~~(early within 4 months)~~
- Length of in-hospital stay (LOS)
- Discharge destination. This may be variably defined in the included studies. We will use the study authors' definitions and report all events without a time limit
- Adverse events

We will group adverse events by relatedness to the implant or fracture, or both. We will report each adverse event type separately for maximum clarity. We anticipate that events may include the following.

Related

- Damage to a nerve, tendon, or blood vessel
- Intraoperative peri-prosthetic fracture
- Postoperative peri-prosthetic fracture
- Loosening of prosthesis
- Wound infection (as defined by trial investigators, often infection is described as deep or superficial)
- Dislocation

Unrelated

- Acute kidney injury
- Blood transfusion
- Cerebrovascular accident
- Chest infection/pneumonia
- Decreased cognitive ability
- Myocardial infarction/acute coronary syndrome
- Sepsis
- Urinary tract infection
- Venous thromboembolic phenomena

Search methods for identification of studies

We will search for all published, unpublished, and ongoing relevant RCTs, without restrictions on language or date. Where possible, we will remove animal studies using the strategy.

We will develop general search strategies for the large bibliographic databases to find records to feed into a number of Cochrane Reviews on hip fracture surgery. We will use three approaches to identify eligible studies. The approaches are described conceptually as:

1. Hip fractures AND RCT filter;
2. Hip replacement AND fractures AND RCT filter;
3. Internal fixtured AND hip fractures AND RCT filter;
4. 1 OR 2 OR 3.

In MEDLINE, we will use the Cochrane Highly Sensitive Search Strategy for identifying randomised trials ([Lefebvre 2018](#)). In Embase, we will use the Cochrane Embase filter to focus on RCTs (www.cochranelibrary.com/central/central-creation).

Electronic searches

We will search the following electronic databases:

- Cochrane Central Register of Controlled Trials (CENTRAL)
- MEDLINE ALL (Ovid)
- Embase (Ovid)
- Science Citation Index (Web of Science)
- Cochrane Database of Systematic Reviews (CDSR) (Cochrane Library)
- Database of Abstracts of Reviews of Effects (DARE) ([CRD website](#))
- Health Technology Assessment (HTA) database ([CRD website](#))
- Epistemonikos (www.epistemonikos.org/)
- ClinicalTrials.gov (<https://clinicaltrials.gov/>)
- WHO ICTRP (www.who.int/ictip/en/).

As CENTRAL is kept fully up-to-date with all records from the BJMT Group's Specialised Register, we do not plan

to search the latter separately.

The search strategies for the above databases will be modelled on the search strategy designed for MEDLINE ([Appendix 1](#)), where possible. Adaptation includes consideration of database interface differences as well as adaptation to different indexing languages. We will document the final search strategies for each of the databases searched in the published review.

Searching other resources

We will search for unpublished research, conference reports, and research reported in the grey literature by searching a range of resources including the following.

- Handsearching abstracts from the following conferences (2016 to present):
 - Fragility Fractures Network Congress
 - British Orthopaedic Association Congress
 - Orthopaedic World Congress (SICOT)
 - Orthopaedic Trauma Association Annual Meeting
 - Bone and Joint Journal Orthopaedic Proceedings
 - American Academy of Orthopaedic Surgeons Annual Meeting
- Proquest Dissertations and Theses
- National Technical Information Service (NTIS, for technical reports)

To identify further studies, we will screen the reference lists of eligible studies and systematic reviews published within the last five years that have been retrieved by the searches. We will screen the reference lists of relevant Cochrane Reviews that are being updated, irrespective of the date they were published.

Data collection and analysis

Any review author who is a co-applicant, author or has or has had an advisory role on any potentially relevant study, will remain independent of study selection decisions, risk of bias assessment and data extraction for their study.

Selection of studies

Two review authors will screen the titles and abstracts of all retrieved bibliographic records using the internet-based systematic reviewing platform, Rayyan ([Ouzzani 2016](#)). Two review authors will retrieve and independently examine the full texts of all potentially eligible records after title/abstract screening using the eligibility criteria (see [Criteria for considering studies for this review](#)). We will perform full-text screening using the internet-based platform Covidence ([Covidence](#)). Disagreements will be resolved by discussion or adjudication by a third review author. Where necessary, we will correspond with study authors where clarification is required to inform study selection. We will exclude duplicates and will collate multiple reports of the same study so that each study, rather than each report, is the unit of interest in the review. We will construct a PRISMA flow-diagram ([Moher 2009](#)), which will outline the study selection process, numbers of records at each stage of selection, and reasons for exclusions of full-text articles. Details will also be recorded in a 'Characteristics of excluded studies' table.

Data extraction and management

Two review authors will independently extract data using a prepiloted structured form to ensure consistency of information and appraisal of each study. They will independently pilot the form on at least one included study before implementation. The two review authors will ascertain that data are entered correctly into the final data set. We will extract data on the following.

Study methodology

Publication type, sponsorship and funding for the trial and any notable conflicts of interest of trial authors, study design, trial phase, number of centres and location(s), size and type of setting (e.g. in-hospital or out-of-hospital or mixed or community), study period and length of follow-up, stated study objectives, study inclusion and exclusion criteria, randomisation method, masking, study disposition (number randomised, number by protocol, number available for analysis).

Population

We will extract information on baseline characteristics of the participants, including age, gender, comorbidities, functional status (such as previous mobility), fracture type and displacement, and cognitive status.

Interventions

We plan to extract data concerning the exact nature of the interventions tested and any reported co-interventions. We will also extract information, where available, on any co-interventions, such as preoperative care (e.g. prophylaxis – antibiotics, venous thromboembolism, delirium), anaesthetic management, and postoperative care (e.g. rehabilitation).

Outcome data

Where possible, we will extract data by arm rather than the summary effect sizes. Outcome worksheets will be in 'one study per row format' and will specify the number of arms for each study and number of participants in each

arm; numbers randomised and analysed for each outcome at each timepoint; number of events in each arm (for rate, binary or categorical data); and means and standard deviations, effect measures, point estimates and confidence limits (for continuous variables). Where available, we will split the outcomes into early and late as described.

Assessment of risk of bias in included studies

We will assess the risk of bias in the included studies using the tool described for standard systematic reviews in the *Cochrane Handbook for Systematic Reviews of Interventions* ([Higgins 2011a](#)). Two review authors will independently perform these assessments and will resolve any discrepancies through discussion or by consulting a third review author. Where details of methods are unclear or not reported, we will contact the study authors for additional information.

We will evaluate the risk of bias in the following domains: selection bias (random sequence generation and allocation concealment), performance bias (blinding of participants and personnel), detection bias (blinding of outcome assessment), attrition bias (incomplete outcome data), reporting bias (selective reporting), and other bias. We will assess each potential source of bias as either high, low, or unclear and will provide a quote from the study report together with a justification for our judgement in the 'Risk of bias' table. The 'Risk of bias' judgements will be summarised across different studies for each of the domains listed.

Assessment of risk of bias is specific to a particular result for a particular outcome (and time point) in the study. However, some domains will apply generally to the whole study (such as random sequence generation and allocation concealment); some will apply mainly to the outcome being measured or measurement method being used (such as blinding of participants and personnel, and blinding of outcome assessment); and some will apply to the specific result (such as selective reporting bias).

In the domains specific to particular outcomes, we will provide considerations of risk of bias for different types of outcomes and will make assessments separately. For example, for participant reported outcomes (e.g. pain, HRQoL), observer reported outcomes not involving judgement (e.g. all-cause mortality), observer reported outcomes involving some judgement (e.g. assessment of a radiograph, clinical examination, and clinical events other than death), outcomes that reflect decisions made by the intervention provider (e.g. hospitalisation, stopping treatment, referral to a different ward and discharge of the patient), and composite outcomes (e.g. major cardiac and cerebrovascular events (MACCE)).

As trials frequently contribute multiple results, mainly through contributing to multiple outcomes, several 'Risk of bias' assessments may be needed for each study. These assessments are likely to align with the outcomes included in a 'Summary of findings' table.

Measures of treatment effect

Where sufficiently reported, we will analyse data on an intention-to-treat (by allocated groups irrespective of compliance) basis.

We will calculate risk ratios (RR) for dichotomous data outcomes with 95% confidence intervals (CIs). Where the number of observed events is small (less than 1% of sample per group or with zero events in either arm), and where trials have balanced treatment groups, we will report the Peto odds ratio (OR) with 95% CI ([Deeks 2017](#)). We will express treatment effects for continuous data outcomes as mean differences (MD) with 95% CI; if the outcomes are measured using different scales, we will use standardised mean differences (SMD) with 95% CI.

Unit of analysis issues

We anticipate that the patient will be the unit of analysis. We do not expect to encounter any within-person randomised trials or cluster-randomised trials. If we do identify any, we will treat them in accordance with the guidance in Chapter 16 of the *Cochrane Handbook for Systematic Reviews of Interventions* ([Higgins 2011b](#)). We expect outcomes to be reported at various timepoints. We have specified short-term and long-term follow-up analyses in the [Types of outcome measures](#) section. We will also treat multi-arm trials in accordance with the advice given in Chapter 16 of the *Cochrane Handbook for Systematic Reviews of Interventions* ([Higgins 2011b](#)).

Dealing with missing data

We will contact the study authors to attempt to obtain missing data or to clarify unclear data. For each included study, we will record the number of participants lost to follow-up. Our primary analysis will be a complete-case analysis; we do not plan to impute missing data. However, if large amounts of data are missing, we will consider exploratory sensitivity analyses to investigate possible effects of the missingness (see [Sensitivity analysis](#)). Where study authors do not report standard deviations, we will try to determine these from standard errors, CIs, or exact P values.

Assessment of heterogeneity

We will use the I^2 statistic, automatically calculated in the Review Manager 5 software ([Review Manager 2014](#)), to quantify the possible degree of heterogeneity of treatment effects between trials. We will assume moderate heterogeneity when the I^2 statistic value is between 30% and 60%; substantial heterogeneity when it is between 50% and 90%; and considerable heterogeneity when it is between 75% and 100%. We will keep in mind that the importance of the I^2 statistic value depends on: 1) magnitude and direction of effects; and 2) strength of evidence

for heterogeneity. If we identify substantial heterogeneity, we will report it and investigate possible causes by following the recommendations in Section 9.6 of the *Cochrane Handbook for Systematic Reviews of Interventions* (Deeks 2017).

We will assess clinical and methodological diversity in terms of participants, interventions, outcomes, effect modifiers, and study characteristics for the included studies to determine whether a meta-analysis is appropriate. We will conduct this by observing these data from the data extraction tables.

We will then visually inspect the forest plot to look at the consistency of intervention effects across included studies. If the studies are estimating the same intervention effect, there should be overlap between the CIs for each effect estimate on the forest plot. However, if overlap is poor or there are outliers, statistical heterogeneity may be likely.

Assessment of reporting biases

Where sufficient numbers of studies (more than 10) are available for meta-analysis, we will construct a funnel plot to identify potential for publication bias as well as to explore possible small study biases. In interpreting funnel plots, we will examine the different reasons possible for funnel plot asymmetry and relate this to the results of the review. We will also undertake formal statistical tests to investigate funnel plot asymmetry, and will follow the recommendations in Section 10.4 of the *Cochrane Handbook for Systematic Reviews of Interventions* (Sterne 2017).

To assess outcome reporting bias, we will check trial protocols against their published reports. We will also screen clinical trial registries for a priori trial protocols. We will evaluate whether selective reporting of outcomes is present.

Data synthesis

We will undertake meta-analyses only where this is meaningful, that is, if the treatments, participants, and the underlying clinical question are similar enough for pooling to make sense. When considered appropriate, we will pool results of comparable groups of trials using random-effects models. We chose this model after careful consideration of the extent to which any underlying effect could truly be thought to be fixed given the complexity of the interventions included in this review. We will present 95% CIs throughout.

We will consider not pooling data where there is considerable heterogeneity (I^2 statistic value of greater than 75%) that cannot be explained by the diversity of methodological or clinical features among trials. We will still present trial data in the analyses or tables for illustrative purposes and report these in the text.

We will not pool multiple component interventions together with studies that report comparisons of only a single intervention. We will group any relevant studies by the combination of interventions (i.e. where the same combination of single categories of intervention is delivered to all participants). We will analyse each combination separately.

When considered appropriate, we will pool data using the generic inverse variance method in Review Manager 5 (Review Manager 2014). This method enables pooling of the adjusted and unadjusted treatment effect estimates reported in the individual studies or that can be calculated from data presented in the published article (see Measures of treatment effect). The generic inverse variance option in Review Manager 5 requires entering the natural logarithm of the treatment effect and its standard error for each trial (Review Manager 2014).

Subgroup analysis and investigation of heterogeneity

If sufficient number of studies meet the inclusion criteria, we will explore (as possible sources of inconsistency or heterogeneity between studies) key confounders, such as age (cut-off 85 years), gender, cognitive impairment, and fracture type (e.g. displaced versus minimally displaced and extracapsular versus intracapsular fractures). We also have some pre-specified comparison-specific subgroups.

We will investigate whether the results of subgroups are significantly different by inspecting overlapping CIs and by performing the test for subgroup differences available in Review Manager 5 (Review Manager 2014), where appropriate.

There is no explicit means of accounting for step changes in co-interventions, and certainly not one that would be applicable to the worldwide totality of the evidence. Therefore we cannot try to explain any heterogeneity by statistical test of subgroups defined by co-intervention we will order forest plots by date of recruitment so we can visually inspect and comment on any temporal trend.

Sensitivity analysis

If a sufficient number of studies meet the inclusion criteria, we will assess the effect of excluding the following types of studies.

- At high risk of bias
- Have substantial amounts of missing data
- Have quasi methods of randomisation
- Have mixed population
- Report interventions that are not currently in clinical use

GRADE assessment and 'Summary of findings' tables

Two review authors will independently assess the certainty of evidence for each of the seven primary ('critical') outcomes using the GRADE approach developed by the international GRADE working group (www.gradeworkinggroup.org/). We will use the software developed by the GRADE working group, [GRADEpro GDT 2015](#), to assess the certainty of the evidence for each outcome, taking into account individual study factors and the meta-analysis results. We will also present the results and the certainty of evidence for the main outcomes in a 'Summary of findings' table using GRADEpro GDT software ([Schünemann 2019](#)).

The chosen outcomes are those we consider to be most clinically important and useful for readers; these are described under 'Critical outcomes' in the [Types of outcome measures](#) section. We will consider the GRADE domains: risk of bias, inconsistency, indirectness, imprecision, and publication bias ([Guyatt 2008](#)). We will resolve any disagreements by consulting a third review author. The certainty of the evidence could be either high, moderate, low, or very low; being downgraded by one or two levels depending on the presence and extent of the five GRADE domains. We will use footnotes to describe the reasons for grading a domain as having 'serious' or 'very serious' problems and we will make further comments to aid the reader's understanding of the 'Summary of findings' table where necessary. We will consider the certainty of the evidence and the effect when making conclusions. We will also provide rationale for the figure used to calculate the assumed risk in a footnote.

We will provide further details on the full methods used for GRADE assessment of outcomes in an appendix in the published review.

Results

Description of studies

Results of the search

Included studies

Excluded studies

Risk of bias in included studies

Allocation (selection bias)

Blinding (performance bias and detection bias)

Incomplete outcome data (attrition bias)

Selective reporting (reporting bias)

Other potential sources of bias

Effects of interventions

Discussion

Summary of main results

Overall completeness and applicability of evidence

Quality of the evidence

Potential biases in the review process

Agreements and disagreements with other studies or reviews

Authors' conclusions

Implications for practice

Implications for research

Acknowledgements

We thank Joanne Elliott, Helen Handoll, and Maria Clarke for editorial base support. We thank Cecilia Rogmark and Emil Schemitsch for their helpful comments at external peer review.

This project was funded and supported by the National Institute for Health Research (NIHR). The views and opinions expressed therein are those of the review authors and do not necessarily reflect those of the Systematic Reviews Programme, the NIHR, the NHS, or the Department of Health.

Contributions of authors

AS (systematic reviewer): drafted the protocol, reviewed, and approved the final protocol.

MP (content expert, Orthopaedics): drafted the protocol, reviewed, and approved the final protocol.

HW (senior information specialist): designed the search strategies, reviewed, and approved the final protocol.

JMG (senior information specialist): designed the search strategies, drafted the protocol, reviewed, and approved

the final protocol.

MY (content expert, Trauma and Orthopaedics): reviewed and approved the final protocol.

JC (statistician): reviewed and approved the final protocol.

XG (content expert, Trauma and Orthopaedics): drafted the protocol, reviewed and approved the final protocol, and is the guarantor of the content.

Contributions of the editorial base

Helen Handoll (Co-ordinating Editor): edited the protocol, advised on methodology and protocol content, and approved the final version for publication.

Joanne Elliott (Managing Editor): coordinated the editorial process, advised on content, and edited the protocol.

Maria Clarke (Information Specialist): checked the search methods section.

Declarations of interest

AS: has no known conflicts of interest.

MP: has received expenses and honorarium from a number of commercial companies and organisations for giving lectures on different aspects of hip fracture treatment. In addition, he has received royalties from BBrown Ltd related to the design and development of an implant used for the internal fixation of intracapsular hip fractures.

HW: has no known conflicts of interest.

JMG: has no known conflicts of interest.

MY: has no known conflicts of interest.

JC: has no known conflicts of interest.

XG is funded by a National Institute for Health Research Clinician Scientist Grant. Further funding from industry and charitable grants are and have been made available to his institution. All decisions relating to the design, conduct, analysis, write-up and publication of research are independent of these funders. He has ongoing expert consultancy with several companies; none involve the development of any implant for use in hip fracture care. He will remain independent of study selection decisions, 'Risk of bias' assessment and any data extraction of any of the studies on which he is an author, co-applicant or has had an advisory role.

Differences between protocol and review

Published notes

Characteristics of studies

Characteristics of included studies

Footnotes

Characteristics of excluded studies

Footnotes

Characteristics of studies awaiting classification

Footnotes

Characteristics of ongoing studies

Footnotes

Summary of findings tables

Additional tables

1 Trochanteric region fractures: type and surgical management (revised AO/OTA classification, January 2018)

Type	Features	Stability	Description
Simple, pertrochanteric fractures (A1)	<ul style="list-style-type: none"> • Isolated pertrochanteric fracture • 2-part fracture • Lateral wall intact 	Stable	The fracture line can begin anywhere on the greater trochanter and end either above or below the lesser trochanter. The medial cortex is interrupted in only 1 place.
Multifragmentary pertrochanteric fractures (A2)	<ul style="list-style-type: none"> • With 1 or more intermediate fragments • Lateral wall may be incompetent 	Unstable	The fracture line can start laterally anywhere on the greater trochanter and runs towards the medial cortex which is typically broken in 2 places. This can result in the detachment of a third fragment which may include the lesser trochanter.
Intertrochanteric fractures (A3)	<ul style="list-style-type: none"> • Simple oblique fracture • Simple transverse fracture • Wedge or multifragmentary fracture 	Unstable	The fracture line passes between the 2 trochanters, above the lesser trochanter medially and below the crest of the vastus lateralis laterally.

Footnotes

AO/OTA: Arbeitsgemeinschaft für Osteosynthesefragen (German for "Association for the Study of Internal Fixation") / Orthopaedic Trauma Association

2 Possible grouping of different types of arthroplasty for hip fracture in adults

Implant category	Variable (articulation/fixation technique)	Implant subcategory	Examples ^a	Description
Total hip arthroplasty	Articulation	Femoral head and acetabular bearing surface materials	<ul style="list-style-type: none"> • Metal-on-polyethylene (MoP) • Ceramic-on-polyethylene (CoP) • Ceramic-on-ceramic (CoC) • Metal-on-metal (MoM) • Polyethylene material • Highly cross linked (HCL) • Not HCL 	Bearing surfaces may be grouped into hard (ceramic and metal) and soft (polyethylene variants). Arthroplasties exist with many of the possible combinations of these bearing surfaces.
		Femoral head size	<ul style="list-style-type: none"> • Large head ≥ 36 mm • Standard small head < 36 mm 	Over the development of hip arthroplasty different sizes of femoral head have been used from 22 mm to very large diameters approximating that of the native femoral head. The size of the head represents a compromise between stability and linear and volumetric wear at the articulation. The optimum size varies by indication and bearing materials. 36 mm is considered as a cut-off between standard and large sizes.
		Acetabular cup mobility	<ul style="list-style-type: none"> • Single • Dual 	A standard THA has a single articulating surface between the femoral head and acetabulum bearing surface. Alternative designs incorporate a further articulation within the structure of the femoral head.
	Fixation technique	Cemented	<ul style="list-style-type: none"> • Exeter Hip System • CPT Hip System 	Both components are cemented with polymethylmethacrylate bone cement that is inserted at the time of surgery. It sets hard and acts a grout between the prosthesis and the bone.
		Modern uncemented	<ul style="list-style-type: none"> • Corail Hip System • Avenir Hip System • Taperloc Hip System 	Neither component is cemented but rely on osseous integration forming a direct mechanical linkage between the bone and the implant. The femoral prosthesis may be coated with a substance such as hydroxyapatite which promotes bone growth into the prosthesis. Alternatively, the surface of the prosthesis may be macroscopically and microscopically roughened so that bone grows onto the surface of the implant. The acetabular component may be prepared similarly and may or may not be augmented with screws fixed into the pelvis.
		Hybrid	Combinations	The femoral stem is cemented and the acetabular cup is uncemented.
		Reverse hybrid	Combinations	The acetabular cup is cemented and the femoral stem is uncemented

Implant category	Variable (articulation/fixation technique)	Implant subcategory	Examples ^a	Description
Hemiarthroplasty	Articulation	Unipolar	<ul style="list-style-type: none"> • Thompson • Austin–Moore • Exeter Trauma Stem • Exeter Unitrax 	A single articulation between the femoral head and the native acetabulum. The femoral component can be a single ‘monoblock’ of alloy or be modular, assembled from component parts during surgery.
		Bipolar	<ul style="list-style-type: none"> • CPT modular bipolar • Exeter modular bipolar • Bateman • Monk 	The object of the second joint is to reduce acetabular wear. This type of prosthesis has a spherical inner metal head with a size between 22 to 36 mm in diameter. This fits into a polyethylene shell, which in turn is enclosed by a metal cap. There are a number of different types of prostheses with different stem designs.
	Fixation technique	First generation uncemented	<ul style="list-style-type: none"> • Thompson • Austin Moore 	These prostheses were designed before the development of polymethylmethacrylate bone cement and were therefore originally inserted as a ‘press fit’. Long-term stability through osseous integration was not part of the design concept.
		Cemented	<ul style="list-style-type: none"> • Thompson • Exeter Trauma Stem • Exeter Hip System • CPT Hip System 	The femoral stem is cemented with polymethylmethacrylate bone cement that is inserted at the time of surgery. It sets hard and acts a grout between the prosthesis and the bone.
		Modern uncemented	<ul style="list-style-type: none"> • Corail • Furlong • Avenir 	The femoral stem relies on osseous integration forming a direct mechanical linkage between the bone and the implant. A prosthesis may be coated with a substance such as hydroxyapatite, which promotes bone growth into the prosthesis. Alternatively, the surface of the prosthesis may be macroscopically and microscopically roughened so that bone grows onto the surface of the implant.

Footnotes

^aThis list is not exhaustive. In the Cochrane Review, we will include any implant tested in included trials not already listed here.

Abbreviations:

CoC: Ceramic-on-ceramic

CoP: Ceramic-on-polyethylene

CPT: collarless polished tapered

HCL: Highly cross linked

MoM: Metal-on-metal

MoP: Metal-on-polyethylene

THA: total hip arthroplasty

References to studies

Included studies

Excluded studies

Studies awaiting classification

Ongoing studies**Other references****Additional references****AAOS 2014**

American Academy of Orthopaedic Surgeons (AAOS). Management of hip fractures in the elderly. Evidence-based clinical practice guideline. Adopted by the American Academy of Orthopaedic Surgeons Board of Directors September 5, 2014. www.aaos.org/research/guidelines/HipFxGuideline.pdf (accessed 29 July 2019).

AO Foundation 2018

AO Foundation. Femur. Journal of Orthopaedic Trauma 2018;32(Suppl 1):S33–44.

Bardsley 2013

Smith P, Ariti C, Bardsley M. Nuffield Trust & Health Foundation: Focus on hip fracture. Trends in emergency admissions for fractured neck of femur, 2001 to 2011. www.nuffieldtrust.org.uk/research/focus-on-hip-fracture (accessed 9 January 2019).

Blundell 1998

Blundell CM, Parker MJ, Pryor GA, Hopkinson–Wooley J, Bonsall SS. An assessment of the AO classification of intracapsular fractures of the proximal femur. Journal of Bone and Joint Surgery – British Volume 1998; 80(4):679–83.

Christie 1994

Christie J, Burnett R, Potts HR, Pell AC. Echocardiography of transatrial embolism during cemented and uncemented hemiarthroplasty of the hip. Journal of Bone and Joint Surgery – British Volume 1994;76(3):409–12.

Cochrane 2018

Cochrane. Glossary. <https://community.cochrane.org/glossary> (accessed 2 November 2018).

Cooper 2011

Cooper C, Cole ZA, Holroyd CR, Earl SC, Harvey NC, Dennison EM, et al. Secular trends in the incidence of hip and other osteoporotic fractures. Osteoporosis International 2011;22(5):1277–88.

Court–Brown 2017

Court–Brown CM, Clement ND, Duckworth AD, Biant LC, McQueen MM. The changing epidemiology of fall-related fractures in adults. Injury 2017;48(4):819–24.

Covidence

Covidence [Computer program]. Version accessed 29 July 2019. Melbourne, Australia: Veritas Health Innovation, 2003. Available at covidence.org.

Deeks 2017

Deeks JJ, Higgins JPT, Altman DG (editors) on behalf of the Cochrane Statistical Methods Group. Chapter 9: Analysing data and undertaking meta-analyses. In: Higgins JPT, Churchill R, Chandler J, Cumpston MS (editors), Cochrane Handbook for Systematic Reviews of Interventions version 5.2.0 (updated June 2017), Cochrane, 2017. Available from www.training.cochrane.org/handbook.

Dhanwal 2011

Dhanwal DK, Dennison EM, Harvey NC, Cooper C. Epidemiology of hip fracture: worldwide geographic variation. Indian Journal of Orthopaedics 2011;45(1):15–22.

Evans 1949

Evans EM. The treatment of trochanteric fractures of the femur. Journal of Bone and Joint Surgery – British Volume 1949;31(B(2)):190–203.

Garden 1961

Garden RS. Low-angle fixation in fractures of the femoral neck. Bone & Joint Journal 1961;43–B(4):647–63.

GRADEpro GDT 2015

GRADEpro GDT [Computer program]. Version (assessed 9 January 2019). Hamilton (ON): McMaster University (developed by Evidence Prime, Inc.), 2015. Available at gradepr.org.

Griffin 2015

Griffin XL, Parsons N, Achten J, Fernandez M, Costa ML. Recovery of health-related quality of life in a United Kingdom hip fracture population: The Warwick Hip Trauma Evaluation—a prospective cohort study. Bone & Joint Journal 2015;97–B(3):372–82.

Guyatt 2008

Guyatt GH, Oxman AD, Vist GE, Kunz R, Falck-Ytter Y, Alonso-Coello P, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 2008;336(7650):924–6.

Haywood 2014

Haywood KL, Griffin XL, Achten J, Costa ML. Developing a core outcome set for hip fracture trials. *Bone and Joint Journal* 2014;96-B(8):1016–23.

Higgins 2011a

Higgins JPT, Altman DG. Chapter 8: Assessing risk of bias in included studies. In: Higgins JPT, Green S, editor(s). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from handbook.cochrane.org.

Higgins 2011b

Higgins JPT, Deeks JJ, Altman DG. Chapter 16: Special topics in statistics. In: Higgins JP, Green S, editor(s). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from handbook.cochrane.org.

Hopley 2010

Hopley C, Stengel D, Ekkernkamp A, Wich M. Primary total hip arthroplasty versus hemiarthroplasty for displaced intracapsular hip fractures in older patients: systematic review. *BMJ (Clinical research ed)* 2010;340:c2332.

Jensen 1980

Jensen JS. Classification of trochanteric fractures. *Acta Orthopaedica Scandinavica* 1980;51(5):803–10.

Johnell 2004

Johnell O, Kanis JA. An estimate of the worldwide prevalence, mortality and disability associated with hip fracture. *Osteoporosis International* 2004;15(11):897–902.

Kanis 2001

Kanis JA, Oden A, Johnell O, Jonsson B, de Laet C, Dawson A. The burden of osteoporotic fractures: a method for setting intervention thresholds. *Osteoporosis International* 2001;12(5):417–27.

Kanis 2012

Kanis JA, Odén A, McCloskey EV, Johansson H, Wahl DA, Cooper C. A systematic review of hip fracture incidence and probability of fracture worldwide. *Osteoporosis International* 2012;23(9):2239–56.

Karantana 2011

Karantana A, Boulton C, Bouliotis G, Shu KS, Scammell BE, Moran CG. Epidemiology and outcome of fracture of the hip in women aged 65 years and under: a cohort study. *Journal of Bone Joint Surgery – British Volume* 2011; 93(5):658–64.

Keating 2010

Keating J. Femoral neck fractures. In: Bucholz RW, Heckman JD, Court-Brown CM, Tornetta P III, editors(s). *Rockwood and Green's Fractures in Adults*. 7 edition. Vol. 2. Philadelphia: Wolters Kluwer Health/Lippincott Williams & Wilkins, 2010.

Lefebvre 2018

Lefebvre C, Glanville J, Briscoe S, Littlewood A, Marshall C, Metzendorf M-I, et al. Chapter 4: Searching for and selecting studies. Draft version (13 September 2018) for inclusion in Higgins JPT, Thomas J, Chandler J, Cumpton MS, Li T, Page MJ, Welch V (editors). *Cochrane Handbook for Systematic Reviews of Interventions*. London: Cochrane.

Lloyd-Jones 2015

Lloyd-Jones G. Trauma x-ray – lower limb. Hip fracture. www.radiologymasterclass.co.uk/tutorials/musculoskeletal/x-ray_trauma_lower_limb/hip_fracture_x-ray (accessed 10 August 2018).

Moher 2009

Moher D, Liberati A, Tetzlaff J, Altman DG; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Medicine* 2009;6(7):e1000097.

Neufeld 2016

Neufeld ME, O'Hara NN, Zhan M, Zhai Y, Broekhuysen HM, Lefevre KA, et al. Timing of hip fracture surgery and 30-day outcomes. *Orthopedics* 2016;39(6):361–8.

NHFD 2017

National Hip Fracture Database (NHFD). The National Hip Fracture Database (NHFD) annual report 2017. www.nhfd.co.uk/ (accessed 30 October 2018).

NICE 2011

National Clinical Guideline Centre (UK). The management of hip fracture in adults. NICE clinical guidelines, no. 124. London: Royal College of Physicians (UK); 2011. www.nice.org.uk/guidance/cg124/evidence/full-guideline-pdf-183081997 (accessed 30 October 2018).

ONS 2018

Office for National Statistics. Mid-2018 population estimates for the UK, England and Wales, Scotland and Northern Ireland. <https://bit.ly/2XBLgy6> (accessed 1 August 2019).

Ouzzani 2016

Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan—a web and mobile app for systematic reviews. *Systematic Reviews* 2016;5(1):210.

Papadimitriou 2017

Papadimitriou N, Tsilidis KK, Orfanos P, Benetou V, Ntzani EE, Soerjomataram I, et al. Burden of hip fracture using disability-adjusted life-years: a pooled analysis of prospective cohorts in the CHANCES consortium. *Lancet Public Health* 2017;2(5):e239–e46.

Parker 1991

Parker MJ, Anand JK. What is the true mortality of hip fractures? *Public Health* 1991;105(6):443–6.

Parker 1993

Parker MJ. Garden grading of intracapsular fractures: meaningful or misleading? *Injury* 1993;24(4):241–2.

Parker 1998

Parker MJ, Dynan Y. Is Pauwels classification still valid? *Injury* 1998;29(7):521–3.

Parker 1999

Parker MJ. Proximal femoral fractures. In: Pynsent PB, Fairbank JCT, Carr AJ, editors(s). *Classification of Musculo-Skeletal Trauma*. Oxford: Butterworth-Heinemann Publications, 1999.

Parker 2010

Parker MJ, Gurusamy KS, Azegami S. Arthroplasties (with and without bone cement) for proximal femoral fractures in adults. *Cochrane Database of Systematic Reviews* 2010, Issue 6. Art. No.: CD001706 DOI: 10.1002/14651858.CD001706.pub4.

Pauwels 1935

Pauwels F. Der Schenkelhalsbruch. Ein mechanisches Problem. In: *Gesammelte Abhandlungen zur funktionellen Anatomie des Bewegungsapparates*. Berlin: Heidelberg: Springer, 1935.

Perry 2016

Perry DC, Metcalfe D, Griffin XL, Costa ML. Inequalities in use of total hip arthroplasty for hip fracture: population based study. *BMJ* 2016;353:i2021.

Review Manager 2014

Review Manager 5 (RevMan 5) [Computer program]. Version 5.3. Copenhagen: Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

Rogmark 2018

Rogmark C, Kristensen MT, Viberg B, Ronnquist SS, Overgaard S, Palm H. Hip fractures in the non-elderly—Who, why and whither? *Injury* 2018;49(8):1445–50.

Sayers 2017

Sayers A, Whitehouse MR, Berstock JR, Harding KA, Kelly MB, Chessier TJ. The association between the day of the week of milestones in the care pathway of patients with hip fracture and 30-day mortality: findings from a prospective national registry – The National Hip Fracture Database of England and Wales. *BMC Medicine* 2017; 15:62.

Schünemann 2019

Schünemann HJ, Higgins JPT, Vist GE, Glasziou P, Akl EA, Skoetz N, et al. Chapter 14: Completing ‘Summary of findings’ tables and grading the certainty of the evidence. Draft version (29 January 2019) for inclusion in: Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA (editors). *Cochrane Handbook for Systematic Reviews of Interventions*. London: Cochrane.

SIGN 2009

Scottish Intercollegiate Guidelines Network (SIGN). Management of hip fracture in older people. A national clinical guideline. www.sign.ac.uk/assets/sign111.pdf (accessed 30 October 2018).

Sterne 2017

Sterne JAC, Egger M, Moher D, Boutron I (editors). Chapter 10: Addressing reporting biases. In: Higgins JPT, Churchill R, Chandler J, Cumpston MS (editors), *Cochrane Handbook for Systematic Reviews of Interventions* version 5.2.0 (updated June 2017), Cochrane, 2017. Available from www.training.cochrane.org/handbook.

Torio 2011

Torio CM, Andrews RM. National inpatient hospital costs: The most expensive conditions by Payer, 2011. Healthcare Cost and Utilization Project Statistical Brief #160. www.hcup-us.ahrq.gov/reports/statbriefs/sb160.pdf (accessed 9 January 2019).

Williamson 2017

Williamson S, Landeiro F, McConnell T, Fulford-Smith L, Javaid MK, Judge A, et al. Costs of fragility hip fractures globally: a systematic review and meta-regression analysis. *Osteoporosis International* 2017;28(10):2791–800.

Other published versions of this review

Classification pending references

Data and analyses

Figures

Sources of support

Internal sources

- National Institute for Health Research (NIHR) Oxford Biomedical Research Centre, Oxford, UK

External sources

- NIHR, UK
NIHR Systematic Review Programme Grant 16/114/15. A programme of high-priority reviews for the management of patients with hip fracture: a collaboration which can inform future healthcare policy guidance.
NIHR Cochrane Infrastructure funding to the Cochrane Bone, Joint and Muscle Trauma Group

Feedback

Appendices

1 MEDLINE search strategy

Ovid interface

- 1 exp Femoral Fractures/
- 2 ((hip or hips or cervical) adj5 (fracture\$ or break\$ or broke\$)).ti,ab,kf.
- 3 ((femoral\$ or femur\$ or acetabul\$) adj5 (fracture\$ or break\$ or broke\$)).ti,ab,kf.
- 4 ((intracapsular or intra-capsular or subcapital or sub-capital or transcervical or trans-cervical or basicervical or basi-cervical) adj5 (fracture\$ or break\$ or broke\$)).ti,ab,kf.
- 5 ((extracapsular or extra-capsular or trochant\$ or subtrochant\$ or pertrochant\$ or intertrochant\$) adj5 (fracture\$ or break\$ or broke\$)).ti,ab,kf.
- 6 (((head or neck or proximal) adj5 (fracture\$ or break\$ or broke\$)) and (femoral\$ or femur\$)).ti,ab,kf.
- 7 or/1–6
- 8 randomized controlled trial.pt.
- 9 controlled clinical trial.pt.
- 10 randomized.ab.
- 11 placebo.ab.
- 12 clinical trials as topic.sh.
- 13 randomly.ab.
- 14 trial.ti.
- 15 8 or 9 or 10 or 11 or 12 or 13 or 14
- 16 7 and 15
- 17 Arthroplasty, Replacement, Hip/ or Hip Prosthesis/
- 18 Arthroplasty, Replacement/ or Hemiarthroplasty/ or Joint Prosthesis/
- 19 (arthroplast\$ or hemiarthroplast\$).ti,ab,kf.
- 20 ((hip or hips) adj5 (replac\$ or prothes\$ or implant\$)).ti,ab,kf.
- 21 ((joint\$1 adj5 (replac\$ or prothes\$ or implant\$)) and (hip or hips or femur\$ or femoral\$ or acetabul\$)).ti,ab,kf.
- 22 or/17–21
- 23 fractures, bone/ or exp fracture dislocation/ or fractures, closed/ or fractures, comminuted/ or fractures,

compression/ or fractures, malunited/ or fractures, multiple/ or fractures, open/ or fractures, spontaneous/ or
 exp fractures, stress/ or fractures, ununited/ or intra-articular fractures/ or osteoporotic fractures/ or
 periprosthetic fractures/ (92406)
 24 fracture\$.ti,ab,kf.
 25 23 or 24
 26 22 and 25 and 15
 27 (pin or pins or nail or nails or screw or screws or plate or plates).ti,ab,kf.
 28 internal fixators/ or bone nails/ or bone plates/ or exp bone screws/
 29 (static adj (device\$1 or implant\$1)).ti,ab,kf.
 30 (dynamic adj (device\$1 or implant\$1)).ti,ab,kf.
 31 or/27-30
 32 ((hip or hips or femur\$ or femoral\$ or acetabul\$) and (fracture\$ or break\$ or broke\$)).ti,ab,kf.
 33 (hip or hips or femur\$ or femoral\$ or acetabul\$).ti,ab,kf. and (fractures, bone/ or exp fracture dislocation/ or
 fractures, closed/ or fractures, comminuted/ or fractures, compression/ or fractures, malunited/ or fractures,
 multiple/ or fractures, open/ or fractures, spontaneous/ or exp fractures, stress/ or fractures, ununited/ or
 intra-articular fractures/ or osteoporotic fractures/ or periprosthetic fractures/)
 34 or/32-33
 35 31 and 34 and 15
 36 16 or 26 or 35
 37 exp animals/ not humans/
 38 36 not 37