

On the categorical meaning of Hausdorff and Gromov distances, I

Andrei Akhvediani^{a,1}, Maria Manuel Clementino^{b,2}, Walter Tholen^{c,*,1}

^a Oxford University Computing Laboratory, Oxford OX1 3QD, United Kingdom

^b Department of Mathematics, University of Coimbra, 3001-454 Coimbra, Portugal

^c Department of Mathematics and Statistics, York University, Toronto, ON M3J 1P3, Canada

ARTICLE INFO

Article history:

Received 5 January 2009

Received in revised form 16 June 2009

Accepted 18 June 2009

MSC:

primary 18E40

secondary 18A99

Keywords:

Lawvere metric space

\mathcal{V} -category

\mathcal{V} -module

Hausdorff metric

Gromov metric

ABSTRACT

Hausdorff and Gromov distances are introduced and treated in the context of categories enriched over a commutative unital quantale \mathcal{V} . The Hausdorff functor which, for every \mathcal{V} -category X , provides the powerset of X with a suitable \mathcal{V} -category structure, is part of a monad on $\mathcal{V}\text{-Cat}$ whose Eilenberg–Moore algebras are order-complete. The Gromov construction may be pursued for any endofunctor K of $\mathcal{V}\text{-Cat}$. In order to define the Gromov “distance” between \mathcal{V} -categories X and Y we use \mathcal{V} -modules between X and Y , rather than \mathcal{V} -category structures on the disjoint union of X and Y . Hence, we first provide a general extension theorem which, for any K , yields a lax extension \bar{K} to the category $\mathcal{V}\text{-Mod}$ of \mathcal{V} -categories, with \mathcal{V} -modules as morphisms.

© 2009 Elsevier B.V. All rights reserved.

1. Introduction

The Hausdorff metric for (closed) subsets of a (compact) metric space has been recognized for a long time as an important concept in many branches of mathematics, and its origins reach back even beyond Hausdorff [10], to Pompeiu [15]; for a modern account, see [2]. It has gained renewed interest through Gromov’s work [9]. The Gromov–Hausdorff distance of two (compact) metric spaces is the infimum of their Hausdorff distances after having been isometrically embedded into any common larger space. There is therefore a notion of convergence for (isometry classes of compact) metric spaces which has not only become an important tool in analysis and geometry, but which has also provided the key instrument for the proof of Gromov’s existence theorem for a nilpotent subgroup of finite index in every finitely-generated group of polynomial growth [8].

By interpreting the (non-negative) distances $d(x, y)$ as $\text{hom}(x, y)$ and, hence, by rewriting the conditions

$$0 \geq d(x, x), \quad d(x, y) + d(y, z) \geq d(x, z) \quad (*)$$

as

$$k \rightarrow \text{hom}(x, x), \quad \text{hom}(x, y) \otimes \text{hom}(y, z) \rightarrow \text{hom}(x, z),$$

Lawvere [13] described metric spaces as categories enriched over the (small and “thin”) symmetric monoidal-closed category $\mathbb{P}_+ = ([0, \infty], \geq, +, 0)$, and in his paper [14] as well as in the Foreword of the electronic “reprint” of [13] he suggested

* Corresponding author.

E-mail addresses: andrei.akhvediani@comlab.ox.ac.uk (A. Akhvediani), mmc@mat.uc.pt (M.M. Clementino), tholen@mathstat.yorku.ca (W. Tholen).

¹ The authors acknowledge partial financial assistance from NSERC.

² The author acknowledges financial support from the Center of Mathematics of the University of Coimbra/FCT.




Fig. 1.

that the Hausdorff and Gromov metrics should be developed for an arbitrary symmetric monoidal-closed category $(\mathcal{V}, \otimes, k)$. In this paper we present notions of Hausdorff and Gromov distance for the case that \mathcal{V} is “thin”. Hence, we replace \mathbb{P}_+ by a commutative and unital *quantale* \mathcal{V} , that is: by a complete lattice which is also a commutative monoid $(\mathcal{V}, \otimes, k)$ such that the binary operation \otimes preserves suprema in each variable. Put differently, we try to give answers to questions of the type: which structure and properties of the (extended) non-negative real half-line allow for a meaningful treatment of Hausdorff and Gromov distances, and which are their appropriate carrier sets? We find that the guidance provided by enriched category theory [12] is almost indispensable for finding satisfactory answers, and that so-called *(bi-)modules* (or *distributors*) between \mathcal{V} -categories provide an elegant tool for the theory which may easily be overlooked without the categorical environment. Hence, our primary motivation for this work is the desire for a better understanding of the true essentials of the classical metric theory and its applications, rather than the desire for giving merely a more general framework which, however, may prove to be useful as well.

Since $(*)$ isolates precisely those conditions of a metric which lend themselves naturally to the hom interpretation, a discussion of the others seems to be necessary at this point; these are:

- $d(x, y) = d(y, x)$ (*symmetry*),
- $x = y$ whenever $d(x, y) = 0 = d(y, x)$ (*separation*),
- $d(x, y) < \infty$ (*finiteness*).

With the distance of a point x to a subset B of the metric space $X = (X, d)$ be given by $d(x, B) = \inf_{y \in B} d(x, y)$, the *non-symmetric Hausdorff distance* from a subset A to B is defined by

$$Hd(A, B) = \sup_{x \in A} d(x, B),$$

from which one obtains the classical *Hausdorff distance*

$$H^s d(A, B) = \max\{Hd(A, B), Hd(B, A)\}$$

by *enforced* symmetrization. But not only symmetry, but also separation and finiteness get lost under the rather natural passage from d to Hd . (If one thinks of $d(x, B)$ as the travel time from x to B , then $Hd(A, B)$ may be thought of as the time needed to evacuate everyone living in the area A to the area B .) In order to save them one usually restricts the carrier set from the entire powerset PX to the closed subsets of X (which makes $H^s d$ separated), or even to the non-empty compact subsets (which guarantees also finiteness). As in [11], we call a \mathbb{P}_+ -category an *L-metric space*, that is a set X equipped with a function $d : X \times X \rightarrow [0, \infty]$ satisfying $(*)$; a \mathbb{P}_+ -functor $f : (X, d) \rightarrow (X', d')$ is a non-expansive map, e.g. a map $f : X \rightarrow X'$ satisfying $d'(f(x), f(y)) \leq d(x, y)$ for all $x, y \in X$. That the underlying-set functor makes the resulting category *Met* *topological* over *Set* (see [7]) provides further evidence that properties $(*)$ are fundamental and are better considered separately from the others, even though symmetry (as a coreflective property) would not obstruct topologicity. But inclusion of (the reflective property of) separation would, and inclusion of (the neither reflective nor coreflective property of) finiteness would make for an even poorer categorical environment.

While from the categorical perspective symmetry seems to be artificially superimposed on the Hausdorff metric, it does play a crucial role for the Gromov distance, which becomes evident already when we look at the most elementary examples. Initially nothing prevents us from considering arbitrary *L-metric spaces* X, Y and putting

$$GH(X, Y) = \inf_Z Hd_Z(X, Y),$$

where Z runs through all *L-metric spaces* Z into which both X and Y are isometrically embedded. But for $X = \{p\}$ a singleton set and $Y = \{x, y, z\}$ three equidistant points, with all distances 1, say, for every $\varepsilon > 0$ we can make $Z = X \sqcup Y$ a (proper) metric space, with $d(p, x) = d(x, p) = \varepsilon$ and all other non-zero distances 1. Then $Hd_Z(X, Y) = \varepsilon$, and $GH(X, Y) = 0$ follows. One has also $GH(Y, X) = 0$ but here one needs non-symmetric (but still separated) structures: put $d(x, p) = d(y, p) = d(z, p) = \varepsilon$, but let the reverse distances be 1. Hence, even a *posteriori* symmetrization leads to a trivial distance between non-isomorphic spaces. However, there are two ways of *a priori* symmetrization which both yield the intuitively desired result $\frac{1}{2}$ for the Gromov distance in this example (see Fig. 1): One way is by restricting the range of

the infimum in the definition of $GH(X, Y)$ to symmetric L -metric spaces, which seems to be natural when X and Y are symmetric. (Indeed, if for our example spaces one assumes $Hd_Z(Y, X) < \frac{1}{2}$ with d_Z symmetric, then the triangle inequality would be violated: $1 \leq d(x, p) + d(p, y) < \frac{1}{2} + \frac{1}{2}$.) The other way “works” also for non-symmetric X and Y ; one simply puts

$$GH^s(X, Y) = \inf_Z H^s d_Z(X, Y),$$

with Z running as in $GH(X, Y)$. (When $Hd_Z(Y, X) \leq \frac{1}{2}$, then

$$\frac{1}{2} = 1 - \frac{1}{2} \leq \min\{d(p, x), d(p, y), d(p, z)\} = Hd_Z(X, Y) \leq H^s d_Z(X, Y),$$

and when $Hd_Z(Y, X) \geq \frac{1}{2}$, then trivially $\frac{1}{2} \leq H^s d_Z(X, Y)$.)

Having recognized H (and H^s) as endofunctors of \mathbf{Met} , these considerations suggest that G is an “operator” on such endofunctors. But in order to “compute” its values, one needs to “control” the spaces Z in their defining formula, and here is where modules come in. (A module between L -metric spaces generalizes a non-expansive map just like a relation generalizes a mapping between sets.) A module from X to Y corresponds to an L -metric that one may impose on the disjoint union $X \sqcup Y$. To take advantage of this viewpoint, it is necessary to extend H from non-expansive maps to modules (leaving its operation on objects unchanged) to become a lax functor \tilde{H} . Hence, $GH(X, Y)$ may then be more compactly defined using an infimum that ranges just over the hom-set of modules from X to Y .

In Section 2 we give a brief overview of the needed tools from enriched category theory, in the highly simplified context of a quantale \mathcal{V} . The purpose of Section 3 is to establish a general extension theorem for endofunctors of $\mathcal{V}\text{-Cat}$, so that they can act on \mathcal{V} -modules rather than just on \mathcal{V} -functors. In Sections 4 and 5 we consider the Hausdorff monad of $\mathcal{V}\text{-Cat}$ and its lax extension to \mathcal{V} -modules, and we determine the Eilenberg–Moore category in both cases. The Gromov “distance” is considered for a fairly general range of endofunctors of $\mathcal{V}\text{-Cat}$ in Section 6, and the resulting Gromov “space” of isomorphism classes of \mathcal{V} -categories is presented as a large colimit. For the endofunctor H^s , in Section 8 this large “space” is shown to carry internal monoid structures in the monoidal category $\mathcal{V}\text{-CAT}$ which allow us to consider H as an internal homomorphism. The effects of symmetrization and the status of separation are discussed in Sections 7 and 9. The fundamental question of transfer of (Cauchy-)completeness from X to HX , as well as the question of completeness of suitable subspaces of the Gromov “space” will be considered in the second part of this paper.

The reader is reminded that, since \mathbb{P}_+ carries the natural \geq as its order, in the context of a general quantale \mathcal{V} the natural infima and suprema of \mathbb{P}_+ appear as joins (\bigvee) and meets (\bigwedge) in \mathcal{V} . While this may appear to be irritating initially, it reflects in fact the logical viewpoint dictated by the elementary case $\mathcal{V} = 2 = (\{\perp < \top\}, \wedge, \top)$, and it translates back well even in the metric case. (For example, if we write the sup-metric d of the real function space $C(X)$ as $d(f, g) = \bigwedge_{x \in X} |f(x) - g(x)|$, then the statement

$$d(f, g) = 0 \iff \text{for all } x \in X: f(x) = g(x)$$

seems to read off the defining formula more directly.)

There are a number of authors who have previously considered the Hausdorff metric from a categorical perspective. We mention here in particular the paper [3] as it triggered subsequent work in theoretical computer science, as well as the papers [16] and [5], preliminary work for which was first presented independently from each other in 2003. While the work presented in this paper first began to take shape when, aimed with her knowledge of the treatment of the Hausdorff metric in [5], Clementino visited Tholen in the Spring of 2008, which then gave rise to a much more comprehensive study by Akhmediani in his Master's thesis [1] that contains many elements of the current work, precursors of it go in fact back to a visit by Richard Wood to Tholen in 2001. However, the attempt to work immediately with a (non-thin) symmetric monoidal-closed category proved to be too difficult at the time.

2. Quantale-enriched categories

Throughout this paper, \mathcal{V} is a commutative, unital quantale. Hence, \mathcal{V} is a complete lattice with a commutative, associative binary operation \otimes and a \otimes -neutral element k , such that \otimes preserves arbitrary suprema in each variable. Our paradigmatic examples

$$2 = (\{\perp < \top\}, \wedge, \top) \quad \text{and} \quad \mathbb{P}_+ = ([0, \infty], \geq, +, 0)$$

were already considered by Lawvere [13]; they serve to provide both an order-theoretic and a metric intuition for the theory.

A \mathcal{V} -relation r from a set X to a set Y , written as $r: X \rightrightarrows Y$, is simply a function $r: X \times Y \rightarrow \mathcal{V}$. Its composition with $s: Y \rightrightarrows Z$ is given by

$$(s \cdot r)(x, z) = \bigvee_{y \in Y} r(x, y) \otimes s(y, z).$$

This defines a category $\mathcal{V}\text{-Rel}$, and there is an obvious functor $\mathbf{Set} \rightarrow \mathcal{V}\text{-Rel}$ which assigns to a mapping $f: X \rightarrow Y$ its \mathcal{V} -graph $f_\circ: X \rightrightarrows Y$ with $f_\circ(x, y) = k$ if $f(x) = y$, and $f_\circ(x, y) = \perp$ otherwise. This functor is faithful only if $k > \perp$, which we will assume henceforth, writing just f for f_\circ . There is an involution $(\)^\circ: \mathcal{V}\text{-Rel}^{\text{op}} \rightarrow \mathcal{V}\text{-Rel}$ which sends

$r : X \rightarrow Y$ to $r^\circ : Y \rightarrow X$ with $r^\circ(y, x) = r(x, y)$. With the pointwise order of its hom-sets, $\mathcal{V}\text{-Rel}$ becomes order-enriched, e.g. a 2-category, and mappings $f : X \rightarrow Y$ become maps in the 2-categorical sense:

$$1_X \leq f^\circ \cdot f, f \cdot f^\circ \leq 1_Y.$$

A \mathcal{V} -category $X = (X, a)$ is a set X with a \mathcal{V} -relation $a : X \rightarrow X$ satisfying $1_X \leq a$ and $a \cdot a \leq a$; elementwise this means

$$k \leq a(x, x) \quad \text{and} \quad a(x, y) \otimes a(y, z) \leq a(x, z).$$

A \mathcal{V} -functor $f : (X, a) \rightarrow (Y, b)$ is a map $f : X \rightarrow Y$ with $f \cdot a \leq b \cdot f$, or equivalently

$$a(x, y) \leq b(f(x), f(y))$$

for all $x, y \in X$. The resulting category $\mathcal{V}\text{-Cat}$ becomes the category Ord of (pre)ordered sets and monotone maps for $\mathcal{V} = 2$ and the category Met of L -metric spaces for $\mathcal{V} = \mathbb{P}_+$.

$\mathcal{V}\text{-Cat}$ has a symmetric monoidal-closed structure, given by

$$(X, a) \otimes (Y, b) = (X \times Y, a \otimes b), \quad X \multimap Y = (\mathcal{V}\text{-Cat}(X, Y), c)$$

with

$$a \otimes b((x, y), (x', y')) = a(x, x') \otimes b(y, y'),$$

$$c(f, g) = \bigwedge_{x \in X} b(f(x), g(x)).$$

Note that $X \otimes Y$ must be distinguished from the Cartesian product $X \times Y$ whose structure is $a \times b$ with

$$a \times b((x, y), (x', y')) = a(x, x') \wedge b(y, y').$$

\mathcal{V} itself is a \mathcal{V} -category with its “internal hom” \multimap , given by

$$z \leq x \multimap y \iff z \otimes x \leq y$$

for all $x, y, z \in \mathcal{V}$. The morphism $2 \rightarrow \mathcal{V}$ of quantales has a right adjoint $\mathcal{V} \rightarrow 2$ that sends $v \in \mathcal{V}$ to \top precisely when $v \geq k$. Hence, there is an induced functor

$$\mathcal{V}\text{-Cat} \rightarrow \text{Ord}$$

which provides a \mathcal{V} -category with the order

$$x \leq y \iff k \leq a(x, y).$$

Since

$$\mathcal{V}\text{-Rel}(X, Y) = \mathcal{V}^{X \times Y} = (X \times Y) \multimap \mathcal{V}$$

is a \mathcal{V} -category (as a product of $(X \times Y)$ -many copies of \mathcal{V} , or as a “function space” with X, Y discrete), it is easy to show that $\mathcal{V}\text{-Rel}$ is $(\mathcal{V}\text{-Cat})$ -enriched, e.g.

$$\begin{aligned} E &\rightarrow \mathcal{V}\text{-Rel}(X, X), & \mathcal{V}\text{-Rel}(X, Y) \otimes \mathcal{V}\text{-Rel}(Y, Z) &\rightarrow \mathcal{V}\text{-Rel}(X, Z), \\ * &\mapsto 1_X, & (r, s) &\mapsto s \cdot r \end{aligned}$$

are \mathcal{V} -functors (where $E = (\{*\}, k)$ is the \otimes -unit in $\mathcal{V}\text{-Cat}$).

3. Modules, extension theorem

For \mathcal{V} -categories $X = (X, a), Y = (Y, b)$ a \mathcal{V} -(bi)module (also: \mathcal{V} -distributor, \mathcal{V} -profunctor) φ from X to Y , written as $\varphi : X \rightleftarrows Y$, is a \mathcal{V} -relation $\varphi : X \rightarrow Y$ with $\varphi \cdot a \leq \varphi$ and $b \cdot \varphi \leq \varphi$, that is

$$a(x', x) \otimes \varphi(x, y) \leq \varphi(x', y) \quad \text{and} \quad \varphi(x, y) \otimes b(y, y') \leq \varphi(x, y')$$

for all $x, x' \in X, y, y' \in Y$. For $\varphi : X \rightleftarrows Y$ one actually has $\varphi \cdot a = \varphi = b \cdot \varphi$, so that $1_X^* := a$ plays the role of the identity morphism in the category $\mathcal{V}\text{-Mod}$ of \mathcal{V} -categories (as objects) and \mathcal{V} -modules (as morphisms). It is easy to show that a \mathcal{V} -relation $\varphi : X \rightleftarrows Y$ is a \mathcal{V} -module if, and only if, $\varphi : X^{\text{op}} \otimes Y \rightarrow \mathcal{V}$ is a \mathcal{V} -functor (see [6]); here $X^{\text{op}} = (X, a^\circ)$ for $X = (X, a)$. Hence,

$$\mathcal{V}\text{-Mod}(X, Y) = (X^{\text{op}} \otimes Y) \multimap \mathcal{V}.$$

In particular $\mathcal{V}\text{-Mod}$ is (like $\mathcal{V}\text{-Rel}$) $\mathcal{V}\text{-Cat}$ -enriched. Also, $\mathcal{V}\text{-Mod}$ inherits the 2-categorical structure from $\mathcal{V}\text{-Rel}$, just via pointwise order.

Every \mathcal{V} -functor $f : X \rightarrow Y$ induces adjoint \mathcal{V} -modules

$$f_* \dashv f^* : Y \rightleftarrows X$$

with $f_* := b \cdot f$, $f^* := f \circ b$ (in $\mathcal{V}\text{-Rel}$). Hence, there are functors

$$(-)_* : \mathcal{V}\text{-Cat} \rightarrow \mathcal{V}\text{-Mod}, \quad (-)^* : \mathcal{V}\text{-Cat}^{\text{op}} \rightarrow \mathcal{V}\text{-Mod}$$

which map objects identically. $\mathcal{V}\text{-Cat}$ becomes order-enriched (a 2-category) via

$$f \leq g \iff f^* \leq g^* \iff \forall x: f(x) \leq g(x).$$

The \mathcal{V} -functor $f : X \rightarrow Y$ is *fully faithful* if $f^* \cdot f_* = 1_X^*$; equivalently, if $a(x, x') = b(f(x), f(x'))$ for all $x, x' \in X$.

Via

$$\frac{\varphi : X \twoheadrightarrow Y}{\frac{X^{\text{op}} \otimes Y \rightarrow \mathcal{V}}{y_\varphi : Y \rightarrow (X^{\text{op}} \multimap \mathcal{V}) =: \hat{X}}},$$

every \mathcal{V} -module φ corresponds to its *Yoneda mate* y_φ in $\mathcal{V}\text{-Cat}$. In particular, $a = 1_X^*$ corresponds to the Yoneda functor

$$y_X = y_{1_X^*} : X \rightarrow \hat{X}.$$

For every \mathcal{V} -functor $f : X^{\text{op}} \rightarrow \mathcal{V}$ and $x \in X$ one has

$$1_{\hat{X}}^*(y_X(x), f) = f(x) \quad (\text{Yoneda lemma}).$$

In particular, $1_{\hat{X}}^*(y_X(x), y_X(x')) = a(x, x')$, i.e. y_X is fully faithful.

The correspondence between φ and y_φ gives:

Proposition 1. $(-)^* : (\mathcal{V}\text{-Cat})^{\text{op}} \rightarrow \mathcal{V}\text{-Mod}$ has a left adjoint $(\hat{-})$, given by

$$\hat{\varphi}(s)(x) = \bigvee_{y \in Y} \varphi(x, y) \otimes s(y)$$

for all $\varphi : X \twoheadrightarrow Y$, $s \in \hat{Y}$, $x \in X$.

Proof. Under the correspondence

$$\frac{\varphi : X \twoheadrightarrow Y}{\Phi : Y \rightarrow \hat{X}}$$

given by $\varphi(x, y) = \Phi(y)(x)$, $\Phi = 1_{\hat{X}}$ gives the unit $\eta_X : X \twoheadrightarrow \hat{X}$ of the adjunction, with

$$\eta_X(x, t) = t(x)$$

for all $x \in X$, $t \in \hat{X}$. Note that one has $\eta_X = (y_X)_*$, by the Yoneda lemma. We must confirm that y_φ is indeed the unique \mathcal{V} -functor $\Phi : Y \rightarrow \hat{X}$ with $\Phi^* \cdot \eta_X = \varphi$. But any such Φ must satisfy

$$\begin{aligned} \varphi(x, y) &= (\Phi^* \cdot (y_X)_*)(x, y) \\ &= \bigvee_{t \in \hat{X}} 1_{\hat{X}}^*(y_X(x), t) \otimes 1_{\hat{X}}^*(t, \Phi(y)) \\ &\leq \bigvee_{t \in \hat{X}} 1_{\hat{X}}^*(y_X(x), \Phi(y)) \\ &\leq \Phi(y)(x) \\ &\leq 1_{\hat{X}}^*(y_X(x), y_X(x)) \otimes 1_{\hat{X}}^*(y_X(x), \Phi(y)) \\ &\leq (\Phi^* \cdot (y_X)_*)(x, y) \\ &= \varphi(x, y) \end{aligned}$$

for all $x \in X$, $y \in Y$. Hence, necessarily $\Phi = y_\varphi$, and the same calculation shows $\varphi = (y_\varphi)^* \cdot \eta_X$. Now, $\hat{\varphi} : \hat{Y} \rightarrow \hat{X}$ is the \mathcal{V} -functor corresponding to $\eta_Y \cdot \varphi$, hence

$$\hat{\varphi}(s)(x) = (\eta_Y \cdot \varphi)(x, s) = \bigvee_{y \in Y} \varphi(x, y) \otimes s(y),$$

for all $s \in \hat{Y}$, $x \in X$. \square

Remarks 1. (1) For $\varphi : X \rightrightarrows Y$, the \mathcal{V} -functor $\hat{\varphi}$ may also be described as the left Kan extension of $y_\varphi : Y \rightarrow \hat{X}$ along $y_Y : Y \rightarrow \hat{Y}$.

(2) The adjunction of Proposition 1 is in fact 2-categorical. It therefore induces a 2-monad $\mathbb{P}_{\mathcal{V}} = (P_{\mathcal{V}}, y, m)$ on $\mathcal{V}\text{-Cat}$, with

$$P_{\mathcal{V}}X = \hat{X} = X \multimap \mathcal{V}, \quad P_{\mathcal{V}}f = \widehat{f^*} : \hat{X} \rightarrow \hat{Y}$$

for $f : X \rightarrow Y = (Y, b)$, where

$$\widehat{f^*}(t)(y) = \bigvee_{x \in X} b(y, f(x)) \otimes t(x)$$

for $t \in \hat{X}$, $y \in Y$. This monad is of Kock–Zöberlein type, i.e. one has

$$\widehat{y_X^*} \leq y_X : \hat{X} \rightarrow \hat{X}.$$

In fact, for all $x, y \in X = (X, a)$, and $t, s \in \hat{X}$ one has $a(x, y) \leq s(x) \multimap s(y)$, hence

$$t(y) \otimes (t(y) \multimap a(x, y)) \otimes s(x) \leq a(x, y) \otimes s(x) \leq s(y),$$

which gives

$$\begin{aligned} \widehat{y_X^*}(s)(t) &= \bigvee_x y_X^*(t, x) \otimes s(x) \\ &= \bigvee_x \bigwedge_y (t(y) \multimap a(x, y)) \otimes s(x) \\ &\leq \bigwedge_y t(y) \multimap s(y) = y_X(s)(t). \end{aligned}$$

(3) The adjunction of Proposition 1 induces also a monad on $\mathcal{V}\text{-Mod}$ which we will not consider further in this paper. But see Section 5 below.

(4) Because of (2), the Eilenberg–Moore category

$$(\mathcal{V}\text{-Cat})^{\mathbb{P}_{\mathcal{V}}}$$

has \mathcal{V} -categories X as objects which come equipped with a \mathcal{V} -functor $\alpha : \hat{X} \rightarrow X$ with $\alpha \cdot y_X = 1_X$ and $1_{\hat{X}} \leq y_X \cdot \alpha$, e.g. \mathcal{V} -categories X for which y_X has a left adjoint. These are known to be the \mathcal{V} -categories that have all *weighted colimits* (see [12]), with α providing a choice of such colimits. Morphisms in $(\mathcal{V}\text{-Cat})^{\mathbb{P}_{\mathcal{V}}}$ must preserve the (chosen) weighted colimits.

(5) In case $\mathcal{V} = 2$, $\mathbb{P}_{\mathcal{V}}X$ can be identified with the set $P_{\downarrow}X$ of down-closed subsets of the (pre)ordered set X , and the Yoneda functor $X \rightarrow P_{\downarrow}X$ sends x to its down-closure $\downarrow x$. Note that $P_{\downarrow}X$ is the ordinary powerset PX of X when X is discrete. $\text{Ord}^{\mathbb{P}_{\downarrow}}$ has complete ordered sets as objects, and its morphisms must preserve suprema. Hence, this is the category *Sup* of so-called *sup-lattices* (with no anti-symmetry condition).

Next we prove a general extension theorem for endofunctors of $\mathcal{V}\text{-Cat}$. While maintaining its effect on objects, we wish extend any functor K defined for \mathcal{V} -functors to \mathcal{V} -modules. To this end we observe that for a \mathcal{V} -module $\varphi : X \rightrightarrows Y$, the left triangle of

$$\begin{array}{ccc} & \hat{Y} & \\ y_Y \nearrow & & \searrow \hat{\varphi} \\ Y & \xrightarrow{y_\varphi} & \hat{X} \end{array} \quad \begin{array}{ccc} & \hat{Y} & \\ y_Y \nearrow & & \searrow \hat{\varphi} \\ Z & \xrightarrow{y_Y \cdot \varphi} & \hat{X} \end{array}$$

commutes, since y_Y is the counit of the adjunction of Proposition 1. More generally, the right triangle commutes for every $\psi : Y \rightrightarrows Z$.

Theorem 1 (Extension theorem). For every functor $K : \mathcal{V}\text{-Cat} \rightarrow \mathcal{V}\text{-Cat}$,

$$\tilde{K}\varphi := (KX \xrightarrow{(Ky_X)^*} K\hat{X} \xrightarrow{(Ky_\varphi)^*} KY)$$

defines a lax functor $\tilde{K} : \mathcal{V}\text{-Mod} \rightarrow \mathcal{V}\text{-Mod}$ which coincides with K on objects. Moreover, if K preserves full fidelity of \mathcal{V} -functors, the diagram

$$\begin{array}{ccc} \mathcal{V}\text{-Mod} & \xrightarrow{\tilde{K}} & \mathcal{V}\text{-Mod} \\ (-)^* \uparrow & & \uparrow (-)^* \\ (\mathcal{V}\text{-Cat})^{\text{op}} & \xrightarrow{K^{\text{op}}} & (\mathcal{V}\text{-Cat})^{\text{op}} \end{array}$$

commutes.

Proof. Lax functoriality of \tilde{K} follows from

$$\begin{aligned} \tilde{K}(1_X^*) &= (K y_X)^* \cdot (K y_X)_* \geq 1_{KX}^*, \\ \tilde{K}(\psi \cdot \varphi) &= (K y_{\psi \cdot \varphi})^* \cdot (K y_X)_* \\ &= (K y_\psi)^* \cdot (K \hat{\varphi})^* \cdot (K y_X)_* \\ &\geq (K y_\psi)^* \cdot (K y_Y)_* \cdot (K y_\varphi)^* \cdot (K y_X)_* \\ &= \tilde{K}\psi \cdot \tilde{K}\varphi, \end{aligned}$$

since $y_\varphi = \hat{\varphi} \cdot y_Y$ implies $(K y_\varphi)^* = (K y_Y)^* \cdot (K \hat{\varphi})^*$, hence $(K \hat{\varphi})^* \geq (K y_Y)_* \cdot (K y_\varphi)^*$ by adjunction.

For a \mathcal{V} -functor $f : X \rightarrow Y$, the triangle

$$\begin{array}{ccc} & Y & \\ f \nearrow & & \searrow y_Y \\ X & \xrightarrow{y_{f*}} & \hat{Y} \end{array}$$

commutes, so that

$$\tilde{K}(f^*) = (K y_{f*})^* \cdot (K y_Y)_* = (K f)^* (K y_Y)^* (K y_Y)_* \geq (K f)^*,$$

and one even has $\tilde{K}(f^*) = (K f)^*$ if K preserves the full fidelity of y_Y . \square

4. The Hausdorff monad on $\mathcal{V}\text{-Cat}$

Let $X = (X, a)$ be a \mathcal{V} -category. Then $\hat{X} = (X^{\text{op}} \multimap \mathcal{V}) = P_{\mathcal{V}}X$ is closed under suprema formed in the product \mathcal{V}^X ; hence, like \mathcal{V} it is a sup-lattice. Consequently, the Yoneda functor $y_X : X \rightarrow \hat{X}$ factors uniquely through the free sup-lattice PX , by a sup-preserving map $\mathcal{Y}_X : PX \rightarrow P_{\mathcal{V}}X$:

$$\begin{array}{ccc} X & \xrightarrow{\{-\}} & PX \\ & \searrow y_X & \downarrow \mathcal{Y}_X \\ & & P_{\mathcal{V}}X \end{array} \quad \begin{array}{c} B \\ \downarrow \\ a(-, B) \end{array}$$

where

$$a(x, B) = \bigvee_{y \in B} a(x, y)$$

for all $x \in X$, $B \subseteq X$. We can provide the set PX with a \mathcal{V} -category structure h_X which it inherits from $P_{\mathcal{V}}X$ (since the forgetful functor $\mathcal{V}\text{-Cat} \rightarrow \text{Set}$ is a fibration, even a topological functor, see [7]). Hence, for subsets $A, B \subseteq X$ one puts

$$h_X(A, B) = \bigwedge_{z \in X} a(z, A) \multimap a(z, B).$$

Lemma 1.

$$h_X(A, B) = \bigwedge_{x \in A} \bigvee_{y \in B} a(x, y).$$

Proof. From $k \leq a(x, A)$ for all $x \in A$ one obtains

$$h_X(A, B) \leq \bigwedge_{x \in A} a(x, A) \multimap a(x, B) \leq \bigwedge_{x \in A} k \multimap a(x, B) = \bigwedge_{x \in A} a(x, B).$$

Conversely, with $v := \bigwedge_{x \in A} \bigvee_{y \in B} a(x, y)$, we must show $v \leq a(z, A) \multimap a(z, B)$ for all $z \in X$. But since for every $x \in A$

$$a(z, x) \otimes v \leq a(z, x) \otimes \bigvee_{y \in B} a(x, y) = \bigvee_{y \in B} a(z, x) \otimes a(x, y) \leq a(z, B),$$

one concludes $a(z, A) \otimes v \leq a(z, B)$, as desired. \square

For a \mathcal{V} -functor $f : X \rightarrow Y = (Y, b)$ one now concludes easily

$$h_X(A, B) \leq \bigwedge_{x \in A} \bigvee_{y \in B} b(f(x), f(y)) = h_Y(f(A), f(B))$$

for all $A, B \subseteq X$. Consequently, with

$$HX = (PX, h_X), \quad Hf : HX \rightarrow HY, \quad A \mapsto f(A),$$

one obtains a (2-)functor H which makes the diagram

$$\begin{array}{ccc} \mathcal{V}\text{-Cat} & \xrightarrow{H} & \mathcal{V}\text{-Cat} \\ \downarrow & & \downarrow \\ \text{Set} & \xrightarrow{P} & \text{Set} \end{array}$$

commute. Actually, one has the following theorem:

Theorem 2. The powerset monad $\mathbb{P} = (P, \{-\}, \bigcup)$ can be lifted along the forgetful functor $\mathcal{V}\text{-Cat} \rightarrow \text{Set}$ to a monad \mathbb{H} of $\mathcal{V}\text{-Cat}$ of Kock–Zöberlein type.

Proof. For a \mathcal{V} -category X , $x \mapsto \{x\}$ gives a fully faithful \mathcal{V} -functor $\{-\} : X \rightarrow HX$. In order to show that

$$\bigcup : HHX \rightarrow HX, \quad \mathcal{A} \mapsto \bigcup \mathcal{A},$$

is a \mathcal{V} -functor, it suffices to verify that for all $x \in A \in \mathcal{A} \in HHX$ and $B \in HHX$ one has

$$h_{HX}(\mathcal{A}, B) \leq a\left(x, \bigcup B\right).$$

But for all $B \in \mathcal{B}$ we have

$$h_X(A, B) \leq a(x, B) \leq a\left(x, \bigcup B\right),$$

so that

$$h_{HX}(\mathcal{A}, B) \leq \bigvee_{B \in \mathcal{B}} h_X(A, B) \leq a\left(x, \bigcup B\right).$$

The induced order of HX is given by

$$A \leq B \iff \forall x \in A: k \leq a(x, B),$$

and that of HHX by

$$\mathcal{A} \leq \mathcal{B} \iff \forall A \in \mathcal{A}: k \leq \bigvee_{B \in \mathcal{B}} h_X(A, B).$$

Hence, from $k \leq a(x, A) = h_X(\{x\}, A)$ for all $A \in HX$ one obtains

$$\{\{x\} \mid x \in A\} \leq \{A\}$$

in HHX , which means $H\{-\}_X \leq \{-\}_{HX}$, i.e., \mathbb{H} is Kock–Zöberlein. \square

Remarks 2. (1) By definition, $h_X(A, B)$ depends only on $a(-, A)$, $a(-, B)$. Hence, if we put

$$\begin{aligned}\Downarrow_X B &:= \{x \in X \mid \{x\} \leq B\} = \{x \in X \mid \downarrow x \leq B\} \\ &= \{x \in X \mid k \leq a(x, B)\},\end{aligned}$$

from $B \subseteq \downarrow_X B$ one trivially has $a(z, B) \leq a(z, \downarrow_X B)$ for all $z \in X$, but also

$$\begin{aligned}a(z, \downarrow_X B) &= \bigvee_{x \in \downarrow_X B} a(z, x) \otimes k \\ &\leq \bigvee_{z \in \downarrow_X B} \bigvee_{y \in B} a(z, x) \otimes a(x, y) \leq a(z, B).\end{aligned}$$

Consequently,

$$h_X(A, B) = h_X(\downarrow_X A, \downarrow_X B).$$

This equation also implies $\downarrow_X \downarrow_X B = \downarrow_X B$.

(2) $\downarrow_X B$ of (1) must not be confused with the down-closure $\downarrow_X B$ of B in X w.r.t. the induced order of X , e.g. with

$$\downarrow_X B = \{x \in X \mid \exists y \in B, x \leq y\} = \{x \in X \mid \exists y \in B (k \leq a(x, y))\}.$$

In general, $B \subseteq \downarrow_X B \subseteq \Downarrow_X B$. While $\downarrow_X B = \Downarrow_X B$ for $\mathcal{V} = 2$, the two sets are generally distinct even for $\mathcal{V} = \mathbb{P}_+$.

(3) In the induced order of HX one has

$$A \leq B \iff A \subseteq \Downarrow_X B.$$

Hence, if we restrict HX to

$$H_{\Downarrow} X := \{B \subseteq X \mid B = \downarrow_X B\},$$

the induced order of $H_{\Downarrow} X$ is simply the inclusion order. H_{\Downarrow} becomes a functor $H_{\Downarrow} : \mathcal{V}\text{-Cat} \rightarrow \mathcal{V}\text{-Cat}$ with

$$(H_{\Downarrow} f)(A) = \downarrow_Y f(A)$$

for all $A \in H_{\Downarrow} X$, and there is a lax natural transformation $\iota : H_{\Downarrow} \rightarrow H$ given by inclusion functions. Like H , also H_{\Downarrow} carries a monad structure, given by

$$\begin{aligned}X &\rightarrow H_{\Downarrow} X, \quad x \mapsto \downarrow_X x = \downarrow_X x, \\ H_{\Downarrow} H_{\Downarrow} X &\rightarrow H_{\Downarrow} X, \quad B \mapsto \downarrow_X \left(\bigcup B \right).\end{aligned}$$

In this way $\iota : \mathbb{H}_{\Downarrow} \rightarrow \mathbb{H}$ becomes a lax monad morphism.

(4) By definition, \mathcal{J}_X is fully faithful. Hence, HX carries the largest \mathcal{V} -category structure making $\mathcal{J}_X : HX \rightarrow P_{\mathcal{V}} X$ a \mathcal{V} -functor. Equivalently, this is the largest \mathcal{V} -category structure making

$$\delta_X : X \rightleftarrows HX$$

with $\delta(x, B) = a(x, B)$ a \mathcal{V} -module.

(5) $\mathcal{J}_X : HX \rightarrow P_{\mathcal{V}} X$ defines a morphism $\mathbb{H} \rightarrow P_{\mathcal{V}}$ of monads. Indeed, the left diagram of

$$\begin{array}{ccc} X & & H H X \\ \swarrow \{-\} & \searrow y_X & \downarrow \cup \\ H X & \xrightarrow{\mathcal{J}_X} & P_{\mathcal{V}} X \end{array} \quad \begin{array}{ccccc} H H X & \xrightarrow{H \mathcal{J}_X} & H P_{\mathcal{V}} X & \xrightarrow{\mathcal{J}_X} & P_{\mathcal{V}} P_{\mathcal{V}} X \\ \downarrow \cup & & \downarrow m_X & & \downarrow m_X \\ H X & \xrightarrow{\mathcal{J}_X} & P_{\mathcal{V}} X & & P_{\mathcal{V}} X \end{array}$$

commutes trivially, and for the right one first observes that $m_X : \hat{\hat{X}} \rightarrow \hat{X}$ is defined by

$$m_X(\tau)(x) = \widehat{\eta}_X(\tau)(x) = \bigvee_{t \in \hat{X}} t(x) \otimes \tau(t)$$

for all $\tau \in \hat{\hat{X}}$, $x \in X$. Hence, for $B \in H H X$ we have

$$\begin{aligned}
(m_X \cdot \mathcal{I}_X \cdot H\mathcal{I}_X)(\mathcal{B})(x) &= \bigvee_{t \in \hat{X}} t(x) \otimes \mathcal{I}_X(\mathcal{I}_X(\mathcal{B})) \\
&= \bigvee_{t \in \hat{X}} t(x) \otimes 1_X^*(t, \mathcal{I}_X(\mathcal{B})) \\
&= \bigvee_{t \in \hat{X}} \bigvee_{B \in \mathcal{B}} t(x) \otimes \left(\bigwedge_{x' \in X} t(x') \multimap a(x', B) \right) \\
&\leq \bigvee_{B \in \mathcal{B}} \bigvee_{t \in \hat{X}} t(x) \otimes (t(x) \multimap a(x, B)) \\
&= \bigvee_{B \in \mathcal{B}} a(x, B) \\
&= \mathcal{I}_X\left(\bigcup \mathcal{B}\right)(x) \\
&\leq a(x, x) \otimes \bigvee_{B \in \mathcal{B}} 1_X^*(y_X(x), a(-, B)) \\
&\leq \bigvee_{t \in \hat{X}} t(x) \otimes 1_X^*(t, \mathcal{I}_X(\mathcal{B})) \\
&= (m_X \cdot \mathcal{I}_X \cdot H\mathcal{I}_X)(\mathcal{B})(x).
\end{aligned}$$

Consequently, there is an induced algebraic functor

$$(\mathcal{V}\text{-Cat})^{\mathbb{H}} \leftarrow (\mathcal{V}\text{-Cat})^{\mathbb{P}\mathcal{V}}$$

of the respective Eilenberg–Moore categories.

We briefly describe the Eilenberg–Moore category

$$(\mathcal{V}\text{-Cat})^{\mathbb{H}}$$

where objects $X \in \mathcal{V}\text{-Cat}$ come equipped with a \mathcal{V} -functor $\alpha : HX \rightarrow X$ satisfying $\alpha \cdot \{-\} = 1_X$ and $1_{HX} \leq \{-\} \cdot \alpha$ (since \mathbb{H} is Kock–Zöberlein). Hence, $\alpha(\{x\}) = x$ for all $x \in X$, and $A \leq \{\alpha(A)\}$ for $A \in HX$, that is:

$$k \leq h_X(A, \{\alpha(A)\}) = \bigwedge_{x \in A} a(x, \alpha(A)).$$

Consequently, $\alpha(A)$ is an upper bound of A in the induced order of X , and for any other upper bound y of A in $X = (X, a)$ one has

$$k \leq \bigwedge_{x \in A} a(x, y) = h_X(A, \{y\}) \leq a(\alpha(A), \alpha(\{y\})) = a(\alpha(A), y)$$

since α is a \mathcal{V} -functor. Hence, $\alpha(A)$ gives (a choice of) a supremum of A in X . Moreover, the last computation shows

$$a\left(\bigvee A, y\right) = \bigwedge_{x \in A} a(x, y) \tag{*}$$

for all $y \in X$, $A \in HX$ (since “ \leq ” holds trivially). Conversely, any \mathcal{V} -category $X = (X, a)$ which is complete in its induced order and satisfies (*) is easily seen to be an object of $(\mathcal{V}\text{-Cat})^{\mathbb{H}}$.

Corollary 1. *The Eilenberg–Moore category of \mathbb{H} has order-complete \mathcal{V} -categories $X = (X, a)$ satisfying (*) as its objects, and morphisms are \mathcal{V} -functors preserving (the chosen) suprema.*

5. The lax Hausdorff monad on $\mathcal{V}\text{-Mod}$

When applying Theorem 1 to the Hausdorff functor $H : \mathcal{V}\text{-Cat} \rightarrow \mathcal{V}\text{-Cat}$ of Theorem 2 we obtain a lax functor $\tilde{H} : \mathcal{V}\text{-Mod} \rightarrow \mathcal{V}\text{-Mod}$ whose value on a \mathcal{V} -module $\varphi : X \rightrightarrows Y$ may be easily computed:

Lemma 2.

$$\tilde{H}\varphi(A, B) = \bigwedge_{x \in A} \bigvee_{y \in B} \varphi(x, y)$$

for all subsets $A \subseteq X$, $B \subseteq Y$.

Proof.

$$\begin{aligned}
 \tilde{H}\varphi(A, B) &= \bigvee_{D \in H\tilde{X}} (Hy_X)_*(A, D) \otimes (Hy_\varphi)^*(D, B) \\
 &= \bigvee_{D \in H\tilde{X}} h_{\tilde{X}}(y_X(A), D) \otimes h_{\tilde{X}}(D, y_\varphi(B)) \\
 &\leq h_{\tilde{X}}(y_X(A), y_\varphi(B)) \\
 &= \bigwedge_{x \in A} \bigvee_{y \in B} 1_{\tilde{X}}^*(y_X(x), y_\varphi(y)) \\
 &= \bigwedge_{x \in A} \bigvee_{y \in B} \varphi(x, y) \quad (\text{Yoneda}) \\
 &= h_{\tilde{X}}(y_X(A), y_\varphi(B)) \\
 &\leq h_{\tilde{X}}(y_X(A), y_X(A)) \otimes h_{\tilde{X}}(y_X(A), y_\varphi(B)) \\
 &\leq \bigvee_{D \in H\tilde{X}} h_{\tilde{X}}(y_X(A), D) \otimes h_{\tilde{X}}(D, y_\varphi(B)) \\
 &= \tilde{H}\varphi(A, B). \quad \square
 \end{aligned}$$

We now prove that \tilde{H} carries a lax monad structure.

Theorem 3. \tilde{H} belongs to a lax monad $\tilde{\mathbb{H}} = (\tilde{H}, \delta, \nu)$ of $\mathcal{V}\text{-Mod}$ such that $\tilde{\mathbb{H}}$ of Theorem 2 is a lifting of $\tilde{\mathbb{H}}$ along $(-)_* : \mathcal{V}\text{-Cat} \rightarrow \mathcal{V}\text{-Mod}$.

Proof. Let us first note that H is a lifting of \tilde{H} along $(-)_*$, in the sense that

$$\begin{array}{ccc}
 \mathcal{V}\text{-Cat} & \xrightarrow{H} & \mathcal{V}\text{-Cat} \\
 (-)_* \downarrow & & \downarrow (-)_* \\
 \mathcal{V}\text{-Mod} & \xrightarrow{\tilde{H}} & \mathcal{V}\text{-Mod}
 \end{array}$$

commutes. Indeed, for $f : X \rightarrow Y = (Y, b)$ in $\mathcal{V}\text{-Cat}$ and $A \in HX$, $B \in HY$ one has

$$\tilde{H}(f_*)(A, B) = \bigwedge_{x \in A} \bigvee_{y \in B} b(f(x), y) = h_Y(f(A), B) = (Hf)_*(A, B).$$

The unit of $\tilde{\mathbb{H}}$, $\delta : 1 \rightarrow \tilde{H}$, is defined by

$$\delta_X = \{-\}_* : X \rightrightarrows HX, \quad \delta_X(x, B) = h_X(\{x\}, B) = a(x, B),$$

for $X = (X, a)$, $x \in X$, $B \in HX$ (see also Remarks 2(2)), and the multiplication $\nu : \tilde{H}\tilde{H} \rightarrow \tilde{H}$ can be given by

$$\nu_X = \bigcup_* : HHX \rightrightarrows HX, \quad \nu_X(\mathcal{A}, B) = h_X\left(\bigcup \mathcal{A}, B\right) = \bigwedge_{A \in \mathcal{A}} h_X(A, B),$$

for $\mathcal{A} \in HHX$, $B \in HX$. The monad conditions hold *strictly* for $\tilde{\mathbb{H}}$, because they hold strictly for \mathbb{H} . For example, $\nu \cdot \tilde{H}\delta = 1$ follows from

$$\nu_X \cdot \tilde{H}\delta_X = \bigcup_* \cdot \tilde{H}(\{-\}_*) = \bigcup_* \cdot (H\{-\})_* = \left(\bigcup \cdot H\{-\}\right)_* = 1_X^*.$$

Surprisingly though, also the naturality squares for both δ_X and ν_X commute *strictly*. Indeed, for $\varphi : X \rightrightarrows Y = (Y, b)$, $x \in X$, $B \in HY$ and $\mathcal{A} \in HHX$ one has

$$\begin{aligned}
 (\tilde{H}\varphi \cdot \delta_X)(x, B) &= \bigvee_{A \in HX} \delta_X(x, A) \otimes \tilde{H}\varphi(A, B) \\
 &= \bigvee_{A \in HX} h_X(\{x\}, A) \otimes \tilde{H}\varphi(A, B) \\
 &= \tilde{H}\varphi(\{x\}, B) \\
 &= \bigvee_{y \in B} \varphi(x, y)
 \end{aligned}$$

$$\begin{aligned}
&= \bigvee_{y \in B} \bigvee_{z \in Y} \varphi(x, z) \otimes b(z, y) \\
&= \bigvee_{z \in Y} \varphi(x, z) \otimes \left(\bigvee_{y \in B} b(z, y) \right) \\
&= \bigvee_{z \in Y} \varphi(x, z) \otimes \delta_Y(z, B) \\
&= (\delta_Y \cdot \varphi)(x, B), \\
(\tilde{H}\varphi \cdot \nu_X)(\mathcal{A}, B) &= \bigvee_{A \in HX} \nu_X(A, A) \otimes \tilde{H}\varphi(A, B) \\
&= \bigvee_{A \in HX} h_X\left(\bigcup \mathcal{A}, A\right) \otimes \tilde{H}\varphi(A, B) \\
&= \tilde{H}\varphi\left(\bigcup \mathcal{A}, B\right) \\
&\leq \left(\bigwedge_{A \in \mathcal{A}} \bigvee_{B' \in HB} \tilde{H}\varphi(A, B') \right) \otimes \bigwedge_{B' \in HB} h_Y(B', B) \quad (\text{since } k \leq h_Y(B', B) \text{ for } B' \in HB) \\
&\leq \bigvee_{B \in HHY} \left(\bigwedge_{A \in \mathcal{A}} \bigvee_{B' \in B} \tilde{H}\varphi(A, B') \right) \otimes \left(\bigwedge_{B' \in B} h_Y(B', B) \right) \\
&= (\nu_Y \cdot \tilde{H}\tilde{H}\varphi)(\mathcal{A}, B) \\
&= \bigvee_{B \in HHY} \tilde{H}\tilde{H}\varphi(\mathcal{A}, B) \otimes \nu_Y(B, B) \\
&\leq \bigvee_{B \in HHY} \bigwedge_{A \in \mathcal{A}} \tilde{H}\varphi\left(A, \bigcup B\right) \otimes h_Y\left(\bigcup B, B\right) \\
&\leq \bigwedge_{A \in \mathcal{A}} \tilde{H}\varphi(A, B) \\
&= (\tilde{H}\varphi \cdot \nu_X)(\mathcal{A}, B). \quad \square
\end{aligned}$$

Remarks 3. (1) We emphasize that, while \tilde{H} is only a lax functor, this is in fact the only defect that prevents \tilde{H} from being a monad in the strict sense.

(2) In addition to the commutativity of the diagram given in the Proof of Theorem 3, since H obviously preserves full fidelity of \mathcal{V} -functors, from Theorem 1 we obtain also the commutativity of

$$\begin{array}{ccc}
(\mathcal{V}\text{-Cat})^{\text{op}} & \xrightarrow{H^{\text{op}}} & (\mathcal{V}\text{-Cat})^{\text{op}} \\
(-)^* \downarrow & & \downarrow (-)^* \\
\mathcal{V}\text{-Mod} & \xrightarrow{\tilde{H}} & \mathcal{V}\text{-Mod}.
\end{array}$$

(3) If \mathcal{V} is *constructively completely distributive* (see [18,5]), then $\tilde{H}\varphi$ for $\varphi : X \multimap Y$ may be rewritten as

$$\tilde{H}\varphi(A, B) = \bigvee \{v \in \mathcal{V} \mid \forall x \in A, \exists y \in B: v \leq \varphi(x, y)\}.$$

In this form the lax functor \tilde{H} was first considered in [5]. In the presence of the Axiom of Choice, so that \mathcal{V} is completely distributive in the ordinary (non-constructive) sense, one can then Skolemize the last formula to become

$$\tilde{H}\varphi(A, B) = \bigvee_{f:A \rightarrow B} \bigwedge_{x \in A} \varphi(x, f(x));$$

here the supremum ranges over arbitrary set mappings $f : A \rightarrow B$. Hence, the $\bigwedge \bigvee$ -formula of Lemma 2 has been transcribed rather compactly in $\bigvee \bigwedge$ -form.

For the sake of completeness we determine the Eilenberg–Moore algebras of \tilde{H} , i.e., those \mathcal{V} -categories $X = (X, a)$ which come equipped with a \mathcal{V} -module $\alpha : HX \multimap X$ satisfying

$$\alpha \cdot \delta_X = 1_X^* (= a), \quad (\dagger)$$

$$\alpha \cdot \nu_X = \alpha \cdot \tilde{H}\alpha. \quad (\ddagger)$$

The left-hand sides of those equations are easily computed as

$$\begin{aligned}
 (\alpha \cdot \delta_X)(x, y) &= \bigvee_{B \in HX} \delta_X(x, B) \otimes \alpha(B, y) \\
 &= \bigvee_{B \in HX} h_X(\{x\}, B) \otimes \alpha(B, y) \\
 &= \alpha(\{x\}, y), \\
 (\alpha \cdot \nu_X)(\mathcal{A}, y) &= \bigvee_{B \in HX} \nu_X(\mathcal{A}, B) \otimes \alpha(B, y) \\
 &= \bigvee_{B \in HX} h_X\left(\bigcup \mathcal{A}, B\right) \otimes \alpha(B, y) \\
 &= \alpha\left(\bigcup \mathcal{A}, y\right),
 \end{aligned}$$

for all $x, y \in X$, $\mathcal{A} \in HX$. Furthermore, if $k \leq \alpha(\{x\}, x)$, for all $x \in X$, then

$$\begin{aligned}
 \alpha\left(\bigcup \mathcal{A}, y\right) &\leq \bigwedge_{A \in \mathcal{A}} \alpha(A, y) \\
 &= \tilde{H}\alpha(\mathcal{A}, \{y\}) \otimes k \\
 &\leq \tilde{H}\alpha(\mathcal{A}, \{y\}) \otimes \alpha(\{y\}, y) \\
 &\leq \bigvee_{B \in HX} \tilde{H}\alpha(\mathcal{A}, B) \otimes \alpha(B, y) \\
 &= \alpha \cdot \tilde{H}\alpha(\mathcal{A}, y).
 \end{aligned}$$

Consequently, (\dagger) and (\ddagger) imply $\alpha(\{x\}, y) = a(x, y)$ and then

$$\begin{aligned}
 \alpha(A, y) &= \alpha\left(\bigcup \{\{x\} \mid x \in A\}, y\right) \\
 &= \bigwedge_{x \in A} \alpha(\{x\}, y) \\
 &= h_X(A, \{y\}) = \{-\}^*(A, y)
 \end{aligned}$$

for all $A \in HX$, $y \in X$. Hence, necessarily $\alpha = \{-\}^*$; conversely, this choice for α satisfies (\dagger) and (\ddagger) .

Corollary 2. The category of strict \tilde{H} -algebras and lax homomorphisms is the category $\mathcal{V}\text{-Mod}$ itself.

Proof. A lax homomorphism is, by definition, a \mathcal{V} -module $\varphi : X \rightharpoonup Y$ with $\varphi \cdot \alpha \leq \beta \cdot \tilde{H}\varphi$ (where α, β denote the uniquely determined structures of X, Y , respectively). A straightforward calculation shows that every \mathcal{V} -module satisfies this inequality. \square

6. The Gromov structure for \mathcal{V} -categories

With \tilde{H} as in Section 5, one is tempted to define

$$GH(X, Y) := \bigvee_{\varphi : X \rightharpoonup Y} \tilde{H}\varphi(X, Y)$$

for all \mathcal{V} -categories X and Y . But, as already suggested by the easy example considered in the Introduction, this definition leads to a trivial distance. Indeed, setting $\varphi(x, y) = \top$ for all $x \in X$, $y \in Y$, gives rise to a \mathcal{V} -module $\varphi : X \rightharpoonup Y$ with

$$\begin{aligned}
 \tilde{H}\varphi(X, Y) &= \bigwedge_{x \in X} \bigvee_{y \in Y} \varphi(x, y) \\
 &= \begin{cases} \perp, & \text{if } X \neq \emptyset, Y = \emptyset; \\ \top, & \text{else.} \end{cases}
 \end{aligned}$$

Consequently, whenever $Y \neq \emptyset$ or $X = \emptyset$

$$\top \leq \tilde{H}\varphi(X, Y) \leq GH(X, Y).$$

Hence, it is essential for us to consider subfunctors of H , rather than just H . More precisely, we consider any *sublifting* $K : \mathcal{V}\text{-Cat} \rightarrow \mathcal{V}\text{-Cat}$ of the powerset functor, by which we mean an endofunctor K with $X \in KX \subseteq HX$ such that the inclusion functions

$$\iota_X : KX \rightarrow HX$$

form a lax natural transformation, e.g., they are \mathcal{V} -functors such that

$$f(A) = (Hf)(A) \leq (Kf)(A)$$

in HY , for all \mathcal{V} -functors $f : X \rightarrow Y$ and $A \in KX$. Our guiding example is $K = H^s$ (see Section 7), but one may also take $K = H_\downarrow$ as in Remarks 2(3). For any such sublifting K of P we put

$$GK(X, Y) := \bigvee_{\varphi : X \twoheadrightarrow Y} \tilde{K}\varphi(X, Y).$$

Lemma 3. *If $f : X' \rightarrow X$ and $g : Y' \rightarrow Y$ are isomorphisms, then $GK(X', Y') = GK(X, Y)$.*

Proof. One can quickly verify the commutativity of the diagrams

$$\begin{array}{ccc} X' & \xrightarrow{y_{X'}} & \widehat{X}' \\ f \downarrow & & \downarrow \widehat{f}^* \\ X & \xrightarrow{y_X} & \widehat{X} \end{array} \quad \begin{array}{ccc} \widehat{X}' & \xleftarrow{y_{g^*} \cdot \varphi \cdot f_*} & Y' \\ \widehat{f}_* \uparrow & & \downarrow g \\ \widehat{X} & \xleftarrow{y_\varphi} & Y \end{array}$$

hence

$$\begin{aligned} \tilde{K}(g^* \cdot \varphi \cdot f_*) &= (Ky_{g^* \cdot \varphi \cdot f_*})^* \cdot (Ky_{X'})_* \\ &= (Kg)^* \cdot (Ky_\varphi)^* \cdot (K\widehat{f}_*)^* \cdot (Ky_{X'})_*, \end{aligned}$$

while

$$\begin{aligned} (Kg)^* \cdot \tilde{K}\varphi \cdot (Kf)_* &= (Kg)^* \cdot (Ky_\varphi)^* \cdot (Ky_X)_* \cdot (Kf)_* \\ &= (Kg)^* \cdot (Ky_\varphi)^* \cdot (K\widehat{f}^*)_* \cdot (Ky_{X'})_*. \end{aligned}$$

When f is an isomorphism, one has $f_*^{-1} = f^*$. Consequently, in this case $(K\widehat{f}_*)^* = (K\widehat{f}^*)_*$, and then

$$\tilde{K}(g^* \cdot \varphi \cdot f_*) = (Kg)^* \cdot \tilde{K}\varphi \cdot (Kf)_*.$$

Thus,

$$\begin{aligned} GK(X', Y') &= \bigvee_{\varphi : X' \twoheadrightarrow Y'} \tilde{K}\varphi(X', Y') \\ &= \bigvee_{\varphi} \tilde{K}(g^* \cdot \varphi \cdot f_*)(X', Y') \\ &= \bigvee_{\varphi} \tilde{K}\varphi(Kf(X'), Kg(Y')) \\ &= \bigvee_{\varphi} \tilde{K}\varphi(f(X'), g(Y')) \\ &= GK(X, Y). \quad \square \end{aligned}$$

We may now prove:

Theorem 4. *GK makes $\mathcal{G} := \text{ob}(\mathcal{V}\text{-Cat})/\cong$ a (large) \mathcal{V} -category, for every sublifting $K : \mathcal{V}\text{-Cat} \rightarrow \mathcal{V}\text{-Cat}$ of the powerset functor.*

Proof. Clearly

$$k \leq 1_{KX}^*(X, X) \leq \tilde{K}1_X^*(X, X) \leq GK(X, X),$$

and

$$\begin{aligned}
GK(X, Y) \otimes GK(Y, Z) &= \bigvee_{\varphi: X \twoheadrightarrow Y, \psi: Y \twoheadrightarrow Z} \tilde{K}\varphi(X, Y) \otimes \tilde{K}\psi(Y, Z) \\
&\leq \bigvee_{\varphi, \psi} \bigvee_{B \in KY} \tilde{K}\varphi(X, B) \otimes \tilde{K}\psi(B, Z) \\
&= \bigvee_{\varphi, \psi} (\tilde{K}\psi \cdot \tilde{K}\varphi)(X, Z) \\
&\leq \bigvee_{\varphi, \psi} \tilde{K}(\psi \cdot \varphi)(X, Z) \\
&\leq \bigvee_{\chi: X \twoheadrightarrow Z} \tilde{K}\chi(X, Z) \\
&= GK(X, Z). \quad \square
\end{aligned}$$

The resulting \mathcal{V} -category

$$\mathcal{G}K := (\mathcal{G}, GK)$$

may, with slightly stronger assumptions on K , be characterized as a colimit. For that purpose we first prove:

Lemma 4. *If $K : \mathcal{V}\text{-Cat} \rightarrow \mathcal{V}\text{-Cat}$ is a 2-functor, then*

$$\tilde{K}(g^* \cdot \varphi \cdot f_*) = (Kg)^* \cdot \tilde{K}\varphi \cdot (Kf)_*$$

for all f, g, φ as above.

Proof. It suffices to prove $(K\hat{f}_*)^* = (K\hat{f}^*)_*$ for all \mathcal{V} -functors $f : X' \rightarrow X$. But since both K and the (contravariant) $(\hat{-})$ preserve the order of hom-sets, from $f_* \dashv f^*$ in $\mathcal{V}\text{-Mod}$ we obtain $K\hat{f}^* \dashv K\hat{f}_*$ in $\mathcal{V}\text{-Cat}$. Now, since for any pair of \mathcal{V} -functors one has

$$h \dashv g \iff g^* \dashv h^* \iff g^* = h_*,$$

the desired identity follows with $h = K\hat{f}^*$ and $g = K\hat{f}_*$. \square

Proposition 2. *For any sublifting K of the powerset functor preserving the order of hom-sets and full fidelity of \mathcal{V} -functors one has*

$$GK(X, Y) = \bigvee_{X \hookrightarrow Z \hookleftarrow Y} 1_{KZ}^*(X, Y) = \bigvee_{X \hookrightarrow (X \sqcup Y, c) \hookleftarrow Y} \tilde{K}c(X, Y)$$

for all \mathcal{V} -categories X and Y .

Here the first join ranges over all \mathcal{V} -categories Z into which X and Y may be fully embedded, and the second one ranges over all \mathcal{V} -category structures c on the disjoint union $X \sqcup Y$ such that X and Y become full \mathcal{V} -subcategories.

Proof. Denoting the two joins by v, w , respectively, we trivially have $w \leq v$, so that $v \leq GK(X, Y) \leq w$ remains to be shown. Considering any full embeddings

$$X \hookrightarrow^{j_X} Z \hookleftarrow^{j_Y} Y$$

and putting $\varphi := j_Y^* \cdot (j_X)_* = j_Y^* \cdot 1_Z^* \cdot (j_X)_*$, because of K 's 2-functoriality and preservation of full fidelity we obtain from Lemma 4 and Theorem 1

$$\tilde{K}\varphi = (Kj_Y)^* \cdot \tilde{K}1_Z^* \cdot (Kj_X)_* = j_{KY}^* \cdot 1_{KZ}^* \cdot (j_{KX})_*$$

and therefore

$$1_{KZ}^*(X, Y) = \tilde{K}\varphi(X, Y) \leq GK(X, Y).$$

Considering any $\varphi : X \twoheadrightarrow Y$, one may define a \mathcal{V} -category structure c on $X \sqcup Y$ by

$$c(z, w) := \begin{cases} 1_X^*(z, w) & \text{if } z, w \in X; \\ \varphi(z, w) & \text{if } z \in X, w \in Y; \\ \perp & \text{if } z \in Y, w \in X; \\ 1_Y^*(z, w) & \text{if } z, w \in Y. \end{cases}$$

Then, with $Z := (X \sqcup Y, c)$, we again have $\varphi = j_Y^* \cdot (j_X)_*$ and obtain

$$\tilde{K}\varphi(X, Y) = \tilde{K}c(X, Y) \leq w. \quad \square$$

Theorem 5. For K as in Proposition 2, $\mathcal{G}K$ is a colimit of the diagram

$$\mathcal{V}\text{-Cat}_{\text{emb}} \longrightarrow \mathcal{V}\text{-Cat} \xrightarrow{K} \mathcal{V}\text{-Cat} \hookrightarrow \mathcal{V}\text{-CAT}.$$

Here $\mathcal{V}\text{-Cat}_{\text{emb}}$ is the category of small \mathcal{V} -categories with full embeddings as morphisms, and $\mathcal{V}\text{-CAT}$ is the category of (possibly large) \mathcal{V} -categories.

Proof. The colimit injection $\kappa_X : KX \rightarrow \mathcal{G}K$ sends $A \subseteq X$ to (the isomorphism class of) A , considered as a \mathcal{V} -category in its own right. Since for $A, B \in KX$ one has full embeddings $A \hookrightarrow X, B \hookrightarrow X$, trivially

$$1_{KX}^* \leq GK(A, B).$$

Hence κ_X is a \mathcal{V} -functor, and $\kappa = (\kappa_X)_X$ forms a cocone. Any cocone given by \mathcal{V} -functors $\alpha_X : KX \rightarrow (\mathcal{J}, J)$ allows us to define a \mathcal{V} -functor $F : \mathcal{G}K \rightarrow \mathcal{J}$ by $FX = \alpha_X(X)$. Indeed, given \mathcal{V} -categories X, Y we may consider any Z into which X, Y may be fully embedded (for example, their coproduct in $\mathcal{V}\text{-Cat}$) and obtain

$$\begin{aligned} 1_{KZ}^*(X, Y) &\leq J(\alpha_Z(X), \alpha_Z(Y)) \\ &\leq J(\alpha_X(X), \alpha_Y(Y)) \\ &= J(FX, FY). \end{aligned}$$

Hence, F is indeed a \mathcal{V} -functor with $F\kappa_X = \alpha_X$ for all X , and it is obviously the only such \mathcal{V} -functor. \square

For the sake of completeness we remark that the assignment

$$K \mapsto \mathcal{G}K$$

is monotone (=functorial): if we order subliftings of the powerset functor by

$$K \leq L \iff \text{there is a nat. tr. } \alpha : K \rightarrow L \text{ given by inclusion functions,}$$

while \mathcal{V} -category structures on $\mathcal{G} = \text{ob}(\mathcal{V}\text{-Cat})/\cong$ carry the pointwise order (as \mathcal{V} -modules), then

$$\mathcal{G} : \text{Sub } H \rightarrow \mathcal{V}\text{-CAT}(\mathcal{G})$$

becomes monotone. Indeed, for every \mathcal{V} -module $\varphi : Z \Rightarrow Y$, naturality of α gives

$$\begin{aligned} \alpha_Y^* \cdot \tilde{L}\varphi \cdot (\alpha_X)_* &= \alpha_Y^* \cdot (Ly_\varphi)^* \cdot (Ly_X)_* \cdot (\alpha_X)_* \\ &= (Ky_\varphi)^* \cdot \alpha_X^* \cdot (\alpha_X)_* \cdot (Ky_X)_* \\ &\geq (Ky_\varphi)^* \cdot (Ky_X)_* = \tilde{K}\varphi. \end{aligned}$$

Consequently,

$$\begin{aligned} \tilde{K}\varphi(X, Y) &\leq (\alpha_Y^* \cdot \tilde{L}\varphi \cdot (\alpha_X)_*)(X, Y) \\ &= \tilde{L}\varphi(\alpha_X(X), \alpha_Y(Y)) \\ &= \tilde{L}\varphi(X, Y), \end{aligned}$$

which gives $GK(X, Y) \leq GL(X, Y)$, for all \mathcal{V} -categories X, Y .

7. Symmetrization

A \mathcal{V} -category X , or just its structure $a = 1_X^*$, is *symmetric* when $a = a^\circ$. This defines the full subcategory $\mathcal{V}\text{-Cat}_s$ of $\mathcal{V}\text{-Cat}$ which is coreflective: the coreflector sends an arbitrary X to $X^s = (X, a^s)$ with $a^s = a \times a^\circ$, that is: $a^s(x, y) = a(x, y) \wedge a(y, x)$ for all $x, y \in X$. By

$$H^s X = (HX)^s = (PX, h_X^s)$$

one can define a sublifting $H^s : \mathcal{V}\text{-Cat} \rightarrow \mathcal{V}\text{-Cat}$ of the powerset functor which (like H) preserves full fidelity, but which (unlike H) fails to be a 2-functor. However its restriction

$$H^s : \mathcal{V}\text{-Cat}_s \rightarrow \mathcal{V}\text{-Cat}_s$$

is a 2-functor.

Remarks 4. (1) $H^s X$ must not be confused with $H(X^s)$. For example, for $\mathcal{V} = 2$ and a set X provided with a separated (= antisymmetric) order, X^s carries the discrete order. Hence, while in $H^s X$ one has $(A \leq B \iff A \subseteq \downarrow B \text{ and } B \subseteq \downarrow A \iff \downarrow A = \downarrow B)$, in $H(X^s)$ one has $(A \leq B \iff A \subseteq B)$.

(2) Even after its restriction to $\mathcal{V}\text{-Cat}_s$ there is no easy way of evaluating $\widetilde{H}^s \varphi(A, B)$ for a \mathcal{V} -module $\varphi : X \rightrightarrows Y$ and $A \subseteq X, B \subseteq Y$, since the computation leading to the easy formula of Lemma 2 does not carry through when H is replaced by H^s .

(3) The following addendum to Proposition 2 suggests how to overcome the difficulty mentioned in (2) when trying to define a non-trivial symmetric Gromov structure: \mathcal{V} -category structures c on the disjoint union $X \sqcup Y$ such that the \mathcal{V} -categories X, Y become full \mathcal{V} -subcategories correspond bijectively to pairs of \mathcal{V} -modules $\varphi : X \rightrightarrows Y, \varphi' : Y \rightrightarrows X$ with

$$\varphi' \cdot \varphi \leq 1_X^*, \quad \varphi \cdot \varphi' \leq 1_Y^*;$$

we write

$$\varphi : X \rightleftarrows Y : \varphi'$$

for such a pair. Under the hypotheses of Proposition 2 we can now write

$$GK(X, Y) = \bigvee_{\varphi : X \rightleftarrows Y : \varphi'} \tilde{K}\varphi(X, Y).$$

Hence, for any sublifting K of P we put

$$G^s K(X, Y) := \bigvee_{\varphi : X \rightleftarrows Y : \varphi'} \tilde{K}\varphi(X, Y) \wedge \tilde{K}\varphi'(Y, X)$$

and obtain easily:

Corollary 3. For any sublifting K of the powerset functor,

$$\mathcal{G}^s K = (\mathcal{G}, G^s K)$$

is a large symmetric \mathcal{V} -category, and when K is a 2-functor preserving full fidelity of \mathcal{V} -functors, then

$$G^s K(X, Y) = \bigvee_{X \hookrightarrow Z \hookleftarrow Y} 1_{KZ}^*(X, Y) \wedge 1_{KZ}^*(Y, X) = \bigvee_{X \hookrightarrow (X \sqcup Y, c) \hookleftarrow Y} \tilde{K}c(X, Y) \wedge \tilde{K}c(Y, X)$$

for all \mathcal{V} -categories X, Y .

Proof. Revisiting the proof of Lemma 3, we just note that

$$\varphi : X \rightleftarrows Y : \varphi', \quad \psi : Y \rightleftarrows Z : \psi' \quad \text{implies} \quad \psi \cdot \varphi : X \rightleftarrows Z : \varphi' \cdot \psi'.$$

A slight adaption of the computation given in Lemma 3 now shows that $G^s K$ is indeed a \mathcal{V} -category structure on $\mathcal{G} = \text{ob}\mathcal{V}\text{-Cat}/\cong$. The given formulae follow as in the proof of Proposition 2. \square

Corollary 4. $G^s H(X, Y) = G(H^s)(X, Y)$, for all \mathcal{V} -categories X, Y .

Extending the notion of symmetry from \mathcal{V} -categories to \mathcal{V} -modules, we call a \mathcal{V} -module $\varphi : X \rightrightarrows Y$ *symmetric* if X, Y are symmetric with $\varphi^\circ \cdot \varphi \leq 1_X^*$ and $\varphi \cdot \varphi^\circ \leq 1_Y^*$; we write

$$\varphi : X \rightleftarrows Y : \varphi^\circ$$

in this situation and define

$$G_s K(X, Y) := \bigvee_{\varphi : X \rightleftarrows Y : \varphi^\circ} \tilde{K}\varphi(X, Y)$$

for every sublifting K of P . Since symmetric \mathcal{V} -modules compose, similarly to Corollary 3 one obtains:

Corollary 5. For any sublifting K of the powerset functor,

$$\mathcal{G}_s K := (\text{ob}\mathcal{V}\text{-Cat}_s/\cong, G_s K)$$

is a large \mathcal{V} -category, and when K is a 2-functor preserving full fidelity of \mathcal{V} -functors, then

$$G_s K(X, Y) = \bigvee_{\substack{X \hookrightarrow Z \hookleftarrow Y \\ Z \text{ symmetric}}} 1_{KZ}^*(X, Y) = \bigvee_{\substack{X \hookrightarrow (X \sqcup Y, c) \hookleftarrow Y \\ c \text{ symmetric}}} \tilde{K}c(X, Y)$$

for all symmetric \mathcal{V} -categories X, Y .

Remarks 5. (1) It is important to note that $G_s K$ is *not* symmetric, even when $K = H$. For $\mathcal{V} = \mathbb{P}_+$, X a singleton and Y 3 equidistant points, we already saw in the Introduction that $G_s H(X, Y) = 0$ while $G_s H(Y, X) = \frac{1}{2}$. Hence it is natural to consider the symmetrization $(G_s K)^s$ of $G_s K$:

$$(G_s K)^s(X, Y) = G_s K(X, Y) \wedge G_s K(Y, X).$$

The same example spaces of the Introduction show that, while $(GH)^s(X, Y) = \max\{GH(X, Y), GH(Y, X)\} = 0$, one has

$$(G_s H)^s(X, Y) = \max\{G_s H(X, Y), G_s H(Y, X)\} = \frac{1}{2}.$$

(2) When the symmetric \mathcal{V} -categories X, Y are fully embedded into some \mathcal{V} -category Z , they are also fully embedded into Z^s . This fact gives

$$G^s H(X, Y) \leq G_s H(X, Y)$$

which, by symmetry of $G^s H$, gives

$$G^s H(X, Y) \leq (G_s H)^s(X, Y).$$

(3) Instead of the coreflector $X \mapsto X^s$ one may consider the *monoidal symmetrization* $X^{\text{sym}} = (X, a^{\text{sym}})$ with $a^{\text{sym}} = a \otimes a^\circ$, that is: $a^{\text{sym}}(x, y) = a(x, y) \otimes a(y, x)$. Hence, replacing \wedge by \otimes one can define $H^{\text{sym}}X$ and $G^{\text{sym}}K$ in complete analogy to $H^s X$ and $G^s X$, respectively. Corollary 3 remains valid when s is traded for sym and \wedge for \otimes .

8. Operations on the Gromov–Hausdorff \mathcal{V} -category

In this section we shall assume that our quantale \mathcal{V} satisfies the frame law. That is, for all $v, v_i \in \mathcal{V}$,

$$v \wedge \bigvee_i v_i = \bigvee_i v \wedge v_i.$$

Proposition 3. With the binary operation $(X, Y) \mapsto X \otimes Y$ the \mathcal{V} -category GH^s becomes a monoid in the monoidal category $\mathcal{V}\text{-CAT}$.

Proof. All we need to show is that

$$\otimes : GH^s \otimes GH^s \rightarrow GH^s$$

is a \mathcal{V} -functor. But for any \mathcal{V} -modules $\varphi : X \rightleftarrows X'$, $\psi : Y \rightleftarrows Y'$ and all $x \in X, y \in Y$ one trivially has

$$\tilde{H}\varphi(X, X') \otimes \tilde{H}\psi(Y, Y') \leq \bigvee_{x' \in X', y' \in Y'} \varphi(x, x') \otimes \psi(y, y'),$$

hence

$$\tilde{H}\varphi(X, X') \otimes \tilde{H}\psi(Y, Y') \leq \tilde{H}(\varphi \otimes \psi)(X \otimes Y, X' \otimes Y'),$$

with the \mathcal{V} -module $\varphi \otimes \psi : X \otimes Y \rightleftarrows X' \otimes Y'$ given by

$$(\varphi \otimes \psi)((x, y), (x', y')) = \varphi(x, x') \otimes \psi(y, y').$$

Consequently,

$$\begin{aligned} GH^s \otimes GH^s((X, Y), (X', Y')) &= GH^s(X, X') \otimes GH^s(Y, Y') \\ &= G^s H(X, X') \otimes G^s H(Y, Y') \\ &= \left(\bigvee_{\varphi, \varphi'} \tilde{H}\varphi(X, X') \wedge \tilde{H}\varphi'(X', X) \right) \otimes \left(\bigvee_{\psi, \psi'} \tilde{H}\psi(Y, Y') \wedge \tilde{H}\psi'(Y', Y) \right) \\ &\leq \bigvee_{\varphi, \psi} \tilde{H}(\varphi \otimes \psi)(X \otimes Y, X' \otimes Y') \wedge \bigvee_{\varphi', \psi'} \tilde{H}(\varphi' \otimes \psi')(X' \otimes Y', X \otimes Y) \\ &\leq \bigvee_{\chi, \chi'} \tilde{H}\chi(X \otimes Y, X' \otimes Y') \wedge \tilde{H}\chi'(X' \otimes Y', X \otimes Y) \\ &= G^s H(X \otimes Y, X' \otimes Y') = GH^s(X \otimes Y, X' \otimes Y'). \quad \square \end{aligned}$$

We note that when the \otimes -neutral element k of \mathcal{V} is its top element \top , then $v \otimes w \leq v \wedge w$ for all $v, w \in \mathcal{V}$ (since $v \otimes w \leq v \otimes k = v$); conversely, this inequality implies $k = \top$ (since $\top = \top \otimes k \leq \top \wedge k = k$).

Proposition 4. *If $k = \top$ in \mathcal{V} , then \mathcal{GH}^s becomes a monoid in the monoidal category $\mathcal{V}\text{-CAT}$ with the binary operation given either by product or by coproduct.*

Proof. We need to show that

$$\times : \mathcal{GH}^s \otimes \mathcal{GH}^s \rightarrow \mathcal{GH}^s \quad \text{and} \quad + : \mathcal{GH}^s \otimes \mathcal{GH}^s \rightarrow \mathcal{GH}^s$$

are both \mathcal{V} -functors. Similarly to the proof of Proposition 3, for the \mathcal{V} -functoriality of \times it suffices to show

$$\tilde{H}\varphi(X, X') \otimes \tilde{H}\psi(Y, Y') \leq \tilde{H}(\varphi \times \psi)(X \times Y, X' \times Y') \quad (\S)$$

for all \mathcal{V} -modules $\varphi : X \rightarrowtail X'$, $\psi : Y \rightarrowtail Y'$, where $\varphi \times \psi : X \times Y \rightarrow X' \times Y'$ is defined by

$$(\varphi \times \psi)((x, y), (x', y')) = \varphi(x, x') \wedge \psi(y, y').$$

(Note that, in this notation, $1_X^* \times 1_Y^*$ is the \mathcal{V} -category structure of the product $X \times Y$ in $\mathcal{V}\text{-Cat}$. The verification that $\varphi \times \psi$ is indeed a \mathcal{V} -module is easy.) But (§) follows just like in Proposition 3 since $k = \top$.

For the \mathcal{V} -functoriality of $+$ it suffices to establish the inequality

$$\tilde{H}\varphi(X, X') \otimes \tilde{H}\psi(Y, Y') \leq \tilde{H}(\varphi + \psi)(X + Y, X' + Y'), \quad (\P)$$

with $\varphi + \psi : X + Y \rightarrowtail X' + Y'$ defined by

$$(\varphi + \psi)(z, z') = \begin{cases} \varphi(z, z') & \text{if } z \in X, z' \in X', \\ \psi(z, z') & \text{if } z \in Y, z' \in Y', \\ \perp & \text{else.} \end{cases}$$

(Again, $1_X^* + 1_Y^*$ is precisely the \mathcal{V} -category structure of the coproduct $X + Y$ in $\mathcal{V}\text{-Cat}$, and the verification of the \mathcal{V} -module property of $\varphi + \psi$ is easy.) To verify (§) we consider $z \in X + Y$; then, for $z \in X$, say, we have

$$\begin{aligned} \tilde{H}\varphi(X, X') \otimes \tilde{H}\psi(Y, Y') &\leq \tilde{H}\varphi(X, X') \wedge \tilde{H}\psi(Y, Y') \\ &\leq \tilde{H}\varphi(X, X') \\ &\leq \bigvee_{x' \in X'} \varphi(z, x') \\ &\leq \bigvee_{z' \in X' + Y'} (\varphi + \psi)(z, z'), \end{aligned}$$

and (§) follows. \square

The previous proof shows that, without the assumption $k = \top$, one has that $+: \mathcal{GH}^s \times \mathcal{GH}^s \rightarrow \mathcal{GH}^s$ is a \mathcal{V} -functor, e.g. that $(\mathcal{GH}^s, +)$ is a monoid in the Cartesian category $\mathcal{V}\text{-CAT}$, but here we will continue to consider the monoidal structure of $\mathcal{V}\text{-CAT}$.

Theorem 6. *If $k = \top$ in \mathcal{V} , then the Hausdorff functor $H : \mathcal{V}\text{-Cat} \rightarrow \mathcal{V}\text{-Cat}$ induces a homomorphism $H : (\mathcal{GH}^s, +) \rightarrow (\mathcal{GH}^s, \times)$ of monoids in the monoidal category $\mathcal{V}\text{-CAT}$.*

Proof. Let us first show that the object-part of the functor $H : \mathcal{V}\text{-Cat} \rightarrow \mathcal{V}\text{-Cat}$ defines indeed a \mathcal{V} -functor $H : \mathcal{GH}^s \rightarrow \mathcal{GH}^s$, so that $\mathcal{GH}^s(X, Y) \leq \mathcal{GH}^s(HX, HY)$ for all \mathcal{V} -categories X, Y . But for every \mathcal{V} -module $\varphi : X \rightarrowtail Y$ and all $A \subseteq X$ one has

$$\tilde{H}\varphi(X, Y) \leq \tilde{H}\varphi(A, Y) \leq \bigvee_{B \subseteq Y} \tilde{H}\varphi(A, B),$$

which implies

$$\tilde{H}\varphi(X, Y) \leq \tilde{H}(\tilde{H}\varphi)(HX, HY)$$

and then the desired inequality.

In order to identify H as a homomorphism, we first note that, as an empty meet, $h_\emptyset(\emptyset, \emptyset)$ is the top element in \mathcal{V} , so that $H\emptyset \cong 1$ is terminal in $\mathcal{V}\text{-Cat}$, e.g. neutral in (\mathcal{GH}, \times) . The bijective map

$$+ : HX \times HY \rightarrow H(X + Y)$$

needs to be identified as an isomorphism in $\mathcal{V}\text{-Cat}$, e.g. we must show

$$(h_X \times h_Y)((A, B), (A', B')) = h_{X+Y}(A + B, A' + B')$$

for all $A, A' \subseteq X$, $B, B' \subseteq Y$. With $a = 1_X^*$ and $b = 1_Y^*$, in the notation of the proof of Proposition 4 one has

$$\bigvee_{z' \in A' + B'} (a + b)(x, z') = \bigvee_{x' \in A'} a(x, x')$$

for all $x \in A$ (since $(a + b)(x, z') = \perp$ when $z' \in B'$). Consequently,

$$\begin{aligned} h_{X+Y}(A + B, A' + B') &= \left(\bigwedge_{x \in A} \bigvee_{z' \in A' + B'} (a + b)(x, z') \right) \wedge \left(\bigwedge_{y \in B} \bigvee_{z' \in A' + B'} (a + b)(y, z') \right) \\ &= \left(\bigwedge_{x \in A} \bigvee_{x' \in A'} a(x, x') \right) \wedge \left(\bigwedge_{y \in B} \bigvee_{y' \in B'} b(y, y') \right) \\ &= h_X(A, A') \wedge h_Y(B, B'), \end{aligned}$$

as desired. \square

Note that $H : \mathcal{G}H^s \rightarrow \mathcal{G}H^s$ is a \mathcal{V} -functor even if the frame law does not hold in \mathcal{V} .

Remarks 6. (1) We may forego the blanket assumption of this section that \mathcal{V} satisfies the frame law, by considering $\mathcal{G}H^{\text{sym}}$ instead of $\mathcal{G}H^s$ (see Remarks 4(3)). All the results of this section still hold in this case.

(2) The $(\mathcal{V}\text{-Cat})$ -isomorphism

$$HX \times HY \cong H(X + Y)$$

exhibited in the proof of Theorem 6 easily extends to the infinite case:

$$\prod_i HX_i \cong H\left(\sum_i X_i\right).$$

(3) Since there is no general concept of a (covariant!) functor transforming coproducts into products, a more enlightening explanation for the formula just encountered seems to be in order, as follows. Since $\mathcal{V}\text{-Cat}$ is an *extensive category* (see [4]), for every (small) family $(X_i)_{i \in I}$ of \mathcal{V} -categories the functor

$$\Sigma : \prod_i \mathcal{V}\text{-Cat}/X_i \rightarrow \mathcal{V}\text{-Cat}/\sum_i X_i$$

is an equivalence of categories. Now, the (isomorphism classes of a) comma category $\mathcal{V}\text{-Cat}/X$ can be made into a (large) \mathcal{V} -category when we define the \mathcal{V} -category structure c by

$$c(f, g) = \bigwedge_{x \in A} \bigvee_{y \in B} 1_X^*(f(x), g(y)) = h_X(f(A), g(B)),$$

for all $f : A \rightarrow X$, $g : B \rightarrow X$ in $\mathcal{V}\text{-Cat}$. In this way the equivalence Σ has become an isomorphism of \mathcal{V} -categories, and since HX is just a \mathcal{V} -subcategory of $\mathcal{V}\text{-Cat}/X$, the $(\mathcal{V}\text{-Cat})$ -isomorphism of (1) is simply a restriction of the isomorphism Σ :

$$\begin{array}{ccc} \prod_i \mathcal{V}\text{-Cat}/X_i & \xrightarrow{\Sigma} & \mathcal{V}\text{-Cat}/\sum_i X_i \\ \uparrow & & \uparrow \\ \prod_i HX_i & \xrightarrow{\sim} & H(\sum_i X_i). \end{array}$$

9. Separation

A \mathcal{V} -category X , or just its structure $a = 1_X^*$, is *separated* when $k \leq a(x, y) \wedge a(y, x)$ implies $x = y$ for all $x, y \in X$. It was shown in [11] (and it is easy to verify) that the separated \mathcal{V} -categories form an epireflective subcategory of $\mathcal{V}\text{-Cat}$: the image of X under the Yoneda functor $y_X : X \rightarrow \hat{X}$ serves as the reflector. Furthermore, there is a closure operator which describes separation of X equivalently by the closedness of the diagonal in $X \times X$. (This description is not needed in what follows, but it further confirms the naturality of the concept.)

In Remarks 2 we already presented a sublifting H_\Downarrow of the powerset functor, and it is easy to check that $\tilde{H}_\Downarrow \varphi(A, B)$ may be computed as $\tilde{H} \varphi(A, B)$ in Lemma 2, e.g. the two values coincide, because of the formula proved in Remarks 2(1). Furthermore, H_\Downarrow is like H a 2-functor which preserves full fidelity of \mathcal{V} -functors. Hence, Proposition 2 is applicable to H_\Downarrow and may in fact be sharpened to:

Corollary 6. For all separated \mathcal{V} -categories X, Y one has

$$GH(X, Y) = GH_{\downarrow}(X, Y) = \bigvee_{\substack{X \hookrightarrow Z \hookleftarrow Y \\ Z \text{ separated}}} h_Z(X, Y) = \bigvee_{\substack{X \hookrightarrow (X \sqcup Y, c) \hookleftarrow Y \\ c \text{ separated}}} \tilde{H}c(X, Y).$$

Proof. The structure c constructed from a \mathcal{V} -module φ as in the proof of Proposition 2 is separated. \square

Remarks 7. (1) From Corollary 3 one obtains

$$\begin{aligned} G^s H(X, Y) &= G^s H_{\downarrow}(X, Y) = \bigvee_{X \hookrightarrow Z \hookleftarrow Y} h_Z(X, Y) \wedge h_Z(Y, X) \\ &= \bigvee_{X \hookrightarrow (X \sqcup Y, c) \hookleftarrow Y} \tilde{H}c(X, Y) \wedge \tilde{H}c(Y, X). \end{aligned}$$

However, here it is *not* possible to restrict the last join to separated structures c : consider the trivial case when $\mathcal{V} = 2$ and X, Y are singleton sets.

(2) \mathcal{V} -category structures c on $X \sqcup Y$ that are both symmetric and separated correspond bijectively to symmetric modules $\varphi : X \multimap Y$ with $k \not\leq \varphi(x, y)$ for all $x \in X, y \in Y$, provided that X and Y are both symmetric and separated. For $\mathcal{V} = 2$, X, Y are necessarily discrete, and the only structure c is discrete as well.

(3) The structure GH on \mathcal{G} is *not* separated, even if we consider only isomorphism classes of separated \mathcal{V} -categories: for $\mathcal{V} = 2$, the order on \mathcal{G} given by GH is chaotic! Likewise when G is traded for G^s .

Acknowledgements

The second- and third-named authors acknowledge encouragement and fertile pointers given by Bill Lawvere over the years, especially after a talk of the third-named author at the Royal Flemish Academy of Sciences in October 2008. This talk also led to a most interesting exchange with Isar Stubbe who meanwhile has carried the theme of this paper into the more general context whereby the quantale \mathcal{V} is traded for a quantaloid \mathcal{Q} (see [17]), a clear indication that the categorical study of the Hausdorff and Gromov metric may still be in its infancy.

References

- [1] A. Akhmediani, Hausdorff and Gromov distances in quantale-enriched categories, Master's thesis, York University, 2008.
- [2] D. Burago, Y. Burago, S. Ivanov, A Course in Metric Geometry, Amer. Math. Soc., Providence, RI, 2001.
- [3] M.M. Bonsangue, F. van Breugel, J.J.M.M. Rutten, Generalized metric spaces: Completion, topology, and powerdomains via the Yoneda embedding, Theoret. Comput. Sci. 193 (1998) 1–51.
- [4] A. Carboni, S. Lack, R.F.C. Walters, Introduction to extensive and distributive categories, J. Pure Appl. Algebra 84 (1993) 145–158.
- [5] M.M. Clementino, D. Hofmann, On extensions of lax monads, Theory Appl. Categ. 13 (2004) 41–60.
- [6] M.M. Clementino, D. Hofmann, Lawvere completeness in topology, Appl. Categ. Structures 17 (2009) 175–210.
- [7] M.M. Clementino, D. Hofmann, W. Tholen, One setting for all: Metric, topology, uniformity, approach structure, Appl. Categ. Structures 12 (2004) 127–154.
- [8] M. Gromov, Groups of polynomial growth and expanding maps, Publ. Math. I.H.E.S. 53 (1981) 53–78.
- [9] M. Gromov, Metric Structures for Riemannian and Non-Riemannian Spaces, Birkhäuser, Boston, MA, 2007.
- [10] F. Hausdorff, Grundzüge der Mengenlehre, Teubner, Leipzig, 1914.
- [11] D. Hofmann, W. Tholen, Lawvere completion and separation via closure, Appl. Categ. Structures, doi:10.1007/s10485-008-9169-9, in press.
- [12] G.M. Kelly, Basic Concepts of Enriched Category Theory, London Math. Soc. Lecture Note Ser., vol. 64, Cambridge University Press, Cambridge, 1982.
- [13] F.W. Lawvere, Metric spaces, generalized logic, and closed categories, Rend. Sem. Mat. Fis. Milano 43 (1973) 135–166; reprints in Theory Appl. Categ. 1 (2002) 1–37.
- [14] F.W. Lawvere, Taking categories seriously, Rev. Colombiana Mat. 20 (1986) 147–178; reprints in Theory Appl. Categ. 8 (2005) 1–24.
- [15] D. Pompeiu, Sur la continuité des fonctions de variables complexes, Doctoral thesis, Paris, 1905.
- [16] V. Schmitt, Completions of non-symmetric metric spaces via enriched categories, Georgian Math. J. 16 (2009) 161–186.
- [17] I. Stubbe, “Hausdorff distance” via conical cocompletion, preliminary report, December 2008.
- [18] R.J. Wood, Ordered sets via adjunctions, in: Categorical Foundations, in: Encyclopedia Math. Appl., vol. 97, Cambridge Univ. Press, Cambridge, 2004, pp. 5–47.