

Dietary patterns and alcohol consumption during pregnancy: secondary analysis of Avon

Longitudinal Study of Parents and Children

Dr Victoria Coathup^{1,2}, Dr Kate Northstone^{3,4}, Dr Ron Gray², Dr Simon Wheeler⁵, Dr Lesley Smith¹

Author affiliations

¹Department of Health and Life Sciences, Oxford Brookes University, Oxford, UK

²National Perinatal Epidemiology Unit, Nuffield Department of Population Health, University of Oxford, Oxford, UK

³School of Social and Community Medicine, University of Bristol, Bristol, UK

⁴The National Institute for Health Research Collaboration for Leadership in Applied Health Research and Care West (NIHR CLAHRC West) at University Hospitals Bristol NHS Foundation Trust, UK

⁵School of Life & Medical Sciences, University of Hertfordshire, Hatfield, UK

Corresponding author

Dr Victoria Coathup

National Perinatal Epidemiology Unit (NPEU)

Nuffield Department of Population Health

University of Oxford, Old Road Campus

Headington, Oxford, OX3 7LF

Victoria.coathup@npeu.ox.ac.uk

01865 617904

Sources of support

The UK Medical Research Council and the Wellcome Trust (Grant ref: 102215/2/13/2) and the University of

Bristol provide core support for ALSPAC. KN is funded by the National Institute for Health Research

Collaboration for Leadership in Applied Health Research and Care (NIHR CLAHRC) West at University Hospitals

Bristol NHS Foundation Trust

Abstract

Background

Large general population surveys show that heavy regular and episodic alcohol consumption are associated with lower intakes of fruits and vegetables, and higher intakes of processed and fried meat. This is of particular concern regarding pregnant women, as both alcohol intake and inadequate maternal nutrition are independently associated with adverse fetal outcomes. The current study aimed to determine associations between maternal dietary patterns and alcohol consumption during pregnancy.

Methods

Secondary analysis of data from the Avon Longitudinal Study of Parents and Children (ALSPAC). Women provided details of alcohol consumption at 18 weeks' gestation and diet at 32 weeks' gestation (n=9,839). Dietary patterns were derived from the food frequency questionnaire data using principal components analysis. Associations between alcohol consumption and dietary patterns were determined using multiple linear regression, adjusted for various socio-demographic and lifestyle factors.

Results

After adjustment, drinking ≥ 1 unit/day during the first trimester; $\beta=0.23$ (95% CI: 0.08, 0.38); $p=0.002$ and binge drinking (≥ 4 units in one day) during the first half of pregnancy; $\beta=0.14$ (95% CI: 0.07, 0.21); $p<0.0001$ were associated with greater adherence to the 'Processed' dietary pattern (high intakes of processed meat and low intakes of fruit and vegetables). Light to moderate alcohol consumption (≤ 1 drink/day) during the first trimester was associated with greater adherence to the 'Health conscious' dietary pattern (high intakes of fruit, vegetables, wholegrains and fish); $\beta=0.09$ (95% CI: 0.04, 0.14); $p<0.0001$.

Conclusions

Two important components of health behaviour during pregnancy appear to be related; greater consumption of processed foods associated with heavier alcohol consumption, and healthier dietary choices associated with light to moderate alcohol intake. Potential synergistic effects of these behaviours may have implications for

57 maternal and fetal health and warrant further investigation. A more holistic approach to addressing health
58 behaviours in women of reproductive age is required.

59

60

61 Keywords: gestation; maternal health; heavy episodic drinking; nutrition; ALSPAC

62

63

64

Introduction

Alcohol consumption and unhealthy dietary choices, characterised by lower intakes of fresh fruit and vegetables and higher consumption of salt, saturated fat and free sugars, are both major health risk behaviours that contribute to the global burden of disease (Murray *et al.*, 2013). Evidence suggests that approximately two thirds of individuals in the UK engage with two or more health risk behaviours, such as poor diet, physical inactivity, alcohol consumption and smoking. However, they are commonly investigated in isolation. Based on 2008 data from the Health Survey for England, approximately 12% of women reported lack of adherence to the recommended level of fruit and vegetable consumption and excessive drinking (Buck and Frosini, 2012).

Previous observational studies involving the general adult population have investigated relationships between diet and alcohol consumption patterns. These suggest that both frequency and quantity of alcohol consumption is associated with particular dietary patterns. Three studies have shown that as mean daily alcohol consumption increased, intakes of fruits, vegetables and dairy products decreased, while red and processed meat and egg intakes increased (Ruf *et al.* 2005; Touvier *et al.* 2014). Another study, which compared dietary habits of current and never drinkers, found never drinkers to have a higher overall Healthy Eating Index (HEI) score (HEI scores represent an individual's adherence to various recommended dietary guidelines in the US), indicative of 'better quality' diet (Breslow *et al.*, 2010). Furthermore, as quantity of alcohol per occasion increased, overall diet quality decreased (Breslow, Guenther and Smothers, 2006). 'Binge' or heavy episodic drinking has also been associated with lower intakes of fruit and vegetables, higher intakes of red and processed meats, and an increased likelihood of skipping meals (Valencia-Martin, Galan and Rodriguez-Artalejo, 2011). Such diets are characterised by lower intakes and plasma concentrations of important micronutrients, in particular folate (Brevik *et al.* 2005).

Concomitant health risk behaviours such as consuming an unhealthy diet and exceeding recommended limits for alcohol consumption might have important implications for maternal and infant health if they persist into pregnancy. Both are independent risk factors for adverse infant and childhood outcomes, including low birth weight (LBW) (Patra *et al.* 2011) and poor cognitive function (Zuccolo *et al.* 2013).

Animal models of Fetal Alcohol Spectrum Disorder (FASD) have indicated that alcohol-induced harm to the fetus is exacerbated by inadequate intake of folate, choline, vitamins E and C and carotenoids (Cohen-kerem and Koren, 2003; Thomas *et al.*, 2010; Ballard, Sun and Ko, 2012; May *et al.*, 2014). Similar findings have been reported in recent studies with human subjects (Avalos *et al.*, 2011; Hutson *et al.*, 2012; Coles *et al.*, 2015). While these results indicate the effects of ethanol are exacerbated in the presence of poor maternal nutrition, they do not explore the patterns between these two health behaviours.

There is evidence to suggest that mothers of children with FASD have significantly lower intakes of key micronutrients compared with mothers of healthy children (May *et al.*, 2014, 2016). However, the dietary data collected as part of that study was collected seven years after birth, and since health-related behaviour may change once a woman becomes pregnant (Crozier *et al.*, 2009), it remains unclear whether the associations between dietary intake and alcohol consumption remain exist during pregnancy, a time of rapid growth and greater maternal demand for micronutrients. If this is indeed the case, then those mothers engaging in multiple health risk behaviours may be particularly at risk of adverse pregnancy outcomes.

To date, the majority of research conducted in this area has explored the mediating effect of single nutrients. Whilst this method can provide valuable insight into the relationships between diet and health, nutrients are consumed as part of a diet, and in various combinations that may be interactive. Dietary patterns provide a broader representation of dietary intake and help to overcome the intercorrelations between foods and nutrients (Hu, 2002). Furthermore, studies have derived dietary patterns that include alcohol as a component, however, less is known about how these two behavioural determinants are associated when alcohol is not included in the dietary pattern analysis and considered separately.

The aim of this study was to determine the association between frequency and quantity of alcohol consumption, binge consumption and dietary patterns during pregnancy using prospectively collected data from the Avon Longitudinal Study of Parents and Children (ALSPAC).

Materials and Methods

Study design and participants

We conducted a secondary analysis of data from the ALSPAC cohort, a population-based study of pregnant women from the West of England and their subsequent children. Participants were followed from eight weeks' gestation to the present day (Golding *et al.*, 2001). ALSPAC recruitment methods have previously been described in detail (Boyd *et al.* 2012). Briefly, women were invited to participate if they resided in a pre-defined area within the county of Avon and their estimated delivery date was between 1st April 1991 and 31st December 1992. Initially, 14,541 pregnant women were recruited into the study; a total of 647 women were excluded, due to unknown outcomes or non-live births, leaving 13,761 unique women enrolled; a total of 14,062 live births. Women were eligible for inclusion in the current analysis if they had a live, singleton birth, provided details of alcohol consumption at 18 weeks' gestation and dietary intake at 32 weeks' gestation (n=11,457).

Ethical approval for the study was obtained from the ALSPAC Ethics and Law Committee and the Local Research Ethics Committee. Please note that the study website contains details of all the data that is available through a fully searchable data dictionary <http://www.bris.ac.uk/alspac/researchers/data-access/data-dictionary/>.

Alcohol consumption

Women completed a questionnaire at 18 weeks' gestation gathering self-reported frequency and quantity of alcohol consumption during the first trimester and episodes of binge drinking during the past month. We explored frequency and quantity of alcohol consumption during the first trimester hereafter referred to as 'regular' consumption as it describes average consumption across a three-month period (approximately). Responses included 'Never', 'Less than 1 glass per week', 'More than 1 glass per week', '1-2 glasses everyday', '3-9 glasses per day', and '10+ glass per day'. Due to low numbers in the higher frequency categories, the three highest categories were grouped together as '1+ glasses per day'. One glass was defined as one pub measure of spirits, one half pint of lager or cider, or one small glass of wine, which equates to approximately one unit (10ml ethanol).

152

153 Binge drinking was also assessed at 18 weeks' gestation with participants reporting the number of days during
154 the past month (14-18 weeks' gestation) when they had drunk the equivalent of two 568ml pints of beer, four
155 125ml glasses of wine, or four 25ml pub measures of spirit, each equating to approximately four standard units
156 (40ml ethanol). The available responses were: 'None', '1-2 days', '3-4 days', '5-10 days', 'More than 10 days'
157 and 'Everyday'. Due to low numbers of women who reported more than 2 occasions of binge drinking during
158 mid-pregnancy, this variable was dichotomized into 'non-binge drinkers' and 'binge drinkers', defined as a
159 woman who reported drinking four or more units on at least one day during the past month. Although binge
160 drinking in the UK is generally defined as the consumption of six or more units during any one occasion (HSCIC,
161 2014), it was defined within ALSPAC as four or more units and is consistent with other secondary analyses
162 using data from the ALSPAC study (Alati et al. 2013). We have included this variable in the current analysis as it
163 represents a pattern of drinking that is associated with higher blood alcohol concentrations (BAC) – an
164 important factor in FASD - (Pierce and West, 1986) compared to average frequency and quantity of alcohol
165 consumed that is also reported in ALSPAC.

166

167 *Dietary assessment*

168

169 At 32 weeks' gestation women completed a food frequency questionnaire (FFQ), reporting how often they
170 were currently consuming 44 common food and drink items. The FFQ was adapted from a previous
171 questionnaire (Yarnell et al. 1983).

172

173 Details of how the FFQ data were prepared have been reported in detail elsewhere (Rogers et al. 1998).
174 Briefly, standard portion sizes were allocated to food and drink items in the FFQ using a UK reference guide
175 (Food Standards Agency, 1988) and weekly intake frequencies were recoded as 0 (Never/Rarely), 0.5 (Once in
176 2 weeks), 2 (1-3 per week), 5.5 (4-7 per week) and 10 (More than 1 per day). For non-alcoholic beverages and
177 bread women recorded the number of servings/slices consumed per day. Cooking methods and types of bread
178 consumed were also reported. Milk quantity was calculated by summing standard amounts from all tea,
179 coffee, cereal, puddings and drinks consumed weekly. Maternal dietary patterns did not include alcohol
180 consumption variables because the sample population comprised of pregnant women, who report lower

alcohol consumption levels compared with the general population; therefore, it is unlikely that heavy or binge drinking would have been a defining characteristic of a dietary pattern.

Potential confounding factors

A wide variety of socio-demographic and lifestyle data were collected at both 8 and 18 weeks' gestation and were explored as potential confounding variables in relation to alcohol consumption and dietary intake during pregnancy. These were: maternal age (<20, 20-24, 25-29 or ≥ 30 years); parity (none or ≥ 1); ethnicity (white, non-white); smoking (current smoker/non-smoker); highest level of maternal education (vocational, O-level, A-level, degree level); housing tenure (owner occupied, council/Housing Association (HA) rented, private rented/other); house crowding index (HCI), defined as the total number of people per household, divided by the total number of rooms (excluding the kitchen and bathrooms) (Melki *et al.*, 2004); living in a single parent household; and depression symptoms measured using the Edinburgh Postnatal Depression Scale (EPDS) score (Cox, Holden and Sagovsky, 1987). Women with EPDS scores of ≥ 13 are more likely to be suffering from depression than those with lower scores (Murray and Cox, 1990). **Table 1** presents details of how each variable is categorised.

Statistical analysis

Dietary patterns were replicated using the methods previously described by Northstone *et al* (2008). Briefly, PCA with a varimax rotation was performed on the 44 standardised food and drink items. Factor loadings represent the correlation between the original dietary variable and the factor (dietary pattern) and food items with factor loadings of ≥ 0.3 or ≤ -0.3 suggested a strong positive or negative association, respectively, and were considered to clearly contribute to that dietary pattern. The five component (dietary pattern) scores represent a participant's adherence to each dietary pattern. Scores have a mean of zero; a value above or below zero indicates stronger or weaker adherence to that dietary pattern, respectively.

Participants with missing data on >10 food and drink items were excluded from the analysis. Those with ≤ 10 missing items were included and the missing data recoded as 0 (Never/Rarely). Unadjusted logistic regression

models were used to assess differences (socio-demographic and lifestyle characteristics) between populations of women with and without dietary data (See supplementary data).

Participant characteristics and alcohol consumption variables are presented as frequencies and percentages. Dietary pattern component scores are presented as means and standard deviations. Unadjusted linear regression models were used to explore associations between alcohol consumption and dietary pattern scores. We fitted separate regression models for each dietary pattern to minimise the risk of multicollinearity. Linear regression models were then adjusted for maternal age, parity, ethnicity, smoking, education, HCI, housing tenure, living in a single parent household and EPDS score. Results are presented as effect sizes with 95% confidence intervals and p-values. All analyses were conducted using STATA 13.1.

Results

A total of 9,839 women were included in the current analysis. Women with missing dietary data were more likely to be younger, smoke and of lower socio-economic status (SES) (see supplementary data). The socio-demographic and lifestyle characteristics of women are presented in **Table 1**. Women of lower socio-economic status (SES) and those of non-white ethnic origin were underrepresented compared to women in the UK during the same time period (Fraser *et al.*, 2013); 13% lived in council or Housing Association rented accommodation; 5% scored >1 on the HCI and 2% were of non-white ethnicity. Approximately 2% of women reported drinking ≥1 unit per day during the first trimester and 7% of women reported binge drinking on at least one day during the previous month when assessed at 18 weeks' gestation.

A full description of the five dietary patterns are provided elsewhere (Northstone *et al.*, 2008). Briefly, the 'Health conscious' component was characterised by greater consumption of wholegrains, cereals, fruits, salad, fish, and lower intake of white bread. 'Traditional' was characterised by greater consumption of vegetables and potatoes. 'Processed' was characterised by greater consumption of white bread, fried foods, processed meats, and lower intakes of wholegrains. 'Confectionery' was characterised by greater consumption of chocolate, crisps, sweets and biscuits. 'Vegetarian' was characterised by low intakes of meat and high intakes of meat substitutes, nuts and pulses. The components accounted for a total of 31.3% of the variation.

'Regular' alcohol consumption

Light-to-moderate alcohol consumption (defined as <1 drink/day) during the first trimester was associated with higher 'Health conscious' dietary pattern scores compared with no drinking during the same period ($\beta=0.12$, 95%CI=0.06, 0.17; $p<0.0001$) (**Table 2**). It was also associated with higher 'Confectionery' scores ($\beta=0.10$, 95%CI=0.04, 0.16; $p<0.0001$). Heavy alcohol consumption (defined as 1+drinks/day) during the first trimester was associated with higher 'Processed' scores ($\beta=0.17$, 95%CI=0.03, 0.31; $p=0.015$) and higher 'Vegetarian' scores ($\beta=0.23$, 95%CI=0.09, 0.37; $p=0.001$), compared with women who reported never drinking alcohol during the first trimester.

After adjustment for confounding (**Table 2**), the association between light-to-moderate alcohol consumption with the 'Health conscious' dietary pattern remained significant ($\beta=0.08$, 95%CI=0.03, 0.13; $p=0.002$); the association with 'Confectionery' was strengthened ($\beta=0.11$, 95%CI=0.05, 0.17; $p=0.001$). Relationships between alcohol consumption during the first trimester and adherence to the 'Processed' dietary pattern remained ($\beta=0.24$, 95%CI=0.09, 0.39; $p=0.002$); as alcohol consumption during the first trimester increased 'Processed' dietary pattern scores increased (**Figure 1**).

Binge drinking

Before adjustment for confounding, at least one episode of binge drinking during 14-18 weeks of pregnancy was associated with higher scores of 'Processed' ($\beta=0.23$, 95%CI=0.17, 0.30; $p<0.0001$) and 'Vegetarian' ($\beta=0.09$, 95%CI=0.03, 0.16; $p=0.007$) dietary patterns, and lower scores of the 'Health conscious' ($\beta=-0.28$, 95%CI=-0.35, -0.21; $p<0.0001$) and 'Confectionery' ($\beta=-0.07$, 95%CI=-0.14, -0.01; $p=0.034$) dietary patterns (**Table 3**). Once adjusted for confounding, however, only the association with the 'Processed' dietary pattern remained ($\beta=0.15$, 95%CI=0.07, 0.22; $p<0.0001$) (**Figure 1**).

Discussion

The aim of this study was to determine the associations between maternal alcohol consumption and dietary patterns during pregnancy. Whilst a number of studies have explored aspects of maternal diet in relation to alcohol consumption, before, during and post pregnancy (Keen *et al.*, 2010; Weiss and Chambers, 2013; May *et al.*, 2014), this is the first study to explore maternal dietary pattern scores in relation to alcohol consumption during the antenatal period. The findings from this secondary analysis have highlighted the associations between two important health risk behaviours during pregnancy – alcohol consumption and dietary intake – and suggest that women who report drinking heavily during pregnancy may also have poorer quality diets, characterised by low intakes of fruit and vegetables and high intakes of processed and fried foods.

After adjusting for potential confounders, associations were evident between drinking one or more alcoholic drinks per day and adherence to the ‘Processed’ dietary pattern. Previous studies have explored associations between individual food groups or nutrients and alcohol consumption in the general population. A large cohort study conducted in France explored the dietary intake of approximately 73,000 adult women and found that fruit and vegetable intakes were lower in those who consumed approximately one or more alcoholic drinks per day (Kesse *et al.*, 2001). Another study conducted in France reported that women consuming alcohol were less likely to eat $\geq 400\text{g}$ of fruit and vegetables per week and more likely to eat $\geq 500\text{g}$ of red meat per week, compared with non-drinkers (Touvier *et al.* 2014).

Clear associations were also observed between binge drinking and adherence to the ‘Processed’ dietary pattern. A study conducted in the USA explored the drinking habits and Healthy Eating Index (HEI) scores of 772 women. The mean HEI score decreased by 5.6 points in women who consumed 3 or more drinks per occasion, compared with women who reported one per occasion, after adjusting for socio-demographic characteristics (Breslow *et al.* 2006). A study conducted in Madrid, Spain, explored binge drinking in a randomly selected sample of approximately 12,000 adults of the general population and found that people who reported heavy episodic drinking were more likely to consume fewer than three portions of fruit and vegetables per day, and more than one serving of meat per day, compared to never drinkers, after adjusting for all socio-demographic characteristics (Valencia-Martin, Galan and Rodriguez-Artalejo, 2011).

Light to moderate alcohol consumption was associated with adherence to the 'Confectionery' and 'Health conscious' dietary patterns. The 'Confectionery' dietary pattern was characterised by high intakes of sweets, cake and biscuits. While two studies have reported a decrease in sweet and sugary foods as alcohol consumption increased (Smith and Smith, 1994; Herbeth *et al.*, 2012), few studies in the general population explored the intakes of confectionery in relation to alcohol consumption. The 'Health conscious' dietary pattern was characterised by high intakes of fruit, salad, wholegrains and fish, and similar patterns during pregnancy have been described in other studies (Crozier *et al.* 2006; Knudsen *et al.* 2007). Evidence from studies in the general population have also reported that patterns of light to moderate alcohol consumption are associated with higher intakes of fruit, vegetables and fish, compared to abstaining or drinking more heavily (Kesse *et al.*, 2001; Barefoot *et al.*, 2002; Valencia-Martin, Galan and Rodriguez-Artalejo, 2011). Breslow *et al.* (2006) found that light and frequent patterns of alcohol consumption were associated with the highest HEI scores compared to abstainers and heavier drinkers.

This study has indicated that women who continue to drink in potentially harmful patterns (binge and 1+/day) are also more likely to have poorer quality diets, characterised by higher intakes of red meat, processed foods and lower intakes of fresh fruits and vegetables compared to women who do not drink. These relationships are particularly important during pregnancy; a time of rapid growth, with greater nutrient demands for the fetus. A study exploring relationships between mean daily micronutrient intakes and the same dietary patterns within the ALSPAC cohort reported that as adherence to the 'Processed' and 'Confectionery' dietary patterns increased, micronutrient intakes decreased, including folate, vitamin B6, vitamin C, vitamin E and carotene (Northstone *et al.* 2008). This may have implications for fetal development, as inadequate intakes of these micronutrients during pregnancy are associated with an increased risk of ethanol-induced fetal harm in experimental models (Gutierrez *et al.* 2007; Naseer *et al.* 2011) and fetal growth restriction and poor cognitive outcomes at 6 months in studies within human populations (Avalos *et al.*, 2011; Coles *et al.*, 2015). Two suggested mechanisms for these relationships are the interference of ethanol in one carbon metabolism (OCM) and the redox state of cells (Cohen-kerem and Koren, 2003; Ballard, Sun and Ko, 2012). If women are drinking heavily and also adhere to dietary patterns characterised by low intakes of fresh fruit and vegetables, this may increase the risk of adverse birth and childhood outcomes.

325 Whilst this is one of the first studies to explore maternal dietary pattern scores in relation to alcohol
326 consumption during pregnancy, other research teams have explored relationships between maternal alcohol
327 consumption and other aspects of diet. Preliminary findings from a trial conducted in the Ukraine found that
328 women who reported alcohol consumption during pregnancy were more likely to have lower plasma zinc and
329 copper concentrations compared to women in the control group (Keen *et al.*, 2010). Whilst this is an
330 interesting finding, the sample was small (n=49) and plasma concentrations do not accurately reflect dietary
331 intake, due to potential biological interactions influencing bioavailability (Moran *et al.*, 2012). Multivitamin
332 supplements were also explored in relation to alcohol consumption during the periconceptional period in a
333 large cross-sectional study conducted in the US and found that as alcohol consumption increased, women
334 were less likely to take multivitamin supplements (Weiss and Chambers, 2013). This finding is particularly
335 interesting in light of the current analysis; if women who binge drink are less likely to consume diets
336 characterised by high intakes of fresh fruit and vegetables, are they also less likely to take folic acid and
337 multivitamin supplements during pregnancy?
338

339 The clustering of risky health behaviours increases the risk of adverse fetal development (Lanting *et al.*, 2009),
340 and may also provide an explanation for why some studies have published findings that suggest women who
341 drink low to moderate amounts during pregnancy have children with better cognitive outcomes: diet may be
342 an overlooked confounder. Negative health behaviours often cluster in populations (French *et al.* 2008;
343 Shankar *et al.* 2010) and a previous study explored relationships between the five dietary patterns described in
344 the ALSPAC cohort study and socio-demographic and lifestyle characteristics. The 'Health conscious' dietary
345 pattern was indicative of higher social affluence; adherence was associated with older age, higher educational
346 attainment, living in an owned or mortgaged property, lower parity, being white, not smoking and having
347 fewer financial difficulties. In contrast, adherence to the 'Processed' dietary pattern was associated with the
348 opposite trends, indicating lower social affluence (Northstone *et al.* 2008). Evidence also suggests that
349 patterns of alcohol consumption during pregnancy are related to socio-demographic characteristics. A
350 systematic review reported that five studies found higher income or social class to be associated with alcohol
351 consumption during pregnancy, but not with binge drinking (Skagerström, Chang and Nilsen, 2011). A study
352 conducted in Sweden found similar results in a non-pregnant population; binge drinking was associated with
353 lower social affluence (Backhans, Lundin and Hemmingsson, 2012).

354

355 Evidence from studies assessing health behaviour change interventions have indicated that when two or more
356 health risk behaviours, such as diet, smoking, alcohol consumption or exercise, are approached in
357 combination, individuals tend to have better outcomes (Jepson, 2000). Since dietary and alcohol intake appear
358 to be related, a more effective way of addressing these health behaviours during pregnancy might be to take a
359 more holistic approach and consider them together (Prochaska and Prochaska, 2011).

360

361 *Strengths and limitations*

362

363 The main strengths of this analysis were the large sample size and the number of respondents reporting
364 alcohol consumption at different frequencies and quantities. Because ALSPAC had collected data on episodic
365 drinking, it was possible to assess the relationships with irregular patterns of alcohol consumption, which may
366 often go undetected when asking questions about average consumption. However, there are also a number of
367 limitations to this study that must be acknowledged. The estimates of alcohol consumption and dietary intake
368 are self-reported, and therefore, vulnerable to recall and social-desirability biases (Davis et al. 2010; Thompson
369 & Subar 2008). In addition to this, the FFQ did not capture portion size data and the validity and reproducibility
370 of the FFQ used to estimate dietary intake are uncertain. However, the values estimated in ALSPAC compared
371 favourably with estimates reported by women in the Dietary and Nutritional Survey (Rogers et al. 1998).
372 Furthermore, the drinking categories were also unbalanced, with a very small proportion of women reporting
373 to drink one or more drinks per day during the first trimester. While this is fairly typical of pregnant
374 populations in the UK it may increase the risk of erroneous results, particularly in multivariate regression
375 models (Button *et al.*, 2013). Moreover, despite the large sample, 98% of the sample population was white,
376 and only 13% lived in property rented by the housing association or council. Low recruitment and retention
377 rates of women from low socio-economic backgrounds are well documented in public health research, and
378 further work should be conducted to evaluate these relationships within populations of women of non-white
379 ethnic origin and lower SES. Whilst a large number of potential confounders have been adjusted for in the
380 analyses, it is possible that these relationships are due to residual confounding from unmeasured SES and
381 other lifestyle factors. Finally, the data included in this analysis was originally collected by ALSPAC in the early
382 1990s and we acknowledge that this threatens external validity due to changes in alcohol (Department of

Health, 2016) and dietary guidelines (SACN, 2011) since that period. Therefore, additional research must be conducted to explore whether the relationships observed in this sample population are present in populations of pregnant women today.

Conclusions

Overall, this study has indicated that the relationships between diet and alcohol that have been previously reported in the general population persist into pregnancy. The findings also suggest the need to address health risk behaviours together, rather in isolation. Alcohol behaviour change interventions during pregnancy may be more successful if tackled as a broader goal, along with diet.

Acknowledgements

We are extremely grateful to all the families who took part in this study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists and nurses. The UK Medical Research Council and the Wellcome Trust (Grant ref: 102215/2/13/2) and the University of Bristol provide core support for ALSPAC. This publication is the work of the authors and KN will serve as guarantor for the contents of this paper. KN is funded by the National Institute for Health Research Collaboration for Leadership in Applied Health Research and Care (NIHR CLAHRC) West at University Hospitals Bristol NHS Foundation Trust. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

Conflicts of interest

The authors declare they have no competing interests.

References

- Alati, R., Davey Smith, G., Lewis, S. J., Sayal, K., Draper, E. S., Golding, J., Fraser, R. and Gray, R. (2013) 'Effect of prenatal alcohol exposure on childhood academic outcomes: contrasting maternal and paternal associations in the ALSPAC study.', *PloS one*, 8(10), p. e74844. doi: 10.1371/journal.pone.0074844.
- Avalos, L. A., Kaskutas, L., Block, G., Abrams, B. and Li, D. K. (2011) 'Does lack of multinutrient supplementation during early pregnancy increase vulnerability to alcohol-related preterm or small-for-gestational-age births?', *Matern Child Health J.* 2010/10/16, 15(8), pp. 1324–1332. doi: 10.1007/s10995-010-0690-8.
- Backhans, M. C., Lundin, A. and Hemmingsson, T. (2012) 'Binge Drinking—A Predictor for or a Consequence of Unemployment?', *Alcoholism: Clinical and Experimental Research*, 36(11), pp. 1983–1990. doi: 10.1111/j.1530-0277.2012.01822.x.
- Ballard, M. S., Sun, M. and Ko, J. (2012) 'Vitamin A, folate, and choline as a possible preventive intervention to fetal alcohol syndrome', *Med Hypotheses*. 2012/01/31, 78(4), pp. 489–493. doi: 10.1016/j.mehy.2012.01.014.
- Barefoot, J. C., Grønbaek, M., Feaganes, J. R., McPherson, R. S., Williams, R. B. and Siegler, I. C. (2002) 'Alcoholic beverage preference, diet, and health habits in the UNC Alumni Heart Study.', *The American journal of clinical nutrition*, 76(2), pp. 466–72.
- Breslow, R. A., Guenther, P. M., Juan, W. and Graubard, B. I. (2010) 'Alcoholic beverage consumption, nutrient intakes, and diet quality in the US adult population, 1999–2006.', *Journal of the American Dietetic Association*, 110(4), pp. 551–62. doi: 10.1016/j.jada.2009.12.026.
- Breslow, R. a, Guenther, P. M. and Smothers, B. a (2006) 'Alcohol drinking patterns and diet quality: the 1999–2000 National Health and Nutrition Examination Survey.', *American journal of epidemiology*, 163(4), pp. 359–66. doi: 10.1093/aje/kwj050.
- Brevik, A., Vollset, S. E., Tell, G. S., Refsum, H., Ueland, P. M. and Loeken, E. B. (2005) 'Plasma concentration of folate as a biomarker for the intake of fruit and vegetables : the Hordaland Homocysteine Study 1 – 3', *American Journal of Clinical Nutrition*, 81(2), pp. 434–439.
- Buck, D. and Frosini, F. (2012) 'Clustering of unhealthy behaviours over time - Implications for policy and practice', *Implications for policy and practice. The Kings Fund: ...*, pp. 1–24.
- Button, K. S., Ioannidis, J. P. a, Mokrysz, C., Nosek, B. a, Flint, J., Robinson, E. S. J. and Munafò, M. R. (2013) 'Power failure: why small sample size undermines the reliability of neuroscience.', *Nature reviews*.

440 *Neuroscience*, 14(5), pp. 365–76. doi: 10.1038/nrn3475.

441 Chiolerio, A., Wietlisbach, V., Ruffieux, C., Paccaud, F. and Cornuz, J. (2006) 'Clustering of risk behaviors with
 442 cigarette consumption: A population-based survey.', *Preventive medicine*, 42(5), pp. 348–53. doi:
 443 10.1016/j.ypmed.2006.01.011.

444 Cohen-kerem, R. and Koren, G. (2003) 'Antioxidants and fetal protection against ethanol teratogenicity I .
 445 Review of the experimental data and implications to humans', 25, pp. 1–9. doi: 10.1016/S0892-
 446 0362(02)00324-0.

447 Coles, C. D., Kable, J. a., Keen, C. L., Jones, K. L., Wertelecki, W., Granovska, I. V., Pashtepa, A. O. and Chambers,
 448 C. D. (2015) 'Dose and Timing of Prenatal Alcohol Exposure and Maternal Nutritional Supplements:
 449 Developmental Effects on 6-Month-Old Infants', *Maternal and Child Health Journal*. Springer US, 19(12), pp. 1–
 450 10. doi: 10.1007/s10995-015-1779-x.

451 Cox, J. L., Holden, J. M. and Sagovsky, R. (1987) 'Detection of postnatal depression. Development of the 10-
 452 item Edinburgh Postnatal Depression Scale.', *The British journal of psychiatry : the journal of mental science*,
 453 150, pp. 782–6.

454 Crozier, S. R., Robinson, S. M., Borland, S. E., Godfrey, K. M. and Cooper, C. (2009) 'Do women change their
 455 health behaviours in pregnancy ? Findings from the Southampton Women ' s Survey', *Europe PMC Funders*
 456 *Group*, 23(5), pp. 446–453. doi: 10.1111/j.1365-3016.2009.01036.x.Do.

457 Crozier, S. R., Robinson, S. M., Borland, S. E. and Inskip, H. M. (2006) 'Dietary patterns in the Southampton
 458 Women's Survey', *Eur J Clin Nutr*. 2006/06/29, 60(12), pp. 1391–1399. doi: 10.1038/sj.ejcn.1602469.

459 Czeizel, E., Petik, D. and Puho, E. (2004) 'Smoking and alcohol drinking during pregnancy. The reliability of
 460 retrospective maternal self-reported information.', *Central European journal of public health*, 12(4), pp. 179–
 461 83.

462 Davis, C. G., Thake, J. and Vilhena, N. (2010) 'Social desirability biases in self-reported alcohol consumption and
 463 harms.', *Addictive behaviors*. England, 35(4), pp. 302–311. doi: 10.1016/j.addbeh.2009.11.001.

464 Department of Health (2016) *How to keep health risks from drinking alcohol to a low level: public consultation*
 465 *on proposed new guideline*. London, UK.

466 Emmett, P., Symes, C., Braddon, F. and Heaton, K. (1992) 'Validation of a new questionnaire for assessing
 467 habitual intakes of starch, non-starch poly-saccharides, sugars and alcohol', *Journal of Human Nutrition and*
 468 *Dietetics*, 5, pp. 245–254.

469 Food Standards Agency (1988) *Food Portion Sizes*. 3rd edn. TSO.

470 Fraser, A., Macdonald-Wallis, C., Tilling, K., Boyd, A., Golding, J., Davey Smith, G., Henderson, J., Macleod, J.,

471 Molloy, L., Ness, A., Ring, S., Nelson, S. M. and Lawlor, D. A. (2013) 'Cohort Profile: the Avon Longitudinal Study

472 of Parents and Children: ALSPAC mothers cohort.', *International journal of epidemiology*, 42(1), pp. 97–110.

473 doi: 10.1093/ije/dys066.

474 French, S., Rosenberg, M. and Knuiman, M. (2008) 'The clustering of health risk behaviours in a Western

475 Australian adult population', *Health Promotion Journal of Australia*. CSIRO PUBLISHING, 19(3), pp. 203–209.

476 doi: 10.1071/HE08203.

477 Golding, J., Pembrey, M., Jones, R. and Team, =ALSPAC Study (2001) 'ALSPAC--the Avon Longitudinal Study of

478 Parents and Children. I. Study methodology', *Paediatric and perinatal epidemiology*. Institute of Child Health,

479 University of Bristol, UK. b.j.stowe@bris.ac.uk, 15(1), pp. 74–87. doi: 10.1046/j.1365-3016.2001.00325.x.

480 Gutierrez, C. M., Ribeiro, C. N. de M., de Lima, G. A., Yanaguita, M. Y. and Peres, L. C. (2007) 'An experimental

481 study on the effects of ethanol and folic acid deficiency, alone or in combination, on pregnant Swiss mice.',

482 *Pathology*. England, 39(5), pp. 495–503. doi: 10.1080/00313020701449290.

483 Herbeth, B., Samara, A., Stathopoulou, M., Siest, G. and Visvikis-Siest, S. (2012) 'Alcohol Consumption,

484 Beverage Preference, and Diet in Middle-Aged Men from the STANISLAS Study.', *Journal of nutrition and*

485 *metabolism*, 2012, p. 987243. doi: 10.1155/2012/987243.

486 HSCIC (2014) *Statistics on Alcohol, England 2014, London: House of Commons*.

487 Hu, F. B. (2002) 'Dietary pattern analysis: a new direction in nutritional epidemiology', *Current Opinion in*

488 *Lipidology*, 13(1).

489 Hutson, J. R., Stade, B., Lehotay, D. C., Collier, C. P. and Kapur, B. M. (2012) 'Folic Acid Transport to the Human

490 Fetus Is Decreased in Pregnancies with Chronic Alcohol Exposure', 7(5), pp. 3–8. doi:

491 10.1371/journal.pone.0038057.

492 Jepson, R. (2000) *The Effectiveness of Intervention to change Health-Related Behaviours: a review of reviews*,

493 *MRC Social & Public Health Sciences Unit*. Glasgow, UK: Medical Research Council.

494 Keen, C. L., Uriu-Adams, J. Y., Skalny, A., Grabeklis, A., Grabeklis, S., Green, K., Yevtushok, L., Wertelecki, W. W.

495 and Chambers, C. D. (2010) 'The plausibility of maternal nutritional status being a contributing factor to the

496 risk for fetal alcohol spectrum disorders: the potential influence of zinc status as an example', *Biofactors*,

497 36(2), pp. 125–135. doi: 10.1002/biof.89.The.

498 Kesse, E., Clavel-Chapelon, F., Slimani, N., van Liere, M. and Group, and the E. (2001) 'Do eating habits differ
 499 according to alcohol consumption? Results of a study of the French cohort of the European Prospective
 500 Investigation into Cancer and Nutrition (E3N-EPIC)', *The American Journal of Clinical Nutrition*, 74(3), pp. 322–
 501 327.

502 Knudsen, V. K., Orozova-Bekkevold, I. M., Mikkelsen, T. B., Wolff, S. and Olsen, S. F. (2007) 'Major dietary
 503 patterns in pregnancy and fetal growth', *Eur J Clin Nutr.* Nature Publishing Group, 62(4), pp. 463–470.

504 Lanting, C. I., Buitendijk, S. E., Crone, M. R., Segaar, D., Gravenhorst, J. B. and van Wouwe, J. P. (2009)
 505 'Clustering of socioeconomic, behavioural, and neonatal risk factors for infant health in pregnant smokers',
 506 *PLoS ONE*, 4(12), pp. 1–6. doi: 10.1371/journal.pone.0008363.

507 May, P. A., Hamrick, K. J., Corbin, K. D., Hasken, J. M., Marais, A. S., Blankenship, J., Hoyme, H. E. and Gossage,
 508 J. P. (2016) 'Maternal nutritional status as a contributing factor for the risk of fetal alcohol spectrum
 509 disorders.', *Reproductive Toxicology*, 59, pp. 101–108. doi: 10.1016/j.reprotox.2015.11.006.MATERNAL.

510 May, P. a, Hamrick, K. J., Corbin, K. D., Hasken, J. M., Marais, A.-S., Brooke, L. E., Blankenship, J., Hoyme, H. E.
 511 and Gossage, J. P. (2014) 'Dietary intake, nutrition, and fetal alcohol spectrum disorders in the Western Cape
 512 Province of South Africa.', *Reproductive toxicology (Elmsford, N.Y.)*. Elsevier Inc., 46C, pp. 31–39. doi:
 513 10.1016/j.reprotox.2014.02.002.

514 Melki, I. S., Beydoun, H. A., Khogali, M., Tamim, H. and Yunis, K. A. (2004) 'Household crowding index: a
 515 correlate of socioeconomic status and inter-pregnancy spacing in an urban setting.', *Journal of epidemiology
 516 and community health*, 58(6), pp. 476–80.

517 Mitchell, J. J., Paiva, M. and Heaton, M. B. (1999) 'Vitamin E and beta-carotene protect against ethanol
 518 combined with ischemia in an embryonic rat hippocampal culture model of fetal alcohol syndrome.',
 519 *Neuroscience letters*, 263(2–3), pp. 189–192. doi: S0304-3940(99)00144-5 [pii].

520 Moran, V. H., Stammers, A. L., Medina, M. W., Patel, S., Dykes, F., Souverein, O. W., Dullemeijer, C., Pérez-
 521 Rodrigo, C., Serra-Majem, L., Nissensohn, M. and Lowe, N. M. (2012) 'The relationship between zinc intake and
 522 serum/plasma zinc concentration in children: A systematic review and dose-response meta-analysis',
 523 *Nutrients*, 4(8), pp. 841–858. doi: 10.3390/nu4080841.

524 Murray, C. J. L., Richards, M. A., Newton, J. N., Fenton, K. A., Anderson, H. R., Atkinson, C., Bennett, D.,
 525 Bernabé, E., Blencowe, H., Bourne, R., Braithwaite, T., Brayne, C., Bruce, N. G., Brugha, T. S., Burney, P.,
 526 Dherani, M., Dolk, H., Edmond, K., Ezzati, M., Flaxman, A. D., Fleming, T. D., Freedman, G., Gunnell, D., Hay, R.

527 J., Hutchings, S. J., Ohno, S. L., Lozano, R., Lyons, R. A., Marcenes, W., Naghavi, M., Newton, C. R., Pearce, N.,
 528 Pope, D., Rushton, L., Salomon, J. A., Shibuya, K., Vos, T., Wang, H., Williams, H. C., Woolf, A. D., Lopez, A. D.
 529 and Davis, A. (2013) 'UK health performance: Findings of the Global Burden of Disease Study 2010', *The Lancet*.
 530 Elsevier Ltd, 381(9871), pp. 997–1020. doi: 10.1016/S0140-6736(13)60355-4.
 531 Murray, D. and Cox, J. L. (1990) 'Screening for depression during pregnancy with the edinburgh depression
 532 scale (EDDS)', *Journal of Reproductive and Infant Psychology*. Routledge, 8(2), pp. 99–107. doi:
 533 10.1080/02646839008403615.
 534 Naseer, M. I., Ullah, I., Ullah, N., Lee, H. Y., Cheon, E. W., Chung, J. and Kim, M. O. (2011) 'Neuroprotective
 535 effect of vitamin C against PTZ induced apoptotic neurodegeneration in adult rat brain.', *Pakistan journal of*
 536 *pharmaceutical sciences*. Pakistan, 24(3), pp. 263–268.
 537 Northstone, K., Emmett, P. M. and Rogers, I. (2008) 'Dietary patterns in pregnancy and associations with
 538 nutrient intakes', *British Journal of Nutrition*, 99(2), pp. 406–415. doi: 10.1017/S0007114507803977.Dietary.
 539 Northstone, K., Emmett, P. and Rogers, I. (2008) 'Dietary patterns in pregnancy and associations with socio-
 540 demographic and lifestyle factors', *European Journal of Clinical Nutrition*, 62(4), pp. 471–479. doi:
 541 10.1038/sj.ejcn.1602741.Dietary.
 542 Patra, J., Bakker, R., Irving, H., Jaddoe, V. W. V, Malini, S. and Rehm, J. (2011) 'Dose-response relationship
 543 between alcohol consumption before and during pregnancy and the risks of low birthweight, preterm birth
 544 and small for gestational age (SGA)-a systematic review and meta-analyses.', *BJOG : an international journal of*
 545 *obstetrics and gynaecology*, 118(12), pp. 1411–21. doi: 10.1111/j.1471-0528.2011.03050.x.
 546 Peng, Y., Kwok, K. H. H., Yang, P. H., Ng, S. S. M., Liu, J., Wong, O. G., He, M. L., Kung, H. F. and Lin, M. C. M.
 547 (2005) 'Ascorbic acid inhibits ROS production, NF- κ B activation and prevents ethanol-induced growth
 548 retardation and microencephaly', *Neuropharmacology*, 48(3), pp. 426–434. doi:
 549 10.1016/j.neuropharm.2004.10.018.
 550 Pierce, D. R. and West, J. R. (1986) 'Blood alcohol concentration: a critical factor for producing fetal alcohol
 551 effects.', *Alcohol (Fayetteville, N.Y.)*, 3(4), pp. 269–272.
 552 Prochaska, J. J. and Prochaska, J. O. (2011) 'A Review of Multiple Health Behavior Change Interventions for
 553 Primary Prevention.', *American journal of lifestyle medicine*, 5(3), pp. 208–221. doi:
 554 10.1177/1559827610391883.
 555 Rogers, I., Emmett, P. and Team, A. study (1998a) 'Diet during pregnancy in a population of pregnant women

556 in South West England', *European journal of clinical nutrition*, 52, pp. 246–250.

557 Rogers, I., Emmett, P. and Team, A. study (1998b) 'Diet during pregnancy in a population of pregnant women

558 in South West England', *European Journal of Clinical Nutrition*, 52, pp. 246–250.

559 Ruf, T., Nagel, G., Altenburg, H.-P., Miller, a B. and Thorand, B. (2005) 'Food and nutrient intake,

560 anthropometric measurements and smoking according to alcohol consumption in the EPIC Heidelberg study.',

561 *Annals of nutrition & metabolism*, 49(1), pp. 16–25. doi: 10.1159/000084173.

562 SACN (2011) *The influence of maternal, fetal and child nutrition on the development of chronic disease in later*

563 *life*.

564 Sayal, K., Heron, J., Golding, J., Alati, R., Smith, G. D., Gray, R. and Emond, A. (2009) 'Binge pattern of alcohol

565 consumption during pregnancy and childhood mental health outcomes: longitudinal population-based study.',

566 *Pediatrics*, 123(2), pp. e289-96. doi: 10.1542/peds.2008-1861.

567 Scholl, T. O., Chen, X., Sims, M. and Stein, T. P. (2006) 'Vitamin E: maternal concentrations are associated with

568 fetal growth', *The American Journal of Clinical Nutrition* , 84(6), pp. 1442–1448.

569 Shankar, A., McMunn, A. and Steptoe, A. (2010) 'Health-related behaviors in older adults relationships with

570 socioeconomic status.', *American journal of preventive medicine*, 38(1), pp. 39–46. doi:

571 10.1016/j.amepre.2009.08.026.

572 Skagerström, J., Chang, G. and Nilsen, P. (2011) 'Predictors of drinking during pregnancy: a systematic review.',

573 *Journal of women's health (2002)*, 20(6), pp. 901–13. doi: 10.1089/jwh.2010.2216.

574 Smith, A. M. and Smith, C. (1994) 'Dietary intake and lifestyle patterns: correlates with socio-economic,

575 demographic and environmental factors', *Journal of Human Nutrition and Dietetics*. Blackwell Publishing Ltd,

576 7(4), pp. 283–294. doi: 10.1111/j.1365-277X.1994.tb00271.x.

577 Streissguth, A. P., Barr, H. M. and Sampson, P. D. (1990) 'Moderate Prenatal Alcohol Exposure: Effects on Child

578 IQ and Learning Problems at Age 7 1/2 Years', *Alcoholism: Clinical & Experimental Research*, 14(5), pp. 662–

579 669.

580 Thomas, J. D., Idrus, N. M., Monk, B. R. and Dominguez, H. D. (2010) 'Prenatal choline supplementation

581 mitigates behavioral alterations associated with prenatal alcohol exposure in rats.', *Birth defects research. Part*

582 *A, Clinical and molecular teratology*. United States: Center for Behavioral Teratology, Department of

583 Psychology, San Diego State University, 6363 Alvarado Court, San Diego, CA 92120, USA.

584 thomas3@mail.sdsu.edu, 88(10), pp. 827–837.

Thompson, F. E. and Subar, A. F. (2008) 'Dietary Assessment Methodology', in Coulton, S., Boushey, C., and Ferruzzi, M. (eds) *Nutrition in Prevention and Treatment of Disease*. 2nd edn. San Diego: Elsevier Inc., pp. 3–11.

Touvier, M., Druet-Pecollo, N., Kesse-Guyot, E., Andreeva, V. a, Galan, P., Hercberg, S. and Latino-Martel, P. (2014) 'Demographic, socioeconomic, disease history, dietary and lifestyle cancer risk factors associated with alcohol consumption.', *International journal of cancer. Journal international du cancer*, 134(2), pp. 445–59. doi: 10.1002/ijc.28365.

Tucker, K. L., Selhub, J., Wilson, P. and Rosenberg, I. (1996) 'Human and Clinical Nutrition Dietary Intake Pattern Relates to Plasma Folate and Homocysteine Concentrations in the Framingham Heart Study', *The Journal of Nutrition*, 126, pp. 3025–3031.

Valencia-Martin, J. L., Galan, I. and Rodriguez-Artalejo, F. (2011) 'The association between alcohol consumption patterns and adherence to food consumption guidelines', *Alcohol Clin Exp Res*. 2011/08/19, 35(11), pp. 2075–2081. doi: 10.1111/j.1530-0277.2011.01559.x.

La Vecchia, C., Negri, E., Franceschi, S., Parazzini, F. and Decarli, a (1992) 'Differences in dietary intake with smoking, alcohol, and education.', *Nutrition and cancer*, 17(3), pp. 297–304. doi: 10.1080/01635589209514199.

Veena, S., Krishnaveni, G., Srinivasan, K., Wills, A., Muthayya, S., Kurpad, A., Yajnik, C. and Fall, C. (2010) 'Higher maternal plasma folate but not vitamin B-12 concentrations during pregnancy are associated with better cognitive function scores in 9-to 10-year-old children in South India', *The Journal of Nutrition*, 140(5), pp. 1014–1022. doi: 10.3945/jn.109.118075.mothers.

Völgyi, E., Carroll, K. N., Hare, M. E., Ringwald-Smith, K., Piyathilake, C., Yoo, W. and Tyllavsky, F. A. (2013) 'Dietary patterns in pregnancy and effects on nutrient intake in the Mid-South: the Conditions Affecting Neurocognitive Development and Learning in Early Childhood (CANDLE) study.', *Nutrients*. Multidisciplinary Digital Publishing Institute, 5(5), pp. 1511–30. doi: 10.3390/nu5051511.

Wang, L. L., Zhang, Z., Li, Q., Yang, R., Pei, X., Xu, Y., Wang, J., Zhou, S. F. and Li, Y. (2009) 'Ethanol exposure induces differential microRNA and target gene expression and teratogenic effects which can be suppressed by folic acid supplementation', *Hum Reprod*. 2008/12/19, 24(3), pp. 562–579. doi: 10.1093/humrep/den439.

Weiss, L. A. and Chambers, C. D. (2013) 'Associations Between Multivitamin Supplement Use and Alcohol Consumption Before Pregnancy: Pregnancy Risk Assessment Monitoring System, 2004 to 2008', *Alcohol Clin Exp Res*, 37(9), pp. 1595–1600. doi: 10.1038/jid.2014.371.

614 Wentzel, P. and Eriksson, U. J. (2006) 'Ethanol-induced fetal dysmorphogenesis in the mouse is diminished by
 615 high antioxidative capacity of the mother.', *Toxicological sciences : an official journal of the Society of*
 616 *Toxicology*. United States, 92(2), pp. 416–422. doi: 10.1093/toxsci/kfl024.
 617 Yarnell, J. W., Fehily, A. M., Milbank, J. E., Sweetnam, P. M. and Walker, C. L. (1983) 'A short dietary
 618 questionnaire for use in an epidemiological survey: comparison with weighed dietary records', *Human*
 619 *nutrition. Applied nutrition*, 37(2), pp. 103–112.
 620 Zuccolo, L., Lewis, S. J., Smith, G. D., Sayal, K., Draper, E. S., Fraser, R., Barrow, M., Alati, R., Ring, S., Macleod,
 621 J., Golding, J., Heron, J. and Gray, R. (2013) 'Prenatal alcohol exposure and offspring cognition and school
 622 performance. A "Mendelian randomization" natural experiment.', *International journal of epidemiology*, 42(5),
 623 pp. 1358–70. doi: 10.1093/ije/dyt172.
 624

625 **Table 1.** Socio-demographic and lifestyle characteristics of sample population (n (%))

	Total	Alcohol consumption during first trimester (n= 9,839)					Binge drinking (n= 9,781)		
		Never	<1 drink/week	1-6 drinks/week	1+ drink/day	p-value*	No binge drinking	≥1 episodes of binge drinking	p-value*
Age (years)									
<20	272 (3)	150 (4)	77 (2)	39 (3)	6 (4)		250 (3)	19 (3)	
20-24	5572 (57)	2697 (61)	2185 (56)	634 (47)	56 (37)		5149 (57)	383 (54)	
25-29	3874 (39)	1502 (34)	1619 (41)	666 (49)	87 (57)		3567 (39)	294 (42)	
30+	121 (1)	52 (1)	49 (1)	16 (1)	4 (3)	<0.0001	110 (1)	9 (1)	0.647
Parity									
Primiparous	4370 (44)	2107 (48)	1621 (41)	567 (42)	75 (49)		4089 (45)	264 (37)	
Multiparous	5469 (56)	2294 (52)	2309 (59)	788 (58)	78 (51)	<0.0001	4987 (55)	441 (63)	<0.0001
Ethnicity									
Non-white	206 (2)	121 (3)	66 (2)	18 (1)	1 (1)		186 (2)	11 (2)	
White	9633 (98)	4280 (97)	3864 (98)	1337 (99)	152 (99)	0.004	8890 (98)	694 (98)	0.373
Maternal smoking									
Smoker	1766 (18)	716 (16)	672 (17)	316 (23)	62 (41)		1520 (17)	238 (34)	
Non-smoker	8073 (82)	3685 (84)	3258 (83)	1039 (77)	91 (59)	<0.0001	7556 (83)	467 (66)	<0.0001
Education									
Vocational	2655 (27)	1276 (29)	971 (25)	359 (26)	49 (32)		2354 (26)	269 (38)	
O-level	3535 (36)	1595 (36)	1450 (37)	442 (33)	48 (31)		3281 (36)	240 (34)	
A-level	2320 (24)	1018 (23)	934 (24)	341 (25)	27 (18)		2163 (24)	148 (21)	
Degree level	1329 (14)	512 (12)	575 (15)	213 (16)	29 (19)	<0.0001	1278 (14)	48 (7)	<0.0001
Single parent household									
No	9334 (95)	4190 (95)	3760 (96)	1250 (92)	134 (88)		8627 (95)	654 (93)	
Yes	505 (5)	211 (5)	170 (4)	105 (8)	19 (12)	<0.0001	449 (5)	51 (7)	0.008
Home ownership									
Owner/occupied	7756 (79)	3433 (78)	3154 (80)	1063 (78)	106 (69)		7224 (80)	490 (70)	
Council/HA rented	1232 (13)	599 (14)	449 (11)	158 (12)	26 (17)		1092 (12)	128 (18)	
Private rent/other	851 (9)	369 (8)	327 (8)	134 (10)	21 (14)	0.001	760 (8)	87 (12)	<0.0001
House Crowding Index									
≤0.5	4379 (45)	1973 (45)	1729 (44)	615 (45)	62 (41)		4105 (45)	253 (36)	
>0.5-0.75	3159 (32)	1381 (31)	1325 (34)	416 (31)	38 (25)		2926 (32)	217 (31)	
>0.75-1	1795 (18)	817 (19)	684 (17)	263 (19)	31 (20)		1607 (18)	174 (25)	
>1	506 (5)	230 (5)	192 (5)	62 (5)	22 (14)	<0.0001	438 (5)	61 (9)	<0.0001
EPDS score									
<13	8467 (86)	3780 (86)	3423 (87)	1147 (85)	117 (76)		7863 (87)	557 (79)	
≥13	1372 (14)	621 (14)	507 (13)	208 (15)	36 (24)	<0.0001	1213 (13)	148 (21)	<0.0001
Total	9839	4401	3930	1355	153		9076	705	

*chi-squared test

626

627

628
629
630

Table 2. Unadjusted and adjusted* beta-coefficients and 95% CI of dietary patterns scores by maternal alcohol consumption during pregnancy

		Health conscious			Traditional			Processed			Confectionery			Vegetarian		
Alcohol consumption		β	95% CI	p	β	95% CI	p	β	95% CI	p	β	95% CI	p	β	95% CI	p
during first trimester																
Unadjusted																
Never			(ref)			(ref)			(ref)			(ref)			(ref)	
<1 drink/week	0.10	(0.06, 0.14)	<0.0001	-0.01	(-0.05, 0.03)	0.511	0.01	(-0.03, 0.05)	0.673	0.07	(0.03, 0.11)	0.001	-0.08	(-0.12, -0.04)	<0.001	
1-6 drinks/week	0.12	(0.06, 0.17)	<0.0001	-0.05	(-0.11, 0.00)	0.074	0.08	(0.03, 0.14)	0.004	0.10	(0.04, 0.16)	<0.0001	0.02	(-0.03, 0.08)	0.391	
1+ drink/day	-0.10	(-0.23, 0.04)	0.178	0.09	(-0.05, 0.22)	0.230	0.17	(0.03, 0.31)	0.015	-0.07	(-0.21, 0.07)	0.353	0.23	(0.09, 0.37)	0.001	
Adjusted*																
Never			(ref)			(ref)			(ref)			(ref)			(ref)	
<1 drink/week	0.03	(-0.01, 0.06)	0.167	-0.04	(-0.08, 0.01)	0.067	0.04	(0.00, 0.08)	0.049	0.07	(0.03, 0.11)	0.002	-0.06	(-0.1, -0.02)	0.009	
1-6 drinks/week	0.08	(0.03, 0.13)	0.002	-0.07	(-0.13, -0.01)	0.031	0.10	(0.04, 0.15)	<0.0001	0.11	(0.05, 0.17)	0.001	0.02	(-0.04, 0.08)	0.575	
1+ drink/day	0.02	(-0.1, 0.15)	0.784	0.08	(-0.08, 0.24)	0.318	0.24	(0.09, 0.39)	0.002	0.03	(-0.13, 0.19)	0.702	0.10	(-0.06, 0.26)	0.242	

*adjusted for maternal age, parity, ethnicity, education, smoking, housing tenure, HCl, single parent household and EPDS score

CI = confidence intervals

631
632
633
634
635

636 **Table 3.** Unadjusted and adjusted* beta-coefficients and 95% CI of dietary patterns scores by maternal binge drinking during pregnancy

Binge drinking	Health conscious			Traditional			Processed			Confectionery			Vegetarian		
	β	95% CI	p	β	95% CI	p	β	95% CI	p	β	95% CI	p	β	95% CI	p
Unadjusted															
No binge drinking		(ref)			(ref)			(ref)			(ref)			(ref)	
≥1 episodes of binge drinking	-0.28	(-0.35, -0.21)	<0.0001	0.05	(-0.02, 0.11)	0.187	0.23	(0.17, 0.30)	<0.0001	-0.07	(-0.14, -0.01)	0.034	0.09	(0.03, 0.16)	0.007
Adjusted*															
No binge drinking		(ref)			(ref)			(ref)			(ref)			(ref)	
≥1 episodes of binge drinking	-0.05	(-0.12, 0.01)	0.103	0.06	(-0.02, 0.14)	0.119	0.15	(0.07, 0.22)	<0.0001	-0.07	(-0.15, 0.00)	0.057	0.05	(-0.03, 0.13)	0.196

*adjusted for maternal age, parity, ethnicity, education, smoking, housing tenure, HCl, single parent household and EPDS score

CI = confidence intervals




Figure 1: Differences in dietary pattern scores between categories of maternal alcohol consumption during pregnancy, relative to 'Never' drinkers (Green: <1 drink/week; Orange: 1-6 drinks/week; Red: ≥1 drinks/day). Differences in dietary pattern scores between 'binge' drinkers (≥4 drinks/day at any time during the previous month) and non-binge drinkers, are shown in blue.

All values are adjusted beta-coefficients and 95% CIs. Estimates are adjusted for: maternal age, parity, ethnicity, education, and smoking, housing tenure, HCl, single parent household and EPDS score.

*P <0.05 ** P<0.01 *** P<0.001

Table Legends

Table 1. Socio-demographic and lifestyle characteristics of sample population

Table 2. Unadjusted and adjusted* beta-coefficients and 95% CI of dietary patterns scores by maternal alcohol consumption during pregnancy

Table 3. Unadjusted and adjusted* beta-coefficients and 95% CI of dietary patterns scores by maternal binge drinking during pregnancy

Figure 1: Differences in dietary pattern scores between categories of maternal alcohol consumption during pregnancy, relative to 'Never' drinkers (Green: <1 drink/week; Orange: 1-6 drinks/week; Red: ≥ 1 drinks/day). Differences in dietary pattern scores between 'binge' drinkers (≥ 4 drinks/day at any time during the previous month) and non-binge drinkers are shown in blue.

Supplementary data

Table 1. Socio-demographic and lifestyle characteristics of included and excluded participants

Supplementary data

Table 1. Socio-demographic and lifestyle characteristics of included and excluded participants

	Included (n= 9,839)		Missing*	
	n	%	n	%
Age (years)				
<20	272	3	230	10
20-24	5572	57	1454	64
25-29	3874	39	554	24
30+	121	1	28	1
Parity				
Primiparous	4370	44	648	44
Multiparous	5469	56	828	56
Ethnicity				
Non-white	206	2	49	7
White	9633	98	701	93
Maternal smoking				
Smoker	1766	18	1078	68
Non-smoker	8073	82	509	32
Education				
Vocational	2655	27	376	48
O-level	3535	36	223	28
A-level	2320	24	116	15
Degree level	1329	14	68	9
Single parent household				
No	9334	95	1626	90
Yes	505	5	186	10
Home ownership				
Owner/occupied	7756	79	1089	57
Council/HA rented	1232	13	537	28
Private rent/other	851	9	290	15
House Crowding Index				
≤0.5	4379	45	533	29
>0.5-0.75	3159	32	533	29
>0.75-1	1795	18	518	28
>1	506	5	252	14
EPDS score				
<13	8467	86	339	77
≥13	1372	14	104	23
Alcohol consumption during 1st trimester				
Never	4401	45	663	11
<1 drink/week	3930	40	459	9
1-6 drinks/week	1355	14	199	11
1+ drink/day	153	2	41	17
Binge drinking				
No binge drinking	9076	93	1178	10
≥1 episodes of binge drinking	705	7	137	14

*Participants with missing dietary data at 32 weeks gestation

Unadjusted logistic regression models indicated that women with missing dietary data were more likely to be younger (OR=0.94, 95%CI=0.93, 0.95; $p<0.0001$), smoke (OR=1.01, 95%CI=1.01, 1.01), live in a single parent household (OR=1.78, 95%CI=1.48, 2.15; $p<0.0001$), live in rented accommodation (OR=1.61, 95%CI=1.51, 1.72; $p<0.0001$) and in more crowded conditions (OR=1.54, 95%CI=1.46, 1.62; $p<0.0001$), drink 1+drinks/day (OR=1.72, 95%CI=1.22, 2.43; $p<0.0001$) and binge drink during pregnancy (OR=1.37, 95%CI=1.12, 1.66; $p=0.002$)