

initiatives showed improvements in child nutrition and education.²² However, to our knowledge, no studies to date in high-income or low-income countries have examined impacts of different combinations of social protection on adolescent HIV risks. In 2014, UNICEF, UNAIDS, UNDP, PEPFAR-USAID, and the World Bank conducted a series of high-level meetings. These aimed to ascertain the evidence needs of Southern and Eastern African governments regarding social protection and HIV prevention (these included the Launch of the World Bank, UNAIDS, and UNICEF Research Network on Structural Drivers and Solutions for HIV. International AIDS Conference, Melbourne 2014, the UNAIDS Planning Coordination Board 2014. Geneva, July, High Level Consultation on Scaling-up Proven Social and Structural Interventions to Prevent HIV Transmission, Johannesburg, July). Policy makers identified 2 key and unanswered requirements: first, to distinguish which specific types of social protection interventions are effective in adolescent HIV-risk reduction; second, to test whether there are cumulative prevention benefits of combination social protection. This study aims to address these questions in a multisite longitudinal sample from South Africa.

METHODS

Participants and Procedures

Three thousand five hundred sixteen adolescents aged 10–18 (56.7% female) were interviewed at baseline (2009–2010) and followed up at 1 year (2011–2012). Refusal rate at baseline was <2.5%, and 1-year retention rate was 96.8%. Two South African provinces were selected in consultation with the South African National Departments of Social Development, Health and Education: Mpumalanga and the Western Cape. Within these provinces, 2 urban and 2 rural health districts with >30% antenatal HIV prevalence were randomly selected. Within each health district, sequentially numbered census enumeration areas were randomly sampled until sample size was reached. In each area, every household was visited and was included in the study if they had a resident adolescent. One adolescent per household was interviewed face to face for 60–70 minutes. Questionnaires and consent forms were translated into Xhosa, Zulu, Sotho, and Tsonga and checked with back translation. Adolescents participated in the language of their choice.

Ethical protocols were approved by the Universities of Oxford, Cape Town, and KwaZulu-Natal and by provincial Health and Education Departments. Adolescents and primary caregivers provided voluntary informed consent. No incentives were given apart from refreshments and certificates of participation. All interviewers were trained in working with vulnerable youth, and confidentiality was maintained except where participants were at risk of significant harm or requested assistance. Where participants reported abuse, rape, or risk of significant harm, referrals were made to child protection, health, and HIV/AIDS services with follow-up support.

Measures

Adolescent HIV-risk behaviors were measured at baseline, and new incidence of HIV-risk behaviors measured

at follow-up, using scales from the National Survey of HIV and Sexual Behavior among Young South Africans and the SA Demographic and Health Survey.²³ Previous evidence suggests that HIV-risk behaviors may cluster, and thus risks were grouped into risks with evidence of poverty, deprivation, and exploitation drivers: “economic sex” (transactional and age-disparate sex) and risks with evidence of adolescent developmental level and social drivers: “incautious sex” (unprotected sex, multiple partners, casual partners, and sex using substances) and “pregnancy” (females only, current and past pregnancy). Transactional sex was past-year incidence of sex in exchange for food, shelter, school fees, transport, or money; age-disparate sex was past-year incidence of having a sexual partner >5 years older than the adolescent.²² Unprotected sex was “sometimes,” “rarely,” or “never” using condoms when having sex in the past year (versus “always” using condoms or no sexual activity). Multiple sexual partners was having ≥ 2 partners in the past year.²⁴ Casual partners was having sexual partners who were not regular boyfriends/girlfriends in the past year. Sex using substances was past-year sex while drunk or using any drug. Pregnancy history and current pregnancy—both a marker of unprotected sex and also a risk factor for increased unprotected sex and sero-conversion for adolescents²⁵—were measured at baseline and follow-up.

Social protection and social care provisions were measured at baseline for 14 components (this study aimed to examine the effects of social protection in real-world Southern African conditions. To maximize utility, we measured cash and care services that are typically provided by governments, nongovernmental organization, and families. These were identified in consultation with the SA National Departments of Social Development, Basic Education and Health, PEPFAR-USAID, UNICEF, and Save the Children). Child-focused cash transfer receipt was measured as household access to either a child support grant or foster child grant.⁷ Pension was within-household access to an old-age grant. School feeding was daily, free lunches provided at school, and provision of free school transport and free school uniform were also measured. Access to food gardens was receiving food from a school or community garden; food parcels or soup kitchen feeding was regular, reliable provision of food parcels to the household or free meals from any organization. Home-based and community-based carer support was at least monthly household visits from a home-based caregiver, nurse, or volunteer providing medical and social support. Teacher social support was social, practical, or emotional support from a teacher, using a standardized scale used previously in South Africa.²⁶ School counselor was past-year school-based counseling. Positive parenting (eg, primary caregiver praise and warmth) and parental monitoring (eg, having set times to be home, parental knowledge of whereabouts) were measured using validated Alabama Parenting Questionnaire subscales.²⁷ Free schooling required both no-fees school and free schoolbooks (many “free” schools required financial contributions for schoolbooks to allow children to take examinations). Evidence suggests that to show effects, social protection requires sustained and predictable duration and also current receipt.²⁸ Consequently, each intervention and care provision were dichotomized into

receipt/no receipt, with positive coding requiring exposure at both baseline and 1-year follow-up.

Preselected covariates included factors potentially affecting HIV-risk or social protection access. Adolescent age, urban/rural location, school enrollment, informal/formal housing, migration, number of children in the home, female primary caregiver, and maternal and paternal death were measured using items adapted from the South African Census. Food insecurity was measured using items from the National Food Consumption Survey⁷ and determined as days in the past week with insufficient food in the home. HIV prevention knowledge at baseline and follow-up was measured using a “free-listing” approach, asking “can you write any things you think a person can do to avoid getting HIV or AIDS?” as preexisting lists of true and false prevention methods can overestimate knowledge levels. Scores were calculated by summed number of accurate methods (eg, use a condom) minus summed inaccurate methods (eg, do not share food with an HIV-positive person). Participants also indicated whether they had a birth certificate, a frequent requirement for accessing social protection programs.

Analyses

Analyses were conducted in 3 stages in SPSS 21.0 and STATA 13.1, using the longitudinal sample ($n = 3401$). Because few adolescents younger than 12 years reported ever having sex ($n = 9$), analyses were limited to the 2668 adolescents aged 12–18 years (male, 1170 and female, 1498). Analyses were gender disaggregated, as previous studies have suggested gender differences in HIV risk and protective factors.¹⁰

First, descriptive statistics for all outcomes, social protection variables, and covariates were calculated (Table 1). Social protection types were excluded where numbers reached were too small for reliable analysis. To check for multicollinearity, social protections and covariates were entered into a linear regression for each of incautious sex, economic sex, and pregnancy. Both tolerance values and the spread of variance proportions on small eigenvalues suggested no multicollinearity.

Second, associations between specific social protections and HIV risk behaviors were assessed in multivariable logistic regressions. To identify social protections with independent effects, 2 models were run.²⁹ In model 1, all interventions and covariates were included with each of the following HIV risk behaviors at time 2 as outcomes: incautious sex (males and females), economic sex (males and females), and pregnancy (females only). Each analysis at this stage controlled for covariates, HIV risk or social protection access: baseline HIV-risk behavior, age, poverty, urban–rural location, school nonenrollment, informal/formal housing, migration, number of children in the home, female primary caregiver, maternal and paternal death, possession of a birth certificate, and HIV knowledge. For model 2, all preselected covariates were included while interventions were selected for inclusion on the basis of $P \leq 0.10$, to prevent potentially important variables from being omitted.²⁸ Covariates were included in both models to control for potential confounding due to the nonrandomized allocation of interventions.

TABLE 1. Characteristics of 2668 Adolescents Aged 12–18 Years Included in Analyses

	All	Girls	Boys
Number	2668	1498	1170
Social protection interventions (received at times 1 and 2)			
Child support or foster child grant	1486 (55.7%)	852 (56.9%)	634 (54.2%)
Pension	136 (5.1%)	81 (5.4%)	55 (4.7%)
Free school and books	1398 (52.4%)	798 (53.3%)	600 (51.3%)
Teaching support	211 (7.9%)	121 (8.1%)	90 (7.7%)
School counselor	98 (3.7%)	55 (3.7%)	43 (3.7%)
Free school meals	1930 (72.3%)	1064 (71.0%)	866 (74.0%)
Food gardens	132 (4.9%)	67 (4.5%)	65 (5.6%)
Food parcels or soup kitchens	13 (0.5%)	10 (0.7%)	3 (0.3%)
Positive parenting	664 (24.9%)	376 (25.1%)	288 (24.6%)
Good parental monitoring	572 (21.4%)	358 (23.9%)	214 (18.3%)
Outcomes (time 2)*			
Economically driven sex	156 (5.8%)	99 (6.6%)	57 (4.9%)
Incautious sex	490 (18.4%)	240 (16.0%)	250 (21.4%)
Ever pregnant	—	71 (4.7%)	—
Covariates†			
Age	14.2 (1.6)	14.3 (1.7)	14.2 (1.6)
No. days without enough food in the household	0.9 (1.6)	0.9 (1.6)	0.8 (1.6)
Informal housing	820 (30.7%)	471 (31.4%)	349 (29.8%)
Urban location	1309 (49.1%)	717 (47.9%)	592 (50.6%)
School nonenrollment	66 (2.5%)	40 (2.7%)	26 (2.2%)
Moved more than twice	935 (35.0%)	558 (37.2%)	377 (32.2%)
No. children in the home	2.6 (1.5)	2.7 (1.5)	2.5 (1.3)
Female primary caregiver	2366 (88.7%)	1354 (90.4%)	1012 (86.5%)
Mother dead	326 (12.2%)	170 (11.3%)	156 (13.3%)
Father dead	588 (22.0%)	350 (23.4%)	238 (20.3%)
Child has birth certificate	2544 (95.4%)	1416 (94.7%)	1128 (96.4%)
HIV knowledge	1.8 (1.4)	1.8 (1.5)	1.8 (1.3)
Economically driven sex	113 (4.2%)	85 (5.7%)	28 (2.4%)
Incautious sex	339 (12.7%)	171 (11.4%)	168 (14.4%)
Ever pregnant	—	46 (3.1%)	—

Data are number of participants, number of participants (%), or mean (SD).

*Reporting both incautious and economic sex: 4% boys and 5% girls.

†All covariates were measured at time 1, except HIV knowledge, which is included from time 2.

Third, we tested whether there are greater HIV prevention effects from combining social protections. All social protection variables and covariates that were included in model 2 were entered into a marginal effects analysis in

STATA using binary logistic regression for each outcome. Each possible combination of included social protection variables on HIV-risk behaviors was tested in each analysis, with all covariates held at their mean values, to determine how the predicted probability of the outcome alters when different interventions (and combinations of interventions) are present.

Role of the Funding Source

The study's sponsors had no role in study design, data collection, analysis, interpretation, or writing of the report. The corresponding author had full access to all data in the study and had final responsibility for the decision to submit for publication.

RESULTS

Sample Characteristics and Social Protection Access

Respondents had a mean age of 14.2 years at follow-up (14.2 males and 14.3 females, both ranged 10–19) (Table 1). They reported a mean of 2.3 days without enough food in the past week. Thirty-one percent lived in informal (shack) housing, and 2.5% had left school before completion. Five percent of boys and 7% of girls reported economically driven sex at follow-up. Twenty-one percent of boys and 16% of girls reported incautious sex at follow-up. Female pregnancy at follow-up was 5%.

Social protection interventions that reached less than 100 adolescents (<0.3% of the sample) were excluded (school counselors, food parcels, and soup kitchens). Remaining social protections were the child support/foster child grant (received by 56% of adolescents), free school meals (72%), pension in the household (5%), free schooling and books (52%), teacher support (8%), school counseling (4%), access to food gardens (5%), positive parenting (25%), and good parental monitoring (21%).

Independent Associations of Social Protection Interventions With HIV-Risk Incidence

Table 2 shows all factors that remained associated at $P < 0.10$ with reduced HIV-risk behaviors in multivariable regressions, independent of all other social protection factors and covariates. For males, reduced incidence of incautious sex was significantly associated ($P < 0.05$) with free schooling [odds ratio (OR): 0.69, CI: 0.48 to 0.99], parental monitoring (OR: 0.51, CI: 0.30 to 0.87), and teacher support (OR: 0.46, CI: 0.21 to 0.99), whereas reduced incidence of economic sex was associated with parental monitoring (OR: 0.10, CI: 0.01 to 0.70). For females, reduced incidence of incautious sex was associated with school feeding (OR: 0.64, CI: 0.44 to 0.93) and reduced incidence of economic sex was associated with child grants (OR: 0.51, CI: 0.32 to 0.83) and free schooling (OR: 0.36, CI: 0.22 to 0.60). Reduced incidence of pregnancy was associated with school feeding (OR: 0.32, CI: 0.15 to 0.67). Although receipt of child grants

met the $P < 0.10$ threshold to be included in model 2 regressions for pregnancy, having children as an adolescent increases the probability of receiving child grants (ie, a reverse causality). Consequently, receipt of child grants was treated as a covariate rather than a predictor of pregnancy in these models.

Associations of Combination Social Protection Interventions With HIV-Risk Behavior Incidence

Where 2 or more social protections were at $P < 0.10$, interval estimates of the probability of the outcome showed which combinations of social protection interventions were significant, whilst controlling for covariates and baseline HIV-risk behavior (Figs. 1, 2). Analyses identified social protections that separately were not significantly associated with reduced HIV-risk behavior incidence but showed significant reductions in outcome probability when combined with other social protections.

For boys, predicted probability of past-year incidence of incautious sex was 18.7% when receiving none of the included social protection interventions of free schooling, teacher support, or parental monitoring. With free school alone, it was 13.7%; with good parental monitoring alone, 10.4%; and with teacher support alone, 9.5%. But with both free school and parental monitoring, boys' incidence of incautious sex was 7.5%; with free school and teacher support, 6.8%; and with parental monitoring and teacher support, 5.1%. Among boys who received all 3 interventions, past-year incidence of incautious sex was 3.5%.

For girls, parental monitoring was not independently associated with incautious sex but, when combined with school feeding, showed stronger associations than school feeding alone. Thus, girls' past-year incidence of incautious sex when receiving none of the included social protection interventions was 15.1%. With good parental monitoring alone, it was 10.5%, and with school feeding alone, 10.2%. When both of these interventions were received, girls' past-year incidence of incautious sex dropped to 6.9%.

Among girls, economically driven sex also showed combination effects of social protection. Girls' past-year incidence of economic sex when receiving none of the included social protection interventions was 10.5%. With parental monitoring alone, incidence was 6.8%; with a child-focused grant alone, 5.7%; and with free schooling alone, 4.1%. But with both parental monitoring and a child grant, girls' incidence of economically driven sex was 3.6%; with parental monitoring and free school, 2.6%; and with a child grant and free school, 2.1%. Among girls who received all 3 interventions, past-year incidence of economically driven sex was 2.1%.

Finally, combination social protection effects were shown on adolescent female pregnancy. Past-year incidence of pregnancy when receiving none of the included social protection interventions was 5.5%. With good parental monitoring alone, it was 3.0%; with free schooling, 2.5%; and with school feeding, 1.9%. But with both parental monitoring and free school, incidence of pregnancy was 1.3%; with parental

TABLE 2. Interventions and Covariates Associated With Incident HIV-Risk Behaviors in Multivariable Binary Logistic Regression Analyses for Girls and Boys

	Girls (n = 1498)					
	Incautious Sex*		Economic Sex†		Pregnancy‡	
	AOR (95% CI)	P	AOR (95% CI)	P	AOR (95% CI)	P
Child grants	—	—	0.51 (0.32 to 0.83)	0.007	1.64 (0.89 to 3.00)	0.110
Free school and books	—	—	0.36 (0.22 to 0.60)	<0.0001	0.47 (0.21 to 1.03)	0.59
School feeding	0.64 (0.44 to 0.93)	0.020	—	—	0.32 (0.15 to 0.67)	0.003
Parental monitoring	0.66 (0.43 to 1.01)	0.056	0.62 (0.34 to 1.15)	0.130	0.55 (0.25 to 1.20)	0.134
Teacher support	—	—	—	—	—	—
Incautious sex, time 1	6.36 (4.24 to 9.54)	<0.0001	—	—	—	—
Economic sex, time 1	—	—	3.78 (2.05 to 6.98)	<0.0001	—	—
Pregnancy, time 1	—	—	—	—	14.95 (6.77 to 33.03)	<0.0001
Age	1.61 (1.44 to 1.81)	<0.0001	1.32 (1.14 to 1.53)	0.0002	1.73 (1.38 to 2.16)	<0.0001
Food insecurity	1.10 (1.00 to 1.21)	0.041	0.93 (0.81 to 1.06)	0.280	1.02 (0.87 to 1.20)	0.827
Informal housing	0.61 (0.41 to 0.91)	0.016	1.33 (0.80 to 2.20)	0.271	0.36 (0.17 to 0.77)	0.008
Urban location	1.01 (0.72 to 1.44)	0.942	1.02 (0.64 to 1.63)	0.922	1.40 (0.76 to 2.59)	0.287
School nonenrollment	0.99 (0.41 to 2.29)	0.981	0.63 (0.22 to 1.78)	0.380	0.84 (0.25 to 2.78)	0.772
Migration	1.60 (1.11 to 2.29)	0.011	1.12 (0.69 to 1.82)	0.638	2.13 (1.15 to 3.93)	0.016
No. children in the home	0.99 (0.89 to 1.09)	0.786	0.94 (0.81 to 1.09)	0.400	0.86 (0.71 to 1.03)	0.099
Female primary caregiver	1.11 (0.65 to 1.92)	0.697	1.66 (0.75 to 3.65)	0.211	2.73 (0.68 to 11.00)	0.157
Mother dead	1.05 (0.65 to 1.69)	0.851	1.86 (1.04 to 3.34)	0.038	0.88 (0.38 to 2.02)	0.760
Father dead	1.05 (0.73 to 1.53)	0.778	1.15 (0.71 to 1.87)	0.560	1.77 (1.00 to 3.15)	0.051
Child has birth certificate	1.01 (0.53 to 1.92)	0.987	0.76 (0.36 to 1.62)	0.479	0.98 (0.37 to 2.60)	0.961
HIV knowledge	0.87 (0.70 to 1.09)	0.216	0.91 (0.69 to 1.22)	0.538	1.28 (0.89 to 1.86)	0.187

	Boys (n = 1170)			
	Incautious Sex§		Economic Sex	
	AOR (95% CI)	P	AOR (95% CI)	P
Child grants	—	—	—	—
Free school and books	0.69 (0.48 to 0.99)	0.046	—	—
School feeding	—	—	—	—
Parental monitoring	0.51 (0.30 to 0.87)	0.013	0.10 (0.01 to 0.70)	0.021
Teacher support	0.46 (0.21 to 0.99)	0.046	—	—
Incautious sex, time 1	3.82 (2.57 to 5.68)	<0.0001	—	—
Economic sex, time 1	—	—	3.32 (1.21 to 9.09)	0.019
Pregnancy, time 1	—	—	—	—
Age	1.67 (1.50 to 1.86)	<0.0001	1.46 (1.22 to 1.75)	<0.0001
Food insecurity	1.04 (0.94 to 1.16)	0.414	1.04 (0.87 to 1.24)	0.662
Informal housing	1.17 (0.80 to 1.70)	0.428	0.72 (0.38 to 1.37)	0.315
Urban location	1.11 (0.77 to 1.58)	0.577	1.02 (0.58 to 1.80)	0.936
School nonenrollment	2.39 (0.60 to 9.55)	0.216	3.27 (0.62 to 17.27)	0.163
Migration	1.60 (1.11 to 2.30)	0.012	1.12 (0.60 to 2.09)	0.715
No. children in the home	0.90 (0.79 to 1.03)	0.113	0.86 (0.68 to 1.08)	0.195
Female primary caregiver	1.29 (0.79 to 2.08)	0.310	0.82 (0.38 to 1.74)	0.602
Mother dead	0.69 (0.40 to 1.16)	0.161	0.49 (0.17 to 1.45)	0.196
Father dead	0.88 (0.58 to 1.33)	0.546	0.60 (0.27 to 1.33)	0.206
Child has birth certificate	1.08 (0.46 to 2.57)	0.856	3.07 (0.38 to 24.97)	0.294
HIV knowledge	1.15 (0.91 to 1.45)	0.258	0.98 (0.65 to 1.46)	0.901

ORs are adjusted for all other variables displayed in the same table column, and statistically significant ORs ($P < 0.05$) are bolded.

*Constant $B = -8.80$.

†Constant $B = -6.27$.

‡Constant $B = -12.41$.

§Constant $B = -9.27$.

||Constant $B = -8.81$.

AOR, adjusted odds ratio.




FIGURE 1. Predicted incidence of incautious sex among adolescent boys ($n = 1170$). All analyses controlled for the following variables at time 1: incautious sex, age, food insecurity, informal housing, urban location, school nonenrollment, migration, number of children in the home, female primary caregiver, maternal and paternal orphanhood, and having a birth certificate. We also controlled for HIV knowledge, measured at time 2. Error bars are 95% CIs.

monitoring and school feeding, 1.0%; and with free school and school feeding, 0.9%. Among girls who received all 3 interventions, pregnancy incidence was less than 0.5%.

DISCUSSION

These findings have 2 key messages for adolescent HIV prevention in South Africa, regarding the impacts of domains of social protection and combinations of types of social protection.

First, they show that specific social protection interventions in 3 domains, cash, psychosocial support, and education (or *cash, care, and classroom*), independently reduce specific HIV-risk behaviors among adolescent boys and girls. In particular, child-focused grants, parental monitoring, free schooling, school feeding, and teacher support each show significant prevention effects, independently of other social interventions and after controlling for covariates and baseline HIV-risk behavior.

Second, findings demonstrate that *combination social protection* interventions can have strong effects on HIV-risk behavior reduction, independently of sociodemographic cofactors and baseline HIV-risk. For example, past-year incidence rates of incautious sex among males reduced more than 3-fold, from 22% with no identified social protection interventions to 6% with a combination of free school, parental monitoring, and teacher support. The predicted probability of incautious sex among females halved from 15% to 7% with combined social protections of parental monitoring and school feeding, and economic sex among females reduced 5-fold from 10% to 2% with combined child grants and free schooling.

These findings not only demonstrate that combination social protection is likely to be more effective than stand-

alone programs but also show that specific combinations should be selected for effects on particular HIV risks and genders. By getting the right combinations of child-focused cash transfers, free schooling, school feeding, parental monitoring, and teacher support, we have potential to reduce adolescent HIV risks.

The study raises further research questions regarding why and how combination social protections reduce HIV risks. We know that sexual decisions have complex behavioral risk factors: It may be that different types of social protection are addressing simultaneous but different sources of risk—for example, a household cash transfer may reduce the financial need for an adolescent girl to have a “sugar daddy,” whereas access to education through free school may reduce her exposure to older men in the community, and good parenting may provide guidance and reduce emotional needs for affection from a high-risk sexual partner. It may also be that some adolescents are particularly vulnerable to 1 type of risk but that not all adolescents are vulnerable to the same type of risk and that combination social protection is necessary to have an impact at a population level in endemic countries. Theoretical models of child and adolescent development and research on risks such as youth offending³⁰ suggest that single interventions may not be powerful enough alone to impact adolescent risk-taking behavior. Thus, it is also possible that multicomponent approaches, providing a compound effect, are required to counter the high-risk environments that characterize these adolescents’ lives. Future research—both qualitative and quantitative—will be of great value in identifying these pathways.

This study has a number of limitations and strengths that should be considered. First, randomized controlled trial




FIGURE 2. Predicted incidence of HIV-risk behaviors among adolescent girls ($n = 1498$): for incautious sex (A), economically driven sex (B), and pregnancy (C). All analyses controlled for the following variables at time 1: sexual risk behavior, age, food insecurity, informal housing, urban location, school nonenrollment, migration, number of children in the home, female primary caregiver, maternal and paternal orphanhood, and having a birth certificate. We also controlled for HIV knowledge, measured at time 2. Analysis of pregnancy as an outcome controlled for receipt of a child support or foster child grant at times 1 and 2. Error bars are 95% CIs.

designs are more reliable in determining causality. However, quasiexperimental panel studies allow simultaneous testing of multiple possible combinations of interventions and can refine these into a smaller number of combinations for future testing in randomized trials. Second, the study only has 2 time points, a year apart. It will be important to examine longer-term associations of social protection and how they may influence HIV-risk behavior as adolescents progress into young adulthood.³¹ However, longitudinal studies have the important advantage of allowing analyses to control for baseline HIV-risk behaviors. This makes it possible to analyze incident HIV-risk behaviors, which allows for stronger causal assumptions. Third, it would also be valuable to examine combination social protection in other countries and in population subgroups not sampled in this study, particularly street youth, and adolescents in prisons and in residential institutions.³ However, this study used a systematic random sampling method, and by follow-up reached the original adolescents across 7 provinces within South Africa. Fourth, although all behavioral outcomes in this study have strong associations with HIV infection among Southern African adolescents, it would be valuable for future studies to use HIV-biomarker endpoints, which is becoming more feasible with accessibility of new assays that can detect time of infection. Finally, this study was restricted to social protection interventions. Future research should examine social protection in combination with biomedical and behavioral interventions, to determine whether cross-sectoral combinations can provide even greater HIV prevention impacts.

This study therefore suggests clear needs for future research. Randomized controlled trials with biomarker HIV incidence outcomes and long-term follow-up would be of great value. Testing of moderator and mediator effects in quasiexperimental and experimental data, combined with qualitative research, could help identify causal pathways by which combination social protection may be impacting HIV risks, and the next steps may be examining potential interactions of social protection with biomedical interventions—such as pre-exposure prophylaxis and antiretroviral therapy for positive adolescents.

This study has one further, and important, advantage. It uses existing interventions that are currently provided by states, nongovernmental organizations, or families at a large scale. It also demonstrates that social protection combinations remain strongly associated with reductions of HIV-risk behavior independently of a range of predictors of HIV and service access. As a result, the findings show not only that combination social protection is effective in adolescent HIV-risk reduction but also that it is demonstrably feasible and scalable in real-world African contexts.

ACKNOWLEDGMENTS

The authors thank Sudhanshu Handa, Patricia Lim Ah Ken, Natalia Winder-Rossi, Jason Wolfe, and Tom Fenn for discussion and input. The authors thank their fieldwork teams and all participants and their families.

REFERENCES

1. UNICEF. *Towards an AIDS-Free Generation—Children and AIDS: Sixth Stocktaking Report*. New York, NY: UNICEF, 2013.
2. Watts C, Seeley J. Addressing gender inequality and intimate partner violence as critical barriers to an effective HIV response in sub-Saharan Africa. *J Int AIDS Soc*. 2014;17:19849.
3. UNDP Discussion paper: *Cash Transfers and HIV-Prevention*. New York, NY: UN Development Program, 2014.
4. Duflo E. Grandmothers and granddaughters: old-age pensions and intrahousehold allocation in South Africa. *World Bank Econ Rev*. 2003;17:1–25.
5. Robertson L, Mushati P, Eaton JW, et al. Effects of unconditional and conditional cash transfers on child health and development in Zimbabwe: a cluster-randomised trial. *Lancet*. 2013;381:1283–1292.
6. Baird SJ, Garfein RS, McIntosh CT, et al. Effect of a cash transfer program for schooling on prevalence of HIV and herpes simplex type 2 in Malawi: a cluster randomised trial. *Lancet*. 2012;379:1320–1329.
7. Department of Social Development, SASSA, UNICEF. *The South African Child Support Grant Impact Assessment: Evidence from a Survey of Children, Adolescents and Their Households*. Pretoria, South Africa: UNICEF South Africa, 2012.
8. Handa S, Halpern CT, Pettifor A, et al. The government of Kenya's cash transfer program reduces the risk of sexual debut among young people age 15–25. *PLoS One*. 2014;9:e85473.
9. Cluver L, Boyes M, Orkin M, et al. Child-focused state cash transfers and adolescent risk of HIV infection in South Africa: a propensity score-matched case-control study. *Lancet Glob Health*. 2013;1:e362–70.
10. Cluver L, Orkin M, Boyes M, et al. Cash plus care: social protection cumulatively mitigates HIV-risk behavior among adolescents in South Africa. *AIDS*. 2014;28(suppl 3):S389–S397.
11. Cluver L, Orkin M, Boyes M, et al. Transactional sex amongst AIDS-orphaned and AIDS-affected adolescents predicted by abuse and extreme poverty. *J Acquir Immune Defic Syndr*. 2011;53:336–343.
12. Adato M, Bassett L. *What is the Potential of Cash Transfers to Strengthen Families Affected by HIV and AIDS? A Review of the Evidence on Impacts and Key Policy Debates*. Washington, DC: International Food Policy Research Institute, 2008.
13. Gillespie S, Kadiyala S, Greener R. Is poverty or wealth driving HIV transmission? *AIDS*. 2007;21(suppl 7):S5–S16.
14. Piot P, Greener R, Russell S. Squaring the circle: AIDS, poverty, and human development. *PLoS Med*. 2007;4:1571–1575.
15. Robinson J, Yeh E. Transactional sex as a response to risk in Western Kenya. *Am Econ J Appl Econ*. 2011;3:35–64.
16. Dinkelmann T, Lam D, Leibbrandt M. Linking poverty and income shocks to risky sexual behavior: evidence from a panel study of young adults in Cape Town. *S Afr J Econ*. 2008;76(suppl 1):S52–S74.
17. Watts CH, Foss AM, Hossain M, et al. Sexual violence and conflict in Africa: prevalence and potential impact on HIV incidence. *Sex Transm Infect*. 2010;86(suppl 3):iii93–99.
18. UNAIDS. *UNAIDS Report on the Global Epidemic 2013*. Geneva, Switzerland: UNAIDS, 2013.
19. Hillis SD, Anda RF, Felitti VJ, et al. Adverse childhood experiences and sexually transmitted diseases in men and women: a retrospective study. *Pediatrics*. 2000;106:E11.
20. UNAIDS. *A New Investment Framework for the Global HIV Response*. Geneva, Switzerland: UN, 2011.
21. Curtis NM, Ronan KR, Bordin CM. Multisystemic treatment: a meta-analysis of outcome studies. *J Fam Psychol*. 2004; 18:411–419.
22. Rivera JA, Sotres-Alvarez D, Habicht JP, et al. Impact of the Mexican program for education, health, and nutrition (Progresá) on rates of growth and anemia in infants and young children: a randomized effectiveness study. *JAMA*. 2004;291:2563–2570.
23. Pettifor A, Rees HV, Kleinschmidt I, et al. Young people's sexual health in South Africa: HIV prevalence and sexual behaviors from a nationally representative household survey. *AIDS*. 2003;19:1525–1534.
24. Shisana O. *South African National HIV Prevalence, HIV Incidence, Behavior and Communication Survey, 2005*. Cape Town, South Africa: HSRC Press, 2005.
25. Christofides NJ, Jewkes R, Dunkle K, et al. Early adolescent pregnancy increases risk of incident HIV infection in the Eastern Cape, South Africa: a longitudinal study. *J Int AIDS Soc*. 2014;17:18585.

26. Van der Merwe A, Dawes A. Prosocial and antisocial tendencies in children exposed to community violence. *J Child Adolesc Ment Health*. 2000;12:19–37.
27. Elgar F, Waschbusch D, Dadds M, et al. Development and validation of a short form of the Alabama parenting questionnaire. *J Child Fam Stud*. 2007;16:243–259.
28. Bailly D. Issues related to consent to healthcare decisions in children and adolescents. *Arch Pediatr*. 2010;17:S7–S15.
29. Hosmer DW Jr, Lemeshow S, Sturdivant R. *Applied Logistic Regression*, ed. New Jersey, NJ: John Wiley, Sons, 2013.
30. Farrington D, Welsh B. *Saving Children from a Life of Crime: Early Risk Factors and Effective Interventions*. Oxford, United Kingdom: Oxford University Press, 2007.
31. Heise L, Lutz B, Ranganathan M, et al. Cash transfers for HIV prevention: considering their potential. *J Int AIDS Soc*. 2013;16:18615.