Homogeneous Catalysis

Synthesis of All-Carbon Disubstituted Bicyclo[1.1.1]pentanes by Iron-Catalyzed Kumada Cross-Coupling**

Jeremy Nugent[†], Bethany R. Shire[†], Dimitri F. J. Caputo, Helena D. Pickford, Frank Nightingale, Ian T. T. Houlsby, James J. Mousseau, and Edward A. Anderson*

Abstract: 1,3-Disubstituted bicyclo[1.1.1]pentanes (BCPs) are important motifs in drug design as surrogates for *p*-substituted arenes and alkynes. Access to all-carbon disubstituted BCPs via cross-coupling has to date been limited to use of the BCP as the organometallic component, which restricts scope due to the harsh conditions typically required for the synthesis of metallated BCPs. Here we report a general method to access 1,3-*C*-disubstituted BCPs from 1-iodo-bicyclo[1.1.1]pentanes (iodo-BCPs) by direct iron-catalyzed cross-coupling with aryl and heteroaryl Grignard reagents. This chemistry represents the first general use of iodo-BCPs as electrophiles in cross-coupling, and the first Kumada coupling of tertiary iodides. Benefiting from short reaction times, mild conditions, and broad scope of the coupling partners, it enables the synthesis of a wide range of 1,3-*C*-disubstituted BCPs including various drug analogues.

1,3-*D*isubstituted bicyclo[1.1.1]pentanes (BCPs) are of high interest in drug discovery as bioisosteres for 1,4-disubstituted arenes and alkynes (Figure 1 a).^[1] Incorporation of these sp³-rich motifs into drug leads often results in pharmacological benefits such as improved solubility, membrane permeability and metabolic stability.^[2] However, access to promising BCP-bearing compounds^[2b,3] can be impeded by lengthy and unscalable reaction sequences, in particular where two carbon substituents are required. These challenges have inspired the development of a number of methods to

synthesize 1,3-*C*-disubstituted BCPs in which installation of the carbon substituents relies on addition of an organometallic nucleophile to the strained C1–C3 σ -bond of [1.1.1]propellane,^[1e,4] followed by palladium-catalyzed cross-coupling of the resulting metallated BCP (Figure 1 b).^[1e,4a,5] While such methods can generate useful products, the harsh conditions required to achieve the initial nucleophilic addition limit the suitability of this chemistry for industrial applications.

1-Iodobicyclo[1.1.1]pentanes (iodo-BCPs, **1**) are attractive substrates for the introduction of carbon substituents on the BCP skeleton. We recently described efficient and functional group-tolerant conditions to access these compounds by atom transfer radical addition of C–I bonds to [1.1.1]propellane, under photoredox catalysis^[6] or using triethylborane as initiator.^[7] As direct palladium-catalyzed cross-coupling of iodo-BCPs can suffer from competing ring fragmentation,^[8] approaches to all-carbon disubstituted BCPs from iodo-BCPs have to date necessitated lithiation of the iodide, followed by cross-coupling as the nucleophilic com-

a: Bicyclo[1.1.1]pentanes in drug discovery

b: Known cross-coupling of metallated BCPs to access 1,3-*C*-disubstituted BCPs

BCP as nucleophile:
Harsh conditions / cryogenic temperatures, not suitable for scale

c: This work: Direct iron-catalyzed Kumada cross-coupling of BCP iodides

BCP as electrophile:
Mild conditions, short reaction times, functional group tolerant

Figure 1. a) Examples of bicyclo[1.1.1]pentanes (BCPs) in medicinal chemistry; b) known cross-coupling of metallated BCPs; c) this work: Direct iron-catalyzed cross-coupling of iodo-BCPs with aryl/ heteroaryl Grignard reagents.

[*] Dr. J. Nugent,^[†] B. R. Shire,^[†] D. F. J. Caputo, H. D. Pickford, F. Nightingale, Prof. E. A. Anderson
Chemistry Research Laboratory, University of Oxford
12 Mansfield Road, Oxford OX1 3TA (UK)
E-mail: edward.anderson@chem.ox.ac.uk

Dr. I. T. T. Houlsby
Syngenta Ltd., Jealott's Hill International Research Centre
Bracknell RG42 6EY (UK)

Dr. J. J. Mousseau
Pfizer Medicine Design
Eastern Point Road, Groton, CT 06340 (USA)

[†] These authors contributed equally to this work.

[**] A previous version of this manuscript has been deposited on a preprint server (<https://doi.org/10.26434/chemrxiv.12009390.v1>).

Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under:
<https://doi.org/10.1002/anie.202004090>.

© 2020 The Authors. Published by Wiley-VCH Verlag GmbH & Co. KGaA. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

ponent (Figure 1b, or reaction with other carbon-based electrophiles).^[1e,4a] However, the conditions needed for lithiation of the iodo-BCP again limit scope and scalability.

We targeted an alternative, catalytic method to access all-carbon disubstituted BCPs (**2**, Figure 1c) directly from iodo-BCPs **1**, under mild conditions and without recourse to organolithium reagents. Iron-catalyzed Kumada cross-couplings of aryl Grignard reagents with secondary alkyl iodides are an efficient means to achieve sp^3 – sp^2 C–C bond formation,^[9] and we questioned whether the tertiary iodide resident in an iodo-BCP could engage in this coupling manifold. While isolated examples of Fe-catalyzed Kumada couplings of tertiary alkyl bromides and chlorides have been described,^[9d,10] the equivalent reaction of tertiary iodides is, to our knowledge, unknown. Here we describe the development of iron-catalyzed cross-coupling reactions of iodo-BCPs with both aryl and heteroaryl Grignard reagents, which represents the first general procedure for the direct cross-coupling of iodo-BCP electrophiles.^[11] The chemistry proceeds under mild conditions and short reaction times, displays wide functional group tolerance, and is applicable to the synthesis of drug-like molecules.

Our studies began with the coupling of iodo-BCP **1a** with *p*-methoxyphenylmagnesium bromide (1.6 equiv, 0.7 mL h^{−1}), which afforded small amounts of coupled product **2a** using Fe(acac)₃ or FeCl₃ (20 mol %) as catalyst (Table 1, Entries 1, 2); the main byproduct was the dehalogenated BCP **3**. Notably, no reaction was seen using Cu(acac)₂ (Entry 3). A number of amine and phosphorus ligands were investigated (Entries 4–9),^[12] among which TMEDA^[9a,b,13] (40 mol %) provided the highest yields of **2a** using either a standard Grignard reagent (PMPBr/Mg, 86 %, Entry 8), or the turbo-Grignard (PMPBr/Mg/LiCl, 79 %, Entry 9). THF was found to be a superior solvent compared to others such as MTBE, 2-MeTHF, and toluene (Entries 10–12).^[9g,14] In spite of the failure of Cu(acac)₂ alone to catalyze the reaction, we were mindful of the potential presence of trace metals in the (97 % purity) Fe(acac)₃.^[15] However, equivalent reaction efficiency was observed using ≥ 99.9 % Fe(acac)₃ (Entry 13), and no reaction with Cu(acac)₂/TMEDA (Entry 14), suggesting this to indeed be an iron-catalyzed transformation. Reducing the loading of catalyst to 10 mol % led to lower yields (Entry 15).

With optimized conditions in hand (Table 1, Entries 8/9), the scope of the aryl Grignard coupling partner was examined (Figure 2) using *p*-trifluoromethylbenzyl (**1a**) and 4-*N*-Boc-piperidyl (**1b**) iodo-BCPs as representative substrates. Both electron-rich and moderately electron-poor Grignard reagents gave good to excellent yields of the coupled products for *para*-substituted aryl organometallics (**2b–2m**, 53–83 %), with the reaction of 4-trimethylsilylphenylmagnesium bromide also delivering an excellent yield of **2m** on 2.7 mmol scale (90 %). Substitution at the *meta* and *ortho* positions was well tolerated (**2n–2r**, 55–86 %), although in the latter cases elevated temperatures (45 °C) were required. Bicyclic and trisubstituted Grignard reagents gave similarly high yields of BCP products **2s–2u** (70–72 %).^[16]

The direct coupling of heteroaromatic Grignard reagents would be of high interest in a pharmaceutical context. To our delight, a variety of heteroaryl Grignards underwent success-

Table 1: Reaction optimization.^[a]

Entry	Catalyst (20 mol %)	Additive (40 mol %)	Solvent	Yield ^[b] (2a : 1a : 3)
1	Fe(acac) ₃	–	THF	13:30:9
2	FeCl ₃	–	THF	4:23:9
3	Cu(acac) ₂	–	THF	0:100:0
4	Fe(acac) ₃	DMEDA	THF	5:22:9
5	Fe(acac) ₃	TMCD	THF	9:76:1
6	Fe(acac) ₃	1,2-DPE	THF	46:33:8
7	Fe(acac) ₃	dcypt	THF	55:20:11
8	Fe(acac) ₃	TMEDA	THF	90:0:2 (86%)
9 ^[c]	Fe(acac) ₃	TMEDA	THF	86:0:4 (79%) (86%)
10	Fe(acac) ₃	TMEDA	2-MeTHF	71:15:2
11	Fe(acac) ₃	TMEDA	MTBE	70:16:2
12	Fe(acac) ₃	TMEDA	toluene	79:0:3
13 ^[d]	Fe(acac) ₃	TMEDA	THF	92:0:2
14	Cu(acac) ₂	TMEDA	THF	0:100:0
15 ^[e]	Fe(acac) ₃	TMEDA	THF	75:7:3

[a] **1a** (0.2 mmol), catalyst (20 mol %), additive (40 mol %), solvent (0.2 mL), 20 °C; then add PMPMgBr (0.8 M in THF, 0.32 mmol) added at 0.7 mL h^{−1}; then stir, 20 °C, 1 h. [b] Yields determined by ¹H NMR spectroscopic analysis using 1,3,5-trimethoxybenzene as an internal standard. Isolated yield in parentheses. [c] Using PMPMgBr·LiCl (1 M in THF). [d] Using ≥ 99.9 % Fe(acac)₃. [e] Using 10 mol % Fe(acac)₃, 20 mol % TMEDA. acac = acetoacetate. DMEDA = *N,N'*-dimethylethylenediamine. dcypt = 3,4-bis(dicyclohexylphosphino)thiophene. 1,2-DPE = 1,2-dipiperidinoethane. MTBE = methyl *tert*-butyl ether. PMP = 4-methoxyphenyl. TMCD = *N,N,N',N'*-trans-tetramethylcyclohexanedi-amine. TMEDA = *N,N,N',N'*-tetramethylethylenediamine.

ful reaction to give heteroaryl-functionalized BCPs **2v–2ad**, albeit warming was required for electron-deficient heterocycles (45 °C). The unsubstituted 3-pyridyl Grignard reagent gave **2v** in a modest 27 % yield, however significant improvement was observed with more electron-rich pyridines (**2w–2y**, 35–55 %), although reactions of 2-pyridyl Grignard reagents were unsuccessful. Cross-couplings of benzofuran, *N*-Boc-indoline, *N*-Boc-indole, *N*-methyl indazole, and *N*-phenyl carbazole organometallics were successful, giving good yields of the heteroarylated products **2z–2ad** (46–67 %).

The scope of the iodo-BCP coupling partner was next investigated in couplings with PMPMgBr. An *N*-Boc azetidine-substituted BCP gave product **2ae** in excellent yield (73 %). Electron-withdrawing groups were well-tolerated, such as sulfone **2af**, and the electrophilic ester **2ag** (67 % and 95 % respectively); in the latter case, no addition of the Grignard to the ester was observed. However, a more electrophilic α,α -difluoro ester proved less successful (**2ah**), likely due to competing addition to the carbonyl. Heteroaryl-substituted iodo-BCPs also proved good substrates, affording the bis-arylated BCP **2ai** in 69 % yield, and double cross-coupled product **2aj** in 64 % yield. The chemistry was applied to more complex iodides, with quinoline **2ak** and nicotinic

Figure 2. Cross-coupling of electron rich, electron neutral, aryl and heteroaryl Grignard reagents with iodo-BCPs. a) Reaction performed at 45 °C. b) 1.0 g scale. c) Reaction performed at 65 °C with rapid addition of the Grignard reagent. d) Reaction performed at 45 °C with 3 equiv of Grignard reagent. e) Using (Het)ArMgBr·LiCl.

acid derivative **2al** being formed in 83% and 40% yields, respectively. These examples emphasize the mild conditions and functional group tolerance of this methodology.

The mechanistic pathways of iron-catalyzed cross-coupling reactions are dependent on a number of factors, including the nature of the Grignard reagent, the rate of its addition, the additive (ligand), and solvent.^[9i,17] The formation of iron nanoparticles may also be observed, in particular under a “rapid addition” (of Grignard) regime.^[18] In our system, a distinctive colour change from orange/red to dark/black was observed during Grignard addition which may indicate nanoparticle formation,^[19] albeit this is not commonly observed under “slow” addition regimes.^[13a] The presence of substoichiometric TMEDA is also clearly beneficial to reaction efficiency,^[9j,20] although its role is unclear given evidence that it may not be ligated to the metal during the coupling process.^[21]

In keeping with couplings carried out using less sterically-hindered aryl Grignard reagents with amine additives, we therefore favour a reaction pathway involving single electron transfer from an Fe^I species such as **4**^[17c,22] to the iodo-BCP (Scheme 1), generating a bicyclopentyl radical **5**.^[23] Reaction of this species with the LArFe(II) complex/ArMgBr liberates the cross-coupled product and results in catalyst turnover;

Scheme 1. Putative reaction mechanism.

however, the precise mechanism by which C–C bond formation occurs (e.g. in cage/out of cage) is not apparent.^[17a,24]

When combined with our previous methods for iodo-BCP synthesis, the Kumada cross-coupling offers a powerful method for the mild and rapid generation of valuable, pharmaceutically-relevant 1,3-C-difunctionalized BCPs. To demonstrate potential utility, we targeted BCP analogues of the anti-inflammatory drug flurbiprofen, and the anti-neoplastic agent brequinar (Scheme 2a). The requisite iodo-

Scheme 2. a) Synthesis of BCP drug analogues. b) Further functionalization. Reagents and Conditions: a) [1.1.1]propellane, Et_3B (10 mol %), Et_2O , 0°C , 15 min; b) $\text{Fe}(\text{acac})_3$ (20 mol %), TMEDA (40 mol %), PhMgBr , THF, rt, 1 h, then NaOH/MeOH ; c) $\text{Ir}(\text{ppy})_3$ (2.5 mol %), [1.1.1]propellane, $t\text{-BuCN}$, blue LEDs, 24 h, 30°C ; d) $\text{Fe}(\text{acac})_3$ (20 mol %), TMEDA (40 mol %), 4-FPhMgBr, THF, rt, 1 h; e) KBr , NCS , AcOH/MeOH , rt; f) ICl , CH_2Cl_2 , 0°C ; g) AcCl , AlCl_3 , CH_2Cl_2 , $0^\circ\text{C} \rightarrow \text{rt}$; then Boc_2O , Et_3N , CH_2Cl_2 , rt. ppy = 2-phenylpyridine.

BCPs **1k** and **1i** were synthesized in excellent yields from reaction of [1.1.1]propellane with commercially available ethyl iodopropanoate **6** (80 %), and iodoquinoline **7** (85 %),^[8] using Et_3B initiation and photoredox catalysis ($\text{Ir}(\text{ppy})_3$ /blue LEDs), respectively. Kumada cross-coupling of iodo-BCP ester **1k** with PhMgBr , followed by hydrolysis, furnished BCP-flurbiprofen **8** in 78 % yield; coupling of 2-pyridyl iodo-BCP **1i** with 4-fluorophenylmagnesium bromide afforded brequinar analogue **9** (66 %).^[25] Finally, access to aryl-BCPs featuring substituents not tolerated under Kumada coupling could be achieved by *ipso*-substitution^[26] of aryl silane **2m** (Scheme 2b), for example with halides suitable for further elaboration by cross-coupling (**10**, **11**),^[27] or an electron-withdrawing acetyl group (**12**).^[28]

In conclusion, we have developed a mild, efficient iron-catalyzed cross-coupling of iodo-BCPs and (hetero)aryl Grignard reagents, which represents the first such example of Kumada cross-coupling of tertiary iodides. The reaction is rapid, exhibits good functional group tolerance, and performs well on gram scale.

Applications to the functionalization of pharmaceutical derivatives, including the synthesis of two BCP drug analogues, demonstrate the potential of this transformation to access highly functionalized 1,3-C-disubstituted BCPs of direct relevance in medicinal chemistry settings.

Acknowledgements

J.N. thanks the Marie Skłodowska-Curie actions for an Individual Fellowship (GA No 786683). B.R.S., D.F.J.C. and H.D.P. thank the EPSRC Centre for Doctoral Training in

Synthesis for Biology and Medicine for studentships (EP/L015838/1), generously supported by AstraZeneca, Diamond Light Source, Defence Science and Technology Laboratory, Evotec, GlaxoSmithKline, Janssen, Novartis, Pfizer, Syngenta, Takeda, UCB and Vertex. E.A.A. thanks the EPSRC for support (EP/S013172/1). F.N. thanks the Royal Society of Chemistry for an Undergraduate Research Bursary. We thank Dr Kirsten E. Christensen and Steven J. Mansfield for assistance with refinement of the X-ray crystal structures.

Conflict of interest

The authors declare no conflict of interest.

Keywords: bicyclopentane · bioisosteres · cross-coupling · homogeneous catalysis · iron

- [1] a) Y. P. Auberson, C. Brocklehurst, M. Furegati, T. C. Fessard, G. Koch, A. Decker, L. La Vecchia, E. Briard, *ChemMedChem* **2017**, *12*, 590; b) A. F. Stepan, C. Subramanyam, I. V. Efremov, J. K. Dutra, T. J. O'Sullivan, K. J. DiRico, W. S. McDonald, A. Won, P. H. Dorff, C. E. Nolan, S. L. Becker, L. R. Pustilnik, D. R. Riddell, G. W. Kauffman, B. L. Kormos, L. Zhang, Y. Lu, S. H. Capetta, M. E. Green, K. Karki, E. Sibley, K. P. Atchison, A. J. Hallgren, C. E. Oborski, A. E. Robshaw, B. Sneed, C. J. O'Donnell, *J. Med. Chem.* **2012**, *55*, 3414; c) N. D. Measom, K. D. Down, D. J. Hirst, C. Jamieson, E. S. Manas, V. K. Patel, D. O. Somers, *ACS Med. Chem. Lett.* **2017**, *8*, 43; d) Y. L. Goh, Y. T. Cui, V. Pendharkar, V. A. Adsool, *ACS Med. Chem. Lett.* **2017**, *8*, 516; e) I. S. Makarov, C. E. Brocklehurst, K. Karaghiosoff, G. Koch, P. Knochel, *Angew. Chem. Int. Ed.* **2017**, *56*, 12774; f) G. A. Patani, E. J. LaVoie, *Chem. Rev.* **1996**, *96*, 3147.
- [2] a) N. A. Meanwell, *Chem. Res. Toxicol.* **2016**, *29*, 564; b) For a review, see: P. K. Mykhailiuk, *Org. Biomol. Chem.* **2019**, *17*, 2839; c) G. M. Locke, S. S. R. Bernhard, M. O. Senge, *Chem. Eur. J.* **2019**, *25*, 4590; d) F. Lovering, J. Bikker, C. Humblet, *J. Med. Chem.* **2009**, *52*, 6752.
- [3] a) For recent examples of the synthesis of bicyclo[1.1.1]pentanes, see: X. Zhang, R. T. Smith, C. Le, S. J. McCarver, B. T. Shireman, N. I. Carruthers, D. W. C. MacMillan, *Nature* **2020**, *580*, 220; b) J. H. Kim, A. Ruffoni, Y. S. S. Al-Faiyz, N. S. Sheikh, D. Leonori, *Angew. Chem. Int. Ed.* **2020**, *59*, 8225; *Angew. Chem.* **2020**, *132*, 8302; c) S. Yu, C. Jing, A. Noble, V. K. Aggarwal, *Angew. Chem. Int. Ed.* **2020**, *59*, 3917; d) R. M. Bär, P. J. Gross, M. Nieger, S. Bräse, *Chem. Eur. J.* **2020**, *26*, 4242; e) M. Kondo, J. Kanazawa, T. Ichikawa, T. Shimokawa, Y. Nagashima, K. Miyamoto, M. Uchiyama, *Angew. Chem. Int. Ed.* **2020**, *59*, 1970; f) for recent reviews of BCP synthesis and previous references, see: X. Ma, L. Nhat Pham, *Asian J. Org. Chem.* **2020**, *9*, 8; g) J. Kanazawa, M. Uchiyama, *Synlett* **2019**, *30*, 1.
- [4] a) M. Messner, S. I. Kozhushkov, A. de Meijere, *Eur. J. Org. Chem.* **2000**, 1137; b) N. Trongsiwat, Y. Pu, Y. Nieves-Quinones, R. A. Shelp, M. C. Kozlowski, P. J. Walsh, *Angew. Chem. Int. Ed.* **2019**, *58*, 13416; c) for a rare example of room temperature anionic addition, see: R. A. Shelp, P. J. Walsh, *Angew. Chem. Int. Ed.* **2018**, *57*, 15857.
- [5] J. D. D. Rehm, B. Ziemer, G. Szeimies, *Eur. J. Org. Chem.* **1999**, 2079.
- [6] J. Nugent, C. Arroniz, B. R. Shire, A. J. Sterling, H. D. Pickford, M. L. J. Wong, S. J. Mansfield, D. F. J. Caputo, B. Owen, J. J. Mousseau, F. Duarte, E. A. Anderson, *ACS Catal.* **2019**, *9*, 9568.

- [7] a) D. F. J. Caputo, C. Arroniz, A. B. Dürr, J. J. Mousseau, A. F. Stepan, S. J. Mansfield, E. A. Anderson, *Chem. Sci.* **2018**, *9*, 5295; b) M. L. J. Wong, J. J. Mousseau, S. J. Mansfield, E. A. Anderson, *Org. Lett.* **2019**, *21*, 2408.
- [8] See the Supporting Information for details.
- [9] a) M. Nakamura, K. Matsuo, S. Ito, E. Nakamura, *J. Am. Chem. Soc.* **2004**, *126*, 3686; b) G. Cahiez, V. Habiak, C. Duplais, A. Moyeux, *Angew. Chem. Int. Ed.* **2007**, *46*, 4364; c) R. R. Chowdhury, A. K. Crane, C. Fowler, P. Kwong, C. M. Kozak, *Chem. Commun.* **2008**, 94; d) T. Hatakeyama, Y.-i. Fujiwara, Y. Okada, T. Itoh, T. Hashimoto, S. Kawamura, K. Ogata, H. Takaya, M. Nakamura, *Chem. Lett.* **2011**, *40*, 1030; e) X. Lin, F. Zheng, F.-L. Qing, *Organometallics* **2012**, *31*, 1578; f) G. Bauer, C. W. Cheung, X. Hu, *Synthesis* **2015**, *47*, 1726; g) D. Parmar, L. Henkel, J. Dib, M. Rueping, *Chem. Commun.* **2015**, *51*, 2111; h) T. Hashimoto, T. Maruyama, T. Yamaguchi, Y. Matsubara, Y. Yamaguchi, *Adv. Syn. Catal.* **2019**, *361*, 4232; for recent reviews, see: i) T. L. Mako, J. A. Byers, *Inorg. Chem. Front.* **2016**, *3*, 766; j) A. Piontek, E. Bisz, M. Szostak, *Angew. Chem. Int. Ed.* **2018**, *57*, 11116; k) B. Barré, L. Gonnard, R. Campagne, S. Reymond, J. Marin, P. Ciapetti, M. Brellier, A. Guérinot, J. Cossy, *Org. Lett.* **2014**, *16*, 6160.
- [10] To our knowledge, the only successful couplings of tertiary bromides and chlorides use adamantyl halides. See Ref. [9d], and S. K. Ghorai, M. Jin, T. Hatakeyama, M. Nakamura, *Org. Lett.* **2012**, *14*, 1066.
- [11] To the best of our knowledge, only a single example of cross-coupling of an iodo-BCP has been described, which involves Cu-catalyzed Sonagashira coupling. See: A. Hazra, M. T. Lee, J. F. Chiu, G. Lalic, *Angew. Chem. Int. Ed.* **2018**, *57*, 5492.
- [12] Entries 4–9 in Table 1 show the optimum results from the ligand screen; see the Supporting Information for full details. TMEDA and TMCD have been used as ligands in related Fe-catalyzed Kumada coupling; see Refs [9a,b] and [9k].
- [13] a) R. B. Bedford, P. B. Brenner, E. Carter, P. M. Cogswell, M. F. Haddow, J. N. Harvey, D. M. Murphy, J. Nunn, C. H. Woodall, *Angew. Chem. Int. Ed.* **2014**, *53*, 1804; b) R. B. Bedford, D. W. Bruce, R. M. Frost, M. Hird, *Chem. Commun.* **2005**, 4161; c) D. Noda, Y. Sunada, T. Hatakeyama, M. Nakamura, H. Nagashima, *J. Am. Chem. Soc.* **2009**, *131*, 6078.
- [14] The concentration, equivalents, and rate of addition of the Grignard reagent also proved important factors in reaction optimization; for full details see the Supporting Information.
- [15] a) S. L. Buchwald, C. Bolm, *Angew. Chem. Int. Ed.* **2009**, *48*, 5586; b) I. Thomé, A. Nijs, C. Bolm, *Chem. Soc. Rev.* **2012**, *41*, 979.
- [16] See the Supporting Information for details of unsuccessful substrates.
- [17] a) A. Hedström, Z. Izakian, I. Vreto, C.-J. Wallentin, P.-O. Norrby, *Chem. Eur. J.* **2015**, *21*, 5946; b) M. L. Neidig, S. H. Carpenter, D. J. Curran, J. C. DeMuth, V. E. Fleischauer, T. E. Iannuzzi, P. G. N. Neate, J. D. Sears, N. J. Wolford, *Acc. Chem. Res.* **2019**, *52*, 140; c) R. B. Bedford, *Acc. Chem. Res.* **2015**, *48*, 1485.
- [18] R. B. Bedford, M. Betham, D. W. Bruce, S. A. Davis, R. M. Frost, M. Hird, *Chem. Commun.* **2006**, 1398.
- [19] For further details and a discussion of mechanistic observations, see the Supporting Information.
- [20] In contrast to the findings of the Nakamura and Caihez groups (Refs [9a,b]) no benefit was observed using superstoichiometric TMEDA (78% yield was obtained using 160 mol % TMEDA). Equally, in contrast to the findings of Bedford et al., (Ref. [13a]), reaction efficiency was severely compromised in the absence of TMEDA.
- [21] R. B. Bedford, P. B. Brenner, D. Elorriaga, J. N. Harvey, J. Nunn, *Dalton Trans.* **2016**, *45*, 15811.
- [22] a) G. Lefèvre, A. Jutand, *Chem. Eur. J.* **2014**, *20*, 4796; b) S. H. Carpenter, T. M. Baker, S. B. Muñoz, W. W. Brennessel, M. L. Neidig, *Chem. Sci.* **2018**, *9*, 7931.
- [23] A. J. Sterling, A. Dürr, R. Smith, E. A. Anderson, F. Duarte, *Chem. Sci.* **2020**, <https://doi.org/10.1039/D0SC01386B>.
- [24] A. Hedström, U. Bollmann, J. Bravidor, P.-O. Norrby, *Chem. Eur. J.* **2011**, *17*, 11991.
- [25] Low-temperature single-crystal X-ray diffraction data for **2a** and **9** were collected with a (Rigaku) Oxford Diffraction SuperNova A diffractometer at 150 K. All data were reduced using CrysAlisPro, solved using SuperFlip [L. Palatinus, G. Chapuis, *J. Appl. Crystallogr.* **2007**, *40*, 786–790] and the structures were refined using CRYSTALS. [P. W. Betteridge, J. R. Carruthers, R. I. Cooper, K. Prout, D. J. Watkin, *J. Appl. Crystallogr.* **2003**, *36*, 1487; R. I. Cooper, A. L. Thompson, D. J. Watkin, *J. Appl. Crystallogr.* **2010**, *43*, 1100–1107]; see the Supporting Information (CIF) for further details. CCDC 1991572, and 1991573 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre.
- [26] a) T. H. Chan, I. Fleming, *Synthesis* **1979**, 761; b) C. Eaborn, *J. Organomet. Chem.* **1975**, *100*, 43.
- [27] X.-H. Gu, R. Zong, N. S. Kula, R. J. Baldessarini, J. L. Neumeyer, *Bioorg. Med. Chem. Lett.* **2001**, *11*, 3049.
- [28] a) C. D. Campbell, R. L. Greenaway, O. T. Holton, H. A. Chapman, E. A. Anderson, *Chem. Commun.* **2014**, *50*, 5187; b) G. Felix, M. Laguerre, J. Dunogues, R. Calas, *J. Org. Chem.* **1982**, *47*, 1423.

Manuscript received: March 19, 2020

Accepted manuscript online: April 28, 2020

Version of record online: ■■■■■, ■■■■■

Communications

Homogeneous Catalysis

J. Nugent, B. R. Shire, D. F. J. Caputo,
H. D. Pickford, F. Nightingale,
I. T. T. Houlsby, J. J. Mousseau,
E. A. Anderson* ———— ■■■■-■■■■

Synthesis of All-Carbon Disubstituted
Bicyclo[1.1.1]pentanes by Iron-Catalyzed
Kumada Cross-Coupling

1,3-C-disubstituted bicyclo-[1.1.1]pentanes (BCPs) are accessed from 1-iodo-BCPs by direct iron-catalyzed cross-coupling with aryl and heteroaryl Grignard reagents. This represents the first general use of iodo-BCPs as cross-coupling electrophiles, and the first Kumada coupling of tertiary iodides in general. Benefiting from short reaction times, mild conditions, and broad scope, it enables the synthesis of a wide range of BCPs including drug analogues.