

Determinants of influenza and pertussis vaccination uptake in pregnancy: a multi-centre questionnaire study of pregnant women and healthcare professionals

Christopher R Wilcox MBBCh ¹, Anna Calvert MRCPh ², Jane Metz MBBS ³, Eliz Kilich BM BCh ², Rachael MacLeod MBChB ³, Kirsten Beadon BSc ⁴, Paul T Heath MRCPh ², Asma Khalil MD ^{5,6}, Adam Finn PhD ³, Matthew D Snape MD ⁴, Tushna Vandrevalla PhD ⁷, Tom Nadarzynski PhD ⁸, Matthew A Coleman MD ⁹, Christine E Jones PhD ¹⁰

1: NIHR Clinical Research Facility, University Hospital Southampton NHS Foundation Trust, Southampton, UK

2: Vaccine Institute, St George's, University of London, London, UK

3: Bristol Children's Vaccine Centre, University of Bristol, Bristol, UK

4: Oxford Vaccine Group, Department of Paediatrics, University of Oxford and the NIHR Oxford Biomedical Research Centre, Oxford, UK

5: Fetal Medicine Unit, Department of Obstetrics and Gynaecology, St George's University Hospitals NHS Foundation Trust, London, UK

6: Vascular Biology Research Centre, Molecular and Clinical Sciences Research Institute, St George's, University of London, London, UK

7: School of Social and Behavioural Sciences, Faculty of Business & Social Sciences, Kingston University, Kingston, London, UK

8: Department of Psychology, University of Westminster, London, UK

9: Department of Fetal Medicine, Princess Anne Hospital, University Hospital Southampton NHS Foundation Trust, Southampton, UK

10: Faculty of Medicine and Institute for Life Sciences, University of Southampton and University Hospital Southampton NHS Foundation Trust, Southampton, UK

Abbreviated title

Improving uptake of maternal vaccination: questionnaire study

Running title

Questionnaire study of maternal vaccination

Corresponding author

- Dr Christopher Wilcox
- NIHR Clinical Research Facility, Southampton Centre for Biomedical Research, C Level West Wing, Mailpoint 218, Southampton General Hospital, Tremona Road, Southampton, SO16 6DY
- Email: christopher.wilcox@soton.ac.uk
- Telephone: 02381204956

Keywords - Vaccination; Pregnancy; Influenza; Pertussis; Vaccine confidence

Abstract

Introduction

Uptake rates of antenatal vaccination remain suboptimal. Our aims were to determine: (1) the acceptability of routine vaccination among pregnant women, (2) the confidence of maternity healthcare professionals (HCPs) discussing vaccination and (3) HCP opinion regarding the optimum healthcare site for vaccine administration.

Methods

Separate questionnaires for pregnant women and HCPs were distributed within four NHS trusts in South England (July 2017-January 2018).

Results

Responses from 314 pregnant women and 204 HCPs (18% obstetricians, 75% midwives, 7% unidentified) were analysed. Previous/intended uptake of influenza and pertussis vaccination was 78% and 92%, respectively. The commonest reason for declining vaccination was feared side-effects for their child. White British women (79%) were significantly more accepting of influenza (85% vs. 61%, OR 3.25, 95% CI: 1.67-6.32) and pertussis vaccination (96% vs. 83%, OR 4.83, 95% CI: 1.77-13.19) compared with non-white-British women. Among HCPs, 25% were slightly or not-at-all confident discussing vaccination. Obstetricians felt significantly more confident discussing pertussis vaccination than midwives (68% vs. 55% were very/moderately confident, OR 2.05, 95% CI: 1.02-4.12). Among HCPs, 53%, 25% and 16% thought vaccines should be administered in primary care (general practice), community midwifery and in hospital, respectively.

Conclusion

Misconceptions exist regarding safety/efficacy of antenatal vaccination, and framing information towards the child's safety may increase uptake. Education of HCPs is essential, and vaccine promotion should be incorporated into routine antenatal care, with an emphasis on women from ethnic minorities. Administration of vaccines in primary care presents logistical barriers however support for alternative sites appears low among HCPs.

Introduction

Both influenza and pertussis result in severe outcomes for pregnant women and their infants (including respiratory illness and death)^{1 2}, and vaccination in pregnancy is an effective means of protection until the period of greatest susceptibility has passed³⁻⁶. In the UK, influenza and pertussis vaccination have been routinely recommended for use in pregnancy since 2010 and 2012, respectively⁷.

Unfortunately, achieving vaccine acceptance among pregnant women and healthcare professionals (HCPs) remains a global challenge⁸. The World Health Organization (WHO) Strategic Advisory Group of Experts on Immunization have called for improved monitoring of vaccine acceptance, and research into the socio-economic determinants of attitudes towards vaccines⁹. The uptake of influenza and pertussis vaccination during pregnancy in England over the September 2016 - January 2017 period was 44.9% and 74.2%, respectively^{10 11}. Pertussis vaccination uptake in the UK has gradually climbed from around 50% since its introduction in 2012, yet influenza vaccine uptake has been relatively static, and remains well below the WHO target of 75%¹⁰. Furthermore, coverage varies significantly between different regions of the UK, with average uptake approximately 10% and 20% lower in London than in northern England for influenza and pertussis, respectively^{9 11}.

Uptake of vaccination could be significantly improved if we are able to fully understand the decision-making processes to acceptance. Furthermore, it is well-acknowledged that encouragement from a familiar HCP significantly improves vaccine acceptance^{12 13}, yet few studies have considered the extent to which HCPs feel confident discussing vaccinations with pregnant women, and the associated factors which might influence this. Optimizing the healthcare site of vaccine administration is also an important issue that may have a considerable impact on vaccine uptake, yet few studies have considered the support of HCPs for alternative approaches. In the UK, vaccination is free-of-charge, and is usually provided within primary care (general practice), and is less commonly available within secondary (hospital-based) care. This may present a logistical barrier if it requires women to arrange extra appointments, and more convenient approach might be to routinely administer vaccination at the time of antenatal appointments.

Ours aims were therefore: (1) to identify factors associated with the acceptance of influenza and pertussis vaccinations in pregnancy, (2) to establish the level of confidence among HCPs in discussing vaccination with pregnant women, as well as the factors which might affect this, and (3) to establish the opinion of HCPs as to the optimum healthcare site for vaccine administration.

Methods

Questionnaire design and development

Two separate anonymized questionnaires were developed for pregnant women and maternity HCPs. These were developed with input from a multi-disciplinary study team including obstetricians, pediatricians, health psychologists, and clinical academic trainees. The questionnaires consisted of closed questions and a single free-text box in which participants could add further comments.

The questions analyzed here (see supplementary file) were nested within a larger questionnaire focussing on the attitudes of pregnant women and HCPs to both routine vaccination in pregnancy and to clinical trials of vaccines in pregnancy. The current paper focuses only on the questions relating to routinely recommended vaccines. Pregnant women were asked whether 1) they had/planned to receive influenza and pertussis vaccination and 2) the motivating reasons for accepting or declining these vaccines. Maternity HCPs were asked whether 1) they felt confident providing advice regarding these two vaccines and 2) their opinion regarding the optimal healthcare site of vaccine administration. Ethical approval was granted (reference 17/LO/0537) and the study was registered on ClinicalTrials.gov prior to recruitment (NCT03096574).

Study population and recruitment

The questionnaire for pregnant women was administered to women (aged ≥ 16 years at the time of completing the questionnaire) attending for routine antenatal care at four study sites in southern England: University Hospital Southampton NHS Foundation Trust, University Hospitals Bristol NHS Foundation Trust, Oxford University Hospitals NHS Foundation Trust, and St George's University Hospitals NHS Foundation Trust, London. These sites were selected because of their high birth rates (all > 4000 births/year)¹⁴, and by distributing our questionnaire across four hospitals, we attempted to increase the demographic diversity of our study population.

The HCP questionnaire was administered to those working in either midwifery or obstetrics at the same four study sites. It should be noted that routine antenatal care in the UK is usually midwife-led (unless a pregnancy is deemed high-risk), and therefore the majority of potential respondents to our questionnaire were midwives, rather than obstetricians. Recruitment of participants took place from July 2017 to January 2018. Pregnant women were recruited in person via opportunistic sampling at antenatal clinics or wards, and given paper questionnaires to complete. Maternity HCPs were either recruited via email (containing a link to an online questionnaire) or face-to-face by opportunistic sampling, in which case they were also given paper questionnaires. The initial response rate from HCPs was promoted by up to two further email reminders. Participation was voluntary and no financial or other incentive was offered. All participants gave informed consent.

Questionnaire data analysis

Questionnaire data was entered at the lead site (Southampton) into iSurvey (www.isurvey.soton.ac.uk). Statistical analysis was performed using IBM SPSS Version 25. Logistic and ordinal regression analyses were performed for pregnant women and HCP responses, respectively, and adjusted odds ratios (ORs) were calculated. P-values < 0.05 were considered as statistically significant. Multicollinearity was examined using the tolerance test and the Variance Inflation Factor (VIF) to ensure variables with a VIF value exceeding 2.5 were not entered into the multivariate regression analysis.

Results

A total of 525 participants completed the questionnaires: 321 pregnant women and 204 HCPs (18% obstetricians, 75% midwives, and 7% unidentified). The numbers of respondents were relatively equally distributed between the four study sites. Eight questionnaires from pregnant women, and five from HCPs, were excluded due to largely incomplete or illegible responses, therefore 513 questionnaires (98%) were included in the analysis. The full characteristics of respondents, including demographic details, are displayed in Table 1.

Responses from pregnant women

Regarding influenza vaccination: of 310 responses, 38% had been vaccinated, 40% were intending to be vaccinated, and 22% were not intending to be vaccinated. Regarding pertussis vaccination: of 302 responses, 56% had been vaccinated, 36% were intending to be vaccinated, and 8% were not intending to be vaccinated. The reasons for declining vaccination are displayed in Figure 1. A similar trend in responses was observed for both vaccines. The most commonly cited reason for declining was concern about possible side effects for their child.

Binary logistic regression analysis (Table 2, supplementary information) demonstrated that women identifying themselves as White British (79% of respondents) were significantly more likely to accept influenza (85% vs. 61%, OR 3.25, 95% confidence interval [CI] 1.67-6.32) and pertussis (96% vs. 83%, OR 4.83, 95% CI 1.77-13.19) vaccination compared to those identifying in all other ethnic groups. In the case of influenza vaccination, study site also had a significant effect, and participants at site B were significantly more likely to accept influenza vaccination than those at site D (91% vs. 64%, OR 4.20, 95% CI 1.47-11.95). Participants' age and whether they had previous children had no significant effect on vaccine uptake. In the qualitative analysis of the free text comments, pregnant women identified further concerns regarding vaccination in pregnancy, including damage to their unborn baby, vaccination being offered too late and insufficient information provided (see supplementary information).

Responses from maternity healthcare professionals

Out of 199 HCPs who responded, they were: extremely (25%), moderately (34%), somewhat (17%), slightly (16%) and not at all (8%) confident providing advice regarding influenza vaccination. For pertussis vaccination, they were: extremely (25%), moderately (32%), somewhat (16%), slightly (15%) and not at all confident (12%). See Figure 2.

Ordinal regression analysis (Table 3, supplementary information) demonstrated that obstetricians were significantly more likely than midwives to feel confident giving advice about the pertussis vaccine (68% vs. 55% were very/moderately confident, OR 2.05, 95% CI 1.02-4.12), however there was no significant difference between either profession for the influenza vaccine. On the other hand, longer experience in maternity care was associated with greater confidence giving advice regarding influenza vaccination, but not pertussis vaccination. Study site was also significantly associated with confidence providing advice for both vaccines, with HCPs from sites B and C being significantly more likely to feel confident than those in site D. Finally, health professional's age and whether or not they had children of their own were not associated with greater confidence in discussing vaccination. No free-text comments from staff relating to influenza/pertussis vaccination were provided for analysis.

218 With regards to the optimal healthcare site for vaccine administration during pregnancy (Figure 3),
219 approximately one-half (53%) of HCP respondents thought that vaccines should be delivered in the
220 primary care setting as part of general practice, 25% thought vaccines should be delivered in by
221 midwives in the community, and 16% thought vaccines should be delivered in secondary care (at
222 the time of antenatal appointments). The remaining 8% either thought that vaccination should be
223 administered in both general practice and community midwifery services (4%) or in all three
224 locations (4%).

Discussion

Vaccination in pregnancy remains a national and international priority for improving healthcare outcomes. Understanding women's and HCP's opinions and attitudes to vaccine acceptance are important in explaining current vaccination attainment levels. Our aims were to identify factors associated with vaccine acceptance and hesitancy among pregnant women, to establish whether HCPs feel confident discussing vaccination with these women, and to establish where HCPs thought these vaccines should be administered.

Uptake of vaccination among pregnant women

Encouragingly, the acceptance of influenza and pertussis vaccination was high among pregnant women in this study. The most common reasons for vaccine hesitancy were concerns about side effects, and doubts regarding the effectiveness and need for vaccination. Perception of possible harm is commonly cited as the primary reason for vaccine refusal among previous studies^{12 15}, and women are usually more concerned about potential risks to their child's health than their own¹⁶. Clearly, important misconceptions still exist regarding the safety of vaccines, including the presence of 'toxins' such as thimerosal (a mercury-containing preservative removed from childhood vaccines in 2001) that was proposed in 2005 to be associated with neurologic conditions, including autism¹⁷. We recommend that vaccine advocacy should emphasise the safety and efficacy of vaccination, specifically towards protection of the baby. Furthermore, accessible alternatives to face-to-face counseling that been successfully used in the past have included social media and webcasts^{18 19}, mobile phone text messages (such as Text4baby)^{20 21} and smart phone apps (such as MatImms²²).

Another important finding was that pregnant women of ethnic minorities were significantly less likely to accept vaccination than those identifying as 'White British'. Previous research has similarly demonstrated lower vaccine acceptance among these groups²³⁻²⁵, and these findings highlight the importance of taking into account possible cultural/religious and language barriers when counselling these women and producing educational materials. The underlying reasons for the difference in vaccine attitudes between ethnic groups remains a significant gap in our knowledge, and future studies in this specific area are needed. Interestingly, we did not find any significant effect of age or having children already in our study, however younger age has been shown to be associated with lower uptake in some previous studies^{23 26}. Study site had no effect on pertussis vaccine acceptance however there was significantly higher influenza vaccine acceptance among pregnant women at site B. These results may be skewed by the recruitment season of this site, however, as recruitment here was all undertaken entirely during the influenza vaccination season (which runs from September to February).

Confidence of healthcare professionals and optimal healthcare site for vaccine administration

Very few previous studies²⁷ have investigated to what extent HCPs feel confident discussing vaccination with pregnant women. This is despite the fact that pregnant women consider their HCP their most trusted source of information, and encouragement from them has been shown to increase intention to receive vaccination by up to 20 times^{13 12}. Conversely, a lack of knowledge of the indications and benefits of vaccination among HCPs has been identified as a barrier to implementation of vaccination recommendations²⁸. Among HCPs in our study, a significant proportion were not confident providing advice to pregnant women. Confidence also varied significantly by study site, suggesting that there is a potential risk of health inequalities based on

differing levels of vaccine confidence and recommendations across the South of England. Further education of multidisciplinary HCPs is essential, and individual barriers to active promotion of these vaccines need to be identified and reduced. Individual sites should aim to establish areas of low confidence within their own working body and push to incorporate active promotion of vaccination into routine antenatal care. Also, while it should be noted that obstetricians, and those with more experience in maternity care, felt more confident giving advice about the pertussis and influenza vaccines, respectively, we suggest that education should not be aimed solely at a particular profession, or those new to maternity care.

Finally, optimizing the healthcare site for vaccine administration is an important and topical issue which may have a considerable impact on vaccine uptake. In the UK, vaccination in pregnancy is usually provided in the primary care setting (within general practice), yet this presents a logistical barrier as it normally requires women to arrange extra primary care appointments. A more convenient and efficient approach might be to routinely offer and administer vaccination at the time of hospital antenatal appointments (such as the fetal anomaly scan at around 20 week's gestation), either by incorporating vaccination directly into these clinics, or providing adjacent vaccination clinics, which women are invited to visit immediately before or after their regular antenatal appointment^{29–31}. Previous studies have demonstrated that vaccinating in secondary care may indeed improve uptake^{29–31}, yet support for this approach appeared to be low (16%) among HCPs surveyed in this study. A lack of staff, lack of a suitable setting and resources, concerns regarding appropriate financial reimbursement, and lack of confidence with vaccine discussion, have all been identified as potential barriers to this approach by HCPs in previous studies^{30–33}. Potential solutions include employing dedicated vaccination staff (including vaccination specialist midwives) and improving vaccine education (as discussed above). Further pragmatic and/or qualitative research is also required to establish the feasibility and effectiveness of this approach, and to establish facilitators and barriers to its acceptance among both pregnant women and HCPs.

Strengths and limitations

This study had significant numbers of respondents, and by distributing our questionnaire at four hospitals in southern England we attempted to maximize the demographic diversity of our study population. That said, the responses to the questionnaire cannot be taken as representative of all pregnant women and maternity HCPs. Reported actual/intended vaccine uptake was higher among our questionnaire respondents than national reports of vaccine uptake, and this may limit the generalisability of our study findings. All of our respondents were recruited from antenatal clinics at tertiary hospitals, and therefore it is possible that our sample was missing subsets of the population that tend to be more anti-vaccination. Future studies would therefore benefit from including a greater number of study sites over a wider geographic area, and recruiting from different types of sites (including smaller non-tertiary hospitals and primary care) and perhaps utilizing online recruitment via popular websites and social media.

Another limitation is that we relied upon self-reported vaccination status/intention, and there is therefore potential reporting bias in our estimations, which may have been improved by verification of women's medical records following delivery; however recent evidence does suggest that self-reported intention correlates well with actual uptake of vaccination^{34 35}. Finally, the number of pregnant women/HCPs approached, and the number who declined participation (as well as their reasons for doing so) was not recorded, and we are therefore unable to report this.

327
328
329
330
331
332
333
334
335

Conclusions

Whilst the high acceptance of vaccination among respondents in this study was encouraging, misconceptions still exist regarding vaccine safety and efficacy. Further education of multidisciplinary HCPs is essential, and active vaccine promotion needs to be incorporated into routine antenatal care, with a particular emphasis on women from ethnic minorities.

Figures

Figure 1: Reasons why the surveyed pregnant women did not intend to receive influenza or pertussis vaccination in pregnancy

Figure 2: Healthcare professionals' confidence providing advice to pregnant women regarding influenza (A) and pertussis (B) vaccination in pregnancy

Figure 3: Healthcare professionals' opinions regarding the optimal healthcare site at which vaccines in pregnancy should be delivered

Tables

Table 1: Characteristics of the respondents to questionnaires (pregnant women and healthcare professionals)

Table 2 [Supplementary information]: Logistic regression analysis of factors predicting pregnant women's intention to receive vaccination

Table 3 [Supplementary information]: Ordinal regression analysis of factors predicting healthcare professionals' confidence in providing advice regarding vaccination in pregnancy

Acknowledgements

The authors would also like to thank all the pregnant women and healthcare staff who took part in the questionnaire, Stephen Yekini for his assistance with data collection in Southampton, and all of the non-study staff that helped facilitate recruitment in the participating sites.

Author Contributions

CW drafted the manuscript. All authors contributed to questionnaire design and critically revised the manuscript. CW, AC, JM, EK, RM, KB, PH, AK, AF, MS, TV, TN, MC and CJ were involved in study set up and data collection at the participating sites. CW, TN and CJ performed the data analysis. CJ conceived the study and was the chief investigator. All authors approved the final version of the manuscript.

Conflict of Interests Statement

CW, AC, JM, KB, PH, AK, AF, MS and CJ are investigators for clinical trials done on behalf of their respective institutions, sponsored by various vaccine manufacturers, but receive no personal funding for these activities. All other authors report no potential conflicts of interest.

Funding

The study was supported by a grant from the British Paediatric Allergy Immunity and Infection Group (BPAIIG). BPAIIG had no role in the study design, data collection, data analysis/interpretation, report writing, or the decision to submit the manuscript for publication.

Clinical trial registration

The questionnaire study was registered on ClinicalTrials.gov prior to recruitment (NCT03096574).

Ethical approval

Ethical approval was granted from the West London & GTAC NHS Research Ethics Committee (reference 17/LO/0537) on 6th April 2017.

References

1. McIntyre P, Wood N. Pertussis in early infancy: disease burden and preventive strategies. *Curr Opin Infect Dis.* 2009;22(3):215–23.
2. Mak T, Mangtani P, Leese J, et al. Influenza vaccination in pregnancy: current evidence and selected national policies. *Lancet Infect Dis.* 2008;8(1):44–52.
3. Poehling KA, Szilagyi PG, Staat MA, et al. Impact of maternal immunization on influenza hospitalizations in infants. *Am J Obstet Gynecol.* 2011;204(6 Suppl 1):S141–8.
4. Amirthalingam G, Andrews N, Campbell H, et al. Effectiveness of maternal pertussis vaccination in England: an observational study. *Lancet.* 2014;384(9953):1521–1528.
5. Zaman K, Roy E, Arifeen SE, et al. Effectiveness of maternal influenza immunization in mothers and infants. *New Engl J Med.* 2008;359(15):1555–64.
6. Madhi SA, Cutland CL, Kuwanda L, et al. Influenza Vaccination of Pregnant Women and Protection of Their Infants. *New Engl J Med.* 2014;371(10):918–931.
7. Jones CE, Calvert A, Le Doare K. Vaccination in Pregnancy – Recent Developments. *Pediatr Infect Dis J.* 2018;37(2):191–193
8. Wilson RJ, Paterson P, Jarrett C, Larson HJ. Understanding factors influencing vaccination acceptance during pregnancy globally: A literature review. *Vaccine.* 2015;33(47):6420–9.
9. Larson H, Jarrett C, Schulz W, et al. Measuring vaccine hesitancy: The development of a survey tool. *Vaccine.* 2015;33(34):4165–75.
10. Public Health England. Seasonal influenza vaccine uptake in GP patients: winter season 2016 to 2017. 2017. Available at: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/613452/Seasonal_influenza_vaccine_uptake_in_GP_patients_winter_season_2016_to_2017.pdf. Accessed on 16/10/2018.
11. Public Health England. Pertussis vaccination programme for pregnant women update: vaccine coverage in England, January to March 2017. *Health Protection Report* 2017;11(19). Available at: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/616198/hpr1917_prntl-prtssVC.pdf. Accessed on 16/10/2018.

12. Wiley KE, Massey PD, Cooper SC, et al. Pregnant women's intention to take up a post-partum pertussis vaccine, and their willingness to take up the vaccine while pregnant: A cross sectional survey. *Vaccine*. 2013;31(37):3972–3978.
13. Wiley K, Massey P, Cooper S, et al. Uptake of influenza vaccine by pregnant women: a cross-sectional survey. *Med J Aust*. 2013;198(7):373–375.
14. NHS Digital. NHS Maternity Statistics, England 2016-2017. 2018. Available at: <https://digital.nhs.uk/data-and-information/publications/statistical/nhs-maternity-statistics/2016-17>. Accessed on 16/10/2018.
15. Sim J, Ulanika A, Katikireddi S, Gorman D. "Out of two bad choices, I took the slightly better one": Vaccination dilemmas for Scottish and Polish migrant women during the H1N1 influenza pandemic. *Public Health*. 2011;125(8):505–511.
16. Wiley K, Cooper S, Wood N, Leask J. Understanding Pregnant Women's Attitudes and Behavior Toward Influenza and Pertussis Vaccination. *Qual Health Res*. 2014;25(3):360–370.
17. Wessel L. Vaccine myths. *Science* 2017;356(6336):368–372.
18. Baxter D. Approaches to the vaccination of pregnant women: experience from Stockport, UK, with prenatal influenza. *Hum Vaccin Immunother*. 2013;9(6):1360–3.
19. Kriss J, Frew P, Cortes M, et al. Evaluation of two vaccine education interventions to improve pertussis vaccination among pregnant African American women: A randomized controlled trial. *Vaccine*. 2017;35(11):1551–1558.
20. Kharbanda E, Vargas CY, Castaño PM, et al. Exploring pregnant women's views on influenza vaccination and educational text messages. *Prev Med*. 2011;52(1):75–77.
21. Gazmararian JA, Elon L, Yang B, et al. Text4baby program: an opportunity to reach underserved pregnant and postpartum women? *Matern Child Health J* 2014;18(1):223–32.
22. Holder B, Borgeau M, Donaldson B, Davies J, Kampmann B. MatImms: A smartphone app to inform and educate women about maternal immunisation. Poster presentation at 4th International Neonatal and Maternal Immunisation Symposium, Brussels, Belgium, 10-12/09/2017.
23. Goldfarb I, Little S, Brown J, Riley L. Use of the combined tetanus-diphtheria and pertussis vaccine during pregnancy. *Am J Obstet Gynecol*. 2014;211(3):299.e1–299.e5.
24. McQuaid F, Jones C, Stevens Z, et al. Antenatal vaccination against Group B streptococcus: Attitudes of pregnant women and healthcare professionals in the UK to participation in clinical trials and routine implementation. *Acta Obstet Gynecol Scand*. 2018;97(3):330-340.
25. Dempsey A, Brewer S, Sevic C, et al. Tdap vaccine attitudes and utilization among pregnant women from a high-risk population. *Hum Vaccin Immunother*. 2016;12(4):872–878.

26. Liu N, Sprague AE, Yasseen AS, et al. Vaccination patterns in pregnant women during the 2009 H1N1 influenza pandemic: a population-based study in Ontario, Canada. *Can J Public Health*. 2012;103(5):e353–8.
27. Vishram B, Letley L, Jan Van Hoek A, et al. Vaccination in pregnancy: Attitudes of nurses, midwives and health visitors in England. *Hum Vaccin Immunother*. 2018;14(1):179–188.
28. MacDougall D, Halperin S. Improving rates of maternal immunization: Challenges and opportunities. *Hum Vaccin Immunother*. 2015;12(4):857–865.
29. Yudin M, Salaripour M, Sgro M. Acceptability and Feasibility of Seasonal Influenza Vaccine Administration in an Antenatal Clinic Setting. *J Obstetrics Gynaecol Can* 2010;32(8):745–748.
30. Webb H, Street J, Marshall H. Incorporating immunizations into routine obstetric care to facilitate Health Care Practitioners in implementing maternal immunization recommendations. *Hum Vaccin Immunother*. 2014;1114–1121.
31. Bonville C, Cibula D, Domachowske J, Suryadevara M. Vaccine attitudes and practices among obstetric providers in New York State following the recommendation for pertussis vaccination during pregnancy. *Hum Vaccin Immunother*. 2015;713–718.
32. Ishola DA, Permalloo, Cordery RJ, Anderson SR. Midwives' influenza vaccine uptake and their views on vaccination of pregnant women. *J Pub Health* 2013;570–577.
33. Leddy MA, Anderson BL, Power ML, et al. Changes in and current status of obstetrician-gynecologists' knowledge, attitudes, and practice regarding immunization. *Obstet Gynecol Surv* 2009;64(12):823–9.
34. Harris K, Maurer J, Lurie N. Do people who intend to get a flu shot actually get one? *J Gen Intern Med* 2009;24(12):1311–3.
35. Mangtani P, Shah A, Roberts J. Validation of influenza and pneumococcal vaccine status in adults based on self-report. *Epidemiol Infect*. 2007;135(1):139–143.

Characteristic	Pregnant women, n=314	Healthcare professionals, n=199
Age		
16-24	34 (11%)	
25-30	107 (34%)	
31-35	92 (29%)	
36-40	58 (19%)	
41-45	13 (4%)	
46+	0 (0%)	
Gestation (weeks)		
<12	8 (2%)	
12-16	37 (12%)	
17-20	31 (10%)	
21-30	55 (18%)	
31-36	93 (30%)	
>37	76 (24%)	
Study site		
A	88 (28%)	43 (22%)
B	77 (25%)	53 (27%)
C	79 (25%)	61 (31%)
D	70 (22%)	42 (21%)
Ethnicity		
Asian (British, Indian, Pakistani, Bangladeshi, Chinese, other)	25 (8%)	4 (2%)
Black (British, African, Caribbean, other)	17 (5%)	4 (2%)
White (British, Irish, other)	248 (79%)	175 (88%)
Mixed (Caribbean, African, Asian, other)	11 (4%)	6 (3%)
Other ethnic group (Arab, other)	3 (1%)	0 (0%)
Did not want to say	1 (0.3%)	1 (1%)
No response	10 (3%)	9 (5%)
Has children		
No	142 (45%)	72 (36%)
Yes	172 (55%)	127 (64%)
Profession		
Obstetrics		37 (19%)
Midwifery		151 (76%)
No response		11 (6%)
Midwifery seniority		
Band 5 (newly-qualified midwife)		8 (4%)
Band 6 (junior midwife)		84 (42%)
Band 7 (senior midwife)		46 (23%)
Band 8 (midwifery manager)		8 (4%)
No response		5 (3%)
Obstetrician seniority		
Specialty training years 1-3 (or equivalent)		8 (22%)
Specialty training years 4-6 (or equivalent)		6 (16%)
Specialty training years 7-8 (or equivalent)		6 (16%)
Consultant		17 (46%)
Time spent working in maternity care (years)		
<2		17 (9%)
2-5		29 (15%)
6-10		37 (19%)
11-15		20 (10%)
16-20		26 (13%)
>21		62 (31%)
No response		8 (4%)

Table 1: Characteristics of the respondents to questionnaires (pregnant women and maternity healthcare professionals)

Variable	Number who had previously received/were intending to receive influenza vaccination (%)	Adjusted odds ratio (95% Confidence interval)	Number who had previously received/were intending to receive pertussis vaccination (%)	Adjusted odds ratio (95% Confidence interval)
Ethnicity				
White British	182/213 (85%)	3.25 (1.67-6.32) ***	203/212 (96%)	4.83 (1.77-13.19) **
Non-White British	51/84 (61%)	1.00 for reference	70/84 (83%)	1.00 for reference
Study site				
Site A	67/85 (79%)	1.36 (0.61-3.04)	73/81 (90%)	0.47 (0.14-1.62)
Site B	68/75 (91%)	4.20 (1.47-11.95) **	67/72 (93%)	0.91 (0.21-3.89)
Site C	58/74 (78%)	1.38 (0.58-3.30)	72/76 (95%)	1.22 (0.26-5.67)
Site D	40/63 (64%)	1.00 for reference	61/67 (91%)	1.00 for reference
Age				
16-24	26/32 (81%)	1.19 (0.40-3.58)	28/29 (97%)	1.96 (0.22-17.66)
25-35	152/188 (81%)	1.24 (0.62-2.49)	173/186 (93%)	1.12 (0.40-3.14)
36-45	49/68 (72%)	1.00 for reference	64/71 (90%)	1.00 for reference
Previous children				
Yes	127/163 (78%)	0.82 (0.44-1.53)	145/160 (91%)	0.47 (0.17-1.32)
No	106/134 (79%)	1.00 for reference	128/136 (94%)	1.00 for reference

*= p<0.05; **= p<0.01; ***= p<0.001;


Table 2: Logistic regression analysis of factors predicting pregnant women's intention to receive/previous receipt of vaccination

Variable	Number who were very or moderately confident providing advice about influenza vaccination (%)	Adjusted odds ratio (95% Confidence interval)	Number who were very or moderately confident providing advice about pertussis vaccination (%)	Adjusted odds ratio (95% Confidence interval)
Professional group				
Obstetrics	24/37 (65%)	2.00 (0.90-4.03)	25/37 (68%)	2.05 (1.02-4.12) *
Midwifery	90/151 (60%)	1.00 for reference	83/151 (55%)	1.00 for reference
Time working in maternity care				
21+ years	42/62 (68%)	3.88 (1.29-11.68) *	36/62 (58%)	1.72 (0.58-5.09)
11-20 years	32/46 (70%)	4.02 (1.33-12.15) *	30/46 (65%)	1.98 (0.67-5.87)
2-10 years	34/66 (52%)	2.83 (1.05-7.66) *	36/66 (55%)	2.22 (0.83-5.95)
<2 years	6/17 (35%)	1.00 for reference	7/17 (41%)	1.00 for reference
Study site				
A	24/43 (56%)	1.27 (0.54-2.99)	20/43 (47%)	1.09 (0.47-2.57)
B	39/53 (74%)	5.05 (2.12-12.01) ***	26/53 (49%)	4.68 (1.98-11.05) ***
C	41/61 (67%)	2.44 (1.13-5.29) *	42/61 (69%)	2.46 (1.14-5.30) *
D	15/42 (36%)	1.00 for reference	16/42 (38%)	1.00 for reference
Has their own children				
Yes	84/127 (66%)	1.23 (0.62-2.42)	76/127 (60%)	1.21 (0.52-3.63)
No	35/72 (49%)	1.00 for reference	38/72 (53%)	1.00 for reference
Ethnicity				
White British	108/175 (62%)	1.75 (0.66-4.66)	101/175 (58%)	1.37 (0.52-3.63)
Non-White British	11/24 (46%)	1.00 for reference	13/24 (54%)	1.00 for reference

*= p<0.05; **= p<0.01; ***= p<0.001

Table 3: Ordinal regression analysis of factors predicting healthcare professionals' confidence in providing advice regarding vaccination in pregnancy

535



536

537

538 **Figure 1:** Reasons why the surveyed pregnant women did not intend to receive influenza or
 539 pertussis vaccination in pregnancy

540

541

542

543

544

545

546

547

548

549

550

551

552

553

554

555

556

557

558




Figure 2: Healthcare professionals' confidence providing advice to pregnant women regarding influenza (A) and pertussis (B) vaccination in pregnancy




Figure 3: Healthcare professionals' opinions regarding the optimal healthcare site at which vaccines in pregnancy should be delivered

Supplementary information

1. Questions for pregnant women analyzed in this study

(1) Have you received either of the following vaccines in this pregnancy?

Flu (influenza) ☐ Yes ☐ No

Whooping cough (pertussis) ☐ Yes ☐ No

(1a) If no, why have you not yet received these vaccines?

For flu (influenza)?

For whooping cough(pertussis)?

I don't intend to receive the vaccine ☐

I haven't been offered the vaccine yet ☐

I haven't got round to getting the vaccine yet ☐

Other (please

specify).....

(1b) If you don't intend to receive these vaccines in this pregnancy, please specify the reasons why. Tick as many apply:

For flu (influenza)?

For pertussis (whooping cough)?

I worry that the injection might be painful ☐

My midwife did not advise it ☐

My obstetrician did not advise it ☐

My GP did not advise it ☐

My family/friends advised against it ☐

I don't believe the vaccine is effective ☐

I worry about potential side effects for my baby ☐

I worry about potential side effects for me ☐

Vaccination was not offered to me ☐

There is not enough safety data ☐

I don't have enough information to decide ☐

I am concerned about information in the media ☐

I don't want to attend extra hospital/GP visits ☐

Religious or other convictions ☐

Other (please specify):

.....

.....

(2) How old are you in years?

16-24 ☐ 25-30 ☐ 31-35 ☐ 36-40 ☐ 41-45 ☐ 46+ ☐

(3) How many weeks pregnant are you?

Less than 12 ☐ 12-16 ☐ 17-20 ☐ 21-30 ☐ 31-36 ☐ 37+ ☐

(4) To what ethnic group do you feel you belong? (Please circle)

White

Black / African / Caribbean / Black British

- English / Welsh / Scottish / Northern Irish - African

- 666 / British Irish - Caribbean
 667 - Gypsy or Irish Traveller - Other (please specify).....
 668 - Other (please specify)
 669
 670 **Mixed/Multiple ethnic groups** **Other ethnic group**
 671 - White and Black Caribbean - Arab
 672 - White and Black African - Other (please specify).....
 673 - White and Asian
 674 - Other (please specify)
 675
 676 **Asian / Asian British** **I'd prefer not to say**
 677 - Indian
 678 - Pakistani
 679 - Bangladeshi
 680 - Chinese
 681 - Other (please specify)
 682
 683

684 **(5) Have you had any children before?**

- 685 ☐ Yes.
 686 If yes, how many?.....
 687 What are their ages?
 688 Child 1: Less than 1 ☐ 1-5 ☐ 6-10 ☐ 11-16 ☐ 17+ ☐
 689 Child 2: Less than 1 ☐ 1-5 ☐ 6-10 ☐ 11-16 ☐ 17+ ☐
 690 Child 3: Less than 1 ☐ 1-5 ☐ 6-10 ☐ 11-16 ☐ 17+ ☐
 691 ☐ No
 692
 693
 694

695 **(6) Optional: Do you have any comments or concerns about any of the issues raised in the questionnaire?**

696
 697
 698
 699
 700
 701
 702
 703

704 **2. Questions for maternity healthcare professionals analyzed in this study**

705
 706
 707 **(1) How confident would you feel about providing advice regarding the flu (influenza) vaccine to women during pregnancy?**

- 708
 709 ☐ Not at all confident
 710 ☐ Slightly confident
 711 ☐ Somewhat confident
 712 ☐ Moderately confident
 713 ☐ Extremely confident
 714
 715

716 **(2) How confident would you feel about providing advice regarding the whooping cough (pertussis) vaccine to women during pregnancy?**

- 717
 718 ☐ Not at all confident
 719 ☐ Slightly confident
 720 ☐ Somewhat confident
 721 ☐ Moderately confident
 722 ☐ Extremely confident
 723

724 **(3) In your opinion, where should these vaccines be delivered to pregnant women?**
725
726 ☐ Primary care (GP practice)
727 ☐ Midwifery services (Community services)
728 ☐ Secondary care at time of antenatal scans or appointments
729 ☐ Other (please specify).....
730
731 **(4) Which healthcare professional group do you belong to?**
732 ☐ Obstetrics
733 ☐ Midwifery
734 ☐ Other (please state)
735
736
737 **(5) How long have you worked in maternity care?**
738 ☐ Under 2 years
739 ☐ 2-5 years
740 ☐ 6-10 years
741 ☐ 11-15 years
742 ☐ 16-20 years
743 ☐ 21+ years
744
745 **(6) What is your grade?**
746 *1. Midwifery/nursing staff*
747 Band 4 ☐ Band 5 ☐ Band 6 ☐ Band 7 ☐ Band 8 ☐ Band 9 ☐
748 *2. Obstetricians*
749 ST 1-3 (or equivalent) ☐ ST 4-6 (or equivalent) ☐ ST 7-8 (or equivalent) ☐ Consultant ☐
750
751 **(7) Have you had any children before?**
752 ☐ Yes.
753 If yes, how many?.....
754 What are their ages?
755 Child 1: Less than 1 ☐ 1-5 ☐ 6-10 ☐ 11-16 ☐ 17+ ☐
756 Child 2: Less than 1 ☐ 1-5 ☐ 6-10 ☐ 11-16 ☐ 17+ ☐
757 Child 3: Less than 1 ☐ 1-5 ☐ 6-10 ☐ 11-16 ☐ 17+ ☐
758 Child 4: Less than 1 ☐ 1-5 ☐ 6-10 ☐ 11-16 ☐ 17+ ☐
759
760 ☐ No
761
762 **(8) To what ethnic group do you feel you belong? (Please circle)**
763
764 **White** **Black / African / Caribbean / Black British**
765 - English / Welsh / Scottish / Northern Irish - African
766 / British Irish - Caribbean
767 - Gypsy or Irish Traveller - Other (please specify).....
768 - Other (please specify)
769
770 **Mixed/Multiple ethnic groups** **Other ethnic group**
771 - White and Black Caribbean - Arab
772 - White and Black African - Other (please specify).....
773 - White and Asian
774 - Other (please specify)
775
776 **Asian / Asian British** **I'd prefer not to say**
777 - Indian
778 - Pakistani
779 - Bangladeshi
780 - Chinese
781 - Other (please specify)

782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806

(9) Optional: Do you have any comments or concerns about vaccination or vaccine research studies during pregnancy?

.....
.....
.....
.....
.....
.....

3. Free text comments (all received from pregnant women)

“Many vaccines contain unsafe levels of mercury, in some cases are produced on human tissue (DNA) and contain various other toxins. I believe a baby is born with a perfect immune system which takes up to three years to fully develop and it’s not healthy injecting a perfectly healthy child with chemicals and toxins (mercury)”

“We were not offered the whooping cough [vaccine] until much later on in the pregnancy – close to it being too late. No flu jab offered – we would have done so otherwise.”

“I would like the opportunity to ask more questions and have more information before agreeing to vaccination”