

Mitochondrial DNA copy number variation and pancreatic cancer risk in the prospective EPIC cohort

Manuel Gentiluomo^{1;2}, Verena Katzke³, Rudolf Kaaks³, Anne Tjønneland^{4;5}, Gianluca Severi^{6;7}, Vittorio Perduca^{6;7;8}, Marie-Christine Boutron-Ruault^{6;7}, Elisabete Weiderpass⁹, Pietro Ferrari⁹, Theron Johnson³, Matthias B. Schulze^{10;11}, Manuela Bergmann¹², Antonia Trichopoulou¹³, Anna Karakatsani^{13;14}, Carlo La Vecchia^{13;15}, Domenico Palli¹⁶, Sara Grioni¹⁷, Salvatore Panico¹⁸, Rosario Tumino¹⁹, Carlotta Sacerdote²⁰, Bas Bueno de Mesquita^{21;22;23;24}, Roel Vermeulen^{25;26}, Torkjel M Sandanger²⁷, J. Ramón Quirós²⁸, Miguel Rodriguez Barranco^{29;30;31}, Pilar Amiano^{32;33}, Sandra Colorado-Yohar^{34;35;36}, Eva Ardanaz^{37;38;39}, Malin Sund⁴⁰, Kay-Tee Khaw⁴¹, Nick Wareham⁴², Julie A. Schmidt⁴³, Paula Jakszyn^{44;45}, Luca Morelli^{46;47}, Federico Canzian^{2#}, Daniele Campa^{1#}.

¹ Department of Biology, University of Pisa, Pisa, Italy

² Genomic Epidemiology Group, German Cancer Research Center (DKFZ), Heidelberg, Germany

³ Division of Cancer Epidemiology, German Cancer Research Center (DKFZ), Heidelberg, Germany.

⁴ Danish Cancer Society Research Center, Strandboulevarden 49, DK 2100 Copenhagen Ø Denmark

⁵ Department of Public Health, Faculty of Health and Medical Sciences, University of Copenhagen

⁶ CESP, Fac. de médecine - Univ. Paris-Sud, Fac. de médecine - UVSQ, INSERM, Université Paris-Saclay, 94805, Villejuif, France

⁷ Gustave Roussy, F-94805, Villejuif, France

⁸ Laboratoire de Mathématiques Appliquées MAP5 (UMR CNRS 8145), Université Paris Descartes, France

⁹ International Agency for Research on Cancer, World Health Organization, Lyon, France

¹⁰ Department of Molecular Epidemiology, German Institute of Human Nutrition Potsdam-Rehbruecke, Nuthetal, Germany

¹¹ Institute of Nutritional Sciences, University of Potsdam, Nuthetal, Germany

¹² Human Study Center, German Institute of Human Nutrition Potsdam-Rehbruecke, Nuthetal, Germany

¹³ Hellenic Health Foundation, Athens, Greece

- ¹⁴ Pulmonary Medicine Dept., School of Medicine, National and Kapodistrian University of Athens, Attikon University Hospital, Haidari, Greece
- ¹⁵ Dept. of Clinical Sciences and Community Health Università degli Studi di Milano
- ¹⁶ Cancer Risk Factors and Life-Style Epidemiology Unit, Institute for Cancer Research, Prevention and Clinical Network - ISPRO, Florence, Italy
- ¹⁷ Epidemiology and Prevention Unit, Fondazione IRCCS Istituto Nazionale dei Tumori di Milano, Milano, Italy
- ¹⁸ Dipartimento di medicina clinica e chirurgia, Federico II University, Naples, Italy
- ¹⁹ Cancer Registry and Histopathology Department, Azienda Sanitaria Provinciale Ragusa (ASP) Ragusa, Italy
- ²⁰ Unit of Cancer Epidemiology, Città della Salute e della Scienza University-Hospital and Center for Cancer Prevention (CPO), Turin, Italy
- ²¹ Former senior scientist, Dept. for Determinants of Chronic Diseases (DCD), National Institute for Public Health and the Environment (RIVM), PO Box 1, 3720 BA Bilthoven, The Netherlands
- ²² Former associate professor, Department of Gastroenterology and Hepatology, University Medical Center, Utrecht, The Netherlands.
- ²³ Former visiting professor, Dept. of Epidemiology and Biostatistics, The School of Public Health, Imperial College London, St Mary's Campus, Norfolk Place, London, W2 1PG London, United Kingdom.
- ²⁴ Former academic / visiting professor, Dept. of Social & Preventive Medicine, Faculty of Medicine, University of Malaya, Pantai Valley, 50603, Kuala Lumpur, Malaysia.
- ²⁵ Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht, the Netherlands.
- ²⁶ Environmental Epidemiology Division, Institute for Risk Assessment Sciences, Utrecht University, Utrecht, the Netherlands.
- ²⁷ Department of Community Medicine, UiT-the Arctic University of Norway, Tromsø, Norway
- ²⁸ Public Health Directorate, Asturias, Spain.
- ²⁹ Andalusian School of Public Health (EASP). Granada, Spain.
- ³⁰ Instituto de Investigación Biosanitaria de Granada (ibs.GRANADA). Universidad de Granada. Granada, Spain
- ³¹ CIBER of Epidemiology and Public Health (CIBERESP). Madrid, Spain"
- ³² Public Health Division of Gipuzkoa, Biodonostia Research Institute, Health Department, 20014 San Sebastian, Spain

- ³³ CIBER de Epidemiología y Salud Pública, CIBERESP, 28029 Madrid, Spain
- ³⁴ Department of Epidemiology, Murcia Regional Health Council, IMIB-Arrixaca, Murcia, Spain
- ³⁵ CIBER Epidemiología y Salud Pública (CIBERESP), Spain
- ³⁶ Research Group on Demography and Health, National Faculty of Public Health, University of Antioquia, Medellín, Colombia
- ³⁷ Navarra Public Health Institute, Pamplona, Spain
- ³⁸ IdiSNA, Navarra Institute for Health Research, Pamplona, Spain
- ³⁹ CIBER Epidemiology and Public Health CIBERESP, Madrid, Spain
- ⁴⁰ Department of Surgical and Perioperative sciences/ Surgery, Umeå University, Sweden
- ⁴¹ University of Cambridge, School of Clinical Medicine Addenbrooke's Hospital, Cambridge, United Kingdom.
- ⁴² MRC Epidemiology Unit, University of Cambridge School of Clinical Medicine, Institute of Metabolic Science, Cambridge, United Kingdom.
- ⁴³ Cancer Epidemiology Unit, Nuffield Department of Population Health, University of Oxford, United Kingdom
- ⁴⁴ Unit of Nutrition and Cancer, Cancer Epidemiology Research Program, Catalan Institute of Oncology-IDIBELL, L'Hospitalet de Llobregat, Barcelona, Spain
- ⁴⁵ Facultat Ciències Salut Blanquerna, Universitat Ramon Llull, Barcelona, Spain
- ⁴⁶ General Surgery, Department of Surgery, Translational and New Technologies, University of Pisa, Pisa, Italy
- ⁴⁷ EndoCAS (Center for Computer Assisted Surgery), University of Pisa, Pisa, Italy
- *#equal contribution

Running title: Mitochondrial DNA copy number and pancreatic cancer risk

Keywords: Pancreatic cancer; mitochondrial copy number; mtDNA copy number, cancer risk, prospective cohort, leukocyte.

Corresponding author: Daniele Campa

Department of Biology

University of Pisa

Via Derna 1

56126 Pisa

Italy

Email daniele.campa@unipi.it

phone +39-050-221510

Abstract

BACKGROUND: Mitochondrial DNA (mtDNA) copy number in peripheral blood has been found to be associated with risk of developing several cancers. However, data on pancreatic ductal adenocarcinoma (PDAC) are very limited.

METHODS: To further our knowledge on this topic we measured relative mtDNA copy number by a quantitative real-time PCR assay in peripheral leukocyte samples of 476 PDAC cases and 357 controls nested within the European Prospective Investigation into Cancer and Nutrition (EPIC) cohort.

RESULTS: We observed lower mtDNA copy number with advancing age ($p=6.54 \times 10^{-5}$) and with a high BMI level ($p=0.004$) and no association with sex, smoking behavior and alcohol consumption. We found an association between increased mtDNA copy number and decreased risk of developing PDAC with an OR=0.35 (95% C.I 0.16-0.79), $p=0.01$ when comparing the 5th quintile with the 1st using an unconditional logistic regression and OR=0.19 (95% C.I 0.07-0.52), $p=0.001$ with a conditional analysis. Analyses stratified by BMI showed an association between high mtDNA copy number and decreased risk in the stratum of normal weight, consistent with the main analyses.

CONCLUSIONS: Our results, suggest a protective effect of a higher number of mitochondria, measured in peripheral blood leukocytes, on PDAC risk.

IMPACT: Our findings highlight the importance of understanding the mitochondrial biology in pancreatic cancer.

Introduction

Pancreatic cancer is a relatively rare disease with an incidence of 17.8/100,000 (crude rate) in Europe, and with a very high mortality rate, considering a 5-year survival lower than 5% (1). Established risk factors are few compared to other cancer types and include smoking, diabetes mellitus, obesity and chronic pancreatitis (2). In the last decade PDAC mortality has continued to increase and it has been estimated that by 2030 it will be the second most frequent cancer for mortality in Europe (3). The poor prognosis is caused by several factors, including the aggressiveness of the disease, lack of effective treatments, lack of knowledge about biological markers for early detection and for risk prediction (4).

A better understanding of risk factors and availability of better risk markers might lead to improved chances of stratifying the population according to their risk and, in the long term, to a faster diagnosis (5–7). In addition, the discovery of novel risk markers could further our knowledge on the biology of this disease.

One such possible risk marker might be mitochondrial DNA. Mitochondria are organelles involved in the regulation of critical cellular functions such as apoptosis, calcium homeostasis and energy production via the oxidative phosphorylation reaction and are responsible for the production of reactive oxygen species (ROS) (8,9). Mitochondria possess own copies of DNA (mtDNA), which are maternally inherited, and in each eukaryotic cell there can be hundreds or thousands of copies of their genomes. mtDNA copy number represents the number of mitochondria contained in each cell, and this number is in a constant range in order to sustain the energetic needs of the cell (10). mtDNA copy number varies by cell type, but, in general, there is a correlation between the amount of mtDNA in different cell types (11). Therefore, mtDNA copy number measured in circulating leukocytes could represent a good and non-invasive indicator of the average amount of mtDNA copy number in other tissues (11). In cells under normal physiological conditions, the amount of mtDNA is relatively stable. Several reports have highlighted that mitochondrial copy number

increases to compensate for mtDNA damage and mitochondrial dysfunction (12) and in addition it could also represent a marker of endogenous and exogenous stressors including oxidative stress (13,14). The number of epidemiologic studies investigating the association of mtDNA copy number measured in leukocytes with cancer risk has been increasing in recent years, summing up to about thirty studies across various tumour types, including breast (11,15–17), colorectal (18–20), prostate (21–23), lymphoma (24,25) and pancreatic cancer (26). The results of these reports are very heterogeneous, with some studies showing an association between high number of mtDNA and increased risk while others showing the opposite. For pancreatic cancer only one prospective study, performed in male smokers in Finland, reported an association between high mtDNA copy number and increased risk to develop pancreatic ductal adenocarcinoma (PDAC) (26).

Given the importance of finding new risk factors for pancreatic cancer we performed a case-control study nested in the European Prospective Investigation into Cancer and Nutrition (EPIC) cohort study (27) investigating the possible association between mtDNA copy number variation in peripheral blood and risk of developing PDAC.

Materials and Methods

Study population

A full description of the EPIC cohort study has been given elsewhere (27). Briefly, EPIC consists of about 520,000 volunteers, recruited between 1992 and 2005 in 10 European countries. After providing informed consent, diet, lifestyle, and personal medical history questionnaires were collected and blood was drawn from participants at recruitment. Diagnosis of cancer determined after recruitment into the cohort was identified through local and national cancer registries (in Italy except the Naples center, Spain, Sweden, Netherlands, United Kingdom, Norway, Denmark) or by a combination of contacts with participants and local tumor registries, national health insurances or physicians and clinics within an active follow-up (in France, Germany, Greece and the Naples center in Italy). In this study 476 incident pancreatic cancer cases and 357 controls from eight countries (France, Germany, Greece, Italy, Spain, Sweden, Netherlands, and the United Kingdom) were used. Of these 833 individuals, 483 and 350, respectively, were women and men in the same proportions in cases and controls. Cases were diagnosed with exocrine pancreatic cancer, mainly pancreatic ductal adenocarcinoma (ICD-10, C25.0–25.3, 25.7–25.9). Due to the different etiology, endocrine pancreatic tumours (CD-10 C25.4,) were not included in this study. For a subgroup of cases (n=301) and an equal number of controls we performed a matching by center, sex, age at recruitment (± 6 months), date at entry in the cohort, and time between blood sampling and time of last consumption of food or drinks (<3, 3-6, and ≥ 6 hours) using an incidence density sampling protocol. The study was conducted in accordance with the Declaration of Helsinki and was approved by the ethics review board of the International Agency for Research on Cancer (IARC) in Lyon.

Sample preparation and DNA extraction

Blood samples from France, Germany, Greece, Italy, the Netherlands, Spain and the United Kingdom were conserved in liquid nitrogen (-196°C) in a central biorepository at IARC. Swedish samples were stored in Sweden in freezers at -70°C. DNA was extracted from leukocytes in batches of 96 with an Autopure instrument with Puregene chemistry. To minimise any possible bias due to differential handling each sample was extracted using the same method. DNA concentration and quality were measured using Qubit 4 Fluorometer (Thermo Fisher, Waltham, Massachusetts, US).

qPCR measurement of mtDNA copy-number

mtDNA copy number was measured in 833 study subjects (476 cases and 357 controls). To measure mtDNA we used a quantitative polymerase chain reaction (qPCR) that quantifies the copy number of the mitochondrial gene NADH dehydrogenase, subunit 1 (*ND1*) using as a reference the nuclear single copy gene albumin (*ALB*). Each reaction was performed in triplicate, in an optical 384-well reaction plate, in a 10 µL reaction volume using 2 µL of 5X HOT FIREPol Probe qPCR Mix Plus with ROX (Solis Bio-Dyne, Tartu, Estonia), 1.5 µM of Syto 9 (Invitrogen, Carlsbad, CA), 5 ng of genomic dried DNA and 8 µL of water. Two primers for *ND1* copy number, and two primers for quantifying *ALB* copy number were used. Each *ALB* primer was modified adding a GC-clamp to the 5' end in order to raise the melting temperature (primer sequences are shown in supplementary materials) (28).

The real-time PCR experiments were carried out using a Viia-7 sequence detection system (Applied Biosystems) using two subsequent (#1 *ND1*; #2 *ALB*) PCR cycling conditions performed in the same plates, to acquire the respective cycle thresholds (Ct) values for copy numbers of *ND1* and *ALB* (control) gene.

The conditions for amplification of *ND1* repeats were 95 °C/ 5 min, 2 cycles of 94 °C/15 sec and 60 °C/1 min, followed by 30 cycles of 85 °C/15 sec with the signal acquisition at 65

°C/1 min. Thermal conditions for *ALB* gene were 35 cycles of 95 °C/15 sec, 85 °C/30 sec, with the signal acquisition at 84 °C/30 sec. The specificity of all amplifications was determined by melting curve analysis done at default settings (95 °C/15 sec, 60 °C/1 min with the continuous signal acquisition at 0.05 °C/sec ramping, 95 °C/15 sec). A serial dilution (1:2) from 20 ng to 0.3 ng of genomic DNA pooled from 50 healthy individuals was included to generate the standard curves for *ND1* and *ALB* genes. The standard curve was used to quantify the *ND1* repeats and *ALB* gene, based on the respective Ct values. For each data point the obtained triplicate values were averaged. Individual values that deviated from the average of the triplicates by more than 5% of the standard deviation were discarded.

Standard curves were graphically represented as a semi-log regression line plot of Ct values and log of standard DNA concentration. The real-time PCR efficiency (E) of each reaction was calculated using standard curve points in the exponential phase according to the equation: $E = 10^{[-1/\text{slope}]}$. Samples whose Ct average was not within the standard curve range were discarded (26 samples) as "out of range" and not included in the analyses.

The mtDNA copy number was expressed as the ratio between *ND1/ALB*, using the Pfaffl method (29), which is best suited to the type of data obtained from a qPCR with efficiencies not perfectly identical between the amplification reactions of *ND1* and *ALB*, using as a calibrator the Ct of the standard curve to the equivalent of 5ng of DNA.

Statistical analysis

Association between mtDNA copy number and potential confounders at baseline, i.e. age, body mass index (BMI, kg/m²), smoking behavior and alcohol consumption, was tested using a generalized linear model (GLM) in the control group.

The mtDNA copy number was categorized into quintiles based on the distribution of mtDNA copy number in the controls and modelled as a categorical variable. The phenotype (pancreatic cancer case or control) was expressed as a dichotomic variable, age was

expressed as a continuous variable, the center of origin was expressed as a categorical variable, the plate of PCR-reactions was expressed as a categorical variable, BMI was expressed as a categorical variable with four values (underweight (BMI<19); normal weight (19≤BMI<25); overweight (25≤BMI<30); obese (BMI≥30), smoking behavior was expressed as a dichotomic variable (never, ever), and the same for alcohol consumption (non-drinker, drinker).

Odds ratios (ORs) and 95% confidence intervals (CIs) of the association between mtDNA copy number and pancreatic cancer risk were estimated using unconditional logistic regression adjusted by sex, age, center, BMI and plate.

Conditional logistic regression was also used on a subgroup of the subjects (n=301 case-control pairs), using center, sex, age at recruitment (±6 months), date at entry in the cohort, interval between blood sampling and time of last consumption of food and drink (<3, 3-6, and ≥6 hours) as matching variables. The analysis was further adjusted for BMI and plate. We also performed a conditional and unconditional analysis adjusting for diabetes and smoking status. However we had this information only in a subgroup of the subjects enrolled in the study.

Analyses were also performed by strata of lag time between blood collection and diagnosis of pancreatic cancer (<7 years, ≥7 years). We also performed analyses stratified by smoking, considering only males (77 pancreatic cancer cases and 80 controls), and the two sexes combined (142 pancreatic cancer cases and 134 controls). These two analyses were performed to compare the results with those of Lynch and collaborators (26). Finally, we performed analyses stratified by classes of BMI and age, because these two covariates are associated with mtDNA copy number, as reported in **Supplementary Table 1**.

A $p<0.05$ was considered statistically significant. All statistical analyses were two-sided.

Results

A detailed description of the study population is shown in **Table 1**.

We tested in controls the possible association between mtDNA copy number in leukocytes and age, BMI, smoking behavior and alcohol consumption. We observed lower mtDNA copy number with advancing age ($p=6.54 \times 10^{-5}$) and with a high BMI level ($p=0.004$). For sex, smoking behavior, diabetes status and alcohol consumption we observed no statistically significant associations (**Supplementary Table 1**).

The results from the unconditional logistic regression show an association between high mtDNA copy number and decreased risk of developing PDAC when analyzing mtDNA copy number categorized in quintiles (**Table 2**). We compared the fifth quintile (highest) with the first (lowest) and obtained OR=0.35 (95% CI 0.16-0.79, $p=0.01$). We also analyzed the quintiles of mtDNA copy number as continuous variable (i.e. the unit of measurement was the increase of one quintile) and obtained OR=0.79 (95% CI 0.66-0.96, $p=0.019$) (**Table 2**).

We also performed a conditional logistic regression analysis using mtDNA copy number categorized in quintiles on a subgroup of matched subjects and the results support the previous observations (OR=0.74, 95% CI 0.59-0.93, $p=0.008$ for the quintiles of mtDNA copy number analyzed as continuous variable, OR=0.19, 95% CI 0.07-0.52, $p=0.001$ comparing the 5th quintile (highest) with the first (lowest) (**Table 2**). The results of the analysis adjusted for diabetes confirmed a protective effect of mtDNA, although reaching statistical significance only in one of the quintiles. These analyses were done on a much smaller number of individuals: a total of 447 subjects for the unconditional analysis and 400 for the conditional analysis (**Supplementary Table 2**). The results of the analysis performed on the subgroup of individuals for which we had smoking history are in general agreement with the unadjusted analyses (**Supplementary Tables 2 and 3**).

We performed analyses stratified by lag time between blood collection and diagnosis of pancreatic cancer (<7 years, ≥ 7 years) and we did not observe any statistically significant

association in either of the strata (Supplementary Table 4). In addition, we also conducted a sensitivity analysis considering smokers (males and females alone and the two sexes combined); in the analyses of men and both sexes, we observed non-significant increases of risk in the highest quintiles (**Supplementary table 5**). Analyses stratified by BMI showed an association between high mtDNA copy number and decreased risk in the stratum of normal weight (**Supplementary table 6**).

Discussion

The association of mtDNA copy number with cancer risk has been studied in many cancer types with heterogeneous results (11,15–26,). In the present study we observed an association between a high level of mtDNA copy number and reduced risk of developing the disease. The association was consistent using both unconditional and conditional analysis (in a subgroup of individuals) and considering mtDNA quintiles as a continuous or categorical variable.

Lynch and colleagues in a nested case-control study performed in 203 PDAC cases who were Finnish male smokers and 656 male smoker controls within the prospective ATBC cohort found, instead, an association between high mtDNA copy number and increased PDAC risk(26). The differences in the results of the two studies may be explained by the differences in study design. Both are prospective studies, however, in the study by Lynch there are only male smokers while ours is considerably larger and composed by both genders and unselected for smoking status. We investigated the association between mtDNA copy number and pancreatic cancer risk in male smokers present in our study (77 pancreatic cancer cases and 80 controls), but we observed non-statistically significant results, although we obtained ORs>1 for the highest quintiles, which agrees with the results of Lynch et al. In addition, we also performed an analysis considering all the current smokers and obtained broadly similar results as in the male smokers.

Our study suggests that peripheral blood mtDNA copy number is inversely associated with BMI, and ageing, in agreement with the literature (30–32). We observed a non-statistically significant association between smoking behavior and lower mtDNA, as recently reported by Wu and colleagues in an all-female population (33).

A possible biological explanation of our findings on PDAC risk can be found in the central role that mitochondria have in regulating global variability in gene expression at cellular level (34). In particular, in a very recent study, Marquez-Jurado and colleagues have shown that

cells with an increased number of mitochondria have a faster response to external stress, increasing apoptotic protein synthesis and triggering the apoptosis process more quickly (35). The authors also observed that high mitochondrial content is associated with increased apoptotic inducers such as TNF alpha, suggesting a key role for mitochondria in discriminating cell fate (35). Additionally Armstrong and colleagues showed a direct effect of altered mitochondrial bioenergetics in pancreatic acinal cells and in the regulation of apoptosis to necrosis switch (36). Triggering a faster apoptotic response could be instrumental in avoiding tissue necrosis, that can cause inflammation that can degenerate into pancreatitis or acute pancreatitis and in turn lead to PDAC development (37,38). In addition several authors observed that there is a correlation between low blood mtDNA copy number with high levels of circulating inflammatory markers such as IL-6, CRP and neutrophil-to-lymphocyte ratio (39–41) that have been investigated as PDAC risk and prognostic markers (42–44). Given that mtDNA copy number measured in leukocytes is considered to be a good proxy for mtDNA content in other tissues (10,11), one can speculate that low levels of mtDNA copy number, measured in blood, may be associated with an increased level of inflammation, that could increase the risk of developing PDAC (45).

This study has clear strengths, such as its prospective nature that reduces the risk of reverse causation bias, and the fact that we have taken into consideration potential confounding factors, such as body mass index, age, smoking, and alcohol consumption. A limitation is its rather small sample size even though it is the largest study on PDAC on mtDNA copy number attempted so far. In addition our study is based on a single measurement of mtDNA which does not reflect the possible fluctuation of mtDNA over long term. However in a previous study we have shown that mtDNA copy numbers are stable over the years (46).

In conclusion, several evidences, including our results, suggest a protective effect of a higher number of mitochondria, measured in peripheral blood leukocytes, on PDAC risk and highlight the importance of understanding the mitochondrial biology in pancreatic cancer.

Disclaimer

Where authors are identified as personnel of the International Agency for Research on Cancer / World Health Organization, the authors alone are responsible for the views expressed in this article and they do not necessarily represent the decisions, policy or views of the International Agency for Research on Cancer / World Health Organization.

Conflict of interest

The authors declare no potential conflicts of interest.

Acknowledgements

We thank the National Institute for Public Health and the Environment (RIVM), Bilthoven, the Netherlands, for their contribution and ongoing support to the EPIC Study.

Funding support

This work was partially supported by intramural funds of University of Pisa and DKFZ, by Fondazione Tizzi and by Fondazione Arpa (www.fondazionearpa.it).

The EPIC-Potsdam study was funded by the Federal Ministry of Education and Research (Germany), the German Cancer Aid, the German Cancer Research Center and the German Institute of Human Nutrition Potsdam-Rehbrücke.

EPIC-Oxford was supported by Cancer Research UK (C8221/A19170) and the Medical Research Council UK (MR/M012190/1).

EPIC-Greece was supported by the Hellenic Health Foundation.

Availability of data and materials

For information on how to submit an application for gaining access to EPIC data and/or biospecimens, please follow the instructions at <http://epic.iarc.fr/access/index.php>.

References

1. Ferlay J, Colombet M, Soerjomataram I, Mathers C, Parkin DM, Piñeros M, et al. Estimating the global cancer incidence and mortality in 2018: GLOBOCAN sources and methods. *Int J Cancer* [Internet]. John Wiley & Sons, Ltd; 2018;ijc.31937.
2. Maisonneuve P, Lowenfels AB. Risk factors for pancreatic cancer: a summary review of meta-analytical studies. *Int J Epidemiol* [Internet]. 2015;44:186–98.
3. Rahib L, Smith BD, Aizenberg R, Rosenzweig AB, Fleshman JM, Matrisian LM. Projecting cancer incidence and deaths to 2030: The unexpected burden of thyroid, liver, and pancreas cancers in the united states [Internet]. *Cancer Res. American Association for Cancer Research*; 2014. page 2913–21.
4. Cros J, Raffenne J, Couvelard A, Poté N. Tumor Heterogeneity in Pancreatic Adenocarcinoma. *Pathobiology* [Internet]. 2018;85:64–71.
5. Ilic M, Ilic I. Epidemiology of pancreatic cancer. *World J Gastroenterol* [Internet]. 2016;22:9694–705.
6. Cooperman AM, Iskandar ME, Wayne MG, Steele JG. Prevention and Early Detection of Pancreatic Cancer. *Surg Clin North Am* [Internet]. 2018;98:1–12.
7. Chiaro M Del, Segersvärd R, Lohr M, Verbeke C. Early detection and prevention of pancreatic cancer: Is it really possible today? *World J Gastroenterol* [Internet]. 2014;20:12118.
8. Mills EL, Kelly B, Logan A, Costa ASH, Varma M, Bryant CE, et al. Succinate Dehydrogenase Supports Metabolic Repurposing of Mitochondria to Drive Inflammatory Macrophages. *Cell* [Internet]. 2016;167:457-470.e13.
9. Zhou R, Yazdi AS, Menu P, Tschopp J. A role for mitochondria in NLRP3 inflammasome activation. *Nature* [Internet]. 2011;469:221–5.
10. Veltri KL, Espiritu M, Singh G. Distinct genomic copy number in mitochondria of different mammalian organs. *J Cell Physiol* [Internet]. Wiley Subscription Services, Inc., A Wiley Company; 1990;143:160–4.
11. Shen J, Platek M, Mahasneh A, Ambrosone CB, Zhao H. Mitochondrial copy number and risk of breast cancer: A pilot study. *Mitochondrion* [Internet]. 2010;10:62–8.
12. Lee HC, Yin PH, Lu CY, Chi CW, Wei YH. Increase of mitochondria and mitochondrial DNA in response to oxidative stress in human cells. *Biochem J* [Internet]. 2000;348 Pt 2:425–32.
13. Chen XJ, Butow RA. The organization and inheritance of the mitochondrial genome. *Nat Rev Genet* 2005 611 [Internet]. Nature Publishing Group; 2005;6:815.
14. Lee HC, Wei YH. Mitochondrial biogenesis and mitochondrial DNA maintenance of mammalian cells under oxidative stress. *Int J Biochem Cell Biol*. 2005;
15. Thyagarajan B, Wang R, Nelson H, Barcelo H, Koh WP, Yuan JM. Mitochondrial DNA Copy Number Is Associated with Breast Cancer Risk. *PLoS One* [Internet]. Public Library of Science; 2013;8:e65968.
16. Lemnrau A, Brook MN, Fletcher O, Coulson P, Tomczyk K, Jones M, et al. Mitochondrial DNA copy number in peripheral blood cells and risk of developing breast cancer. *Cancer Res*. 2015;75:2844–50.
17. Campa D, Barrdahl M, Santoro A, Severi G, Baglietto L, Omichessan H, et al. Mitochondrial DNA copy number variation, leukocyte telomere length, and breast cancer risk in the European Prospective Investigation into Cancer and Nutrition (EPIC) study. *Breast Cancer Res* [Internet]. BioMed Central; 2018;20:29.
18. Thyagarajan B, Wang R, Barcelo H, Koh WP, Yuan JM. Mitochondrial copy number is associated with colorectal cancer risk. *Cancer Epidemiol Biomarkers Prev* [Internet]. NIH Public Access; 2012;21:1574–81.
19. Huang B, Gao YTY-T, Shu XOX-O, Wen W, Yang G, Li G, et al. Association of Leukocyte Mitochondrial DNA Copy Number with Colorectal Cancer Risk: Results

- from the Shanghai Women's Health Study. *Cancer Epidemiol Biomarkers Prev* [Internet]. NIH Public Access; 2014;23:2357–65.
20. Kumar B, Bhat ZI, Bansal S, Saini S, Naseem A, Wahabi K, et al. Association of mitochondrial copy number variation and T16189C polymorphism with colorectal cancer in North Indian population. *Tumor Biol.* 2017;39.
 21. Zhou W, Zhu M, Gui M, Huang L, Long Z, Wang L, et al. Peripheral Blood Mitochondrial DNA Copy Number Is Associated with Prostate Cancer Risk and Tumor Burden. *PLoS One* [Internet]. Public Library of Science; 2014;9:e109470.
 22. Tu H, Gu J, Meng QH, Kim J, Davis JW, He Y, et al. Mitochondrial DNA copy number in peripheral blood leukocytes and the aggressiveness of localized prostate cancer. *Oncotarget.* 2015;
 23. Moore A, Lan Q, Hofmann JN, Liu C-S, Cheng W-L, Lin T-T, et al. A prospective study of mitochondrial DNA copy number and the risk of prostate cancer. *Cancer Causes Control* [Internet]. Springer International Publishing; 2017;28:529–38.
 24. Hosnijeh FS, Lan Q, Rothman N, San Liu C, Cheng W-L, Nieters A, et al. Mitochondrial DNA copy number and future risk of B-cell lymphoma in a nested case-control study in the prospective EPIC cohort. *Blood* [Internet]. 2014;124:530–5.
 25. Lan Q, Lim U, Liu CS, Weinstein SJ, Chanock S, Bonner MR, et al. A prospective study of mitochondrial DNA copy number and risk of non-Hodgkin lymphoma. *Blood.* 2008;112:4247–9.
 26. Lynch SM, Weinstein SJ, Virtamo J, Lan Q, Liu C-S, Cheng W-L, et al. Mitochondrial DNA Copy Number and Pancreatic Cancer in the Alpha-Tocopherol Beta-Carotene Cancer Prevention Study. *Cancer Prev Res* [Internet]. NIH Public Access; 2011;4:1912–9.
 27. Riboli E, Hunt K, Slimani N, Ferrari P, Norat T, Fahey M, et al. European Prospective Investigation into Cancer and Nutrition (EPIC): study populations and data collection. *Public Health Nutr* [Internet]. Cambridge University Press; 2002;5:1113.
 28. Cawthon RM. Telomere length measurement by a novel monochrome multiplex quantitative PCR method. *Nucleic Acids Res* [Internet]. Oxford University Press; 2009;37:e21.
 29. Pfaffl MW. A new mathematical model for relative quantification in real-time RT-PCR [Internet]. *Nucleic Acids Res.* 2001.
 30. Skuratovskaia DA, Sofronova JK, Zatolokin PA, Popadin KY, Vasilenko MA, Litvinova LS, et al. Additional evidence of the link between mtDNA copy number and the body mass index [Internet]. *Mitochondrial DNA Part A DNA Mapping, Seq. Anal.* 2018. page 1240–4.
 31. Meng S, Wu S, Liang L, Liang G, Giovannucci E, De Vivo I, et al. Leukocyte mitochondrial DNA copy number, anthropometric indices, and weight change in US women. *Oncotarget* [Internet]. Impact Journals, LLC; 2016;7:60676–86.
 32. Mengel-From J, Thinggaard M, Dalgård C, Kyvik KO, Christensen K, Christiansen L. Mitochondrial DNA copy number in peripheral blood cells declines with age and is associated with general health among elderly. *Hum Genet* [Internet]. Springer Berlin Heidelberg; 2014;133:1149–59.
 33. Wu S, Li X, Meng S, Fung T, Chan AT, Liang G, et al. Fruit and vegetable consumption, cigarette smoke, and leukocyte mitochondrial DNA copy number. *Am J Clin Nutr* [Internet]. Narnia; 2019;109:424–32.
 34. Guantes R, Rastrojo A, Neves R, Lima A, Aguado B, Iborra FJ. Global variability in gene expression and alternative splicing is modulated by mitochondrial content. *Genome Res* [Internet]. Cold Spring Harbor Laboratory Press; 2015;125:633–44.
 35. Márquez-Jurado S, Díaz-Colunga J, Das Neves RP, Martínez-Lorente A, Almazán F, Guantes R, et al. Mitochondrial levels determine variability in cell death by modulating apoptotic gene expression. *Nat Commun* [Internet]. Nature Publishing Group;

2018;9:389.

36. Armstrong JA, Cash NJ, Ouyang Y, Morton JC, Chvanov M, Latawiec D, et al. Oxidative stress alters mitochondrial bioenergetics and modifies pancreatic cell death independently of cyclophilin D, resulting in an apoptosis-to-necrosis shift. *J Biol Chem* [Internet]. American Society for Biochemistry and Molecular Biology; 2018;293:8032–47.
37. Criddle DN. Reactive oxygen species, Ca²⁺ stores and acute pancreatitis; a step closer to therapy? [Internet]. *Cell Calcium*. Churchill Livingstone; 2016. page 180–9.
38. Gukovsky I, Pandol SJ, Gukovskaya AS. Organellar Dysfunction in the Pathogenesis of Pancreatitis. *Antioxid Redox Signal* [Internet]. Mary Ann Liebert, Inc.; 2011;15:2699–710.
39. Knez J, Marrachelli VG, Cauwenberghs N, Winckelmans E, Zhang Z, Thijs L, et al. Peripheral blood mitochondrial DNA content in relation to circulating metabolites and inflammatory markers: A population study. Samuels DC, editor. *PLoS One* [Internet]. Public Library of Science; 2017;12:e0181036.
40. Fisic E, Poropat G, Bilic-Zulle L, Licul V, Milic S, Stimac D. The Role of IL-6, 8, and 10, sTNF α , CRP, and Pancreatic Elastase in the Prediction of Systemic Complications in Patients with Acute Pancreatitis. *Gastroenterol Res Pract* [Internet]. Hindawi; 2013;2013:282645.
41. Imamura T, Tanaka S, Yoshida H, Kitamura K, Ikegami A, Takahashi A, et al. Significance of measurement of high-sensitivity C-reactive protein in acute pancreatitis. *J Gastroenterol* [Internet]. 2002;37:935–8.
42. Grote VA, Kaaks R, Nieters A, Tjønneland A, Halkjær J, Overvad K, et al. Inflammation marker and risk of pancreatic cancer: a nested case-control study within the EPIC cohort. *Br J Cancer* [Internet]. Nature Publishing Group; 2012;106:1866–74.
43. Arima K, Okabe H, Hashimoto D, Chikamoto A, Tsuji A, Yamamura K, et al. The diagnostic role of the neutrophil-to-lymphocyte ratio in predicting pancreatic ductal adenocarcinoma in patients with pancreatic diseases. *Int J Clin Oncol* [Internet]. 2016;21:940–5.
44. Sierzega M, Lenart M, Rutkowska M, Surman M, Mytar B, Matyja A, et al. Preoperative Neutrophil-Lymphocyte and Lymphocyte-Monocyte Ratios Reflect Immune Cell Population Rearrangement in Resectable Pancreatic Cancer. *Ann Surg Oncol* [Internet]. 2017;24:808–15.
45. Padoan A, Plebani M, Basso D. Inflammation and Pancreatic Cancer: Focus on Metabolism, Cytokines, and Immunity. *Int J Mol Sci* [Internet]. Multidisciplinary Digital Publishing Institute (MDPI); 2019;20:676.
46. Campa D, Barrdahl M, Santoro A, Severi G, Baglietto L, Omichessan H, et al. Mitochondrial DNA copy number variation, leukocyte telomere length, and breast cancer risk in the European Prospective Investigation into Cancer and Nutrition (EPIC) study. *Breast Cancer Res. England*; 2018;20:29.

Table 1. Participant characteristics. Distribution across countries and characteristics of the cases and controls participating in the European Prospective Investigation into Cancer and Nutrition (EPIC) study.

Country	Controls	Cases	Total
France	11	10	21
Germany	55	90	145
Greek	21	32	53
Italy	33	56	89
Spain	37	49	86
Sweden	119	117	236
Netherlands	39	58	97
United Kingdom	42	64	106
Total	357	476	833
Sex			
Men	155 (43%)	195 (41%)	350 (42%)
Women	202 (57%)	281 (59%)	483 (58%)
Mean age (years)	57	57	57
Median age at recruitment	60	58	59
(25 th -75 th percentile)	36-75	35-75	30-75
BMI (median-mean)	25 - 25.95	26 – 26.76	26 – 26.41
BMI (%)			
Underweight (BMI<19)	10 (3%)	1 (<1%)	11 (1%)
Healthy weight (19<BMI<25)	141 (42%)	182 (40%)	324 (41%)
Overweight (25<BMI<30)	131 (39%)	192 (42%)	323 (41%)
Obese (BMI>30)	54 (16%)	85 (18%)	139 (17%)
Cigarette smokers			
Never	130 (49%)	174 (45%)	304 (47%)
Ever	137 (51%)	211 (55%)	348 (53%)
- Current	49 (18%)	109 (28%)	158 (24%)
- Former	88 (33%)	102 (27%)	190 (29%)
Alcohol use			
Non-drinker	41 (14%)	49 (18%)	90 (16%)
Drinker	251 (86%)	223 (82%)	474 (84%)
Diabetes status			
Diabetic	13 (5%)	29 (8%)	42 (7%)
Non-diabetic	227 (95%)	317 (92%)	544 (93%)

Table 2. Association between mtDNA copy number and PDAC risk using an unconditional and a conditional analysis.

Unconditional analysis	Controls	Cases	Total	OR	95% C.I.	P-value
Continuous variable log-transformed	357	476	833	0.47	0.28; 0.79	0.004
Continuous variable divided in quintiles	357	476	833	0.79	0.66, 0.96	0.019
Analysis by quintiles						
1 st quintile	76	126	202	ref	-	-
2 nd quintile	70	105	175	0.91	0.57, 1.56	0.809
3 rd quintile	68	117	185	0.85	0.51, 1.52	0.639
4 th quintile	74	71	145	0.49	0.24, 1.08	0.077
5 th quintile	69	57	126	0.35	0.16, 0.79	0.010
Conditional analysis						
Continuous variable log-transformed	301	301	602	0.35	0.19; 0.64	0.001
Continuous variable divided in quintiles	301	301	602	0.74	0.59; 0.93	0.008
Analysis by quintiles						
1 st quintile	61	68	129	ref	-	-
2 nd quintile	60	56	116	0.81	0.47, 1.40	0.448
3 rd quintile	55	69	124	0.94	0.52, 1.68	0.827
4 th quintile	61	63	124	0.34	0.13, 0.87	0.025
5 th quintile	64	45	109	0.19	0.07, 0.52	0.001

Unconditional logistic regression performed using mitochondrial copy number variable (Pfaffl) categorized in quintiles and analyzed as continuous variable log-transformed, and continuous variable divided in quintiles (whereby the unit of measurement is a single quintile). Analyses were adjusted for sex, age, BMI, plate and recruitment center. Individual matching in conditional analysis was done by center, gender, age at recruitment (± 6 months), date at entry in the cohort, time between blood sampling, and time of last consumption of food and drink (<3 , $3-6$, and ≥ 6 hours). This analysis was adjusted for plate and BMI.

Supplementary materials

Primers

Primers for *ND1* copy number: forward primer 5'-CCCTAAAACCCGCCACATCT-3' and reverse primer 5'- GAGCGATGGTGAGAGCTAAGGT-3' amplify a 69-bp amplicon.

Primers for quantifying *ALB* copy number, modified adding of a GC-clamp (in upper-case letters) to the 5' end of each primer to raise the melting temperature: forward primer 5'- cggcggcgggcggcgcgggctgggcgg|CCATGCTTTTCAGCTCTGCAAGTC-3' and reverse primer 5'-gcccggcccggccgccccgtcccggcg|AGCATTAAGCTCTTTGGCAACGTAGGTTTC-3'

Supplementary Table 1. Association between mtDNA copy number and PDAC risk factors.

Variable	Coeff.*	95% C.I.	P-value
Age (expressed in years)	-0.03	-0.04, -0.01	0.005
- Age >60years vs <60years	-0.62	-0.92, -0.31	6.54×10⁻⁵
Sex	-0.26	-0.56, 0.04	0.089
BMI	-0.29	-0.49, -0.10	0.004
- healthy weight vs underweight	-0.92	-1.83, -0.11	0.047
- overweight vs underweight	-1.13	-2.04, -0.22	0.015
- obese vs underweight	-1.41	-2.37, -0.45	0.004
- underweight vs healthy weight	0.92	0.01, 1.83	0.047
- obese vs healthy weight	-0.49	-0.93, -0.04	0.033
Alcohol (no/yes)	-0.53	-1.09, 0.03	0.065
Smoking (no/yes)	-0.19	-0.66, 0.29	0.439
Smoking (never; former; current)	-0.12	-0.33, 0.10	0.293
Diabetes (no/yes)	-0.79	-1.58, 0.01	0.051

* regression coefficient.

Body mass index (BMI) was divided in four class: underweight (BMI<19); normal weight ((19≤BMI<25); overweight (25≤BMI<30); obese (BMI≥30); For sex the reference class is females and risk is referred to males.

Supplementary Table 2. Association between mtDNA copy number and pancreatic cancer adjusted for diabetes condition and smoking status.

	Controls	Cases	Total	OR	95% C.I.	P-value
Adjusted by diabetes						
Unconditional analysis						
Continuous variable log-trasformed	226	221	447	0.68	0.38; 1.22	0.197
Continuous variable expressed in quintiles	226	221	447	0.91	0.73 ;1.13	0.389
1 st quintile	67	65	132	ref	-	-
2 nd quintile	53	59	112	1.18	0.68; 2.08	0.553
3 rd quintile	46	51	97	1.06	0.58; 1.96	0.844
4 th quintile	31	20	51	0.44	0.18; 1.09	0.076
5 th quintile	29	26	55	0.67	0.26; 1.77	0.421
Conditional analysis						
Continuous variable log-trasformed	200	200	400	0.52	0.26; 1.05	0.067
Continuous variable expressed in quintiles	200	200	400	0.83	0.64; 1.07	0.154
1 st quintile	55	57	112	ref		
2 nd quintile	47	47	94	0.92	0.49; 1.72	0.799
3 rd quintile	38	48	86	1.01	0.52; 1.98	0.972
4 th quintile	28	20	48	0.28	0.09; 0.91	0.034
5 th quintile	32	28	60	0.30	0.09; 1.01	0.052
Adjusted by smoking status (never;former;current)						
Unconditional analysis						
Continuous variable log-trasformed	267	385	652	0.62	0.35; 1.11	0.107
Continuous variable expressed in quintiles	267	385	652	0.88	0.72; 1.09	0.255
1 st quintile	74	121	195	ref		
2 nd quintile	62	101	163	1.10	0.65; 1.86	0.736
3 rd quintile	57	100	157	1.00	0.57; 1.77	0.990
4 th quintile	37	28	65	0.38	0.16; 0.90	0.029
5 th quintile	37	35	72	0.61	0.24; 1.55	0.303
Conditional analysis						
Continuous variable log-trasformed	222	222	444	0.50	0.25; 0.99	0.049
Continuous variable expressed in quintiles	222	222	444	0.84	0.66; 1.06	0.145
1 st quintile	59	64	123	ref		
2 nd quintile	55	52	107	0.89	0.50; 1.56	0.676
3 rd quintile	47	58	105	0.99	0.54; 1.81	0.972
4 th quintile	29	20	49	0.27	0.09; 0.84	0.024
5 th quintile	32	28	60	0.28	0.08; 0.93	0.038

Unconditional logistic regression performed using mitochondrial copy number variable (Pfaffl) categorized in quintiles and analyzed as continuous variable log-transformed, continuous variable divided in quintiles (whereby the unit of measurement is a single quintile). Analyses were adjusted for sex, age, BMI, plate recruitment center and separately the analysis were adjusted for diabetes condition and smoking status. Individual matching in conditional analysis was done by center, gender, age at recruitment (± 6 months), date at entry in the cohort, time between blood sampling, and time of last consumption of food and drink (<3, 3-6, and ≥ 6 hours). This analysis was adjusted for plate and BMI.

Supplementary Table 3. Association between mtDNA copy number and pancreatic cancer adjusted for number of cigarette packs smoking for year .

	Controls	Cases	Total	OR	95% C.I.	P-value
Smoking (pack-years)						
Continuous variable log-transformed	111	120	231	0.90	0.65; 1.25	0.544
Continuous variable expressed in quintiles	111	120	231	0.76	0.29; 1.97	0.569
1 st quintile	37	40	77	ref		
2 nd quintile	29	38	67	1.08	0.52; 2.28	0.829
3 rd quintile	29	27	56	0.72	0.32; 1.63	0.435
4 th quintile	10	7	17	0.63	0.14; 1.93	0.559
5 th quintile	6	8	14	1.03	0.17; 6.12	0.973

Unconditional logistic regression performed using mitochondrial copy number variable (Pfaffl) categorized in quintiles and analyzed as continuous variable log-transformed and continuous variable divided in quintiles (whereby the unit of measurement is a single quintile). Analyses were adjusted for sex, age, BMI, plate recruitment center and cigarettes smoking packs.

Supplementary Table 4 Analyses stratified by lag time between blood collection and diagnosis of pancreatic cancer (<7 years, ≥7 years)

	Controls	Cases	Total	OR	95% C.I.	P-value
Conditional analysis cases <7 years from diagnosis						
Continuous variable log-transformed	136	136	272	0.58	0.25; 1.33	0.198
Continuous variable expressed in quintiles	136	136	272	0.83	0.62; 1.12	0.219
1 st quintile	41	48	89	ref		
2 nd quintile	42	37	79	0.73	0.37; 1.44	0.369
3 rd quintile	26	30	56	0.92	0.45; 1.88	0.820
4 th quintile	14	9	23	0.20	0.04; 1.05	0.057
5 th quintile	13	12	25	0.23	0.04; 1.40	0.110
Conditional analysis cases ≥7 years from diagnosis						
Continuous variable log-transformed	74	74	148	0.47	0.15; 1.52	0.208
Continuous variable expressed in quintiles	74	74	148	0.83	0.54; 1.28	0.402
1 st quintile	17	14	31	ref		
2 nd quintile	8	11	19	1.96	0.50; 7.76	0.336
3 rd quintile	17	24	41	1.99	0.50; 7.93	0.332
4 th quintile	14	11	25	0.53	0.08; 3.27	0.495
5 th quintile	18	14	32	0.47	0.07; 3.20	0.443

Unconditional logistic regression performed using mitochondrial copy number variable (Pfaffl) categorized in quintiles and analyzed as continuous variable log-transformed and continuous variable divided in quintiles (whereby the unit of measurement is a single quintile). Analyses were adjusted for sex, age, BMI, plate and recruitment center. Individual matching in conditional analysis was done by center, gender, age at recruitment (±6 months), date at entry in the cohort, time between blood sampling, and time of last consumption of food and drink (<3, 3-6, and ≥6 hours). This analysis was adjusted for plate and BMI.

Supplementary Table 5. Association between mtDNA copy number and pancreatic cancer stratified by smoking status.

	Controls	Cases	Total	OR	95% C.I.	P-value
Male smokers						
Continuous variable log-transformed	80	77	157	0.92	0.28; 2.99	0.887
Continuous variable expressed in quintiles	80	77	157	0.97	0.63; 1.49	0.892
1 st quintile	28	26	54	ref.		-
2 nd quintile	16	20	36	1.27	0.45; 3.54	0.650
3 rd quintile	21	19	40	1.00	0.37; 2.73	0.994
4 th quintile	9	6	15	0.56	0.08; 3.76	0.554
5 th quintile	6	6	12	1.03	0.10; 11.00	0.981
Female smokers						
Continuous variable log-transformed	54	65	119	0.06	0.04; 1.08	0.061
Continuous variable expressed in quintiles	54	65	119	0.68	0.40; 1.14	0.146
1 st quintile	10	16	26	ref.		-
2 nd quintile	17	20	37	0.48	0.14; 1.68	0.249
3 rd quintile	14	13	27	0.24	0.06; 1.08	0.063
4 th quintile	6	3	9	0.16	0.01; 1.74	0.132
5 th quintile	7	13	20	0.44	0.04; 4.59	0.490
Ever smokers (both sexes)						
Continuous variable log-transformed	134	142	276	0.79	0.33; 1.86	0.585
Continuous variable expressed in quintiles	134	142	276	0.94	0.70; 1.26	0.668
1 st quintile	38	42	80	ref.		-
2 nd quintile	33	40	73	0.99	0.48; 2.03	0.979
3 rd quintile	35	32	67	0.73	0.34; 1.56	0.416
4 th quintile	15	9	24	0.47	0.12; 1.76	0.261
5 th quintile	13	19	32	1.38	0.32; 5.89	0.662
Never smokers (both sexes)						
Continuous variable log-transformed	130	174	304	0.40	0.16; 0.97	0.043
Continuous variable expressed in quintiles	130	174	304	0.74	0.53; 1.05	0.094
1 st quintile	36	50	86	ref.		
2 nd quintile	29	46	75	0.87	0.35; 2.19	0.768
3 rd quintile	22	48	70	1.35	0.50; 3.65	0.553
4 th quintile	19	16	35	0.31	0.08; 1.22	0.093
5 th quintile	24	14	38	0.21	0.04; 1.00	0.050

Logistic regression performed using mitochondrial copy number variable (Pfaffl) categorized in quintiles and analyzed as continuous variable log-transformed, and continuous variable divided in quintiles (whereby the unit of measurement is a single quintile). Analyses were adjusted for age, BMI, plate and center of origin.

Supplementary Table 6. Association between mtDNA copy number and pancreatic cancer risk stratified by BMI.

	Controls	Cases	Total	OR	95% C.I.	P-value
Healthy weight (19<BMI<25)						
Continuous variable log-trasformed	141	183	324	0.25	0.09; 0.63	0.003
Continuous variable expressed in quintiles	141	183	324	0.58	0.42; 0.81	0.001
1 st quintile	25	47	72	ref.		
2 nd quintile	26	42	68	0.88	0.34; 2.24	0.783
3 rd quintile	30	42	72	0.34	0.12; 0.97	0.044
4 th quintile	33	28	61	0.13	0.04; 0.51	0.003
5 th quintile	27	24	51	0.12	0.03; 0.47	0.002
Overweight (25<BMI<30)						
Continuous variable log-trasformed	131	192	323	0.33	0.15; 0.74	0.007
Continuous variable expressed in quintiles	131	192	323	0.80	0.58; 1.10	0.171
1 st quintile	33	52	85	ref.		
2 nd quintile	26	38	64	0.67	0.29; 1.55	0.347
3 rd quintile	25	49	74	0.82	0.33; 2.04	0.671
4 th quintile	19	30	49	0.87	0.26; 2.94	0.825
5 th quintile	28	23	51	0.35	0.09; 1.30	0.116
Obese (BMI>30)						
Continuous variable log-trasformed	54	85	139	1.46	0.33; 6.45	0.621
Continuous variable expressed in quintiles	54	85	139	1.14	0.63; 2.08	0.658
1 st quintile	16	23	39	ref.		
2 nd quintile	13	24	37	0.58	0.14; 2.40	0.448
3 rd quintile	8	22	30	1.87	0.39; 8.94	0.435
4 th quintile	11	10	21	0.61	0.05; 7.23	0.697
5 th quintile	6	6	12	1.32	0.09; 20.03	0.840

Logistic regression stratified by BMI performed using mitochondrial copy number variable (Pfaffl) categorized in quintiles and analyzed as continuous variable log-transformed and continuous variable divided in quintiles (whereby the unit of measurement is a single quintile). Analyses were adjusted for sex, age, plate and center of origin. The number of underweight subjects (1 case and 10 controls) was too small to perform an analysis in this stratum.