



# Cognitive and behavioural processes in adolescents with social anxiety disorder

Eleanor Leigh<sup>a,\*</sup>, Ray Percy<sup>b</sup>, David M. Clark<sup>a</sup>, Cathy Creswell<sup>c</sup>, Polly Waite<sup>c</sup>

<sup>a</sup> Department of Experimental Psychology, University of Oxford, UK

<sup>b</sup> School of Psychology and Clinical Language Sciences, University of Reading, UK

<sup>c</sup> Department of Experimental Psychology and Department of Psychiatry, University of Oxford, UK

## ABSTRACT

**Background:** A better understanding of the processes that maintain social anxiety disorder (SAD) in adolescents could improve treatment outcomes. This study aimed to establish whether cognitive and behavioural processes known to be important in the maintenance of adult SAD are observed in adolescent populations and whether they are specific to SAD.

**Methods:** We recruited three groups of adolescents (12–18y): (1) 90 adolescents with a SAD diagnosis, (2) 58 adolescents with an anxiety disorder that was not SAD, and (3) 45 community-based adolescents. Participants completed measures of negative social cognitions, social attitudes, safety behaviours, self-focused attention, and social anxiety, anxiety, and depression symptoms.

**Results:** The clinical SAD sample endorsed higher levels of negative social cognitions, attitudes, and safety behaviours compared to both control groups. Self-focused attention was higher in the clinical SAD sample compared to the anxiety clinical control group but not compared to the community control group.

**Conclusions:** This study provided evidence of SAD-specific mechanisms including negative social cognitions, attitudes and safety behaviours in adolescents. The study did not provide evidence of disorder-specific mechanisms of self-focused attention but this may have been due to methodological limitations. These findings indicate avenues for further research and point to the potential value of Cognitive Therapy for the treatment of adolescent SAD.

## Funding information

Eleanor Leigh was funded by a Wellcome Trust Clinical Research Training Fellowship (102176/Z/13/Z) and is now funded by a MRC Clinician Scientist Fellowship (MR/W02389X/1). Polly Waite was funded by an NIHR Postdoctoral Research Fellowship (PDF-2016-09-092) and is now funded by an NIHR Development and Skills Enhancement award (NIHR302843). David M Clark was funded by the Wellcome Trust (WT069777). Cathy Creswell was funded by an NIHR Research Professorship (RP-2014-04-018) and receives support from the Oxford and Thames Valley NIHR Applied Research Collaboration and the Oxford Health NIHR Biomedical Research Centre. The study was sponsored by the University of Reading. For the purpose of Open Access, the author has applied a CC BY public copyright licence to any Author Accepted Manuscript version arising from this submission.

## 1. Introduction

Anxiety disorders are among the most common psychological disorders in young people, with global community prevalence rates of around 6.5% (Polanczyk, Salum, Sugaya, Caye, & Rohde, 2015). The

most common anxiety disorder in adolescents is social anxiety disorder (SAD), with lifetime prevalence rates of around 7% (Fehm, Pelissolo, Furmark, & Wittchen, 2005). SAD is characterised by persistent and disproportionate fear of social situations where embarrassment may occur. It is associated with an increased risk of other mental health problems, including other anxiety and depressive disorders, substance abuse and psychosis (Beesdo et al., 2007; Chartier, Walker, & Stein, 2003; Schutters et al., 2012). Adolescents with SAD also have an increased risk of leaving school early and educational underachievement (Vilaplana-Pérez et al., 2020). Furthermore, it has the lowest natural recovery rate of all anxiety disorders (Bruce et al., 2005).

We have effective treatments for SAD in adults. For example, Cognitive Therapy for SAD (CT-SAD) achieves high recovery rates (of up to 84%; Mayo-Wilson et al. (2014)) and is one of the first-line treatments recommended in the National Institute for Health and Care Excellence (2013) guidelines. The treatment was designed to reverse the specific cognitive and behavioural processes invoked in the cognitive model of Clark and Wells (1995). According to this model, social anxiety is driven by excessively negative beliefs about social interactions, which are maintained by self-focused attention, maladaptive safety behaviours, and negative self-imagery. There is substantial empirical support for the

\* Corresponding author.

E-mail address: [eleanor.leigh@psy.ox.ac.uk](mailto:eleanor.leigh@psy.ox.ac.uk) (E. Leigh).

<https://doi.org/10.1016/j.brat.2023.104416>

Received 24 May 2023; Received in revised form 30 September 2023; Accepted 9 October 2023

Available online 14 October 2023

0005-7967/© 2023 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

model in adult SAD (see Leigh and Clark (2018) for a summary).

SAD typically starts in adolescence (median 13 years), with most people developing the condition before they reach their twenties (Beesdo-Baum et al., 2012; Kessler et al., 2005). SAD-specific treatments such as those of Masia-Warner, Fisher, Shrout, Rathor, and Klein (2007) and Beidel et al. (2007) have been developed and show promising outcomes. However, these treatments do not show a strong focus on the psychological processes specified in the Clark and Wells' model, and direct evidence of superiority to generic CBT is currently lacking. In practice, adolescents with SAD and other common anxiety disorders are often treated in the same way, with a broad form of Cognitive Behaviour Therapy (CBT). Unfortunately the majority of children and adolescents with SAD continue to meet diagnostic criteria after treatment with broad-based CBT (Evans, Clark, & Leigh, 2021). For example, in one US RCT of generic CBT, recovery among those with SAD was 41% compared to 72% for those with other anxiety disorders (Ginsburg et al., 2011). Therefore, the question arises as to whether the Clark & Wells model is relevant to adolescents, because this would indicate that CT-SAD may be a useful treatment approach. This cannot be assumed given the distinctive neurocognitive, emotional, and social factors in operation at this stage of life (Blakemore, 2008) and many of the processes invoked in the Clark & Wells model are underpinned by cognitive and emotional factors still in development. For example, self-consciousness develops through adolescence and is intimately linked to self-focused attention (Rankin, Lane, Gibbons, & Gerrard, 2004). Addressing this question has the potential to improve outcomes for youth with SAD. Two clinical trials, both of which have compared a broad-based form of CBT with a version of SAD-specific CBT, are relevant (Rapee et al., 2023; Spence, Donovan, March, Kenardy, & Hearn, 2017). Both trials developed an SAD-specific version of CBT that included social skills training, attention training, reduction of safety behaviours, cognitive challenging and coping skills, with neither demonstrating differences between treatment arms at post-treatment or follow-up. Whilst it is possible that this indirectly counters the idea that these processes are implicated in SAD in youth, as suggested by Rapee et al. (2023) and Leigh (2023) it seems more likely that it is because the augmented CBT tested in these two trials does not maximize opportunities to modify maintenance processes.

A number of empirical studies with community samples have tested whether the components of the Clark & Wells model are relevant to adolescents. Cross-sectional observational studies such as those of Hodson, McManus, Clark, and Doll (2008) and Schreiber, Höfling, Stangier, Bohn, and Steil (2012) support an association between negative thoughts, safety behaviours, negative imagery, and self-focused attention and social anxiety symptoms in adolescents (see Leigh and Clark (2018) for a review), and more recently prospective and experimental studies have pointed to a causal role for these processes (Chiu, Clark, & Leigh, 2021; Leigh et al., 2020, 2021). Taken together the findings indicate that targeting these processes in treatment may be beneficial in reducing social anxiety in youth. However, so far, studies have been largely carried out with community rather than clinical populations. Whilst a small number of studies have been undertaken with clinical samples of adolescents (see Leigh and Clark (2018) for a review), to our knowledge none have examined the various processes of interest simultaneously or their specificity to SAD. Therefore, this study aimed to establish whether key cognitive and behavioural processes specified in the Clark and Wells (1995) model are observed in clinical populations and whether they are specific to SAD. To do this, we compared a clinical sample of adolescents with SAD with a clinical sample of adolescents with a non-SAD anxiety disorder and with a community sample of adolescents.

We hypothesized that adolescents with SAD would show significantly higher levels of (i) negative social cognitions and attitudes, (ii) self-focused attention, and (iii) safety behaviours, compared to adolescents with other anxiety disorders and with community adolescents.

## 2. Methods

### 2.1. Study design

The present study was cross-sectional and comprised five self-report questionnaires administered to three groups of adolescents. Questionnaires were selected to measure symptoms of social anxiety, symptoms of anxiety and depression, social cognitions and attitudes, self-focused attention, and safety behaviours.

### 2.2. Participants

Three groups of participants were recruited for the study. The first group ('SAD' group) consisted of 90 adolescents with a SAD diagnosis, either as their primary or secondary diagnosis (if secondary, the primary disorder was required to be another anxiety disorder). The second group ('Anxiety Control' group) consisted of 58 adolescents with one or more anxiety disorder that was not SAD (i.e., Generalised Anxiety Disorder, Separation Anxiety Disorder, a Specific Phobia, Panic Disorder or Agoraphobia). The SAD and Anxiety Control groups were recruited through a clinical service. The adolescent was required to meet diagnostic criteria for SAD or another anxiety disorder using the ADIS-CP; Silverman & Albano, 1996) and an anxiety disorder had to be identified as the young person's primary problem. The third group ('Community Control' group) consisted of 45 adolescents recruited through schools. Due to resourcing limitations, we were not able to conduct diagnostic assessments with this group. Participants in all three groups were required to be aged between 12 and 18 years. They were not eligible to take part if they had an intellectual/learning disability (based on school/clinic/parent information) that may interfere with their ability to read and understand the questions.

### 2.3. Procedure

The study procedures were in accordance with ethical standards and were approved by the National Research Ethics Committee South Central – Berkshire B (reference: 15/SC/0081) and the University of Reading Research Ethics Committee (reference: 15/27). Young people aged 12–15 years provided written assent and their parents/carers provided written consent, while those aged 16–18 years provided written consent.

The two groups with a diagnosed anxiety disorder (SAD and Anxiety Control groups) were recruited through a local NHS-commissioned child and adolescent mental health service between December 2015 and February 2020. To be eligible for referral to the service, young people had to be experiencing symptoms of anxiety and/or depression at a level that was causing distress and/or interference (i.e., likely to meet diagnostic criteria). However, they were not eligible if they had an established autistic spectrum disorder, were taking medication for the treatment of anxiety/depression, had immediate suicidal intent, or had been identified by social care as currently 'at risk' due to significant child protection concerns. For those referred, if the referral indicated that they had symptoms of anxiety and were likely to meet the study eligibility criteria, the adolescent and their parent/carer were sent the study information sheets. They then undertook a diagnostic assessment, which involved the young person (and where possible, the parent/carer) being interviewed separately. All assessments were carried out by trained assessors (either clinicians working in the service or placement students, under the supervision of an experienced clinician). All assessors were trained to high levels of reliability and received supervision for every assessment from a clinical psychologist (or equivalent) with extensive experience of delivering and supervising diagnostic assessments and proven reliability. If the diagnostic assessment confirmed that the young person had an anxiety disorder as the main presenting problem, the study was then discussed with them at the subsequent clinical appointment, and they were invited to take part. If the

adolescent and their parent agreed, they were then given the set of self-report measures to complete. This could be done as a hard copy or online, through a secure system.

The Community Control group was recruited through schools between March and August 2016. Head teachers were asked to assist with recruitment and if they agreed, the study was advertised through school newsletters and/or invitation letters sent out to students and their parents/carers. For those who indicated they were interested in taking part, adolescents and their parents/carers were sent information sheets about the study. They were then contacted by the research team to address any questions they might have about the study and, if they wished to take part, complete consent/assent forms. Questionnaires could be completed on paper or online. At the end of the questionnaires, participants were told that if they had any concerns or worries to get in touch (or ask their parent/guardian to get in touch) with the research team and were also given further information about support, advice, and resources. Participants in this group were given a £10 gift voucher to reimburse them for their time.

## 2.4. Measures

The measures were given to all participants, except for the diagnostic assessment which was only administered to the SAD and Anxiety Control groups.

**Diagnostic assessment.** The diagnostic assessment was conducted to determine eligibility to the clinical service and the study. It used sections from two validated semi-structured interview schedules to determine whether the young person met diagnostic criteria for an anxiety/depressive disorder; the Anxiety Disorders Interview Schedule (ADIS-IV-C/P; Silverman et al., 1996) (adapted for DSM-5) and the Kiddie Schedule for Affective Disorders and Schizophrenia Present and Lifetime (K-SAD-S-PL; Kaufman et al., 1997). The ADIS was used to assess DSM-5 diagnoses of anxiety and externalising disorders in children and adolescents. It has excellent interrater reliability (Lyneham et al., 2007) and concurrent validity (Wood et al., 2002). In the current study, interrater reliability was good overall (see Supplementary Information for further details). The K-SAD-S-PL was used to assess DSM-V Major and Persistent Depressive Disorder and Mania. It has good concurrent validity of screens and K-SAD-S-PL diagnoses (Kaufman et al., 1997). Both interrater reliability and test-retest reliability are excellent (Kaufman et al., 1997), suggesting that the K-SAD-S-PL produces both reliable and valid psychiatric diagnoses. Using information from both the young person and their parent/carer, the clinician assigns a Clinical Severity Ratings (CSRs) to diagnoses (as is standard practice with the ADIS) using a nine-point scale (0–8), with ‘0’ indicating no impairment and ‘8’ indicating severe impairment. Impairment is defined as affecting the young person’s life and/or creating significant distress. A CSR  $\geq 4$  signifies a clinical diagnosis. The disorder with the highest CSR was allocated as the primary diagnosis.

**Demographic information.** For all groups, information about the young person (age, gender, ethnicity) was collected.

**Anxiety and depression symptoms.** The Revised Child Anxiety and Depression Scale (RCADS; Chorpita, Yim, Moffitt, Umemoto, and Francis (2000)) is a self-report measure that assesses symptoms of separation anxiety disorder, social anxiety disorder, generalised anxiety disorder, panic disorder, obsessive compulsive disorder and major depressive disorder. There are 47 items and respondents are asked to rate how often each item applies on a scale of 0 (‘never’) to 3 (‘always’). It has been shown to have robust psychometric properties in children and young people from 7 to 18 years of age (Chorpita, Moffitt, & Gray, 2005). For the current study, we used the total RCADS score and all subscale scores. The internal consistency for the total score in our clinical sample (SAD and Anxiety Control participants) and community sample was excellent (clinical sample  $\alpha = 0.91$ ; community sample  $\alpha = 0.95$ ). The internal consistency was excellent ( $\alpha > 0.90$ ) for all subscales in the clinical and community samples.

**Social anxiety symptoms.** The Liebowitz Social Anxiety Scale for Children and Adolescents (LSAS-C/A; Masia, Klein, & Liebowitz, 1999) is a measure of social anxiety symptoms that was used in self-report version for the present study. It includes 24 items, rated on a scale from 0 (‘none’) to 3 (‘severe’) for fear and avoidance of social interaction and performance. The total score includes both the fear and avoidance ratings and scores range from 0 to 144. It has well established psychometric properties when administered to children and young people from 7 to 18 years of age (Leigh & Clark, 2021b; Masia-Warner et al., 2003). The internal consistency for both samples was excellent (clinical sample  $\alpha = 0.98$ ; community sample  $\alpha = 0.98$ ).

**Social cognitions.** The Child & Adolescent Social Cognitions Questionnaire (CASCQ; Leigh & Clark, 2021a) is a self-report questionnaire that assesses common social anxiety-related cognitions and was adapted from the adult Social Cognitions Questionnaire (Clark, 2003), for example “People will stare at me” and “People think I am boring”. There are 27 items and respondents are asked firstly to rate how frequently they experience each thought when they are nervous or frightened, from 1 (‘thought never occurs’) to 5 (‘thought always occurs when I am nervous’). They are then asked to rate how much they believe the thought, from 0 (‘I do not believe this thought’) to 100 (‘I am completely convinced this thought is true’). A mean frequency and belief score is calculated. The measure has good internal consistency and convergent validity in an adolescent community sample (Leigh & Clark, 2021a). The internal consistency for both samples was excellent (clinical sample  $\alpha = 0.97$ ; community sample  $\alpha = 0.97$ ) for both frequency and belief ratings.

**Social attitudes.** The Child & Adolescent Social Attitudes Questionnaire (CASAQ) is a self-report questionnaire that assesses common social anxiety-related attitudes or beliefs. There are 52 items and respondents are asked how much they agree or disagree with each statement on a 7-point scale from ‘totally agree’ to ‘totally disagree’. A mean CASAQ score is calculated, with lower scores indicating a more negative attitude. The measure was originally developed for adults and has good internal consistency and discriminant validity (Clark, 2003). The internal consistency for both samples was excellent (clinical sample  $\alpha = 0.92$ ; community sample  $\alpha = 0.93$ ).

**Self-focused attention.** Two items from the Child and Adolescent Social Summary Weekly Rating Scale (CASSWRS) (Clark, 2003) were used to measure self-focused attention over the previous week. One item assessed self-focused attention for social situations that the participant found difficult (“For social situations that you found difficult, please choose a number from the scale below to show how much your mind was focused on yourself (‘self-focused’) or on other people and what was going on around you (‘externally focused’) in the last week”). The other item assessed self-focused attention for social situations in general (“For social situations in general, please choose a number from the scale below to show how much your mind was focused on yourself (‘self-focused’) or on other people and what was going on around you (‘externally focused’) in the last week”). Both items are scored from 0 (‘entirely externally focused’) to 8 (‘entirely self-focused’), and a mean is obtained. The internal consistency for our clinical sample was acceptable ( $\alpha = 0.76$ ) but for our community sample it was questionable ( $\alpha = 0.65$ ).

**Social behaviours.** The Child and Adolescent Social Behaviour Questionnaire (CASBQ; Evans, Chiu, Clark, Waite, & Leigh, 2021) is a self-report questionnaire consisting of 28 items that reflect a range of safety behaviours in social situations. For each item, the respondent is asked to rate how often they undertake the behaviour when they are anxious in or before a social situation, from 0 (‘never’) to 3 (‘always’). An overall CASBQ score is calculated as a mean of responses, with a range of 0 (no safety behaviours) to 3 (very frequent safety behaviours). The original version of the questionnaire designed for use with adults (Social Behaviour Questionnaire; Clark (2003)) has good psychometrics properties (Clark, 2005). Sound psychometric properties of the youth version have been found with adolescents (Evans, Chiu, Clark, Waite, & Leigh, 2021). The internal consistency was good in our clinical sample

( $\alpha = 0.89$ ) and excellent in our community sample ( $\alpha = 0.92$ ).

## 2.5. Analysis plan

Analysis was undertaken in SPSS and R version 2022.02.3. Mean substitution was performed when less than 10% of items were missing in each questionnaire. Questionnaires with more than 10% of missing items were treated as missing variables at participant level with complete cases used in the analysis. For the clinical sample, at variable level, 5% of data was missing for the RCADS total, 8% for the LSAS-CA, 8% for the CASCQ-F, 11% for the CASCQ-B, 5% for the CASBQ, 9% for the CASAQ and 7% for the CASPWSS. For the community sample, 4% of data was missing for the RCADS total, 0% for the LSAS-CA, 2% for the CASCQ-F, 2% for the CASCQ-B, 2% for the CASBQ, 0% for the CASAQ and 0% for the CASPWSS.

Descriptive statistics are reported. A series of ANCOVA were undertaken with Group (SAD vs. Anxiety Control and SAD vs. Community Control) as between-subjects factors, and age and gender as covariates were undertaken to examine differences in symptom scores and process scores. We ran separate ANCOVAs to compare the SAD group to the Anxiety Control group and then the SAD group to the Community Control group because we did not compare the two control groups directly. We report generalised eta-square as an estimate of the effect size.

## 3. Results

### 3.1. Descriptive statistics

For the 90 participants in the SAD group, this was the primary problem for 62% of participants [ $n = 56$ ] and the secondary problem for 38% [ $n = 34$ ] (amongst these participants, GAD was the most common primary problem). 90.47% of the SAD group were White British. For the 58 participants in the Anxiety Control group, the three most common primary diagnoses were: GAD ( $n = 19$ ; 33%), Separation Anxiety Disorder ( $n = 13$ ; 22%), and specific phobia ( $n = 13$ ; 22%). 79.31% of this group were White British. Data on the ethnicity of the Community sample was not available.

Table 1 provides demographic and clinical information on the participants. Comparing the SAD group and Community Control group, the SAD group had significantly more girls ( $\chi^2(1) = 17.60, p < .001$ ) and was significantly younger ( $t(133) = 2.00, p = .024$ ). The SAD and Anxiety Control groups did not significantly differ in terms of gender ( $\chi^2(1) = 0.34, p = .560$ ), but the SAD group was significantly older than the Anxiety Control group ( $t(146) = 4.14, p < .001$ ).

### 3.2. Symptom measures: group comparisons

Symptom measures for the SAD, Anxiety Control, and Community Control groups are shown in Table 2 along with the ANCOVA test statistics. The SAD group scored significantly higher than the Anxiety Control group on both measures of social anxiety symptoms and on the RCADS total scale, with no differences between the groups on the other subscales of the RCADS. The SAD group scored significantly higher than the Community Control group on all symptom measures.<sup>1 2</sup>

<sup>1</sup> Results remained unchanged when we applied the Benjamini Hochberg method with a 10% false discovery rate to control for Type 1 errors due to multiple comparisons.

<sup>2</sup> As a sensitivity analysis, analyses were rerun with a primary SAD only SAD group (i.e., participants with a secondary diagnosis of SAD were excluded). Results are reported in the Supplementary Information.

**Table 1**

Descriptive statistics for SAD and anxiety control clinical groups and community control group.

|                                 | SAD          | Anxiety control | Community control |
|---------------------------------|--------------|-----------------|-------------------|
| Total N                         | 90           | 58              | 45                |
| Gender, n (%)                   |              |                 |                   |
| Girls/young women               | 69 (76.7%)   | 42 (72.4%)      | 18 (40.0%)        |
| Boys/young men                  | 21 (23.3%)   | 16 (27.6%)      | 27 (60.0%)        |
| Non-binary                      | 0            | 0               | 0                 |
| Age Mean (SD)                   | 14.06 (2.27) | 12.40 (2.73)    | 14.89 (2.30)      |
| Primary anxiety disorder, n (%) |              |                 |                   |
| SAD                             | 56 (62.2%)   | 0               | –                 |
| GAD                             | 26 (28.9%)   | 19 (32.8%)      | –                 |
| Separation anxiety disorder     | 3 (3.3%)     | 13 (22.4%)      | –                 |
| Specific phobia                 | 2 (2.2%)     | 13 (22.4%)      | –                 |
| Panic disorder                  | 1 (1.1%)     | 6 (10.3%)       | –                 |
| Agoraphobia                     | 1 (1.1%)     | 5 (8.6%)        | –                 |
| Illness anxiety disorder        | 1 (1.1%)     | 0               | –                 |
| AD-NOS                          | 0            | 2 (3.4%)        | –                 |
| Presence of diagnoses, n (%)    |              |                 |                   |
| SAD                             | 90 (100%)    | 0               | –                 |
| GAD                             | 70 (77.8%)   | 30 (51.7%)      | –                 |
| Separation anxiety disorder     | 15 (16.7%)   | 18 (31.0%)      | –                 |
| Specific phobia                 | 13 (14.4%)   | 18 (31.0%)      | –                 |
| Panic disorder                  | 10 (11.1%)   | 7 (12.1%)       | –                 |
| Agoraphobia                     | 10 (11.1%)   | 9 (15.5%)       | –                 |
| Illness anxiety disorder        | 1 (1.1%)     | 0               | –                 |
| AD-NOS                          | 0            | 3 (5.2%)        | –                 |
| OCD                             | 2 (2.2%)     | 1 (1.7%)        | –                 |
| PTSD                            | 1 (1.1%)     | 0               | –                 |
| Dysthymia                       | 3 (3.3%)     | 0               | –                 |
| MDD                             | 5 (5.6%)     | 3 (5.2%)        | –                 |
| ADIS reporter, n (%)            |              |                 |                   |
| Adolescent + parent report      | 81 (90.0%)   | 54 (93.1%)      | –                 |
| Adolescent-only report          | 9 (10.0%)    | 4 (6.9%)        | –                 |

Notes: AD-NOS = anxiety disorder – not otherwise specified; GAD = generalised anxiety disorder; MDD = major depressive disorder; OCD = obsessive compulsive disorder; PTSD = post-traumatic stress disorder; SAD = social anxiety disorder; SD = standard deviation.

### 3.3. Process measures: group comparisons

Mean and standard deviations for the cognitive and behavioural process measures in the SAD, Anxiety Control, and Community Control groups are shown in Tables 2 and in Fig. 1.

The ANCOVA test statistics are shown in Table 2. As can be seen, measures of negative social cognitions (CASCQ both frequency and belief ratings) and safety behaviours (CASBQ) were significantly higher in the SAD group compared to both the Anxiety and Community Control samples. Similarly, social attitudes scores (CASAQ) were significantly lower (more negative) in the SAD group compared to both the Anxiety and Community Control samples. Self-focused attention was significantly higher in the SAD group compared to the Anxiety Control group but not the Community Control group.<sup>1, 2 and 3</sup>

## 4. Discussion

This study aimed to examine whether key cognitive and behavioural processes specified in the Clark and Wells (1995) model of SAD (in adults) are observed in adolescents. It builds on existing studies by examining this in clinical and community populations and determining

<sup>3</sup> ANCOVAs were rerun with depression symptoms (as measured by the child reported RCADS depression subscale) as an additional covariate. There were no differences in the pattern of results.




**Table 2**

Symptom and process measure scores for SAD and Anxiety Control Clinic Samples and Community Sample.

|                         | SAD Mean (SD)        | Anxiety Controls Mean (SD) | Statistical Test SAD vs Anxiety Controls <sup>†</sup> | $\eta^2G$ | Community Controls Mean (SD) | Statistical Test SAD vs Community Controls <sup>†</sup> | $\eta^2G$ |
|-------------------------|----------------------|----------------------------|---|-----------|------------------------------|---|-----------|
| <b>Symptom Measures</b> |                      |                            |   |           |                              |   |           |
| RCADS-Total             | <b>67.33 (22.17)</b> | 53.58 (21.02)              | $F(1, 132) = 8.33, p < .01$                           | 0.059     | 35.36 (20.35)                | $F(1, 124) = 41.25, p < .001$                           | 0.250     |
| RCADS-MDD               | <b>13.35 (5.38)</b>  | 10.45 (6.22)               | $F(1, 132) = 2.26, p = .135$                          | 0.017     | 7.95 (5.76)                  | $F(1, 124) = 20.56, p < .001$                           | 0.142     |
| RCADS-GAD               | <b>9.32 (3.63)</b>   | 8.76 (3.89)                | $F(1, 131) = 0.46, p = .497$                          | 0.004     | 5.74 (3.33)                  | $F(1, 123) = 15.43, p < .001$                           | 0.111     |
| RCADS-SEP               | <b>8.02 (4.09)</b>   | 7.78 (4.86)                | $F(1, 131) = 2.33, p = .129$                          | 0.017     | 2.37 (2.41)                  | $F(1, 123) = 45.00, p < .001$                           | 0.268     |
| RCADS-PD                | <b>11.53 (5.92)</b>  | 10.12 (6.06)               | $F(1, 132) = 0.25, p = .621$                          | 0.002     | 4.64 (4.36)                  | $F(1, 124) = 27.45, p < .001$                           | 0.181     |
| RCADS-OCD               | <b>5.64 (3.64)</b>   | 5.59 (3.38)                | $F(1, 131) = 0.21, p = .648$                          | 0.002     | 3.84 (3.24)                  | $F(1, 123) = 4.68, p < .05$                             | 0.037     |
| RCADS-SAD               | <b>19.32 (6.57)</b>  | 10.881 (6.65)              | $F(1, 132) = 35.61, p < .001$                         | 0.212     | 10.86 (6.20)                 | $F(1, 124) = 38.71, p < .001$                           | 0.238     |
| LSAS-CA                 | <b>87.92 (30.88)</b> | 39.35 (29.30)              | $F(1, 130) = 62.78, p < .001$                         | 0.326     | 50.70 (35.18)                | $F(1, 125) = 33.81, p < .001$                           | 0.213     |
| <b>Process Measures</b> |                      |                            |   |           |                              |   |           |
| CASCQ-F                 | <b>2.95 (0.85)</b>   | 1.96 (0.91)                | $F(1, 131) = 27.92, p < .001$                         | 0.172     | 2.22 (0.90)                  | $F(1, 122) = 11.85, p < .001$                           | 0.089     |
| CASCQ-B                 | <b>47.69 (21.54)</b> | 21.87 (21.40)              | $F(1, 125) = 32.59, p < .001$                         | 0.207     | 27.60 (24.35)                | $F(1, 119) = 14.96, p < .001$                           | 0.112     |
| CASAQ                   | <b>2.18 (0.83)</b>   | 3.29 (1.13)                | $F(1, 103) = 24.52, p < .001$                         | 0.192     | 3.27 (1.09)                  | $F(1, 113) = 24.01, p < .001$                           | 0.175     |
| SFA                     | <b>4.07 (1.87)</b>   | 3.27 (1.73)                | $F(1, 131) = 2.99, p < .05$                           | 0.032     | 4.10 (1.42)                  | $F(1, 123) = 0.11, p = .744$                            | 0.001     |
| CASBQ                   | <b>1.39 (0.43)</b>   | 0.83 (0.58)                | $F(1, 134) = 27.31, p < .001$                         | 0.169     | 1.03 (0.50)                  | $F(1, 125) = 15.61, p < .001$                           | 0.111     |

Notes: CASAQ = Child & Adolescent Social Attitudes Questionnaire; CASCQ-F = Child & Adolescent Social Cognitions Questionnaire - Frequency; CASCQ-B = Child & Adolescent Social Cognitions Questionnaire - Belief; CASAQ = Child & Adolescent Social Attitudes Questionnaires; SFA = Child and Adolescent Social Summary Weekly Rating Scale – Self-Focused Attention items; LSAS-CA The Liebowitz Social Anxiety Scale for Children and Adolescents; RCADS-Total = The Revised Child Anxiety and Depression Scale – Total Score; RCADS-SAD = The Revised Child Anxiety and Depression Scale – Social Anxiety Disorder Subscale Score; SAD = social anxiety disorder; MDD = depression; SEP = separation anxiety disorder; GAD = generalised anxiety disorder; PD = panic disorder; OCD = obsessive compulsive disorder; SD = standard deviation;  $\eta^2G$  = generalised eta squared; <sup>†</sup> One way ANCOVA controlling for age and sex.

**Fig. 1.** Process Measure Scores for SAD Clinical Group, Anxiety Control Group and Community Control Group (means with 95% Confidence Intervals displayed).

the extent to which the processes are specific to SAD. Consistent with our hypotheses, we found that adolescents with SAD showed significantly more negative social cognitions and social attitudes and more safety behaviours compared to adolescents with other anxiety disorders and with community adolescents. Self-focused attention was significantly

higher in the SAD group than the anxiety control group, but contrary to our hypothesis it was not significantly higher in the SAD group compared to the community control group.

Our finding that adolescents with SAD showed significantly more negative social cognitions, social attitudes and more safety behaviours

compared to adolescents with other anxiety disorders and with community adolescents is consistent with previous literature and promising findings from recent clinical trials using treatments based on Clark and Wells' cognitive model. Studies with mainly community-based adolescents have shown associations between negative thoughts, safety behaviours, and self-focused attention, and social anxiety symptoms using cross-sectional (e.g., Hodson et al. (2008); Schreiber et al. (2012)) as well as prospective and experimental designs (Chiu et al., 2021; Leigh et al., 2020, 2021). Taken together, these findings indicate that targeting these processes in treatment for adolescent SAD is likely to be beneficial. This is indeed borne out by the findings of two trials (Ingul, Aune, & Nordahl, 2014; Leigh & Clark, 2023), which have demonstrated that CT-SAD is associated with large controlled treatment effects and furthermore the trial of Leigh & Clark (2022) found that the beneficial effect of CT-SAD was mediated by changes in these processes.

Although the SAD group endorsed more self-focused attention in social situations compared to the Anxiety Control group, we were surprised to find that there was no significant difference between the SAD group and the Community group. A number of studies with community samples have reported an association between self-focused attention and social anxiety symptoms (Blote, Miers, Heyne, Clark, & Westenberg, 2014; Chiu et al., 2021; Hodson et al., 2008; Leigh et al., 2021; Schreiber et al., 2012). It may be that there is no difference in self-focused attention between these groups, or it could be that it is a mechanism relevant across anxiety disorders more broadly. Alternatively, the failure to find an effect may in part be methodological. We created an average of two items (taken from the SPWSS) to measure self-focused attention. However, the internal consistency for the measure in the Community group was below the accepted level of 0.7 (Nunnally & Bernstein, 1978). Measures with one or two items are prone to random measurement errors and may underestimate associations and so it is possible that this methodological limitation explains the finding. Use of a multi-item measure such as the Focus of Attention Questionnaire (Woody, Chambless, & Glass, 1997) to capture self-focused attention will increase reliability and mitigate against this limitation in future studies.

The findings add to the existing evidence base and suggests that not only are these cognitive and behavioural processes sensitive markers of SAD but they are also specific SAD process variables. With this study we addressed what Garber and Hollon (1991) called the question of broad specificity, in that we examined whether these cognitive and behavioural processes could differentiate young people with SAD from those with a range of other anxiety problems. It will be informative to now examine the question of narrow specificity, or in other words whether the processes can differentiate SAD from other specific anxiety disorders such as compared to those with generalised anxiety disorder or panic disorder.

Our findings indicate that the core processes that are implicated in the maintenance of adult SAD are also relevant to adolescents. However we note the broad age range assessed in the present study which precludes a fine-grained analysis of how these processes vary across the adolescent period. But this would be a valuable avenue of future research because recent studies suggest that there may be differences. For example, it was found that the types of safety behaviours used differed between younger and older adolescents, with increasing use of more cognitively sophisticated behaviours with age (Evans, Chiu, et al., 2021).

The findings have important clinical implications. They help us to understand why outcomes for adolescents with SAD from traditional broad-based anxiety treatments may be disappointing and point to the value of disorder specific interventions for this population. They also indicate that we may achieve better outcomes by carefully targeting these processes in therapy and this is best done in an integrated fashion (Leigh, 2023). One such treatment is Cognitive Therapy. Cognitive Therapy is comprised of a series of techniques addressing negative social cognitions and beliefs, self-focused attention, and unhelpful safety behaviours. Core techniques include video feedback and behavioural

experiments. In support of this assertion, two randomised controlled trials with adolescents with SAD have found the treatment to be associated with large controlled effect sizes (Ingul et al., 2014; Leigh & Clark, 2023).

This study has strengths. It examines maintenance mechanisms in clinical samples that allow us to understand the extent to which they are distinct to SAD. Adolescents in the clinical groups were all referred for treatment within routine services and all received a gold standard diagnostic assessment, conducted by assessors with high levels of training and supervision. In terms of limitations, we were not able to carry out diagnostic assessments with the group of adolescents recruited from the community. Although, as expected, the group's mean scores on all symptom measures were significantly lower than those of the SAD group, it is possible that this group included young people who met diagnostic criteria for an anxiety disorder and potentially other disorders. However, the fact that we found the differences between the groups with this more conservative bar makes the findings potentially more robust. Because we wanted to examine the extent to which differences between the two clinical groups was accounted for by SAD, our Anxiety Control group did not include anyone with SAD. It is important to note that as a result this group is not a representative clinical sample because SAD is common as both a primary and secondary anxiety disorder in adolescents.

We note minor demographic differences between groups; the SAD group included significantly more girls than the Community Control group. On average the SAD group was significantly older than the Anxiety Control group and younger than the Community Control group. However, both variables were controlled for in the analysis. We were able to collect ethnicity data for our two clinical groups, and it is important to acknowledge that the SAD sample was predominantly White British and therefore findings may not be generalisable to more diverse groups. In addition, we were not able to collect ethnicity data from the community control group and therefore there may be important differences that are not accounted for.

To conclude, the findings from this study establish that key cognitive and behavioural processes specified in the Clark and Wells (1995) model are observed in adolescent clinical populations and support the suggestion that they are specific to SAD. Historically, adolescent SAD has been associated with poorer treatment outcomes. Our findings add to the growing evidence base that suggests that interventions that single-mindedly target these cognitive and behavioural processes may yield better outcomes for this population.

#### CRediT authorship contribution statement

**Eleanor Leigh:** Conceptualization, Methodology, Formal analysis, Writing – original draft. **Ray Percy:** Investigation, Writing – review & editing. **David M. Clark:** Conceptualization, Methodology, Writing – review & editing. **Cathy Creswell:** Conceptualization, Methodology, Writing – review & editing. **Polly Waite:** Conceptualization, Methodology, Formal analysis, Writing – original draft.

#### Declaration of competing interest

None

#### Data availability

Data will be made available on request.

#### Acknowledgements

We would like to thank the young people who took part in the study, as well as clinicians, staff and students from the AnDY Research Clinic at the University of Reading who helped in collecting the data. We would also like to thank the young people who provided feedback on the

measures. The views expressed in this publication are those of the author and not necessarily those of the NIHR or the Department of Health and Social Care. The materials for the study can be accessed by contacting the corresponding author. All questionnaires are freely available to download from [oxcadatresources.com](https://oxcadatresources.com).

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.brat.2023.104416>.

## References

- Beesdo, K., Bittner, A., Pine, D. S., Stein, M. B., Höfler, M., Lieb, R., et al. (2007). Incidence of social anxiety disorder and the consistent risk for secondary depression in the first three decades of life. *Archives of General Psychiatry*, 64(8), 903–912. <https://doi.org/10.1001/archpsyc.64.8.903>
- Beesdo-Baum, K., Knappe, S., Fehm, L., Höfler, M., Lieb, R., Hofmann, S. G., et al. (2012). The natural course of social anxiety disorder among adolescents and young adults. *Acta Psychiatrica Scandinavica*, 126(6), 411–425. <https://doi.org/10.1111/j.1600-0447.2012.01886.x>
- Beidel, D. C., Turner, S. M., Sallee, F. R., Ammerman, R. T., Crosby, L. A., & Pathak, S. (2007). SET-C versus fluoxetine in the treatment of childhood social phobia [Article]. *Journal of the American Academy of Child & Adolescent Psychiatry*, 46(12), 1622–1632. <https://doi.org/10.1097/chi.0b013e318154bb57>
- Blakemore, S. J. (2008). Development of the social brain during adolescence. *Quarterly Journal of Experimental Psychology*, 1, 40–49. <https://doi.org/10.1080/17470210701508715>
- Blote, A. W., Miers, A. C., Heyne, D. A., Clark, D. M., & Westenberg, P. M. (2014). The relation between social anxiety and audience perception: Examining Clark and Wells' (1995) model among adolescents. *Behavioural and Cognitive Psychotherapy*, 42(5), 555–567. <https://doi.org/10.1017/s1352465813000271>
- Bruce, S. E., Yonkers, K. A., Otto, M. W., Eisen, J. L., Weisberg, R. B., Pagano, M., et al. (2005). Influence of psychiatric comorbidity on recovery and recurrence in generalized anxiety disorder, social phobia, and panic disorder: A 12-year prospective study. *American Journal of Psychiatry*, 162(6), 1179–1187. <https://doi.org/10.1176/appi.ajp.162.6.1179>
- Chartier, M. J., Walker, J. R., & Stein, M. B. (2003). Considering comorbidity in social phobia. *Social Psychiatry and Psychiatric Epidemiology: The International Journal for Research in Social and Genetic Epidemiology and Mental Health Services*, 38(12), 728–734. <https://doi.org/10.1007/s00127-003-0720-6>
- Chiu, K., Clark, D. M., & Leigh, E. (2021). Cognitive predictors of adolescent social anxiety. *Behaviour Research and Therapy*, 137, Article 103801. <https://doi.org/10.1016/j.brat.2020.103801>
- Chorpita, B. F., Moffitt, C. E., & Gray, J. (2005). Psychometric properties of the revised child anxiety and depression scale in a clinical sample. *Behaviour Research and Therapy*, 43(3), 309–322. <https://doi.org/10.1016/j.brat.2004.02.004>
- Chorpita, B. F., Yim, L., Moffitt, C., Umemoto, L. A., & Francis, S. E. (2000). Assessment of symptoms of DSM-IV anxiety and depression in children: A revised child anxiety and depression scale. *Behaviour Research and Therapy*, 38(8), 835–855. <https://doi.org/10.1016/j.brat.2004.02.004>
- Clark, D. M. (2003). *Social anxiety process measures*. <https://oxcadatresources.com>.
- Clark, D. M., & Wells, A. (1995). A cognitive model of social phobia. In G. Heimberg, M. R. M. R. Liebowitz, D. Hope, & F. Scheier (Eds.), *Social phobia: Diagnosis, assessment, and treatment* (pp. 69–93). The Guilford Press.
- Evans, R., Chiu, K., Clark, D. M., Waite, P., & Leigh, E. (2021). Safety behaviours in social anxiety: An examination across adolescence. *Behaviour Research and Therapy*, Article 103931. <https://doi.org/10.1016/j.brat.2021.103931>
- Evans, R., Clark, D. M., & Leigh, E. (2021). Are young people with primary social anxiety disorder less likely to recover following generic CBT compared to young people with other primary anxiety disorders? A systematic review and meta-analysis. *Behavioural and Cognitive Psychotherapy*, 49(3), 352–369. <https://doi.org/10.1017/S135246582000079X>
- Fehm, L., Pelissolo, A., Furmark, T., & Wittchen, H.-U. (2005). Size and burden of social phobia in Europe. *European Neuropsychopharmacology*, 15(4), 453–462. <https://doi.org/10.1016/j.euroneuro.2005.04.002>
- Garber, J., & Hollon, S. D. (1991). What can specificity designs say about causality in psychopathology research? *Psychological Bulletin*, 110, 129–136. <https://doi.org/10.1037/0033-2909.110.1.129>
- Ginsburg, G. S., Kendall, P. C., Sakolsky, D., Compton, S. N., Piacentini, J., Albano, A. M., et al. (2011). Remission after acute treatment in children and adolescents with anxiety disorders: Findings from the CAMS. *Journal of Consulting and Clinical Psychology*, 79(6), 806–813. <https://doi.org/10.1037/a0025933>
- Hodson, K. J., McManus, F. V., Clark, D. M., & Doll, H. (2008). Can Clark and Wells' (1995) cognitive model of social phobia be applied to young people? *Behavioural and Cognitive Psychotherapy*, 36(Special Issue 04), 449–461. <https://doi.org/10.1017/S1352465808004487>
- Ingul, J. M., Aune, T., & Nordahl, H. M. (2014). A randomized controlled trial of individual cognitive therapy, group cognitive behaviour therapy and attentional placebo for adolescent social phobia. *Psychotherapy and Psychosomatics*, 83(1), 54–61. <https://doi.org/10.1159/000354672>
- Kessler, R. C., Berglund, P., Demler, O., Jin, R., Merikangas, K. R., & Walters, E. E. (2005). Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the national comorbidity survey replication. *Archives of General Psychiatry*, 62(6), 593–602. <https://doi.org/10.1001/archpsyc.62.6.593>
- Leigh, E. (2023). Editorial: Treating social anxiety disorder in children and young people: Challenges and opportunities. *Journal of the American Academy of Child & Adolescent Psychiatry*, 62(6), 621–623. <https://doi.org/10.1016/j.jaac.2022.12.002>
- Leigh, E., Chiu, K., & Clark, D. M. (2020). The effects of modifying mental imagery in adolescent social anxiety. *PLoS One*, 15(4), Article e0230826-e0230826. <https://doi.org/10.1371/journal.pone.0230826>
- Leigh, E., Chiu, K., & Clark, D. M. (2021). Self-focused attention and safety behaviours maintain social anxiety in adolescents: An experimental study. *PLoS One*, 16(2), Article e0247703. <https://doi.org/10.1371/journal.pone.0247703>
- Leigh, E., & Clark, D. M. (2018). Understanding social anxiety disorder in adolescents and improving treatment outcomes: Applying the cognitive model of Clark and Wells (1995). *Clinical Child and Family Psychology Review*, 21(3), 388–414. <https://doi.org/10.1007/s10567-018-0258-5>
- Leigh, E., & Clark, D. M. (2021a). Development and preliminary validation of the child & adolescent social cognitions questionnaire. *Child psychiatry & human development*. <https://doi.org/10.1007/s10578-021-01163-0>
- Leigh, E., & Clark, D. M. (2021b). Establishing the psychometric properties of the self-report Liebowitz Social Anxiety Scale for Children and Adolescents in a general population adolescent sample. *European Journal of Psychological Assessment*. <https://doi.org/10.1027/1015-5759/a000659>
- Leigh, E., & Clark, D. M. (2023). Internet-delivered therapist-assisted cognitive therapy for adolescent social anxiety disorder (OSCA): A randomised controlled trial addressing preliminary efficacy and mechanisms of action. *Journal of Child Psychology and Psychiatry*. <https://doi.org/10.1111/jcpp.13680>
- Masia-Warner, C., Fisher, P. H., Shrout, P. E., Rathor, S., & Klein, R. G. (2007). Treating adolescents with social anxiety disorder in school: An attention control trial. *Journal of Child Psychology and Psychiatry*, 48(7), 676–686. <https://doi.org/10.1111/j.1469-7610.2007.01737.x>
- Masia-Warner, C., Storch, E. A., Pincus, D. B., Klein, R. G., Heimberg, R. G., & Liebowitz, M. R. (2003). The Liebowitz social anxiety scale for children and adolescents: An initial psychometric investigation. *Journal of the American Academy of Child & Adolescent Psychiatry*, 42(9), 1076–1084. <https://doi.org/10.1097/01.CHL.0000070249.24125.89>
- Masia, C. L., Klein, R. G., & Liebowitz, M. R. (1999). *The Liebowitz social anxiety scale for children and adolescents*. LSAS-CA.
- Mayo-Wilson, E., Dias, S., Mavranzeouli, I., Kew, K., Clark, D. M., Ades, A., et al. (2014). Psychological and pharmacological interventions for social anxiety disorder in adults: A systematic review and network meta-analysis. *The Lancet Psychiatry*, 1, 368–376. [https://doi.org/10.1016/S2215-0366\(14\)70329-3](https://doi.org/10.1016/S2215-0366(14)70329-3)
- National Institute for Health and Care Excellence. (2013). *Social anxiety disorder: Recognition, assessment and treatment of social anxiety disorder*. Clinical guideline 159) <https://guidance.nice.org.uk/CG159>
- Nunnally, J. C., & Bernstein, I. H. (1978). *Psychometric theory* (2nd ed.). McGraw-Hill.
- Polanczyk, G. V., Salum, G. A., Sugaya, L. S., Caye, A., & Rohde, L. A. (2015). Annual research review: A meta-analysis of the worldwide prevalence of mental disorders in children and adolescents. *Journal of Child Psychology and Psychiatry*, 56(3), 345–365. <https://doi.org/10.1111/jcpp.12381>
- Rapee, R. M., McLellan, L. F., Carl, T., Trompeter, N., Hudson, J. L., Jones, M. P., et al. (2023). Comparison of transdiagnostic treatment and specialized social anxiety treatment for children and adolescents with social anxiety disorder: A randomized controlled trial. *Journal of the American Academy of Child & Adolescent Psychiatry*, 62(6), 646–655. <https://doi.org/10.1016/j.jaac.2022.08.003>
- Schreiber, F., Höfling, V., Stangier, U., Bohn, C., & Steil, R. (2012). A cognitive model of social phobia: Applicability in a large adolescent sample. *International Journal of Cognitive Therapy*, 5(3), 341–358. <https://doi.org/10.1521/ijct.2012.5.3.341>
- Schutters, S. I., Dominguez, M. D., Knappe, S., Lieb, R., van Os, J., Schruers, K. R., et al. (2012). The association between social phobia, social anxiety cognitions and paranoid symptoms. *Acta Psychiatrica Scandinavica*, 125(3), 213–227. <https://doi.org/10.1111/j.1600-0447.2011.01787.x>
- Spence, S. H., Donovan, C. L., March, S., Kenardy, J. A., & Hearn, C. S. (2017). Generic versus disorder specific cognitive behavior therapy for social anxiety disorder in youth: A randomized controlled trial using internet delivery. *Behaviour Research and Therapy*, 90, 41–57. <https://doi.org/10.1016/j.brat.2016.12.003>
- Vilaplana-Pérez, A., Pérez-Vigil, A., Sidorchuk, A., Brander, G., Isomura, K., Hesselmark, E., et al. (2020). Much more than just shyness: The impact of social anxiety disorder on educational performance across the lifespan. *Psychological Medicine*, 1–9. <https://doi.org/10.1017/S0033291719003908>
- Woody, S. R., Chambless, D. L., & Glass, C. R. (1997). Self-focused attention in the treatment of social phobia. *Behaviour Research and Therapy*, 35(2), 117–129. [https://doi.org/10.1016/s0005-7967\(96\)00084-8](https://doi.org/10.1016/s0005-7967(96)00084-8)