



The space of barcode bases for persistence modules

Emile Jacquard¹ · Vidit Nanda¹ · Ulrike Tillmann¹

Received: 17 March 2022 / Revised: 29 April 2022 / Accepted: 2 May 2022 /
Published online: 13 July 2022
© The Author(s) 2022, corrected publication 2022

Abstract

The barcode of a persistence module serves as a complete combinatorial invariant of its isomorphism class. Barcodes are typically extracted by performing changes of basis on a persistence module until the constituent matrices have a special form. Here we describe a new algorithm for computing barcodes which also keeps track of, and outputs, such a change of basis. Our main result is an explicit characterisation of the group of transformations that sends one barcode basis to another. Armed with knowledge of the entire space of barcode bases, we are able to show that any map of persistence modules can be represented via a partial matching between bars provided that neither source nor target admits nested bars in its barcode. We also generalise the algorithm and results described above to work for zigzag modules.

Keywords Persistence module · Barcode basis · Quiver representation

Mathematics Subject Classification 55N31 · 16G20

1 Introduction

Persistence modules appear in different forms and guises across many areas of mathematics. In recent years, particular interest and focus has come from their extensive use in topological data analysis (TDA) in general and persistent homology in particular.

✉ Vidit Nanda
nanda@maths.ox.ac.uk
Emile Jacquard
jacquard@maths.ox.ac.uk
Ulrike Tillmann
tillmann@maths.ox.ac.uk

¹ Mathematical Institute, University of Oxford, Oxford, UK

In this paper, we examine persistence modules of the form (V_\bullet, f_\bullet) :

$$V_0 \xrightarrow{f_1} V_1 \xrightarrow{f_2} \cdots \xrightarrow{f_{\ell-1}} V_{\ell-1} \xrightarrow{f_\ell} V_\ell, \quad (1)$$

where each V_i is a vector space of finite dimension n_i (over an underlying field \mathbb{F}) and each $f_i : V_{i-1} \rightarrow V_i$ is a linear map. We study three different aspects.

1.1 Computing barcode bases

The first is the central question of finding a barcode basis for (V_\bullet, f_\bullet) . This amounts to a choice of basis for each V_i with respect to which the linear maps f_i have a particularly nice form — they admit at most a single 1 in each row and column, with all other entries being 0. The existence of such bases and matrix representations is well known (Gabriel 1972). We say that the matrices are in *barcode form* and the corresponding basis of $\bigoplus_{i=0}^{\ell} V_i$ is a *barcode basis*, since the barcodes familiar from TDA can easily be extracted.

Algorithms to compute barcode bases in TDA typically take as input a filtered chain complex as in Zomorodian and Carlsson (2005), where one has recourse to matrix representations of the boundary operators. Algorithms for general persistence modules include the well-known (Carlsson and de Silva 2010) and much more recently, (Hang and Mio 2020) and (Henselman-Petrusek 2017). Here we present a new algorithm that takes as its input a matrix representation $A_\bullet = (A_1, \dots, A_\ell)$ relative to some initial basis of the persistence module (V_\bullet, f_\bullet) and outputs a sequence $g = (g_0, \dots, g_\ell)$ of change of basis matrices g_i for each of the V_i so that the new matrix representation $A'_\bullet = (A'_1, \dots, A'_\ell)$ with $A'_i = g_i \cdot A_i \cdot g_{i-1}^{-1}$ is in barcode form. Our algorithm in Sect. 3.2 is explicit and elementary in the sense that every intermediate step amounts to performing standard (row or column) operations on the constituent A_i 's. The key difficulty here is that column operations on A_i often force new matrix operations on A_k for $k < i$, and similarly row operations on A_i often require changes in A_k for $k > i$.

1.2 The space of barcode bases

Our second goal is to describe the set of all barcode bases¹ of (V_\bullet, f_\bullet) . We show that this set can naturally be identified as the stabiliser of a matrix representation A_\bullet of (V_\bullet, f_\bullet) , and hence as a subgroup of the product

$$G := \mathrm{GL}(n_0; \mathbb{F}) \times \cdots \times \mathrm{GL}(n_\ell; \mathbb{F}),$$

where $\mathrm{GL}(d; \mathbb{F})$ indicates the general linear group of invertible $d \times d$ matrices with entries in \mathbb{F} . Writing

$$[i_1, j_1] \leq [i_2, j_2] \quad \text{whenever} \quad i_1 \leq i_2 \leq j_1 \leq j_2$$

¹ This set has a natural topology when working over a field with topology such as \mathbb{R} or \mathbb{C} .

and $\text{Mat}(m \times n; \mathbb{F})$ for the set of $m \times n$ matrices with coefficients in \mathbb{F} , we show in Theorem 4.4 that the set of barcode bases is in one-to-one correspondence with

$$\prod_{0 \leq i \leq j \leq \ell} \text{GL}(d_{ij}; \mathbb{F}) \times \prod_{[i_1, j_1] \preceq [i_2, j_2]} \text{Mat}(d_{i_1 j_1} \times d_{i_2 j_2}; \mathbb{F}).$$

Here d_{ij} is the multiplicity of the interval $[i, j]$ in the barcode of (V_\bullet, f_\bullet) . If the matrix representation is already in barcode form, then the elements in $\text{GL}(d_{ij}; \mathbb{F})$ correspond to changes of basis for the sub-vector space of V_i spanned by the basis elements corresponding to the bars $[i, j]$, and the elements in $\text{Mat}(d_{i_1 j_1} \times d_{i_2 j_2}; \mathbb{F})$ represent the changes to basis vectors in V_{i_2} corresponding to intervals $[i_2, j_2]$ obtained by adding vectors from V_{i_2} which correspond to intervals $[i_1, j_1]$.

1.3 Simplifying maps of persistence modules

We now turn attention to our third problem. Given a map of persistence modules

$$\phi : (V_\bullet, f_\bullet) \rightarrow (W_\bullet, h_\bullet),$$

we seek barcode bases for source and target in terms of which ϕ assumes its simplest form, in the sense that we now specify. An interval $[i, j]$ represents a submodule canonically isomorphic to the interval module $\mathbf{I}[i, j]_\bullet$. It is an elementary observation that such a module can be mapped non-trivially to another interval module $\mathbf{I}[i', j']_\bullet$ if and only if $[i', j'] \preceq [i, j]$; and in this case the non-zero map is unique up to a non-zero scalar. In the simplest case, ϕ induces a partial matching where each bar in the source is mapped to exactly one in the target or mapped to zero. Surprisingly, we show in Theorem 5.3 that such a partial matching exists (after a change of barcode bases) whenever neither source nor target admit a pair of strictly nested intervals in their respective barcode decompositions.² In an example we also show that these conditions are necessary.

1.4 Zigzag modules

Finally, we generalise the algorithm and theorems described above to zigzag modules of a fixed type τ : the linear maps of the persistence module (1) can go either forward $V_{i-1} \xrightarrow{f_i} V_i$ or backward $V_{i-1} \xleftarrow{g_i} V_i$ according to pattern fixed by τ . Such modules are also classified in terms of sums of interval modules. Our algorithm can be adapted to compute barcode bases of zigzag modules of any type. Next, we introduce a generalisation of the order \preceq that depends on the type τ . This order takes into account that the order \preceq has to be reversed when all the arrows in (1) are reversed. With this order in place, we can once again classify the set of all barcode bases, see Theorem 6.11. Similarly when considering maps of zigzag modules, we generalise the

² Two bars $[i, j]$ and $[i', j']$ are strictly nested if $i < i'$ and $j' < j$.

notion of strictly nested bars, which once again depends on the type τ . Excluding such nested bars, we are able to obtain barcode bases of the source and target zigzag modules in terms of which the map is described by a partial matching on the set of bars; see Theorem 6.14.

1.5 Outline

We take the view that a persistence module (V_\bullet, f_\bullet) is a quiver representation. In Sect. 3, we provide a constructive proof (and concomitant algorithm) of Gabriel's decomposition theorem for persistence modules. In Sect. 4 we identify the set of all barcode bases with a stabiliser of the action of the group G on the set of all possible matrix representations of (V_\bullet, f_\bullet) . In Sect. 5 we study maps between persistence modules by viewing them as representations of ladder quivers with relations. Gabriel's theorem no longer applies here; however, using similar arguments as in Asashiba et al. (2018), we are able to prove a finite decomposition when source and target have no nested bars. Finally, in Sect. 6 we extend our results to any quiver of type A, that is zigzag persistence modules.

1.6 Related work

As mentioned above, there are several well known algorithms which compute barcodes of persistence modules. Some start with chains on a filtered simplicial complex, some deal with more general persistence modules and in some cases zigzag modules.

There are two algorithms that explicitly deal with computing the barcode bases associated to the interval decomposition. In Carlsson et al. (2019) the authors use matrix factorisation techniques to obtain bases in which the matrices are in echelon form. This technique also applies to zigzag modules. In Gregorio et al. (2021) the authors inductively compute interval bases using basis completions techniques at each step, but they do not deal with the zigzag case. Neither of these papers attempts to compute the set of barcode bases associated with the persistence module, and the algorithm we describe here takes a different approach to reducing matrices in barcode form. Most recently, the authors of Hang et al. (2021) use U-match matrix factorisation to reduce computational complexity and memory storage in computing barcodes.

It was shown in Escolar and Hiraoka (2016) that maps between persistence modules of length less than 5 admit a tractable classification in the sense that the associated ladder persistence modules are always of finite type. In contrast, the authors of Buchet and Escolar (2018) find an infinite class of indecomposable non-isomorphic ladder persistence module whenever the length is greater than 5. In Asashiba et al. (2018) the authors outline an algorithm which computes the decomposition into a sum of indecomposables for ladder persistence modules of length < 5 .

2 Persistence modules and barcode bases

A **persistence module**, for the purposes of this paper, is a finite collection (V_\bullet, f_\bullet) of finite-dimensional vector spaces V_i over a field \mathbb{F} along with \mathbb{F} -linear maps f_i arranged as follows:

$$V_0 \xrightarrow{f_1} V_1 \xrightarrow{f_2} \dots \xrightarrow{f_{\ell-1}} V_{\ell-1} \xrightarrow{f_\ell} V_\ell.$$

The number $\ell + 1$ is called the *length* of (V_\bullet, f_\bullet) . The *direct sum* of (V_\bullet, f_\bullet) with another persistence module (W_\bullet, h_\bullet) of the same length is defined pointwise — in other words, the vector space at its i -th position is $V_i \oplus W_i$ for each admissible index i , and similarly the corresponding linear map is given by $f_i \oplus h_i$. We call (V_\bullet, f_\bullet) *isomorphic* to (W_\bullet, h_\bullet) if there are invertible linear maps $\phi_i : V_i \rightarrow W_i$ so that the square

$$\begin{array}{ccc} V_{i-1} & \xrightarrow{f_i} & V_i \\ \phi_{i-1} \downarrow \sim & & \sim \downarrow \phi_i \\ W_{i-1} & \xrightarrow{h_i} & W_i \end{array}$$

commutes for each index i in $\{1, \dots, \ell\}$. Isomorphisms from (V_\bullet, f_\bullet) to itself are called *automorphisms*, and these evidently form a group under composition. We denote this group by $\text{Aut}(V_\bullet, f_\bullet)$.

The **interval module** corresponding to a pair of non-negative integers $i \leq j$ is the persistence module $\mathbf{I}[i, j]_\bullet$ given by

$$0 \rightarrow \dots \rightarrow 0 \rightarrow \mathbb{F} \rightarrow \dots \rightarrow \mathbb{F} \rightarrow 0 \rightarrow \dots \rightarrow 0,$$

where the contiguous string of \mathbb{F} 's spans $\{i, i + 1, \dots, j - 1, j\}$, all intermediate $\mathbb{F} \rightarrow \mathbb{F}$ maps are identities, and all other vector spaces are trivial. The importance of interval modules stems from the following result (Zomorodian and Carlsson 2005).

Theorem 2.1 *For each persistence module (V_\bullet, f_\bullet) of length $\ell + 1$, there exists a finite set of non-negative integer pairs*

$$\text{Bar}(V_\bullet, f_\bullet) := \{i_1 \leq j_1, \dots, i_k \leq j_k\},$$

(with $[i_p, j_p] \subset [0, \ell]$ for all $1 \leq p \leq k$), called the **barcode** of (V_\bullet, f_\bullet) , and an integer **multiplicity** $d_{i_p, j_p} > 0$ so that (V_\bullet, f_\bullet) is isomorphic to a direct sum of interval modules:

$$(V_\bullet, f_\bullet) \simeq \bigoplus_{p=1}^k \mathbf{I}[i_p, j_p]_\bullet^{d_{i_p, j_p}}. \tag{2}$$

Here the i -th summand on the right side is to be interpreted as the d_{i_p, j_p} -fold direct sum of the interval module $\mathbf{I}[i_p, j_p]_\bullet$ with itself.

This *interval decomposition* theorem follows from Gabriel’s foundational result on the decomposability of quiver representations (Gabriel 1972) — since (V_\bullet, f_\bullet) is a representation of a type- $\mathbf{A}_{\ell+1}$ quiver. Our goal here is to provide an explicit algorithm which not only furnishes such the isomorphism (2), but can also be readily implemented on a computer.

To this end, fix a persistence module (V_\bullet, f_\bullet) of length $\ell + 1$ and set $n_i := \dim_{\mathbb{F}} V_i$ for each $i \in \{0, \dots, \ell\}$. Without loss of generality, we may select a *basis family*

$$\mathcal{B} := \{B_i \subset V_i \mid 0 \leq i \leq \ell\},$$

where each B_i forms an ordered basis for the vector space V_i . This choice amounts to fixing an isomorphism $V_i \simeq \mathbb{F}^{n_i}$ for each i . Thus, every linear map $f_i : V_{i-1} \rightarrow V_i$ can be represented (in terms of the chosen bases B_{i-1} and B_i from \mathcal{B}) as a matrix A_i of size $n_i \times n_{i-1}$ with entries in \mathbb{F} ; consequently, (V_\bullet, f_\bullet) is isomorphic to

$$\mathbb{F}^{n_0} \xrightarrow{A_1} \mathbb{F}^{n_1} \xrightarrow{A_2} \dots \xrightarrow{A_{\ell-1}} \mathbb{F}^{n_{\ell-1}} \xrightarrow{A_\ell} \mathbb{F}^{n_\ell}. \tag{3}$$

In light of Theorem 2.1, we are particularly interested in a special class of basis families.


Definition 2.2 An $m \times n$ matrix A of rank r is in **barcode form** if there exists a strictly increasing function $c : \{1, \dots, r\} \rightarrow \{1, \dots, n\}$ so that

$$A_{ij} = \begin{cases} 1 & \text{if } j = c(i), \\ 0 & \text{otherwise.} \end{cases}$$

Thus, a matrix is in barcode form whenever its entries lie in $\{0, 1\}$, with at most one non-zero term in each row and column, and the r non-zero terms appear in the first r rows and in strictly increasing column order.

A basis family \mathcal{B} is called an **barcode basis** for (V_\bullet, f_\bullet) if all of the A_i are in barcode form. The natural basis arising from an interval decomposition of a persistence module is a barcode basis.

Example 2.3 Consider, for instance, the persistence module of length 4 given by the barcode containing $0 \leq 3$ along with $0 \leq 1$ and $1 \leq 3$, each with multiplicity one:



With respect to the basis family obtained by ordering these intervals from top to bot-

tom, the matrices A_i are given by

$$A_1 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad A_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad A_3 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix},$$

and all three are evidently in barcode form. Conversely, one can also recover the interval decomposition immediately from these three matrices.

To put our quest for a constructive proof of Theorem 2.1 on a firm algebraic footing, let X be the set of all the possible matrix-sequences A_\bullet which can arise in (3). It is a (strict) subset of the product of matrices of the appropriate dimensions:

$$X \subset \prod_{i=1}^{\ell} \text{Mat}(n_i \times n_{i-1}; \mathbb{F}). \tag{4}$$

Writing $\text{GL}(n; \mathbb{F})$ for the group of all $n \times n$ invertible matrices over \mathbb{F} , consider the product

$$G := \prod_{i=0}^{\ell} \text{GL}(n_i; \mathbb{F}), \tag{5}$$

which acts naturally via a change-of-basis action on X : the group element $g := (g_0, \dots, g_\ell)$ sends each matrix-sequence A_\bullet in X to the new sequence $(gA)_\bullet$ given by

$$(gA)_i := g_i \cdot A_i \cdot g_{i-1}^{-1} \tag{6}$$

for each admissible index i . This is equivalent to replacing the original basis family $\mathcal{B} = \{B_i\}$ with the new basis family $g\mathcal{B} = \{g_i B_i\}$. Thus, X is the free orbit of A_\bullet under this G -action. So our first task, solved in Sect. 3, translates to discovering some $g \in G$ that transforms a given basis family \mathcal{B} of (V_\bullet, f_\bullet) to a barcode basis.

3 Constructing a barcode basis

Throughout this section, we fix a persistence module (V_\bullet, f_\bullet) expressed as a sequence of matrices A_\bullet as in (3) with respect to an arbitrary (i.e., not necessarily barcode) basis family \mathcal{B} .

3.1 Barcode bases via elementary matrix operations

To conveniently describe relevant elements of G , we fix notation for matrices which implement certain fundamental row and column operations.

Definition 3.1 For each dimension $n > 0$, distinct indices $1 \leq p, q \leq n$, and scalar $\lambda \in \mathbb{F}$, let $e_{p,q}^n(\lambda)$ denote the **elementary matrix** in $\text{GL}(n; \mathbb{F})$ which has 1's all along its diagonal, λ in the (p, q) -th position, and zeros everywhere else.

Since the dimension n will be clear from context, we omit it from the superscript and simply write $e_{p,q}(\lambda)$ to indicate the relevant elementary matrix. The following standard facts about such matrices will be freely used in the sequel —

- (1) multiplying a matrix on the left by $e_{p,q}(\lambda)$ implements the following **elementary row operation**

$$\text{Row}(p) \leftarrow \text{Row}(p) + \lambda \cdot \text{Row}(q)$$

which we denote $\mathbf{R}_{p \leftarrow q}(\lambda)$; similarly,

- (2) multiplying a matrix on the right by $e_{p,q}(\lambda)$ implements the following **elementary column operation**

$$\text{Col}(q) \leftarrow \text{Col}(q) + \lambda \cdot \text{Col}(p),$$

which we denote $\mathbf{C}_{q \leftarrow p}(\lambda)$; and finally,

- (3) the inverse of $e_{p,q}(\lambda)$ is $e_{p,q}(-\lambda)$.

Remark 3.2 Consider the element $g = (g_0, \dots, g_\ell) \in G$ for which $g_i = e_{p,q}(\lambda)$ and all the other g_j are identity matrices. The action of this g on a given matrix sequence A_\bullet is to simultaneously perform $\mathbf{C}_{p \leftarrow q}(\lambda)$ on A_i and $\mathbf{R}_{q \leftarrow p}(-\lambda)$ on A_{i-1} while leaving all the other A_j 's invariant.

The following result plays an essential part in our constructive proof of Theorem 2.1. In its statement and beyond, we will use $A(p, q)$ to indicate the entry in the p -th row and q -th column of a given matrix A .

Lemma 3.3 *Assume that the first $\ell - 1$ matrices $\{A_i \mid 1 \leq i < \ell\}$ of (3) are in barcode form, and that the last matrix A_ℓ has a pivot in the (r, q) position, i.e., $A_\ell(r, q) = 1$ and all other entries in the q -th column are zero. If there is a nonzero entry $\alpha := A_\ell(r, p)$ in the same row r but some other column $p > q$, then there exists $g \in G$ with $g_\ell = \text{Id}$ so that $(gA)_\bullet$ equals A_\bullet except A_ℓ where the α entry is replaced by zero.*

Proof We proceed by induction on ℓ , noting that the case $\ell = 1$ is immediately true since there is only one matrix in sight. Assume that the statement holds up to $\ell - 1$. The r -th row of A_ℓ contains a pivot 1 in the q -th column and some $\alpha \neq 0$ in the p -th column. To eliminate this offending α , we perform $\mathbf{C}_{p \leftarrow q}(-\alpha)$ on A_ℓ by performing the basis change $e_{p,q}(-\alpha)$ on $V_{\ell-1}$. Since $A_\ell(r, q)$ is assumed to be a pivot, the only resulting difference in A_ℓ is that the (r, p) -th entry changes from α to 0. But by Remark 3.2, we are also compelled to perform $\mathbf{R}_{q \leftarrow p}(\alpha)$ on the preceding matrix $A_{\ell-1}$. This results in a new matrix $A'_{\ell-1}$, and there are now 2 cases to consider, of which only the second requires the inductive hypothesis:

Case 1: if the p -th row of $A_{\ell-1}$ is identically zero, then our row operation has had no effect whatsoever; thus, $A'_{\ell-1} = A_{\ell-1}$ is still in barcode form and we have arrived at the desired result.

Case 2: If the p -th row of $A_{\ell-1}$ is nonzero, then since $q < p$ and $A_{\ell-1}$ is in barcode form, we see that the q -th row of $A_{\ell-1}$ must also be non-zero. Then by Definition 2.2 they must have pivot ones in distinct columns, say c and d respectively, and furthermore $c < d$. Thus, after we have performed $\mathbf{R}_{q \leftarrow p}(\alpha)$ on $A_{\ell-1}$, the resulting matrix $A'_{\ell-1}$ has the form

$$A'_{\ell-1} = \begin{bmatrix} & c & & & d & & \\ 0 & 1 & 0 & \cdots & 0 & \alpha & 0 \\ & 0 & & & & 0 & \\ & \vdots & & & & \vdots & \\ & 0 & & & & 0 & \\ 0 & 0 & 0 & \cdots & & 1 & 0 \end{bmatrix} \begin{matrix} q \\ \\ \\ \\ p \end{matrix}$$

By induction, there exists a $g \in \prod_{i=0}^{\ell-1} \text{GL}_{n_i}(\mathbb{F})$ with $g_{\ell-1} = \text{Id}$ so that $g_i A_i g_{i-1}^{-1}$ is still in barcode form for $1 \leq i \leq \ell - 2$, and

$$g_{\ell-1} A_{\ell-1} g_{\ell-2}^{-1} = \begin{bmatrix} & c & & & d & & \\ 0 & 1 & 0 & \cdots & 0 & 0 & 0 \\ & 0 & & & & 0 & \\ & \vdots & & & & \vdots & \\ & 0 & & & & 0 & \\ 0 & 0 & 0 & \cdots & & 1 & 0 \end{bmatrix} \begin{matrix} q \\ \\ \\ \\ p \end{matrix}$$

is again in barcode form. Furthermore, since $g_{\ell-1} = \text{Id}$, the matrix A_ℓ is left unchanged by this change of basis. The desired basis change is $(g_0, g_1, \dots, g_{\ell-2}, e_{p,q}(-\alpha), \text{Id})$. \square

Proposition 3.4 *Given the sequence of matrices A_\bullet as in (6), there is a $g \in G$ such that $(gA)_\bullet$ has all its matrices in barcode form.*

Proof When $\ell = 1$, we may diagonalise the matrix A_1 via standard row and column operations. Proceeding by induction for $\ell > 1$, assume the existence of some group element

$$g' = (g_0, \dots, g_{\ell-1}) \in \prod_{i=0}^{\ell-1} \text{GL}_{n_i}(\mathbb{F})$$

satisfying the following property: the matrices $g_i A_i g_{i-1}^{-1}$ are in barcode form for $1 \leq i \leq \ell - 1$.

Consider $g = (g', \text{Id}_{n_\ell})$, which evidently lies in G . Replacing A_\bullet by $(gA)_\bullet$ if necessary, we may assume that A_\bullet has its first $\ell - 1$ matrices in barcode form. Performing row operations on A_ℓ has no impact on the previous matrices, as it corresponds to multiplying A_ℓ on the left by some g_ℓ . Thus, we may assume without loss of generality that all previous matrices are in barcode form while A_ℓ itself is in reduced row echelon

form. By Lemma 3.3, there is a basis change $g \in G$ which zeroes out each non-pivot entry whilst maintaining the barcode form of the previous matrices. Applying these basis changes gets us to the desired barcode basis. \square

Remark 3.5 If \mathcal{B} is the basis family with respect to which (V_\bullet, f_\bullet) has matrix form A_\bullet , then $g\mathcal{B}$ is a barcode basis where $g \in G$ is as in Proposition 3.4. We may therefore regard it as a constructive analogue of Theorem 2.1.

3.2 Algorithms

Here we describe algorithms which implement the constructions of Lemma 3.3 and Proposition 3.4. In particular, the main algorithm **CompPers** described below accepts as input an initial sequence of matrices A_\bullet as in (3) and puts them in barcode form. The sub-computations which we require frequently have been isolated into concomitant subroutines, described as follows.

- (1) The first subroutine **ColOp** implements the inductive strategy underlying our proof of Lemma 3.3; in particular, this algorithm acts as step k of the inductive procedure described in the proof of that lemma.
- (2) The second subroutine **Reduce** takes as input a sequence A_\bullet for which the first $\ell - 1$ matrices are in barcode form together with an invertible matrix $g \in \prod_{i=0}^{\ell-1} \text{GL}(n_i; \mathbb{F})$. It then reduces the final matrix A_ℓ until it is in barcode form, *while maintaining the barcode form of all previous matrices* and suitably updating the basis change g .
- (3) Finally, the main algorithm **CompPers**(A_\bullet) takes as input an arbitrary sequence of matrices A_\bullet and produces as output $g \in G$ together with $(gA)_\bullet$ in barcode form. From these matrices we can directly access all intervals in barcode of (V_\bullet, f_\bullet) .

Algorithm 1: ColOp

Input: A_\bullet, g, k, r, q, p
Output: Updated A_\bullet and basis change g , zeroing out $A_k(r, p)$

- 1 $\mathbf{C}_{p \leftarrow q}(-\alpha)$ on A_k
- 2 $\mathbf{R}_{q \leftarrow p}(\alpha)$ on A_{k-1}
- 3 $g_k = e_{p,q}(-\alpha)$
- 4 **if** $k = 0$ or p -th row of $A_{k-1} = 0$ **then**
- 5 | return (A_\bullet, g)
- 6 **end**
- 7 **else**
- 8 | Find pivot columns $c < d$ of the pivot rows $q < p$ of A_{k-1}
- 9 | return **(ColOp** $(A_\bullet, g, k - 1, q, c, d)$)
- 10 **end**

Remark 3.6 The computational complexity of **CompPers**(A_\bullet) can be expressed in terms of $n = \max_{0 \leq i \leq \ell} n_i$ and ℓ . The cost of placing all the A_i in reduced row echelon form via Gaussian elimination is $O(n^3\ell)$. Furthermore, performing column

Algorithm 2: Reduce

Input: A_\bullet, g , where A_\bullet has its first $\ell - 1$ matrices in reduced form
Output: A_\bullet in reduced barcode form and updated g

- 1 row reduce (A_ℓ)
- 2 Append g with corresponding g_ℓ
- 3 **while** there are $A_\ell(r, p) \neq 0$ terms with $A_\ell(r, q) = 1$ a pivot **do**
- 4 | $A_\bullet, h = \text{ColOp}(A_\bullet, g, \ell, r, q, p)$
- 5 | $g = hg$
- 6 **end**
- 7 return (A_\bullet, g)

Algorithm 3: CompPers

Input: A_\bullet
Output: Reduced A_\bullet with corresponding change of basis g

- 1 $A'_\bullet = \{A_1\}$
- 2 $g = (\text{Id}_{n_0})$
- 3 **for** $1 \leq i \leq \ell - 1$ **do**
- 4 | $A'_\bullet, g = \text{Reduce}(A'_\bullet, g)$
- 5 | $A'_\bullet.\text{append}(A_{i+1}g_i^{-1})$
- 6 **end**
- 7 return ($\text{Reduce}(A'_\bullet, g)$)

operations to further reduce these matrices requires at most $O(n^2)$ operations on each matrix. And column operations on A_i will, in the worst case, require downstream column operations on $A_{i-1} \dots A_1$. Thus, for column operations, we have a $O(n^2 \sum_{i=1}^\ell i) = O(n^2 \frac{\ell(\ell-1)}{2}) = O(n^2 \ell^2)$ complexity. Combining these factors, the total complexity of the algorithm is

$$O(n^3 \ell + n^2 \ell^2).$$

At each step of the algorithm, we perform an elementary basis change on a single vector space V_i , which amounts to multiplying a matrix $g_i \in \text{GL}(V_i)$ by an elementary matrix. This incurs an $O(n)$ cost; thus, if we also wish to keep track of the basis changes, then the total complexity of **CompPers** becomes

$$O(n^4 \ell + n^3 \ell^2).$$

We conclude with an illustrative example of how **CompPers** acts on a sequence of input matrices.

Example 3.7 Consider

$$A_\bullet = \left(\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{bmatrix} \right)$$

To put A_3 in barcode form, we must zero out the terms $A_3(1, 3)$ and $A_3(2, 3)$ using column operations. At each step, we will be performing row and column operations on matrices of A_\bullet , amounting to basis changes on the vectors spaces V_0, V_1 and V_2 . For conciseness sake, we will not keep track of the basis changes done along the way, and will simply be performing operations on the matrices to put them in barcode form.

We begin by zeroing out the $A_3(2, 3)$ term.

1: $C_{3 \leftarrow 2}(-1)$ on A_3 , inducing $R_{2 \leftarrow 3}(1)$ on A_2 , giving us matrices

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

2: $C_{3 \leftarrow 2}(-1)$ on A_2 , inducing $R_{2 \leftarrow 3}(1)$. We see here that the third row of A_1 is zero, so we are in **Case 1** of Lemma 3.3, and so we are done.

We have achieved our goal of zeroing out $A_3(2, 3)$ whilst keeping the previous matrices in barcode form, making no other changes to A_3 . It remains to zero out the $A_3(1, 3)$ term.

1: $C_{3 \leftarrow 1}(-1)$ on A_3 , inducing $R_{1 \leftarrow 3}(1)$ on A_2 giving us matrices

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

2: $C_{3 \leftarrow 1}(-1)$ on A_2 , inducing $R_{1 \leftarrow 3}(1)$ on A_1 . Since the third row of A_1 is zero, this operations has no impact on A_1 , giving us matrices

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

and so we are done.

4 The space of barcode bases

Let $A_\bullet \in X$ be a sequence of matrices as in (3) arising from an arbitrary choice of basis for some persistence module (V_\bullet, f_\bullet) . Consider the group G from (5), recalling that G acts on X via change of basis. Our quest to describe all possible barcode bases for (V_\bullet, f_\bullet) begins with a formula for the stabiliser of the chosen matrices A_\bullet under this G -action. Namely, we seek the subgroup of G given by

$$\text{Stab}(A_\bullet) := \left\{ g \in G \mid g_i \cdot A_i \cdot g_{i-1}^{-1} = A_i \text{ for all } 1 \leq i \leq \ell \right\}. \tag{7}$$

To describe $\text{Stab}(A_\bullet)$, we employ two binary relations on the set of all intervals which might possibly arise in the barcode decomposition of (V_\bullet, f_\bullet) à la Theorem 2.1.

Definition 4.1 Let \preceq be the binary relation on $\{[i, j] \in \mathbb{Z}^2 \mid 0 \leq i \leq j \leq \ell\}$ given by

$$[a, b] \preceq [c, d] \text{ whenever } a \leq c \leq b \leq d.$$

(Although this relation \preceq is reflexive and anti-symmetric on its domain, it is not transitive and hence does not form a partial order.)

The second binary relation is the standard lexicographic order.

Definition 4.2 Let \trianglelefteq be the lexicographic ordering on $\{[i, j] \in \mathbb{Z}^2 \mid 0 \leq i \leq j \leq \ell\}$, given by

$$[a, b] \trianglelefteq [c, d] \iff a < c \text{ or } a = c \text{ and } b \leq d.$$

This yields a total order on the set of all possible bars in the interval decomposition of our persistence module (V_\bullet, f_\bullet) .

Remark 4.3 The binary relation \preceq is compatible with the lexicographical order \trianglelefteq in the sense that $[a, b] \preceq [c, d]$ implies $[a, b] \trianglelefteq [c, d]$.

Given a barcode basis \mathcal{B} , we may totally order its bars using the lexicographic order, arbitrarily ordering bars with the same start and end point. This in turn yields a natural ordering of the bases B_i . The matrix representation of such bases is unique. Indeed, the first d_{0i} basis vectors of B_i are part of $[0, i]$ bars, then the next $d_{0,i+1}$ basis vectors are those part an $[0, i + 1]$ bar, and so on following the lexicographic ordering until finally the $d_{i,\ell}$ basis vectors part of an $[i, \ell]$ bar. This yields a matrix representation for which A_i is of the form

$$A_i = \begin{bmatrix} M_0 & & & & \\ & M_1 & & & \\ & & \ddots & & \\ & & & & M_i \end{bmatrix}, \tag{8}$$

where

$$M_j = \begin{bmatrix} & & \text{Id}_{d_{j,i}} & & & \\ & 0 & & & & \\ & & \text{Id}_{d_{j,i+1}} & & & \\ & & & \ddots & & \\ & & & & & \text{Id}_{d_{j,\ell}} \end{bmatrix}$$

is a matrix of dimension $(\sum_{k=i}^{\ell} d_{jk}) \times (\sum_{k=i-1}^{\ell} d_{jk})$. As such, bases that have been ordered in the above way are barcode bases in the usual sense; we call these **ordered** barcode bases of (V_\bullet, f_\bullet) and devote the remainder of this section to completely characterising them.

Theorem 4.4 For each pair $[i, j]$ in $\{0, 1, \dots, \ell\}$ with $i \leq j$, let d_{ij} equal the multiplicity of $i \leq j$ in the barcode of (V_\bullet, f_\bullet) , with the understanding that $d_{ij} = 0$ whenever $[i, j]$ is not in $\mathbf{Bar}(V_\bullet, f_\bullet)$. Then there is a bijection of sets:

$$\text{Stab}(A_\bullet) \cong \prod_{[i,j]} \text{GL}(d_{ij}; \mathbb{F}) \times \prod_{[i_1,j_1] \preceq [i_2,j_2]} \text{Mat}(d_{i_1j_1} \times d_{i_2j_2}; \mathbb{F}).$$

(The induced group structure on the right side is given in Corollary 4.5 below)

Proof Elements in the same orbit have isomorphic stabilisers, so without loss of generality we may assume A_\bullet is given by the matrix representation of the linear maps in a ordered barcode basis. An element $g = (g_0, \dots, g_\ell)$ of G lies in $\text{Stab}(A_\bullet)$ if and only if we have an equality of matrix products

$$g_i \cdot A_i = A_i \cdot g_{i-1}$$

for each $i \in \{1, \dots, \ell\}$. Set $k_i := \text{rank } A_i$ and note that since A_i is in barcode form, there is a strictly increasing function $c_i : \{1, \dots, k_i\} \rightarrow \{1, \dots, n_{i-1}\}$ so that the unique nonzero entry in the p -th row of A_i lies in column $c_i(p)$. The product $g_i \cdot A_i$ on the left side of our equality has as its q -th column either the $c_i^{-1}(q)$ -th column of g_i (if q lies in the image of c_i), or is identically zero otherwise. Conversely, for $p \leq k_i$ the matrix $A_i \cdot g_{i-1}$ on the right side has as its p -th row the $c(p)$ -th row of g_{i-1} , and its rows corresponding to $p > k_i$ are identically zero.

Therefore, requiring these two products to be equal amounts to imposing three types of constraints on the entries of g_{i-1} and g_i :

- (1) $g_i(p, q) = 0$ whenever $p > k_i \geq q$.
- (2) $g_{i-1}(p, q) = 0$ whenever $p \in \text{Img}(c_i)$ and $q \notin \text{Img}(c_i)$.
- (3) $g_{i-1}(c_i(p), c_i(q)) = g_i(p, q)$ whenever both p and q are $\leq k_i$.

Recalling that (V_\bullet, f_\bullet) is the persistence module represented by A_\bullet , we have a bijection

$$\left[\begin{array}{l} \text{intervals } [i, j] \text{ in the} \\ \text{barcode of } (V_\bullet, f_\bullet) \end{array} \right] \xleftrightarrow{\cong} \left[\begin{array}{l} \text{sequences } \{p_k \mid i \leq k \leq j\} \text{ with} \\ c_k(p_k) = p_{k-1} \text{ for } i + 1 \leq k \leq j \end{array} \right]$$

Let $[i_1, j_1]$ and $[i_2, j_2]$ be two intervals in the barcode decomposition of (V_\bullet, f_\bullet) , and denote their corresponding sequences by $\{p_\bullet\}$ and $\{q_\bullet\}$. It follows from constraint (3) above that $g_k(p_k, q_k)$ remains constant whenever k ranges over the indices in $[i, j] := [i_1, j_1] \cap [i_2, j_2]$. In other words, we have

$$g_k(p_k, q_k) = g_{k'}(p_{k'}, q_{k'}) \text{ for all } k, k' \in [i, j]. \tag{9}$$

The following observation is crucial.

Claim: The entry $g_k(p_k, q_k)$ is zero for all $k \in [i, j]$ whenever $[i_1, j_1] \not\preceq [i_2, j_2]$.

To prove this claim, note that if $i_2 < i_1$ holds then $p_{i_1} > k_{i_1} \geq q_{i_1}$, so $g_{i_1}(p_{i_1}, q_{i_1}) = 0$ by constraint (1) above. Thus the claim extends to all k in $[i, j]$ by (9). Similarly, if $j_2 < j_1$ then $p_{j_2} \in \text{Img}(c_{j_2})$ but $q_{j_2} \notin \text{Img}(c_{j_2})$, whence $g_{j_2}(p_{j_2}, q_{j_2}) = 0$ by

constraint (2). Once again, this extends to all $k \in [i, j]$ by (9), and so the claim is proved.

Returning to the main argument, for each pair $[i_1, j_1] \preceq [i_2, j_2]$ of intervals in the barcode of (V_\bullet, f_\bullet) , we may select some $k \in [i_1, j_1] \cap [i_2, j_2]$. We denote by $g_{[i_1, j_1]}^{[i_2, j_2]}$ the submatrix of g_k spanned by all entries $g_k(p_k, q_k)$ for which $\{p_\bullet\}$ and $\{q_\bullet\}$ are sequences corresponding to intervals of type $[i_1, j_1]$ and $[i_2, j_2]$ respectively. Thus, $g_{[i_1, j_1]}^{[i_2, j_2]}$ has exactly $d_{i_1 j_1}$ rows and $d_{i_2 j_2}$ columns; and from (9) we know that it forms a submatrix of g_k for all k in $[i_1, j_1] \cap [i_2, j_2]$. It follows from our claim that each g_k is block upper-triangular:

$$g_k = \begin{bmatrix} g_{[0,k]}^{[0,k]} & g_{[0,k]}^{[0,k+1]} & \dots & g_{[0,k]}^{[0,\ell]} & 0 & \dots & \dots & 0 \\ 0 & g_{[0,k+1]}^{[0,k+1]} & \dots & g_{[0,k+1]}^{[0,\ell]} & 0 & \dots & \dots & 0 \\ 0 & 0 & \ddots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \vdots & \vdots & 0 & \dots & \dots & \dots & 0 & g_{[k,\ell-1]}^{[k,\ell-1]} & g_{[k,\ell-1]}^{[k,\ell]} \\ 0 & 0 & \dots & \dots & \dots & \dots & 0 & 0 & g_{[k,\ell]}^{[k,\ell]} \end{bmatrix} \tag{10}$$

The fact that g_k must be invertible forces the diagonal blocks to be invertible, while the off-diagonal blocks remain entirely unconstrained. The map

$$\text{Stab}(A_\bullet) \rightarrow \prod_{[i,j]} \text{GL}(d_{ij}; \mathbb{F}) \times \prod_{[i_1, j_1] \preceq [i_2, j_2]} \text{Mat}(d_{i_1 j_1} \times d_{i_2 j_2}; \mathbb{F}),$$

which sends each g to this distinguished collection of invertible $g_{[i,j]}^{[i,j]}$ and arbitrary $g_{[i_1, j_1]}^{[i_2, j_2]}$ furnishes the desired bijection. □

Using the block upper triangular form of the matrices g_k described in the argument above, we may immediately obtain the group structure of $\text{Stab}(A_\bullet)$.

Corollary 4.5 *For $g, h \in \text{Stab}(A_\bullet)$, we have*

$$(gh)_{[i_1, j_1]}^{[i_2, j_2]} = \sum_{[a,b]} g_{[i_1, j_1]}^{[a,b]} h_{[a,b]}^{[i_2, j_2]},$$

with the sum being indexed over intervals that satisfy $[i_1, j_1] \preceq [a, b] \preceq [i_2, j_2]$.

If \mathbb{F} is the field of real or complex numbers, as a subgroup of $\prod_{i=0}^\ell \text{GL}_{n_i}(\mathbb{F})$, the group $\text{Stab}(A_\bullet)$ is a Lie group. Theorem 4.4 immediately allows us to obtain its dimension.

Corollary 4.6 *If \mathbb{F} is the field of real or complex numbers, $\text{Stab}(A_\bullet)$ is a Lie group of dimension*

$$\sum_{[i_1, j_1] \preceq [i_2, j_2]} d_{i_1 j_1} d_{i_2 j_2}$$

As stated at the start of this section, our task here is to determine the space of all possible ordered barcode bases for a given persistence module (V_\bullet, f_\bullet) .

We denote by $\mathbb{B} = \{\mathcal{B} = (B_i)_{0 \leq i \leq \ell} \mid B_i \subset V_i \text{ is an ordered basis}\}$ the set of all possible ordered bases of (V_\bullet, f_\bullet) . Having fixed an initial basis $\mathcal{B} \in \mathbb{B}$, we know the group G from (5) acts freely and transitively on the set \mathbb{B} , so that any element of \mathbb{B} may be expressed as $g\mathcal{B}$ for some unique $g \in G$. Thus, once we have fixed an initial basis \mathcal{B} , the set \mathbb{B} may be identified with G . Then as subset of \mathbb{B} , the set of all possible ordered barcode bases of (V_\bullet, f_\bullet) can be identified as a subset of the group G .

Recall, X is the set of all possible matrix-sequences as defined in (4). For each possible basis $\mathcal{B} \in \mathbb{B}$, we define $A(\mathcal{B})_\bullet \in X$ to be the matrix representation of the linear maps f_\bullet in the chosen basis \mathcal{B} . This assignment prescribes the *matrix representation map*

$$A(\cdot)_\bullet : \mathbb{B} \mapsto X,$$

and finding all possible ordered barcode bases for (V_\bullet, f_\bullet) amounts to determining all bases $\mathcal{B} \in \mathbb{B}$ for which $A(\mathcal{B})_\bullet$ is as in (8). Furthermore, this map is equivariant in the sense that $A(g\mathcal{B})_\bullet = (gA(\mathcal{B}))_\bullet$ for each $g \in G$, where g acts via the basis action defined in (6).

The following result makes the link between the stabiliser of A_\bullet and the set of ordered barcode bases of a persistence module (V_\bullet, f_\bullet) .

Proposition 4.7 *Given a persistence module (V_\bullet, f_\bullet) together with an ordered barcode basis \mathcal{B} with matrix representation A_\bullet , the set of all ordered barcode bases is given by the orbit $\text{Stab}(A_\bullet)\mathcal{B}$.*

Proof Let \mathcal{B}' be another ordered barcode basis. As seen above, two ordered barcode bases have the same matrix representations (8), so that $A(\mathcal{B}')_\bullet = A_\bullet$. As previously stated, G acts freely and transitively on \mathbb{B} so that there exists a unique $g \in G$ for which $\mathcal{B}' = g\mathcal{B}$. We then have $A_\bullet = A(\mathcal{B})_\bullet = A(g\mathcal{B}')_\bullet = gA(\mathcal{B})_\bullet = gA_\bullet$, so that $g \in \text{Stab}(A_\bullet)$ which implies $\mathcal{B}' \in \text{Stab}(A_\bullet)\mathcal{B}$. \square

As such, we may identify the set of all ordered barcode bases of a persistence module (V_\bullet, f_\bullet) with $\text{Stab}(A_\bullet)$, which was fully characterised in Theorem 4.4.

Remark 4.8 The automorphism group $\text{Aut}_Q(M)$ of a representation M of a general quiver Q has been described, for instance in (Brion 2012, Section 2.2). It is known that $\text{Aut}_Q(M)$ is a semi-direct product of the form

$$U \rtimes \prod_{i=1}^r \text{GL}(m_i, \mathbb{F}).$$

Here $M = \bigoplus_{i=1}^r M_i^{m_i}$ is a decomposition of M into indecomposable summands M_i , while U is unipotent normal subgroup of $\text{Aut}_Q(M)$ (see (Brion 2012, Prop 2.2.1)). Viewed from this context, the main content of Theorem 4.4 is an explicit description of U in the special case where Q is a type-A quiver. In particular, U is generated by matrices which have the form (10), but with identity blocks along the diagonal. This

explicit description of U in the type-A case plays a crucial role in subsequent results which appear in this paper.


5 Simplifying maps of persistence modules

We now shift our interest to maps of persistence modules $\phi_\bullet : (V_\bullet, f_\bullet) \mapsto (W_\bullet, h_\bullet)$, where V_\bullet and W_\bullet are persistence modules of length $\ell + 1$. We recall that each such ϕ_\bullet is a collection of linear maps $\phi_i : V_i \mapsto W_i$ satisfying $\phi_i \circ f_i = h_i \circ \phi_{i-1}$. In other words, the following diagram commutes:

$$\begin{array}{ccccccc}
 V_0 & \xrightarrow{f_1} & V_1 & \xrightarrow{f_2} & \dots & \xrightarrow{f_{\ell-1}} & V_{\ell-1} & \xrightarrow{f_\ell} & V_\ell \\
 \downarrow \phi_0 & & \downarrow \phi_1 & & & & \downarrow \phi_{\ell-1} & & \downarrow \phi_\ell \\
 W_0 & \xrightarrow{h_1} & W_1 & \xrightarrow{h_2} & \dots & \xrightarrow{h_{\ell-1}} & W_{\ell-1} & \xrightarrow{h_\ell} & W_\ell
 \end{array}$$


Our objective here is to show that if neither V_\bullet nor W_\bullet admits a pair of strictly nested bars in its barcode, then there exist barcode bases of V_\bullet and W_\bullet in which ϕ induces a *partial matching* of the bars. To make this precise, we first describe the nestedness condition.

Definition 5.1 A bar $[i_2, j_2]$ is strictly nested in a bar $[i_1, j_1]$, denoted $[i_2, j_2] \subset [i_1, j_1]$ if $i_1 < i_2 \leq j_2 < j_1$. This is best represented as



Note that two intersecting bars are strictly nested if and only if they are not related by \preceq .

Before stating the main result, we remark that the data of our map ϕ_\bullet can be interpreted as a representation of the *rectangle* quiver of length $\ell + 1$:



Such representations have been called **ladder** persistence modules in the literature (Escobar and Hiraoka 2016). When treating ϕ as a ladder persistence module, we will denote it $(V_\bullet, W_\bullet, \phi_\bullet)$. We may therefore seek to decompose ϕ_\bullet into a direct sum of indecomposable ladder persistence modules.

Definition 5.2 Three families of ladder persistence modules are defined below:

- (1) Given intervals $[i_1, j_1] \preceq [i_2, j_2]$, denote by $\mathbf{R}_{[i_1, j_1]}^{[i_2, j_2]}$ the ladder persistence module where V_\bullet is the interval module $\mathbf{I}[i_2, j_2]_\bullet$ while W_\bullet is the interval module $\mathbf{I}[i_1, j_1]_\bullet$; all vertical maps are 1's whenever possible and 0 otherwise:

$$\begin{array}{ccccccc} & & \mathbb{F} & \longrightarrow & \mathbb{F} & \longrightarrow & \dots & \longrightarrow & \mathbb{F} & \longrightarrow & \dots & \longrightarrow & \mathbb{F} \\ & & \downarrow & & \downarrow & & & & \downarrow & & & & \\ \mathbb{F} & \longrightarrow & \dots & \longrightarrow & \mathbb{F} & \longrightarrow & \mathbb{F} & \longrightarrow & \dots & \longrightarrow & \mathbb{F} & & \end{array}$$

- (2) Given an interval $[i, j]$, let $\mathbf{I}^+[i, j]_\bullet$ denote the ladder persistence module for which V_\bullet is $\mathbf{I}[i, j]_\bullet$ and W_\bullet is 0, with all vertical maps necessarily being 0:

$$\begin{array}{ccccccc} & & \mathbb{F} & \longrightarrow & \mathbb{F} & \longrightarrow & \dots & \longrightarrow & \mathbb{F} \\ & & \downarrow & & \downarrow & & & & \downarrow \\ 0 & \longrightarrow & \dots & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & \dots & \longrightarrow & 0 & \longrightarrow & \dots & \longrightarrow & 0 \end{array}$$

- (3) And finally, given an interval $[i, j]$, let $\mathbf{I}^-[i, j]_\bullet$ be the ladder persistence module for which V_\bullet is trivial while W_\bullet is $\mathbf{I}[i, j]_\bullet$, so once again all vertical maps are 0:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \dots & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & \dots & \longrightarrow & 0 & \longrightarrow & \dots & \longrightarrow & 0 \\ & & & & \downarrow & & \downarrow & & & & \downarrow & & & & \\ & & & & \mathbb{F} & \longrightarrow & \mathbb{F} & \longrightarrow & \dots & \longrightarrow & \mathbb{F} & & & & \end{array}$$

It is readily seen that these three families of ladder persistence modules are mutually non-isomorphic and indecomposable. Therefore, by the Krull-Schmidt theorem, if ϕ_\bullet were to decompose as a direct sum of modules sourced from these three families, then such a decomposition would be unique. From such a decomposition, we can obtain the desired partial matching of the source and target bars: the presence of each $\mathbf{R}_{[i_1, j_1]}^{[i_2, j_2]}$ summand matches a bar $[i_2, j_2]$ of V_\bullet to a bar $[i_1, j_1]$ of W_\bullet , whilst the existence of $\mathbf{I}^+[i, j]_\bullet$ (or $\mathbf{I}^-[i, j]_\bullet$) summands reveals bars $[i, j]$ in V_\bullet (or W_\bullet) that are matched to 0. With this in mind, here is the main result of this section.

Theorem 5.3 *Let $(V_\bullet, W_\bullet, \phi_\bullet)$ be a ladder persistence module of length $\ell + 1$ where neither V_\bullet nor W_\bullet admit a pair of strictly nested bars. Then there are integers $r_{[i_1, j_1]}^{[i_2, j_2]}$ and $d_{ij}^\pm \in \mathbb{N}$ for which:*

$$(V_\bullet, W_\bullet, \phi_\bullet) \simeq \bigoplus_{[i_1, j_1] \preceq [i_2, j_2]} \left(\mathbf{R}_{[i_1, j_1]}^{[i_2, j_2]} \right)^{r_{[i_1, j_1]}^{[i_2, j_2]}} \oplus \bigoplus_{i \leq j} (\mathbf{I}^+[i, j]_\bullet)^{d_{ij}^+} \oplus \bigoplus_{i \leq j} (\mathbf{I}^-[i, j]_\bullet)^{d_{ij}^-}$$

(In Example 5.4 below we show that the assumption precluding nested bars is necessary.)

Proof The argument proceeds along three basic steps.

Step 1: Representing ϕ . Consider ordered barcode bases

$$\mathcal{B}_V := \{B_{V,i} \subset V_i \mid 0 \leq i \leq \ell\} \text{ and } \mathcal{B}_W := \{B_{W,i} \subset W_i \mid 0 \leq i \leq \ell\}$$

of V_\bullet and W_\bullet . Let $d_{i,j}^V$ and $d_{i,j}^W$ be the multiplicity of $[i, j]$ bars in the barcodes of V_\bullet and W_\bullet . We denote by b_\bullet the matrix representations of the maps ϕ_\bullet in these chosen bases. As in the proof Theorem 4.4, let $[i_1, j_1]$ and $[i_2, j_2]$ be two intervals in the barcode decomposition of (V_\bullet, f_\bullet) , and denote their corresponding sequences by $\{p_\bullet\}$ and $\{q_\bullet\}$. Using the commuting relations $\phi_k \circ f_k = h_k \circ \phi_{k-1}$, we have $\phi_k(p_k, q_k) = \phi_{k'}(p_{k'}, q_{k'})$, for all $k, k' \in [i, j] = [i_1, j_1] \cap [i_2, j_2]$, with this coefficient being zero unless $[i_1, j_1] \leq [i_2, j_2]$. Assuming this order relation holds, define $X_{[i_1, j_1]}^{[i_2, j_2]}$, to be the submatrix of b_i obtained by taking the d_{i_2, j_2}^V columns corresponding to basis vectors part of an $[i_1, j_1]$ bar of V , and the d_{i_1, j_1}^W rows corresponding to basis vectors part of an $[i_2, j_2]$ bar of W . From the above observation, $X_{[i_1, j_1]}^{[i_2, j_2]}$ is a submatrix of b_i, b_{i+1}, \dots, b_j , and is of dimension $d_{i_1, j_1}^W \times d_{i_2, j_2}^V$. Thus, the matrices b_\bullet are completely determined by the matrices $X_{[i_1, j_1]}^{[i_2, j_2]}$ and may therefore be represented as a single block matrix

$$\begin{bmatrix} X_{[0,0]}^{[0,0]} & X_{[0,0]}^{[0,1]} & \dots & X_{[0,0]}^{[0,\ell]} & 0 & \dots & 0 \\ 0 & X_{[0,1]}^{[0,1]} & \dots & \dots & X_{[0,1]}^{[1,\ell]} & \dots & 0 \\ \vdots & \vdots & \ddots & & & & \\ & & & \ddots & & & \\ & & & & \ddots & & \\ & & & & & X_{[\ell,\ell-1]}^{[\ell,\ell-1]} & X_{[\ell,\ell-1]}^{[\ell,\ell]} \\ 0 & 0 & & & & 0 & X_{[\ell,\ell]}^{[\ell,\ell]} \end{bmatrix}$$

This is a matrix of size $(\sum d_{i,j}^W) \times (\sum d_{i,j}^V)$.

Step 2: Admissible Operations. Given two ordered barcode bases \mathcal{B}_V and \mathcal{B}_W we may define a block matrix as above, which we denote $b(\mathcal{B}_V, \mathcal{B}_W)$. By Proposition 4.7, the set of ordered all barcode bases of (V_\bullet, f_\bullet) coincides precisely with the orbit $\text{Stab}(A(\mathcal{B}_1)_\bullet)\mathcal{B}_1$, where \mathcal{B}_1 is an ordered barcode basis. Then given $(h, k) \in \text{Stab}(A(\mathcal{B}_V)_\bullet) \times \text{Stab}(A(\mathcal{B}_W)_\bullet)$, we may consider $b(h\mathcal{B}_V, k\mathcal{B}_W)$. As seen in the proof of Theorem 4.4, we are able to completely characterise an element $h \in \text{Stab}(A(\mathcal{B}_V)_\bullet)$ in terms of the submatrices $h_{[i_1, j_1]}^{[i_2, j_2]}$, so that h may be represented as a single block matrix

$$\begin{bmatrix} h_{[0,0]}^{[0,0]} & h_{[0,0]}^{[0,1]} & \dots & h_{[0,0]}^{[0,\ell]} & 0 & \dots & 0 \\ 0 & h_{[0,1]}^{[0,1]} & \dots & \dots & h_{[0,1]}^{[1,\ell]} & \dots & 0 \\ \vdots & \vdots & \ddots & & & & \\ & & & \ddots & & & \\ & & & & h_{[\ell,\ell-1]}^{[\ell,\ell-1]} & h_{[\ell,\ell-1]}^{[\ell,\ell]} \\ 0 & 0 & & & 0 & h_{[\ell,\ell]}^{[\ell,\ell]} \end{bmatrix}$$

of size $(\sum d_{i,j}^V) \times (\sum d_{i,j}^W)$. The same is true for $k \in \text{Stab}(A(\mathcal{B}_W)_\bullet)$, whence $b(h\mathcal{B}_V, k\mathcal{B}_W)$ equals the (matrix) product $k \cdot b(\mathcal{B}_V, \mathcal{B}_W) \cdot h^{-1}$. Thus, we are only allowed to perform the following legal operations on $b(\mathcal{B}_V, \mathcal{B}_W)$:

- (1) Using invertible block diagonal elements $h_{[i_1, j_1]}^{[i_1, j_1]}$ and $k_{[i_1, j_1]}^{[i_1, j_1]}$, we may perform any operations between column and rows corresponding to $[i_1, j_1]$ bars in our block matrix.
- (2) Using the block matrices $h_{[i_1, j_1]}^{[i_2, j_2]}$, we see we may modify columns corresponding to $[i_2, j_2]$ bars using columns corresponding to $[i_1, j_1]$ in our block matrix, whenever $[i_1, j_1] \leq [i_2, j_2]$.
- (3) Using the block matrices $k_{[i_1, j_1]}^{[i_2, j_2]}$, we see we may modify rows corresponding to $[i_1, j_1]$ bars using columns corresponding to $[i_2, j_2]$ in our block matrix, whenever $[i_1, j_1] \leq [i_2, j_2]$.

Step 3: Matrix Reduction. We wish to find ordered barcode bases for which the corresponding matrix $b(\mathcal{B}_V, \mathcal{B}_W)$ admits at most one non-zero term 1 in each row and column. From this form, the desired decomposition can be easily extracted: every 1 in a row corresponding to an $[i_1, j_1]$ bar and column corresponding to an $[i_2, j_2]$ bar (with $[i_1, j_1] \leq [i_2, j_2]$) corresponds to an $\mathbf{R}_{[i_1, j_1]}^{[i_2, j_2]}$ summand. Similarly, zero rows and columns then yield $\mathbf{I}^-[i, j]_\bullet$ and $\mathbf{I}^+[i, j]_\bullet$ summands respectively.

Let $\mathcal{B}_V, \mathcal{B}_W$ be ordered barcode bases of V_\bullet, W_\bullet so that

$$b(\mathcal{B}_V, \mathcal{B}_W) = \begin{bmatrix} X_{[0,0]}^{[0,0]} & X_{[0,0]}^{[0,1]} & \dots & X_{[0,0]}^{[0,\ell]} & 0 & \dots & 0 \\ 0 & X_{[0,1]}^{[0,1]} & \dots & \dots & X_{[0,1]}^{[1,\ell]} & \dots & 0 \\ \vdots & \vdots & \ddots & & & & \\ & & & \ddots & & & \\ & & & & \ddots & & \\ & & & & & X_{[\ell,\ell-1]}^{[\ell,\ell-1]} & X_{[\ell,\ell-1]}^{[\ell,\ell]} \\ 0 & 0 & & & 0 & X_{[\ell,\ell]}^{[\ell,\ell]} \end{bmatrix}$$

We seek stabiliser elements (h, k) so that $B(h\mathcal{B}_V, k\mathcal{B}_W) = kB(\mathcal{B}_V, \mathcal{B}_W)h^{-1}$ is in barcode form. To this end, we perform basis changes using legal operations of type (1), (2) and (3). We process the column-blocks of this matrix from left to right, in each case starting from the diagonal block and working our way upwards. We will denote each treated matrix that has been put in adequate form by $P_{[i_1, j_1]}^{[i_2, j_2]}$.


That is, we start with $X_{[0,0]}^{[0,0]}$, putting it in Smith normal form using basis changes $h_{[0,0]}^{[0,0]}$ and $k_{[0,0]}^{[0,0]}$. Now assume we wish to treat $X_{[i_1, j_1]}^{[i_2, j_2]}$, where all matrices below it and to its left have been treated. That is, we have

$$\begin{bmatrix} 0 & \dots & P_{[i_1, j_1]}^{[i_1, j_1]} & \dots & X_{[i_1, j_1]}^{[i_2, j_2]} \\ & & & & \vdots \\ & & & & P_{[i_2, \ell]}^{[i_2, j_2]} \\ & & & & 0 \\ & & & & \vdots \\ & & & & 0 \end{bmatrix}$$

Given non-intersecting bars $[a, b] \preceq [c, d]$, we have either $[a, b] \preceq [c, d]$ or $[a, b] \subset [c, d]$. Then by hypothesis, given $P_{[i_1, j_1]}^{[a, b]}$ to the left of $X_{[i_1, j_1]}^{[i_2, j_2]}$, we have $[a, b] \preceq [i_2, j_2]$. So we may zero out rows of $X_{[i_1, j_1]}^{[i_2, j_2]}$ in which $P_{[i_1, j_1]}^{[a, b]}$ has 1's using operations of type (2). Similarly, given $P_{[a, b]}^{[i_2, j_2]}$ below $X_{[i_1, j_1]}^{[i_2, j_2]}$, we may zero out corresponding columns using operations of type (3). The non-zero columns of the resulting matrix $\tilde{X}_{[i_1, j_1]}^{[i_2, j_2]}$ have 0's below them, and non-zero rows have 0's to their left.

Then using basis changes $h_{[i_2, j_2]}^{[i_2, j_2]}$ and $k_{[i_1, j_1]}^{[i_1, j_1]}$, we may put A in Smith normal form, without adding non-zero terms in any rows below and column to its left, preserving the desired structure. □

Example 5.4 We illustrate the difficulties imposed by strictly nested bars in the context of Theorem 5.3. Consider the map $\phi_\bullet : V_\bullet \mapsto W_\bullet$ where V_\bullet has barcode



and W_\bullet has barcode




with associated ordered barcode bases $\mathcal{B}_V, \mathcal{B}_W$, and ϕ_\bullet is given by the block matrix representation

$$b(\mathcal{B}_V, \mathcal{B}_W) = \begin{bmatrix} [1, 4] & [2, 3] \\ [1 & 1] & [0, 3] \end{bmatrix}$$

Since $[2, 3] \subset [1, 4]$, stabiliser changes of basis for \mathcal{B}_V are invertible diagonal matrices $\begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix}$, and stabiliser changes of basis for \mathcal{B}_W are invertible matrices $\begin{bmatrix} c \end{bmatrix}$. As such, $(V_\bullet, W_\bullet, \phi_\bullet)$ which will never be expressible as a direct sum of modules as in Theorem 5.3, since there is no change of basis which will allow us to transform this matrix into either $\begin{bmatrix} 0 & 1 \end{bmatrix}$ or $\begin{bmatrix} 1 & 0 \end{bmatrix}$. The same is true if V_\bullet has barcode



and W_\bullet has barcode



with associated ordered barcode bases $\mathcal{B}_V, \mathcal{B}_W$, and ϕ_\bullet is given by the block matrix representation

$$\begin{bmatrix} [1, 4] \\ 1 \\ 1 \end{bmatrix} \begin{bmatrix} [0, 3] \\ [1, 2] \end{bmatrix}$$

Thus, if the nested condition is violated on either V_\bullet or W_\bullet , a decomposition as in Theorem 5.3 is not always possible.

Remark 5.5 The non-nestedness hypothesis on the bars of V_\bullet and W_\bullet from Theorem 5.3 is quite restrictive. There are, however, several scenarios of interest where it is satisfied:

- (1) The 0-th persistent homology of a point cloud satisfies the hypothesis because all bars have left endpoint 0.
- (2) Similarly, the 1st persistent homology of a filtered graph admits no strictly nested bars because all bars have right endpoint ∞ .
- (3) More generally, the n -th persistent homology of an n -dimensional filtered complex satisfies the non-nestedness criterion. One may consider, for instance, the *Linial-Meshulam* model of random simplicial complexes (Linial and Meshulam 2006). A random simplicial complex chosen from this model on a given vertex set consists of every possible simplex of dimension $< n$, with candidate n -simplices being included independently with uniform probability $p \in [0, 1]$. The n -th persistent homology of any filtration of such a random complex satisfies the hypothesis.

Theorem 5.3 applies in all such cases.

6 Zigzag modules

In this section, we wish to generalise the previous work to representations of any type A quiver, in other words zigzag persistence modules. The nomenclature we adopt here is extracted from (Carlsson and de Silva 2010). A zigzag module is given by a sequence

$$V_0 \xleftarrow{p_1} V_1 \xleftarrow{p_2} \dots \xleftarrow{p_{\ell-1}} V_{\ell-1} \xleftarrow{p_\ell} V_\ell$$

where each $\xleftarrow{p_i}$ is either a forward map $\xrightarrow{f_i}$ or a backwards map $\xleftarrow{q_i}$. The direction of the arrows define the **type** τ of the zigzag module, which is the direction of the arrows of the underlying type A quiver. For example,

$$V_0 \xrightarrow{f_1} V_1 \xleftarrow{q_2} V_2$$

has type $\tau = fq$. We denote zigzag modules as $(V_\bullet, p_\bullet, \tau)$.

The **interval module** of type τ corresponding to a pair of non-negative integers $i \leq j$ is the zigzag module $\mathbf{I}_\tau[i, j]_\bullet$ given by

$$0 \longleftrightarrow \dots \longleftrightarrow 0 \longleftrightarrow \mathbb{F} \longleftrightarrow \dots \longleftrightarrow \mathbb{F} \longleftrightarrow 0 \longleftrightarrow \dots \longleftrightarrow 0,$$

where the contiguous string of \mathbb{F} 's spans $\{i, i + 1, \dots, j - 1, j\}$, all intermediate $\mathbb{F} \longleftrightarrow \mathbb{F}$ maps are identities in the direction depending on τ , and all other vector spaces are trivial. Zigzag modules also decompose into interval modules: this follows from the main result of Gabriel (1972), and is established more directly in Carlsson and de Silva (2010). In particular, every zigzag module $(V_\bullet, p_\bullet, \tau)$ is isomorphic to a direct sum

$$(V_\bullet, p_\bullet, \tau) \cong \bigoplus_{0 \leq i \leq j \leq \ell} (\mathbf{I}_\tau[i, j])^{d_{ij}},$$

for some uniquely determined $d_{ij} \in \mathbb{N}$.

Definition 6.1 An $m \times n$ matrix A of rank r is in **reversed barcode form** if there exists a strictly increasing function $c : \{m - r + 1, \dots, m\} \rightarrow \{1, \dots, n\}$ so that

$$A_{ij} = \begin{cases} 1 & \text{if } j = c(i), \\ 0 & \text{otherwise.} \end{cases}$$

Thus, a matrix is in reversed barcode form whenever its entries lie in $\{0, 1\}$, with at most one non-zero term in each row and column, and satisfies the following additional requirement: the r non-zero terms appear in the last r columns, with increasing row order. (We warn the reader that if a matrix A is in barcode form, then its transpose will not in general be in reversed barcode form; however, the off-diagonal transpose of A will be in reversed barcode form).

Definition 6.2 A basis family \mathcal{B} for a zigzag module is called an **barcode basis** for $(V_\bullet, p_\bullet, \tau)$ if all of the A_i for which $V_{i-1} \xrightarrow{f_i} V_i$ are in barcode form, and all the A_i for which $V_{i-1} \xleftarrow{q_i} V_i$ are in reversed barcode form. The corresponding matrices A_\bullet are then said to be in **zigzag barcode form**.

We note that this notion coincides with Definition 2.2 whenever all maps in sight are forward. Barcode bases for zigzag modules are the natural bases arising from their decomposition into interval modules.

6.1 Algorithm for zigzag modules

We wish to generalise our algorithm from Sect. 3 to treat the case of zigzag modules. Instead of presenting concrete algorithms, we will adapt our proof of Proposition 3.4 in Sect. 3 to the more general setting. Implementing the algorithm is then achieved in similar fashion to what was done in Sect. 3 for classical persistence modules. To this end, we fix a zigzag module $(V_\bullet, p_\bullet, \tau)$ expressed as a sequence of matrices A_\bullet with respect to an arbitrary (i.e., not necessarily barcode) basis family \mathcal{B} :

$$\mathbb{F}^{n_0} \xleftarrow{A_1} \mathbb{F}^{n_1} \xleftarrow{A_2} \dots \xleftarrow{A_{\ell-1}} \mathbb{F}^{n_{\ell-1}} \xleftarrow{A_\ell} \mathbb{F}^{n_\ell} \tag{11}$$

The matrices A_i are of dimension either $n_i \times n_{i-1}$ or $n_{i-1} \times n_i$, depending on the type τ of V_\bullet . Analogously to Sect. 2, we may define X_τ to be the set of all the possible matrix-sequences A_\bullet which can arise in (11). It is a (strict) subset

$$X_\tau \subset \prod_{i=1}^\ell Y_i, \quad \text{where } Y_i = \begin{cases} \text{Mat}(n_i \times n_{i-1}; \mathbb{F}) & \text{if } i-1 \longrightarrow i \\ \text{Mat}(n_{i-1} \times n_i; \mathbb{F}) & \text{if } i-1 \longleftarrow i \end{cases}$$

Changes of bases for zigzag modules are obtained by a new action of the group G from (5) on the set X_τ ; the major difference between this action and the one treated in Sect. 3 is that the (direction of) conjugation now depends on the type τ . Explicitly, if $V_{i-1} \xrightarrow{f_i} V_i$ points forward, then it gets sent to $g_i \circ f_i \circ g_{i-1}^{-1}$ as before; and conversely, if $V_{i-1} \xleftarrow{q_i} V_i$ points backwards, then it is sent to $g_{i-1} \circ q_i \circ g_i^{-1}$. For this action, we obtain the following analogue of Lemma 3.3.

Lemma 6.3 Assume that the first $\ell - 1$ matrices of $\{A_i \mid 1 \leq i < \ell\}$ from (11) are in zigzag barcode form.

- If $f_\ell : V_{\ell-1} \rightarrow V_\ell$, and the last matrix A_ℓ has a pivot in the (r, q) position³, together with a nonzero entry $\alpha := A_\ell(r, p)$ in the same row r but some other column $p > q$, then there exists $g \in G$ with $g_\ell = \text{Id}$ so that $(gA)_\bullet$ equals A_\bullet except A_ℓ where the α entry is replaced by zero.

³ i.e., we have $A_\ell(r, q) = 1$ while all other entries in the q -th column of A_ℓ are zero.

- If $q_\ell : V_{\ell-1} \leftarrow V_\ell$, and the last matrix A_ℓ has a pivot in the (r, q) position, together with a nonzero entry $\alpha := A_\ell(s, q)$ in the same column q but some other row $s < r$, then there exists $g \in G$ with $g_\ell = \text{Id}$ so that $(gA)_\bullet$ equals A_\bullet except A_ℓ where the α entry is replaced by zero.

Proof The proof is done by induction, in similar fashion to that of Lemma 3.3. The case $\ell = 1$ remains trivial. To prove the induction, 4 cases should now be considered.

Case 1: $V_{\ell-2} \rightarrow V_{\ell-1} \rightarrow V_\ell$

This case is precisely that of Lemma 3.3, and the proof remains, the same.

Case 2: $V_{\ell-2} \leftarrow V_{\ell-1} \rightarrow V_\ell$

Performing $\mathbf{C}_{p \leftarrow q}(-\alpha)$ on A_ℓ induces the same operation on $A_{\ell-1}$. If the q -th column of $A_{\ell-1}$ is identically zero, this operation leaves $A_{\ell-1}$ unchanged. Otherwise, since $p > q$ and $A_{\ell-1}$ is in reversed barcode form, if the q -th column isn't zero then the p -th column is also non-zero. Then the resulting matrix $A'_{\ell-1}$ has the form

$$A'_{\ell-1} = \begin{bmatrix} & q & & & & p & \\ 0 & 1 & 0 & \cdots & 0 & -\alpha & 0 \end{bmatrix} \begin{matrix} c \\ \\ \\ \\ \\ d \end{matrix}$$

where we may again use induction.

Case 3: $V_{\ell-2} \rightarrow V_{\ell-1} \leftarrow V_\ell$

We must perform the row operation $\mathbf{R}_{s \leftarrow r}(-\alpha)$ on A_ℓ , inducing the same row operation on $A_{\ell-1}$. Again, if the r -th row of $A_{\ell-1}$ is identically zero, we are done. Otherwise, since A_ℓ is in barcode form and $s < r$, if the r -th row isn't zero then the s -th row isn't zero, and so the resulting matrix $A'_{\ell-1}$ has the form

$$A'_{\ell-1} = \begin{bmatrix} & c & & & & d & \\ 0 & 1 & 0 & \cdots & 0 & -\alpha & 0 \end{bmatrix} \begin{matrix} s \\ \\ \\ \\ \\ r \end{matrix}$$

where we may again use induction.

Case 4: $V_{\ell-2} \leftarrow V_{\ell-1} \leftarrow V_\ell$

We perform the row operation $\mathbf{R}_{s \leftarrow r}(-\alpha)$ on A_ℓ , inducing the column operation $\mathbf{C}_{r \leftarrow s}(\alpha)$ on $A_{\ell-1}$. Again, if the s -th column of $A_{\ell-1}$ is zero, we are done. Otherwise, since $A_{\ell-1}$ is in barcode form and $s < r$, the r -th column is also non-zero so that the

resulting matrix $A'_{\ell-1}$ has the form

$$A'_{\ell-1} = \begin{bmatrix} & s & & & r & & \\ 0 & 1 & 0 & \cdots & 0 & \alpha & 0 \\ & 0 & & & & 0 & \\ & \vdots & & & & \vdots & \\ 0 & 0 & 0 & \cdots & & 0 & \\ 0 & 0 & 0 & \cdots & & 1 & 0 \end{bmatrix} \begin{matrix} c \\ \\ \\ \\ d \end{matrix}$$

where we may again use induction. □

The zigzag-compatible avatar of Proposition 3.4 is given as follows. As before, we regard this result as the 'matrix version' of the decomposition theorem for zigzag persistence modules.

Proposition 6.4 *Given the sequence of matrices A_\bullet as in (11), there is a $g \in G$ such that $(gA)_\bullet$ has all its matrices in zigzag barcode form.*

Proof We proceed by induction. The case $\ell = 1$ is trivial. We can reduce ourselves by induction hypothesis to the case where the first $\ell - 1$ matrices are in zigzag barcode form. If $V_{\ell-1} \rightarrow V_\ell$ we perform row operations on A_ℓ to put it in reduced row echelon form. Otherwise, if $V_{\ell-1} \leftarrow V_\ell$, we perform column operations to put A_ℓ in reversed reduced column echelon form. That is, we put A_ℓ in the form

0	★	★	★
	1	0	0
	⋮				
0	★	★	★		
	1	0	0		
0			★	★	
			1	0	
0				★	
				1	
	0				

This is achieved through a slight tweak to the standard Gaussian algorithm for placing matrices in column echelon form, where one starts with the last row and works upwards. Finally, we may apply Lemma 6.3 to the non-zero ★ term of A_ℓ obtain the desired result. □

6.2 Barcode bases of zigzag modules


In order to characterise the set of barcode bases of a zigzag module, we must again attempt to characterise the set $\text{Stab}(A_\bullet)$ corresponding to linear maps of a zigzag module

in a barcode basis. In Sect. 4, we defined an order \leq on the set of intervals that allowed us to classify the stabiliser as

$$\text{Stab}(A_\bullet) \cong \prod_{[i,j]} \text{GL}(d_{ij}; \mathbb{F}) \times \prod_{[i_1,j_1] \leq [i_2,j_2]} \text{Mat}(d_{i_1,j_1} \times d_{i_2,j_2}; \mathbb{F}).$$

The $\text{Mat}(d_{i_1,j_1} \times d_{i_2,j_2}; \mathbb{F})$ came from the fact that in standard persistence, the interval module $\mathbf{I}[i_1, j_1]$ can be mapped non-trivially to the interval $\mathbf{I}[i_2, j_2]$ if and only if $[i_1, j_1] \leq [i_2, j_2]$. In zizag persistence, this is no longer true.

Example 6.5 One checks that in the following scenario



the interval $[1, 4]$ may be non-trivially mapped to the interval $[2, 3]$.

To then obtain a similar classification result for zizag modules, we see we must adapt our partial order \leq to the type τ of our zizag module.

Definition 6.6 Let τ be a type of zizag module, defining an orientation on the standard length ℓ quiver. Let \leq_τ be the binary relation on $\{[i, j] \in \mathbb{Z}^2 \mid 0 \leq i \leq j \leq \ell\}$ given by

$$[i_1, j_1] \leq_\tau [i_2, j_2] \Leftrightarrow \begin{cases} [i_1, j_1] \cap [i_2, j_2] = [i, j] \neq \emptyset \text{ and} \\ i_1 \leq i_2 \text{ if } i - 1 \rightarrow i, \quad i_2 \leq i_1 \text{ if } i - 1 \leftarrow i \text{ and} \\ j_1 \leq j_2 \text{ if } j \rightarrow j + 1, \quad j_2 \leq j_1 \text{ if } j \leftarrow j + 1 \end{cases}$$

We observe that when all maps point forward, i.e., when $\tau = ff \dots f$, we have $\leq_\tau = \leq$. And if all maps point backwards, ie $\tau = qq \dots q$, then \leq_τ is the reverse of \leq in the sense that

$$[i_1, j_1] \leq_\tau [i_2, j_2] \Leftrightarrow [i_2, j_2] \leq [i_1, j_1].$$

Our description of the stabiliser from Theorem 4.4 relied on the compatibility of \leq with the lexicographical total order \leq on the bars. Having produced a τ -analogue of \leq , we must now construct the zigzag version of \leq . To this end, we define two new auxiliary total orders on the set of all possible endpoints $\{0, \dots, \ell\}$. Note that any such order $<$ amounts to a choice of element σ_ℓ lying in the permutation group $S_{\ell+1}$ via the identification

$$\sigma(0) < \sigma(1) < \dots < \sigma(\ell).$$

Definition 6.7 Let \leq_τ be the total order on $\{0, 1 \dots \ell\}$ given by the permutation σ_ℓ defined inductively as follows. Assuming we have ordered $\{0, \dots, i\}$ with corresponding permutation σ_i , we order $\{0, \dots, i + 1\}$ with permutation σ_{i+1} , setting

$$\sigma_{i+1}(k) = \begin{cases} \sigma_i(k) & \text{if } i \longrightarrow i + 1 \text{ for } k \in \{0, \dots, i\} \\ i + 1 & \text{if } i \longrightarrow i + 1 \text{ for } k = i + 1 \\ \sigma_i(k) + 1 & \text{if } i \longleftarrow i + 1 \text{ for } k \in \{0, \dots, i\} \\ 0 & \text{if } i \longleftarrow i + 1 \text{ for } k = i + 1 \end{cases}$$

Definition 6.8 Let \leq_τ^* be the total order on $\{0, 1 \dots, \ell\}$ given by the permutation σ_ℓ^* defined inductively as follows. Assuming we have ordered $\{j + 1, \dots, \ell\}$ with corresponding permutation σ_{j+1}^* , we order $\{j, \dots, \ell\}$ with permutation σ_j^* , setting

$$\sigma_j^*(k) = \begin{cases} \sigma_{j+1}^*(k) & \text{if } j \longrightarrow j + 1 \text{ for } k \in \{j + 1, \dots, \ell\} \\ j & \text{if } j \longrightarrow j + 1 \text{ for } k = j \\ \sigma_{j+1}^*(k) - 1 & \text{if } j \longleftarrow j + 1 \text{ for } k \in \{j + 1, \dots, \ell\} \\ \ell & \text{if } j \longleftarrow j + 1 \text{ for } k = j \end{cases}$$

If $\tau = ff \dots f$, both of the above total orders coincide with the standard ordering \leq on $\{0, 1 \dots \ell\}$.

Example 6.9 Consider zigzag modules of length 4 with type $0 \longleftarrow 1 \longrightarrow 2 \longleftarrow 3$. For such modules, the two total orders defined above are

$$3 \leq_\tau 1 \leq_\tau 0 \leq_\tau 2, \text{ and} \\ 1 \leq_\tau^* 3 \leq_\tau^* 2 \leq_\tau^* 0.$$

Definition 6.10 Let τ be a type of length- ℓ zigzag module. The corresponding total order \leq_τ on $\{[i, j] \in \mathbb{Z}^2 \mid 0 \leq i \leq j \leq \ell\}$ is given by

$$[a, b] \leq_\tau [c, d] \Leftrightarrow a <_\tau c \text{ or } a = c \text{ and } b \leq_\tau^* d.$$

By construction, if $\tau = ff \dots f$, this coincides with the lexicographic order \leq from Sect. 4, and $[i_1, j_1] \leq_\tau [i_2, j_2] \Leftrightarrow [i_1, j_1] \leq_\tau [i_2, j_2]$. Using this total order on the bars, we may generalise our notion of **ordered barcode bases** to zigzag modules, by ordering the bars with the order \leq_τ . We now have all the necessary tools to adapt our results from Sect. 4 to the case of zigzag modules.


Theorem 6.11 For each pair $[i, j]$ in $\{0, 1, \dots, \ell\}$ with $i \leq j$, let d_{ij} be the multiplicity in the barcode of $(V_\bullet, p_\bullet, \tau)$, with the understanding that $d_{ij} = 0$ whenever $[i, j]$ is not in $\mathbf{Bar}(V_\bullet, p_\bullet, \tau)$. Then there is a bijection of sets:

$$\text{Stab}(A_\bullet) \cong \prod_{[i, j]} \text{GL}(d_{ij}; \mathbb{F}) \times \prod_{[i_1, j_1] \leq_\tau [i_2, j_2]} \text{Mat}(d_{i_1 j_1} \times d_{i_2 j_2}; \mathbb{F}).$$

Proposition 6.12 *Given a zigzag module $(V_\bullet, p_\bullet, \tau)$ together with an ordered barcode basis \mathcal{B} with matrix representation A_\bullet , the set of all ordered barcode bases is given by the orbit $\text{Stab}(A_\bullet)\mathcal{B}$.*

6.3 Maps of zigzag modules

We now turn our attention to morphisms $(V_\bullet, p_\bullet, \tau) \rightarrow (W_\bullet, \tilde{p}_\bullet, \tau)$ between zigzag persistence modules of the same type τ . Each such morphism is determined by linear maps $\phi_i : V_i \mapsto W_i$ along with the requirement that the evident squares commute. That is, if $p_i = f_i$ (and so $\tilde{p}_i = \tilde{f}_i$), we have $\phi_i \circ f_i = \tilde{f}_i \circ \phi_{i-1}$, and if $p_i = q_i$ (and so $\tilde{p}_i = \tilde{q}_i$), we have $\phi_{i-1} \circ q_i = \tilde{q}_i \circ \phi_i$. This is best represented through the following diagram



where each natural square commutes.

As in Sect. 5, such maps may be identified as ladder persistence module on the rectangle quiver with orientation τ . We denote such modules $(V_\bullet, W_\bullet, \phi_\bullet, \tau)$. We may analogously define a special class of modules, namely $\mathbf{R}_{\tau[i_1, j_1]^\bullet}^{[i_2, j_2]}$, $\mathbf{I}_\tau^+[i, j]_\bullet$ and $\mathbf{I}_\tau^-[i, j]_\bullet$. Here is the zigzag analogue of Definition 5.1

Definition 6.13 *Given a type τ of zigzag module, we say a bar $[i_2, j_2]$ is strictly nested in a bar $[i_1, j_1]$ with regards to τ , denoted $[i_2, j_2] \subset_\tau [i_1, j_1]$, if they are non-intersecting with $[i_1, j_1] \leq_\tau [i_2, j_2]$ but $[i_1, j_1] \not\leq_\tau [i_2, j_2]$.*

As in Sect. 5, maps of zigzag module may be compactly represented as block upper triangular matrices using ordered barcode basis for the source and target and zigzag modules. Having excluded strictly nested bars, we are free to perform operations of type (1), (2) and (3) à la Theorem 5.3 and obtain the following result.

Theorem 6.14 *Let $(V_\bullet, W_\bullet, \phi_\bullet, \tau)$ be a ladder persistence module of length $\ell + 1$ and type τ , where neither V_\bullet nor W_\bullet admit a pair of strictly nested bars with regards to τ . Then there are integers $r_{[i_1, j_1]^\bullet}^{[i_2, j_2]}$, $d_{ij}^\pm > 0$ for which :*

$$(V_\bullet, W_\bullet, \phi_\bullet, \tau) \simeq \bigoplus_{[i_1, j_1] \leq_\tau [i_2, j_2]} (\mathbf{R}_{\tau[i_1, j_1]^\bullet}^{[i_2, j_2]})^{r_{[i_1, j_1]^\bullet}^{[i_2, j_2]}} \oplus \bigoplus_{i \leq j} (\mathbf{I}_\tau^+[i, j]_\bullet)^{d_{ij}^+} \oplus \bigoplus_{i \leq j} (\mathbf{I}_\tau^-[i, j]_\bullet)^{d_{ij}^-}$$

Acknowledgements The authors are members of the Centre for Topological Data Analysis, funded by the EPSRC grant EP/R018472/1. We are grateful to the two anonymous referees for their helpful comments and corrections.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give

appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Asashiba, H., Escolar, E.G., Hiraoka, Y., Takeuchi, H.: Matrix method for persistence modules on commutative ladders of finite type. *Jpn. J. Ind. Appl. Math.* **36**(1), 97–130 (2018)
- Buchet, M., Escolar, E.G.: Realizations of indecomposable persistence modules of arbitrarily large dimension. In: 34th International Symposium on Computational Geometry, volume 99 of LIPIcs. Leibniz Int. Proc. Inform., pages Art. No.15, 13. Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern, (2018)
- Brion, M.: Representations of quivers. *Geometric methods in representation theory I*, 104–144 (2012)
- Carlsson, G., Dwaraknath, A., Nelson, B.J.: Persistent and zigzag homology: a matrix factorization viewpoint. [arXiv:1911.10693](https://arxiv.org/abs/1911.10693), (2019)
- Carlsson, G., de Silva, V.: Zigzag persistence. *Found. Comput. Math.* **10**(4), 367–405 (2010)
- Escolar, E.G., Hiraoka, Y.: Persistence modules on commutative ladders of finite type. *Discrete Comput. Geom.* **55**(1), 100–157 (2016)
- Gabriel, P.: Unzerlegbare darstellungen I. *Manuscripta mathematica* **6**, 71–104 (1972)
- Gregorio, A De., Guerra, M., Scaramuccia, S., Vaccarino, F.: Parallel decomposition of persistence modules through interval bases, (2021)
- Hang, H., Giusti, C., Ziegelmeier, L., Henselman-Petrusek, G.: U-match factorization: sparse homological algebra, lazy cycle representatives, and dualities in persistent (co)homology, (2021)
- Hang, H., Mio, W.: Correspondence modules and persistence sheaves: a unifying perspective on one-parameter persistent homology. [arXiv:2006.08557](https://arxiv.org/abs/2006.08557), (2020)
- Henselman-Petrusek, G.: Matroids and canonical forms: theory and applications. [arXiv:1710.06084](https://arxiv.org/abs/1710.06084), (2017)
- Linial, N., Meshulam, R.: Homological connectivity of random 2-complexes. *Combinatorica* **26**, 475–487 (2006)
- Zomorodian, A., Carlsson, G.E.: Computing persistence homology. *Discrete Comput Geom*, 33(2), (2005)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.