



EMPIRICALLY GROUNDED CLINICAL GUIDANCE PAPER

# Ten misconceptions about CBT for psychosis

Katherine Newman-Taylor<sup>1,2</sup> , Louise Isham<sup>3,4</sup>  and Alison Brabban<sup>5</sup>

<sup>1</sup>Psychology Department, University of Southampton, Southampton, UK, <sup>2</sup>Psychology Department, Southern Health NHSF Trust, Southampton, UK, <sup>3</sup>Oxford Cognitive Therapy Centre, Oxford Health NHS Foundation Trust, Oxford, UK, <sup>4</sup>Oxford Cognitive Approaches to Psychosis, Department of Experimental Psychology, University of Oxford, Oxford, UK and <sup>5</sup>Psychology Department, Tees, Esk and Wear Valleys NHS Foundation Trust, Durham, UK

**Corresponding author:** Katherine Newman-Taylor; Email: [k.newman-taylor@soton.ac.uk](mailto:k.newman-taylor@soton.ac.uk)

(Received 25 May 2025; revised 25 July 2025; accepted 28 July 2025)

## Abstract

CBT for psychosis is an established and evolving psychological therapy. Historical controversies about the nature of psychosis persist, and more recent debates about the outcome literature lack precision, muddying the waters further. Based on our experience as clinicians, teachers and supervisors, and following NHS and national lead roles, we describe ten common misconceptions about CBT for psychosis. These include misconceptions about the evidence, the focus of therapy, ‘thinking positively’, and the nature of collaboration and the therapeutic relationship. We refute these misconceptions based on current theory, research, and best practice guidelines. We highlight the need to get out of the clinic room, measure the impact of therapy on personal recovery and autonomy, and meet training and governance requirements. It is essential that clinicians, service leads, and our professional bodies uphold core standards of care if people with psychosis are to have access to high quality CBT of the standard we would be happy to see offered to our own family and friends.

## Key learning aims

- (1) To recognise common misconceptions about CBT for psychosis.
- (2) To counter these misconceptions theoretically and empirically – to inform ourselves, colleagues and service leads committed to ensuring high quality CBT for psychosis.
- (3) To highlight statutory and professional body responsibilities to ensure parity of esteem for people with psychosis, who deserve high quality, ‘full dose’ treatments delivered by appropriately trained clinicians, and supported by robust governance systems, just as we would expect for people with physical health conditions.

**Keywords:** CBT; psychosis; training

## Introduction

Developments in the psychological understanding and treatment of psychosis represent one of the most significant advances in mental health care of the last century. Jaspers (1959) viewed psychosis as ‘*un-understandable*’ – on the other side of the ‘*abyss*’ that separates what we can and cannot comprehend. Psychosis was distinguished from the neuroses, and deemed a purely neurological condition unrelated to a person’s life experiences, requiring medical intervention. Psychological therapy was considered inappropriate and potentially harmful – clinicians were

advised not to talk to people about their voices and delusions as ‘colluding’ could worsen the illness. Cases where psychotic experiences might be linked to adversity or trauma were (and often still are) not considered to be true psychosis.

A considerable body of evidence now contradicts these early assumptions and shows that we can understand and treat psychosis psychologically; Cognitive Behavioural Therapy (CBT) and Family Intervention (FI) are recommended for people with psychosis in many Western countries (e.g. Galletly *et al.*, 2016; National Institute for Health and Care Excellence [NICE], 2014). Nevertheless, historical views live on, and people often assume purely biomedical explanations for psychosis as caused by chemical imbalances in the brain amenable only to medication. Clinical practice lags well behind the evidence base, despite empirical support for psychological theories and therapies accumulating over the last 30 years.

Following an early case study (Beck, 1952), evidence for CBT for psychosis (CBTp) emerged in the 1980s and 1990s. Initial studies showed that contrary to the prevailing clinical wisdom, (1) it is safe to talk to people about their psychosis (Kingdon and Turkington, 1991), and (2) psychological therapy can be beneficial (Chadwick and Birchwood, 1994; Garety *et al.*, 1994; Nelson, 1997; Tarrier *et al.*, 1993). From this point, CBTp evolved rapidly, from promoting coping strategies to developing a detailed understanding (formulation) of the psychological processes likely to contribute to the development and maintenance of distressing voices, images and delusional beliefs that could be targeted in therapy.

CBTp focuses on reducing distress and pre-occupation with psychotic symptoms, and improving functioning and quality of life. Offering therapy to a person who experiences the world, rather than the self, as changed, is a delicate business, and requires us to shape therapy and therapeutic interactions to the person rather than expecting the person to fit expectations about ‘suitability’. For this reason, CBTp is explicitly flexible, targeting therapeutic goals through different levels of belief change – from no change in delusional appraisals (where the focus is on improving mood and coping skills) to complete change in belief conviction (see Johns *et al.*, 2014). This flexibility to work within the person’s belief system means that CBTp can benefit many people with psychosis, including those who are in hospital, use substances, or ‘lack insight’. Indeed, at the start of the twenty-first century, the evidence was strong enough for the UK treatment guideline on schizophrenia (NICE, 2002) to recommend that everyone with psychosis be offered CBTp, the only exceptions being those unable to engage in an informed discussion about their treatment.

A quarter of a century on, the availability of CBTp remains sparse. The UK NHS Mental Health Services Data Set indicates that less than 2% of people supported by community mental health teams in England have accessed *any NICE recommended therapy*, including CBTp. Notwithstanding quality issues common to nationally mandated datasets, this is a damning indictment of current healthcare provision. Furthermore, also in the UK, people from many minoritised ethnic groups are less likely to be offered CBTp than those who are White British (Schlief *et al.*, 2023). Our failure to deliver evidence-based treatments is likely to be due to a combination of limited resource (a shortage of appropriately trained therapists with time protected to deliver therapy), systemic service barriers (Birchwood *et al.*, 2013; Schlief *et al.*, 2023), and people’s own hesitancy to seek and accept help, often linked to stigma and other psychosocial factors (Skrobinska *et al.*, 2024). When people do access services, the quality of psychological therapy provided is variable – CBTp done well can be life changing while CBTp done poorly may be harmful (Goldsmith *et al.*, 2015). Would we be satisfied with limited provision and sub-optimal doses of chemotherapy for cancer patients? If we are to ensure parity of esteem for people with severe mental ill-health, in terms of both availability and quality of provision, we need to address the individual, social and structural barriers to routine access to high quality CBTp and other therapies likely to benefit people with psychosis.

### Misconception 1: CBTp doesn't work

The UK NICE guideline (2002; 2009; 2014) recommends CBTp be offered to everyone with psychosis, regardless of the phase of illness. Despite meta-analyses supporting this recommendation, and a recent umbrella review (Solmi *et al.*, 2023), there is a persistent distrust of the guidance and a perception that the evidence has been overstated (e.g. Jauhar *et al.*, 2014). This scepticism is likely to be due in large part to the variation in effect sizes across meta-analyses conducted over the last two decades. With effects ranging from very small or negligible to moderate, some have interpreted this variability as indicating that the evidence is unreliable or exaggerated. This has been discussed at length (e.g. Birchwood *et al.*, 2014b; Peters, 2014), with proponents of CBTp attributing the range of outcomes to heterogeneity of (1) symptoms that have been grouped together (e.g. hallucinations, delusions, and negative symptoms), (2) therapies (e.g. first generation CBTp<sup>1</sup>, avatar therapy, and problem-specific CBTp), and (3) primary outcomes – do we focus on symptom reduction, as is typical in meta-analyses, or associated distress and personal recovery, which are targeted in psychological therapies? Additionally, we'd argue that psychological therapies should be evaluated in addition to medication effects as most people receiving CBTp also take anti-psychotics.

Taking account of these issues yields more nuanced insights. For example, CBTp tends to be more effective for delusions than hallucinations or negative symptoms (Sitko *et al.*, 2020), when interventions target mechanisms underlying specific symptoms (e.g. Freeman *et al.*, 2021), and when assessing emotional and behavioural outcomes (e.g. Birchwood *et al.*, 2014a). For example, an early trial of CBT for command hallucinations showed that while participants continued to hear voices, compliance reduced considerably; at 18 months, just 28% of the CBTp group still fully complied with their voices, compared with 46% of the control group (OR 0.45, 95% CI 0.23–0.88,  $p = 0.021$ ). In further support of the cognitive model, this key behaviour change was mediated by a reduction in strength of belief about the power of the voice (Birchwood *et al.*, 2014a).

A meta-analyses completed in 2020 showed that CBTp was an effective treatment with a small, pooled effect size (e.g. Cohen's  $d = 0.26$  for hallucinations, 0.36 for delusions; Sitko *et al.*, 2020), comparable to some anti-psychotics (Huhn *et al.*, 2019). While this may seem unimpressive compared with CBT for other presentations such as social anxiety and PTSD which yield large effects (NICE, 2005; 2013), CBTp is still a relatively young intervention and will evolve as our understanding of psychosis deepens. For example, recent research aimed at reducing delusional conviction and distress by targeting specific mechanisms contributing to persecutory delusions yielded large effects at end of treatment (Cohen's  $d = 0.86$  and 1.20, respectively) compared with befriending (Freeman *et al.*, 2021).<sup>2</sup>

Targeting distinct causal mechanisms for different psychotic experiences holds much promise. Developments in the broader psychotherapy outcome research could also improve CBTp treatment effects. Analyses of large datasets reveal heterogenous outcomes in therapies for depression and anxiety based on pre-therapy factors (e.g. Lorenzo-Luaces *et al.*, 2017). Personalised therapy decisions informed by machine learning could clarify what works best for whom (cf. Paul, 1967), and may dramatically change psychotherapy outcome research in the near future.

<sup>1</sup>First generation CBTp describes approaches that formulate and address a range of psychotic experiences, and *problem-specific CBTp* formulates and targets specific psychotic experiences, e.g. paranoia.

<sup>2</sup>Effect sizes comparable to clozapine (Leucht *et al.*, 2013).

### Misconception 2: CBTp is about ‘thinking positively’ or correcting people’s thinking

CBTp involves working closely with a person to develop an understanding of the cognitive, behavioural, and affective processes contributing to the development and maintenance of their psychosis and linked distress. An individualised formulation forms the basis for reflection on early learning and self-perpetuating cycles of thinking, feeling, and responding that maintain distress and disability. For example, recognising how a derogatory voice triggers and reflects self-critical thoughts and beliefs learnt early in life, which affect mood and willingness to do what’s important to the individual, leading to further self-criticism; or how anxiety drives expectations of threat from others and avoidance, leading to further anxiety when next in the feared situation. Based on an individualised formulation, we can support people to make changes in line with personal goals, through a process of guided discovery (Padesky, 1993; Padesky and Kennerley, 2023); for example, re-evaluating self-critical appraisals, or gradually building confidence in feared situations even when feeling anxious.

In addition to formulating, we seek to understand how an individual knows themselves and the world – their *epistemic style* (Royce *et al.*, 1964; Royce *et al.*, 1978). Moorey (2024) describes three epistemic styles linked to change in cognitive behavioural therapies – people with a *rational/empirical* style prioritise logical consistency and objective validity (e.g. evidence for and against), people with a more *pragmatic* style benefit from considering impact and utility (e.g. ‘*does it help you to . . . ?*’), and those with a *constructivist* style benefit from generating alternative perspectives (e.g. ‘*how would you like to be in this situation?*’). Guided discovery is likely to be most effective when drawing on the person’s preferred epistemic style(s). For example, someone with a rational/empirical style who is fearful of the power of a voice might benefit from testing whether the voice can cause harm (e.g. by gradually postponing compliance). If they had a pragmatic style, we might encourage them to consider the impact of spending hours ruminating on the voice, given their goal to spend more time on their artwork. If they had a more constructivist style, we might support them to develop a coherent narrative of past traumas to help make sense of core self- and other-beliefs reflected in the voice. In practice, we draw on all these approaches; the skilful CBTp therapist recognises people’s epistemic style(s) and adapts therapy accordingly.

The mistaken notion that CBTp is about ‘thinking positively’ or correcting people’s thinking is sometimes advanced by people who have not been trained in CBTp. A more subtle version of this argument sees CBT as paternalistic (doing to), or simplistic (correcting rather than guiding discovery). It is not our role to decide what people should or should not think, or to educate people in ‘correct thought’. Indeed, we are likely to work with many people whose opinions and views we disagree with, and may sometimes find objectionable. Our focus is on the problems for which the person is seeking help, and supporting their goals for personal recovery and autonomy (Hutton and Morrison, 2013). Our role is to *create the conditions for change*<sup>3</sup> that enable a person to step back and re-evaluate self-defeating appraisals, try out different behaviours, and develop ways to manage difficult emotions. The fallacy of Orwellian ‘thought police’ is also undermined by the fact that changing what are typically over-learned and emotionally driven responses to voices, paranoia and other psychotic experiences do not simply follow corrective information or encouragement to ‘think positively’.

CBTp is not about thinking in any particular way. It is about creating the conditions for people to make changes that are important to them, if they so choose. Skilful CBTp involves an awareness of individuals’ epistemic styles (and our own) to support therapeutic change in line with the person’s formulation and personal goals for recovery and autonomy.

<sup>3</sup>Thanks to Sean Harper for clarifying our thinking regarding CBT therapists’ role in *creating conditions for change*.

### Misconception 3: CBTp doesn't recognise the therapeutic relationship (or it's all about the therapeutic relationship)

The therapeutic relationship is the basis for all CBT; '*The aspiring cognitive therapist must be, first, a good psychotherapist*' (Beck, 1979; p. 22), and the most common mistake made by those new to or unfamiliar with CBT is '[s]lighting the therapeutic relationship' (p. 32). Despite this clear emphasis from the earliest CBT texts, many criticise CBT and therefore CBTp for being mechanistic in delivery and effecting superficial change, and (paradoxically) that any benefit is due solely to relationship factors (cf. Proctor, 2003). The suggestion that CBTp does not recognise the therapeutic relationship is a violation of the core principles and practice of CBT (e.g. Brabban *et al.*, 2017). The suggestion that the impact of CBTp is entirely due to the relationship is inconsistent with the evidence (e.g. Freeman *et al.*, 2021).

In CBTp, as in CBT, we articulate and evaluate the therapeutic relationship in terms of *interpersonal effectiveness* and *collaboration* (Blackburn *et al.*, 2001). Following Rogers (1957), interpersonal effectiveness describes the ability to communicate genuine regard, empathy, and warmth. Collaboration describes the ability to engage a person in the active task of therapeutic discovery and change. If we fail to attend to the therapeutic relationship, we are not delivering CBT. Like any therapy, CBTp done poorly could cause harm (cf. Parry *et al.*, 2016; Proctor, 2008).

In line with the well-established argument for the role of common interpersonal factors across psychological therapies (Rosenzweig, 1936; Wampold, 2001; Wampold and Imel, 2015), meta-analyses consistently indicate a modest effect of relationship quality on clinical outcomes, irrespective of modality and presenting problem (e.g. Martin *et al.*, 2000). Meta-analyses also evidence this for psychosis specifically, showing that relationship quality has a small to moderate impact on therapy engagement and outcomes, including in CBTp (Bourke *et al.*, 2021; Browne *et al.*, 2021). Furthermore, in a secondary analysis of a CBTp trial, Goldsmith *et al.* (2015) showed that the therapeutic relationship is likely to have a *causal* effect on outcomes; in the context of a strong relationship, more sessions led to better outcomes ( $\beta = -2.91$ ), whereas with a poor relationship, more sessions had a detrimental effect ( $\beta = 7.74$ ).

The consistent finding that the relationship contributes to therapeutic outcomes is particularly important in CBTp given that people with psychosis typically delay accessing services for 1–2 years (Birchwood *et al.*, 2013), and about a third then disengage following initial contact (Doyle *et al.*, 2014; Kreyenbuhl *et al.*, 2009) including from CBTp (though estimates vary, e.g. Peters *et al.*, 2015; Richardson *et al.*, 2019). The quality of the relationship early in CBTp may therefore determine whether someone remains engaged and receives 'full dose' therapy.

In CBTp, we assume that the quality of the relationship is necessary but not sufficient to effect therapeutic change (cf. Beck, 2020). Meta-analyses of first generation CBTp indicate small effects compared with (increasingly improved) treatment as usual, and active therapeutic controls such as supportive counselling (Solmi *et al.*, 2023). More recent trials of problem-specific CBTp show large effects compared with befriending (e.g. Freeman *et al.*, 2021). Both supportive counselling and befriending focus primarily on fostering a strong and supportive therapeutic relationship, and the outcomes are not as good as for CBTp. Collectively, the evidence indicates that CBTp is likely to be most beneficial when based on a sound therapeutic relationship, *and* targeting the psychological mechanisms maintaining specific psychotic experiences.

### Misconception 4: CBTp focuses only on psychotic symptoms

Rarely is someone referred for CBTp with a single, discrete problem or diagnosis. Most people have a history of adversity or trauma, multiple co-existing symptoms, and a range of social needs. CBTp often faces criticism for being too narrowly focused to benefit people who have complex presentations, who may not be asking for help with discrete problems such as voices or paranoia, or who are more focused on finding a job or coping with the effects of early childhood abuse.

While CBTp is most developed for voices and paranoia, the assumption that therapy focuses only on these symptoms, disregarding other problems and goals that the individual wishes to address, is incorrect. The misunderstanding that CBTp is unsuitable for people with additional needs discourages services from offering an evidence-based, potentially transformative intervention to people who could benefit from the opportunity to address a range of psychological and social challenges.

During CBTp assessment and formulation, individuals are encouraged to reflect on what is most problematic for them, how they would like things to be different, and what they hope to achieve in therapy. These objectives typically extend beyond symptom reduction to aspects of personal recovery and autonomy, such as improving relationships, being ready to find meaningful employment, and self-acceptance. As part of the assessment, we collaborate with the person to agree personally meaningful therapy goals that align with what CBTp can offer. Addressing these goals typically involves focusing on a range of presenting problems, including psychosis, trauma symptoms, mood and sleep disturbance, social withdrawal, and relationship issues.

Some people prioritise understanding the past or managing fears about the future. Where people want to make sense of complex and confusing experiences, past and present, a developmental formulation can give context and coherence to a complex personal history and ongoing impact on current experience and relationships. For many, past episodes of psychosis and linked treatments have themselves been traumatic, prompting Gumley and Schwannauer (2006) to develop a framework to formulate and guide treatment for fear of relapse and linked iatrogenic harms.

'Negative symptoms' describe reductions in usual affect, cognition, motivation, social interaction, and behavioural expression. Whether these are due to primary neurocognitive changes or secondary to other factors (such as social deprivation, trauma, substance misuse, and medication side-effects), cognitive models assume that repeated setbacks and failures foster low expectations about performance (e.g. *'I can't do it so there's no point trying'*), relationships (*'I'm not normal so no one will want me'*), pleasure and success (e.g. *'If I get a job I won't enjoy it'*), and resources (e.g. *'my memory is terrible'*). These maintain disengagement and avoidance, functioning to protect the person from expected pain and rejection (Beck *et al.*, 2018; Beck *et al.*, 2020; Perivoliotis and Cather, 2009). CBTp for negative symptoms draws on the theory and practice of CBT for depression, targeting these appraisals, and utilising behavioural activation by working on small steps towards personal goals, while accommodating neurocognitive impairments and engagement difficulties (e.g. in planning, concentration, and memory; Beck *et al.*, 2018; Beck *et al.*, 2020; Perivoliotis and Cather, 2009). CBTp is less well developed in this area, there are far fewer trials, and reviews indicate small pooled effects, usually as secondary outcomes (Aleman *et al.*, 2017; Lutgens *et al.*, 2017; Riehle *et al.*, 2020). It will be interesting to see whether, like CBTp for voices and paranoia, future research adopts a more symptom specific approach, targeting likely maintenance mechanisms, and if this leads to improved outcomes. Broader psychosocial interventions such as exercise, music, dance, and cognitive remediation also show promise in this area (Cella *et al.*, 2023; Lutgens *et al.*, 2017).

The breadth of people's needs requires therapists to be skilled in delivering CBT for a wide range of issues including but not limited to 'positive symptoms' of psychosis. The UK CBTp training curriculum, based on the UCL competence framework (Roth and Pilling, 2013), is designed to reflect this, equipping therapists with the knowledge and skills to address common affective presentations, including depression, social anxiety and PTSD, as well as psychosis. Trainees' competencies are rigorously assessed throughout their training, so that when qualified, therapists are able to provide high quality, evidence-based treatment for a range of presenting problems.

### Misconception 5: Collaboration means never being direct

A common issue we encounter when training clinicians to deliver CBTp is novice therapists equating collaboration with passivity. The Cognitive Therapy Scale-Revised (CTS-R) describes collaboration as encouraging *active engagement* in therapy, leading to *productive teamwork* (Blackburn *et al.*, 2001). Skilful collaboration involves the therapist facilitating a shared

responsibility for the therapeutic process through the use of non-verbal prompts, as well as joint decision making and problem solving. While a questioning approach is important, therapist passivity is highlighted as an obstacle to effective collaboration.

As an example, CBTp is a goal-oriented therapy and so at assessment we discuss what the person would like to get out of the sessions in order to agree personalised therapy goals. This does not mean that the therapist should *only* ask the individual about their goals, without *also* explaining what CBTp can offer and making suggestions, given their training and experience, as to what might be helpful. People with psychosis often say they would like to get rid of their voices or the images they see, or request help reporting neighbours or others interfering in their lives to the police. Others may struggle to articulate goals or have such low expectations that they cannot imagine how therapy could be beneficial. We spend time supporting people to think through possible goals and provide information about how we may be able to help make sense of unshared experiences, cope with them more effectively, and address key maintenance processes (e.g. worry and sleep in the case of persecutory delusions). We would also want to provide information about other evidence-based treatments where these align better with the person's needs and goals, for example family intervention for psychosis to optimise caregiving when day-to-day arguments make things even more difficult for the person, and exacerbate their distress further (see Jolley and Grice, 2024).

For any of us, when we go to see a healthcare professional, unless we have specific training in that field, we don't know what we don't know. We expect the clinician to share with us information about the nature of the problem, evidence-based treatment options, and likely outcomes, in order to make informed treatment decisions. Similarly, collaboration in CBTp includes being explicit about what we can and cannot offer and what, from our perspective, may be helpful, in order that the person can make an informed decision about whether to pursue therapy, and if so, the goals they would like to prioritise.

This may all seem common sense. However, many trainees worry that providing information, guiding a conversation, making suggestions, and sometimes interrupting people, will be invalidating and disrupt the therapeutic relationship. Behavioural experiments in (or agreed in) supervision to experience being given information, guided in a discussion, and respectfully interrupted (and observing patients' responses to similar) can be very effective in addressing these unhelpful appraisals. When delivering training in the Feeling Safe approach (Freeman *et al.*, 2021) the trainers (including L.I.) are sometimes asked whether offering therapy with the explicit goal of '*feeling safer, happier, and getting back to doing more of the things you want to do*' is too directive. Clearly, insisting that the person *must* engage in a specific approach would be highly uncollaborative, but providing information that allows people to decide if this is something they would like to pursue is consistent with therapeutic collaboration.

Finally, it is important to note that CBTp often requires a degree of assertive outreach when making an offer of therapy. Some people may not initially recognise that they have a difficulty as described in typical psychiatric language (e.g. schizoaffective disorder), may not know what psychological therapy involves, or may not see that therapy has anything to offer, and so not seek referral. As clinicians, we therefore sometimes need to be proactive in making an offer of therapy and ensure we remain up-to-date with the evidence base so we can give clear and accurate information to support genuine collaboration and informed decision making.

### Misconception 6: CBTp isn't trauma informed

Early adversity, particularly childhood interpersonal trauma, is associated with the development, severity and persistence of psychosis (Bailey *et al.*, 2018; De Vries and Goggin, 2018; Varese *et al.*, 2012), and shapes voices and other hallucinatory experiences phenomenologically (Peach *et al.*, 2021; van den Berg *et al.*, 2022). People with psychosis report higher levels of trauma and PTSD than the general population, including psychosis-related PTSD (Fornells-Ambrojo *et al.*, 2016). In a large Dutch sample, de Bont *et al.* (2015) found that over 78% of people with psychosis reported

at least one traumatic event in their lives, and 16% were likely to meet criteria for PTSD (in contrast to 0.5% reported in clinical notes). The literature indicates that psychosis is likely to be both a response to and cause of trauma (Gumley and MacBeth, 2007; Hardy, 2017; Larkin and Read, 2008; Morrison *et al.*, 2003), which is often not recognised, and more often not recorded (de Bont *et al.*, 2015). CBTp therapists must therefore be willing and able to assess and treat trauma and the consequences of trauma, drawing on CBTp as a *trauma-informed therapy* and directly with *trauma-focused CBTp* (Hardy *et al.*, 2023).

Over the last decade, *trauma-informed care* has gained widespread recognition as a foundational component of recovery focused mental health services. Trauma-informed care incorporates the *realisation* of how trauma can affect individuals and communities, a *recognition* of the signs of trauma, organisational policy and procedural *responses*, and avoidance of iatrogenic *re-traumatisation* (e.g., Substance Abuse and Mental Health Services Administration, 2014). To be consistent with these principles, CBTp assessment must involve asking sensitively about early and subsequent adversity and trauma (see Read (2007) for guidance on asking people with psychosis about past trauma and responding effectively). Formulation then involves linking current psychotic experiences and associated appraisals, emotions, and behavioural responses, with beliefs and assumptions that have typically been learnt early in life, to help make sense of the development and maintenance of distressing psychosis. For many, this is a validating process, recognising, often for the first time, that psychosis and responses to psychosis can be understood in the context of earlier life experiences.

Where people report trauma symptoms, show signs of trauma-related mechanisms (e.g. dissociation), or meet criteria for PTSD, they are likely to need a trauma-focused therapy, such as trauma-focused CBTp, Imagery Rescripting, EMDR and dialogic approaches, to process trauma-related memories, beliefs and emotions directly (Hardy *et al.*, 2023). Here, again, clinical practice falls well behind the evidence base and best practice guidelines which recommend trauma interventions for people with psychosis and PTSD (NICE, 2014). This is likely to be due in large part to clinicians' caution about re-traumatising people, and increasing risks to self or others, despite evidence that PTSD can safely and effectively be treated in people with psychosis (Sin and Spain, 2016). The current STAR trial incorporates CBT for PTSD and CBTp for people with psychosis (Peters *et al.*, 2022), and is likely to shape the field.

### Misconception 7: CBTp is simple and doesn't require specialist or extensive training

A common mistake is to assume that interventions based on a simple premise are simple to put into practice. While CBTp is based on the arguably simple premise that people can effect emotional and behavioural change through cognitive change, in practice this requires considerable expertise.

To deliver competent CBTp, a therapist must have a critical knowledge of the relevant theoretical and research literature, and the clinical competence to translate this knowledge into practice, for a wide range of presentations (see Misconception 4). The UK CBTp training curriculum is based on the UCL competence framework (Roth and Pilling, 2013), which describes the knowledge, skills, and meta-competences required. These include fundamental principles of CBT theory; problem specific models and protocols for depression, anxiety presentations, trauma, bipolar and psychosis; and broader CBTp relevant constructs (e.g. how memory works). Clinical application requires the ability to develop and maintain good therapeutic relationships (including with people who find it hard to trust others and may incorporate the therapist in their psychosis); to conduct CBT assessments and develop collaborative, accessible formulations that draw on multiple models where needed; to deliver culturally competent, individualised interventions with respect for the person's values, culture and social differences; to work within a person's belief system as required; and to utilise standardised and idiosyncratic measures to monitor and support therapeutic progress.

The depth of knowledge, skill and experience required to deliver good quality CBTp is considerable. Almost all clinicians attending our one- and two-year postgraduate training courses in CBTp, including qualified practitioner psychologists, do not meet competency standards on therapy recordings at the start of training. This is to be expected (though often disappointing to trainees) and the purpose of CBTp training. CBTp skills can be acquired (the majority meet and exceed these standards by the end of training), but doing so requires specialist teaching, close supervision, and practise over an extended period. Competency in CBT and CBTp is not acquired by attending a few workshops, and the failure to recognise this risks patients being given poor quality interventions which are not, in fact, CBTp, but labelled as such. This in turn can result in poor outcomes, iatrogenic harms, people being less willing to engage in future therapy, and CBTp developing a poor reputation with referrers and the general public. Current UK guidance on Early Intervention in Psychosis services requires that services only record delivery of CBTp if the clinician has completed one- or two-year specialist CBTp postgraduate training, evidencing competence, or equivalent.

Poor public understanding of CBTp, and reports that some people have found (mislabelled) ‘CBTp’ to be simplistic, invalidating, or ineffective, behove our professional bodies to maintain and publicise rigorous standards of care, particularly in terms of training and ‘full dose’ therapy. In medicine, we expect clinicians to be appropriately trained and qualified, and their practice held to account by relevant governance systems. Likewise, prescription of medicines without the active ingredients, or in insufficient quantities, would be considered unethical and malpractice. While not everyone with psychosis will choose or benefit from CBTp, as is the case for most treatments, it is essential that we are clear about what CBTp is and is not, and distinguish evidence-based CBTp from widely used ‘CBT-based’ practice if we are to ensure people can make informed decisions about treating and managing their mental health.

### Misconception 8: This person is too complex (too ill, too recently discharged . . .) for CBTp

Most of the people we see in UK secondary care services present with complex needs, including additional diagnoses, substance use, and/or risk (usually to themselves). The NICE guideline (2014) recommends CBTp for everyone with psychosis including in the acute phase.

Early suitability criteria for CBT were based on predictors of outcome following short-term therapy. These criteria emphasised awareness of thoughts and feelings, ability to engage in a therapeutic relationship, shorter-term problems, and personal responsibility for change (Safran *et al.*, 1993). People with psychosis may initially struggle to articulate their internal experiences, be cautious about trusting a therapist, present with complex and long-standing difficulties, and be ambivalent about change. This is understandably daunting for novice therapists.

It is essential that senior clinicians uphold best practice guidelines, recognise when junior colleagues are hesitant to offer therapy in these contexts, and support access to training and supervision as needed. Equally, service leads should not use criteria developed before the evidence for CBTp evolved to manage scarce resources. Where we have insufficient provision, for CBTp and other evidence-based therapies, we need to be transparent about this and not suggest that any lack of offer is due to the person being too unwell, too recently discharged, or otherwise too complex. For example, a recent review found that CBT can be delivered in acute care and is likely to be beneficial (Wood *et al.*, 2020).

Barton and colleagues (2017, 2023) describe *problem complexity*, when people have multiple problems that interact with each other, and *relational complexity*, when complications in the therapeutic relationship or service context interfere with effective collaboration. In CBTp, we are likely to see both problem and relational complexity. People with psychosis often present with a mix of psychosis (e.g. voices, paranoia, social withdrawal), affective (e.g. depression, anxiety) and other difficulties (e.g. substance misuse, risk). Additionally, psychosis can have a direct impact on the therapeutic relationship, e.g. voices often worsen prior to or during therapy, may criticise the

person for coming to sessions, and undermine the therapist. Social withdrawal and confusion can make it hard for people to attend and engage actively in therapy. Paranoia and grandiosity can make it difficult to trust and talk to a therapist. In a qualitative study of grandiose delusions, participants emphasised how their beliefs made sense of anomalous or adverse experiences, but that these were often hard to put into words (Isham *et al.*, 2021). Problem and relational complexity need to be recognised, formulated, and addressed in therapy (and supervision) for CBT to be effective (Barton *et al.*, 2017; Barton *et al.*, 2023).

As an example, a young man with a history of disrupted attachments, substance misuse and various previous diagnoses presented with voices, paranoia, social withdrawal, anxiety, anger and some ongoing drug use (problem complexity). He did well to simply get to therapy sessions (and occasionally didn't). When he attended, he appeared distracted by his voices and wary of the therapist (K.N.-T.), consistent with a pervasive paranoia about people in authority (relational complexity). Supervision and some flexibility in service expectations were helpful in clarifying (1) the focus for therapy in the context of problem complexity – first to make sense of his experiences (formulation), and then to address fears about seeing family, friends and working (his goals), and (2) the impact of relational complexity – the ways in which his psychosis, anxiety, and substance use were likely to effect the therapeutic relationship, sensitively anticipating this with him, and problem solving together when these issues arose.

### Misconception 9: CBTp can be done entirely in the clinic room

A core aspect of CBT involves taking therapy out of the clinic room and into the real world (Kennerley *et al.*, 2016). When working with someone with depression, we might go for a brisk walk together to find out what impact this has on the person's mood. In social anxiety, we might visit a café, order a coffee, and observe whether, as the person predicts, the waitress notices and responds negatively to their anxiety. In treating OCD, we might go with someone to use a public restroom whilst refraining from compulsive hand washing afterwards, to discover whether they become physically unwell or experience an intolerable level of anxiety, as feared.

Such *in vivo* work offers many advantages over and above therapy conducted solely in the confines of the therapy room by:

- Allowing the therapist to provide immediate encouragement and support when the person tries a feared or avoided activity for the first time.
- Identifying key information that might not be recognised if the therapist was not present (e.g. subtle safety seeking behaviours).
- Providing an opportunity for modelling, practising, and refining new strategies, and problem-solving difficulties as they arise.
- Allowing the therapist to support the person to articulate new appraisals in the moment of discovery (e.g. *'I can go out safely even when feeling afraid'*).
- Building momentum in therapy – providing more opportunities for key learning that can be built on with between-session tasks.
- Strengthening the therapeutic alliance, by being alongside the person when commencing the difficult process of behaviour change.

These benefits are just as relevant when working with people with psychosis, and recommended in CBTp manuals and approaches (e.g. Morrison, 2017). Despite this, clinicians are often reluctant to get out of the clinic room. In a recent training event, more than 100 CBTp practitioners were asked what proportion of their patients they had routinely been out with as part of their therapy; 92% reported having done this with fewer than half their patients, and 33% said they never did this.

Therapists cite a number of reasons for their reluctance to leave the clinic. Some have not tried *in vivo* experiments before and are anxious about them ‘going wrong’. Others are nervous about managing risk away from the team base. These anxieties are understandable but likely to impede therapy. Accessing supervision to support the planning and execution of within session activities, and robust (but not excessively cautious) risk assessment and planning, should ensure these not be default reasons to avoid *in vivo* work. Most commonly, however, therapists say they simply do not have time to leave the clinic with patients, reporting systemic pressures, such as needing to see people ‘back-to-back’ and in sessions of no more than 50 minutes. This is particularly concerning; systematically excluding a core tenet of CBT risks diluting the intervention and reducing likely treatment effectiveness. We suggest it is unethical to tell people they are receiving a NICE recommended treatment and then giving a substandard ‘dose’. Once more, this would not be tolerated in physical health settings (e.g. providing half the recommended dosage of diabetes medication), and should not be tolerated in mental health care.

### Misconception 10: Measures are too burdensome (and don’t get to what’s really important)

Routine outcome measurement is an integral part of CBT, and can benefit the person with psychosis, the therapist, and the service (Faija *et al.*, 2022). The wider psychotherapy outcome literature shows that taking measures improves engagement and outcomes (Lambert *et al.*, 2003; De Jong *et al.*, 2014). Indeed, most people with psychosis find measures useful as a way of ‘*feeling understood, valuing opportunities to reflect, expressing feelings, and tracking progress towards goals*’ (Fornells-Ambrojo *et al.*, 2017; p. 254).

At the start of their training, many of our CBTp trainees are using few or no regular measures. Some are uncertain what to measure or which scales to use, but most are concerned that people will find measures aversive or reductive, and that this will hamper engagement. It is of note that this assumed threat to engagement is also given as a reason for excluding other core aspects of CBT, such as setting an agenda and agreeing homework tasks. Whilst we wholeheartedly support continuous attention to, and prioritisation of, the therapeutic relationship (see Misconception 3), this does not necessitate omitting core components of CBT when the evidence shows that done well, routine outcome measurement enhances both engagement and outcomes.

*How to introduce measures?* Outcome measurement is introduced as part of CBTp assessment, with a clear rationale in line with the person’s goals. Potential benefits include further understanding of the problem(s) for which they are seeking help (and linked mechanisms), identifying treatment targets, and capturing progress (Kennerley *et al.*, 2016). This is usually highly valued as a way to keep therapy on track in line with personal goals for recovery (cf. Waller and Turner, 2016). Without a clear and shared rationale, measures may be seen as irrelevant, and this can certainly hinder engagement (Rao *et al.*, 2010).

*Which measures to use?* CBTp utilises a limited combination of standardised measures, brief idiosyncratic sessional measurements (e.g. subjective units of distress), and within session tools (e.g. belief ratings in behavioural experiments and activity diaries). The choice of measures is based on the person’s presentation, needs and goals, considering:

- Is the measure useful? What will it tell us and does the person agree it will be helpful? Is this the best option?
- For a standardised measure – is it valid and reliable (psychometric properties)?
- Is the measure accessible? This takes account of the person’s literacy and belief system. Occasionally, we may change a word if likely to be confusing or unacceptable (e.g. replacing ‘delusions’ with ‘concerns others are trying to harm me’). While better not to change standardised questionnaires, minor adaptations are sometimes necessary, and preferable to omitting measures altogether.

*How often and in what format?* Brief measures can be used sessionally, whereas longer (typically standardised) measures are used less frequently at regular intervals. The best combination of measures varies from person to person, taking account of their problems, priorities and capacity. Some people are happy to complete measures between sessions, others prefer taking a few minutes to run through these with the therapist each week, and some need to spread just a few measures across sessions to reduce burden.

In this way, we come to a shared agreement about which measures to use, why, how often, and in what format, ensuring the combination is relevant, useful, and not burdensome. Too often, clinicians fail to take any measures, having assumed without discussion that doing so would be arduous and without benefit, when a limited number of carefully selected measures is both possible and likely to be helpful.

Routine outcome measurement is built into UK NHS Talking Therapies<sup>4</sup> services to monitor impact at individual, service, and national levels. This is likely to play a significant role in ongoing funding decisions. While national guidelines recommend similar for people with severe mental ill-health (Royal College of Psychiatrists, 2024), adherence is very patchy. A minimum co-produced dataset could support and shape services, and provide the large scale data needed to advance personalised psychological therapies for people with psychosis (Newman-Taylor and Bentall, 2024).

## Conclusion

As CBTp therapists, service leads, and professional bodies, we have a collective responsibility to ensure that people with psychosis and their families understand what best practice CBTp involves, and that this is one of a number of evidence-based treatment options that should be available to them. We also have a collective duty to address the misinformation and stigma that can deter people from seeking help, and tackle the structural service barriers that delay timely access to culturally sensitive treatments (Birchwood *et al.*, 2013; Schlieff *et al.*, 2023). As psychological therapists, our role is to deliver CBTp of the standard we would genuinely be happy to see offered to our own family and friends. This paper discusses ten common misconceptions about CBTp, to support clinicians, service leads, and our professional bodies to implement best practice mental health care for people with psychosis.

### Key practice points

- (1) Common and pervasive misconceptions about psychosis constitute a major barrier to providing best practice mental health care.
- (2) If we address any misconceptions that we hold ourselves, we will be in a stronger position to offer high quality CBT to people with psychosis.
- (3) If we find the courage to challenge these misconceptions with colleagues, service leads and the public, we may be able to influence the wider health care system that so often continues to deliver sub-standard treatments to people with serious mental health conditions.

## Further reading

- Brabban, A., Byrne, R., Longden, E., & Morrison, A. P.** (2017). The importance of human relationships, ethics and recovery-orientated values in the delivery of CBT for people with psychosis. *Psychosis*, 9, 157–166.
- Freeman, D.** (2024). Developing psychological treatments for psychosis. *British Journal of Psychiatry*, 224, 147–149.
- Newman-Taylor, K., & Bentall, R.** (2024). Cognitive behavioural therapy for psychosis: the end of the line or time for a new approach? *Psychology and Psychotherapy: Theory, Research and Practice*, 97, 4–18.

<sup>4</sup>Previously ‘Improving Access to Psychological Therapies’ (IAPT) – the first freely accessible national psychological therapies service.

Solmi, M., Croatto, G., Piva, G., Rosson, S., Fusar-Poli, P., Rubio, J.M., Carvalho, A.F., Vieta, E., Arango, C., DeTore, N.R., Eberlin, E.S., Mueser, K.T., & Correll, C.U. (2023). Efficacy and acceptability of psychosocial interventions in schizophrenia: systematic overview and quality appraisal of the meta-analytic evidence. *Molecular Psychiatry*, 28, 354–368.

**Data availability statement.** Data availability is not applicable to this article as no new data were created or analysed in this study.

**Acknowledgements.** We thank the many people with psychosis who have helped us understand their often highly distressing and isolating experiences, and how CBT can be most useful to them. This paper also reflects the work of our clinical colleagues who skilfully support people with psychosis towards personal recovery and autonomy. Our curious and committed CBT and DClin trainees have helped shape our thinking about how people learn the art of CBT, placing the person with psychosis at the heart of this work. Finally, thanks go to Hannah Murray and colleagues for *Ten Misconceptions About Trauma-Focused CBT for PTSD* which inspired this paper.

**Author contributions.** **Katherine Newman-Taylor:** Conceptualization (equal), Writing - original draft (equal), Writing - review & editing (equal); **Louise Isham:** Conceptualization (equal), Writing - original draft (equal), Writing - review & editing (equal); **Alison Brabban:** Conceptualization (equal), Writing - original draft (equal), Writing - review & editing (equal).

**Financial support.** This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

**Competing interests.** Katherine Newman-Taylor is associate editor for *the Cognitive Behaviour Therapist* and *Behavioural and Cognitive Psychotherapy* journals. She was not involved in the review process for this paper. Louise Isham is a trainer on the Feeling Safe Training Programme 5-day course, and a director of the Ox Psychology Group limited company. Alison Brabban is National Clinical Advisor, Adult Mental Health, for NHS England. We have no other competing interests to declare.

**Ethical standards.** The authors have abided by the Ethical Principles of Psychologists and Code of Conduct as set out by the BABCP and BPS.

## References

- Aleman, A., Lincoln, T. M., Bruggeman, R., Melle, I., Arends, J., Arango, C., & Knegteling, H. (2017). Treatment of negative symptoms: where do we stand, and where do we go? *Schizophrenia Research*, 186, 55–62.
- Bailey, T., Alvarez-Jimenez, M., Garcia-Sanchez, A. M., Hulbert, C., Barlow, E., & Bendall, S. (2018). Childhood trauma is associated with severity of hallucinations and delusions in psychotic disorders: a systematic review and meta-analysis. *Schizophrenia Bulletin*, 44, 1111–1122. <https://doi.org/10.1093/schbul/sbx161>
- Barton, S., Armstrong, P., Wicks, L., Freeman, E., & Meyer, T. D. (2017). Treating complex depression with cognitive behavioural therapy. *the Cognitive Behaviour Therapist*, 10, e17.
- Barton, S. B., Armstrong, P. V., Robinson, L. J., & Bromley, E. H. (2023). CBT for difficult-to-treat depression: self-regulation model. *Behavioural and Cognitive Psychotherapy*, 51, 543–558.
- Beck, A. T. (1952). Successful outpatient psychotherapy of a chronic schizophrenic with a delusion based on borrowed guilt. *Psychiatry*, 15, 305–312. <https://doi.org/10.1080/00332747.1952.11022883>
- Beck, A. T. (1979). *Cognitive Therapy and the Emotional Disorders*. Penguin.
- Beck, A. T., & Grant, P. M., Inverso, E., Brinen, A. P., & Perivoliotis, D. (2020). *Recovery-Oriented Cognitive Therapy for Serious Mental Health Conditions*. Guilford Press.
- Beck, A. T., Himelstein, R., Bredemeier, K., Silverstein, S. M., & Grant, P. (2018). What accounts for poor functioning in people with schizophrenia: a re-evaluation of the contributions of neurocognitive v. attitudinal and motivational factors. *Psychological Medicine*, 48, 2776–2785.
- Beck, J. S. (2020). *Cognitive Behavior Therapy: Basics and Beyond*. Guilford Publications.
- Birchwood, M., Connor, C., Lester, H., Patterson, P., Freemantle, N., Marshall, M., Fowler, D., Lewis, S., Jones, P., Amos, T., Everard, L., & Singh, S. P. (2013). Reducing duration of untreated psychosis: Care Pathways to Early Intervention in psychosis services. *British Journal of Psychiatry*, 203, 58–64. <https://doi.org/10.1192/bjp.bp.112.125500>
- Birchwood, M., Michail, M., Meaden, A., Tarrier, N., Lewis, S., Wykes, T., *et al.* (2014a). Cognitive behaviour therapy to prevent harmful compliance with command hallucinations (COMMAND): a randomised controlled trial. *Lancet Psychiatry*, 1, 23–33.
- Birchwood, M., Shiers, D., & Smith, J. (2014b). CBT for psychosis: not a ‘quasi-neuroleptic’. *British Journal of Psychiatry*, 204, 488–489.
- Blackburn, I.-M., James, I. A., Milne, D. L., Baker, C., Standart, S., Garland, A., & Reichelt, F. K. (2001). The revised Cognitive therapy scale (CTS-R): psychometric properties. *Behavioural and Cognitive Psychotherapy*, 29, 431–446. <https://doi.org/10.1017/s1352465801004040>

- Bourke, E., Barker, C., & Fornells-Ambrojo, M.** (2021). Systematic review and meta-analysis of therapeutic alliance, engagement, and outcome in psychological therapies for psychosis. *Psychology and Psychotherapy: Theory, Research and Practice*, *94*, 822–853.
- Brabban, A., Byrne, R., Longden, E., & Morrison, A. P.** (2017). The importance of human relationships, ethics and recovery-orientated values in the delivery of CBT for people with psychosis. *Psychosis*, *9*, 157–166.
- Browne, J., Wright, A. C., Berry, K., Mueser, K. T., Cather, C., Penn, D. L., & Kurtz, M. M.** (2021). The alliance-outcome relationship in individual psychosocial treatment for schizophrenia and early psychosis: a meta-analysis. *Schizophrenia Research*, *231*, 154–163.
- Cella, M., Roberts, S., Pillny, M., Riehle, M., O'Donoghue, B., Lyne, J., . . . & Preti, A.** (2023). Psychosocial and behavioural interventions for the negative symptoms of schizophrenia: a systematic review of efficacy meta-analyses. *British Journal of Psychiatry*, *223*, 321–331.
- Chadwick, P., & Birchwood, M.** (1994). The omnipotence of voices: a cognitive approach to auditory hallucinations. *British Journal of Psychiatry*, *164*, 190–201.
- de Bont, P. A., van den Berg, D. P., van der Vleugel, B. M., de Roos, C., de Jongh, A., van der Gaag, M., & van Minnen, A.** (2015). Predictive validity of the Trauma Screening Questionnaire in detecting post-traumatic stress disorder in patients with psychotic disorders. *British Journal of Psychiatry*, *206*, 408–416.
- De Jong, K., Timman, R., Hakkaart-Van Roijen, L., Vermeulen, P., Kooiman, K., Passchier, J., & Busschbach, J. V.** (2014). The effect of outcome monitoring feedback to clinicians and patients in short and long-term psychotherapy: a randomized controlled trial. *Psychotherapy Research*, *24*, 629–639. <https://doi.org/10.1080/10503307.2013.871079>
- De Vries, I., & Goggin, K. E.** (2018). The impact of childhood abuse on the commercial sexual exploitation of youth: a systematic review and meta-analysis. *Trauma, Violence, & Abuse*, *21*, 886–903. <https://doi.org/10.1177/1524838018801332>
- Doyle, R., Turner, N., Fanning, F., Brennan, D., Renwick, L., Lawlor, E., & Clarke, M.** (2014). First-episode psychosis and disengagement from treatment: a systematic review. *Psychiatric Services*, *65*, 603–611. <https://doi.org/10.1176/appi.ps.201200570>
- Fajja, C. L., Bee, P., Lovell, K., Lidbetter, N., Gellatly, J., Ardern, K., Rushton, K., Brooks, H., McMillan, D., Armitage, C. J., Woodhouse, R., & Barkham, M.** (2022). Using routine outcome measures as clinical process tools: maximising the therapeutic yield in the IAPT programme when working remotely. *Psychology and Psychotherapy: Theory, Research and Practice*, *95*, 820–837. <https://doi.org/10.1111/papt.12400>
- Fornells-Ambrojo, M., Gracie, A., Brewin, C. R., & Hardy, A.** (2016). Narrowing the focus on the assessment of psychosis-related PTSD: a methodologically orientated systematic review. *European Journal of Psychotraumatology*, *7*. <https://doi.org/10.3402/ejpt.v7.32095>
- Fornells-Ambrojo, M., Johns, L., Onwumere, J., Garety, P., Milosh, C., Iredale, C., . . . & Jolley, S.** (2017). Experiences of outcome monitoring in service users with psychosis: findings from an Improving Access to Psychological Therapies for people with Severe Mental Illness (IAPT-SMI) demonstration site. *British Journal of Clinical Psychology*, *56*, 253–272.
- Freeman, D., Emsley, R., Diamond, R., Collett, N., Bold, E., Chadwick, E., *et al.*** (2021) Comparison of a theoretically driven cognitive therapy (the feeling safe programme) with befriending for the treatment of persistent persecutory delusions: a parallel, single-blind, randomised controlled trial. *Lancet Psychiatry*, *8*, 696–707.
- Galletly, C., Castle, D., Dark, F., Humberstone, V., Jablensky, A., Killackey, E., . . . & Tran, N.** (2016). Royal Australian and New Zealand College of Psychiatrists clinical practice guidelines for the management of schizophrenia and related disorders. *Australian & New Zealand Journal of Psychiatry*, *50*, 410–472.
- Garety, P. A., Kuipers, L., Fowler, D., Chamberlain, F., & Dunn, G.** (1994). Cognitive behavioural therapy for drug-resistant psychosis. *British Journal of Medical Psychology*, *67*, 259–271.
- Goldsmith, L. P., Lewis, S. W., Dunn, G., & Bentall, R. P.** (2015). Psychological treatments for early psychosis can be beneficial or harmful, depending on the therapeutic alliance: an instrumental variable analysis. *Psychological Medicine*, *45*, 2365–2373.
- Gumley, A., & Schwannauer, M.** (2006). *Staying Well After Psychosis: A Cognitive Interpersonal Approach to Recovery and Relapse Prevention*. John Wiley & Sons.
- Gumley, A. I., & MacBeth, A.** (2007). A trauma-based model of relapse in psychosis. In *Trauma and Psychosis* (pp. 295–316). Routledge.
- Hardy, A.** (2017). Pathways from trauma to psychotic experiences: a theoretically informed model of posttraumatic stress in psychosis. *Frontiers in Psychology*, *8*. <https://doi.org/10.3389/fpsyg.2017.00697>
- Hardy, A., Keen, N., van den Berg, D., Varese, F., Longden, E., Ward, T., & Brand, R. M.** (2023). Trauma therapies for psychosis: a state-of-the-art review. *Psychology and Psychotherapy: Theory, Research and Practice*, *97*, 74–90. <https://doi.org/10.1111/papt.12499>
- Huhn, M., Nikolakopoulou, A., Schneider-Thoma, J., Krause, M., Samara, M., Peter, N., Arndt, T., Bäckers, L., Rothe, P., Cipriani, A., Davis, J., Salanti, G. & Leucht, S.** (2019). Comparative efficacy and tolerability of 32 oral antipsychotics for the acute treatment of adults with multi-episode schizophrenia: a systematic review and network meta-analysis. *Lancet*, *394*, 10202, 939–951.
- Hutton, P., & Morrison, A. P.** (2013). Collaborative empiricism in cognitive therapy for psychosis: a practice guide. *Cognitive and Behavioral Practice*, *20*, 429–444.

- Isham, L., Griffith, L., Boylan, A., Hicks, A., Wilson, N., Byrne, R., Sheaves, B., Bentall, R. P., & Freeman, D. (2021). Understanding, treating, and renaming grandiose delusions: a qualitative study. *Psychology and Psychotherapy: Theory, Research and Practice*, 94, 119–140. <https://doi.org/10.1111/papt.12260>
- Jaspers, K. (1959). *Truth and Symbol*. Bloomsbury Publishing PLC.
- Jauhar, S., McKenna, P. J., Radua, J., Fung, E., Salvador, R., & Laws, K. R. (2014). Cognitive-behavioural therapy for the symptoms of schizophrenia: systematic review and meta-analysis with examination of potential bias. *British Journal of Psychiatry*, 204, 20–29.
- Johns, L., Jolley, S., Keen, N., & Peters, E. (2014). CBT with people with psychosis. In A. Whittington & N. Grey (eds), *How to Become a More Effective CBT Therapist: Mastering Metacompetence in Clinical Practice* (pp. 191–207). John Wiley & Sons.
- Jolley, S., & Grice, S. (2024). State of the art in psychological therapies for psychosis: family interventions for psychosis. *Psychology and Psychotherapy: Theory, Research and Practice*, 97, 19–33.
- Kennerley, H., Kirk, J., & Westbrook, D. (2016). *An Introduction to Cognitive Behaviour Therapy: Skills and Applications* (3rd edn). Sage.
- Kingdon, D. G., & Turkington, D. (1991). The use of cognitive behavior therapy with a normalizing rationale in schizophrenia: preliminary report. *Journal of Nervous and Mental Disease*, 179, 207–211.
- Kreyenbuhl, J., Nossel, I. R., & Dixon, L. B. (2009). Disengagement from mental health treatment among individuals with schizophrenia and strategies for facilitating connections to care: a review of the literature. *Schizophrenia Bulletin*, 35, 696–703. <https://doi.org/10.1093/schbul/sbp046>
- Lambert, M. J., Whipple, J. L., Hawkins, E. J., Vermeersch, D. A., Nielsen, S. L., & Smart, D. W. (2003). Is it time for clinicians to routinely track patient outcome? A meta-analysis. *Clinical Psychology: Science and Practice*, 10, 288.
- Larkin, W., & Read, J. (2008). Childhood trauma and psychosis: evidence, pathways, and implications. *Journal of Postgraduate Medicine*, 54, 287–293.
- Lorenzo-Luaces, L., DeRubeis, R. J., van Straten, A., & Tiemens, B. (2017). A prognostic index (PI) as a moderator of outcomes in the treatment of depression: a proof of concept combining multiple variables to inform risk-stratified stepped care models. *Journal of Affective Disorders*, 213, 78–85.
- Lutgens, D., Garipey, G., & Malla, A. (2017). Psychological and psychosocial interventions for negative symptoms in psychosis: systematic review and meta-analysis. *British Journal of Psychiatry*, 210, 324–332.
- Martin, D. J., Garske, J. P., & Davis, M. K. (2000). Relation of the therapeutic alliance with outcome and other variables: a meta-analytic review. *Journal of Consulting and Clinical Psychology*, 68, 438–450. <https://doi.org/10.1037//0022-006x.68.3.438>
- Moorey, S. (2024). Characteristics of Cognitive Therapy. In *Seminars in General Adult Psychiatry* (p. 147). Cambridge University Press.
- Morrison, A. P. (2017). A manualised treatment protocol to guide delivery of evidence-based cognitive therapy for people with distressing psychosis: learning from clinical trials. *Psychosis*, 9, 271–281.
- Morrison, A. P., Frame L., & Larkin, W. (2003) Relationships between trauma and psychosis: a review and integration. *British Journal of Clinical Psychology*, 42, 331–353.
- National Institute for Health and Care Excellence (2002). *Schizophrenia*. NICE. <https://www.nice.org.uk/guidance/cg1>
- National Institute for Health and Care Excellence (2005). *Post-traumatic stress disorder*. NICE. <https://www.nice.org.uk/guidance/ng116/evidence/full-guideline-march-2005-6602623597?tab=evidence>
- National Institute for Health and Care Excellence (2009). *Psychosis and schizophrenia: management*. NICE. <https://www.nice.org.uk/guidance/cg82>
- National Institute for Health and Care Excellence (2013). *Social anxiety disorder: recognition, assessment and treatment*. NICE. <https://www.nice.org.uk/guidance/cg159>
- National Institute for Health and Care Excellence (2014). *Psychosis and schizophrenia in adults: prevention and management*. NICE. <https://www.nice.org.uk/guidance/cg178>
- Nelson, H. E. (1997). *Cognitive Behavioural Therapy with Schizophrenia: A Practice Manual*. Nelson Thornes.
- Newman-Taylor, K., & Bentall, R. (2024). Cognitive behavioural therapy for psychosis: the end of the line or time for a new approach? *Psychology and Psychotherapy: Theory, Research and Practice*, 97, 4–18.
- Padesky, C. A. (1993). Socratic questioning: changing minds or guiding discovery. In *A keynote address delivered at the European Congress of Behavioural and Cognitive Therapies*, London (vol. 24, p. 44).
- Padesky, C. A., & Kennerley, H. (eds) (2023). *Dialogues for Discovery: Improving Psychotherapy's Effectiveness*. Oxford University Press.
- Parry, G., Crawford, M., & Duggan, C. (2016). Iatrogenic harm from psychological therapies – time to move on. *British Journal of Psychiatry*, 208, 210–212.
- Paul, G. L. (1967). Strategy of outcome research in psychotherapy. *Journal of Consulting Psychology*, 31, 109–118. <https://doi.org/10.1037/h0024436>

- Peach, N., Alvarez-Jimenes, M., Cropper, S. J., Sun, P., Halpin, E., O'Connell, J. & Bendall, S. (2021). Trauma and the content of hallucinations and post-traumatic intrusions in first episode psychosis. *Psychology and Psychotherapy*, 9, 223–241.
- Perivoliotis, D., & Cather, C. (2009). Cognitive behavioral therapy of negative symptoms. *Journal of Clinical Psychology*, 65, 815–830.
- Peters, E. (2014). An oversimplification of psychosis, its treatment, and its outcomes? *British Journal of Psychiatry*, 205, 159–160.
- Peters, E., Hardy, A., Dudley, R., Varese, F., Greenwood, K., Steel, C., Emsley, R., Keen, N., Bowe, S., Swan, S., Underwood, R., Longden, E., Byford, S., Potts, L., Heslin, M., Grey, N., Turkington, D., Fowler, D., Kuipers, E., & Morrison, A. (2022). Multisite randomised controlled trial of trauma-focused cognitive behaviour therapy for psychosis to reduce post-traumatic stress symptoms in people with co-morbid post-traumatic stress disorder and psychosis, compared to treatment as usual: study protocol for the star (study of trauma and recovery) trial. *Trials*, 23. <https://doi.org/10.1186/s13063-022-06215-x>
- Peters, M. D. J., Godfrey, C. M., Khalil, H., McInerney, P., Parker, D., & Soares, C. B. (2015). Guidance for conducting systematic scoping reviews. *International Journal of Evidence-Based Healthcare*, 13, 141–146. <https://doi.org/10.1097/xeb.000000000000050>
- Proctor, G. (2003). CBT: Collaboration or compliance? *Clinical Psychology*, 25, 14–16.
- Proctor, G. (2008). CBT: the obscuring of power in the name of science. *European Journal of Psychotherapy and Counselling*, 10, 231–245.
- Rao, A. S., Hendry, G., & Watson, R. (2010). The implementation of routine outcome measures in a tier 3 psychological therapies service: the process of enhancing data quality and reflections of implementation challenges. *Counselling and Psychotherapy Research*, 10, 32–38.
- Read, J. (2007). Why promoting biological ideology increases prejudice against people labelled 'schizophrenic'. *Australian Psychologist*, 42, 118–128. <https://doi.org/10.1080/00050060701280607>
- Richardson, T., Dasyam, B., Courtney, H., White, L., Tedbury, J., Butt, J., & Newman-Taylor, K. (2019). Predictors of disengagement from cognitive behavioural therapy for psychosis in a National Health Service setting: a retrospective evaluation. *British Journal of Clinical Psychology*, 58, 440–451.
- Riehle, M., Böhl, M. C., Pillny, M., & Lincoln, T. M. (2020). Efficacy of psychological treatments for patients with schizophrenia and relevant negative symptoms: a meta-analysis. *Clinical Psychology in Europe*, 2, e2899.
- Rogers, C. R. (1957). The necessary and sufficient conditions of therapeutic personality change. *Journal of Consulting Psychology*, 21, 95–103. <https://doi.org/10.1037/h0045357>
- Rosenzweig, S. (1936). Some implicit common factors in diverse methods of psychotherapy. *American Journal of Orthopsychiatry*, 6, 412–415. <https://doi.org/10.1111/j.1939-0025.1936.tb05248.x>
- Roth, A. D., & Pilling, S. (2013). *A Competence Framework for Psychological Interventions with People with Psychosis and Bipolar Disorder*. [www.ucl.ac.uk/pals/research/clinical-educational-and-health-psychology/research-groups/core/competence-frameworks](http://www.ucl.ac.uk/pals/research/clinical-educational-and-health-psychology/research-groups/core/competence-frameworks)
- Royal College of Psychiatrists. (2024). *Patient Reported Outcome Measures for People With Severe Mental Illness in Community Mental Health Settings Implementation Guidance*. [www.rcpsych.ac.uk](http://www.rcpsych.ac.uk)
- Royce, J. R. (1978). How we can best advance the construction of theory in psychology. *Canadian Psychological Review*, 19, 259–276. <https://doi.org/10.1037/h0081480>
- Royce, J. R., & Smith, W. A. (1964). A note on the development of the Psycho-Epistemological Profile (PEP). *Psychological Reports*, 14, 297–298. <https://doi.org/10.2466/pr0.1964.14.1.297>
- Safran, J. D., Segal, Z. V., Vallis, T. M., Shaw, B. F., & Samstag, L. W. (1993). Assessing patient suitability for short-term cognitive therapy with an interpersonal focus. *Cognitive Therapy and Research*, 17, 23–38.
- Schlief, M., Rich, N., Rains, L. S., Baldwin, H., Rojas-Garcia, A., Nyikavaranda, P., . . . & Johnson, S. (2023). Ethnic differences in receipt of psychological interventions in Early Intervention in Psychosis services in England—a cross-sectional study. *Psychiatry Research*, 330, 115529.
- Sin, J., & Spain, D. (2016). Psychological interventions for trauma in individuals who have psychosis: a systematic review and meta-analysis. *Psychosis*, 9, 67–81. <https://doi.org/10.1080/17522439.2016.1167946>
- Sitko, K., Bewick, B., Owens, D. & Masterson, C. (2020). Meta-analysis and meta-regression of cognitive behavioral therapy for psychosis (CBTp) across time: the effectiveness of CBTp has improved for delusions. *Schizophrenia Bulletin Open*, 1, 10. [10.1093/schizbullopen/sgaa023](https://doi.org/10.1093/schizbullopen/sgaa023).
- Skrobinska, L., Newman-Taylor, K., & Carnelley, K. (2024). Psychosis and help-seeking behaviour – a systematic review of the literature. *Psychology and Psychotherapy: Theory, Research and Practice*, 97, 583–605.
- Solmi, M., Croatto, G., Piva, G., Rosson, S., Fusar-Poli, P., Rubio, J. M., Carvalho, A. F., Vieta, E., Arango, C., DeTore, N. R., Eberlin, E. S., Mueser, K. T., & Correll, C. U. (2023). Efficacy and acceptability of psychosocial interventions in schizophrenia: systematic overview and quality appraisal of the meta-analytic evidence. *Molecular Psychiatry*, 28, 354–368.

- Substance Abuse and Mental Health Services Administration** (2014) SAMHSA's Concept of Trauma and Guidance for a Trauma-Informed Approach. HHS Publication No. (SMA) 14-4884. Rockville, MD: Substance Abuse and Mental Health Services Administration.
- Tarrier, N., Beckett, R., Harwood, S., Baker, A., Yusupoff, L., & Ugarteburu, I.** (1993). A trial of two cognitive-behavioural methods of treating drug-resistant residual psychotic symptoms in schizophrenic patients: I. Outcome. *British Journal of Psychiatry*, *162*, 524-532.
- van den Berg, M., Magaraggia, I., Schreiber, R., Hillhouse, T. M., & Porter, J. H.** (2022). How to account for hallucinations in the interpretation of the antidepressant effects of psychedelics: a translational framework. *Psychopharmacology*, *239*, 1853-1879. <https://doi.org/10.1007/s00213-022-06106-8>
- Varese, F., Smeets, F., Drukker, M., Lieverse, R., Lataster, T., Viechbauer, W., Read, J., van Os, J. & Bentall, R. P.** (2012). Childhood adversities increase the risk of psychosis: a meta-analysis of patient control, prospective and cross-sectional cohort studies. *Schizophrenia Bulletin*, *38*, 661-671.
- Waller, G., & Turner, H.** (2016). Therapist drift redux: why well-meaning clinicians fail to deliver evidence-based therapy, and how to get back on track. *Behaviour Research and Therapy*, *77*, 129-137. <https://doi.org/10.1016/j.brat.2015.12.005>
- Wampold, B.** (2001). Contextualizing psychotherapy as a healing practice: culture, history, and methods. *Applied and Preventive Psychology*, *10*, 69-86
- Wampold, B. E., & Imel, Z. E.** (2015). *The Great Psychotherapy Debate: The Evidence for What Makes Psychotherapy Work*. Routledge.
- Wood, L., Williams, C., Billings, J., & Johnson, S.** (2020). A systematic review and meta-analysis of cognitive behavioural informed psychological interventions for psychiatric inpatients with psychosis. *Schizophrenia Research*, *222*, 133-144.