The impact of cognitive behavioural therapy for insomnia on objective sleep parameters: A meta-analysis and systematic review

Mitchell, L.¹, Bisdounis, L.^{1,2}, Ballesio A.³, Omlin, X.,¹ Kyle, S.D*.¹

¹Sleep and Circadian Neuroscience Institute, Nuffield Department of Clinical Neurosciences, University of Oxford, Oxford, UK

²Department of Clinical Psychology, University of Amsterdam, Amsterdam, the Netherlands

³Department of Psychology, Sapienza University of Rome, Italy

*Correspondence should be addressed to Dr Simon Kyle:

simon.kyle@ndcn.ox.ac.uk

+44 (0) 1865 618675

Sleep and Circadian Neuroscience Institute (SCNi), Nuffield Department of Clinical Neurosciences, Sir William Dunn School of Pathology, University of Oxford, South Parks Road, Oxford, United Kingdom, OX1 3RE

Acknowledgements

The authors disclose that they have no potential conflict of interests relating to the material presented in this manuscript. This research was supported by the National Institute for Health Research (NIHR) Oxford Biomedical Research Centre (BRC). The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health.

LB was supported by an Erasmus+ internship grant.

Author contributions

LM and SDK conceived the original idea and designed the data selection criteria. LM conducted the literature search and paper screening. LB and AB extracted and cross-validated data under the supervision of SDK. LB planned and performed the statistical analysis under the supervision of SDK. AB cross-checked the analysis scripts. LM, LB and SDK drafted the manuscript. All authors contributed to and approved the final manuscript.

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

Summary

It is well-established that cognitive behavioural therapy for insomnia (CBT-I) improves self-reported sleep disturbance, however the impact on objective sleep is less clear. This meta-analysis aimed to quantify the impact of multi-component CBT-I on objective measures of sleep, indexed via polysomnography (PSG) and actigraphy. Fifteen studies met inclusion criteria. Following appraisal for risk of bias, extracted data were meta-analysed using random-effects models. The quality of the literature was generally high, although reporting of methodological detail varied markedly between studies. Meta-analyses found no evidence that CBT-I reliably improves PSG-defined sleep parameters. Actigraphy evidence was more mixed; with a small positive effect for reduction in sleep onset latency (Hedge's $g = -0.28$ [95% confidence interval (CI) -0.51 to -0.05], $p = 0.018$) and a moderate effect for reduction in total sleep time (TST) (Hedge's $g = -0.51$ [95% CI -0.75 to -0.26], $p < 0.001$). In contrast, and consistent with recent meta-analyses, CBT-I was associated with robust improvements in diary measures of sleep initiation and maintenance (Hedge's g range = 0.50 to 0.79) but not TST. While the literature is small and still developing, the sleep benefits of CBT-I are more clearly expressed in the subjective versus objective domain.

Keywords: Cognitive behavioural therapy; sleep; insomnia; actigraphy; polysomnography; meta-analysis.

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON
OBJECTIVE SLEEP

List of abbreviations

ACT = Actigraphy

BBTI = brief Behavioural Treatment for Insomnia

CBT-I = cognitive Behavioural Therapy for Insomnia

CC = control condition

CI = confidence interval

CMI = comorbid insomnia

CR = cognitive restructuring

CT = cognitive therapy

DSM(IV-TR) = Diagnostic and Statistical Manual of Mental Disorders (IV-Text Revision)

f. = female

f/up = follow up

IC = information control

ICBT-I = internet cognitive behavioural therapy for insomnia

ICSD = International Classification of Sleep Disorders

ISR = intensive sleep retraining

MCI = multi-component intervention

Min = minutes

Mo = month(s)

PE = psychoeducation

PI = primary insomnia

PSG = polysomnography

PT = placebo therapy (psychological)

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON
OBJECTIVE SLEEP

RCTs = randomised controlled trials

RDC = research diagnostic criteria

REL = relaxation

RP = relapse prevention

RT = relaxation therapy

SC = sleep compression

SCT = stimulus control therapy

SD = standard deviation

SE = sleep efficiency

SH = sleep hygiene

SOL = sleep onset latency

SRT = sleep restriction therapy

SS = sleep seminar

TCC = Tai Chi Chih

TST = total sleep time

TWT = total wake time at night

WASO = wake after sleep onset

Wk(s) = week(s)

Yrs = year(s)

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

Introduction

Insomnia is characterised by perceived difficulty initiating or maintaining sleep, which in turn has a negative influence on daytime functioning [1]. Reports on the prevalence of insomnia are variable [2], with one estimate suggesting that one third of the adult population displays at least one symptom of insomnia, while 6% meet diagnostic criteria [3]. It is estimated that the prevalence of insomnia is steadily increasing [4]. Chronic insomnia is associated with substantial impairment to quality of life, loss of functioning and increased health risk [5, 6]. It can also be a risk factor or trigger for mental health difficulties such as depression [7], psychotic experiences [8, 9] and episodes of mania or depression in bipolar disorder [10]. Treatment of insomnia is important, both as a condition in its own right [11] and as a comorbid condition in those with psychiatric or medical diagnoses [12].

The most common clinical response to insomnia is the prescription of hypnotic drugs [13], despite cognitive behavioural therapy for insomnia (CBT-I) being recommended as the first-line treatment in clinical guidelines [14-16]. CBT-I is considered to have well-established efficacy [17-19] and its effects may be more long-lasting than medication [20]. A number of previous reviews demonstrate the efficacy of CBT-I when measured by sleep diaries [11, 19, 21, 22], however relatively few studies collect or report on objective sleep outcomes, measured by polysomnography (PSG) and actigraphy. As such, the impact of CBT-I on objective sleep has been less thoroughly investigated [18]. This is in contrast to literature on the pharmacological management of insomnia [23-25] or treatments for other common sleep disorders, like obstructive sleep apnoea, where PSG measurement is routine in intervention trials [26].

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

While contemporary diagnostic criteria [1, 27] clearly consider insomnia to be a disorder based on the perception of sleep, research guidelines emphasise the assessment of *both* objective and subjective sleep [28]. Consistent with this, meta-analyses show that insomnia is characterised by objective sleep impairment relative to normal sleeping controls (e.g., in PSG-defined SOL, SE, and TST) [29]. Assessment of objective sleep is also important because insomnia with objective short sleep may be particularly linked to poorer health outcomes and impaired cognitive performance [30], as well as blunted response to CBT-I [31].

Assuming that objective sleep disturbance is a reliable feature of insomnia disorder, a logical extension of the above literature is that a) if CBT-I truly “treats” insomnia we should observe amelioration of objective sleep impairment; and b) CBT-I is likely to confer greater health benefits if it can improve objective indices of sleep (particularly for those who exhibit pronounced baseline sleep deficits). One previous meta-analysis, published in 2011, reported on changes in objectively-defined sleep (PSG and/or actigraphy). This review observed medium effect sizes at post-treatment for wake-time after sleep onset and sleep efficiency, with no evidence of improvement in objective total sleep time [18]. However, the authors combined actigraphy and PSG when computing effect size estimates - limiting precision - and did not perform quality appraisal of studies. Moreover, since 2011, several published studies have included objective measures of sleep [32-38]. Clearly the field requires a contemporary assessment of the effects of CBT-I on objective sleep to advance understanding and guide both future studies and treatment refinement. We set out to undertake a systematic search of the insomnia treatment literature with the aim of quantitatively and qualitatively

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP
 appraising randomised controlled trials (RCTs) of multi-component CBT-I that report on objective measures of night-time sleep.

Methods

Search strategy

This systematic review and meta-analysis was conducted in line with PRISMA guidelines [39]. Published studies were identified via medical and allied health online databases EMBASE, PsycINFO, MEDLINE, CINAHL and AMED. Databases were searched from inception to 13th March 2018 using terms for cognitive behavioural therapy, insomnia and objective measures of sleep (PSG and actigraphy). The following search terms were used with the OVID database platform: "cognitive behav* therap*" or "cognitive behav* intervention" or "cognitive therap*" or "behav* therap*" or "behav* modification" or "CBT" or "CBT-I" or "ICBT" or "sleep hygiene" or "stimulus control" or "relaxation" or "sleep restriction" or "psychotherap*" AND "actigraph*" or "actimetry" or "ACT" or "polysomnograph*" or "PSG" or "objective measures" or "objective sleep" or "sleep parameters" AND "insomnia" or "chronic insomnia" or "sleeplessness" or "sleep disorder*" or "sleep initiation" or "sleep maintenance" or "poor sleep" or "sleep problem" or "sleep disturbance".

In addition, bibliographies from previous reviews and retrieved articles were hand searched [11]. Grey literature was not included. All titles and abstracts were screened by the first author to identify studies providing a CBT-I intervention to adults with insomnia disorder. If the abstract indicated that the study would potentially meet the inclusion/

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

exclusion criteria outlined, the full-text version was obtained for data extraction. Where there was uncertainty, an inclusion decision was reached by author consensus.

Study Selection

Included papers were peer reviewed and written in English. Research trials were included in the review if they met the following criteria: i) a randomised controlled trial (RCT) design, ii) the population sampled were adults (≥ 18) who had insomnia disorder, as diagnosed using either research [40] or clinical diagnostic criteria (the diagnostic and statistical manual of mental disorders (DSM) [1] or the international classification of sleep disorders (ICSD) [27] criteria) iii) the intervention was CBT-I, defined as multimodal therapy delivered on at least two occasions and incorporating at least two of the five most widely accepted components of CBT-I: cognitive therapy, stimulus control, sleep restriction, sleep hygiene (psychoeducation), and relaxation therapy. All delivery modalities (e.g. face-to-face, web-based) were eligible for inclusion and there was no requirement for the treatment to combine cognitive and behavioural components, iv) outcomes included mean differences in sleep parameters measured using validated objective measures of sleep (PSG or actigraphy), and v) studies had at least one passive control arm (i.e. wait list control, sleep hygiene education, or psychological placebo condition).

Studies were excluded if co-morbid psychiatric, medical or sleep disorders besides insomnia were listed as the *main focus* of the study, or as a participant inclusion criterion. However, studies were not excluded if the frequency of comorbid conditions in the sample were reported. Patients with insomnia seen in clinical practice are likely to have a number of

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

co-morbid conditions, therefore it was not considered pragmatic to exclude studies based on this parameter. Studies were also excluded if CBT-I was combined with alternative treatment for sleep disorders (e.g. CBT-I combined with medication) and when the control condition was focussed on pharmacologic intervention (i.e. medication withdrawal/placebo).

Data Extraction

Details of the included studies were recorded by the first author (LM) within a standardised data-extraction form. The form was constructed to extract the following study characteristics: 1) citation of the publication, 2) geographic location, 3) total number of participants and the number of female participants, 4) mean age (and standard deviation when applicable), 5) social and cultural characteristics of the population, 6) method of recruitment, 7) diagnostic criteria and mean duration of insomnia, 8) medication status, 9) co-morbidity, 10) inclusion and exclusion criteria for participation - including screening for objective sleep problems - , 11) components, format -individual or group, face to face or phone- frequency and duration of the treatment intervention and who delivered it, 12) type, frequency and duration of the control condition and who delivered it, 13) objective and subjective sleep measurements used, 14) the duration of PSG and actigraphy when applicable and 15) the format and timing of the follow-up.

Findings regarding the effect of CBT-I on objective measures at post-treatment were extracted and coded for data analysis. Although some studies reported differences within groups before and after treatment, this meta-analysis only considered differences between treatment and control groups (i.e. between-group differences) as robust evidence to assess the

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

effect of CBT-I on objective sleep. Additionally, the meta-analysis was interested in the main, super-ordinate features of objective sleep. As such, we extracted outcomes regarding the following commonly reported sleep parameters: sleep onset latency (SOL), wake-time after sleep onset (WASO), sleep efficiency (SE) and total sleep time (TST). As a secondary aim, and to permit comparison with objective dependent variables, we also extracted and calculated effects sizes for diary-reported variables (SOL, WASO, SE, TST) where available. Note, reporting of sleep diary variables was not considered a criterion for study inclusion.

Quality assessment

This meta-analysis considered whether the conclusions made were reliable and valid based on the methodological quality of the summarised research. Therefore, all included studies were independently appraised by the first and second authors using the Cochrane Collaboration's tool for assessing risk of bias [41]. This involves appraising bias in selection, performance, detection, attrition and reporting domains. In each domain, studies were given a rating of low, high or unclear risk. In the case of a discrepancy in appraisal of risk of bias between the two authors', a consensus was reached after consultation with the remaining authors.

Data synthesis and analysis

All analyses were conducted in R, version 3.5.1 [42], using the tidyverse [43] and metafor [44] packages. Effect sizes and 95% confidence intervals (CI) were calculated for between-group comparisons at post-treatment. The effect size statistic selected for this meta-analysis was Hedge's *g* [45], which is a measure of standardised mean difference. Hedge's *g*

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

is similar to Cohen's d in the sense that both metrics operate on the assumption of equal population variance. However, Cohen's d can be positively biased especially in the context of smaller sample sizes [45]. Hedge's g reduces this upward bias by weighting each group's standard deviation (SD) by its sample size and therefore using pooled weighted standard deviations (SDs). This makes Hedge's g more appropriate for studies including smaller sample sizes [46], a common characteristic for studies in clinical psychology. In this meta-analysis Hedge's g was the quotient of the difference between the mean of the experimental group against the mean of the control group, divided by the pooled weight standard deviation incorporating Bessel's correction:

$$g = \frac{\mu_{exp} - \mu_{ctrl}}{s}$$

, where μ_{exp} and μ_{ctrl} denote the mean values of the experimental and control group respectively, whilst s denotes the pooled weighted standard deviation, calculated as follows:

$$s = \frac{\sqrt{(n_{exp} - 1)SD_{exp}^2 + (n_{ctrl} - 1)SD_{ctrl}^2}}{n_{exp} + n_{ctrl} - 2}$$

, where n_{exp} and n_{ctrl} indicate the number of participants and SD_{exp} and SD_{ctrl} the standard deviation points for the experimental and control group respectively. Effect sizes of 0-0.32 are interpreted as small, effect sizes of 0.33-0.55 are moderate and effect sizes of 0.56-1.2 are assumed to be large [47]. For the effect size calculations, we used the available means and SDs as published in the included papers. Some studies reported other measures of dispersion, such as CIs [36] and standard errors [33, 34, 37, 38, 48, 49]. These metrics were transformed to SDs using the following formulas:

$$SD = ES * \sqrt{n}$$

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

$$SD = \frac{(\mu - CI.lb)}{1.96} * \sqrt{n}$$

, where CI.lb denotes the lower bound of the confidence intervals.

Two papers [37, 50] reported on two controlled post-treatment analyses. One of them [50] divided participants according to insomnia profile (i.e. primary or co-morbid insomnia) and assigned them to either the treatment or control arm. The other study [37] classified participants across both trial arms into long (≥ 6 hrs) and short (<6 hrs) sleepers based on one night of baseline PSG. These four comparisons were independently coded in the dataset and entered in the main analysis.

For each measurement and outcome, heterogeneity analyses were conducted to test for the implementation of a fixed-effects model. Cochran's Q-statistic was utilised for the heterogeneity tests. Cochran's Q is calculated as the weighted sum of squared differences between individual study effects and the pooled effect across studies, with the weights being those used in the pooling method. According to the associated Q-statistic, variance in a considerable amount of effect size estimates was larger than what would be predicted based on subject-level sampling variability. Therefore, a fixed-effects model was implemented in the sensitivity analysis for the appropriate outcomes and random effects modelling was utilised for the main analysis. Furthermore, to account for small sample sizes and achieve nominal coverage probabilities, heterogeneity was also assessed based on I^2 and the accompanying CIs as calculated using the Q-profile method [51]. I^2 indicates the percentage of total variability in effect sizes that reflects heterogeneity among the true effects and not sampling error. An I^2 value of 0 signifies 0% heterogeneity; the greater the I^2 value the greater the observed heterogeneity.

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

Sensitivity analysis was performed to assess the effect of model choice (i.e. fixed vs random modelling) on the effect size estimations. This was achieved by implementing fixed effects models to the applicable outcomes and measurements and comparing the results with those provided from random effects models. Additionally, sensitivity analysis was conducted to investigate the effect of double comparisons within a single study [37, 50]. This was achieved by re-calculating the main analysis each time including only one comparison per outcome and measurement from each study.

Publication bias was assessed via visual inspection of the funnel plots and Egger's weighted regression test for funnel plot asymmetry [52]. Wherever publication bias was detected, the Duval and Tweedie trim-and-fill method [53] was implemented to produce a model after accounting for any publication bias. This method also yields a numerical estimate of the missing studies due to the congregation of the most extreme results on one side of the funnel plots [43].

Results

After removing duplicate records (n=628), titles and abstracts of 1,701 records were screened for inclusion. This procedure resulted in the exclusion of 1,638 references and a full-text inspection of 63 citations followed. A total of 15 published studies met the inclusion criteria for this meta-analysis (see Figure 1.).

[Insert Figure 1. PRISMA flow chart of included studies]

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

Study characteristics

Table 1 presents study characteristics for all included studies. The earliest study was published in 1993 and the latest in 2017, with one study published before 2000 (6.67%), five studies published between 2000 and 2010 (33.33%), and nine studies published from 2011 onwards (60%). Three studies solely analysed PSG outcomes, 11 studies solely analysed actigraphy outcomes and two studies [49] analysed both PSG and actigraphy outcomes. Out of the 15 studies, eleven were conducted in the USA, one in the UK and three in Australia. These studies recruited a total of 1,541 participants, with sample sizes ranging between 24 and 201. Studies recruited adults and older adults from the community (n=10), patients attending a medical practice (n=1), military veterans (n= 2), active army personnel (n=1) and college students (n=1) with a mean age across studies of 57 years. A total of 47% of the sample was female; the relatively low percentage of females can be explained by the predominantly male veteran and army population in three studies.

[Insert Table 1]

In all studies, insomnia criteria (DSM or ICSD) was ascertained via clinical interview and/or sleep diaries. Ten studies excluded sleep disorders other than insomnia using clinical interview and PSG screening [32, 34, 35, 37, 50, 54], three studies used other in-home monitoring devices [36, 55, 56], whereas two studies used clinical interview only [38, 57]. No studies used objective sleep criteria for study entry, although Edinger and colleagues excluded individuals who met clinical and PSG criteria for sleep state misperception in one

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

trial [48] and in another study used data from one-night diagnostic PSG to exclude individuals for whom objective sleep time was ≥ 2 times higher than their subjectively estimated sleep time [54]. One trial [37] used baseline PSG to classify participants into short and long sleep duration groups. Hypnotic medication was allowed in five studies [36, 38, 49, 50, 57]. All studies, with one exception [32], used sleep restriction therapy, alongside an educational element of either sleep hygiene (n=5), psychoeducation (n=6), or both (n=4). Other components included cognitive techniques (n=8), relaxation (n=3) and relapse prevention (n=3). Only three RCTs did not include SCT [34, 37, 55].

The RCTs delivered weekly treatment, with the exception of one study, which delivered sessions bi-weekly [50]. A range of methods were used to deliver the intervention; face-to-face individual (n=8) and group sessions (n=6) or a comparison of both (n=1); face-to-face group sessions with individual follow up phone calls (n=1), face-to-face individual sessions with follow up phone calls (n=1) and a comparison of both face-to-face and online individual formats (n=1). In all but two studies treatment was four-to-six sessions in length. One study [33] included 16 weekly CBT sessions compared with an equal amount of Tai Chi Chi and a sleep education control group, whilst another study [54] compared one, two, four and eight sessions of CBT-I with a wait-list control. All studies indicated the use of a structured protocol, with the majority of studies specifically referencing a manualised approach or describing methods to ensure treatment fidelity such as training, supervision, evaluation of recordings and checklists. Only two studies did not reference the use of a treatment manual or measures to ensure treatment fidelity [37, 55]. The control arms included

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

wait-list (n=6), treatment as usual at a GP practice (n=1), minimal contact control (n=1), relaxation therapy (n=1), psychological placebo (n=1) and educational control (n=5).

Regarding PSG outcomes, two studies conducted PSG in a sleep lab and reported mean outcomes over two nights at each assessment time point [33, 58], with one study including an additional night for adaptation [33]. The other two studies conducted ambulatory PSG in the home environment for one night [48] or two nights [49] at each assessment point. Sleep stages were scored using standardised criteria. Only one study [58] reported that reliability checks were performed, with inter-rater agreement being greater than 85% for sleep-stage scoring.

The actigraphy outcomes were typically calculated using the mean score of one or two weeks of actigraphy data at each assessment time point. In and out of bed times were recorded via an event marker on the actigraph hardware, or via concurrent use of sleep diaries. Studies reported using scoring algorithms associated with their chosen software, however it was unclear how these may have been programmed differently. Only one study reported manually adjusting sensitivity criteria [32]. The majority of studies did not report the frequency at which the actigraph records movement, although the studies that did report this indicated either 30-second or one-minute epochs. Comprehensiveness of the description of actigraph recordings varied widely between studies, which may be expected considering objective measures were often reported as secondary outcomes.

Quality assessment

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

Appraisal of risk of bias (see table S1) yielded a mixed picture with regards to how comprehensively methodological details were reported, with only two studies reporting sufficient detail to perform a complete appraisal. However, in general, the studies were judged to have low risk of bias in most domains.

Almost all studies (n=14; 93%) generated an adequate randomisation sequence and in only one study (7%) the randomisation protocol was unclear. In five studies the allocation was adequately concealed (33%), whereas in the rest of the studies (n=10; 67%) the concealment procedure was unclear. The inherent risk of bias in psychotherapy research due to difficulty blinding participants and researchers was considered unavoidable by some authors. However, three out of the 15 studies (20%) attempted to minimise this bias by using minimally active control conditions and informing all participants that they would receive a behavioural treatment for insomnia [36, 48, 50]. The remaining studies either reported that blinding was impossible (n=6; 40%) or did not provide enough details to reach a conclusion (n=6; 40%).

Regarding blinding of outcome assessors scoring objective sleep, eleven studies (73%) did not provide enough details, three studies (20%) reported a sufficient blinding with one study (7%) reporting that blinding of assessors was not performed. Both actigraphy and PSG require analysis and interpretation [59], therefore blinding of outcome assessors is important to report. Since objective measures features as secondary outcomes it may be that full reporting on these details was overlooked. In total, 14 studies (93%) described the completeness of the outcome data and provided a good amount of details concerning their

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

attrition rates. Only one study (7%) was unclear regarding the handling of incomplete data points. In 14 studies (93%) there was no evidence for selective reporting. However, there was evidence for selective reporting in one study (7%). In this particular study [36], although individual and group CBT-I protocols were independently administered, they were merged together in the analysis with sleep measurements reported collectively from both protocols. Selective reporting was only assessed on the basis of objective measurements, meaning that selective reporting in relation to diary measurements was not accounted for.

Heterogeneity analyses

Heterogeneity analyses revealed that approximately one half of effect size estimates showed pronounced heterogeneity, while the other half were more homogeneous. Specifically, for PSG variables: SOL [$Q(3)=1.59, p=0.661$ for SOL], WASO [$Q(3) = 3.08, p=0.379$], SE [$Q(4) = 10.534, p= 0.032$] and TST [$Q(4) = 8.726, p= 0.068$]. For actigraphy variables: SOL [$Q(10) = 22.241, p= 0.014$], WASO [$Q(11) = 16.817, p= 0.113$], SE [$Q(13) = 22.103, p= 0.054$] and TST [$Q(11) = 24.385, p= 0.011$]. For sleep diary variables: SOL [$Q(15) = 33.132, p= 0.005$], WASO [$Q(15) = 37.313, p= 0.001$], SE [$Q(16) = 49.764, p < 0.001$] and TST [$Q(15) = 24.345, p= 0.060$]. Due to the pattern of heterogeneity in effect size estimates across sleep parameters, the more conservative random effects model was implemented for the main analysis. When appropriate, fixed effects modelling was also implemented as part of the sensitivity analysis.

Effects on PSG-defined sleep variables

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

See Table 2 for effect size summary and Figures 2a-d for forest plots grouped by sleep parameter. For PSG, effect sizes were predominantly small and non-significant [SE ($N_c=5$; $g=0.33$ [95% CI= -0.14 to 0.80], $p=0.169$); WASO ($N_c=4$; $g=-0.10$ [95% CI= -0.39 to 0.19, $p=0.509$); SOL ($N_c=4$; $g=-0.07$ [95% CI= -0.36 to 0.22], $p=0.641$; and TST ($N_c=5$; $g=0.04$ [95% CI= -0.37 to 0.45], $p=0.848$)]. It is noteworthy that the highest number of available comparisons (N_c) was just five.

[Insert Table 2]

Effects on actigraphy-defined sleep variables

A small, reliable effect was observed for a decrease in actigraphy-defined SOL ($N_c=11$; $g=-0.28$ [95% CI= -0.51 to -0.05], $p=0.018$), while a negative effect of moderate size was observed for TST ($N_c=12$; $g=-0.51$ [95% CI= -0.75 to -0.26], $p<0.001$), indicating a *reduction* of sleep time in the experimental versus control arm. There was no significant effect for WASO ($N_c=12$; $g=-0.14$ [95% CI= -0.33 to 0.05], $p=0.138$) or SE ($N_c=14$; $g=0.14$ [95% CI= -0.04 to 0.31], $p=0.137$).

Effects on sleep diary variables

Moderate-to-large effects were found for SOL ($N_c=16$; $g=-0.50$ [95% CI= -0.70 to -0.30], $p<0.001$), SE ($N_c=17$; $g=0.79$ [95% CI= 0.57 to 1.02], $p<0.001$) and WASO ($N_c=16$; $g=0.75$ [95% CI= -0.96 to -0.55], $p<0.001$), while there was no reliable difference for TST ($N_c=16$; $g=0.06$ [95% CI= -0.12 to 0.23], $p=0.537$).

[Insert Figure 2a-d]

Fig 2. Forest plots showing individual study and global effect size estimates for comparisons between CBT-I and control groups on (a) sleep efficiency; (b) sleep onset latency; (c) total sleep time; and (d) wake-time after sleep onset as measured by polysomnography, actigraphy and sleep diary.

Sensitivity analyses

Sensitivity analysis employing fixed-effect modelling was conducted for the post-treatment outcome measures where no significant heterogeneity was detected, yielding comparable results (see table S2). Sensitivity analysis was also conducted to control for the effect of double comparisons within a study for the same outcome (as quantified by the same measurement). The re-calculations of the random effects model resulted in highly comparable values as those reported in the main analysis.

Publication bias analysis

Visual inspection of the funnel plots indicated that there was no considerable evidence of publication bias across all outcomes and measurements for post-treatment. Egger's regression tests confirmed the results from the visual inspections, with no study yielding a z-score larger than 1.50 and a p-value of less than 0.135.

Discussion

This meta-analysis sought to evaluate the extent to which multicomponent CBT-I improves objective measures of sleep in insomnia disorder, whilst simultaneously appraising the quality of the evidence-base. Our results find no evidence that CBT-I reliably improves

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

PSG sleep parameters relative to control groups at post-treatment, although it must be noted that very few studies incorporated PSG measurement ($n=5$). For actigraphy, where we had more comparisons, there was a small positive effect for SOL ($g= -0.28$) and a moderate (negative) effect for TST ($g= -0.51$). In contrast, and while not a primary aim of our study, there was clear evidence that CBT-I results in robust improvements in diary measures of SOL ($g= -0.50$), WASO ($g= -0.75$) and SE ($g= 0.79$), but not TST ($g= 0.06$), which is consistent with other recent meta-analyses [11] reporting effect sizes of $g=0.57$ (SOL), $g=0.63$ (WASO), $g=0.71$ (SE) and $g=0.16$ (TST).

The most robust effects for objective sleep in this review were found for TST, reflecting a *reduction* in actigraphy-defined sleep time of approximately 30 minutes relative to the control group. This decrease may reflect an *initial* reduction in sleep time driven by sleep restriction therapy and/or stimulus control in order to consolidate sleep [60], however it is noteworthy that we did not observe reliable reductions in WASO or improvements in SE (for either actigraphy or PSG) and only a small relative reduction of approximately 6 mins for actigraphy-defined sleep latency. The assumption is that any reduction in TST is transient, however we conducted exploratory (and by extension, cautionary) meta-analyses of studies with follow-up data at three months and found a sustained effect for reduced (actigraphy-defined) total sleep time ($N_c=4$; $g=-0.56$, $p<.001$ see Table S3). Future studies, particularly those employing both PSG and actigraphy, are needed to systematically track changes in sleep *during* CBT treatment and over long-term follow-up.

While design differences between psychotherapy and pharmacotherapy trials make it difficult to directly compare treatment effect size estimates (across meta-analyses) it appears

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

that hypnotics have more pronounced effects on PSG-defined sleep than CBT. A meta-analysis of both published and unpublished trials of z drugs reported a small, but reliable effect size (SMD) of -0.36 (CI: -0.57 to -0.16) for PSG-defined SOL relative to placebo [23]. This effect, of course, is greater than what we observed in the present analysis ($g=-0.07$ CI: -0.36 to 0.22). A recent Cochrane review of eszopiclone versus placebo also showed reliable reductions in PSG-defined SOL and WASO, and increased TST, although number of comparisons was small [25]. In contrast to our findings, hypnotics seem to have similar effects across both objective and subjective sleep parameters, while we see pronounced effects on subjective but not PSG-defined variables. However, results must be interpreted cautiously because our review clearly shows that objective sleep measurement is rarely applied in trials of CBT-I and sample sizes are typically small (contrasting with studies of hypnotics).

A recent comprehensive meta-analysis reported on sleep diary data from 87 studies [11]. While our meta-analysis, in comparison, included a much smaller number of studies with sleep diary data, results show a similar pattern and magnitude of effects, supporting the view that CBT-I principally modifies subjective perceptions of sleep (see Figures 2a-d). Indeed, previous research indicates that CBT-I “corrects” the discrepancy between self-reported sleep and objective sleep, while having no significant impact on PSG-defined sleep at post-treatment [61]. Conceivably, CBT may modify key psychobiological processes that maintain *perceptions* of poor sleep and daytime impairment in insomnia, such as dysfunctional arousal, sleep incompatible behaviours and key cognitive processes, without having pronounced therapeutic effects on *gross* measures of objective sleep [62, 63]. However, it should be noted that we did not assess number of awakenings, sleep architecture or microstructure in the

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

present review, instead focusing on more commonly reported sleep continuity measures. It is possible that CBT modifies EEG indices of arousal (e.g., beta and gamma spectral power) and sleep intensity (delta power) but given the paucity of PSG studies, the evidence-base for these variables is assumed to be limited. Future, adequately-powered and pre-registered studies, are needed to a) test the effects of CBT on sleep physiology and b) investigate associations with important clinical outcomes.

Ability to show change in objective sleep likely depends on sample characteristics. Studies in our review recruited participants based on self-report data - consistent with diagnostic criteria - and not on objective sleep. In contrast, trials of hypnotics often recruit participants on the basis of PSG-confirmed sleep disruption [64, 65]. It is possible, therefore, that in the studies we reviewed there was relatively little scope to change objective sleep. One recent, uncontrolled study of inpatient CBT-I [66], found that patients with objective sleep impairment (TST<6hrs) at baseline were more likely to show PSG improvements in WASO at 6 month follow-up, compared to those with relatively normal sleep (defined as ≥ 6 hrs sleep duration). Interestingly, those with normal sleep duration were more likely to report a post-treatment *decrease* in PSG-defined total sleep time, consistent with actigraphy observations in the present review, as well as results from a recent secondary analysis of a randomised trial [31]. To fully understand the impact of CBT-I on objective sleep, studies may need to subgroup based on baseline sleep profiles and assess differential change at post-treatment and follow-up.

Study quality

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

Overall, reviewed studies were judged to have low risk of bias in most domains, lending credibility to our review findings. However, our review revealed a mixed picture regarding reporting of methodological details, which made it more difficult to establish potential bias in objective measurement. Comprehensiveness of the description of PSG and actigraphy recordings and scoring algorithms varied widely between studies. This may be expected considering objective measures were often reported as secondary outcomes. For example, only two RCTs reported their definition of PSG SOL, which were clearly different from one another (latency to stage 2 sleep [58], and the first 10 minutes of sleep containing no more than 2 minutes of wake time, stage 1, or movement time [48]). Moreover, studies rarely reported details on blinding status of outcome assessors in relation to objective measurement, which is a potentially important source of bias because actigraphy and PSG require analysis and interpretation [59]. In order to facilitate consistency between trials and to appraise sources of bias, studies should more fully describe procedures for the set-up and recording of sleep (e.g., how the recording period was defined in PSG studies), as well as the analysis and interpretation (including blinding status) of objective data.

Limitations

The narrow inclusion criteria of this review may be considered a limitation, since this led to the exclusion of two recent, potentially relevant studies, as a result of their insufficiently stringent diagnostic criteria [67, 68]. Findings may not generalise to insomnia complaints in the general population, where a significant proportion experience symptoms of insomnia but do not meet diagnostic criteria and/or experience significant comorbid medical or psychiatric disorders [3]. Most studies included in this review used a highly selected

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

sample via the exclusion of other sleep disorders, dementia/mild cognitive impairment and uncontrolled physical or psychiatric difficulties. Although this reduces the influence of potentially confounding factors, it also limits the generalisability of review findings. Future work should aim to synthesise evidence on the objective sleep effects of CBT-I in comorbid insomnia samples. This review sought to make conclusions based on a relatively homogeneous set of studies; therefore, narrow inclusion criteria were considered appropriate. Despite this, differences between the studies included in this review should also be considered, particularly given observed heterogeneity for some of the treatment effect estimates. First, it must be noted that studies with very small samples (for example in the case of Friedman et al, $n=8$ for CBT vs $n=4$ for control) can strongly influence estimates from random-effects models and contribute to heterogeneity. Second, treatment protocols included in this meta-analysis differed with respect to component composition, treatment duration and format, and therapist background. While this grouping is common in CBT meta-analyses, in part because such variables appear to have limited impact on clinical outcomes [11], the effect on objective measures should be systematically interrogated in pre-defined sub-group analyses as the field matures. A final point is that hypnotic medication was *permitted* in five studies [36, 38, 49, 50, 57], which may impact objective sleep outcomes, particularly if there are between-group differences in the proportion taking medication at post-treatment. We found that only one of these studies reported hypnotic medication use by study arm at post-treatment; future studies should report such information in order to aid interpretation of effects and permit sub-group analyses when synthesising across studies.

Conclusion

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

Insomnia is a highly prevalent condition that has a considerable negative impact on daytime functioning and health. Objective measurement of sleep is relevant since insomnia is characterised by reliable impairments in objective sleep, and such impairments may confer risk for adverse medical and cognitive outcomes. While the literature is small and still developing, CBT-I appears to have limited impact on objective sleep continuity, a small effect for decreasing actigraphy-defined sleep initiation and a robust effect (of moderate strength) for decreasing actigraphy-defined sleep duration. Future, well-designed studies, with low risk of bias and transparent reporting are needed to specifically test the impact of CBT-I on objective sleep, including sleep microstructure.

Practice Points

- 1) Objective sleep outcomes are rarely investigated in trials of CBT-I
- 2) While the literature is small, this meta-analysis finds no evidence that CBT-I modifies PSG measures of sleep initiation, maintenance or duration. Conversely, actigraphy shows a small reduction in sleep onset latency and moderate reduction in total sleep time.
- 3) The sleep benefits of CBT-I are more clearly expressed in the subjective versus objective domain.

Research Agenda

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

- 1) Studies should clearly report methodological details of actigraphic and polysomnographic assessments, as well as information on use of prescription sleep aids during these assessments
- 2) Studies should include objective parameters (including SOL, WASO, SE, TST, number of awakenings and arousals), alongside subjective sleep outcomes, which should be tracked *throughout treatment* and assessed at both post-treatment and at follow-up.
- 3) Prospective randomised trials, stratifying participants by objective sleep profiles (e.g., normal and short sleep duration) are needed to understand effects on post-treatment objective sleep.
- 4) Studies should determine whether degree of change in objective sleep, pre-to-post CBT-I, mediates improvement in relevant health and disease outcomes.
- 5) Objective measures beyond gross sleep architecture or continuity, such as quantitative EEG metrics and sleep microstructure, should be examined in future RCTs (and potentially through secondary analyses of completed studies).

References

- [1] American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders. 5th ed. Washington, DC: American Psychiatric Association; 2013.
- [2] Grewal R, Doghramji K. Epidemiology of insomnia. In: Attarian HP, Schuman C, editors. Clinical handbook of insomnia. 2nd ed: Humana Press; 2010. p. 13-22.
- [3] Ohayon MM. Epidemiology of insomnia: What we know and what we still need to learn. Sleep Med Rev. 2002;6:97-111.

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

- [4] Calem M, Bisla J, Begum A, Dewey M, Bebbington PE, Brugha T, et al. Increased prevalence of insomnia and changes in hypnotics use in England over 15 Years: Analysis of the 1993, 2000, and 2007 national psychiatric morbidity surveys. *Sleep*. 2012;35:377-84.
- [5] Roth T. Insomnia: Definition, prevalence, etiology, and consequences. *J Clin Sleep Med*. 2007;3:7-10.
- [6] Kyle SD, Morgan K, Espie CA. Insomnia and health-related quality of life. *Sleep Med Rev*. 2010;14:69-82.
- [7] Baglioni C, Battagliese G, Feige B, Spiegelhalder K, Nissen C, Voderholzer U, et al. Insomnia as a predictor of depression: A meta-analytic evaluation of longitudinal epidemiological studies. *J Affect Disord*. 2011;135:10-9.
- [8] Reeve S, Sheaves B, Freeman D. The role of sleep dysfunction in the occurrence of delusions and hallucinations: A systematic review. *Clin Psychol Rev*. 2015;42:96-115.
- [9] Mulligan LD, Haddock G, Emsley R, Neil ST, Kyle SD. High resolution examination of the role of sleep disturbance in predicting functioning and psychotic symptoms in schizophrenia: A novel experience sampling study. *J Abnorm Psychol*. 2016;125:788-97.
- [10] Lewis KS, Gordon-Smith K, Forty L, Di Florio A, Craddock N, Jones L, et al. Sleep loss as a trigger of mood episodes in bipolar disorder: individual differences based on diagnostic subtype and gender. *Br J Psychiatry*. 2017;211:169-74.
- [11] *van Straten A, van der Zweerde T, Kleiboer A, Cuijpers P, Morin CM, Lancee J. Cognitive and behavioral therapies in the treatment of insomnia: a meta-analysis. *Sleep Med Rev*. 2017.
- [12] Ballesio A, Aquino MRJV, Feige B, Johann AF, Kyle SD, Spiegelhalder K, et al. The effectiveness of behavioural and cognitive behavioural therapies for insomnia on depressive

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

and fatigue symptoms: a systematic review and network meta-analysis. *Sleep Med Rev.*

2018;37:114-29.

[13] Siriwardena AN, Apekey T, Tilling M, Dyas JV, Middleton H, Ørner R. General practitioners' preferences for managing insomnia and opportunities for reducing hypnotic prescribing. *J Eval Clin Pract.* 2010;16:731-7.

[14] National Institute for Health and Care Excellence. Clinical Knowledge Summaries: Insomnia. National Institute for Health and Care Excellence; 2015.

[15] Qaseem A, Kansagara D, Forciea MA, Cooke M, Denberg TD. Management of chronic insomnia disorder in adults: a clinical practice guideline from the American college of physicians management of chronic insomnia disorder in adults. *Ann Intern Med.* 2016;165:125-33.

[16] Riemann D, Baglioni C, Bassetti C, Bjorvatn B, Dolenc Groselj L, Ellis JG, et al. European guideline for the diagnosis and treatment of insomnia. *J Sleep Res.* 2017;26:675-700.

[17] Irwin MR, Cole JC, Nicassio PM. Comparative meta-analysis of behavioral interventions for insomnia and their efficacy in middle-aged adults and in older adults 55+years of age. *Health Psychol.* 2006;25:3-14.

[18] Okajima I, Komada Y, Inoue Y. A meta-analysis on the treatment effectiveness of cognitive behavioral therapy for primary insomnia. *Sleep Biol Rhythms.* 2011;9:24-34.

[19] Trauer JM, Qian MY, Doyle JS, Rajaratnam SW, Cunnington D. Cognitive behavioral therapy for chronic insomnia: A systematic review and meta-analysis. *Ann Intern Med.* 2015;163:191-204.

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

- [20] Mitchell MD, Gehrman P, Perlis M, Umscheid Ca. Comparative effectiveness of cognitive behavioral therapy for insomnia: a systematic review. *BMC Fam Pract.* 2012;13:40.
- [21] Montgomery P, Dennis JA. Cognitive behavioural interventions for sleep problems in adults aged 60+. *Cochrane Database Syst Rev.* 2003;1.
- [22] Koffel EA, Koffel JB, Gehrman PR. A meta-analysis of group cognitive behavioral therapy for insomnia. *Sleep Med Rev.* 2015;19:6-16.
- [23] Huedo-Medina TB, Kirsch I, Middlemass J, Klonizakis M, Siriwardena AN. Effectiveness of non-benzodiazepine hypnotics in treatment of adult insomnia: meta-analysis of data submitted to the Food and Drug Administration. *BMJ : British Medical Journal.* 2012;345.
- [24] Buscemi N, Vandermeer B, Friesen C, Bialy L, Tubman M, Ospina M, et al. The efficacy and safety of drug treatments for chronic insomnia in adults: a meta-analysis of RCTs. *J Gen Intern Med.* 2007;22:1335-50.
- [25] Rosner S, Englbrecht C, Wehrle R, Hajak G, Soyka M. Eszopiclone for insomnia. *Cochrane Database Syst Rev.* 2018;10:Cd010703.
- [26] Jordan AS, McSharry DG, Malhotra A. Adult obstructive sleep apnoea. *Lancet (London, England).* 2014;383:736-47.
- [27] American Academy of Sleep Medicine. International classification of sleep disorders. 3rd Edition ed. Rochester: American Academy of Sleep Medicine; 2014.
- [28] *Buysse DJ, Ancoli-Israel S, Edinger JD, Lichstein KL, Morin CM. Recommendations for a standard research assessment of insomnia. *Sleep.* 2006;29:1155-73.

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

- [29] *Baglioni C, Regen W, Teghen A, Spiegelhalder K, Feige B, Nissen C, et al. Sleep changes in the disorder of insomnia: A meta-analysis of polysomnographic studies. *Sleep Med Rev.* 2014;18:195-213.
- [30]*Vgontzas AN, Fernandez-Mendoza J, Liao D, Bixler EO. Insomnia with objective short sleep duration: the most biologically severe phenotype of the disorder. *Sleep Med Rev* 2013;4:241-254..
- [31]*Bathgate CJ, Edinger JD, Krystal AD. Insomnia patients with objective short sleep duration have a blunted response to cognitive behavioral therapy for insomnia. *Sleep.* 2017;40:123-32.
- [32] Harris J, Lack L, Kemp K, Wright H, Bootzin R. A randomized controlled trial of intensive sleep retraining (ISR): a brief conditioning treatment for chronic insomnia. *Sleep.* 2012;35:49-60.
- [33] Irwin MR, Olmstead R, Carrillo C, Sadeghi N, Breen EC, Witarama T, et al. Cognitive behavioral therapy vs. tai chi for late life insomnia and inflammatory risk: a randomized controlled comparative efficacy trial. *Sleep.* 2014;37:1543-52.
- [34] Lovato N, Lack L, Wright H, Kennaway DJ. Evaluation of a brief treatment program of cognitive behavior therapy for insomnia in older adults. *Sleep.* 2014;37:117-26.
- [35] Taylor DJ, Zimmerman MR, Gardner CE, Williams JM, Grieser EA, Tatum JI, et al. A pilot randomized controlled trial of the effects of cognitive-behavioral therapy for insomnia on sleep and daytime functioning in college students. *Behav Ther.* 2014;45:376-89.
- [36] Alessi C, Martin JL, Fiorentino L, Fung CH, Dzierzewski JM, Rodriguez Tapia JC, et al. Cognitive behavioral therapy for insomnia in older veterans using nonclinician sleep coaches: randomized controlled trial. *J Am Geriatr Soc.* 2016;64:1830-8.

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

- [37] Lovato N, Lack L, Kennaway DJ. Comparing and contrasting therapeutic effects of cognitive-behavior therapy for older adults suffering from insomnia with short and long objective sleep duration. *Sleep Med.* 2016;22:4-12.
- [38] Taylor DJ, Peterson AL, Pruiksma KE, Young-McCaughan S, Nicholson K, Mintz J, et al. Internet and in-person cognitive behavioral therapy for insomnia in military personnel: A randomized clinical trial. *Sleep.* 2017;40.
- [39] Moher D, Shamseer L, Clarke M, Gherzi D, Liberatî A, Petticrew M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Systematic Reviews.* 2015;4.
- [40] Edinger JD, Bonnet MH, Bootzin RR, Doghramji K, Dorsey CM, Espie CA, et al. Derivation of research diagnostic criteria for insomnia: report of an American academy of sleep medicine work group. *Sleep.* 2004;27:1567-96.
- [41] Higgins JPT, Green S. *Cochrane handbook for systematic reviews of interventions.* 4th ed. Chichester: Wiley-Blackwell; 2011.
- [42] Team RC. *R: A Language and Environment for Statistical Computing.* Foundation for Statistical Computing. 2018.
- [43] Viechtbauer W. Conducting meta-analyses in R with the metafor package. *Journal of statistical software.* 2010;36.
- [44] Wickham H. *Tidyverse: Easily install and load'tidyverse'packages.* R package version. 2017;1.
- [45] Hedges LV. Distribution theory for Glass's estimator of effect size and related estimators. *journal of Educational Statistics.* 1981;6:107-28.

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

[46] Grissom RJ, Kim JJ. Effect sizes for research: A broad practical approach: Lawrence

Erlbaum Associates Publishers; 2005.

[47] Lipsey MW, Wilson DB. The efficacy of psychological, educational, and behavioral

treatment: Confirmation from meta-analysis. *Am Psychol.* 1993;48:1181.

[48] Edinger JD, Wohlgemuth WK, Radtke RA, Marsh GR, Quillian RE. Cognitive

behavioral therapy for treatment of chronic primary insomnia: a randomized controlled trial. *J*

Am Med Assoc. 2001;285:1856-64.

[49] Buysse DJ, Germain A, Moul DE, Franzen PL, Brar LK, Fletcher ME, et al. Efficacy of

brief behavioral treatment for chronic insomnia in older adults. *Arch Intern Med.*

2011;171:887-95.

[50] Edinger JD, Olsen MK, Stechuchak KM, Means MK, Lineberger MD, Kirby A, et al.

Cognitive behavioral therapy for patients with primary insomnia or insomnia associated predominantly with mixed psychiatric disorders: a randomized clinical trial. *Sleep.*

2009;32:499-510.

[51] Viechtbauer W. Confidence intervals for the amount of heterogeneity in meta-analysis.

Stat Med. 2007;26:37-52.

[52] Egger M, Smith GD, Schneider M, Minder C. Bias in meta-analysis detected by a

simple, graphical test. *BMJ.* 1997;315:629-34.

[53] Duval S, Tweedie R. Trim and fill: a simple funnel-plot-based method of testing and

adjusting for publication bias in meta-analysis. *Biometrics.* 2000;56:455-63.

[54] Edinger JD, Wohlgemuth WK, Radtke RA, Coffman CJ, Carney CE. Dose-response

effects of cognitive-behavioral insomnia therapy: a randomized clinical trial. *Sleep.*

2007;30:203-12.

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

- [55] Friedman L, Benson K, Noda A, Zarcone V, Wicks DA, O'Connell K, et al. An actigraphic comparison of sleep restriction and sleep hygiene treatments for insomnia in older adults. *J Geriatr Psychiatry Neurol.* 2000;13:17-27.
- [56] Epstein DR, Sidani S, Bootzin RR, Belyea MJ. Dismantling multicomponent behavioral treatment for insomnia in older adults: a randomized controlled trial. *Sleep.* 2012;35:797-805.
- [57] Espie CA, MacMahon KM, Kelly HL, Broomfield NM, Douglas NJ, Engleman HM, et al. Randomized clinical effectiveness trial of nurse-administered small-group cognitive behavior therapy for persistent insomnia in general practice. *Sleep.* 2007;30:574-84.
- [58] Morin CM, Kowatch RA, Barry T, Walton E. Cognitive-behavior therapy for late-life insomnia. *J Consult Clin Psychol.* 1993;61:137-46.
- [59] Miller CB, Kyle SD, Melehan KL, Bartlett DJ. Methodology for the Assessment of Sleep. In: Feldner MT, editor. *Sleep and Affect.* San Diego: Academic Press; 2015. p. 65-90.
- [60] *Kyle SD, Miller CB, Rogers Z, Siriwardena AN, Macmahon KM, Espie CA. Sleep restriction therapy for insomnia is associated with reduced objective total sleep time, increased daytime somnolence, and objectively impaired vigilance: implications for the clinical management of insomnia disorder. *Sleep.* 2014;37:229-37.
- [61] Lund HG, Rybarczyk BD, Perrin PB, Leszczyszyn D, Stepanski E. The discrepancy between subjective and objective measures of sleep in older adults receiving CBT for comorbid insomnia. *J Clin Psychol.* 2013;69:1108-20.
- [62] *Maurer LF, Espie CA, Kyle SD. How does sleep restriction therapy for insomnia work? A systematic review of mechanistic evidence and the introduction of the Triple-R model. *Sleep Med Rev.* 2018;42:127-38.

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

[63] Buysse DJ, Germain A, Hall M, Monk TH, Nofzinger EA. A Neurobiological Model of Insomnia. *Drug Discov Today Dis Models*. 2011;8:129-37.

[64] Zammit GK, McNabb LJ, Caron J, Amato DA, Roth T. Efficacy and safety of eszopiclone across 6-weeks of treatment for primary insomnia. *Curr Med Res Opin*. 2004;20:1979-91.

[65] McCall WV, Erman M, Krystal AD, Rosenberg R, Scharf M, Zammit GK, et al. A polysomnography study of eszopiclone in elderly patients with insomnia. *Curr Med Res Opin*. 2006;22:1633-42.

[66] Cronlein T, Wetter TC, Rupprecht R, Spiegelhalder K. Cognitive behavioral treatment for insomnia is equally effective in insomnia patients with objective short and normal sleep duration. *Sleep Med*. 2018, in press. doi: 10.1016/j.sleep.2018.10.038.

[67] Martin JL, Song Y, Hughes J, Jouldjian S, Dzierzewski JM, Fung CH, et al. A four-session sleep intervention program improves sleep for older adult day health care participants: results of a randomized controlled trial. *Sleep*. 2017;40.

[68] Schiller H, Söderström M, Lekander M, Rajaleid K, Kecklund G. A randomized controlled intervention of workplace-based group cognitive behavioral therapy for insomnia. *Int Arch Occup Environ Health*. 2018;91:413-24.

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON
OBJECTIVE SLEEP

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

Table 1

Randomised controlled trials of cognitive behavioural therapy for insomnia which report post treatment changes in objective night time sleep

Author (year) <i>Location</i>	Measure	Sample size (f), mean age (SD) Recruitment population	Group Allocation n N: randomised (complete d)	Diagnosis/ Definition of Insomnia <i>(determined by)</i> Mean duration (SD)	CBT-I components	Hypnotic medication status	Therapy Delivery Therapist Qualifications	Number of nights' data at each assessment <i>(Time points)</i>
Morin et al. (1993) [58] <i>USA</i>	PSG	N=24 (17 f) 67.1yrs (5.3) Older adult (60+) community sample	CBT-I: 12 WLC/ delayed treatment: 12	Persistent Psychophysiological Insomnia - ICSD WASO > 30min, ≥ 3n/wk, ≥ 6 month + daytime imp. <i>(Clinical interview and sleep log at baseline).</i> 13yrs (14.1)	PE, SRT, SCT, CR	Medication free	8 sessions; face-to-face group (n = 4-6); weekly; 90 min. clinical psychologist	Average of 2 nights <i>(Pre-post)</i>

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

Eding er et al. (2007)[54] <u>USA</u>	Actigr aphy f) 55.4yrs (9.7) Middle age and older adult communit y sample	N= 86 (43 (9) 55.4yrs (9.7) Middle age and older adult communit y sample	WL:11 (9) CBT-I (No. of Sessions) \ (1): 16 (16) (2): 18 (17) (4): 24 (21) (8): 17 (15)	Primary Insomnia (DSM- IV) WASO > 60 minutes, > 6 months with onset after age 10, ≥1 sleep disruptive practice e.g. napping <u>(Clinical</u> <u>interview and</u> <u>sleep log at</u> <u>baseline).</u>	PE, SCT, SRT	Medic ation free	1 (week 1), 2 (week 1 & 5), 4 (week 1, 3, 5 & 7) & 8 sessions; face- to-face individual; first session 45-60mins, follow up sessions 15-30mins. Clinical psychologists	2 weeks <u>(Pre-post,</u> <u>3 months,</u> <u>6 months)</u>
---	--	--	--	---	--------------------	------------------------	---	--

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

Espie et al. (2007)[⁵⁷] <i>Scotla nd, Unite d Kingd om</i>	Actigr aphy	N = 201 (137 f) 54yrs Patients attending General Medical Practice	CBT-I: 107(107) TAU: 94(94)	Insomnia (ICSD- R/ DSM-IV) SOL $\geq$ 30 min and/or WASO $\geq$ 30 min, $\geq$ 3 nights per week, $\geq$ 6 months and negative impact. <i>(Clinical interview and sleep log at baseline)</i> CBT: 11.6yrs (9.79) TAU: 10.6yrs (12.2)	PE, SH, SCT, SRT, CT	Medic ation allowe d	5 sessions; face-to-face group (n=4-6) weekly, 1 hr. Trained and supervised health visitors	2 weeks <i>(Pre- post)</i>
Eding er et al. (2009)[⁵⁰] <i>USA</i>	Actigr aphy	N = 81 (11 f) 54.2yrs (13.7) Veteran outpatient s with a history of sleep disturbanc e	CBT-I CMI: 21(20) CBT-I PI: 20(16) SH CMI: 20(15) SH PI: 20(18)	Insomnia Disorder (RDC) and/or DSM-IV- TR Primary Insomnia SOL + WASO $\geq$ 60 mins per night <i>(sleep log at baseline and diagnostic interview)</i> 10.0yrs (11.0)	PE, SCT, SRT	Medic ation allowe d	4 sessions; bi- weekly; individual; 30-60 mins. Clinical psychologists	2 weeks <i>(Pre- post, 6 months)</i>

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

Buyss e et al. (2011)[49] <u>USA</u>	PSG and Actigr aphy	N = 82 (54 f) 71.7yrs Older adult communit y sample and 1 primary care clinic.	BBT-I: 42(39) IC: 40(40)	Primary insomnia (DSM- IV-TR) minus comorbidity criteria and general insomnia (ICSD-2) Sleep complaint ≥ 1 month + daytime impairment. <u>(Self-report questionnaire and clinical interview).</u>	PE, SRT, SCT	Medic ation allowe d	4 contacts; weekly; 2 individual face-to-face (initial 45-60mins, 30min f/up at 2 wks.) 2 20min telephone calls (wk 1 and 3). Masters level nurse practitioner	PSG: Average of 2 nights Actigraph y: 2 weeks <u>(Pre-post)</u>
Epstei n et al. (2012)[56] <u>USA</u>	Actigr aphy	N=179 (115 f) 68.9yrs Older adult (55+) communit y sample	SCT: 44 (38) SRT: 44 (39) MCI: 41 (39) WLC: 50 (38)	Chronic primary insomnia SOL ≥45min for ≥3nights/wk, ≥6 months + daytime impairment. <u>(Clinical interview and sleep log at baseline).</u> SCT: 8.23yrs SRT: 9.82yrs MCI: 9.94yrs WLC: 11.7yrs	PE, SH, SCT, SRT	Medic ation free	6 sessions; wk 1-4 group (n =4-6), week 5-6 telephone; weekly; 60-120 min groups, 15min telephone. Mental health clinical nurse specialist (Masters Level)	2 weeks <u>(Pre-post, 3months, 1yr)</u>

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

Harris et al. (2012)] ^[32] <u>Austr</u> <u>alia</u>	Actigr aphy	N = 79 (56 f) 40.9yrs (13.8) Communit y sample	ISR: 19(16) SCT: 20(17) SCT + ISR: 20(20) SH: 20(18)	Sleep onset insomnia (+/- sleep maintenance difficulties) SOL > 30min, ≥3 nights/wk, ≥6months + daytime impairment. <u>(Clinical</u> <u>interview and</u> <u>sleep log at</u> <u>baseline).</u> 16yrs (13.6)	SCT, SH, RP ation free Clinical psychologist	Medic ation free Clinical psychologist	5 sessions; Individual face-to-face; weekly, 30min. Clinical psychologist	2 weeks <u>(Pre-post,</u> <u>2 months,</u> <u>6months)</u>
Irwin et al. (2014)] ^[33] <u>USA</u>	PSG	N=123 (88 f) 65.7yrs Older adult (55+) communit y sample	CBT-I: 50(48) TCC: 48(40) SS: 25(24)	Primary insomnia (DSM- IV-TR) and general insomnia (ICSD-2) Difficulty initiating/ maintaining sleep ≥1 month. + Daytime impairment. <u>(Clinical</u> <u>interview).</u>	PE, CR, SCT, SRT, REL, RP, mood intervent ion	Medic ation free Clinical psychologist	16 sessions: face-to-face group (n=7-10) 120min class weekly, 4months Clinical psychologist supervised by PhD specialist	Average of 2 nights after 1 night adaptatio n <u>(Pre-post)</u>

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

Lovat o et al. (2014) ^[34] <i>Austr alia</i>	Actigr aphy	N=188 (63 f) 63.76 yrs (6.45) Older adult communit y sample	CBT-I: 86 (78) WLC: 32(31)	Sleep maintenance insomnia WASO <30min, for >3night/wk, ≥6mo + daytime impairment: <i>(Clinical interview and sleep log at baseline)</i> Mean duration not reported (73% >5yrs)	SRT, CT, PE, SH, RP	Medic ation free	4 sessions, weekly; group (n=4-5); 60 mins. Five trainee psychologists	1 week <i>(Pre-post, 3months)</i>
Taylo r et al. (2014) ^[35] <i>USA</i>	Actigr aphy	N = 34 (14 f) 19.71yrs (2.10) College students (age 18-24yrs) from University of North Texas	CBT-I: 17(16) WLC: 17(13)	Insomnia (DSM-5) SOL or WASO ≥30mins, 3 nights/wk, ≥3mo + daytime impairment: <i>(Clinical interview and sleep log at baseline)</i>	SCT, SRT, SH, REL, CR	Medic ation free	6 sessions; face-to-face individual; weekly Three doctoral level graduate students	1 week <i>(Pre-post)</i>

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

Alessi et al. (2016) ^[36] <u>USA</u>	Actigraphy	N=159 (5 f) 72.2 yrs (7.7) Veteran (age 60+) community sample	CBT-I total: 106(97) (Individual al: 54 Group: 52) PE: 53 (53)	Chronic Insomnia Disorder (ICSD-2) ≥3months <u>(Postal survey:</u> <u>ICSD-2 insomnia</u> <u>disorder)</u> Mean duration not reported (90.6% ≥12months)	SH, SCT, SRT, CT	Medication allowe d	5 sessions; face-to-face individual or group (n=3-5); weekly; 60 min Masters level Sleep-coaches	1 week <u>(Pre-post,</u> <u>6months,</u> <u>12months</u> <u>)</u>
--	------------	---	--	--	------------------------	---------------------------	--	---

Lovato et al. (2016) ^[37] <u>Australia</u>	Actigraphy	N=91 (48 f) 63.34yrs (6.41) Community “young” older adults (65-74) targeted but no age exclusion criteria used	CBT-I: 63(63) (Sleep duration: Short: 30; Long: 32) WLC: 28(28) (Short: 9; Long: 19)	Sleep maintenance insomnia WASO <30min, for >3night/wk, ≥6months + daytime impairment. <u>Clinical</u> <u>interview</u> Not reported (76.92% <5yrs)	SRT, CT, PE	Medication free	4 sessions, weekly; group (n=4-5); 60 mins.	1 week <u>(Pre-post,</u> <u>3months)</u>
--	------------	--	---	--	----------------	--------------------	--	--

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

Taylor et al. (2017) [38]	Actigraphy	N=100 (17 f) 32.73yrs (7.73)	ICBT-I: 34 (27) CBT-I: 33 (30)	Chronic Insomnia (DSM-5)	SCT, SRT, SH, REL, CR	Medication allowed (if stable use for 4wk)	6 sessions; weekly; 60mins; individual online or face-to-face Clinical psychologist, postdoctoral fellows, social worker.	1 week (Pre-post, 6months)
---------------------------	------------	---------------------------------	-----------------------------------	--------------------------	-----------------------	--	---	-------------------------------

Abbreviations: BBTI = Brief Behavioural Treatment for Insomnia; CBT-I = Cognitive Behavioural Therapy for Insomnia; CC = control condition; CMI = comorbid insomnia; CR = Cognitive Restructuring; CT = Cognitive Therapy; DSM(IV-TR)= Diagnostic and Statistical Manual of Mental Disorders (IV-Text Revision); f. = Female; f/up = follow up; IC = information control; ICBT-I = Internet Cognitive Behavioural Therapy for Insomnia; ICSD = International Classification of Sleep Disorders; ISR = Intensive Sleep Retraining; ITT = Intention To Treat; MCI = multi-component intervention; Min. = minutes; Mo = Month; n = Nights; PE = Psychoeducation; PI: primary insomnia; PSG = Polysomnography; PT: placebo therapy (psychological); RDC = research diagnostic criteria; REL = Relaxation; RT = relaxation therapy; RP = Relapse Prevention; SC = Sleep Compression; SCT = Stimulus Control Therapy; SD = Standard Deviation; SE = Sleep Efficiency; SH = Sleep Hygiene; SOL = Sleep Onset Latency; SRT = Sleep Restriction Therapy; SS = Sleep Seminar; TCC = Tai Chi Chih; TST = Total Sleep Time; TWT = total wake time at night; WASO = Wake After Sleep Onset; Wk(s)=Week(s) Yrs = Year

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

Table 2

Effect sizes for post-treatment sleep differences between patients assigned to CBT-I versus control.

Outcome	Polysomnography				Actigraphy				Sleep Diary			
	N _c	g (95%CI)	I ² (95% CI)	p	N _c	g (95%CI)	I ² (95% CI)	p	N _c	g (95%CI)	I ² (95% CI)	p
Sleep												
Onset	4	-0.07 (-0.36 to 0.22)	0 (0-86)	0.641	11	-0.28 (-0.51 to -0.05)	52 (5-86)	0.018	16	-0.50 (-0.70 to -0.30)	56 (17-84)	<0.001
Latency (SOL)												
Wake												
After												
Sleep	4	-0.10 (-0.39 to 0.19)	0 (0-97)	0.509	12	-0.14 (-0.33 to 0.05)	38 (0-69)	0.138	16	-0.75 (-0.96 to -0.55)	56 (18-81)	<0.001
Onset (WASO)												
Sleep												
Efficiency	5	0.33 (-0.14 to 0.80)	63 (0-97)	0.169	14	0.14 (-0.04 to 0.31)	42 (0-69)	0.137	17	0.79 (0.57 to 1.02)	66 (36-85)	<0.001
(SE)												
Total												
Sleep	5	0.04 (-0.37 to 0.45)	53 (0-97)	0.848	12	-0.51 (-0.75 to -0.26)	54 (9-84)	<0.001	16	0.06 (-0.12 to 0.23)	41 (0-72)	0.537
Time (TST)												

95% CI = 95% confidence interval; N_c = number of comparisons

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON
OBJECTIVE SLEEP

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON
OBJECTIVE SLEEP
Supplementary information

Table S1

*Assessment of risk of bias using the Cochrane Collaboration's tool for assessing risk of bias
in randomised trials*

Study (author, year)	Random Sequence Generation	Allocation concealment	Blinding of participants and personnel	Blinding of outcome assessment	Incomplete outcome data	Selective reporting
Morin et al (1993) ^[58]	Unclear	Unclear	Unclear	Unclear	Unclear	Low
Friedman et al. (2000) ^[55]	Low	Unclear	High	Unclear	Low	Low
Edinger et al. (2001) ^[48]	Low	Unclear	Low	Low	Low	Low
Edinger et al. (2007) ^[54]	Low	Unclear	High	Unclear	Low	Low
Espie et al. (2007) ^[57]	Low	Low	High	Unclear	Low	Low
Edinger et al. (2009) ^[50]	Low	Unclear	Low	Unclear	Low	Low
Buysse et al. (2011) ^[49]	Low	Unclear	Unclear	Unclear	Low	Low
Epstein et al. (2012) ^[56]	Low	Unclear	High	Unclear	Low	Low
Harris et al. (2012) ^[32]	Low	Unclear	Unclear	Unclear	Low	Low
Irwin et al. (2014) ^[33]	Low	Low	Unclear	Low	Low	Low
Lovato et al. (2014) ^[34]	Low	Low	High	Unclear	Low	Low
Taylor et al. (2014) ^[35]	Low	Low	High	High	Low	Low
Alessi et al. (2016) ^[36]	Low	Low	Low	Low	Low	High
Lovato et al. (2016) ^[37]	Low	Unclear	Unclear	Unclear	Low	Low
Taylor et al. (2017) ^[38]	Low	Unclear	Unclear	Unclear	Low	Low

Key: High = High risk of bias; Low = Low risk of bias; Unclear = Insufficient information to allow judgement

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

Table S2

*Sensitivity analyses: Fixed effects modelling for outcomes showing no evidence of
heterogeneity*

Outcome	Polysomnography			Actigraphy			Sleep Diary		
	N _c	g (95%CI)	p	N _c	g (95%CI)	p	N _c	g (95%CI)	p
Sleep Onset	4	-0.07 (-0.36 to 0.22)	0.6	-	-	-	-	-	-
Latency(SOL)	4	-0.10 (-0.39 to 0.19)	0.5	1	-0.12 (-0.26 to 0.02)	0.0	-	-	-
Wake After Sleep Onset (WASO)	4	-	-	2	0.11 (-0.02 to 0.24)	0.0	-	-	-
Sleep Efficiency (SE)	-	-	-	4	-	-	-	-	-
Total Sleep Time (TST)	5	0.07 (-0.19 to 0.33)	0.6	-	-	-	1	0.05 (-0.08 to 0.18)	0.4
							6		64

95% CI = 95% confidence interval; N_c = number of comparisons

Table S3

*Exploratory meta-analysis of PSG and actigraphy outcomes in CBT versus control for studies
with 3-month follow-up data*

Polysomnography	Actigraphy
-----------------	------------

RUNNING HEAD: EFFECTS OF COGNITIVE BEHAVIOURAL THERAPY ON OBJECTIVE SLEEP

Outcome	N _c	g (95%CI)	I ² (95%CI)	p	N _c	g (95%CI)	I ² (95%CI)	p
Sleep Onset Latency (SOL)	1	0.31 (-1.02 to 1.64)	NA	0.64 7	3	0.33 (-0.11 to 0.78)	20 (0-97)	0.14 3
Wake After Sleep Onset (WASO)	1	0.01 (-1.32 to 1.33)	NA	0.99 2	4	-0.23 (-0.51 to 0.05)	0 (0-94)	0.10 6
Sleep Efficiency (SE)	1	-0.45 (-1.79 to 0.89)	NA	0.51 1	4	-0.07 (-0.35 to 0.22)	0 (0-57)	0.64 8
Total Sleep Time (TST)	1	0.00 (-1.33 to 1.33)	NA	0.99 9	4	-0.56 (-0.85 to -0.28)	0 (0-89)	<0.0 01

95% CI = 95% confidence interval; N_c = number of comparisons