


Accepted Manuscript

Cross-sectional study of the prevalence, causes and management of hospital-onset diarrhoea



Damian Mawer, Fiona Byrne, Sarah Drake, Claire Brown, Alison Prescott, Ben Warne, Rachel Bousfield, Jordan P. Skittrall, Isobel Ramsay, Donald Somasunderam, Moira Bevan, Julie Coslett, Jyothi Rao, Philip Stanley, Adrian Kennedy, Rosemarie Dobson, Sandra Long, Tobi Obisanya, Taher Esmailji, Christina Petridou, Kordo Saeed, Kelly Brechany, Karen Davis-Blue, Helen O'Horan, Bruce Wake, Jessica Martin, Jennifer Featherstone, Charlotte Hall, Joanna Allen, Greta Johnson, Cheryl Hornigold, Nurul Amir, Kathleen Henderson, Catriona McClements, Ignatius Liew, Ashutosh Deshpande, Elen Vink, Debbie Trigg, Jane Guilfoyle, Matthew Scarborough, Claire Scarborough, TH Nicholas Wong, Timothy Walker, Nicola Fawcett, Gayti Morris, Kim Tomlin, Caroline Grix, Emma O'Cofaigh, David McCaffrey, Michael Cooper, Kim Corbett, Kathryn French, Susan Harper, Claire Hayward, Matthew Reid, Vanessa Whatley, Jodie Winfield, Selina Hoque, Lynn Kelly, Isobel King, Amy Bradley, Bernie McCullagh, Carleen Hibberd, Monica Merron, Caroline McCabe, Samantha Horridge, Johnathan Taylor, Sharon Koo, Fadwa Elsanousi, Rosalind Saunders, Felicia Lim, Amy Bond, Sheldon Stone, Iain D. Milligan, Damien JF. Mack, Aaron Nagar, Robert M. West, Mark H. Wilcox, Andrew Kirby, Jonathan AT. Sandoe

PII: S0195-6701(19)30192-6

DOI: <https://doi.org/10.1016/j.jhin.2019.05.001>

Reference: YJHIN 5720

To appear in: *Journal of Hospital Infection*

Received Date: 6 April 2019

Accepted Date: 1 May 2019

Please cite this article as: Mawer D, Byrne F, Drake S, Brown C, Prescott A, Warne B, Bousfield R, Skittrall JP, Ramsay I, Somasunderam D, Bevan M, Coslett J, Rao J, Stanley P, Kennedy A, Dobson R, Long S, Obisanya T, Esmailji T, Petridou C, Saeed K, Brechany K, Davis-Blue K, O'Horan H, Wake B, Martin J, Featherstone J, Hall C, Allen J, Johnson G, Hornigold C, Amir N, Henderson K, McClements C, Liew I, Deshpande A, Vink E, Trigg D, Guilfoyle J, Scarborough M, Scarborough C, Wong TN, Walker T, Fawcett N, Morris G, Tomlin K, Grix C, O'Cofaigh E, McCaffrey D, Cooper M, Corbett K, French K,

Harper S, Hayward C, Reid M, Whatley V, Winfield J, Hoque S, Kelly L, King I, Bradley A, McCullagh B, Hibberd C, Merron M, McCabe C, Horridge S, Taylor J, Koo S, Elsanousi F, Saunders R, Lim F, Bond A, Stone S, Milligan ID, Mack DJ, Nagar A, West RM, Wilcox MH, Kirby A, Sandoe JA, Cross-sectional study of the prevalence, causes and management of hospital-onset diarrhoea, *Journal of Hospital Infection*, <https://doi.org/10.1016/j.jhin.2019.05.001>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Title: Cross-sectional study of the prevalence, causes and management
of hospital-onset diarrhoea**

Authors: Damian Mawer, Fiona Byrne, Sarah Drake, Claire Brown, Alison Prescott, Ben Warne, Rachel Bousfield, Jordan P Skittrall, Isobel Ramsay, Donald Somasunderam, Moira Bevan, Julie Coslett, Jyothi Rao, Philip Stanley, Adrian Kennedy, Rosemarie Dobson, Sandra Long, Tobi Obisanya, Taher Esmailji, Christina Petridou, Kordo Saeed, Kelly Brechany, Karen Davis-Blue, Helen O’Horan, Bruce Wake, Jessica Martin, Jennifer Featherstone, Charlotte Hall, Joanna Allen, Greta Johnson, Cheryll Hornigold, Nurul Amir, Kathleen Henderson, Catriona McClements, Ignatius Liew, Ashutosh Deshpande, Elen Vink, Debbie Trigg, Jane Guilfoyle, Matthew Scarborough, Claire Scarborough, TH Nicholas Wong, Timothy Walker, Nicola Fawcett, Gayti Morris, Kim Tomlin, Caroline Grix, Emma O’Cofaigh, David McCaffrey, Michael Cooper, Kim Corbett, Kathryn French, Susan Harper, Claire Hayward, Matthew Reid, Vanessa Whatley, Jodie Winfield, Selina Hoque, Lynn Kelly, Isobel King, Amy Bradley, Bernie McCullagh, Carleen Hibberd, Monica Merron, Caroline McCabe, Samantha Horridge, Johnathan Taylor, Sharon Koo, Fadwa Elsanousi, Rosalind Saunders, Felicia Lim, Amy Bond, Sheldon Stone, Iain D Milligan, Damien JF Mack, Aaron Nagar, Robert M West, Mark H Wilcox, Andrew Kirby, Jonathan AT Sandoe

Authors’ affiliations:

Damian Mawer

Trust grade microbiologist, Department of Microbiology, Leeds Teaching Hospitals NHS Trust, Leeds, LS9 7TF, UK.

Fiona Byrne

Specialty registrar, Department of Microbiology, Leeds Teaching Hospitals NHS Trust, Leeds, LS9 7TF, UK.

Sarah Drake

28 Specialty registrar, Department of Microbiology, Leeds Teaching Hospitals NHS Trust, Leeds,
29 LS9 7TF, UK.

30 Claire Brown

31 Research nurse, Department of Microbiology, Leeds Teaching Hospitals NHS Trust, Leeds,
32 LS9 7TF, UK.

33 Alison Prescott

34 Specialty registrar, Department of Microbiology, Leeds Teaching Hospitals NHS Trust, Leeds,
35 LS9 7TF, UK.

36 Ben Warne

37 Specialty registrar, Department of Infectious Diseases, Cambridge University Hospitals NHS
38 Foundation Trust, Cambridge, CB2 0QQ, UK.

39 Rachel Bousfield

40 Specialty registrar, Department of Infectious Diseases, Cambridge University Hospitals NHS
41 Foundation Trust, Cambridge, CB2 0QQ, UK.

42 Jordan P Skittrall

43 NIHR Academic Clinical Fellow in Infectious Diseases, Royal Papworth Hospital NHS
44 Foundation Trust, Papworth Everard, Cambridge, CB23 3RE, UK.

45 Isobel Ramsay

46 Specialty registrar, Department of Infectious Diseases, Cambridge University Hospitals NHS
47 Foundation Trust, Cambridge, CB2 0QQ, UK.

48 Donald Somasunderam

49 Specialty registrar, Department of Infectious Diseases, Cambridge University Hospitals NHS
50 Foundation Trust, Cambridge, CB2 0QQ, UK.

51 Moira Bevan

52 Lead nurse for Infection Prevention, Department of Infection Prevention, Royal Gwent
53 Hospital, Newport, NP20 2UB, UK.

54 Julie Coslett

55 Senior Infection Prevention nurse, Department of Infection Prevention, Royal Gwent
56 Hospital, Newport, NP20 2UB, UK.

57 Jyothi Rao

58 Consultant microbiologist, Department of Microbiology, Barnsley Hospital NHS Foundation
59 Trust, Barnsley, S75 2EP, UK.

60 Philip Stanley

61 Director of Infection Prevention and Control, Bradford Teaching Hospitals NHS Foundation
62 Trust, Bradford, BD9 6RJ, UK.

63 Adrian Kennedy

64 Specialty registrar, Department of Medicine, Bradford Teaching Hospitals NHS Foundation
65 Trust, Bradford, BD9 6RJ, UK.

66 Rosemarie Dobson

67 Lead infection control nurse, Department of Infection Prevention & Control, Bradford
68 Teaching Hospitals NHS Foundation Trust, Bradford, BD9 6RJ, UK.

69 Sandra Long

70 Consultant microbiologist, Department of Microbiology, East Lancashire Hospitals NHS
71 Trust, Blackburn, BB2 3HH, UK.

72 Tobi Obisanya

73 Foundation Year 1 doctor, Department of Microbiology, East Lancashire Hospitals NHS
74 Trust, Blackburn, BB2 3HH, UK.

75 Taher Esmailji

76 Foundation year 2 doctor, East Lancashire Hospitals NHS Trust, Blackburn, BB2 3HH, UK.

77 Christina Petridou

78 Specialty registrar, Department of Microbiology, Hampshire Hospitals NHS Foundation
79 Trust, Winchester, SO22 5DG, UK.

80 Kordo Saeed

81 Consultant microbiologist, Department of Microbiology, Hampshire Hospitals NHS
82 Foundation Trust, Winchester, SO22 5DG, UK.

83 Karen Davis-Blue

84 Acting lead infection prevention nurse, Department of Infection Prevention and Control,
85 Hampshire Hospitals NHS Foundation Trust, Winchester, SO22 5DG, UK.

86 Kelly Brechany

87 Infection control nurse, Department of Infection Prevention and Control, Hampshire
88 Hospitals NHS Foundation Trust, Winchester, SO22 5DG, UK.

89 Helen O'Horan

90 Vascular access nurse, Department of Infection Prevention and Control, Hampshire
91 Hospitals NHS Foundation Trust, Winchester, SO22 5DG, UK.

92 Bruce Wake

93 Infection surveillance coordinator, Department of Infection Prevention and Control,
94 Hampshire Hospitals NHS Foundation Trust, Winchester, SO22 5DG, UK.

95 Jessica Martin

96 Specialty registrar, Department of Microbiology, Harrogate and District NHS Foundation
97 Trust, Harrogate, HG2 7SX, UK.

98 Jennifer Featherstone

99 Lead Infection Control nurse, Department of Infection Prevention & Control, Harrogate and
100 District NHS Foundation Trust, Harrogate, HG2 7SX, UK.

101 Charlotte Hall

102 Specialty registrar, Department of Infectious Diseases, Hull and East Yorkshire Hospitals NHS
103 Trust, Hull, HU3 2JZ, UK.

- 104 Joanna Allen
- 105 Specialty registrar, Department of Infectious Diseases, Hull and East Yorkshire Hospitals NHS
- 106 Trust, Hull, HU3 2JZ, UK.
- 107 Greta Johnson
- 108 Lead Nurse, Department of Infection Prevention & Control, Hull & East Hospitals NHS Trust,
- 109 Hull, HU3 2JZ, UK
- 110 Cheryl Hornigold
- 111 Infection prevention and control nurse, Department of Infection Prevention & Control, Hull
- 112 and East Yorkshire Hospitals NHS Trust, Hull, HU3 2JZ, UK.
- 113 Nurul Amir
- 114 Consultant microbiologist, Department of Microbiology, Mid Yorkshire Hospitals NHS Trust,
- 115 Wakefield, WF1 4DG, UK.
- 116 Kathleen Henderson
- 117 Core medical trainee, Inverclyde Royal Hospital, Greenock, PA16 0XN, UK.
- 118 Catriona McClements
- 119 Foundation Year 2 doctor, Inverclyde Royal Hospital, Greenock, PA16 0XN, UK.
- 120 Ignatius Liew
- 121 Foundation Year 1 doctor, Inverclyde Royal Hospital, Greenock, PA16 0XN, UK.
- 122 Ashutosh Deshpande
- 123 Consultant microbiologist, Department of Microbiology, Inverclyde Royal Hospital,
- 124 Greenock, PA16 0XN, UK.
- 125 Elen Vink
- 126 Specialty registrar, Department of Microbiology, Royal Infirmary of Edinburgh, Edinburgh,
- 127 EH16 4SA, UK.
- 128 Debbie Trigg

129 Matron, Department of Infection Prevention & Control, Nottingham University Hospitals
130 NHS Trust, Nottingham, NG7 2UH, UK.

131 Jane Guilfoyle

132 Infection control nurse, Department of Infection Prevention & Control, Nottingham
133 University Hospitals NHS Trust, Nottingham, NG7 2UH, UK.

134 Matthew Scarborough

135 Consultant in infectious diseases & microbiology, Department of Infectious Diseases, Oxford
136 University Hospitals NHS Trust, Oxford, OX3 9DU, UK.

137 Claire Scarborough

138 Clinical research fellow, Nuffield Department of Medicine, University of Oxford, OX3 7FZ,
139 UK.

140 TH Nicholas Wong

141 Specialty registrar, Department of Infectious Diseases, Oxford University Hospitals NHS
142 Trust, Oxford, OX3 9DU, UK.

143 Timothy Walker

144 Specialty registrar, Department of Infectious Diseases, Oxford University Hospitals NHS
145 Trust, Oxford, OX3 9DU, UK.

146 Nicola Fawcett

147 Specialty registrar, Department of Medicine, Oxford University Hospitals NHS Trust, Oxford,
148 OX3 9DU, UK.

149 Gayti Islam

150 Specialty registrar, Department of Microbiology, Sheffield Teaching Hospitals NHS
151 Foundation Trust, Sheffield, S10 2JF.

152 Kim Tomlin

153 Infection control nurse, Department of Infection Prevention & Control, Sheffield Teaching
154 Hospitals NHS Foundation Trust, Sheffield, S10 2JF.

155 Caroline Grix
156 Infection control nurse, Department of Infection Prevention & Control, Sheffield Teaching
157 Hospitals NHS Foundation Trust, Sheffield, S10 2JF.

158 Emma O’Cofaigh
159 Consultant in infectious diseases, Department of Medicine, Friarage Hospital, South Tees
160 Hospital NHS Foundation Trust, Northallerton, DL6 1JG, UK.

161 David McCaffrey
162 Lead infection control nurse, Department of Infection Prevention & Control, James Cook
163 University Hospital, South Tees Hospital NHS Foundation Trust, Middlesbrough, TS4 3BW.
164 UK.

165 Michael Cooper
166 Consultant microbiologist, Department of Microbiology, The Royal Wolverhampton NHS
167 Trust, Wolverhampton, WV10 0QP, UK.

168 Kim Corbett
169 Infection control nurse, Department of Infection Prevention & Control, The Royal
170 Wolverhampton NHS Trust, Wolverhampton, WV10 0QP, UK.

171 Kathryn French
172 Specialty registrar, Department of Microbiology, The Royal Wolverhampton NHS Trust,
173 Wolverhampton, WV10 0QP, UK.

174 Susan Harper
175 Senior infection control nurse, Department of Infection Prevention & Control, The Royal
176 Wolverhampton NHS Trust, Wolverhampton, WV10 0QP, UK.

177 Claire Hayward
178 Infection control nurse, Department of Infection Prevention & Control, The Royal
179 Wolverhampton NHS Trust, Wolverhampton, WV10 0QP, UK.

180 Matthew Reid

181 Infection prevention nurse manager, Department of Infection Prevention & Control, The
182 Royal Wolverhampton NHS Trust, Wolverhampton, WV10 0QP, UK.

183 Vanessa Whatley

184 Head of nursing, Corporate Support Services, The Royal Wolverhampton NHS Trust,
185 Wolverhampton, WV10 0QP, UK.

186 Jodie Winfield

187 Infection prevention nurse manager, Department of Infection Prevention & Control, The
188 Royal Wolverhampton NHS Trust, Wolverhampton, WV10 0QP, UK.

189 Selina Hoque

190 Director of Infection Prevention and Control, Department of Microbiology, Torbay and
191 South Devon Healthcare NHS Foundation Trust, Torquay, TQ2 7AA, UK.

192 Lynn Kelly

193 Lead infection control nurse, Department of Infection Prevention & Control, Torbay and
194 South Devon Healthcare NHS Foundation Trust, Torquay, TQ2 7AA, UK.

195 Isobel King

196 Infection prevention and control lead, Department of Infection Prevention & Control, Ulster
197 Hospital, South Eastern Health and Social Care Trust, Belfast, BT16 1RH, UK.

198 Amy Bradley

199 Senior infection prevention and control nurse, Department of Infection Prevention &
200 Control, Ulster Hospital, South Eastern Health and Social Care Trust, Belfast, BT16 1RH, UK.

201 Bernie McCullagh

202 Lead antimicrobial pharmacist, Pharmacy Department, Ulster Hospital, South Eastern Health
203 and Social Care Trust, Belfast, BT16 1RH, UK.

204 Carleen Hibberd

205 Antimicrobial pharmacist, Pharmacy Department, Ulster Hospital, South Eastern Health and
206 Social Care Trust, Belfast, BT16 1RH, UK.

207 Monica Merron

208 Infection prevention and control lead, Department of Infection Prevention & Control, Ulster
209 Hospital, South Eastern Health and Social Care Trust, Belfast, BT16 1RH, UK.

210 Caroline McCabe

211 Infection prevention and control nurse, Department of Infection Prevention and Control,
212 Ulster Hospital, South Eastern Health and Social Care Trust, Belfast, BT16 1RH, UK.

213 Samantha Horridge

214 Clinical scientist, Department of Microbiology, University Hospital Coventry, University
215 Hospitals of Coventry and Warwickshire, Warwick, CV2 2DX, UK.

216 Johnathan Taylor

217 Clinical scientist, Department of Virology and Molecular Pathology, University Hospital
218 Coventry, University Hospitals of Coventry and Warwickshire, Warwick, CV2 2DX, UK.

219 Sharon Koo

220 Specialty registrar, Department of Microbiology, University Hospitals of Leicester NHS Trust,
221 Leicester, LE1 5WW, UK.

222 Fadwa Elsanousi

223 Specialty registrar, Department of Microbiology, University Hospitals of Leicester NHS Trust,
224 Leicester, LE1 5WW, UK.

225 Rosalind Saunders

226 Specialty registrar, Department of Microbiology, University Hospitals of Leicester NHS Trust,
227 Leicester, LE1 5WW, UK.

228 Felicia Lim

229 Specialty registrar, Department of Microbiology, University Hospitals of Leicester NHS Trust,
230 Leicester, LE1 5WW, UK.

231 Amy Bond

- 232 Specialty registrar, Department of Microbiology, York Teaching Hospital NHS Foundation
233 Trust, York, YO31 8HE, UK.
- 234 Sheldon Stone
- 235 Senior lecturer, Stroke physician and consultant physician for older people, Royal Free
236 Campus, University College Medical School, London, NW3 2QG, UK.
- 237 Iain D Milligan
- 238 Specialty registrar, Department of Microbiology, Royal Free Hospital, University College
239 London Hospitals NHS Foundation Trust, London, NW3 2QG, UK.
- 240 Damien Mack
- 241 Consultant microbiologist, Department of Microbiology, Royal Free Hospital, Royal Free
242 London NHS Foundation Trust, London, NW3 2QG, UK.
- 243 Aaron Nagar,
- 244 Consultant microbiologist, Department of Microbiology, Antrim Area Hospital, Northern
245 Health and Social Care Trust, Bush Road, Antrim, BT41 2RL, UK.
- 246 Robert M West
- 247 Professor of biostatistics, Leeds Institute of Health Sciences, University of Leeds, Leeds, LS2
248 9JT, UK.
- 249 Mark H Wilcox
- 250 Professor of medical microbiology, Leeds Institute of Biomedical and Clinical Sciences,
251 University of Leeds, Leeds, LS2 9JT, UK.
- 252 Andrew Kirby
- 253 Associate clinical professor in microbiology, Leeds Institute of Medical Research, University
254 of Leeds, Leeds, LS2 9JT, UK.
- 255 Jonathan AT Sandoe
- 256 Associate clinical professor in microbiology, Leeds Institute of Biomedical and Clinical
257 Sciences, University of Leeds, Leeds, LS2 9JT, UK.

Corresponding author contact details:

Damian Mawer. Consultant microbiologist, Department of Microbiology, York Hospital, Wigginton Road, York, YO31 8HE, UK. Telephone: +44 1904 726188; Fax: +44 1904 725991; E-mail: damian.mawer@nhs.net

Running Title:

Prevalence, causes and management of hospital-onset diarrhoea.

Summary

Background: The National Health Service in England advises hospitals collect data on hospital-onset diarrhoea (HOD). Contemporaneous data on HOD are lacking.

Aim: To investigate prevalence, aetiology and management of HOD on medical, surgical and elderly-care wards.

Methods: A cross-sectional study in a volunteer sample of UK hospitals, which collected data on one winter and one summer day in 2016. Patients admitted ≥ 72 hours were screened for HOD (definition: ≥ 2 episodes of Bristol Stool Type 5-7 the day before the study, with diarrhoea-onset > 48 hours after admission). Data on HOD aetiology and management were collected prospectively.

Findings: Data were collected on 141 wards in 32 hospitals (16 acute, 16 teaching). Point-prevalence of HOD was 4.5% (230/5142 patients; 95% CI 3.9-5.0%). Teaching hospital HOD prevalence (5.9%, 95% CI 5.1-6.9%) was twice that of acute hospitals (2.8%, 95% CI 2.1-3.5%; odds ratio 2.2, 95% CI 1.7-3.0). At least one potential cause was identified in 222/230 patients (97%): 107 (47%) had a relevant underlying condition, 125 (54%) were taking antimicrobials, and 195 (85%) other medication known to cause diarrhoea. 9/75 tested patients were *Clostridium difficile* toxin positive (4%). 80 (35%) patients had a documented medical assessment of the diarrhoea. Documentation of HOD in medical notes correlated

with testing for *C. difficile* (78% of those tested versus 38% not tested, $p < 0.001$). 144 (63%) patients were not isolated following diarrhoea onset.

Conclusion: HOD is a prevalent symptom affecting thousands of patients across the UK health system each day. Most patients have multiple potential causes of HOD, mainly iatrogenic, but only a third had medical assessment. Most were not tested for *C. difficile* and were not isolated.

Key words: Diarrhoea; nosocomial; hospital onset; hospital acquired; *Clostridium difficile*

Abbreviations:

BSC – Bristol stool chart

CDI – *Clostridium difficile* infection

HOD – hospital-onset diarrhoea

NHS – National Health Service

Introduction:

Guidelines from the National Health Service (NHS) in England encourage hospitals to collect data on the prevalence of hospital-onset diarrhoea (HOD), to facilitate detection of unexpected variations [1]. However, NHS hospitals generally do not collect such data and contemporaneous prevalence and risk factor information in developed countries are lacking.

HOD is known to affect patients and health services [2]. Causes such as *Clostridium difficile* infection (CDI) are associated with morbidity and mortality [3]. Management of HOD also requires staff time and hospital resources [3,4]. Single-centre cohort studies from the 1990s suggested it was common, affecting 22-32% of patients during admission, but were performed in settings with high CDI rates [5,6]. Whilst the prevalence was lower in more recent publications, they were performed in diverse ward/hospital settings using different

definitions of HOD, making results difficult to generalise [7-12]. Studies of the aetiology and management of HOD focus mainly on specialist areas (e.g. intensive care, oncology) [13,14]; there are few data from general medical, surgical and elderly-care settings, although they represent over half of adult hospital admissions in England [15]. To address these knowledge gaps and comply with NHS guidelines [1] we undertook a cross-sectional study on general medical, surgical and elderly-care wards across the UK to describe the prevalence of HOD, its potential causes and management.

Methods

Ethics

The NHS Health Research Authority advised the study constituted a service evaluation, not requiring written patient consent or formal review by a research ethics committee. Appropriate local approval was required at participating sites.

Design

Prospective, observational, cross sectional study. The protocol was developed by the UK's National Infection Trainee Collaborative for Audit and Research, following the STROBE statement (checklist in supplementary material) [16].

Setting

The study was advertised through the British Infection Association, Healthcare Infection Society and Infection Prevention Society in autumn 2015. All acute and teaching NHS hospital trusts/health boards (administrative units typically covering 1-3 hospitals) in the UK were eligible to participate. Participating sites, which were self-selected, chose one day in each of two periods (11-22 January and 6-17 June 2016) for data collection. Data were collected in winter and summer to allow for potential seasonal variation. Sites identified at least two wards for data collection from a pre-specified specialty list (supplementary material, Table S1).

Participants

On each data collection day all patients on study wards were eligible for inclusion, except those receiving end-of-life care. Patients specifically declining to be involved were excluded.

Definitions and variables

HOD was defined as ≥ 2 episodes of unformed stools (Bristol Stool Chart (BSC) type 5-7) in the day before data collection, with diarrhoea onset >48 hours after hospital admission [7,17]. In patients with a stoma, HOD was defined as an acute increase in daily faecal effluent above that normally expected (by the patient or clinical team). If known, an effluent volume of $>500\text{mL}/24\text{hr}$ for a colostomy or $>1000\text{mL}/24\text{hr}$ for an ileostomy was considered significant.

Variables assessed included: diarrhoea frequency and duration, previous HOD episodes during that admission and possible causes of HOD (underlying medical conditions associated with diarrhoea, gastrointestinal infections, antimicrobials and other medication associated with diarrhoea).

Common causes of HOD were identified by literature search (see supplementary material), though investigators could report any condition, medication or intervention they believed was causing diarrhoea. For an iatrogenic exposure to be considered a possible cause of HOD it had to fulfil these criteria: exposure had to precede diarrhoea onset with exposure >48 hours, or two or more doses of a new medication (with either the second of these, or a subsequent dose, given in the 24 hours before diarrhoea onset) and the diarrhoea resolving once exposure ceased (if known on the study day) [5]. For enemas and bowel preparation a single dose was considered sufficient. Earlier medication exposure was not studied. Results from microbiological testing of faecal samples submitted ≤ 72 hours after data collection were included.

Data were also collected on hospital characteristics: size (small (<500 beds), medium (500-1000 beds) and large (>1000 beds)), type, isolation capacity, testing protocols for faecal samples and infection control policies. Definitions from mandatory reporting were used for hospital type ("acute" being medium-sized, general hospitals providing services for local populations, and "teaching" larger hospitals, providing medical training and local general services, plus specialised regional services).

Data sources, collection and analysis

Standardised data collection forms were developed and piloted at three hospitals (supplementary material Appendices 1-3). Investigators received telephone-based training and guidance notes for data collection. On each study day patients admitted >72 hours were questioned to ascertain whether they had HOD, using the BSC to ensure correct identification of diarrhoea. Evidence for HOD was also obtained through questioning of ward staff and chart review. If more than one source indicated a patient had HOD, further data were collected from the patient, ward staff, medical charts/records and laboratory results systems. All possible causes of HOD were documented; where multiple causes were identified no attempt was made to rank their importance.

Hospitals, wards and patients with HOD were given unique identifiers at each participating site. Anonymised patient data were then uploaded on to a standardised database (Microsoft Excel) and verified locally. Each site submitted their database for analysis in Leeds.

Sample size

Based on published studies the point prevalence of HOD was estimated to be ~10% [5-12]. It was assumed there would be ~25 patients per ward and 5 wards per hospital. There was no information about clustering effects to be anticipated, therefore an intra-class correlation coefficient of 0.01 was used for patients within wards and wards within hospitals. Without any adjustment for clustering, to achieve a 95% confidence interval of $\pm 1\%$, assuming a binomial model, would need 3600 patients. The design effect for wards was 1.24 and that for hospitals was 1.04. The sample size required to achieve $\pm 1\%$ was therefore $1.24 \times 1.04 \times 3600 = 4643$ patients.

Statistical Analysis

The primary outcome measure was the point-prevalence of HOD. Secondary outcomes were hospital characteristics associated with HOD, the proportion of cases with a potential cause(s) and the proportion of cases tested for CDI. Standards for the evaluation of HOD, including testing for CDI, were taken from national guidance [1].

Appropriate summary statistics were reported for continuous (normally and non-normally distributed) and categorical variables. As HOD patients could potentially influence ward

prevalence, which in turn could affect hospital prevalence, a multilevel model was used with patients clustered within wards and wards within hospitals. Random intercepts for ward and hospital were included in a variance components binomial regression. Clustering of hospitals within NHS trusts/health boards was not considered, since most trusts/health boards had one hospital represented in the study and none had more than two.

To seek explanation of the variance at different levels, several plausible explanatory variables were considered in a three-level binomial regression model and then a parsimonious model was presented featuring only statistically significant terms with random intercepts retained for wards and for hospital, to account for clustering. Prevalence estimates were presented from these results, complete with confidence intervals.

Fixed effects terms were included for season and speciality and were to be included for ward and hospital characteristics as appropriate, unless little clustering was seen by ward or hospital. Analyses were performed using R version 3.3.2 employing the lme4 library version 1.1-12 [18,19].

Results

Settings & Recruitment

Thirty-two hospitals (16 acute, 16 teaching) at 25 NHS trusts/health boards participated (listed in supplementary material). One hundred and forty-one wards were included: 63 (45%) medical, 52 (37%) surgical and 26 (18%) elderly-care (Table S1). Data were collected from 116 wards in both rounds, 20 just in winter, and 5 in summer only. Patient recruitment and results are summarized in Figure 1.

Multilevel model output

The multilevel model revealed minimal contribution to clustering either at ward or hospital level. Patient level variance was 3.28987, ward level 0.07115, and hospital level 0.00004. Likelihood ratio tests confirmed this finding, so simple single-level models were used.

Prevalence of HOD and influencing factors

The crude point-prevalence of HOD was 4.5% (230/5142 patients; 95% confidence interval (CI) 3.9-5.0%). Another 395 patients had one episode of unformed stool in the day before data collection.

HOD prevalence in teaching hospitals was 5.9% (163/2748, 95% CI 5.1-6.9%), more than twice the 2.8% prevalence in acute hospitals (67/2394, 95% CI 2.1-3.5%; odds ratio 2.2, 95% CI 1.7-3.0). Prevalence was unaffected by season, hospital size, or ward characteristics (specialty, number of beds and side-rooms).

Patient and HOD characteristics

The median age of HOD patients was 77 (IQR 66-85years); 52% were female. Median duration of hospitalisation at HOD onset was eight days (IQR 5-15 days). Patients reported a median of three episodes of diarrhoea the day before data collection; median symptom duration was two days (IQR 1-6 days). Sixty (26%) patients had experienced more than one previous episode of HOD during their admission and 14 (6%) had a history of CDI.

Potential causes of HOD

Nearly all HOD patients (222, 97%) had at least one potential cause of diarrhoea. The majority (196 patients, 85%) had multiple possible causes (median 3; IQR 2-5). Almost half the patients, 107 (47%) had underlying conditions associated with diarrhoea (Table I). The commonest was constipation with overflow diarrhoea (39/230, 17%). Seven patients had no relevant diagnosed condition but reported longstanding diarrhoea.

Just over half (125, 54%) of HOD patients were exposed to ≥ 1 antimicrobial within the 24 hours before diarrhoea onset (Table II). In total, 123/201 (61%) antimicrobials administered were intravenous; 184 (91%) were started in hospital. Antibiotics are listed by group and stratified by risk of CDI in Supplementary Material, Table S2. The commonest groups were: beta lactam-beta lactamase inhibitor combinations (61/230, 27%), macrolides (17, 7%) and carbapenems (15, 7%).

Most patients with HOD (195/230, 85%) were prescribed ≥ 1 non-antimicrobial drug that can cause diarrhoea (median 2, IQR 1-3; Table II). Diarrhoea is reported as a "very common" (occurring with a frequency ≥ 1 in 10) or "common" side effect (occurring with a frequency ≥ 1 in 100) for all drugs listed, except steroids and levothyroxine, for which the frequency is

unknown [20]. The commonest were laxatives (150/230, 65%), proton-pump inhibitors (100, 44%) and selective serotonin-reuptake inhibitors (30, 13%). In total, 247/442 (56%) of all medications were started before admission. Medication started pre-admission was the only potential cause of HOD in 21 (9%) of cases.

Microbiological investigation of HOD

Eighty patients with HOD (35%) had stool microbiology testing following diarrhoea onset; in 56/80 patients (71%) the stool sample was obtained within 24 hours. Seventy-five HOD patients (33%) were tested for CDI; this included 48 of the 125 (38%) patients exposed to antimicrobials. The proportion of patients receiving ≥ 1 antimicrobial associated with moderate to high risk of CDI (defined in Table S2) who were tested for CDI was only slightly higher (42/103, 41%). All laboratories followed a two-stage testing algorithm, consistent with UK guidance [21]. Nine HOD patients were faecal-toxin positive (4%); a further four were GDH positive with negative toxin assays, suggesting *C. difficile* colonization. There was no difference between patients who were tested for CDI (including those with the infection), and those not tested, in terms of frequency of diarrhoea, the median number and distribution of potential causes of HOD per patient (Table III). Only documentation of HOD in medical notes had a significant association with CDI testing (78% of those tested versus 38% not tested, $p < 0.001$). Norovirus testing was undertaken in 16 patients; three were positive.

Patient management

Overall, 218/230 (95%) HOD patients had documentation of bowel movements; most commonly on a BSC (173, 79%). In 143 (62%) patients the number of diarrhoeal episodes in the 24 hours before the study day was documented. In 80 cases the number of patient-reported episodes could be compared to the number documented: 21 patients (26%) reported more, 6 (8%) reported fewer, whilst for 53 (66%) there was no difference. The presence of HOD was documented in the medical notes in 115/230 (50%) of cases. For 94/115 (82%) it was within two days of diarrhoea onset. There was documented evidence of medical assessment of the diarrhoea for 80/230 (35%) patients. In 64/80 (80%) cases this took place either on the day of diarrhoea onset or the following day.

Not including faecal sampling, 93 (40%) patients had ≥ 1 additional investigation for HOD (range 1-4; Table S3). One or more treatments was stopped, started or adjusted owing to

HOD in 61 patients (27%; Table S4). Laxatives were stopped in 30/150 (20%) of those receiving them.

At the time of developing HOD, 40/230 (17%) of patients were already in isolation; a further 46 (20%) were placed in a side-room following diarrhoea onset. The remaining 144 (63%) HOD patients were not isolated. This included 25 patients who were tested for *C. difficile* (one of whom was toxin positive), though a significantly higher proportion of those tested were isolated (27/52 (52%) tested patients v 11/129 (9%) untested patients, $p < 0.001$).

Discussion

Key findings

The point prevalence of HOD on general medical, surgical and elderly-care wards was 4.5%. HOD was associated with a prolonged hospital admission at symptom-onset and was recurrent in over a quarter of cases. Almost all patients had at least one identifiable potential cause of HOD; 85% had multiple potential causes. Only 35% of patients overall were tested for CDI, including only 38% of those receiving antimicrobials. Nine patients were *C. difficile* toxin positive (4%).

The rate of diarrhoea documentation was high, though a quarter of patients reported more episodes than documented. Only 35% of patients had a documented medical assessment of the diarrhoea. Of the patients not already in a side-room just 20% were isolated following diarrhoea onset. 48% of those tested for *C. difficile* were not isolated.

Strengths and limitations

This multicentre study is the largest published investigation of HOD, with data collected from multiple hospitals of different types across the UK. The narrow confidence intervals indicate the results could be used to estimate the prevalence of HOD on general wards in similar settings.

There are several limitations. The design precluded an assessment of the incidence and duration of HOD, which would require a cohort study. The hospitals involved were self-selected; their practices may differ from other hospitals in the NHS. Industrial action during

the winter data collection period led to a national reduction in the number of elective admissions. This may have led to under-representation of elective patients and lower bed occupancy rates for this period.

Patient recall bias or unrecognised cognitive impairment may have led to an underestimate of HOD prevalence, although to minimise this risk questions about diarrhoea were limited to the 24 hours before the study day and multiple information sources were reviewed for evidence of HOD. Data were mostly collected from the notes/electronic records, inaccuracies in which may have impacted the findings.

The proportion of cases of HOD caused by gastrointestinal infections, particularly viral pathogens, may have been underestimated as most patients were not tested for CDI, no hospitals cultured faecal samples from patients admitted over 72 hours and only one routinely tested such samples for norovirus. Newly available, rapid molecular diagnostic assays, that can simultaneously identify multiple pathogens in faecal samples, might increase the identification of infections in HOD patients [22].

Comparison with other studies

The prevalence of HOD was less than reported in most other studies, though meaningful comparison is difficult, owing to the diverse range of settings, lack of contemporaneous reports and varied definitions of HOD used [5-12]. It may reflect our exclusion of specialist units, where HOD prevalence is typically higher, e.g. intensive care, and the lower rate of CDI than some studies [5-7]. HOD prevalence was significantly higher in teaching hospitals, but unaffected by hospital and ward size, specialty, number of side-rooms and season. This suggests that patient factors and the complexity of their condition influences the risk of HOD, rather than differences in systems or processes, although unmeasured factors may have had an impact. The low level of norovirus activity in the UK during the study may explain why there were no seasonal differences [23].

Identifying the potential causes of HOD facilitates recognition of modifiable causes. Known causes of HOD include numerous medications, especially laxatives and antibiotics [24,25]. Almost two thirds of HOD patients were on laxatives. In only 20% were they stopped following the onset of diarrhoea, consistent with a previous study [8]. Constipation with overflow diarrhoea was the commonest underlying condition causing HOD. These related

findings suggest that improvements in bowel care could reduce the occurrence of HOD. Antibiotic treatment was a predisposing factor in half of HOD patients; since the incidence of diarrhoea varies between antibiotic used, there may be an opportunity to modify this risk through better antimicrobial stewardship.

Most patients were not tested for CDI. This may reflect close adherence to UK guidance, which advises testing HOD for CDI is unnecessary if there is an alternative cause, and in almost all patients at least one potential cause was evident [21]. Alternatively, in hospitals where the definition of HOD included a minimum daily frequency of diarrhoea, not testing for CDI may reflect failure to identify and document all episodes of diarrhoea in some patients. Failing to diagnose infectious causes of HOD, such as CDI and norovirus, has negative consequences for patients and hospitals. Patients can undergo unnecessary investigations, whilst treatment and instigation of appropriate infection control measures may be delayed [26,27]. In this study, forty percent of patients had at least one investigation and 27% had some alteration to their treatment as a result of HOD, adding to patient morbidity and hospital costs.

Almost two thirds of patients had no documented medical assessment of their diarrhoea. It is unclear whether this reflected a lack of awareness of HOD amongst clinicians (a known problem [8]) or that they did not consider the diarrhoea to be significant. A previous study found HOD receives little attention after CDI is excluded [7].

The failure to isolate most patients with diarrhoea is contrary to published guidelines and may increase the risk of faecal pathogen transmission [28,29]. Reasons for not isolating those tested for CDI merit further investigation but may reflect limited availability of single-room accommodation in many NHS hospitals.

Implications

The overall economic impact of HOD has not been evaluated, though CDI and norovirus are known to place a financial burden on healthcare organisations [3,4]. The cost of HOD is likely to be significant, given the large number of patients involved. In England there are approximately 100,000 acute hospital beds in the included specialties [30]. Applying the 4.5% prevalence result, this equates to approximately 3600 patients with HOD per day (1.3 million patient days of HOD/year). Given the scope of the problem, recommendations that

healthcare organisations monitor their prevalence of HOD has significant implications for infection control teams and may not be feasible without additional resources.

All organisations in this study had policies requiring source isolation of patients with diarrhoea but compliance was poor, with most patients not isolated, perhaps because of insufficient isolation capacity. Better management of HOD would facilitate more efficient use of side rooms.

UK guidelines advise against CDI testing when there is an alternative cause of HOD [20]. As the number and distribution of potential causes of HOD did not differ between patients tested for *C. difficile* (including those with CDI) and those not tested, a more inclusive testing strategy should be considered, to avoid missing CDI cases and contributing to onward transmission [27, 30]. If the proportion of patients with CDI (4%) was the same in the cohort of untested HOD patients (n=155) six cases were missed. As documentation of HOD in medical notes was associated with CDI testing, mandating medical assessment in all cases of HOD, could help to identify potentially infectious diarrhoea and encourage CDI testing and prompt isolation, consistent with published recommendations [29]. It could also shorten HOD duration, by ensuring appropriate management of non-infectious causes. Of note, NHS Improvement has given notice of its intention, from 2020, to examine both diarrhoea sampling and CDI testing rates in NHS hospitals [31].

Conclusions

The prevalence of HOD on general medical, surgical and elderly-care wards in NHS hospitals was 4.5%, suggesting there are over a million patient days of HOD on these wards in English hospitals each year. Multiple potential causes of HOD can be identified in most patients, complicating its management, including decisions around CDI testing. Work to reduce impact on patients and healthcare organisations should focus on ensuring patients with HOD undergo medical assessment focussed on the need for CDI testing and isolation and identifying and managing other common, modifiable causes.

Acknowledgements

The authors would like to thank the following for their contribution to this study. Graziella Kontkowski (from UK *C. difficile* Support) provided insights in to the importance of HOD from a patient perspective. In Leeds, Dermot Burke, Clare Donnellan and Claire Berry. In Papworth, Margaret Gillham, Helen Wickenden and Victoria Stoneman. In Belfast, Michelle Copeland. In Hampshire Hospitals NHS Foundation Trust, Hazel Gray.

Conflict of interest statement

Damian Mawer received a fellowship grant from the Healthcare Infection Society to undertake the work.

Mark Wilcox reports grants and personal fees from Actelion, grants and personal fees from Cubist, grants and personal fees from Astellas, grants and personal fees from Merck, grants and personal fees from Sanofi-Pasteur, grants and personal fees from Summit, grants and personal fees from Seres, grants and personal fees from Biomerieux, grants from Da Volterra, personal fees from Valneva, personal fees and other from Alere, grants and personal fees from Qiagen, grants and personal fees from Pfizer, other from Astellas Pharma , non-financial support from Astellas Pharma , personal fees from Ferring, personal fees from Synthetic Biologics, during the conduct of the study; personal fees from Astra Zeneca, personal fees from Nabriva, personal fees from Pfizer, personal fees from Roche, personal fees from The Medicine Company, grants and personal fees from Abbott, personal fees from Basilea, grants and personal fees from European Tissue Symposium, personal fees from Bayer, personal fees from Allergan, personal fees from Menarini, personal fees from Motif Biosciences, grants and personal fees from Paratek, personal fees from AiCuris, personal fees from Antabio, personal fees from Spero, grants and personal fees from Tetraphase, grants and personal fees from Surface Skins, outside the submitted work.

None of the other authors have any conflicts of interest to declare.

Funding sources

Damian Mawer was funded by the Healthcare Infection Society during the work.

628 Jordan Skittrall is funded by a National Institute for Health Research Academic Clinical Fellowship.
629 This paper presents independent research funded by the National Institute for Health Research
630 (NIHR). The views expressed are those of the authors and not necessarily those of the NHS, the NIHR
631 or the Department of Health.

632

References

1. Patient Safety Domain, NHS England. Clostridium difficile infection objectives for NHS organisations in 2015/16 and guidance on sanction implementation. London: NHS England, 2015. Available at: <https://www.england.nhs.uk/wp-content/uploads/2015/07/cdClostridium-difficile-infection-objectives-for-NHS-organisations.pdf> [Accessed 3 September 2018].
2. Polage CR, Solnick JV, Cohen SH. Nosocomial diarrhea: evaluation and treatment of causes other than Clostridium difficile. Clin Infect Dis. 2012;55:982-9
3. Wiegand PN, Nathwani D, Wilcox MH, Stephens J, Shelbaya A, Haider S. Clinical and economic burden of Clostridium difficile infection in Europe: a systematic review of healthcare-facility-acquired infection. J Hosp Infect. 2012;81:1-14
4. Lopman BA, Reacher MH, Vipond IB, et al. Epidemiology and cost of nosocomial gastroenteritis, Avon, England, 2002-2003. Emerg Infect Dis. 2004;10:1827-34
5. McFarland LV. Epidemiology of infectious and iatrogenic nosocomial diarrhea in a cohort of general medicine patients. Am J Infect Control. 1995;23:295-305
6. Samore MH, DeGirolami PC, Tlucko A, Lichtenberg DA, Melvin ZA, Karchmer AW. Clostridium difficile colonization and diarrhea at a tertiary care hospital. Clin Infect Dis. 1994;18:181-7
7. Garey KW, Graham G, Gerard L, et al. Prevalence of diarrhea at a university hospital and association with modifiable risk factors. Annal pharmacother. 2006;40:1030-4
8. Kyne L, Moran A, Keane C, O'Neill D. Hospital-acquired diarrhoea in elderly patients: epidemiology and staff awareness. Age Ageing. 1998;27:339-43
9. Bhuiyan MU, Luby SP, Zaman RU, et al. Incidence of and risk factors for hospital-acquired diarrhea in three tertiary care public hospitals in Bangladesh. Am J Trop Med Hyg. 2014;91:165-72
10. Zaidi M, Ponce de Leon S, Ortiz RM, et al. Hospital-acquired diarrhea in adults: a prospective case-controlled study in Mexico. Infect Control Hosp Epidemiol. 1991;12:349-55
11. Schwaber MJ, Simhon A, Block C, Roval V, Ferderber N, Shapiro M. Factors associated with nosocomial diarrhea and Clostridium difficile-associated disease on the adult wards of an urban tertiary care hospital. Eur J Clin Microbiol Infect Dis. 2000;19:9-15
12. McErlean A, Kelly O, Bergin S, Patchett SE, Murray FE. The importance of microbiological investigations, medications and artificial feeding in diarrhoea evaluation. Irish J Med Sci. 2005;174: 21-5
13. Wiesen P, Van Gossum A, Preiser JC. Diarrhoea in the critically ill. Curr Opin Crit Care. 2006;12:149-54
14. Cox GJ, Matsui SM, Lo RS, et al. Etiology and outcome of diarrhea after marrow transplantation: a prospective study. Gastroenterol. 1994;107:1398-407
15. Health and Social Care Information Centre. Hospital Episode Statistics, Admitted Patient Care - England, 2014-15: Main specialties [Microsoft Excel spreadsheet]. Available at: <http://content.digital.nhs.uk/searchcatalogue?productid=19420&q=title%3a%22Hospital+Episode+Statistics%2c+Admitted+patient+care+-+England%22&sort=Relevance&size=10&page=1#top> [Accessed 5 February 2017].

16. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. STROBE Initiative. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *BMJ*. 2007;335:806-8.
17. Lewis SJ, Heaton KW. Stool form scale as a useful guide to intestinal transit time. *Scand J Gastroenterol*. 1997;32:920-4
18. R Core Team (2016). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Available at: <https://www.R-project.org/> [Accessed 8 March 2017].
19. Bates D, Maechler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using lme4. *J Stat Softw*. 2015;67:1-48
20. Joint Formulary Committee. British National Formulary (online) London: BMJ Group and Pharmaceutical Press. Available at: <http://www.medicinescomplete.com> [Accessed 4 February 2018].
21. Department of Health. Updated guidance on the diagnosis and reporting of *Clostridium difficile*. London: Department of Health, 2012. Available at: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/215135/dh_133016.pdf [Accessed 5 January 2015].
22. Macfarlane-Smith LR, Ahmed S, Wilcox MH. Molecular versus culture-based testing for gastrointestinal infection. *Curr Opin Gastroenterol*. 2018;34:19-24.
23. Public Health England. PHE national norovirus and rotavirus report. Summary of surveillance of norovirus and rotavirus. 11 January 2017 – data to week 52. Public Health England, January 2017. Available at: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/591226/Norovirus_update_2016_week_52.pdf [Accessed 7 March 2018].
24. Abraham B, Sellin JH. Drug-induced diarrhea. *Curr Gastroenterol Rep*. 2007;9:365-72
25. Cote GA, Buchman AL. Antibiotic-associated diarrhoea. *Expert Opin Drug Saf*. 2006;5:361-72
26. Beersma MF, Sukhrie FH, Bogerman J, et al. Unrecognized norovirus infections in health care institutions and their clinical impact. *J Clin Microbiol* 2012;50: 3040-5.
27. Frenz MB, McIntyre AS. 2003. Reducing delays in the diagnosis and; treatment of *Clostridium difficile* diarrhoea. *QJM*. 2003;96:579–582.
28. McDonald LC, Derding DN, Johnson S, et al. Clinical practice guidelines for *Clostridium difficile* infection in adults and children: 2017 update by the Infectious Diseases Society of America (IDSA) and Society for Healthcare Epidemiology of America (SHEA). *Clin Infect Dis*. 2018;66:e1-e48.
29. Health Protection Agency and Department of Health. UK. *Clostridium difficile* infection: How to deal with the problem. 2008. Available at: <https://www.gov.uk/government/publications/clostridium-difficile-infection-how-to-deal-with-the-problem> [Accessed: 13 February 2019].
30. NHS England. Bed availability and occupancy data – overnight. Data for NHS organisations in England, Quarter 3, 2017-18. Available at: <https://www.england.nhs.uk/statistics/statistical-work-areas/bed-availability-and-occupancy/bed-data-overnight/> [Accessed 7 March 2018].
31. NHS Improvement. *Clostridium difficile* infection objectives for NHS organisations in 2019/20 and guidance on the intention to review financial sanctions and sampling rates from 2020/21 Available at:

718 [https://improvement.nhs.uk/documents/808/CDI objectives for NHS organisation](https://improvement.nhs.uk/documents/808/CDI_objectives_for_NHS_organisations_in_2019_Feb19.pdf)
719 [s in 2019 Feb19.pdf](https://improvement.nhs.uk/documents/808/CDI_objectives_for_NHS_organisations_in_2019_Feb19.pdf) [Accessed 5 March 2019].
720

ACCEPTED MANUSCRIPT

721 Tables

	N	% (n=230)
Number of underlying conditions		
Any	107	47
1	84	37
2	18	8
3	4	2
4	0	-
5	0	-
6	1	<1
Underlying condition		
Constipation with overflow diarrhoea	39	17
Previous bowel resection or recent abdominal surgery	35	15
Diverticular disease (including diverticulitis)	18	8
Severe sepsis (non-gastrointestinal source)	11	5
Colorectal cancer (in the past 5 years)	8	4
Inflammatory bowel disease	8	4
Stoma	8	4
Gastrointestinal bleed (within the last 72 hours)	7	3
Liver disease	6	3
Pancreatitis	4	2
Biliary sepsis	2	1
Irritable bowel syndrome	2	1
Short bowel syndrome	2	1
Other*	20	9

*Alcohol withdrawal, anxiety, bile acid malabsorption, bowel perforation, colonoscopy with extensive irrigation, faecal incontinence, fruit induced (2 cases), gastric cancer, graft versus host disease, high-output stoma, ileal ulceration, intravenous drug use, ischaemic colitis, non-compliance with medication, pancreatic insufficiency, peritoneal dialysis, rectal prolapse, small bowel fibrotic stricture, thyrotoxicosis

Table I: Number (percentage) of underlying or pre-existing conditions associated with diarrhoea in HOD patients

730

Medication	Total		Started pre-admission		Started post-admission	
	n	% of all HOD patients (n=230)	n	% of patients on that medication	n	% of patients on that medication
Laxatives	150	65	67	45	83	55
Antimicrobials*	125	54	17	8	184	92
Proton pump inhibitors	100	44	81	81	19	19
Selective serotonin reuptake inhibitors	30	13	28	93	2	7
Steroids	25	11	14	56	11	44
Iron (oral)	22	10	14	64	8	36
Metformin	21	9	20	95	1	5
Enteral feed	18	8	4	22	14	78
Enema (in the preceding 24hr)	14	6	0		14	100
Glycerine suppositories	10	4	1	10	9	90
Metoclopramide	10	4	1	10	9	90
Immunosuppressants other than steroids	7	3	5	71	2	29
Bowel preparation	5	2	0		5	100
Donepezil	3	1	3	100	0	0
Nutritional supplements	3	1	0		3	100
Chemotherapy	2	1	1	50	1	50
Levothyroxine	2	1	2	100	0	0
Magnesium salts	2	1	0	0	2	100
Phosphate and potassium salts	2	1	0	0	2	100
Other**	15	7	5	38	8	50

*Figures (and percentages) shown for the timing of when antimicrobials were started are based on the total number of antimicrobials prescribed (n=201), as some patients were on a combination of antibiotics started both before and after admission at the time of diarrhoea onset.

**Other medication judged by investigator to be a potential cause of HOD (whether started before or after admission not known in 2 cases): allopurinol, bisoprolol, calcium/vitamin D, colchicine, furosemide, Gastrografin, lithium, iopanoic acid, opiates (patient reported), pancrelipase (2 cases), quetiapine, ranitidine, sevelamer, simvastatin.

731 **Table II: Prescriptions of medications associated with diarrhoea in patients with HOD**

732


	No CDI test, No. (%)	CDI test, No. (%)	Adjusted OR (95% CI)	P value
Total	155	75	-	-
Age (mean \pm SD)	73 \pm 17	73 \pm 15	1.00 (0.98, 1.03)	0.80
Sex (m)	76 (49)	34 (45)	0.85 (0.45, 1.63)	0.70
No. of potential causes of HOD/ patient (median)	3	3	-	-
No. of diarrhoea episodes in 24 hr before the survey (median)	3	3	1.10 (0.94, 1.29)	0.26
Any underlying condition associated with diarrhoea	74 (48)	33 (44)	0.77 (0.40, 1.49)	0.70
Receiving antimicrobials	78 (50)	47 (63)	1.73 (0.89, 3.37)	0.11
Any other medication that can cause diarrhoea	130 (84)	65 (87)	1.38 (0.52, 3.62)	0.72
Medication started pre-admission only potential cause of HOD	17 (11)	4 (5)	0.42 (0.11, 1.59)	0.25
HOD documented in medical notes	59 (38)	58 (78)	6.47 (3.31, 12.66)	<0.001

733

734 Table III: Clinical features and potential causes of HOD in patients tested for CDI versus those
 735 not tested for CDI.

736

737

738 **Figures**

739

740 **Figure 1: Flow diagram showing patient recruitment and results, both overall and by season and**
 741 **hospital type.**