

European Journal of Obstetrics & Gynecology and Reproductive Biology

Pregnancy outcomes following different types of bariatric surgery: a national cohort study

--Manuscript Draft--

Manuscript Number:	EJOGRB-21-23633
Article Type:	Full Length Article
Section/Category:	Obstetrics
Keywords:	Pregnancy; bariatric surgery; bypass; banding; sleeve; cohort
Corresponding Author:	Katie Cornthwaite, MRCOG University of Bristol Bristol, UNITED KINGDOM
First Author:	Katie Cornthwaite, MRCOG
Order of Authors:	Katie Cornthwaite, MRCOG Chetan Prajapati, PhD Erik Lenguerrand, PhD Marian Knight, DPhil, FRCPE Natalie Blencowe, MRCS, PhD Andrew Johnson, MD, FRCP Tim Draycott, MD, FRCOG Dimitrios Siassakos, MD, FRCOG
Abstract:	<p>Objective</p> <p>To assess the impact of type of bariatric surgery on pregnancy outcomes.</p> <p>Study design</p> <p>This is a national prospective observational study using the UK Obstetric Surveillance System (UKOSS). Data collection was undertaken in 200 consultant-led NHS maternity units between November 2011 and October 2012 (gastric banding), and April 2014 and March 2016 (gastric bypass and sleeve gastrectomy). Participants were pregnant women following gastric banding (n=127), gastric bypass (n=134) and sleeve gastrectomy (n=29).</p> <p>Maternal and perinatal outcomes were compared using generalised linear and linear mixed models. Maternal outcomes included gestational weight gain, pre-eclampsia, gestational diabetes, anaemia, surgical complications. Perinatal outcomes included birthweight, small / large for gestational age (SGA/LGA), preterm birth, stillbirth.</p> <p>Results</p> <p>Maternal</p> <p>Women pregnant after gastric banding and sleeve gastrectomy had a lower risk of anaemia compared with gastric bypass (banding (16%) vs bypass (39%): p=0.002, sleeve (21%) vs bypass: p=0.04). Gestational diabetes risk was lower after gastric banding compared with gastric bypass (7% vs 16%, p=0.03) despite women with banding having significantly greater weight at booking as well as gestational weight gain. Women pregnant after gastric banding and sleeve gastrectomy had a lower risk of surgical complications than after gastric bypass (banding (0.9%) vs bypass (11.4%): p=0.03, sleeve (0.0%) vs bypass: p=0.06).</p> <p>Perinatal</p>

	<p>Infants born to mothers after gastric banding had a higher birthweight than those born to mothers after gastric bypass (mean difference=260g (125-395), $p<0.001$). Infants were more likely to be LGA if their mothers had gastric banding compared with gastric bypass or sleeve gastrectomy (banding (21%) vs bypass (5%): $p=0.006$; banding vs sleeve (3%): $p=0.03$). Risk of preterm birth was higher in women with gastric banding compared with gastric bypass (13% vs 8%, $p=0.04$).</p> <p>Conclusions</p> <p>Women planning bariatric surgery should be counselled regarding the differing impacts of different types of procedure on any future pregnancy. Pre-existing gastric bypass is associated with higher rates of potentially serious surgical complications during pregnancy.</p>
Suggested Reviewers:	Sarah Stock Sarah.Stock@ed.ac.uk
	Lucy Chappell lucy.chappell@kcl.ac.uk
Opposed Reviewers:	

Translational Health Sciences
University of Bristol
Department of Women's Health,
Southmead Hospital,
Southmead Road,
Bristol,
BS10 5NB,
UK
kc17995@bristol.ac.uk
+447834597324

22nd January 2021

Dear Editor-in-Chief,

Re: Manuscript submission: Pregnancy outcomes following different types of bariatric surgery: a national cohort study

Thank you for considering the above article for publication in EJOG. We hope that this will be a topic of interest to your readers.

Over a **quarter of a million** bariatric surgery procedures are performed throughout the world each year and over half of these are in women of childbearing age (IFSO Global Registry). This has led to a dramatic increase in pregnancies following weight loss surgery and rates are only expected to rise as the global burden of obesity increases. Several different bariatric procedures are available to women considering weight loss surgery, each with distinct mechanisms of action. We have included a table in our paper with a summative literature review which highlights a historic tendency to group bariatric procedures together when evaluating pregnancy outcomes.

In this study, we used a nationwide obstetric surveillance system (UKOSS) to investigate the impact of the most widely used bariatric procedures (gastric bypass, sleeve gastrectomy and gastric banding) on pregnancy outcomes. Our study is **prospective, conducted over a short time frame**, and has good numbers of participants representing the majority of women in the UK during the period of study. Contrary to previous studies that included women over long periods and conflated different types of surgery to achieve statistical significance and publication, this study is unique in comparing outcomes between distinct types of surgery.

A key finding, reported for the first time in the literature, is that women with gastric bypass have a rate of **serious surgical complications** of 11.4% during pregnancy, significantly higher (both clinically and statistically) than women with banding (0.9%) and sleeve gastrectomy (0%).

This has clear **implications for consultations** between surgeons, physicians, general practitioners and obstetricians, with women before and during pregnancy.

We also found evidence of increased risk of maternal anaemia and lower birthweight in infants born to women who had undergone gastric bypass, whilst the risk of preterm birth was higher amongst women who had gastric banding. These findings may influence a woman's preference for a specific procedure if considering a pregnancy following bariatric surgery. Obstetricians and obstetric physicians must also be vigilant regarding the distinct risks associated with these pregnancies.

We believe that this study will be of considerable interest to the readers of EJOG and are grateful for your consideration.

We confirm that this manuscript has not been published elsewhere and is not under consideration by another journal.

Many thanks.
Yours faithfully,

Katie Cornthwaite & Dimitrios Siassakos, on behalf of the authors

Pregnancy outcomes following different types of bariatric surgery: a national cohort study

Katie Cornthwaite, MRCOG^{a, b}, Chetan Prajapati, PhD^b, Erik Lenguerrand, PhD^b, Marian Knight, DPhil, FRCPE^c, Natalie Blencowe, MRCS, PhD^d, Andrew Johnson, MD, FRCP^e, Tim Draycott, MD, FRCOG^a, Dimitrios Siassakos, MD, FRCOG^{f, g, h}

^a Women's Health Department, North Bristol NHS Trust, UK

^b Translational Health Sciences, University of Bristol, UK

^c National Perinatal Epidemiology Unit, University of Oxford, UK

^d Population Health Sciences, University of Bristol, UK

^e Diabetes and Endocrinology Department, North Bristol NHS Trust, UK

^f Elizabeth Garrett Anderson Institute for Women's Health, University College London, UK

^g NIHR Biomedical Research Centre at University College London Hospital

^h Wellcome/EPSRC centre for Surgical and Interventional Sciences (WEISS)

Corresponding author

Dr Katie Cornthwaite

The Chilterns

North Bristol NHS Trust

Westbury-on-Trym

Bristol

BS10 5NB

Tel: 07834597324

Email: kc17995@bristol.ac.uk

Abstract

Objective

To assess the impact of type of bariatric surgery on pregnancy outcomes.

Study design

This is a national prospective observational study using the UK Obstetric Surveillance System (UKOSS). Data collection was undertaken in 200 consultant-led NHS maternity units between November 2011 and October 2012 (gastric banding), and April 2014 and March 2016 (gastric bypass and sleeve gastrectomy).

Participants were pregnant women following gastric banding (n=127), gastric bypass (n=134) and sleeve gastrectomy (n=29).

Maternal and perinatal outcomes were compared using generalised linear and linear mixed models. Maternal outcomes included gestational weight gain, pre-eclampsia, gestational diabetes, anaemia, surgical complications. Perinatal outcomes included birthweight, small / large for gestational age (SGA/LGA), preterm birth, stillbirth.

Results

Maternal: Women pregnant after gastric banding and sleeve gastrectomy had a lower risk of anaemia compared with gastric bypass (banding (16%) vs bypass (39%): $p=0.002$, sleeve (21%) vs bypass: $p=0.04$). Gestational diabetes risk was lower after gastric banding compared with gastric bypass (7% vs 16%, $p=0.03$) despite women with banding having significantly greater weight at booking as well as gestational weight gain. Women pregnant after gastric banding and sleeve gastrectomy had a lower risk of surgical complications than after gastric bypass (banding (0.9%) vs bypass (11.4%): $p=0.03$, sleeve (0.0%) vs bypass: $p=0.06$).

Perinatal: Infants born to mothers after gastric banding had a higher birthweight than those born to mothers after gastric bypass (mean difference=260g (125-395),

p<0.001). Infants were more likely to be LGA if their mothers had gastric banding compared with gastric bypass or sleeve gastrectomy (banding (21%) vs bypass (5%): p=0.006; banding vs sleeve (3%): p=0.03). Risk of preterm birth was higher in women with gastric banding compared with gastric bypass (13% vs 8%, p=0.04).

Conclusions

Women planning bariatric surgery should be counselled regarding the differing impacts of different types of procedure on any future pregnancy. Pre-existing gastric bypass is associated with higher rates of potentially serious surgical complications during pregnancy.

Keywords

Pregnancy, bariatric surgery, bypass, banding, sleeve, cohort

1. Introduction

One in five pregnant women in the UK have a BMI >30 and at least 5% have a BMI >35.[1,2] The adverse consequences of obesity on maternal and perinatal health are well established.[3–6] Weight loss can mitigate these risks and is increasingly being achieved with bariatric surgery.[7] In the UK, over 75% of weight loss operations are in women of childbearing age.[8] Pregnancy after bariatric surgery is now more common,[9] but the preference for type of surgery has changed over time, with sleeve gastrectomy increasingly popular.[10,11]

Evidence from observational studies and systematic reviews has highlighted the benefits of bariatric surgery in reducing obesity-related pregnancy complications such as gestational diabetes.[9,12–14] However, improvements in some maternal outcomes may be at the expense of fetal wellbeing, with higher rates of small for gestational age (SGA) infants and preterm labour in pregnancies following bariatric surgery.[9,12–14] Most studies have conflated different types of bariatric surgery when evaluating pregnancy outcomes, or examined retrospectively cohorts of women spanning several years and likely receiving different care over time (Table 1). However, there are mechanistic differences between bariatric procedures (Figure 1), which are likely to have distinct effects on the mother and fetus. The impact of different types of bariatric surgery on pregnancy outcomes are unclear and conflicting.[9,14–16]

It is essential that healthcare professionals involved in the care of women of reproductive age understand the impact of different types of bariatric surgery on future pregnancy outcomes.[17] Using a nationwide obstetric surveillance system

(UKOSS), we investigated the procedure-related impact of bariatric surgery on pregnancy outcomes in distinct national cohorts.

2. Materials and Methods

2.1. Study design

National prospective observational study of women pregnant after gastric band, bypass and sleeve. Women were identified through the UK Obstetric Surveillance System (UKOSS)[18] in which all 200 consultant-led UK maternity units participate. Nominated clinicians in each unit reported cases and completed a customised, anonymous, data collection form using routinely collected data (<https://www.npeu.ox.ac.uk/ukoss/methodology>).

2.2. Study population

A cohort study of women pregnant with gastric banding was undertaken between November 2011 and October 2012. Outcomes of these pregnancies, according to band management (inflation versus deflation), are published in detail elsewhere.[19]

To understand pregnancy outcomes after other types of bariatric surgery and given the change in preference for type of surgery, a separate cohort of pregnant women who had undergone bypass surgery was collected between April 2014 and March 2016. This cohort was divided into women with gastric bypass (Roux-en-Y, loop and unspecified bypass) and sleeve. Women with undefined bariatric surgery were excluded. Cases of biliopancreatic diversion and duodenal switch were removed since they are associated with more profound malabsorption and are now rarely performed.[20] We also removed cases with missing outcome data from comparative

analyses, as well as multifetal pregnancies due to their differing gestation, fetal growth and complication rates.

2.3. Outcomes and covariates

Maternal outcomes were gestational weight gain (GWG) (kg), BMI change (kg/m²), gestational hypertension, pre-eclampsia, gestational diabetes (GDM), anaemia, induced labour, mode of birth (vaginal or caesarean), surgical complications, major medical complications, and maternal death. GWG and BMI change were derived from the difference between the first (booking) and late third trimester measurements. Hypertensive disorders and anaemia diagnosed during pregnancy were as defined by the reporting hospital. Surgical complications included surgical procedure in pregnancy, band rupture, band slipping, internal hernia, bowel obstruction, incisional hernia, cholelithiasis and gastric dumping syndrome. Major medical complications included cardiac arrest, acute respiratory distress syndrome, disseminated intravascular coagulation, septicaemia, thrombotic event and intensive care unit (ICU) admission.

Perinatal outcomes were birthweight (grams), low birthweight (<2.5kg), macrosomia (>4kg), SGA infants, large for gestational age (LGA) infants, gestational age at birth (completed weeks), preterm birth (<37 weeks), neonatal unit admission, low Apgar score (<7 at 5 minutes), congenital abnormalities, stillbirth and neonatal death.

Birthweight percentiles for gestation (z scores) were calculated using UK-WHO growth reference charts.[21] SGA was defined as birthweight <10th percentile for gestational age and LGA as birthweight >90th percentile for gestational age.

1 Confounding factors were included in the multivariable model if biologically plausible.
2
3 Covariates included age, parity, ethnicity, early pregnancy BMI, smoking status,
4
5 employment status and pre-existing diabetes.
6
7
8
9

10 *2.4. Statistical methods*

11
12 Procedure-specific prevalence rates were derived using number of maternities after
13
14 each type of bariatric surgery in one-year as the nominator and total number of UK
15
16 maternities (in 2012 for gastric banding and 2015 for gastric bypass and sleeve)
17
18 recorded in NHS maternity statistics as the denominator.[22]
19
20
21
22
23

24 Frequencies and proportions, or means and medians with respective standard
25
26 deviations and interquartile ranges are reported, depending on data distribution. The
27
28 three bariatric groups could not be compared within the same modelling framework
29
30 since data in the sleeve group was sparse and unbalanced. The following inferential
31
32 statistics were performed:
33
34
35
36
37
38

39 *2.4.1. Comparisons between maternities after gastric banding and bypass*

40
41 Univariable and multivariable comparisons of categorical outcomes were performed
42
43 using modified Poisson regression[23] and results reported as RRs and 95%CI.
44
45 GWG and BMI change were compared using linear mixed regression to account for
46
47 repeated measurements within the same individuals. To account for missing data on
48
49 third trimester weight and BMI, we used multiple imputation (20 sets using chain
50
51 equations under missing at random (MAR) assumption)[24] and Rubin's combination
52
53 rules.[25] Sensitivity analyses were performed for three scenarios using single value
54
55 imputation under missing not at random (MNAR) assumption: weights or BMIs equal
56
57
58
59
60
61
62
63

1 to the 25th, 50th and 75th percentiles of third trimester observed weights or BMIs for
2 each surgical group were imputed. Birthweights were compared using generalized
3 least square model, and gestational age using quantile regression due to skewed
4 distribution.
5
6
7
8
9

10 11 *2.4.2. Comparisons of maternities after sleeve gastrectomy with gastric banding and* 12 *bypass* 13 14 15

16 Comparisons were unadjusted due to the small number of maternities in the sleeve
17 group. Categorical outcomes were compared using unconditional Barnard test,
18 which tests the null hypothesis of no difference in proportion between groups. This
19 test is more appropriate than Fisher's exact test when outcomes are rare and
20 unbalanced.[26] Results reported as differences in proportion and 95%CI.
21
22
23
24
25
26

27 Continuous outcomes were analyzed using the methods described above, including
28 imputation of missing data for GWG and BMI change.
29
30
31
32
33
34
35

36 *2.4.3. Subgroup analysis within gastric bypass group* 37 38

39 Outcomes were compared between maternities after Roux-en-Y gastric bypass and
40 unknown type of bypass. Loop and laparoscopic bypass were uncommon and not
41 analyzed.
42
43
44
45
46
47

48 Analyses were performed using Stata version 14.2, and R 3.5.1.
49
50
51
52

53 *2.5. Ethical approval* 54 55 56 57 58 59 60 61 62

1 Data were collected with Research Ethics Committee approvals (gastric band: NRES
2 11/SW/0227, gastric bypass: NRES 14/LO/0491). UKOSS methodology has
3
4 Research Ethics Committee approval.[27]
5
6
7
8
9

10 **3. Results**

11
12 UKOSS data collection was complete for 94% of notified women during the collection
13 periods. We collected data for 333 maternities following bariatric surgery (Figure 2).
14
15
16

17
18 Between November 2011 and October 2012, there were 127 cases of women
19 pregnant after gastric banding in an estimated 721,574 maternities[22] (UK
20 prevalence: 17.6 per 100,000 maternities (95%CI [14.7-21.0])). Between April 2014
21 and March 2016, there were 134 cases of pregnancy following gastric bypass (UK
22 prevalence: 9.7 per 100,000 maternities (95%CI [8.2-11.5])) and 29 cases of sleeve
23 gastrectomy (UK prevalence: 2.1 per 100,000 maternities (95%CI [1.4-3.1])) in an
24 estimated 1,377,097 maternities.[22]
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

40 Details of participant characteristics for pregnancies after gastric band, bypass and
41 sleeve are presented in Table 2 and for subgroups of gastric bypass in Table S1.
42
43
44
45
46
47
48
49

50 *3.1. Maternal outcomes*

51 *3.1.1. Gestational weight gain (GWG) and BMI change (Figure 3 and Table S2)*

52 Booking weight was higher in women pregnant with gastric banding than those with
53 bypass (+8.0kg, 95%CI [2.4-13.5]) and sleeve (+10.9kg, 95%CI [2.9-19.0]). There
54 was evidence of greater GWG in the gastric banding compared with bypass group
55
56
57
58
59
60
61
62
63
64
65

when missing data were accounted for with multiple imputation under MAR assumption (+3.6kg, 95%CI [0.24-7.0]). Sensitivity analysis using single imputation under MNAR assumption further strengthened these findings.

GWG was comparable between pregnancies after sleeve gastrectomy and gastric banding or bypass when multiple imputation under MAR assumption was used for missing data. However, imputation of observed median and 75th percentile weights suggested evidence of greater GWG amongst women who had gastric banding compared with sleeve (scenario III (75th percentile): +9.3kg, 95%CI [1.7-16.9]).

3.1.2. Obstetric and medical outcomes (Tables 3 and S3)

Risk of anaemia during pregnancy following gastric banding (15.9%) and sleeve (21%) was lower compared to those who had bypass (38.6%) (band vs bypass: adjusted RR=0.44 [0.27-0.72]; sleeve vs bypass: difference in proportion=0.20, [0.01-0.35]).

Risk of gestational diabetes (GDM) was lower after gastric banding (7%) than after bypass (16%) (adjusted RR=0.35 [0.13-0.92]). In the gastric band cohort, 84% of women had GDM screening, compared with 63% in the bypass cohort. Type of screening was variable, with 45% of women with gastric bypass having oral glucose tolerance testing.

Other obstetric and medical outcomes were comparable between bariatric surgery types.

3.1.3. Surgical and major medical complications (Table 3)

Surgical complications were lower in pregnancies after gastric banding (0.9%) compared with bypass (11.4%) (adjusted RR=0.08 [0.008-0.70]). There were no significant differences in major medical complications between different types of surgery.

In the gastric bypass group, surgical complications included: 1 incisional hernia, 2 intussusceptions, 1 bowel obstruction, 3 cases of cholelithiasis and 9 cases of gastric dumping syndrome. One pregnant woman died from surgical complications related to her gastric bypass. Major medical complications included: 1 case of septicaemia, 1 thrombotic event and 4 cases of ICU admission.

In the gastric band group, one woman suffered band slippage necessitating total parenteral nutrition and laparoscopic removal. Major medical complications included: 2 thrombotic events and 1 ICU admission, unrelated to band complications.

3.2. Perinatal outcomes (Tables 4 and S4)

3.2.1. Fetal growth

Infants born to mothers who had gastric banding had a higher birthweight (mean=3380g) than those born to mothers who had bypass (mean=3159g) (mean difference=+260g, [125-395]). Infants born to mothers who had gastric banding were more likely to be LGA than those who had bypass (adjusted RR=4.74 [1.54-14.6]) or sleeve (difference in proportion=+0.17 [0.02-0.27]). The risk of an SGA infant was comparable between groups.

3.2.2. Preterm birth

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Infants of mothers who had gastric banding were more likely to be born preterm (13.1%) than those born to mothers who had bypass (8.3%) (adjusted RR=2.27 [1.02-5.03]). Rates of elective preterm birth >34 weeks was similar between groups (banding: 3.7%, bypass: 3.0%). The risk of preterm birth was comparable between women who had sleeve and bypass / banding.

3.2.3. Morbidity and mortality

Risk of neonatal unit admission, low Apgar score, congenital abnormalities, stillbirth and perinatal death were similar between groups.

3.3. Subgroup analysis (Tables S5 and S6)

There was a higher risk of macrosomia in pregnancies following unspecified gastric bypass compared with Roux-en-Y (difference in proportion=-0.07[-0.17 - -0.01]). All other maternal and perinatal outcomes were similar between bypass subgroups.

4. Discussion

Pregnancies after gastric bypass were associated with increased risk of surgical complications and maternal anaemia compared with those after gastric banding or sleeve. Infants born to women who had gastric bypass had a lower birthweight but no increased risk of being SGA compared with infants of mothers who had banding. However, there was a higher risk of LGA infants in the gastric banding compared with bypass and sleeve groups. The risk of preterm birth was also higher in maternities following gastric banding compared with bypass.

4.1. Strengths and limitations

1 This is the first national prospective study to compare pregnancy outcomes between
2 distinct cohorts of women with different types of bariatric surgery. UKOSS is a well-
3 established surveillance system with comprehensive coverage and high
4 ascertainment. Data were collected prospectively eliminating recall bias. The
5 UKOSS cohorts spanned a total period of four and a half years. As opposed to
6 studies examining pregnancies across different decades (Table 1), comparisons
7 between the UKOSS cohorts are less likely to be affected by temporal changes in
8 healthcare practice. For comparisons between gastric bypass and banding, we
9 performed multivariable analyses adjusting for factors that may influence maternal
10 and perinatal outcomes. However, there may be residual confounding from
11 behaviour and lifestyle factors.

12 We did not perform power calculations as there was no single primary outcome on
13 which to base one and there is evidence against post-hoc sample size
14 calculation.[28] However, the relatively modest sample size limited our ability to
15 detect differences in rare outcomes and following sleeve gastrectomy. We undertook
16 steps to mitigate the impact of unbalanced and rare events and performed sensitivity
17 analyses to minimise the impact of missing data. We acknowledge that there may be
18 some residual heterogeneity in our comparison groups. For example, women
19 pregnant following gastric banding included those with an inflated and deflated
20 band.[19] Many women in the gastric bypass cohort had unspecified bypass surgery.
21 However, subgroup analyses revealed no significant differences in important
22 outcomes following Roux-en-Y and unspecified gastric bypass.

23 *4.2. Interpretation*

Differences in outcomes between restrictive and malabsorptive procedures may relate to macronutrient deficiencies, for which GWG can be used as a surrogate marker, and micronutrient deficiencies, for example, anaemia.[29] We found evidence of higher booking BMI and increased GWG in maternities after gastric banding compared with bypass. In contrast, pregnancies after gastric bypass were associated with a higher risk of anaemia compared to other bariatric procedures. Maternal anaemia is associated with an increased risk of low birthweight infants.[30] Existing evidence for risk of fetal growth restriction in pregnancies following malabsorptive compared with restrictive procedures is variable[15,16,30] In this study, women with gastric bypass had a lower infant birthweight compared to women with banding but we found no evidence of a difference in SGA risk.

Previous studies comparing outcomes with non-surgical controls, have found higher incidences of preterm birth in pregnancies following bariatric surgery[14] and specifically, gastric banding.[31] We identified a higher risk of spontaneous preterm birth in pregnancies after gastric banding compared with bypass. This may be related to the higher rates of residual obesity in the gastric band group, although this should in part be mitigated by adjustment for BMI.

The increased risk of GDM in women pregnant after gastric bypass compared with banding is surprising. It may be due to the type of diabetes screening performed: In women with gastric bypass, the conventional oral glucose tolerance test can trigger gastric dumping syndrome and variable glucose levels following the glucose load[32]. This can impair interpretation of results and lead to unpleasant symptoms

and potential harm to mother and baby. Therefore, alternative GDM screening methods are recommended for women pregnant after gastric bypass.[32,33]

This study highlights the increased risk of surgical complications in pregnancies following gastric bypass. Rising intra-abdominal pressure during pregnancy increases the risk of internal hernia[34] and small bowel obstruction.[35] These complications can have devastating consequences including maternal or fetal death.[36] In women of reproductive age, mesenteric defects should be closed at primary bypass surgery to reduce the risk of internal hernia.[37] Obstetricians must also be mindful of potential surgical complications and involve surgeons early in women with gastric bypass presenting with abdominal pain.[38]

5. Conclusion

The increased risk of maternal morbidity related to surgical complications after gastric bypass is concerning. There is currently no consensus regarding the type of bariatric procedure preferable to women planning a future pregnancy. Further studies are needed to establish if restrictive procedures should be used in preference to malabsorptive for women planning to have children[39].

Funding

North Bristol NHS Trust – researchers worked independently from the funder. There was no involvement of the funder nor the sponsor in the study design, data collection, analysis, writing of report or decision to publish.

Conflict of interest

None.

Acknowledgements

The authors would like to thank the UKOSS team for their input with data collection.

As specified and agreed in the UKOSS letter of understanding, the views expressed do not necessarily represent those of the UKOSS Steering Committee. We would also like to thank the clinicians reporting to UKOSS, without whom this work would not have been possible.

References

- [1] Enquiries C for M and C, Health CE into M and C. Maternal Obesity in the UK. 2010.
- [2] Giles J. Maternity Services Monthly Statistics 2018:1–41.
- [3] Heslehurst N, Rankin J, Wilkinson JR, Summerbell CD. A nationally representative study of maternal obesity in England, UK: trends in incidence and demographic inequalities in 619[[thinsp]]323 births, 1989[[ndash]]2007. *International Journal of Obesity and Related Metabolic Disorders : Journal of the International Association for the Study of Obesity* 2010;34:420–8. <https://doi.org/10.1038/ijo.2009.250>.
- [4] Knight M, Kurinczuk JJ, Spark P, Brocklehurst P, System UOS. Extreme obesity in pregnancy in the United Kingdom. *Obstet Gynecol* 2010;115:989–97. <https://doi.org/10.1097/aog.0b013e3181da8f09>.
- [5] Knight M, Bunch K, Tuffnell D, Jayakody H, Shakespeare J, Kotnis R, et al. Saving Lives, Improving Mothers' Care Lessons learned to inform future maternity care from the UK and Ireland Confidential Enquiries into Maternal Deaths and National Perinatal Epidemiology Unit 2018.
- [6] Aune D, Saugstad OD, Henriksen T, Tonstad S. Maternal Body Mass Index and the Risk of Fetal Death, Stillbirth, and Infant Death: A Systematic Review and Meta-analysis. *JAMA* 2014;311:1536–46. <https://doi.org/10.1001/jama.2014.2269>.
- [7] Sjöström L, Lindroos A-K, Peltonen M, Torgerson J, Bouchard C, Carlsson B, et al. Lifestyle, diabetes, and cardiovascular risk factors 10 years after bariatric surgery. *New England Journal of Medicine* 2004;351:2683–93. <https://doi.org/10.1056/nejmoa035622>.
- [8] The United Kingdom National Bariatric Surgery Registry 2016:1–15.
- [9] Kwong W, Tomlinson G, Feig DS. Maternal and neonatal outcomes after bariatric surgery; a systematic review and meta-analysis: do the benefits outweigh the risks? *American Journal of Obstetrics and Gynecology* 2018;218:573–80. <https://doi.org/10.1016/j.ajog.2018.02.003>.
- [10] Welbourn R, Hollyman M, Kinsman R, Dixon J, Liem R, Ottosson J, et al. Bariatric Surgery Worldwide: Baseline Demographic Description and One-Year Outcomes from the Fourth IFSO Global Registry Report 2018 2019:1–14. <https://doi.org/10.1007/s11695-018-3593-1>.
- [11] English WJ, DeMaria EJ, Brethauer SA, Mattar SG, Rosenthal RJ, Morton JM. American Society for Metabolic and Bariatric Surgery estimation of metabolic and bariatric procedures performed in the United States in 2016. *Surgery for Obesity and Related Diseases* 2018;14:259–63. <https://doi.org/10.1016/j.soard.2017.12.013>.
- [12] Galazis N, Docheva N, Simillis C, Nicolaidis KH. Maternal and neonatal outcomes in women undergoing bariatric surgery: a systematic review and meta-analysis. *European Journal of Obstetrics, Gynecology, and Reproductive Biology* 2014;181C:45–53. <https://doi.org/10.1016/j.ejogrb.2014.07.015>.
- [13] Johansson K, Cnattingius S, Näslund I, Roos N, Lagerros YT, Granath F, et al. Outcomes of pregnancy after bariatric surgery. *New England Journal of Medicine* 2015;372:814–24. <https://doi.org/10.1056/nejmoa1405789>.
- [14] Roos N, Neovius M, Cnattingius S, Lagerros YT, Sääf M, Granath F, et al. Perinatal outcomes after bariatric surgery: nationwide population based matched cohort study. *BMJ* 2013;347:f6460–f6460. <https://doi.org/10.1136/bmj.f6460>.

- [15] Ducarme G, Parisio L, Santulli P, Carbillon L, Mandelbrot L, Luton D. Neonatal outcomes in pregnancies after bariatric surgery: a retrospective multi-centric cohort study in three French referral centers. *The Journal of Maternal-Fetal & Neonatal Medicine* 2012. <https://doi.org/10.3109/14767058.2012.735723>.
- [16] Facchiano E, Iannelli A, Santulli P, Mandelbrot L, Msika S. Pregnancy after laparoscopic bariatric surgery: comparative study of adjustable gastric banding and Roux-en-Y gastric bypass. *Surg Obes Relat Dis* 2012;8:429–33. <https://doi.org/10.1016/j.soard.2011.06.006>.
- [17] Cornthwaite K, Jefferys A, Lenguerrand E, Haase A, Lynch M, Johnson A, et al. Pregnancy after weight loss surgery: a commentary. *BJOG: An International Journal of Obstetrics & Gynaecology* 2016;123:165–70. <https://doi.org/10.1111/1471-0528.13791>.
- [18] Knight M, Kurinczuk JJ, Tuffnell D, Brocklehurst P. The UK Obstetric Surveillance System for rare disorders of pregnancy. *BJOG: An International Journal of Obstetrics & Gynaecology* 2005;112:263–5. <https://doi.org/10.1111/j.1471-0528.2005.00609.x>.
- [19] Jefferys A, Lenguerrand E, Cornthwaite K, Johnson A, Lynch M, Hyde J, et al. Deflation versus maintained inflation of gastric band in pregnancy: A national cohort study. *Obstetric Medicine: The Medicine of Pregnancy* 2017;10:67–73. <https://doi.org/10.1177/1753495x16681200>.
- [20] Slater C, Morris L, Ellison J, Syed A. Nutrition in Pregnancy Following Bariatric Surgery. *Nutrients* 2017;9:1338–11. <https://doi.org/10.3390/nu9121338>.
- [21] Cole TJ, Williams AF, Wright CM. Revised birth centiles for weight, length and head circumference in the UK-WHO growth charts. *Annals of Human Biology* 2010;38:7–11. <https://doi.org/10.3109/03014460.2011.544139>.
- [22] Statistics O of N. Birth Summary Tables, England and Wales. 2019.
- [23] Zou G. A Modified Poisson Regression Approach to Prospective Studies with Binary Data. *American Journal of Epidemiology* 2004;159:702–6. <https://doi.org/10.1093/aje/kwh090>.
- [24] Buuren S van. mice v2. 30. RdocumentationOrg 2011.
- [25] Rubin DB. Inference and Missing Data. *Biometrika* 1976;63:581–92.
- [26] Mehta CR, Corporation PS-CS, 2003. Conditional versus unconditional exact tests for comparing two binomials. *ResearchgateNet* 2003.
- [27] O'Connor M, Kurinczuk J, Knight M. UKOSS Annual Report 2018. 2018:1–44.
- [28] Walters SJ. Consultants' forum: should post hoc sample size calculations be done? *Pharm Stat* 2009;8:163–9. <https://doi.org/10.1002/pst.334>.
- [29] Hammeken LH, Betsagoo R, Jensen AN, Sørensen AN, Overgaard C. Nutrient deficiency and obstetrical outcomes in pregnant women following Roux-en-Y gastric bypass: A retrospective Danish cohort study with a matched comparison group. *Eur J Obstet Gyn R B* 2017;216:56–60. <https://doi.org/10.1016/j.ejogrb.2017.07.016>.
- [30] MD AC, PhD GKM, MD MC, MD NL, PhD SMM, MD LM. Impact of bariatric surgery on fetal growth restriction: experience of a perinatal and bariatric surgery center. *The American Journal of Obstetrics & Gynecology* 2016:1–7. <https://doi.org/10.1016/j.ajog.2015.11.017>.
- [31] Lapolla A, Marangon M, Dalfrà MG, Segato G, Luca MD, Fedele D, et al. Pregnancy Outcome in Morbidly Obese Women Before and After Laparoscopic Gastric Banding. *Obesity Surgery* 2010;20:1251–7. <https://doi.org/10.1007/s11695-010-0199-7>.

- [32] Harreiter J, Schindler K, Bancher-Todesca D, Göbl C, Langer F, Prager G, et al. Management of Pregnant Women after Bariatric Surgery. *J Obes* 2018;2018:1–14. <https://doi.org/10.1155/2018/4587064>.
- [33] Shawe J, Ceulemans D, Akhter Z, Neff K, Hart K, Heslehurst N, et al. Pregnancy after bariatric surgery: Consensus recommendations for periconception, antenatal and postnatal care. *Obes Rev* 2019;20:1507–22. <https://doi.org/10.1111/obr.12927>.
- [34] Santulli P, Mandelbrot L, Facchiano E, Dussaux C, Ceccaldi P-F, Ledoux S, et al. Obstetrical and Neonatal Outcomes of Pregnancies following Gastric Bypass Surgery: A Retrospective Cohort Study in a French Referral Centre. *Obesity Surgery* n.d.;20:1501–8. <https://doi.org/10.1007/s11695-010-0260-6>.
- [35] Stenberg E, Chen R, Hildén K, Fall K. Pregnancy As a Risk Factor for Small Bowel Obstruction After Laparoscopic Gastric Bypass Surgery. *Annals of Surgery* 2018:1–5. <https://doi.org/10.1097/sla.0000000000003163>.
- [36] Moore KA, Ouyang DW, Whang EE. Maternal and fetal deaths after gastric bypass surgery for morbid obesity. *New England Journal of Medicine* 2004;351:721–2. <https://doi.org/10.1056/nejm200408123510722>.
- [37] Stenberg E, Szabo E, Ågren G, Ottosson J, Lancet RMT, 2016. Closure of mesenteric defects in laparoscopic gastric bypass: a multicentre, randomised, parallel, open-label trial. Elsevier n.d.
- [38] Wax JR, Pinette MG, Cartin A. Roux-en-Y gastric bypass-associated bowel obstruction complicating pregnancy-an obstetrician's map to the clinical minefield. *American Journal of Obstetrics and Gynecology* 2013;208:265–71. <https://doi.org/10.1016/j.ajog.2012.08.014>.
- [39] Akhter Z, Rankin J, Ceulemans D, Ngongalah L, Ackroyd R, Devlieger R, et al. Pregnancy after bariatric surgery and adverse perinatal outcomes: A systematic review and meta-analysis. *Plos Med* 2019;16:e1002866. <https://doi.org/10.1371/journal.pmed.1002866>.

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Tables and figures captions

Table 1 Summary of key studies comparing pregnancy outcomes following different types of bariatric surgery published in the last decade

Figure 1 Types of bariatric surgery: a) Gastric banding: an adjustable band is placed around the upper portion of the stomach to create a small upper pouch and a narrow opening into the main body of the stomach. b) Gastric bypass: Food intake is restricted by creating a small pouch at the gastric fundus, whilst the main body of the stomach, duodenum and length of jejunum are bypassed (Roux-en-Y). c) Sleeve gastrectomy: the greater curvature of the stomach is removed, resulting in a narrow gastric sleeve.

Reproduced from : ‘What is the most effective operation for adults with severe and complex obesity’ J Blazeby, J Byrne, R Welbourn; 348; 4-5;2014 with permission from BMJ Publishing Group Ltd

Figure 2 Flowchart of the gastric banding and bypass / sleeve cohorts

Table 2 Participant characteristics

Figure 3 Weight change during pregnancies after gastric band, bypass and sleeve

Table 3 Comparison of maternal outcomes between pregnancies after gastric banding, bypass and sleeve

Table 4 Comparison of perinatal outcomes between pregnancies after gastric banding, bypass and sleeve

Supplementary tables and figure legends

Table S1 Participant characteristics for subgroups of bypass cases

Table S2 Analysis of weight gain and BMI change including missing data imputation

Table S3 Comparison of all maternal outcomes after gastric band, bypass and sleeve

Table S4 Comparison of all perinatal outcomes after gastric band, bypass and sleeve

Table S5 Subgroup analysis of maternal outcomes according to type of bypass surgery

Table S6 Subgroup analysis of perinatal outcomes according to type of bypass surgery

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1 Summary of key studies comparing pregnancy outcomes following different types of bariatric surgery published in the last decade

Study	Study design	Period of observation	No. cases	Comparison ^a	Types of Surgery	Distinct or conflated comparisons ^b
This study	Cohort ^c	5 years	290	Bypass vs banding vs sleeve	Bypass / Banding / Sleeve	Distinct
Shai et al 2013	Case-control	22 years ^d	326	Different obese women without BS	Bariatric surgery (undifferentiated)	Conflated
Aricha-Tamir et al 2012	Case-control	20 years ^d	144	Same women before BS	Bypass / Banding / Gastroplasty	Conflated
Grandfils et al 2019	Cohort	14 years ^d	337	Insufficient vs adapted vs excessive gestational weight gain	Bypass / Banding / Sleeve	Conflated
Lapolla et al 2010	Cohort	12 years ^e	83	A. Different obese women without BS B. Same women before BS	Banding	Conflated
Jonasson et al 2011	Cohort	10 years ^f	681	A. Different women without BS B. Same women before BS	Bypass / Banding / Gastroplasty	Conflated
Kjaer et al 2013	Case-control	6 years ^d	339	Different women without BS (BMI matched)	Bypass / Banding	Conflated
Ansalem et al 2014	Case-control	6 years ^d	109	Same women before BS	Banding / Gastroplasty	Conflated
Jonasson et al 2015	Case-control	5 years ^d	670	Different women without BS (BMI matched)	Bypass / Banding / Other	Conflated
Bennett et al 2010	Cohort	4 years ^g	585	Same women before BS	Bypass / Banding / Other	Conflated
Burke et al 2010	Cohort	4 years ^e	354	Different obese women without BS	Bypass / Banding	Conflated
Lesko et al 2012	Case-control	4 years ^d	70	Different women without BS (BMI matched)	Bypass / Banding	Conflated
Belogolovkin et al 2012	Cohort	3 years ^d	293	Different obese women without BS	Bariatric surgery (undifferentiated)	Conflated
Balestrin et al 2019	Case-control	2 years ^d	93	Different obese women without BS	Bypass / Restrictive procedures	Conflated
Parker et al 2015	Cohort	1 year ^d	1585	Different obese women without BS	Bariatric surgery (undifferentiated)	Conflated
Shiner et al 2009	Cohort	20 years ^e	449	Between different surgery types A. Bypass vs banding vs gastroplasty B. Malabsorptive vs restrictive	Bypass / Banding / Gastroplasty	Distinct
Roos et al 2013	Case-control	17 years ^d	2,562	A. Different women without BS (BMI matched) B. Between subgroups of different surgery types	Bypass / Banding / Gastroplasty	Distinct
Berlac et al 2014	Cohort	15 years ^e	415	Different women without BS (BMI matched)	Bypass	Distinct
Gonzalez et al 2014	Cohort	14 years ^d	168	Restrictive vs malabsorptive	Bypass / Banding / Sleeve / Gastroplasty / Biliopancreatic diversion	Grouped according to malabsorptive vs restrictive
Goupaye et al 2018	Cohort	13 years ^d	123	Bypass vs sleeve	Bypass / Sleeve	Distinct
Chevrot et al 2016	Case-control	9 years ^e	139	A. Different women without BS (BMI matched) B. Malabsorptive vs restrictive	Bypass / Banding / Sleeve	Grouped according to malabsorptive vs restrictive
Watanabe et al 2019	Case-control	9 years ^e	24	Bypass vs banding vs sleeve	Bypass / Banding / Sleeve	Distinct
Sanulli et al 2010	Cohort	6 years ^d	24	Different women without BS (BMI matched)	Bypass	Distinct
Ducarme et al 2012	Cohort	5 years ^d	94	Bypass vs banding	Bypass / Banding	Distinct
Facchiano et al 2012	Cohort	4 years ^d	42	Bypass vs banding	Bypass / Banding	Distinct
Hammeken et al 2017	Cohort	3 years ^d	151	Different women without BS (BMI matched)	Bypass	Distinct

^aBS = Bariatric surgery, ^bPregnancy outcomes analyzed with different types of procedure grouped together (conflated) or by procedure type (distinct). ^cProspective study. All other studies retrospective.
^dDuring which births occurred, ^eDuring which surgeries performed, ^fDuring which women born, ^gDuring which claims made

Table 2 Participant characteristics

		Overall (n=268)	Type of surgery		
			Gastric Banding (n= 107, 40%)	Gastric Bypass (n=132, 49%)	Sleeve Gastrectomy (n=29,11%)
Maternal age (years)	Mean (SD)	32.9 (5.2)	31.8 (4.9)	33.5 (5.2)	34.2 (5.8)
	Missing (n)	0	0	0	0
BMI at booking	Mean (SD)	34.5 (7.0)	36.4 (7.3)	33.6 (6.8)	32.0 (5.3)
	Missing (n)	2	1	0	1
Weight at booking (kg)	Mean (SD)	96 (20.1)	101.6 (20.9)	92.5 (19.7)	90.2 (15)
	Missing (n)	2	1	0	1
Parity^a n (%)	Nulliparous	99	50 (47)	37 (28)	12 (41)
	Parous	166	57 (53)	92 (70)	17 (59)
	Grandparous	3	0 (0)	3 (2)	0 (0)
	Missing (n)	0	0	0	0
Ethnicity n (%)	White	246	100 (93)	122 (92)	24 (83)
	Asian	11	4 (4)	6 (5)	1 (3)
	Black African	3	1 (1)	2 (2)	0 (0)
	Mixed, other	8	2 (2)	2 (2)	4 (14)
	Missing (n)	0	0	0	0
Employment n (%)	Employed	178	78 (73)	84 (64)	16 (55)
	Unemployed	82	26 (24)	44 (33)	12 (41)
	Missing (n)	8	3	4	1
Smoking during pregnancy n (%)	Yes	38	15 (14)	18 (14)	5 (17)
	No	227	92 (86)	112 (85)	23 (79)
	Missing (n)	3	0	2	1
Pre-existing Diabetes n (%)	Yes	33	7 (7)	20 (15)	6 (21)
	No	235	100 (93)	112 (85)	23 (79)
	Missing (n)	0	0	0	0
Pre-existing hypertension n (%)	Yes	26	11 (10)	12 (9)	3 (10)
	No	242	96 (90)	120 (91)	26 (90)
	Missing (n)	0	0	0	0
^a Nulliparous = 0 previous maternities; Parous = 1-4 previous maternities; Grandparous = 5-8 previous maternities Inferential measures and significance tests not performed, as per STROBE guidelines					

Table 3 Comparison of maternal outcomes between pregnancies after gastric banding, bypass and sleeve

		Gastric banding (n=107)	Gastric bypass (n=132)	Sleeve gastrectomy (n=29)	Banding vs Bypass				Banding vs Sleeve		Bypass vs Sleeve	
					Unadjusted analyses		Adjusted analyses ^a		Difference in proportion (95%CI) ^b	p value	Difference in proportion (95%CI) ^b	p value
					RR (95%CI)	p value	RR (95%CI)	p value				
Pre-eclampsia (n, %)	Yes	3 (2.8)	4 (3.0)	0 (0)	0.90 (0.21,3.94)	0.91	1.39 (0.48,4.00)	0.54	0.03 (-0.10,0.08)	0.47	0.03 (-0.10,0.07)	0.43
	No	99 (92.5)	119 (90.2)	29 (100)								
	Missing	5 (4.7)	9 (6.8)	0 (0)								
Gestational Diabetes ^c (n, %)	Yes	7 (7)	18 (16)	1 (4)	0.42 (0.18,0.96)	0.03	0.35 (0.13,0.92)	0.03	0.03 (-0.15,0.11)	0.82	0.14 (-0.06,0.24)	0.09
	No	90 (90)	87 (78)	22 (96)								
	Missing	3 (3)	7 (6)	0 (0)								
Anaemia (n, %)	Yes	17 (15.9)	51 (38.6)	6 (21)	0.40 (0.25,0.65)	<0.001	0.44 (0.27,0.72)	0.002	-0.03 (-0.21,0.11)	0.74	0.20 (0.01,0.35)	0.04
	No	85 (79.4)	72 (54.5)	23 (79)								
	Missing	5 (4.7)	9 (6.8)	0 (0)								
Surgical complications (n, %)	Yes	1 (0.9)	15 (11.4)	0 (0)	0.08 (0.01,0.60)	0.001	0.08 (0.008,0.70)	0.03	0.009 (-0.12,0.05)	0.73	0.11 (-0.03,0.18)	0.06
	No	101 (94.4)	108(81.8)	29 (100)								
	Missing	5 (4.7)	9 (6.8)	0 (0)								
Major medical complications (n, %)	Yes	3 (2.8)	5 (3.8)	0 (0)	0.61 (0.16,2.40)	0.73	0.63 (0.10,3.71)	0.60	0.03 (-0.10,0.08)	0.46	0.04 (-0.07,0.10)	0.28
	No	99 (92.5)	118 (89.4)	29 (100)								
	Missing	5 (4.7)	9 (6.8)	0 (0)								
Cases with missing observations (outcome or covariates) removed prior to analysis.												
^a Adjusted analyses using Poisson regression model.												
^b Unadjusted analyses using Barnard exact test. Adjustment not possible due to sparse and unbalanced data.												
^c Cases with pre-existing diabetes excluded.												

Table 4 Comparison of perinatal outcomes between pregnancies after gastric banding, bypass and sleeve

		Gastric banding (n=107)	Gastric bypass (n=132)	Sleeve gastrectomy (n=29)	Banding vs Bypass				Banding vs Sleeve		Bypass vs Sleeve	
					Unadjusted analyses		Adjusted analyses		Mean difference / Difference in proportion (95%CI) ^{b,c}	p value	Mean difference / Difference in proportion (95%CI) ^{b,c}	p value
					Mean difference / RR (95%CI) ^a	p value	Mean difference / RR (95%CI) ^a	p value				
Birth weight (g) ^d	Mean	3380	3159	3199	221 (64,380)	0.006	260 (125,395)	<0.001	138 (-97,373)	0.25	-90 (-320,140)	0.44
	SD	641	530	487								
	Missing	7 (6.5)	10 (7.6)	3 (10)								
Small for gestational age ^e (n, %)	Yes	7 (7)	14 (11)	1 (3)	0.66 (0.27,1.58)	0.48	0.46 (0.16,1.33)	0.15	0.03 (-0.11,0.11)	0.60	0.07 (-0.07,0.15)	0.26
	No	98 (92)	117 (89)	28 (97)								
	Missing	2 (2)	1 (0)	1 (0)								
Large for gestational age ^e (n, %)	Yes	22 (21)	6 (5)	1 (3)	4.27 (1.79,10.17)	<0.001	4.74 (1.54,14.6)	0.006	0.17 (0.02,0.27)	0.03	0.01 (-0.14,0.07)	0.89
	No	83 (78)	125 (95)	28 (97)								
	Missing	2 (2)	1 (0)	1 (0)								
Preterm birth ^e (n, %)	Yes	14 (13)	12 (9)	4 (14)	1.54 (0.73,3.24)	0.29	2.27 (1.02,5.03)	0.04	-0.006 (-0.18,0.11)	0.98	-0.05 (-0.22,0.06)	0.50
	No	92 (86)	119 (90)	25 (86)								
	Missing	1 (1)	1 (1)	0 (0)								
Stillbirth ^e (n, %)	Yes	1 (0.9)	1 (0.8)	0 (0)	-		-		-		-	
	No	104 (97.2)	131 (99.2)	29 (100)								
	Missing	2 (0)	0 (0)	0 (0)								

Cases with missing observations (outcome or covariates) removed prior to analysis

^aMean difference reported for continuous outcome (birth weight) and RR reported for categorical outcomes

^bMean difference reported for continuous outcome (birth weight) and difference in proportion reported for categorical outcomes.

^cUnadjusted analyses using Barnard exact test. Adjustment not possible due to sparse and unbalanced data.

^dAdjusted analyses for banding vs bypass comparison using generalised least square model

^eAdjusted analyses for banding vs bypass comparison using Poisson regression model

Supplementary Tables

Table S1 Participant characteristics for subgroups of bypass cases

		All Bypass cases (n=132)	Subgroup of bypass surgery			
			Roux-en-Y (n=67, 42%)	Loop gastric bypass (n=2, 6%)	Laparoscopic gastric bypass (unspecified) (n=8, 1%)	Bypass (type unknown) (n=55, 51%)
Maternal age (years)	Mean (SD)	33.5 (5.2)	33 (4.4)	32.5 (2.1)	35.9 (6.1)	33.8 (5.9)
	Missing (n)	0	0	0	0	0
Body Mass Index at booking	Mean (SD)	33.6 (6.8)	34.6 (6.8)	29 (2.8)	32.2 (9.2)	32.8 (6.3)
	Missing (n)	0	0	0	0	0
Weight at booking (Kg)	Mean (SD)	92.5 (19.7)	96.1 (18.6)	82.3 (1.8)	87.7 (28)	89.3 (19.6)
	Missing (n)	0	0	0	0	0
Parity ^a n (%)	Primiparous	37 (28)	21 (31)	0 (0)	1 (13)	15 (27)
	Multiparous	92 (70)	45 (67)	2 (100)	7 (88)	38 (69)
	Grandparous	3 (2)	1 (1)	0 (0)	0 (0)	2 (4)
	Missing (n)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Ethnicity n (%)	White	122(92)	64 (96)	2 (100)	6 (86)	50 (92)
	Asian	6 (5)	2 (3)	0 (0)	1 (13)	3 (5)
	Black African	2 (2)	1 (1)	0 (0)	1(13)	0 (0)
	Mixed, other	2 (2)	0 (0)	0 (0)	0 (0)	2 (4)
	Missing (n)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Employment n (%)	Employed	84 (64)	46 (69)	0 (0)	5 (63)	33 (60)
	Unemployed	44 (33)	20 (30)	2 (100)	2 (3)	20 (36)
	Missing (n)	4 (3)	1 (1)	0 (0)	1 (13)	2 (4)
Smoking during pregnancy n (%)	Yes	18 (14)	7 (10)	1 (50)	1 (13)	9 (16)
	No	112 (85)	58 (87)	1 (50)	7 (88)	46 (84)
	Missing (n)	2 (2)	2 (3)	0 (0)	0 (0)	0 (0)
Pre-existing Diabetes n (%)	Yes	20 (15)	11 (16)	0 (0)	1 (13)	8 (15)
	No	112 (85)	56 (84)	2 (100)	7 (88)	47 (85)
	Missing (n)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Maternal Anaemia n (%)	Yes	54 (41)	25 (37)	1 (50)	3 (38)	25 (45)
	No	77 (58)	41 (61)	1 (50)	5 (63)	30 (55)
	Missing (n)	1 (1)	1 (1)	0 (0)	0 (0)	0 (0)
Pre-existing hypertension n (%)	Yes	12 (9)	3 (4)	0 (0)	2 (3)	7 (13)
	No	120 (91)	64 (96)	2 (100)	6 (86)	48 (87)
	Missing (n)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)

^a Primiparous= 0 previous maternities; Parous = 1-4 previous maternities; Grandparous = 5-8 previous maternities

Table S2 Analysis of weight gain and BMI change including missing data imputation

			Estimated weight gain and BMI change in each group ^a			Difference in weight gain and BMI change between groups					
			Gastric banding Estimate (95% CI)	Gastric bypass Estimate (95% CI)	Sleeve gastrectomy Estimate (95% CI)†	Banding vs Bypass ^b		Banding vs Sleeve ^c		Bypass vs Sleeve ^c	
						Difference (95% CI)	p value	Difference (95% CI)	p value	Difference (95% CI)	p value
Gestational weight gain	Original unadjusted model		10.2 (8.1,12.3)	7.2 (5.3,9.0)	8.2 (2.4,14.0)	3.0 (0.2,5.4)	0.03	2.1 (-4.0,8.6)	0.48	-1.1 (-5.4,3.2)	0.62
	Original adjusted model		9.9 (7.7,12.0)	7.3 (5.3,9.2)	-	2.6 (-0.3,5.5)	0.08	-		-	
	Missing at random (MAR) ^c		10.9 (8.2,13.5)	7.2 (5.0,9.4)	7.1 (0.6,13.5)	3.6 (0.2,7.0)	0.04	4.3 (-2.6,11.2)	0.21	0.7 (-6.0,7.4)	0.83
	Missing not at random (MNAR) ^d	I	8.5 (5.8,11.3)	1.8 (-0.7,4.2)	3.3 (-2.6,9.2)	6.8 (3.1,10.4)	<0.001	4.8 (-0.9,10.5)	0.10	-1.9 (-8.4,4.5)	0.56
		II	9.9 (7.7,12.0)	7.3 (5.3,9.2)	6.2 (0.9,11.6)	2.6 (-0.3,5.5)	0.08	9.5 (3.2,15.7)	0.003	-0.4 (-6.3,5.5)	0.90
		III	20.9 (17.7,24.1)	11.1 (8.3,14.0)	12.2 (6.7,17.8)	9.7 (5.5,14.1)	<0.001	9.3 (1.7,16.9)	0.02	-0.9 (-7.1,5.2)	0.76
BMI change	Original unadjusted model		3.6 (2.8,4.3)	2.6 (2.0,3.3)	3.1 (1.0,5.0)	0.9 (-0.1,1.9)	0.07	0.5 (-1.8,2.8)	0.67	-0.5 (-2.1,1.0)	0.50
	Original adjusted model		3.5 (2.7,4.3)	2.7 (2.0,3.4)	-	0.8 (-0.2,1.8)	0.14	-		-	
	Missing at random (MAR) ^d		3.6 (2.9,4.3)	2.5 (1.9,3.1)	2.8 (0.9,4.8)	1.1 (0.2,2.0)	0.02	0.9 (-1.2,2.9)	0.42	-0.4 (-1.9,1.1)	0.62
	Missing not at random (MNAR) ^e	I	3.3 (2.3,4.3)	0.7 (-0.4,1.6)	2.3 (0.5,4.2)	2.6 (1.2,4.0)	<0.001	0.8 (-1.2,2.8)	0.41	-1.4 (-3.4,0.7)	0.19
		II	5.1 (4.1,6.1)	2.4 (1.5,3.3)	4.2 (2.4,5.9)	2.7 (1.3,4.0)	<0.001	0.7 (-1.4,2.9)	0.50	-1.6 (-3.6,0.3)	0.09
		III	6.7 (5.6,7.8)	3.4 (2.4,4.4)	6.6 (4.8,8.5)	3.3 (1.8,4.8)	<0.001	-0.1 (-2.6,2.4)	0.94	-3.1 (-5.2, -1.1)	0.003
^a Difference between 1 st and 3 rd trimester measurements											
^b Adjusted for age, parity, ethnicity, early pregnancy BMI and weight, smoking status, employment status and pre-existing diabetes.											
^c Adjustment not performed for sleeve gastrectomy											
^d Multiple imputation using chain equations, under MAR assumption (adjusted model for banding vs bypass). Twenty imputed datasets created and combined using Rubin's rule.											
^e Sensitivity analysis using single value imputation, under MNAR assumption (adjusted model for banding vs bypass): women with missing weight / BMI measurements at the third trimester visit were assumed to have a weight / BMI equal to the group-specific 25 th (I), 50 th (II), and 75 th (III) percentile of observed weights / BMIs at the third trimester visit.											

Table S3 Comparison of all maternal outcomes after gastric band, bypass and sleeve

		Gastric banding (n=107)	Gastric bypass (n=132)	Sleeve gastrectomy (n=29)	Banding vs Bypass				Banding vs Sleeve		Bypass vs Sleeve	
					Unadjusted RR (95%CI)	Unadjusted p value	Adjusted RR (95%CI) ^a	Adjusted p value	Difference in proportion (95%CI) ^b	p value	Difference in proportion (95%CI) ^b	p value
Gestational hypertension^c (n, %)	Yes	5 (5)	18(62)	0 (0)	2.19 (0.61,7.92)	0.32	2.19 (0.61,7.92)	0.32	0.05 (-0.08,0.12)	0.26	0.02 (-0.11,0.07)	0.50
	No	86 (90)	0 (0)	26 (100)								
	Missing	5 (5)	26 (100)	0 (0)								
Pre-eclampsia (n, %)	Yes	3 (2.8)	0 (0)	0 (0)	1.39 (0.48,4.00)	0.54	1.39 (0.48,4.00)	0.54	0.03 (-0.10,0.08)	0.47	0.03 (-0.10,0.07)	0.43
	No	99 (92.5)	0 (0)	29 (100)								
	Missing	5 (4.7)	29 (100)	0 (0)								
Gestational Diabetes^d (n, %)	Yes	7 (7)	0 (0)	1 (4)	0.35 (0.13,0.92)	0.03	0.35 (0.13,0.92)	0.03	0.03 (-0.15,0.11)	0.82	0.14 (-0.06,0.24)	0.09
	No	90 (90)	1 (4)	22 (96)								
	Missing	3 (3)	22 (96)	0 (0)								
Anaemia (n, %)	Yes	17 (15.9)	0 (0)	6 (21)	0.44 (0.27,0.72)	0.002	0.44 (0.27,0.72)	0.002	-0.03 (-0.21,0.11)	0.74	0.20 (0.01,0.35)	0.04
	No	85 (79.4)	6 (21)	23 (79)								
	Missing	5 (4.7)	23 (79)	0 (0)								
Induced labor (n, %)	Yes	39 (36.4)	0 (0)	10 (34)	0.78 (0.55,1.11)	0.16	0.78 (0.55,1.11)	0.16	0.04 (-0.17,0.22)	0.73	0.06 (-0.14,0.23)	0.58
	No	61 (57)	10 (34)	19 (66)								
	Missing	7 (6.5)	19 (66)	0 (0)								
Mode of birth (n, %)	Caesarean	47 (43.9)	0 (0)	13 (45)	1.26 (0.91,1.73)	0.16	1.26 (0.91,1.73)	0.16	0.05 (-0.16,0.25)	0.66	-0.06 (-0.26,0.13)	0.55
	Vaginal	53 (49.5)	13 (45)	16 (55)								
	Missing	7 (6.5)	16 (55)	0 (0)								
Surgical complications (n, %)	Yes	1 (0.9)	0 (0)	0 (0)	0.08 (0.008,0.70)	0.03	0.08 (0.008,0.70)	0.03	0.009 (-0.12,0.05)	0.73	0.11 (-0.03,0.18)	0.06
	No	101 (94.4)	0 (0)	29 (100)								
	Missing	5 (4.7)	29 (100)	0 (0)								
Major medical complications (n, %)	Yes	3 (2.8)	0 (0)	0 (0)	0.63 (0.10,3.71)	0.60	0.63 (0.10,3.71)	0.60	0.03 (-0.10,0.08)	0.46	0.04 (-0.07,0.10)	0.28
	No	99 (92.5)	0 (0)	29 (100)								
	Missing	5 (4.7)	29 (100)	0 (0)								
Maternal death (n, %)	Yes	0 (0)	0 (0)	0 (0)	-	-	-	-	-	-	-	-
	No	107 (100)	0 (0)	29 (100)								
	Missing	0 (0)	29 (100)	0 (0)								

Cases with missing observations (outcome or covariates) removed prior to analysis

^aAdjusted analyses using Poisson regression model

^bUnadjusted analyses using Barnard exact test. Adjustment not possible due to sparse and unbalanced data

^cCases with pre-existing hypertension excluded

^dCases with pre-existing diabetes excluded

Table S4 Comparison of all perinatal outcomes after gastric band, bypass and sleeve

		Gastric banding (n=107)	Gastric bypass (n=132)	Sleeve gastrectomy (n=29)	Banding vs Bypass				Banding vs Sleeve		Bypass vs Sleeve	
					Mean difference / Unadjusted RR (95%CI)*	Unadjusted p value	Mean difference / Adjusted RR (95%CI) ^a	Adjusted p value	Mean difference / Difference in proportion (95%CI) ^{a, b}	p value	Mean difference / Difference in proportion (95%CI) ^{a, b}	p value
Birth weight (g)^c	Mean	3380	3159	3199	221 (64,380)	0.006	260 (125,395)	<0.001	138 (-97,373)	0.25	-90 (-320,140)	0.44
	SD	641	530	487								
	Missing	7 (6.5)	10 (7.6)	3 (10)								
Gestational age at birth^d (n, %)	Median	39	39	38.8	-0.1 (-0.5,0.6)	0.57	-0.7 (-1.43,0.06)	0.07	0.23 (-0.74,1.20)	0.64	0.32 (-0.62,1.27)	0.49
	IQR	(38,40)	(38.1,38.9)	(37.5,39.4)								
	Missing	6 (5.6)	10 (7.6)	3(10)								
Low birth weight^e (n, %)	Yes	8 (7.5)	8 (6.1)	3 (10)	1.2 (0.47,3.13)	0.79	1.89 (0.66,1.71)	0.23	-0.03 (-0.20,0.07)	0.82	-0.03 (-0.20,0.05)	0.69
	No	92 (86)	114 (86.4)	26 (90)								
	Missing	7 (6.5)	10 (7.6)	0 (0)								
Macrosomia^e (n, %)	Yes	17 (15.9)	6 (4.5)	1 (3)	3.46 (1.42,8.44)	0.004	3.59 (1.45,8.90)	0.005	0.14 (-0.03,0.23)	0.06	0.01 (-0.14,0.07)	0.88
	No	83 (77.6)	116 (87.9)	28 (97)								
	Missing	7 (6.5)	10 (7.6)	0 (0)								
Small for gestational age (SGA)^e (n, %)	Yes	7 (6.5)	13 (9.8)	1 (3)	0.66 (0.27,1.58)	0.48	0.46 (0.16,1.33)	0.15	0.03 (-0.11,0.11)	0.60	0.07 (-0.07,0.15)	0.26
	No	93 (86.9)	109 (82.6)	28 (97)								
	Missing	7 (6.5)	10 (7.6)	0 (0)								
Large for gestational age (LGA)^e (n, %)	Yes	21 (19.6)	6 (4.5)	1 (3)	4.27 (1.79,10.17)	0.0003	4.74 (1.54,14.6)	0.006	0.17 (0.02,0.27)	0.03	0.01 (-0.14,0.07)	0.89
	No	79 (73.8)	116 (87.9)	28 (97)								
	Missing	7 (5.5)	10 (7.6)	0 (0)								
Preterm birth^e (n, %)	Yes	14 (13.1)	11 (8.3)	4 (14)	1.54 (0.73,3.24)	0.29	2.27 (1.02,5.03)	0.04	-0.006 (-0.18,0.11)	0.98	-0.05 (-0.22,0.06)	0.50
	No	87 (81.3)	111 (84.1)	25 (86)								
	Missing	6 (5.6)	10 (7.6)	0 (0)								
Neonatal unit admission^e (n, %)	Yes	11 (10.3)	11 (8.3)	1 (3)	1.23 (0.56,2.72)	0.66	0.62 (0.26,1.51)	0.29	0.07 (-0.07,0.15)	0.26	0.05 (-0.10,0.12)	0.44
	No	88 (82.2)	111 (84.1)	28 (97)								
	Missing	8 (7.5)	10 (7.6)	0 (0)								
Low Apgar score^e (n, %)	Yes	2 (1.9)	2 (1.5)	1 (3)	1.3 (0.19,9.06)	0.80	-	-	-0.02 (-0.16,0.04)	0.82	-0.02 (-0.17,0.03)	0.59
	No	93 (86.9)	119 (90.2)	27 (93)								
	Missing	12 (11.2)	11 (8.3)	1 (3)								
Congenital abnormalities^e (n, %)	Yes	3 (2.8)	3 (2.3)	0 (0)	1.89 (0.32,11.10)	0.65	-	-	0.03 (-0.10,0.08)	0.47	0.02 (-0.10,0.06)	0.50
	No	95 (88.8)	119 (90.2)	29 (100)								
	Missing	9 (8.4)	10 (7.6)	0 (0)								
Stillbirth^e (n, %)	Yes	1 (0.9)	1 (0.8)	0 (0)	-	-	-	-	-	-	-	-
	No	104 (97.2)	131 (99.2)	29 (100)								
	Missing	0 (0)	0 (0)	0 (0)								

Cases with missing observations (outcome or covariates) removed prior to analysis

^aMean difference reported for continuous outcomes and RR or difference in proportion reported for remaining categorical outcomes

^bUnadjusted analyses using Barnard exact test. Adjustment not possible due to sparse and unbalanced data

^cAdjusted analyses using generalised least square model for banding versus bypass comparisons

^dAdjusted analyses using quantile regression model for banding versus bypass comparisons

^eAdjusted analyses using Poisson regression model for banding versus bypass comparisons

Table S5 Subgroup analysis of maternal outcomes according to type of bypass surgery


		Roux-en-Y (n=67, 42%)	Unspecified bypass (n=55, 51%)	Mean diff (95% CI) / Difference in proportion (95% CI) ^a	p value (unadjusted)
Gestational weight gain (kg) ^b	Mean	7.5	6.4	-0.7 (-4.9, 3.5)	0.74
	SD	7.1	6.0		
	Missing (n, %)	31 (46)	29 (53)		
BMI change ^b	Mean	2.7	2.4	-0.5 (-1.5,0.5)	0.34
	SD	2.6	2.2		
	Missing (n, %)	31 (46)	29 (53)		
Gestational hypertension ^c (n, %)	Yes	1 (2)	0 (0)	0.02 (-0.06, 0.08)	0.52
	No	63 (98)	48 (100)		
	Missing	0 (0)	0 (0)		
Pre-eclampsia (n, %)	Yes	1 (1)	2 (4)	-0.02 (-0.11,0.05)	0.59
	No	66(99)	53 (96)		
	Missing	0 (0)	0 (0)		
Gestational diabetes ^d (n, %)	Yes	14 (25)	7 (15)	0.10 (-0.06,0.26)	0.22
	No	42 (75)	40 (85)		
	Missing	0 (0)	0 (0)		
Anaemia (n, %)	Yes	25 (37)	25 (45)	-0.08 (-0.25,0.10)	0.41
	No	41 (62)	30 (55)		
	Missing	1 (1)	0 (0)		
Induced labor (n, %)	Yes	29 (43)	21(38)	0.05 (-0.13,0.22)	0.58
	No	38 (57)	34 (62)		
	Missing	0 (0)	0 (0)		
Mode of birth (n, %)	Caesarean	23 (34)	25 (45)	-0.11 (-0.29,0.06)	0.21
	Vaginal	44 (66)	30 (55)		
	Missing	0 (0)	0 (0)		
Surgical complications (n, %)	Yes	10 (15)	5 (9)	0.06 (-0.07,0.18)	0.36
	No	57 (85)	50 (91)		
	Missing	0 (0)	0 (0)		
Major medical complications (n, %)	Yes	4 (6)	2 (4)	0.02 (-0.07,0.11)	0.63
	No	63 (94)	53 (96)		
	Missing	0 (0)	0 (0)		
Maternal death (n, %)	Yes	1 (1)	0 (0)	0.015 (-0.05,0.08)	0.51
	No	66 (99)	55 (100)		
	Missing	0 (0)	0 (0)		

^aMean difference reported for continuous outcomes and difference in proportion reported for categorical outcomes
^bMean difference reported after multiple imputation for missing data under missing at random (MAR) assumption
^cCases with pre-existing hypertension excluded
^dCases with pre-existing diabetes excluded

Table S6 Subgroup analysis of perinatal outcomes according to type of bypass surgery

		Roux-en-Y (n=67, 42%)	Unspecified bypass (n=55, 51%)	Mean diff (95% CI) / Difference in proportion (95% CI) ^a	p value (unadjusted)
Birth weight (g)	Mean	3194	3116	-77 (-256,104)	0.40
	SD	384	619		
	Missing	0 (0)	0 (0)		
Low birth weight (n, %)	Yes	1 (1)	6 (11)	-0.05 (-0.12,0.001)	0.06
	No	99 (99)	89 (89)		
	Missing	0 (0)	0 (0)		
Macrosomia (n, %)	Yes	0 (0)	4 (0)	-0.07 (-0.17, -0.01)	0.03
	No	67 (100)	51 (93)		
	Missing	0 (0)	0 (0)		
Small for gestational age (SGA) (n, %)	Yes	6 (9)	6 (11)	-0.018 (-0.14,0.10)	0.78
	No	60 (90)	49 (89)		
	Missing	1 (0)	0 (0)		
Large for gestational age (LGA) (n, %)	Yes	2 (3)	3 (5)	-0.02 (-0.12,0.06)	0.63
	No	64 (96)	52 (95)		
	Missing	1 (1)	0 (0)		
Gestational age at birth (n, %)	Median	39	38.9	-0.15 (-0.67,0.37)	0.56
	IQR	(38.1,40)	(38.2,40)		
	Missing	1 (1)	0 (0)		
Preterm birth (n, %)	Yes	4 (6)	7 (13)	-0.07 (-0.19,0.04)	0.23
	No	62 (93)	48 (87)		
	Missing	1 (1)	0 (0)		
Neonatal Unit admission (n, %)	Yes	5 (93)	5 (89)	-0.02 (-0.13,0.10)	0.76
	No	62 (7)	49 (9)		
	Missing	0 (0)	1 (2)		
Low Apgar score (n, %)	Yes	0 (0)	1 (2)	-0.02 (-0.10,0.04)	0.45
	No	67 (100)	53 (96)		
	Missing	0 (0)	1 (2)		
Congenital abnormalities (n, %)	Yes	2 (3)	1 (2)	0.01 (-0.07,0.10)	0.82
	No	65 (97)	53 (96)		
	Missing	0 (0)	1 (2)		
Stillbirth (n, %)	Yes	0 (0)	1(2)	-0.02 (-0.10,0.04)	0.45
	No	67 (100)	54 (98)		
	Missing	0 (0)	0 (0)		
Perinatal death (n, %)	Yes	1 (0.9)	1 (0.8)	-0.02 (-0.10,0.04)	0.45
	No	104 (97.2)	131 (99.2)		
	Missing	2 (1.9)	0 (0)		

^aMean difference reported for continuous outcome (birth weight) and difference in proportion reported for categorical outcomes







Figure 3

[Click here to access/download;Figure;Figure 3.png](#)



Declaration of interests

☒ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☐The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

STROBE Statement—Checklist of items that should be included in reports of *cohort studies*

	Item No	Recommendation	Page No
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	1
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	2
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	4
Objectives	3	State specific objectives, including any prespecified hypotheses	5
Methods			
Study design	4	Present key elements of study design early in the paper	5
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	5
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up (b) For matched studies, give matching criteria and number of exposed and unexposed	5
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	6
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	5-6
Bias	9	Describe any efforts to address potential sources of bias	7-8
Study size	10	Explain how the study size was arrived at	
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	7-8
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding (b) Describe any methods used to examine subgroups and interactions (c) Explain how missing data were addressed (d) If applicable, explain how loss to follow-up was addressed (e) Describe any sensitivity analyses	7-8 8 7-8 8
Results			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed (b) Give reasons for non-participation at each stage (c) Consider use of a flow diagram	9 Fig 2
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders (b) Indicate number of participants with missing data for each variable of interest (c) Summarise follow-up time (eg, average and total amount)	9 9
Outcome data	15*	Report numbers of outcome events or summary measures over time	9-12

Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included (b) Report category boundaries when continuous variables were categorized (c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	9-12
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	12
Discussion			
Key results	18	Summarise key results with reference to study objectives	12
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	13
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	14-15
Generalisability	21	Discuss the generalisability (external validity) of the study results	14-15
Other information			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	15

*Give information separately for exposed and unexposed groups.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at <http://www.strobe-statement.org>.

EJOG Highlights

Pregnancy outcomes following different types of bariatric surgery: a national cohort study

- Maternal and perinatal outcomes differ according to type of bariatric surgery
- Serious surgical complications are more common in women with gastric bypass than gastric banding
- Choice of bariatric surgery should take into account the impact of surgery on future pregnancy