

EVIDENCE-BASED SYSTEMATIC REVIEWS

Ten-Year Implant Survival and Functional Outcomes Following Combined Medial Unicompartmental Knee Arthroplasty and Anterior Cruciate Ligament Reconstruction

A Systematic Review and Call for High-Quality Comparative Trials

Fardis Vosoughi, MD, Iman Menbari Oskouie, MD, MPH, Amir Kasaeian, PhD, Masoud Imani, Tianyi David Luo, MD, PhD, Abtin Alvand, PhD, FRCS, and Pooya Vahedi, MD

Investigation performed at Shariati Hospital, Tehran University of Medical Sciences (TUMS), Tehran, Iran

Background: Managing medial compartment knee osteoarthritis (OA) with anterior cruciate ligament (ACL) deficiency, particularly in younger, active patients, remains challenging. Medial unicompartmental knee arthroplasty (UKA) combined with ACL reconstruction (ACLR) (UKACL) has gained interest, yet outcomes remain incompletely defined. This systematic review aims to evaluate the clinical effectiveness, implant survivorship, complications, and patient-reported outcomes after combined UKA and ACLR in end-stage medial OA with ACL deficiency.

Methods: We systematically searched PubMed, Embase, Scopus, and Web of Science. Studies reporting outcomes of UKA performed with ACLR were included; case reports, technical notes, and biomechanical studies were excluded. Risk of bias was assessed with ROBINS-I V2. We extracted clinical outcomes, implant survival, complications, radiographic findings, and validated functional scores.

Results: Fourteen studies comprising 353 patients met the inclusion criteria. Reported survivorship consistently exceeded 90% at 10 years. Ten revisions were reported, most commonly for lateral OA progression. Overall complication rate was 9.06% with no difference between mobile-bearing and fixed-bearing designs. Mobile-bearing implants had a slightly higher bearing dislocation risk, whereas fixed-bearing designs showed marginally higher polyethylene wear. Functional outcomes improved across studies.

Conclusion: Combined UKA and ACLR appears effective for younger, active patients with isolated medial OA and ACL deficiency, yielding high survivorship and consistent functional gains. Given heterogeneity among studies, high-quality, long-term randomized trials are needed to refine patient and implant selection.

Level of Evidence: Level IV, systematic review of nonrandomized studies. See Instructions for Authors for a complete description of levels of evidence.

Introduction

Anterior cruciate ligament (ACL) injury increases knee osteoarthritis (OA) risk 4-fold to 6-fold¹. ACL tears predominantly affect active individuals younger than 30 years and

predispose to early OA in the 30 to 50 age range with pain, functional impairment, and reduced quality of life^{2,3}. Given the high incidence of ACL rupture, clarifying factors that drive postinjury OA is essential⁴.

Disclosure: The **Disclosure of Potential Conflicts of Interest** forms are provided with the online version of the article (<http://links.lww.com/JBJSOA/B92>).

Copyright © 2026 The Authors. Published by The Journal of Bone and Joint Surgery, Incorporated. All rights reserved. This is an open access article distributed under the [Creative Commons Attribution License 4.0 \(CCBY\)](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Unicompartmental knee arthroplasty (UKA) is a well-established and excellent option for end-stage medial OA, with advantages over total knee arthroplasty (TKA)⁵. Optimal management of concurrent medial OA and ACL deficiency remains unsettled⁶. Goodfellow et al. reported worse outcomes and lower survival rates following mobile-bearing UKA in ACL-deficient (ACLD) knees⁷. Traditional contraindications to UKA in the ACLD knee reflect this concern⁸. More recent studies suggest similar results for medial UKA with and without ACL deficiency^{8,9}.

Younger high-demand patients present a treatment challenge for knee surgeons¹⁰. Several therapeutic approaches are suggested for management, including conservative care¹¹, HTO alone or with ACLR^{12,13}, UKA with ACLR (UKACL)¹⁴, isolated UKA¹⁵, and TKA¹⁶, yet consensus is lacking^{6,17}. The link between ACL deficiency and worse outcomes after UKA is primarily reported in studies of mobile-bearing UKAs published over 30 years ago⁸.

UKACL has gained popularity^{14,18,19}, and the choice of fixed versus mobile bearings remains debated^{20,21}. Considering the potential advantages of this technique and the growing interest in its applications, the purpose of this study was to perform a systematic review of the literature to evaluate the outcomes of combined medial UKA and ACLR in patients with end-stage symptomatic medial compartment OA and ACLD.

Methods

Search Strategy

A systematic literature search was conducted across PubMed, Embase, Scopus, and Web of Science from inception to July 2025 in accordance with Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines²². (Fig. 1) Search terms combined “unicompartmental knee arthroplasty,” “medial osteoarthritis,” and “ACL,” with database-specific strategies (Supplementary Table I). References of included studies were screened. The protocol was registered at the International Prospective Register Of Systematic Reviews (PROSPERO) (CRD420250652298). The search was conducted primarily by 2 independent reviewers (P.V. and I.M.), followed by full-text screening with disagreements resolved by a senior reviewer (F.V.).

Data Extraction and Eligibility Criteria

This review included all English and non-English studies published in peer-reviewed journals that reported on UKA performed concomitantly with ACLR in patients with medial knee osteoarthritis and ACL deficiency. Case reports, conference abstracts, symposiums, technical notes, letters, reviews, cadaveric and animal studies, biomechanical studies, and studies with unexplained treatment approaches were excluded. A specified data extraction form was used to extract pertinent details from included articles, encompassing study characteristics, patient data, treatment approaches, and outcome measures. Outcome assessments comprised the following factors:¹ implantation survival², bearing and graft type³, clinical evaluations including all subjective functional scores⁴, reported radiological data, and⁵ complications.

Risk of Bias (quality) Assessment

Two authors (P.V. and I.M.) independently assessed study quality using the Risk Of Bias In Non-randomized Studies—of Interventions version 2 (ROBINS-I V2) tool developed by Cochrane²³. Across domains of confounding, intervention classification, participant selection, deviations from intended interventions, missing data, outcome measurement, and selective reporting, disagreements were adjudicated by a third author (F.V.).

Data Analyses

All statistical analyses were performed using R software (version 4.0; R Foundation for Statistical Computing, Vienna, Austria)²⁴. Heterogeneity was evaluated with I^2 ²⁵. Forest plots were generated to visually display the pooled results, while funnel plots evaluated potential publication bias²⁶. The difference between mean preoperative and postoperative values was reported as mean differences (MD). Preoperative versus postoperative differences were reported as MD. Survival data were extracted from Kaplan–Meier curves using Plot Digitizer²⁷. Data synthesis and analysis, whenever possible, were performed using the “Meta Survival” package in R 4.2.1^{28,29}, to construct combined survival curves. Given the non-randomized design, heterogeneity, risk of bias, and methodological variability among the included studies, a meta-analysis was not performed for the overall outcomes.

Ethics Approval and Consent to Participate

This was a systematic review study using publicly accessible documents as evidence.

Availability of Data and Materials

The authors confirm that the data supporting the findings of this study are available within the article [and/or] its supplementary materials.

Result

Study Selection and Characteristics of the Included Studies

A total of 1,190 records (PubMed 226; Scopus 362; Embase 355; Web of Science 247) articles were identified (Fig. 1). After duplicate removal, 520 records underwent screening, yielding 31 full texts. Ten were excluded (4 with unclear treatment descriptions; 6 with insufficient or unavailable data). This resulted in 21 studies initially eligible for inclusion. Seven studies were found to involve overlapping patients resulting in 14 included studies.

Quality Assessment and Publication Bias

ROBINS-I V2 assessments are summarized in Fig. 1 of the supplementary files, and funnel plots are shown in Supplementary Fig. 2.

Participants Characteristics

Across 14 studies, 353 patients were included (Table I). Pain with instability was the most common presentation. All studies focused on isolated ACL deficiency with medial compartment OA; additional ligamentous injuries (posterior cruciate ligament, medial collateral ligament, lateral collateral ligament) were either not reported or listed as exclusion criteria. There




Fig. 1 PRISMA flow diagram for the study selection process *Seven studies were excluded due to overlapping patient populations with other included studies from the same institutions and recruitment periods; only one representative study from each overlapping group was retained.

was large heterogeneity with respect to study design. The mean age ranged from 46 to 58.9 years and follow-up of 1.7 to 14.6 years. The reported mean BMI ranged from 24.1 to 28.2 kg/m². The mean sample size was 25.2 (range 10-75).

Implant Survival

Three studies reported survivorship explicitly. Jaber et al.¹⁴ observed 100% at 5 years, 95.7% at 10 years, and 91.4% at 14.5 years. Kennedy et al.³⁴ reported 97.0% at 5 years, 92.3% at 10, and 15 years. Weston-Simons et al.³⁹ reported 92.7% at 5 and 8 years (Fig. 2). Of the remaining studies, 3 reported revision counts^{33,35,41}. Among the 6 studies with revision data^{14,33-35,39,41}, the 5-year cumulative revision rate was 3.7% (6 revisions among 161 patients), corresponding to a 5-year survival of

96.3%. Studies with less than 2 years of follow-up reported early function but were insufficient to address durability.

Complications

Ten UKA revisions occurred among 353 cases (Table II). Revision indications were lateral OA progression (5, 50%)^{14,34,39}, periprosthetic joint infection (3 cases, 30%)^{34,39,41}, and aseptic loosening (2, 20%)^{33,35}. Reported nonrevision complications included 3 bearing dislocations (mobile-bearing)^{38,39}, 2 arthroscopic releases^{21,40}, 1 lateral meniscus injury³⁵, 1 thrombosis⁴⁰, 9 hemarthroses⁴⁰, and 1 arthroscopic loose-body removal³⁷. Five graft failures were reported in one study³². Additional revisions noted by Legnani et al.¹⁹ and Ventura et al.⁴² were not duplicated in our table due to cohort overlap with Ventura et al.³⁷. Foissey et al.⁴³

TABLE I Characteristics of the Included Studies							
First Author/Publication Year	Study design	Patients (M:F)	Mean age	Age (SD)	Mean Follow-Up	Follow-Up (SD)	Mean BMI
Tang et al., 2024 ³⁰	Retrospective cohort	13 (3:10)	58.9	4.2	7.14	2.45	NR
Jaber et al., 2023 ¹⁴	Retrospective case series	23 (18:5)	48	6.25	10	2.12	NR
Aslan and Çevik, 2022 ³¹	Retrospective case series	12	NR	NR	3.8	0.36	NR
Kurien et al., 2022 ³²	Retrospective case series	24 (16:8)	48.8	8.17	5.1	2.87	24.1
Derreveaux et al., 2022 ³³	Retrospective cohort	20 (7:13)	58	6.7	2.5	1.41	26.5
Kennedy et al., 2019 ³⁴	Prospective case series	75 (59:16)	52.6	8.75	6.4	3.5	28.2
Iriberrri et al., 2019 ³⁵	Retrospective case series	8 (5:3)	52	4.5	14.6	2.93	NR
Tecame et al., 2019 ³⁶	Retrospective case series	24 (20:4)	Mobile: 47.8 Fixed: 48.4	Mobile: 3 Fixed: 2.75	Mobile:4.41 Fixed:3.5	0.69 0.55	NR
Ventura et al., 2017 ³⁷	Retrospective case series	14 (9:5)	55	3.5	2.22	0.33	NR
Tian et al., 2015 ³⁸	Prospective Case Series	32 (12:20)	50	NR	4.58	2.92	NR
Weston-Simons et al., 2012 ³⁹	Prospective case series	51 (40:11)	51	7.75	5	0.18	NR
Tinius et al., 2007 ⁴⁰	Prospective Case Series	32 (25:7)	46	4.25	2.58	0.58	NR
Dervin et al., 2007 ²¹	Prospective case series	10 (5:5)	52	6	1.7	0.72	NR
Pandit et al., 2006 ⁴¹	Prospective cohort	15 (13:2)	49.8	6	2.8	0.45	NR

F = female, M = male, NR = not reported, and SD = standard deviation. Age and follow-up values are reported in years.

reported 2 arthroscopic arthrolyses for stiffness, excluded from the complication table to avoid duplication with Derreveaux et al³³.

Surgical Techniques

Six studies used a fixed-bearing configuration, 7 used mobile-bearing, and 1 incorporated both. Mean operative time,




Fig. 2 This figure shows the survival curves extracted from the included studies. The red line represents the data from Kennedy et al., the blue line corresponds to Weston-Simons et al., and the green line depicts the findings by Jaber et al.

TABLE II Complications and Revisions

First Author/ Publication Year	Follow-up/Last Follow-up (Year)	Total Patients Revision	UKA/ACL Revisions	UKA Revision Rate (%)	Implant Survival (%)	Cause of Revision	Year of Revision	Other Complications
Tang et al., 2024 ³⁰	Mean 7.14/11	13	0/0	0	100	None	None	None
Jaber et al., 2023 ¹⁴	Average 10/14.5	23	2/0	8.69	91.30	Trauma & lateral POA	6 and 12 years postoperatively	None
Aslan and Çevik 2022 ³¹	Mean 3.8/4.33	12	0/0	0	100	None	None	None
Kurien et al., 2022 ³²	Mean 5.1/12.8	24	0/5	0	79.2	None	None	1 patient had a failed Gortex ACL graft, 1 had a failed Carbon fiber graft, and the remaining 3 had failed hamstring grafts at 28 months post-UKR
Derreveau et al., 2022 ³³	Mean 2.5/5.09	20	1/0	5	95	aseptic loosening	at 5 years follow-up	None
Kennedy et al., 2019 ³⁴	Mean 6.4/15	75	3/0	4	96	2 lateral POA/1 PJI	one after 5 months/ 1 after 4 years/ 1 after 9.9 years	None
Iriberri et al., 2019 ³⁵	Mean 14.6/21.5	8	1/0	12.5	87.5	aseptic loosening	after 117 months	1 lateral meniscus tear repair
Tecame et al., 2019/M group ³⁶	Mean 4.41/NR	9	0/0	0	100	None	None	None
Tecame et al., 2019/F group ³⁶	Mean 3.5 years/ NR	15	0/0	0%	100%	None	None	None
Ventura et al., 2017 ³⁷	Mean 2.22/3.33	14	0/0	0%	100	None	None	1 arthroscopic loose body removal (16 months post-op)
Tian et al., 2015 ³⁸	Mean 4.58/9.3	32	0/0	0	100	None	None	2 Mobile bearing dislocation (exchanged)
Weston-Simons et al., 2012 ³⁹	Mean 5/10	51	2/0	3.84	96.15	1 PJI/ 1 lateral POA	NR	1 bearing dislocation (exchanged)
Tinius et al., 2007 ⁴⁰	Mean 2.58/3.16	32	0/0	0	100	None	None	9 hemarthrosis (aspirated), 1 thrombosis in the second postoperative year, 1 arthroscopic release
Dervin et al., 2007 ²¹	Median 1.7/9	10	0/0	0	100	None	None	1 arthroscopic release
Pandit et al., 2006 ⁴¹	Mean 2.8/4.3	15	1/0	6.66	93.33	1 PJI	NR	None

ACL = anterior cruciate ligament, and UKA = unicompartmental knee arthroplasty.

reported in 6 studies, ranged from 88.5 to 105 minutes^{21,30,37,40}. Most prostheses were cemented; 1 study included cemented, cementless, and hybrid fixation³¹, and 2 did not specify cementation^{21,34}.

ACLR was most commonly used hamstring autograft (11 studies)^{14,21,30,31,33-38,40}; 2 studies used hamstring or bone–patellar tendon–bone autograft^{39,41}, and 1 used allograft³²). Eight studies performed single-stage UKACL^{14,30-32,35-38}, 4 used staged

procedures^{21,33,39,40}; 2 studies tailored the choice of staged versus single-stage surgery to patient-specific clinical factors^{34,41}.

Common implant systems were Smith & Nephew (7 studies)^{21,33,35-38,41}, Zimmer (4)^{30-32,34}, and Oxford (2)^{14,39}. Surgical sequencing varied across reports: some surgeons performed ACLR after UKA, others started with ACLR, and several used a combined or overlapping approach. In 7 studies, the tibial tunnel guide angle was 55°^{21,30,31,35-38}. Intra-articular tunnel position was described as slightly lateral in 3 studies^{31,36,37} and anterior in 2 studies^{14,33}. Graft fixation most commonly used a femoral Endo Button with tibial interference screw (Table III).

Rehabilitation

Most rehabilitation protocols were brace-free with immediate or day 1 knee motion. Early goals emphasized full extension and progressive flexion; isometric quadriceps exercises began within 6 to 24 hours. Partial weight-bearing from day 1 typically continued for 4 to 6 weeks. Some protocols limited flexion to 90° initially, then advanced as tolerated. Proprioceptive training often began around week 4. Programs were generally UKA-based with adjustments for return to sport; continuous passive motion and pain-adapted physiotherapy were variably used.

Radiographic Outcomes

Radiographic data showed changes in knee alignment and stability. Tang et al.³⁰ noted hip-knee ankle angle improvement from 174.07° ± 2.06 to 177.79° ± 1.25 and a posterior tibial slope (PTS) decrease from 7.93° ± 1.07 to 4.57° ± 0.94. Tecame et al.³⁶ likewise observed decreases in PTS and varus angles with both bearing types. Aslan et al.³¹ reported varus reduction from 3.6° ± 1° to 2.6° ± 1°, while Tian et al.³⁸ reported maintained varus and valgus angles and posterior tibial prosthesis tilt at the last follow-up. Derreveau³³ described preoperative mechanical femorotibial angles averaging 178° ± 6°.

Functional Outcomes and PROMs

Validated PROMs included the American Knee Society Score (AKSS; objective and functional), the Oxford Knee Score (OKS), the Knee Society Score (KSS; knee and function scores), the Tegner Activity Scale, and the Western Ontario and McMaster Universities Osteoarthritis Index. Three studies (80 patients) reported objective AKSS improvements (MD 32-44; I² = 33.4%). The functional AKSS component improved by 10 to 13 points across 3 studies (80 patients) with no heterogeneity (I² = 0.0%). For OKS, 3 fixed-bearing studies (51 patients) reported MD 14.2 to 26; 5 mobile-bearing studies (185 patients) reported MD 12 to 19.1. Three studies (84 patients) reported KSS improvements: knee score MD 32.8 to 54.2 (I² = 93.6%) and function score MD 28.8 to 48.6 (I² = 95%). Tegner improved by 0.8 to 2.2 points across 7 studies (234 patients) with high heterogeneity (I² = 80.4%) (Supplementary Fig. 3).

Beyond PROMs, multiple studies showed significant functional improvements across KOOS subscales (except Sports), OKS, EQ-5D, EQ-VAS, Lysholm, UCLA, IKDC-

2000, Tegner activity, and VAS pain^{14,31}. Kurien et al. also reported improved knee stability³². Objective testing supported restored ligament function: Tecame et al. found negative Lachman and anterior drawer signs³⁶, and Ventura et al. observed clinically acceptable KT-1000 side-to-side differences³⁷. Overall, UKACL improved pain and function while restoring objective stability (Supplementary Table II).

Discussion

This systematic review evaluated medial UKA performed in conjunction with ACLR (UKACL), assessing implant survivorship, functional, radiological, and patient-reported outcomes. The main finding of this study was that UKACL is associated with implant survival consistently exceeding 90% at 10 years in appropriately selected patients. Kennedy et al. and Jaber et al. reported 10-year survivorship of 92.3% and 95.7%, respectively^{14,34}, and Weston-Simons et al. reported 92.7% at 5 to 8 years³⁹. Across 6 studies, the 5-year cumulative revision rate was 3.7% (5-year survival 96.3%). These results are comparable with those of isolated UKA in ACL-intact (ACLI) knees (5- and 10-year survival, 90.5% and 83.5%, respectively)⁴⁴, challenging the presumption that ACL deficiency necessitates TKR^{8,45}.

Historically, ACL deficiency was considered a contraindication to UKA³ due to concerns over increased anteroposterior and rotational instability leading to early implant failure. Fixed-bearing designs may mitigate sagittal laxity, but rotational laxity persists⁴⁶. As noted by Zumbunn et al.⁴⁶, isolated UKA in ACLD patients—especially older, low-demand individuals—may be successful only in the presence of clinical rotational stability. However, this conclusion derives from an elderly cohort (mean age >60 years), creating a selection bias in the existing literature.

By contrast, all 14 UKACL studies in this review reported mean ages <60 years. Mancuso et al.⁶ found superior midterm survival with UKACL (97%) versus UKA in untreated ACLD knees (88%), noting that ACLD-UKA cohorts were older (>60 years) while UKACL cohorts were younger (<55 years). Du et al. reported similar outcomes for ACLD-UKA and ACLI-UKA in 8 studies; however, all included populations had mean ages older than 60 years⁹. Thus, the literature supports ACLD-UKA primarily in older patients, whereas the current review supports UKACL in younger individuals. Potential explanations include stabilizing effects of osteophytes and advanced OA in the elderly and better tolerance of ACL deficiency in low-demand populations.

This study shows consistent improvements across several PROMs (AKSS, KSS, Tegner). Despite variability in reporting functional scores, meaningful conclusions about PROMs remain evident. Eight studies showed consistent OKS improvements of 12 to 26 points^{30-32,34,37-39,41}, and 7 studies reported Tegner gains of 0.8 to 2.2 points^{14,32,34,37-39,41} (Supplementary Fig. 3). These results align with prior literature and highlight the clinical value of restoring ligamentous stability in these patients.

TABLE III Surgical Techniques

First Author/Publication Year	Bearing Type	Graft Type	Cementation	Staged	Guide Angle	Type of Prosthesis	Mean Surgery Duration (Min)
Tang et al., 2024 ³⁰	Fixed	Hamstring	Cemented	Single	55	Zimmer Biomet	88.5
Jaber et al., 2023 ¹⁴	Mobile	Hamstrings	Cemented	Single	NR	Oxford UKA	NR
Aslan and Çevik, 2022 ³¹	Mobile	Hamstrings	Cementless (5) Cemented (4) Hybrid (3)	Single	55	Zimmer	NR
Kurien et al., 2022 ³²	Fixed	Hamstrings BPTB Allograft	Cemented	Single	NR	Physica ZUK (Lima Corporate, Udine, Italy)	NR
Derreveau et al., 2022 ³³	Fixed	Hamstrings	Cemented	11 single 9 staged	NR	Journey II Uni (Smith and Nephew) in 26 cases HLS Uni evolution (Tornier) in 8 cases	NR
Kennedy et al., 2019 ³⁴	Mobile	Hamstrings	Cementless (NR) Cemented (NR)	Single or Staged, based on the complaint	NR	Zimmer Biomet	NR
Iriberry et al., 2019 ³⁵	Fixed	Hamstrings	Cemented	Single	55	Smith & Nephew	NR
Tecame et al., 2019 ³⁶	Fixed: 15 Mobile: 9	Hamstrings	Cemented	Single	55	Smith & Nephew	NR
Ventura et al., 2017 ³⁷	Fixed	Hamstrings	Cemented	Single	55	Smith & Nephew	98
Tian et al., 2015 ³⁸	Mobile	Hamstrings	Cemented	Single	55	Smith & Nephew	NR
Weston-Simons et al., 2012 ³⁹	Mobile	BPTB (?) Hamstring (?)	Cemented	33 single 18 staged	NR	Oxford UKA	NR
Tinius et al., 2007 ⁴⁰	Fixed	Hamstrings	Cemented	21 single 11 staged	NR	Repicci-II; Eius	95
Dervin et al., 2007 ²¹	Mobile	Hamstrings	NR	9 single 1 staged	55	Smith & Nephew	105
Pandit et al., 2006 ⁴¹	Mobile	Hamstring (11) 3 BPTB (3)	Cemented	4 patients simultaneously 11 patients had ACL reconstruction first, and arthroplasty later if needed	NR	Smith & Nephew	NR

BPTB = bone patellar tendone bone autograft, NR = not reported, SD = standard deviation, and UKA = unicompartmental knee arthroplasty.

Regarding implant design, although biomechanical studies suggest mobile-bearing UKA may better replicate normal kinematics and reduce wear^{20,47}, our review did not find consistent superiority of mobile-bearing implants in terms of clinical outcomes. This suggests that in the setting of UKACL, the theoretical biomechanical advantages of mobile bearings may not translate into clinically meaningful gains, possibly due to the remaining laxity after ACLR. Collectively, current evidence supports UKACL as a viable option for younger, active patients with isolated medial OA and ACL deficiency³⁴, with several studies reporting excellent subjective and objective outcomes up to 8 to 10 years^{19,42,48-50}. Kennedy et al.³⁴ also reported low revision and conversion rates with both staged and simultaneous procedures (Supplementary Fig. 4).

Despite promising findings, the current review has serious limitations. Heterogeneity in surgical techniques, implant designs, and patient selection criteria hindered meta-analytic synthesis. Publication bias remains a concern despite comprehensive searching. Long-term, adequately powered randomized controlled trials comparing UKACL to isolated UKA are needed. Future work should incorporate preoperative stability, age and activity level, implant design, and radiographic alignment into predictive models to refine indications and optimize outcomes.

Conclusion


Combined medial UKA and ACLR is associated with sustained functional improvement (AKSS, OKS, KSS, Tegner)

and exceeded 90% 10-year survivorship in appropriately selected patients. For younger, active individuals with isolated medial OA and ACL deficiency, UKACL represents a viable surgical option. Further high-quality prospective studies, including randomized trials, are warranted to establish optimal patient selection criteria, refine surgical technique, and benchmark outcomes against alternative strategies.

Funding

The research presented in this manuscript received no external form of funding.

Appendix

 Supporting material provided by the authors is posted with the online version of this article as a data supplement at [jbjs.org \(http://links.lww.com/JBJSOA/B93\)](http://links.lww.com/JBJSOA/B93). This content was not copyedited or verified by JBJS. ■

Fardis Vosoughi, MD¹
Iman Menbari Oskouie, MD, MPH²
Amir Kasaeian, PhD³

Masoud Imani⁴
Tianyi David Luo, MD, PhD⁵
Abtin Alvand, PhD, FRCS⁶
Pooya Vahedi, MD⁷

¹Department of Orthopedic and Trauma Surgery, Shariati Hospital and School of Medicine, Tehran University of Medical Sciences (TUMS), Tehran, Iran

²Urology Research Center, Tehran University of Medical Sciences, Tehran, Iran

³Clinical Research Development Unit, Shariati Hospital, Tehran University of Medical Sciences, Tehran, Iran

⁴Department of Biostatistics, School of Public Health, Iran University of Medical Sciences, Tehran, Iran

⁵Indiana Joint Replacement Institute, Fort Wayne, Indiana

⁶Nuffield Department of Orthopaedics (NOC), University of Oxford, Oxford, United Kingdom

⁷Student Research Committee, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

E-mail address for P. Vahedi: Vahedi.pooya@gmail.com

References

- Poulsen E, Goncalves GH, Bricca A, Roos EM, Thorlund JB, Juhl CB. Knee osteoarthritis risk is increased 4-6 fold after knee injury—A systematic review and meta-analysis. *Br J Sports Med.* 2019;53(23):1454-63.
- Friel NA, Chu CR. The role of ACL injury in the development of posttraumatic knee osteoarthritis. *Clin Sports Med.* 2013;32(1):1-12.
- Lohmander LS, Englund PM, Dahl LL, Roos EM. The long-term consequence of anterior cruciate ligament and meniscus injuries: osteoarthritis. *Am J Sports Med.* 2007;35(10):1756-69.
- Rodríguez-Merchan EC, Encinas-Ullan CA. Knee osteoarthritis following anterior cruciate ligament reconstruction: frequency, contributory elements, and recent interventions to modify the route of degeneration. *Arch Bone Jt Surg.* 2022;10(11):951-8.
- Wilson HA, Middleton R, Abram SGF, Smith S, Alvand A, Jackson WF, Bottomley N, Hopewell S, Price AJ. Patient relevant outcomes of unicompartmental versus total knee replacement: systematic review and meta-analysis. *BMJ.* 2019;364:l352.
- Mancuso F, Hamilton TW, Kumar V, Murray DW, Pandit H. Clinical outcome after UKA and HTO in ACL deficiency: a systematic review. *Knee Surg Sports Traumatol Arthrosc.* 2016;24(1):112-22.
- Goodfellow JW, Kershaw CJ, Benson MK, O'Connor JJ. The Oxford Knee for unicompartmental osteoarthritis. The first 103 cases. *The J Bone Joint Surg Br.* 1988;70-B(5):692-701.
- Kozinn SC, Scott R. Unicompartmental knee arthroplasty. *J Bone Jt Surg.* 1989;71(1):145-50.
- Du G, Qiu H, Zhu J, Wang H, Xiao Q, Zhang Z, Lin X, Zheng G. No difference unicompartmental knee arthroplasty for medial knee osteoarthritis with or without anterior cruciate ligament deficiency: a systematic review and meta-analysis. *J Arthroplasty.* 2023;38(3):586-93.e1.
- van der List JP, Chawla H, Villa JC, Pearle AD. The role of patient characteristics on the choice of unicompartmental versus total knee arthroplasty in patients with medial osteoarthritis. *J Arthroplasty.* 2017;32(3):761-6.
- van der Woude JAD, Wiegant K, van Heerwaarden RJ, Spruijt S, van Roermund PM, Custers RJH, Mastbergen SC, Lafeber F. Knee joint distraction compared with high tibial osteotomy: a randomized controlled trial. *Knee Surg Sports Traumatol Arthrosc.* 2017;25(3):876-86.
- Herman BV, Giffin JR. High tibial osteotomy in the ACL-deficient knee with medial compartment osteoarthritis. *J Orthop Traumatol.* 2016;17(3):277-85.
- Jin C, Song EK, Jin QH, Lee NH, Seon JK. Outcomes of simultaneous high tibial osteotomy and anterior cruciate ligament reconstruction in anterior cruciate ligament deficient knee with osteoarthritis. *BMC Musculoskelet Disord.* 2018;19(1):228.
- Jaber A, Kim CM, Barié A, Streit M, Schmitt H, Clarius M, Merle C, Bangert Y. Combined treatment with medial unicompartmental knee arthroplasty and anterior cruciate ligament reconstruction is effective on long-term follow-up. *Knee Surg Sports Traumatol Arthrosc.* 2023;31(4):1382-7.
- Kagan R, Anderson MB, Bailey T, Hofmann AA, Pelt CE. Ten-Year survivorship, patient-reported outcomes, and satisfaction of a fixed-bearing unicompartmental knee arthroplasty. *Arthroplasty Today.* 2020;6(2):267-73.
- van der List JP, Sheng DL, Kleeblad LJ, Chawla H, Pearle AD. Outcomes of cementless unicompartmental and total knee arthroplasty: a systematic review. *Knee.* 2017;24(3):497-507.
- Vajapey SP, Alvarez PM, Greco NJ, Chonko DJ. Medial osteoarthritis in an ACL-deficient knee: a critical analysis review. *JBJS Rev.* 2021;9(3):e20.
- Adravanti P, Budhiparama NC, Berend KR, Thienpont E. ACL-deficient knee and unicompartmental OA: state of the art. *J ISAKOS.* 2017;2(3):162-70.
- Legnani C, Borgo E, Macchi V, Terzaghi C, Ventura A. Unicompartmental knee replacement combined with anterior cruciate ligament reconstruction provides comparable results to total knee replacement with no increased risk of complications. *SICOT J.* 2024;10:10.
- Ko YB, Gujarathi MR, Oh KJ. Outcome of unicompartmental knee arthroplasty: a systematic review of comparative studies between fixed and Mobile bearings focusing on complications. *Knee Surg Relat Res.* 2015;27(3):141-8.
- Dervin GF, Conway AF, Thurston P. Combined anterior cruciate ligament reconstruction and unicompartmental knee arthroplasty: surgical technique. *Orthopedics.* 2007;30(5 suppl):39-41.
- Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L, Tetzlaff JM, Akl EA, Brennan SE, Chou R, Glanville J, Grimshaw JM, Hróbjartsson A, Lalu MM, Li T, Loder EW, Mayo-Wilson E, McDonald S, McGuinness LA, Stewart LA, Thomas J, Tricco AC, Welch VA, Whiting P, Moher D. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ.* 2021;372:n71.
- Sterne JA, Hernán MA, Reeves BC, Savović J, Berkman ND, Viswanathan M, Henry D, Altman DG, Ansari MT, Boutron I, Carpenter JR, A-W Chan, Churchill R, Deeks JJ, Hróbjartsson A, Kirkham J, Jüni P, Loke YK, Pigott TD, Ramsay CR, Regidor D, Rothstein HR, Sandhu L, Santaguida PL, Schünemann HJ, Shea B, Shrier I, Tugwell P, Turner L, Valentine JC, Waddington H, Waters E, Wells GA, Whiting PF, Higgins JP. ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ.* 2016;355:i4919.
- Team RDC. R: A language and environment for statistical computing. 2010.
- Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med.* 2002;21(11):1539-58.

26. Vosoughi F, Menbari Oskouie I, Rahimdoost N, Kasaeian A, Sherafat Vaziri A. Intrarater and inter-rater reliability of tibial Plateau fracture classifications: systematic review and meta-analysis. *JBJS Open Access*. 2024;9(4):e23.00181.
27. Drevon D, Fursa SR, Malcolm AL. Inter-coder reliability and validity of Web-PlotDigitizer in extracting graphed data. *Behav Modif*. 2017;41(2):323-39.
28. Shahzad M, Iqbal Q, Amin MK, Kasaeian A, Oskouie IM, Warraich SZ, Yu J, Anwar I, Jaglal M, Mushtaq MU. Outcomes of allogeneic hematopoietic stem cell transplantation in patients aged 70 years and older: a systematic review and meta-analysis. *Transplant Cell Ther*. 2025;31(3):172.e1-172.e13.
29. Shahzad M, Amin MK, Khalid MF, Kasaeian A, Oskouie IM, Yu J, Warraich SZ, Basharat A, Butt A, Zaidi M, Anwar I, Jaglal MV, Mushtaq MU. Outcomes with allogeneic hematopoietic stem cell transplantation in therapy related myeloid neoplasms: a systematic review and meta-analysis. *Clin Lymphoma Myeloma Leuk*. 2025;25(5):e319-e335.
30. Tang H, Shan PC, Zhang HM. Mid-term effects of unicompartmental knee arthroplasty combined with anterior cruciate ligament reconstruction for the medial compartment osteoarthritis with anterior cruciate ligament deficiency by proportioning with the pure medial compartment osteoarthritis [in Chinese]. *Zhonghua Yi Xue Za Zhi*. 2024;104(5):350-6.
31. Asian H, Çevik HB. Outcomes of combined unicompartmental knee arthroplasty and anterior cruciate ligament reconstruction. *J Knee Surg*. 2022;35(10):1087-90.
32. Kurien T, Stragier B, Senevirathna S, Geutjens G. Excellent outcomes with combined single stage Physica ZUK medial unicompartment knee replacement and anterior cruciate ligament reconstruction results in young, active patients with instability and osteoarthritis with a mean follow up of 5 years. *Knee*. 2022;36:114-9.
33. Derreveau V, Schmidt A, Shatrov J, Sappey-Marinière E, Bataillier C, Servien E, Lustig S. Combined procedures with unicompartmental knee arthroplasty: high risk of stiffness but promising concept in selected indications. *SICOT J*. 2022;8:4.
34. Kennedy JA, Molloy J, Mohammad HR, Mellon SJ, Dodd CAF, Murray DW. Mid- to long-term function and implant survival of ACL reconstruction and medial Oxford UKR. *Knee*. 2019;26(4):897-904.
35. Iriberrí I, Suau S, Payán L, Aragón JF. Long-term deterioration after one-stage unicompartmental knee arthroplasty and anterior cruciate ligament reconstruction. *Musculoskelet Surg*. 2019;103(3):251-6.
36. Tecame A, Savica R, Rosa MA, Adevranti P. Anterior cruciate ligament reconstruction in association with medial unicompartmental knee replacement: a retrospective study comparing clinical and radiological outcomes of two different implant design. *Int Orthop*. 2019;43(12):2731-7.
37. Ventura A, Legnani C, Terzaghi C, Iori S, Borgo E. Medial unicompartmental knee arthroplasty combined to anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc*. 2017;25(3):675-80.
38. Tian S, Wang B, Liu L, Ha C, Yuanhe W, Liu J, Li Q, Yang X, Sun K. Anterior cruciate ligament reconstruction in treatment of osteoarthritis and anterior cruciate ligament injury [in Chinese]. *Zhongguo Xiu Fu Chong Jian Wai Ke Za Zhi*. 2015;29(12):1457-61.
39. Weston-Simons JS, Pandit H, Jenkins C, Jackson WFM, Price AJ, Gill HS, Dodd CAF, Murray DW. Outcome of combined unicompartmental knee replacement and combined or sequential anterior cruciate ligament reconstruction. *J Bone Joint Surg Br*. 2012;94-B(9):1216-20.
40. Tinius M, Ecker TM, Klima S, Tinius W, Josten C. [Minimally invasive unicompartmental knee arthroplasty with simultaneous ACL reconstruction: treatment of medial compartment osteoarthritis and cruciate ligament defect]. *Unfallchirurg*. 2007;110(12):1030-8.
41. Pandit H, Beard DJ, Jenkins C, Kimstra Y, Thomas NP, Dodd CA, Murray DW. Combined anterior cruciate reconstruction and Oxford unicompartmental knee arthroplasty. *J Bone Joint Surg Br*. 2006;88-B(7):887-92.
42. Ventura A, Legnani C, Terzaghi C, Macchi V, Borgo E. Unicompartmental knee replacement combined to anterior cruciate ligament reconstruction: mid-term results. *J Knee Surg*. 2020;33(11):1152-6.
43. Foissey C, Bataillier C, Shatrov J, Servien E, Lustig S. Is combined robotically assisted unicompartmental knee arthroplasty and anterior cruciate ligament reconstruction a good solution for the young arthritic knee? *Int Orthop*. 2023;47(4):963-71.
44. Ekhtiari S, Bozzo A, Madden K, Winemaker MJ, Adili A, Wood TJ. Unicompartmental knee arthroplasty: survivorship and risk factors for revision: a population-based cohort study with minimum 10-Year Follow-up. *J Bone Jt Surg*. 2021;103(23):2170-6.
45. Hamilton TW, Pandit HG, Jenkins C, Mellon SJ, Dodd CAF, Murray DW. Evidence-based indications for mobile-bearing unicompartmental knee arthroplasty in a consecutive cohort of thousand knees. *J Arthroplasty*. 2017;32(6):1779-85.
46. Zumbunn T, Schütz P, von Knoch F, Preiss S, List R, Ferguson SJ. Medial unicompartmental knee arthroplasty in ACL-Deficient knees is a viable treatment option: in vivo kinematic evaluation using a moving fluoroscope. *Knee Surg Sports Traumatol Arthrosc*. 2020;28(6):1765-73.
47. Banks SA, Fregly BJ, Boniforti F, Reinschmidt C, Romagnoli S. Comparing in vivo kinematics of unicompartmental and bi-unicompartmental knee replacements. *Knee Surg Sports Traumatol Arthrosc*. 2005;13(7):551-6.
48. Tinius M, Hepp P, Becker R. Combined unicompartmental knee arthroplasty and anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc*. 2012;20(1):81-7.
49. Pandit H, Van Duren BH, Gallagher JA, Beard DJ, Dodd CA, Gill HS, Murray DW. Combined anterior cruciate reconstruction and Oxford unicompartmental knee arthroplasty: in vivo kinematics. *Knee*. 2008;15(2):101-6.
50. Tian S, Wang B, Wang Y, Ha C, Liu L, Sun K. Combined unicompartmental knee arthroplasty and anterior cruciate ligament reconstruction in knees with osteoarthritis and deficient anterior cruciate ligament. *BMC Musculoskelet Disord*. 2016;17(1):327.